CAN THE DARK-MATTER DEFICIT IN THE HIGH-REDSHIFT GALAXIES EXPLAIN THE PERSISTENT DISCREPANCY IN HUBBLE CONSTANTS?

Yurii V. Dumin^{1,2}

¹P.K. Sternberg Astronomical Institute of M.V. Lomonosov Moscow State University Universitetskii prosp., 13, 119234, Moscow, Russia ²Space Research Institute of the Russian Academy of Sciences Profsoyuznaya str. 84/32, 117997, Moscow, Russia dumin@yahoo.com, dumin@sai.msu.ru

Abstract: One of hot topics in the last years is a systematic discrepancy in the determination of Hubble parameter by various methods. Namely, the values derived "directly" from the distance scale based on Cepheids and supernovae—and referring to the relatively "local" part of the Universe—are about 10% greater than the ones following from the analysis of the cosmic microwave background (CMB) radiation, which refers to the "global" scales. The most popular interpretation of this discord, widely discussed nowadays, is variation of the dark-energy equation-of-state parameter w. However, there might be a much simpler explanation, following from the recent observations of the rotation curves in the high-redshift galaxies. Namely, it was found that they have much smaller dark-matter halos than galaxies in the vicinity of us [Genzel, et al. Nature 543 (2017), 397]. Since both the dark and luminous matter possess the same dust-like equation of state and, therefore, their average cosmological densities evolve by the same way, our local neighborhood is not quite typical but rather overfilled with the dark matter. Then, the local value of the Hubble constant should be greater than the global one. Roughly speaking, a twofold excess of the dark matter in our local Universe would give just the above-mentioned 10% increase in the value of Hubble parameter.

Keywords: Hubble constant, dark matter, high-redshift galaxies

PACS: 98.80.Es, 95.35.+d, 95.36.+x, 98.62.Gq

Determination of the Hubble parameter H_0 is a long-standing problem in cosmology, lasting for almost a century; and the corresponding values varied in this period by an order of magnitude, 50 to 500 km s⁻¹ Mpc⁻¹ (see, for example, [1, 2] and references therein). Despite of considerable improvements, some discrepancies persist till now. The most notable of them is that the value of H_0 derived from the distance scale based on Cepheids and supernovae is 73.24 ± 1.74 km s⁻¹ Mpc⁻¹ and, for some calibration, can even be as large as 76.18 ± 2.37 km s⁻¹ Mpc⁻¹ [3]. On the other hand, the analysis based on measurements of the cosmic microwave background (CMB) by *Planck* satellite under assumption of the Λ CDM cosmological model leads to the values $H_0 = 66.88\pm0.91$ to 67.31 ± 0.96 km s⁻¹ Mpc⁻¹, depending on the method of data processing [4]. So, these numbers are about 10% less than in the first case.

The above-mentioned discrepancy between the "local" (by Cepheids) and "global" (by CMB) measurements of H_0 was clearly recognized in the recent years, and it is commonly attributed now either to the systematic errors (such as degeneracy between different quantities in the analysis of CMB) or to the uncertainty in the fitting parameters (e.g., the number and masses of neutrinos, etc.) [5, 6]. Yet another popular explanation is a modification of the dark-energy equation-of-state parameter w(where $p = w\rho$) [7, 8, 9, 10]; though the resulting values w < -1 look quite suspicious from the viewpoint of general physical principles.¹

However, from our point of view, the spread in values of H_0 can have a much more straightforward explanation, following from the recent observations of the rotation curves in distant galaxies [12, 13]: it was found that the amount of dark matter is considerably less in the vicinity of galaxies located at large redshifts, z = 0.6-2.6. Next, it should be kept in mind that due to the same dust-like equation of state $(w \approx 0)$ both for the luminous and dark matter, the ratio of their densities does not change with cosmological time. So, we have to conclude that this ratio should be substantially variable in space and, thereby, the Hubble parameter should be scale-dependent.

Really, according to the standard Friedmann equation [14]:

$$H_0 = \left[\frac{8\pi G}{3}\right]^{1/2} \left[\rho_{\rm de} + \langle \rho_{\rm dm} \rangle + \langle \rho_{\rm lm} \rangle\right]^{1/2},\tag{1}$$

where ρ_{de} is density of the dark energy, which is assumed to be perfectly uniform in space (i.e., described by the Λ -term), ρ_{dm} and ρ_{lm} are densities of the dark and luminous (baryonic) matter, and the angular brackets denote averaging over the given spatial scale. Then, ratio of the Hubble parameters at the local and global scales should be:²

$$\frac{H_0^{(\text{loc})}}{H_0^{(\text{glob})}} = \left[\frac{\Omega_{\text{de}} + \langle \Omega_{\text{dm}} \rangle^{(\text{loc})} + \langle \Omega_{\text{lm}} \rangle}{\Omega_{\text{de}} + \langle \Omega_{\text{dm}} \rangle^{(\text{glob})} + \langle \Omega_{\text{lm}} \rangle}\right]^{1/2},\tag{2}$$

where $\Omega_i = \rho_i / \rho_c$ are the corresponding densities normalized to the critical density at the global scale; and we assume that the luminous matter distribution is sufficiently uniform.

¹For example, the values of w somewhat greater than -1 (i.e., |w| < 1) could be easily attributed to the small-scale irregularities of the scalar field representing the "dynamic" dark energy [11], but such an effect cannot result in w < -1.

²For simplicity, we ignore here the curvature term that might appear at the local scales due to the non-uniform dark matter distribution.



Figure 1: Ratio of the Hubble parameters at the local and global scales $H_0^{(\text{loc})}/H_0^{(\text{glob})}$ as function of the local dark-matter density normalized to the global critical density $\langle \Omega_{\text{dm}} \rangle^{(\text{loc})}$ (diagonal black curve). Horizontal red and vertical blue strips show the range of observable values. Horizontal and vertical green lines correspond to the trivial case when $\langle \Omega_{\text{dm}} \rangle^{(\text{loc})} = \langle \Omega_{\text{dm}} \rangle^{(\text{glob})}$.

This ratio of the Hubble parameters is plotted in Fig. 1 as function of the local dark-matter density at the standard cosmological parameters: $\Omega_{\rm de} = 0.69$, $\langle \Omega_{\rm dm} \rangle^{\rm (glob)} = 0.26$, and $\langle \Omega_{\rm lm} \rangle = 0.05$. The range of observed values of $H_0^{\rm (loc)}/H_0^{\rm (glob)}$ is shown by the horizontal red strip. Then, the corresponding normalized densities of the dark matter in our local cosmological neighborhood should be in the range $\langle \Omega_{\rm dm} \rangle^{\rm (loc)} = 0.44 - 0.56$ (vertical blue strip), i.e. about two times greater than globally.

In fact, Genzel, et al. [12] already emphasized that at the global scales the dark matter should play a smaller part than in the local Universe. So, from our point of view, the systematic discrepancy between the "local" and "global" values of the Hubble parameter is just a direct consequence of the above-mentioned fact.

Finally, let us mention that a number of papers published in the recent years made just the opposite statement as compared to [12]: namely, that there is a considerable deficit of luminous and dark matter in our local cosmological neighborhood. For example, Makarov & Karachentsev [15] and Karachentsev [16] found $\Omega_{\rm dm} + \Omega_{\rm lm} =$ 0.08 ± 0.02 in the sphere of radius $z \sim 0.01$ around us, which is over three times smaller than the standard value in the Λ CDM model. Unfortunately, their analysis involved a lot of model assumptions. On the other hand, the work by Genzel, et al. [12], which is based solely on the galaxy rotation curves, seems to be much less model-dependent; and the corresponding results on the deficit of dark matter in the high-redshift (rather than local) galaxies look more reliable.

Acknowledgements

I am grateful to Yu.V. Baryshev, S.M. Kopeikin, M. Křížek, A. Maeder, and M. Nowakowski for valuable discussions of the problem of small-scale Hubble expansion, as well as to the referee for a few important remarks.

References

- [1] Freedman, W.L.: Cosmology at a crossroads. Nature Astron. 1 (2017), 0121.
- [2] Ryden, B.: A constant conflict. Nature Phys. 13 (2017), 314.
- [3] Riess, A.G., Macri, L.M., Hoffmann, S.L., et al.: A 2.4% determination of the local value of the Hubble constant. Astrophys. J. 826 (2016), 56.
- [4] Aghanim, N., Ashdown, M., Aumont, J., et al. (Planck Collaboration): Planck intermediate results: XLVI. Reduction of large-scale systematic effects in HFI polarization maps and estimation of the reionization optical depth. Astron. Astrophys. 596 (2016), A107.
- [5] Verde, L., Protopapas, P., and Jimenez, R.: Planck and the local Universe: Quantifying the tension. Phys. Dark Univ. **2** (2013), 166.
- [6] Bernal, J.L., Verde, L., and Riess, A.G.: The trouble with H_0 . J. Cosmol. Astropart. Phys. **10** (2016), 019.
- [7] Huang, Q.-G. and Wang, K.: How the dark energy can reconcile Planck with local determination of the Hubble constant. Eur. Phys. J. C 76 (2016), 506.
- [8] Liu, Z.-E., Yu, H.-R., Zhang, T.-J., and Tang, Y.-K.: Direct reconstruction of dynamical dark energy from observational Hubble parameter data. Phys. Dark Univ. 14 (2016), 21.
- [9] Di Valentino, E.: Crack in the cosmological paradigm. Nature Astron. 1 (2017), 569.
- [10] Zhao, G.-B., Raveri, M., Pogosian, L., et al.: Dynamical dark energy in light of the latest observations. Nature Astron. 1 (2017), 627.
- [11] Linde, A.D.: The inflationary Universe. Rep. Prog. Phys. 47 (1984), 925.
- [12] Genzel, R., Förster Schreiber, N.M., Übler, H., et al.: Strongly baryondominated disk galaxies at the peak of galaxy formation ten billion years ago. Nature 543 (2017), 397.

- [13] Swinbank, M.: Distant galaxies lack dark matter. Nature 543 (2017), 318.
- [14] Olive, K.A. and Peacock, J.A.: Big-bang cosmology. In: Particle Data Group: *Review of Particle Physics.* Chin. Phys. C 40 (2016), 100001, p. 355.
- [15] Makarov, D. and Karachentsev, I.: Galaxy groups and clouds in the local $(z \sim 0.01)$ Universe. Mon. Not. R. Astron. Soc. **412** (2011), 2498.
- [16] Karachentsev, I.D.: Missing dark matter in the local Universe. Astrophys. Bull. 67 (2012), 123.