

# A Simple Generalized Gradient Approximation for the Non-interacting Kinetic Energy Density Functional

K. Luo,<sup>1,\*</sup> V.V. Karasiev,<sup>2,†</sup> and S.B. Trickey<sup>3,‡</sup>

<sup>1</sup>Quantum Theory Project, Department of Physics, University of Florida, Gainesville, FL 32611

<sup>2</sup>Laboratory for Laser Energetics, University of Rochester, Rochester, NY 14623

<sup>3</sup>Quantum Theory Project, Department of Physics and Department of Chemistry, University of Florida, Gainesville, FL 32611

(Dated: 22 June 2018)

A simple, novel, non-empirical, constraint-based orbital-free generalized gradient approximation (GGA) non-interacting kinetic energy density functional is presented along with illustrative applications. The innovation is adaptation of constraint-based construction to the essential properties of pseudo-densities from the pseudo-potentials that are essential in plane-wave-basis *ab initio* molecular dynamics. This contrasts with constraining to the qualitatively different Kato-cusp-condition densities. The single parameter in the new functional is calibrated by satisfying Pauli potential positivity constraints for pseudo-atom densities. In static lattice tests on simple metals and semiconductors, the new LKT functional outperforms the previous best constraint-based GGA functional, VT84F (Phys. Rev. B **88**, 161108(R) (2013)), is generally superior to a recently proposed meta-GGA, is reasonably competitive with parametrized two-point functionals, and is substantially faster.

*Introduction.* Hohenberg-Kohn density functional theory (DFT) [1, 2] has come to prominence mainly in Kohn-Sham (KS) orbital form [3]. However, driving *ab initio* molecular dynamics (AIMD) [4–7] with KS DFT exposes a computational cost-scaling burden. The KS computational cost scales no better than  $N_e^3$  with  $N_e$  the number of electrons or number of thermally occupied bands. Additionally there is reciprocal space sampling cost or equivalent costs from large real-space unit cells used with  $\Gamma$ -point sampling. In contrast, orbital-free DFT (OF-DFT) offers linear scaling with system size [8, 9] for use of AIMD on arbitrarily large systems.

The long-standing barrier to widespread use of OF-DFT has been the lack of reliable non-empirical approximate kinetic energy density functionals (KEDFs). In terms of the KS orbitals  $\varphi_j$ , the exact, positive definite KS kinetic energy (KE) density is

$$t_s[n] = t_s^{orb} \equiv \frac{1}{2} \sum_{j=1}^{N_e} |\nabla \varphi_j|^2, \quad (1)$$

in Hartree atomic units with  $n(\mathbf{r})$  the electron number density (and unit occupation for simplicity). Two types of approximate KEDFs have been explored, semi-local (one-point)

$$T_s[n] = \int d\mathbf{r} t_s[n(\mathbf{r}), \nabla n(\mathbf{r}), \dots] \quad (2)$$

and two-point with a non-local term

$$T_{NL}[n] = c_{TF} \int \int d\mathbf{r} d\mathbf{r}' n^\alpha(\mathbf{r}) K[n(\mathbf{r}), n(\mathbf{r}'), \mathbf{r}, \mathbf{r}'] n^\beta(\mathbf{r}') \quad (3)$$

with  $c_{TF} = \frac{3}{10}(3\pi^2)^{\frac{2}{3}}$ . For a dimensionless  $K$ ,  $\alpha + \beta = 8/3$ . Most approximate  $K$ s are parametrized; see Refs. [9–13] for details as well as brief discussion below. In this communication, we propose a novel non-empirical one-point KEDF and show that it is competitive with current two-point KEDFs, generally better than other one-point functionals, more transferable, and notably faster.

*Generalized Gradient Approximations.* The simplest one-point functionals are Thomas-Fermi [14–16]

$$T_{TF}[n] := \int d\mathbf{r} t_{TF}(\mathbf{r}), \quad t_{TF}(\mathbf{r}) := c_{TF} n^{\frac{5}{3}}(\mathbf{r}), \quad (4)$$

and von-Weizsäcker [17]

$$T_W[n] := \frac{1}{8} \int d\mathbf{r} \frac{|\nabla n(\mathbf{r})|^2}{n(\mathbf{r})} \equiv \int d\mathbf{r} t_W(\mathbf{r}). \quad (5)$$

Neither is satisfactory as a general KEDF. As with approximate exchange-correlation (XC) functionals [18], the gradient expansion of the weakly inhomogeneous electron gas KE leads to consideration of generalized gradient approximations (GGA) for  $T_s$ ,

$$T_s^{GGA}[n] = \int d\mathbf{r} t_{TF}(\mathbf{r}) F_t(s(\mathbf{r})). \quad (6)$$

Here  $F_t(s)$  is the GGA KE enhancement factor, a function of the dimensionless reduced density gradient  $s := \frac{|\nabla n|}{2nk_f} \equiv \frac{1}{2(3\pi^2)^{1/3}} \frac{|\nabla n|}{n^{4/3}}$  familiar from GGA X functionals. GGA KEDFs so constructed automatically satisfy  $T_s$  uniform scaling requirements [19]. In GGA form the von Weizsäcker KE becomes  $F_W(s) = \frac{5}{3}s^2$ .

From the Pauli term decomposition [8, 20, 21],

$$T_s[n] = T_W[n] + T_\theta[n], \quad (7)$$

three constraints follow[22],

$$T_\theta[n] \geq 0, \quad (8)$$

$$v_\theta(\mathbf{r}) \geq 0 \quad \forall \mathbf{r}, \quad (9)$$

$$v_\theta(\mathbf{r}) \geq \frac{t_\theta(\mathbf{r})}{n(\mathbf{r})} \quad \forall \mathbf{r}, \quad t_\theta := t_s^{orb} - t_W, \quad (10)$$

with the Pauli potential defined as  $v_\theta(\mathbf{r}) := \delta T_\theta[n] / \delta n(\mathbf{r})$  and the Pauli enhancement factor is  $F_\theta(s) = F_t(s) - F_W(s)$ .

To date, perhaps the best constraint-based GGA KEDF is VT84F (evaluated at T=0 K of course)[23].

It is successful in finite-T AIMD simulations [24] and is the only non-empirical GGA KEDF that yields reasonable binding in simple solids. It was constrained to satisfy Eqs. (8) and (9) for physical atom densities, i.e., those that obey the Kato cusp condition [25]. VT84F also was constrained to respect  $\lim_{s \rightarrow \infty} F_\theta(s)/F_W(s) = 0$ . This comes from the one-electron tail region of a many-electron atom [26] where  $t_\theta/t_W$  must vanish, hence  $t_s \rightarrow t_W$  [27].

In terms of the universal Hohenberg-Kohn-Levy density functional, such a physically motivated constraint is non-universal: the Kato cusp condition is specific to an external Coulomb potential. Such non-universality is rational for material and molecular property calculations. But the ubiquitous use of pseudo-potential plane-wave basis methods in AIMD simulations means that it is not the optimal non-universality for them. OF-DFT calculations in fact require a local pseudo-potential (LPP). The OF-DFT Euler equation then implies that  $v_\theta$  is closely related to the LPP  $v_{\text{ext}}^{\text{pseudo}}$  and that  $v_\theta$  is evaluated with the corresponding pseudo-density. Thus any constraint based on density characteristics should be specific to a particular type or class of pseudo-potential.

Ref. [28] explored some elementary consequences for constraint satisfaction (or violation) with non-Kato densities. Difficulties with simpler one-point KEDFs (linear combinations of  $T_{TF}$  and  $T_W$ ) used with orbital-free projector augmented-wave pseudo-densities also have been reported [29]. So far as we know, no approximate KEDF has been constructed by explicit satisfaction of the foregoing constraints, Eqs. (8)-(9), for a specified type of pseudo-densities. Nor has Eq. (10) been used.

*New GGA KEDF.* We resolve this pseudopotential AIMD deficiency by devising a GGA KEDF constrained to satisfy Eqs. (8) and (9) for pseudo-densities of a particular kind and show that in most spatial regions its  $v_\theta$  satisfies Eq. (10) as well. The new GGA KEDF enhancement factor is

$$F_t^{LKT}(s) = \frac{1}{\cosh(as)} + \frac{5}{3}s^2 \quad (11)$$

with parameter  $a > 0$ . Fig. 1 compares  $F_\theta^{LKT}$  with the VT84F and APBEK [30] enhancement factors. It satisfies the obvious homogeneous electron gas constraint  $\lim_{s \rightarrow 0} F_{t/\theta}(s) = 1$  and obeys  $0 \leq F_\theta^{LKT} \leq 1$  so as to satisfy the bound conjectured by Lieb [31, 32]

$$T_s \leq T_{TF} + T_W. \quad (12)$$

$F_t^{LKT}$  also satisfies [22, 26, 33]  $t_\theta([n]; \mathbf{r}) \geq 0 \forall \mathbf{r}$ , thus  $T_\theta^{LKT} \geq 0$ .

The sole parameter  $a = 1.3$  was determined as follows. A set of pseudo-densities was generated for the atoms H through Ne with a typical Hamann norm-conserving non-local pseudo-potential (NLPP) scheme [34] using default radii in the APE code [35] and the Perdew-Zunger




FIG. 1. Pauli enhancement factors for LKT ( $a = 1.3$ ) (red dot-dashed), VT84F (blue dashed), and APBEK (orange dotted).

(PZ) XC local density approximation (LDA) [36]. Then  $a$  was found such that all the post-scf Pauli potentials from those pseudo-densities satisfied  $v_\theta \geq 0 \forall r$ . Importantly, as long as an  $a$  value gave  $v_\theta \geq 0$  for the H atom, positivity also was met for all the heavier atoms. For Li  $a < 1.4$  is required, while for H,  $a \leq 1.3$  is needed to get a post-scf  $v_\theta \geq 0$ . For He, the  $a$  value does not seem to matter within the range tested. While the  $a$  value is non-universal, we expect reasonable transferability to those other pseudo-potential types for which the pseudo-densities are similar, specifically those with nearly flat pseudo-densities near the nucleus. The expectation is confirmed by post-scf and scf calculations for atoms.

Though reference atom set, H-Ne, encompasses 1-8 pseudo-electrons, equally good performance for other elements is not assured. Post-scf determination of  $a$  also is distinct from self-consistent calculation, which might vitiate the supposedly constrained behavior. Atomic tests are the first line of investigating these issues. For a given pseudo-potential and XC approximation, self-consistent solution of the KS equation provides the exact KS  $t_\theta$  and the ingredients to construct the exact KS Pauli  $v_\theta$  (see Eq. (35) in Ref. [20]). Those are the standards against which to judge  $t_\theta$  and  $v_\theta$  from an approximate KEDF. In anticipation of the OF-DFT calculations on periodically bounded systems reported below, we focused upon the bulk-derived LPP (BLPS) [37, 38] for two atoms, Al and Li. Here we discuss Al because it was not in the  $a$  calibration. Li discussion is in the Supplemental Material [39]. (The Li pseudo-atom is challenging because it is a one-orbital system ( $2s^1$ ) for which  $T_\theta$  should vanish.) Again the XC functional is PZ.

Fig. 2 displays the exact  $t_\theta/n$  and  $v_\theta$  for the BLPS Al pseudo-atom in the  $3s^2 3p^1$  configuration and the post-scf results with that pseudo-density for both VT84F and LKT. Note several features. Though VT84F was constructed to satisfy  $v_\theta^{VT84F} \geq 0$  near a nucleus for Kato-cusped densities, it also satisfies that constraint arbitrarily close to the nucleus for the cusp-less pseudo-density.




FIG. 2. Upper left: Al BLPS as function of radial position (inset: KS pseudo density); Upper right: exact KS  $v_\theta$  and  $t_\theta/n$ . Lower left: post-scf  $v_\theta$  (solid) and  $t_\theta/n$  for VT84F (dashed). Lower right: Same for LKT.

However,  $v_\theta^{VT84F}$  becomes negative near  $r = 0.1$  bohr, a clear example of the crucial non-universality. LKT does not have that problem. Second,  $v_\theta^{LKT}$  is much smoother than  $v_\theta^{VT84F}$ , though not as smooth as  $v_\theta^{KS}$ . Third, except for a small region around  $r = 1.8$  bohr,  $v_\theta^{LKT}$  respects the Pauli potential inequality, Eq. (10), whereas  $v_\theta^{VT84F}$  violates it in four regions that span much of the significant density magnitude.

Note also that, unlike some other GGA KEDFs, e.g. E00[40], PBE2 [21], and APBEK,  $v_\theta^{LKT}(r)$  decays correctly to zero asymptotically for an atom. This may be useful in the AIMD simulation of low-density regions of matter. Though  $v_\theta^{VT84F}$  decays similarly, its rapid oscillations in the dominant density region might slow scf convergence rates as well as cause other difficulties.

Self-consistent OF calculations for the BLPS Al pseudo-atom show that  $v_\theta^{LKT}$  stays positive, though it exhibits oscillations quite similar to those seen in the post-scf case; see Fig. 3. The inequality of Eq. (10) is violated only around  $r = 1.8$  bohr as in the post-scf case. However, the LKT Pauli energy per particle is far from the KS value.

*Performance on Solids.* Validation of the new functional for AIMD requires accuracy tests on extended systems. We therefore did KS-DFT and OF-DFT calculations on simple metals and semiconductors. Conventional KS calculations were done with ABINIT [41] and the OF-DFT calculations used PROFESS[42] and/or PROFESS@QUANTUM-ESPRESSO [43]. Again the PZ LDA XC functional and BLPS were used. For comparison we included the Wang-Govind-Carter (WGC) [10], Huang-Carter (HC) [11], and Constantin et al. KGAP [13] two-point KEDFs and the one-point Constantin et al. SOF-CFD [44] meta-GGA (Laplacian-dependent)




FIG. 3. Top: Al KS (solid, red) and LKT(dashed, blue) pseudo-densities as function of radial position. Bottom: KS vs. LKT  $v_\theta$  (solid red vs. dash-dotted blue, upper pair) and similarly  $t_\theta/n$  (dashed red v. dotted blue; lower pair).

KEDF. Technical details and parameter values are in the Supplemental Material [39].

Note that WGC was parametrized for main-group metals and yields poor binding curves for semiconductors, while HC was parametrized for semiconductors. KGAP is parametrized to experimental direct band gaps. Results from the one-point functionals E00, APBEK, and PBE2 are omitted because of unrealistic binding curves for the former two and instability problems for the latter one. KGAP comparisons are from Tables I and II of Ref. [13]. SOF-CFD values are from Table I of Ref. [44]. Equilibrium volumes, energies, and bulk moduli for other functionals were generated by varying  $\pm 5\%$  around the equilibrium volume to obtain 11 energy-volume points, which then were fitted to the Birch-Murnaghan equation of state [45].

The metals were Li, Mg, and Al in the simple cubic, body-centered cubic, face-centered cubic, and hexagonal close-packed structures. Nine III-V semiconductors in zinc-blende structures were treated: AlP, AlAs, AlSb, GaP, GaAs, GaSb, InP, InAs, and InSb.

TABLE I. KEDF performance on solid metals and semiconductors: MARE of equilibrium volumes  $V_0$ , energies  $E_0$ , and bulk moduli  $B_0$ , as percentages. See text for notation.

KEDF	Metals			Semiconductors		
	$V_0$	$E_0$	$B_0$	$V_0$	$E_0$	$B_0$
WGC	0.7	0.0	2.7	-	-	-
HC	5.5	0.6	12.3	1.5	0.5	4.9
KGAP <sup>a</sup>	4.0	-	5.1	3.0	-	16.2
VT84F	6.0	0.1	11.6	10.5	3.6	56.4
SOF-CFD <sup>a</sup>	5.2	0.6	8.5	3.4	0.9	10.0
LKT	4.0	0.2	7.7	2.1	2.8	4.3

<sup>a</sup> Note: only metals with cubic symmetry were included and PBE XC was used.

With KS quantities as references, Table I shows the mean absolute relative error (MARE) percentages for equilibrium volume  $V_0$ , energy  $E_0$  per atom (for metals) or per cell (for semiconductors), and bulk moduli  $B_0$  from WGC, HC, KGAP, VT84F, SOF-CFD, and LKT. These are calculated as  $|(Q_{\text{OF}} - Q_{\text{KS}})/Q_{\text{KS}}| \times 100/N_{\text{systems}}$ , where  $Q$  is  $V_0$ ,  $E_0$ , or  $B_0$ . (More detailed tabulations are in the Supplemental Material [39].) For  $V_0$  and  $B_0$ , LKT is a significant improvement over VT84F. The  $V_0$  and  $B_0$  MAREs are reduced by 33% in metals. The reduction is more dramatic in the semi-conductors, a factor of 5 for  $V_0$  and 13 for  $B_0$ . The semiconductor  $E_0$  MARE is reduced by 22% but worsened slightly from 0.1% to 0.2% for the metals. Except for performance on semiconductor  $E_0$ , it also is clear that the LKT GGA is superior to the more-complicated non-empirical SOF-CFD meta-GGA KEDF.

Regarding the two-point functionals, WGC outperforms all the other functionals on the metals but is inapplicable on semiconductors, recall above [10]. Conversely, HC with averaged parameters exhibits balanced error, with all three MAREs within 5% (except  $B_0$  for metals). KGAP does well on volumes in both classes but not  $B_0$ . Remarkably LKT exhibits performance competitive with both HC and KGAP in prediction of equilibrium volumes for both material classes. Moreover, LKT outperforms HC for  $B_0$  and is much more balanced than KGAP for  $B_0$ . (Comparison with the recent MGP two-point functional is of no avail, since its parametrization is tuned to match KS results for each system [12].)

For the case of AIP, we found that LKT converges for relatively smaller energy cutoff than needed with VT84F and HC. Typically LKT also requires fewer self-consistent iterations for solution to a given tolerance than are needed by either HC or VT84F and each LKT iteration is typically about one-fifth the time of an HC iteration. Thus the one-point LKT is more useful as a broadly applicable functional than the highly parametrized two-point HC KEDF or the experimentally parametrized two-point KGAP KEDF yet is simpler, faster, and mostly better than the SOF-CFD one-point KEDF. LKT seems therefore to be currently the most promising candidate for general AIMD OF-DFT use or with small-box algorithms [46]. Though it remains to be tested, we anticipate the finite-T generalization [47] of LKT will be of value for warm dense matter simulations.

As to limitations, LKT does not yield a good value of  $V_0$  for bcc Li with a 3-electron LPP. So far as we know, all GGA KEDFs developed so far share this limitation. The extent of transferability to another distinct class of pseudo-potential, along with the post-scf determination of  $a$ , remains to be examined.

KL and SBT were supported by U.S. Dept. of Energy grant de-sc 0002139. VVK acknowledges support by the Dept. of Energy National Nuclear Security Administration under Award Number de-na0001944.

- 
- \* kluo@ufl.edu  
† vkarasev@lle.rochester.edu  
‡ trickey@qtp.ufl.edu
- [1] P. Hohenberg and W. Kohn, Phys. Rev. **136**, B864 (1964).
  - [2] M. Levy, Proc. Natl. Acad. Sci. USA **76**, 6062 (1979).
  - [3] W. Kohn and L.J. Sham, Phys. Rev. **140**, A1133 (1965).
  - [4] R.N. Barnett and U. Landman, Phys. Rev. B **48**, 2081 (1993).
  - [5] D. Marx and J. Hutter in *Modern Methods and Algorithms of Quantum Chemistry*, J. Grotendorst ed., John von Neumann Institute for Computing, (Jülich, NIC Series, Vol. 1, 2000) 301 and refs. therein.
  - [6] J.S. Tse, Annu. Rev. Phys. Chem. **53**, 249 (2002).
  - [7] *Ab Initio Molecular Dynamics: Basic Theory and Advanced Methods*, D. Marx and J. Hutter, (Cambridge University Press, Cambridge, 2009) and refs. therein.
  - [8] V.V. Karasiev, D. Chakraborty, and S.B. Trickey, in *Many-electron Approaches in Physics, Chemistry, and Mathematics: A Multidisciplinary View*, L. Delle Site and V. Bach eds. (Springer, Heidelberg, 2014) 113 and refs. therein.
  - [9] W.C. Witt, B.G. del Rio, J.M. Dieterich, and E.A. Carter, J. Mat. Res. **33**, 777 (2018) and refs. therein.
  - [10] Y.A. Wang, N. Govind and E.A. Carter, Phys. Rev. B **60** 16350 (1999); erratum Phys. Rev. B **64**(E), 089903 (2001).
  - [11] C. Huang and E.A. Carter, Phys. Rev. B **81**, 045206 (2010).
  - [12] W. Mi, A. Genova, and M. Pavanello, J. Chem. Phys. **148**, 184107 (2018).
  - [13] L.A. Constantin, E. Fabiano, and F. Della Sala, Phys. Rev. B **97**, 205137 (2018).
  - [14] L.H. Thomas, Proc. Camb. Phil. Soc. **23**, 542 (1927).
  - [15] E. Fermi, Rend. Accad. Naz. Lincei **6**, 602 (1927).
  - [16] E. Fermi, Z. Phys. **48**, 73 (1928).
  - [17] C.F. von Weizsäcker, Z. Phys. **96**, 431 (1935).
  - [18] J.P. Perdew, Phys. Lett. A **165**, 79 (1992).
  - [19] M. Levy and J.P. Perdew, Phys. Rev. A **32**, 2010 (1985).
  - [20] V.V. Karasiev, R.S. Jones, S.B. Trickey, and F.E. Harris, Phys. Rev. B **80**, 245120 (2009).
  - [21] V.V. Karasiev, R.S. Jones, S.B. Trickey, and F.E. Harris, in *New Developments in Quantum Chemistry*, J.L. Paz and A.J. Hernández, eds. (Transworld Research Network, Kerala, India, 2009) p. 25.
  - [22] M. Levy, and H. Ou-Yang, Phys. Rev. A **38**, 625 (1988).
  - [23] V.V. Karasiev, D. Chakraborty, O.A. Shukruto, and S.B. Trickey, Phys. Rev. B **88**, 161108(R) (2013).
  - [24] V.V. Karasiev, L. Calderín, and S.B. Trickey, Phys. Rev. E **93**, 063207 (2016).
  - [25] T. Kato, Commun. Pure Appl. Math. **10**, 151 (1957).
  - [26] M. Levy, J.P. Perdew, and V. Sahni, Phys. Rev. A **30**, 2745 (1984).
  - [27] R.M. Dreizler and E.K.U. Gross, *Density Functional Theory* (Springer-Verlag, Berlin, 1990).
  - [28] V.V. Karasiev and S.B. Trickey, Adv. Quantum Chem. **71**, 221 (2015).
  - [29] J. Lehtomäki, I. Makkonen, M.A. Caro, and O. Lopez-Acevedo, J. Chem. Phys. **141**, 234102 (2014).
  - [30] L.A. Constantin, E. Fabiano, S. Laricchia, and F. Della Sala, Phys. Rev. Lett. **106**, 186406 (2011).

- [31] E.H. Lieb, Lecture Notes in Physics **116**, 91 (1980).
- [32] J.L. Gázquez and J. Robles, J. Chem. Phys. **76**, 1467 (1982).
- [33] C. Herring, Phys. Rev. A **34**, 2614 (1986).
- [34] D.R. Hamann, M. Schlüter, and C. Chiang, Phys. Rev. Lett. **43**, 1494 (1979) .
- [35] M.J.T. Oliveira and F. Nogueira, Comput. Phys. Commun. **178**, 524 (2008).
- [36] J.P. Perdew, and A. Zunger, Phys. Rev. B **23**, 5048 (1981).
- [37] B. Zhou, Y.A. Wang, and E.A. Carter, Phys. Rev. B **69**, 125109 (2004).
- [38] C. Huang and E. A. Carter, Phys. Chem. Chem. Phys., **10**, 7109 (2008).
- [39] Supplemental Material link here.
- [40] M. Ernzerhof, J. Mol. Struct.: THEOCHEM **501-502**, 59 (2000).
- [41] X. Gonze *et al.*, Computer Phys. Commun. **180**, 2582 (2009).
- [42] L. Hung, C. Huang, I. Shin, G.S. Ho, V.L. Lignères, and E.A. Carter, Comput. Phys. Commun. **181**, 2208 (2010).
- [43] V.V. Karasiev, T. Sjöström, and S.B. Trickey, Comput. Phys. Commun. **185**, 3240 (2014).
- [44] L.A. Constantin, E. Fabiano, and F. Della Sala, arXiv:1802.02889 [cond-mat.mtrl-sci]. This functional was denoted “SOF” (semi-local orbital-free) but that properly is the name of a class of functionals. For specificity we append the authors’ initials, “SOF-CFD”.
- [45] F. Birch, Phys. Rev. **71**, 809 (1947).
- [46] M. Chen, X. Jiang, H. Zhuang, L. Wang, and E.A. Carter, J. Comp. Theor. Comp., **12**, 2950 (2016).
- [47] V.V. Karasiev, T. Sjöström, and S.B. Trickey, Phys. Rev. B **86**, 115101 (2012).