

Comment on scalar glueball mass

Amir H. Fariborz ^a and Renata Jora ^b

^a *Department of Mathematics/Physics, SUNY Polytechnic Institute, Utica, NY 13502, USA and*

^b *National Institute of Physics and Nuclear Engineering PO Box MG-6, Bucharest-Magurele, Romania*

We study the interaction of scalar glueball with quark-antiquark and four-quark spinless meson fields within the framework of a generalized linear sigma model in which the trace anomaly of QCD is exactly realized. To determine the pure scalar glueball mass (m_h), we consider the decoupling limit in which the scalar glueball decouples from the meson fields. We find the exact relationship $m_h = 2h_0$ where h_0 is the vacuum expectation value of the scalar glueball field independent of the properties of the framework used.

The glueballs are the inherent byproducts of self-interacting gluons in QCD [1]. Since they can mix with meson fields of the same quantum numbers, determination of their properties becomes naturally challenging. The scalar glueballs are particularly difficult to study due to the fact that they have the quantum numbers of QCD vacuum and can induce spontaneous chiral symmetry breaking by developing vacuum expectation values. There have been an extensive investigation of glueballs from different approaches, including lattice QCD [2], QCD sum-rules [3, 4] and chiral models [5].

We probe scalar glueballs within the framework of the generalized linear sigma model of [6, 7] in which the properties of scalar and pseudoscalar mesons below and above 1 GeV and their underlying quark-antiquark and four-quark mixings have been extensively studied. In this work we are specifically interested in properties of the lightest pure scalar glueball, hence, we consider a limit of the generalized linear sigma model in which the glueballs decouple from the quark-antiquark and four-quark states. The model is constructed in terms of 3×3 matrix chiral nonet fields:

$$M = S + i\phi, \quad M' = S' + i\phi', \quad (1)$$

where M and M' transform in the same way under chiral SU(3) transformations

$$\begin{aligned} M &\rightarrow U_L M U_R^\dagger, \\ M' &\rightarrow U_L M' U_R^\dagger, \end{aligned} \quad (2)$$

but transform differently under U(1)_A transformation properties

$$\begin{aligned} M &\rightarrow e^{2i\nu} M, \\ M' &\rightarrow e^{-4i\nu} M'. \end{aligned} \quad (3)$$

M contains the ‘‘bare’’ (unmixed) quark-antiquark scalar nonet S and pseudoscalar nonet ϕ , while M' contains ‘‘bare’’ two quarks and two antiquarks scalar nonet S' and pseudoscalar nonet ϕ' . The model distinguishes M from M' through the U(1)_A transformation.

The Lagrangian density has the following general structure in terms of chiral nonets M and M' as well as scalar glueball h and pseudoscalar glueball g :

$$\begin{aligned} \mathcal{L} &= -\frac{1}{2}\text{Tr}(\partial^\mu M \partial_\mu M^\dagger) - \frac{1}{2}\text{Tr}(\partial^\mu M' \partial_\mu M'^\dagger) - V, \\ -V &= f + f_A + f_S + f_{\text{SB}}. \end{aligned} \quad (4)$$

Here f is a general function of fields M , M' , h and g and is chiral, U(1)_A and scale invariant; f_A and f_S exactly mock up the axial and trace anomalies and f_{SB} introduces explicit breaking of the chiral symmetry due to quark masses. The leading choice of terms corresponds to eight or fewer underlying quark plus antiquark lines at each effective vertex [6, 7] and are as follows:

$$\begin{aligned} f(M, M', g, h) &= - (u_1 h^2 \text{Tr}[MM^\dagger] + u_2 \text{Tr}[MM^\dagger MM^\dagger] \\ &\quad + u_3 h^2 \text{Tr}[M'M'^\dagger] \\ &\quad + u_4 h (\epsilon_{abc} \epsilon^{def} M_d^a M_e^b M_f^c + h.c.) \\ &\quad + u_5 h^4 + u_6 h^2 g^2 + \dots), \end{aligned} \quad (5)$$

$$f_A = i \frac{G}{12} \left[\gamma_1 \ln \left(\frac{\det M}{\det M^\dagger} \right) + \gamma_2 \ln \left(\frac{\text{Tr}(MM^\dagger)}{\text{Tr}(M'M'^\dagger)} \right) \right], \quad (6)$$

where $G = h^3 g$ and γ_1 and γ_2 are arbitrary parameters that must satisfy the constraint $\gamma_1 + \gamma_2 = 1$.

$$\begin{aligned} f_S &= -H \left\{ \lambda_1 \ln \left(\frac{H}{\Lambda^4} \right) \right. \\ &\quad + \lambda_2 \left[\ln \left(\frac{\det M}{\Lambda^3} \right) + \ln \left(\frac{\det M^\dagger}{\Lambda^3} \right) \right] \\ &\quad \left. + \lambda_3 \left[\ln \left(\frac{\text{Tr}MM^\dagger}{\Lambda^2} \right) + \ln \left(\frac{\text{Tr}M'M'^\dagger}{\Lambda^2} \right) \right] \right\}. \end{aligned} \quad (7)$$

where Λ is the characteristic scale, $H = h^4$ and λ_1 , λ_2 and λ_3 satisfying the constraint $4\lambda_1 + 6\lambda_2 + 4\lambda_3 = 1$. The two terms f_A and f_S ensure that the effective Lagrangian exactly satisfies the U(1)_A and the trace anomalies according to:

$$\begin{aligned} \partial^\mu J_\mu^5 &= \frac{g^2}{16\pi^2} N_F \tilde{F} F = G, \\ \theta_\mu^\mu &= \partial^\mu D_\mu = -\frac{\beta(g^2)}{2g} F F = H, \end{aligned} \quad (8)$$

where F is the SU(3)_C field tensor, \tilde{F} is its dual, N_F is the number of flavors, $\beta(g^2)$ is the beta function for the coupling constant, J_μ^5 is the axial current and D_μ is the dilatation current. The symmetry breaking term takes the simple form:

$$f_{\text{SB}} = 2\text{Tr}[AS], \quad (9)$$

where $A = \text{diag}(A_1, A_2, A_3)$ is a matrix proportional to the three light quark masses.

The isosinglet scalar states $f_1 \cdots f_5$ are a linear combination of the following five components: two- and four-quark building blocks as well as a glueball component:

$$\begin{aligned}
(i) \quad f_a &= \frac{S_1^1 + S_2^2}{\sqrt{2}} && \propto & n\bar{n}, \\
(ii) \quad f_b &= S_3^3 && \propto & s\bar{s}, \\
(iii) \quad f_c &= \frac{S_1^1 + S_2^2}{\sqrt{2}} && \propto & ns\bar{n}s, \\
(iv) \quad f_d &= S_3^3 && \propto & nn\bar{n}\bar{n}, \\
(v) \quad h & && &
\end{aligned} \tag{10}$$

where the non-strange (n) and strange (s) quark content for each basis state has been listed at the end of each line above. The VEV of the scalar fields

$$\begin{aligned}
\alpha_a &= \langle S_a^a \rangle, \\
\beta_a &= \langle S'^a_a \rangle, \\
h_0 &= \langle h \rangle,
\end{aligned} \tag{11}$$

with $a = 1 \cdots 3$ spontaneously break chiral symmetry.

The minimum equations that determine the vacuum of the model are obtained from equations:

$$\begin{aligned}
\left\langle \frac{\partial V}{\partial S_1^1} \right\rangle_0 &= 0, \\
\left\langle \frac{\partial V}{\partial S_3^3} \right\rangle_0 &= 0, \\
\left\langle \frac{\partial V}{\partial S_1^1} \right\rangle_0 &= 0, \\
\left\langle \frac{\partial V}{\partial S_3^3} \right\rangle_0 &= 0, \\
\left\langle \frac{\partial V}{\partial h} \right\rangle_0 &= 0,
\end{aligned} \tag{12}$$

where brackets with subscript zero represent evaluation of each derivative at VEV values of (11).

In order to extract the properties of the scalar glueball, we consider the decoupling limit in which the scalar

glueball decouples from quark mesons:

$$\begin{aligned}
\left\langle \frac{\partial^2 V}{\partial S_1^1 \partial h} \right\rangle_0 &= 4\sqrt{2}u_4[\beta_1\alpha_3 + \alpha_1\beta_3] + 4\sqrt{2}u_1h_0\alpha_1 \\
&\quad + \frac{8\sqrt{2}h_0^3\lambda_2}{\alpha_1} + \frac{8\sqrt{2}h_0^3\lambda_3\beta_1}{2\alpha_1\beta_1 + \alpha_3\beta_3} = 0, \\
\left\langle \frac{\partial^2 V}{\partial S_3^3 \partial h} \right\rangle_0 &= 8u_4\alpha_1\beta_1 + 4u_1h_0\alpha_3 + \frac{8h_0^3\lambda_2}{\alpha_3} \\
&\quad + \frac{8h_0^3\lambda_3\beta_3}{2\alpha_1\beta_1 + \alpha_3\beta_3} = 0, \\
\left\langle \frac{\partial^2 V}{\partial S_1^1 \partial h} \right\rangle_0 &= 4\sqrt{2}\alpha_1\alpha_3u_4 + 4\sqrt{2}u_3h_0\beta_1 \\
&\quad + \frac{8\sqrt{2}h_0^3\lambda_3\alpha_1}{2\alpha_1\beta_1 + \alpha_3\beta_3} = 0, \\
\left\langle \frac{\partial^2 V}{\partial S_3^3 \partial h} \right\rangle_0 &= 4\alpha_1^2u_4 + 4u_3h_0\beta_3 \\
&\quad + \frac{8h_0^3\lambda_3\alpha_3}{2\alpha_1\beta_1 + \alpha_3\beta_3} = 0.
\end{aligned} \tag{13}$$

The first two minimum equations in (12) can be used to solve for A_1 and A_3 in terms of other parameters. The third and fourth equations in (12) give:

$$\begin{aligned}
4\alpha_1^2\alpha_3\beta_1u_4 + 2\alpha_3^2\beta_3\alpha_1u_4 + 2\beta_1^2\alpha_1h_0u_3 \\
+ h_0^3\lambda_3\alpha_1 + \alpha_3\beta_1\beta_3h_0u_3 &= 0, \\
4\alpha_1^3\beta_1u_4 + 2\alpha_1^2\alpha_3\beta_3u_4 + 2\alpha_1\beta_1\beta_3h_0u_3 \\
+ \alpha_3\beta_3^2h_0u_3 + \alpha_3h_0^3\lambda_3 &= 0.
\end{aligned} \tag{14}$$

Solving this system results in relationships:

$$\begin{aligned}
\alpha_3 &= \alpha_1 = \alpha, \\
\beta_3 &= \beta_1 = \beta,
\end{aligned} \tag{15}$$

and consequently $A_1 = A_3 = A$, and shows that in the decoupling limit the chiral symmetry breaks into its $SU(3)_V$ subgroup. Furthermore, taking the decoupling equations (13) into account, we find additional relationships between the model parameters:

$$\begin{aligned}
u_3 &= -\frac{3\alpha^2u_4}{\beta h_0}, \\
\lambda_2 &= -\frac{\alpha^2(h_0u_1 + 4\beta u_4)}{2h_0^3}, \\
\lambda_3 &= \frac{3\alpha^2\beta u_4}{h_0^3}.
\end{aligned} \tag{16}$$

The scalar glueball mass is determined from

$$m_h^2 = \left\langle \frac{\partial^2 V}{\partial^2 h} \right\rangle_0, \tag{17}$$

where, in the decoupling limit, becomes

$$\begin{aligned}
m_h^2 &= 6u_1\alpha^2 + 6u_3\beta^2 + 12u_5h_0^2 + 28h_0^2\lambda_1 \\
&\quad + 12h_0^2 \left[\lambda_1 \ln \left(\frac{h_0^4}{\Lambda^4} \right) + 2\lambda_2 \ln \left(\frac{\alpha^3}{\Lambda^3} \right) \right. \\
&\quad \left. + 2\lambda_3 \ln \left(\frac{3\alpha\beta}{\Lambda^2} \right) \right].
\end{aligned} \tag{18}$$

The fifth equation in (12) can be used to solve for u_5 in terms of other parameters, which upon substituting this solution back into (18), the glueball mass squared simplifies to:

$$m_h^2 = \frac{1}{h_0} (-36 \alpha^2 \beta u_4 - 12 u_3 h_0 \beta^2 - 12 u_1 h_0 \alpha^2 + 16 h_0^3 \lambda_1). \quad (19)$$

The first two terms cancel out exactly using the first relation in (16). Using the other two relations in (16), simplifies the third and the fourth terms in (19):

$$\begin{aligned} m_h^2 &= -12 u_1 \alpha^2 + 16 h_0^2 \lambda_1 \\ &= -12 \alpha^2 \left(\frac{-2 h_0^2 \lambda_2}{\alpha^2} - \frac{4 \beta u_4}{h_0} \right) + 16 h_0^2 \lambda_1 \\ &= 24 h_0^2 \lambda_2 + 48 \frac{\beta \alpha^2 u_4}{h_0} + 16 h_0^2 \lambda_1 \\ &= 4 h_0^2 (4 \lambda_1 + 6 \lambda_2 + 4 \lambda_3). \end{aligned} \quad (20)$$

The parenthesis is equal to 1 [see (7)] leading to a simple relationship for the pure glueball mass.

$$m_h = 2 h_0. \quad (21)$$

Note that this result which was derived for a specific potential is model independent. This can be shown by directly examining the trace anomaly for a generic potential V which contains the fields M , M' , h and g . Requiring that the scale transformation of V exactly gives the second equation of (8), yields

$$\left[M_{ij} \frac{\partial V}{\partial M_{ij}} + M'_{ij} \frac{\partial V}{\partial M'_{ij}} + h.c. \right] + \frac{\partial V}{\partial h} h + \frac{\partial V}{\partial g} g - 4V = h^4. \quad (22)$$

Differentiating with respect to the field h and taking the vacuum expectation values results in

$$\begin{aligned} &\left[\langle M_{ij} \rangle_0 \left\langle \frac{\partial^2 V}{\partial M_{ij} \partial h} \right\rangle_0 + \langle M'_{ij} \rangle_0 \left\langle \frac{\partial^2 V}{\partial M'_{ij} \partial h} \right\rangle_0 + h.c. \right] \\ &+ \left\langle \frac{\partial^2 V}{\partial h^2} \right\rangle_0 h_0 + \left\langle \frac{\partial V}{\partial h} \right\rangle_0 + \left\langle \frac{\partial^2 V}{\partial g \partial h} \right\rangle_0 \langle g \rangle_0 - \left\langle 4 \frac{\partial V}{\partial h} \right\rangle_0 \\ &= 4 h_0^3. \end{aligned} \quad (23)$$

In the decoupling limit of (16), the square bracket vanishes. The first term after the square bracket is the scalar glueball mass squared times h_0 , terms including $\langle \partial V / \partial h \rangle_0$ vanish because of minimum condition [the last Eq. in (12)] and $\langle g \rangle_0$ vanishes because of parity, resulting in the same conclusion as in (21).

Although relationship (21) can be derived model independently, computation of h_0 requires specific modeling of the potential. In first approximation,

$$h_0^4 \approx h_x^4 = \langle h^4 \rangle = \langle H \rangle = \left\langle -\frac{\beta(g)}{2g} G^2 \right\rangle = \frac{9}{8\pi} \langle \alpha_s G^2 \rangle, \quad (24)$$

where the gluon condensate can be imported from QCD sum-rules or lattice QCD. However, the relationship between h_0 and h_x should be established more carefully. The partition function for our model reads:

$$Z = \int dh dg dM dM' \exp[i \int d^4 x \mathcal{L}(h, g, M, M')]. \quad (25)$$

Making a change of variables from h and g to $H = h^4$ and $G = h^3 g$, with the Jacobian of the transformation

$$J(x-y) = \det \left[H(x)^{-3/2} \delta(x-y) \right], \quad (26)$$

the minimum equation for H then becomes:

$$\frac{\partial iS}{\partial H(z)} - \frac{3}{2} \frac{1}{H(z)} \delta(0) = 0. \quad (27)$$

Here S is the action and the second term comes from the differentiation of the determinant in Eq. (26). Eq. (27) can be rewritten as:

$$\int d^4 y \frac{\partial iS}{\partial h(y)} \frac{\partial h(y)}{\partial H(z)} - \frac{3}{2} \frac{1}{H(z)} \delta(0) = 0, \quad (28)$$

which becomes:

$$-i \frac{\partial V(z)}{\partial h(z)} \frac{1}{4H(z)^{3/4}} - \frac{3}{2} \frac{1}{H(z)} \delta(0) = 0. \quad (29)$$

Note that this time the derivative is not taken in the functional sense. The δ function in Eq. (29) needs regularization. Since we are interested in a small range of values around the vacuum condensate we can write:

$$\begin{aligned} \delta(0) &= \lim_{x \rightarrow y} \delta(x-y) \\ &= \int \frac{d^4 k}{(2\pi)^4} \exp\left(\frac{\partial^2}{m_c^4}\right) \exp[-ik(x-y)] = i \frac{m_c^4}{16\pi^2}, \end{aligned} \quad (30)$$

where the integral is computed in the Euclidean space and m_c is an appropriate cut-off scale for a linear sigma model with three flavors. Then the vacuum equation (29) becomes:

$$\frac{\partial V}{\partial h} \Big|_{h=h_x} \frac{1}{4h_x^3} + \frac{6}{64\pi^2} \frac{m_c^4}{h_x^4} = 0. \quad (31)$$

We multiply the above equation by 4 and subtract from it the vacuum equation for h_0 divided by h_0^3 to get:

$$K(h_0) = \frac{\partial V}{\partial h} \Big|_{h=h_x} \frac{1}{h_x^3} + \frac{6}{16\pi^2} \frac{m_c^4}{h_x^4} - \frac{\partial V}{\partial h} \Big|_{h=h_0} \frac{1}{h_0^3} = 0. \quad (32)$$

In the decoupling limit that we considered here, this equation can be rewritten in terms of the model parameters

$$\begin{aligned} K(h_0) &= 4\lambda_1 \ln \left(\frac{h_x^4}{h_0^4} \right) + (6\alpha^2 u_1 + 6\beta^2 u_3) \left(\frac{1}{h_x^2} - \frac{1}{h_0^2} \right) \\ &+ 12u_4 \alpha^2 \beta \left(\frac{1}{h_x^3} - \frac{1}{h_0^3} \right) + \frac{6}{16\pi^2} \frac{m_c^4}{h_x^4} = 0. \end{aligned} \quad (33)$$

In this equation, h_x is an input, computed from Eq. (24) with the value of the gluon condensate taken from QCD sum-rules or lattice QCD. (There is a large discrepancy between the values of gluon condensate given by lattice QCD and QCD sum-rules.) The cut off m_c is adjusted such that $K = 0$ occurs at a stable point (the local minimum of K at which there is no sensitivity to determination of h_0). For example, from QCD sum-rule analysis of [4]:

$$\langle \alpha_s G^2 \rangle = (0.070 \pm 0.013) \text{ GeV}^4, \quad (34)$$

which gives $h_x^{\min} = 0.057 \text{ GeV}$ and $h_x^{\max} = 0.083 \text{ GeV}$, function K vanishes at a local minimum if $m_c = 1.04 \text{ GeV}$ and 1.14 GeV , respectively, as shown in Fig. 1. At these local minima h_0 is 0.887 GeV and 0.974 GeV , respectively. These values of h_0 , when entered into Eq. (21), result in a pure scalar glueball mass of 1.77 GeV and 1.95 GeV , overlapping with some of the estimates in the literature [2, 5].

In summary, this work presented a model independent relationship between the mass and the condensate of a pure scalar glueball. This relationship can be of a key importance in low-energy QCD analyses by bridging frameworks involving quarks and gluons to hadronic models of low-energy QCD.




FIG. 1. Function $K(h_0)$ [Eq. (33)] versus h_0 with the two values of h_x , its minimum (dotted line) and its maximum (solid line) extracted from QCD sum-rule analysis of [4]. For each case, the cut off values are adjusted such that $K = 0$ occurs at its minimum which is insensitive to extraction of h_0 . At the two minima, the values of h_0 are 0.887 and 0.974 GeV corresponding to the minimum and maximum values of h_x , respectively.

ACKNOWLEDGMENTS

A.H.F. gratefully acknowledges the support of College of Arts and Sciences of SUNY Poly in Spring 2018.

-
- [1] C. Patrignani et al. (Particle Data Group), *Chin. Phys. C*, **40**, 100001 (2016); E. Klempt and A. Zaitsev, *Phys. Rept.* **454**, 1 (2007).
- [2] C.J. Morningstar and M.J. Peardon, *Phys. Rev. D* **60**, 034509 (1999); W.J. Lee and D. Weingarten, *Phys. Rev. D* **61**, 014015 (2000); G. S Bali et al., *Phys. Lett. B* **309**, 379 (1993); C. McNeile *et al.* [UKQCD Collaboration], *Phys. Rev. D* **63**, 114503 (2001); L.C. Gui *et al.* [CLQCD Collaboration], *Phys. Rev. Lett.* **110**, no. 2, 021601 (2013).
- [3] M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *Nucl. Phys. B* **147**, 385, 448 (1979); S. Narison and G. Veneziano, *Int. J. Mod. Phys. A* **4**, 2751 (1981); S. Narison, *Z. Phys. C* **26**, 209 (1984).
- [4] S. Narison, *Phys. Lett. B* **706**, 412-422 (2012).
- [5] M. Albaladejo and J.A. Oller, *Phys. Rev. Lett.* **101**, 252002 (2008); F. Giacosa, T. Gutsche, V.E. Lyubovitskij and A. Faessler, *Phys. Rev. D* **72**, 094006 (2005); M. Chanowitz, *Phys. Rev. Lett.* **95**, 172001 (2005); A.H. Fariborz, *Int. J. Mod. Phys. A* **19**, 2095 (2004); A.H. Fariborz, *Phys. Rev. D* **74**, 054030 (2006); C. Amsler and F.E. Close, *Phys. Rev. D* **53**, 295 (1996).
- [6] A.H. Fariborz, R. Jora and J. Schechter, *Phys. Rev. D* **72**, 034001 (2005); *Phys. Rev. D* **76**, 014011 (2007); *Phys. Rev. D* **76**, 114001 (2007); *Phys. Rev. D* **77**, 034006 (2008); *Phys. Rev. D* **79**, 074014 (2009).
- [7] A.H. Fariborz and R. Jora, *Phys. Rev. D* **96**, 096021 (2017); *Phys. Rev. D* **95**, 116003 (2017); *Phys. Rev. D* **95**, 114001 (2017).