

Gallai-Ramsey number of odd cycles with chords

Fangfang Zhang^{1,*}, Zi-Xia Song^{2,†}, Yaojun Chen^{1,‡}

¹*Department of Mathematics, Nanjing University, Nanjing 210093, China*

²*Department of Mathematics, University of Central Florida, Orlando, FL 32816, USA*

Abstract

A Gallai coloring is a coloring of the edges of a complete graph without rainbow triangles, and a Gallai k -coloring is a Gallai coloring that uses at most k colors. For an integer $k \geq 1$, the Gallai-Ramsey number $GR_k(H)$ of a given graph H is the least positive integer N such that every Gallai k -coloring of the complete graph K_N contains a monochromatic copy of H . Let C_m denote the cycle on $m \geq 4$ vertices and let Θ_m denote the family of graphs obtained from C_m by adding an additional edge joining two non-consecutive vertices. We prove that $GR_k(\Theta_{2n+1}) = n \cdot 2^k + 1$ for all $k \geq 1$ and $n \geq 3$. This implies that $GR_k(C_{2n+1}) = n \cdot 2^k + 1$ all $k \geq 1$ and $n \geq 3$. Our result yields a unified proof for the Gallai-Ramsey number of all odd cycles on at least five vertices.

Keywords: Gallai coloring, Ramsey theory, cycles, rainbow triangle

2020 Mathematics Subject Classification: 05C55; 05D10; 05C15

1 Introduction

In this paper we consider graphs that are finite, simple and undirected. We use P_m , C_m and K_m to denote the path, cycle and complete graph on m vertices, respectively. For $m \geq 4$, let Θ_m denote the family of graphs obtained from C_m by adding an additional edge joining two non-consecutive vertices. For any positive integer k , we write $[k]$ for the set $\{1, 2, \dots, k\}$. Given an integer $k \geq 1$ and graphs H_1, \dots, H_k , the classical *Ramsey number* $R(H_1, \dots, H_k)$ is the least integer N such

*This work was done while the first author visited the University of Central Florida as a visiting student. The hospitality of the hosting institution is greatly acknowledged. The visit was funded by the Chinese Scholarship Council. E-mail address: fangfangzh@smail.nju.edu.cn.

†Partially supported by the National Science Foundation of China under Grant No. DMS-1854903. E-mail address: Zixia.Song@ucf.edu.

‡Yaojun Chen and Fangfang Zhang are partially supported by the National Natural Science Foundation of China under grant numbers 11671198 and 11871270. E-mail address: yaojunc@nju.edu.cn.

that every k -coloring of the edges of K_N contains a monochromatic copy of H_i in color i for some $i \in [k]$. When $H = H_1 = \dots = H_k$, we simply write $R_k(H)$ to denote the k -color Ramsey number of H . In the seminal paper of Ramsey [29], it is shown that Ramsey numbers are finite. This was rediscovered independently by Erdős and Szekeres [11]. Since the 1970s, Ramsey theory has grown into one of the most active areas of research in combinatorics, overlapping variously with graph theory, number theory, geometry and logic. However, determining Ramsey numbers is notoriously difficult in general.

Let (G, τ) denote a k -colored complete graph, where G is a complete graph and $\tau : E(G) \rightarrow [k]$. We say (G, τ) is \mathcal{F} -free if G does not contain a monochromatic copy of a graph in a given family \mathcal{F} under the k -coloring τ ; when $\mathcal{F} = \{F\}$, we simply say (G, τ) is F -free. By abusing notation, we say (G, τ) contains a *monochromatic copy of \mathcal{F}* if G contains a monochromatic copy of a graph in \mathcal{F} under τ . One of the earliest and well-known problems is that of determining the Ramsey number $R_k(C_m)$. For odd cycles C_{2n+1} , Erdős and Graham [10] observed that one can naturally construct a C_{2n+1} -free (G, τ) by induction. Indeed when $k = 1$, simply take (G, τ) to be a 1-colored K_{2n} ; for $k > 1$, let (G, τ) be obtained by taking two disjoint copies of the construction for $k - 1$ and color every edge between the two copies with a new color. This construction showed that $R_k(C_{2n+1}) \geq n \cdot 2^k + 1$ for all $k \geq 1$ and $n \geq 2$, which led Bondy and Erdős [1] to make the following conjecture.

Conjecture 1.1 ([1]). *For all $k \geq 1$ and $n \geq 2$, $R_k(C_{2n+1}) = n \cdot 2^k + 1$.*

Recently, Jenssen and Skokan [22] showed that Conjecture 1.1 is true for all fixed k and all n sufficiently large. However, for all fixed n and all k sufficiently large, Day and Johnson [9] recently showed that $R_k(C_{2n+1}) > 2n \cdot (2 + \epsilon)^{k-1}$ for some constant $\epsilon = \epsilon(n) > 0$, and so Conjecture 1.1 is false when n is small with respect to k . For further results in this direction, we refer the reader to Graham, Rothchild and Spencer [16] and Radziszowski [26] for a dynamic survey.

In this paper we investigate Ramsey numbers of odd cycles and odd cycles with chords under Gallai colorings, where a *Gallai coloring* is a coloring of the edges of a complete graph without rainbow triangles (that is, a triangle with all its edges colored differently). Gallai colorings naturally arise in several areas including: information theory [23]; the study of partially ordered sets, as in Gallai's original paper [15] (his result was restated in [19] in the terminology of graphs); and the study of perfect graphs [5]. There are now a variety of papers which consider Ramsey-type problems in Gallai colorings (see, e.g., [2, 4, 7, 14, 17, 18, 20, 24, 27, 28]). More information on this topic can be found in [12, 13].

A *Gallai k -coloring* is a Gallai coloring that uses at most k colors. Given an integer $k \geq 1$ and graphs H_1, \dots, H_k , *Gallai-Ramsey number* $GR(H_1, \dots, H_k)$ is defined to be the least integer N such that every Gallai k -coloring of K_N contains a monochromatic copy of H_i in color i for some

$i \in [k]$. When $H = H_1 = \dots = H_k$, we simply write $GR_k(H)$. Clearly, $GR_k(H) \leq R_k(H)$ for all $k \geq 1$ and $GR(H_1, H_2) = R(H_1, H_2)$. Theorem 1.2 below is a result of Gyárfás, Sárközy, Sebő and Selkow [18] which characterizes the general behavior of $GR_k(H)$.

Theorem 1.2 ([18]). *Let H be a fixed graph with no isolated vertices and let $k \geq 1$ be an integer. Then $GR_k(H)$ is exponential in k if H is not bipartite, linear in k if H is bipartite but not a star, and constant (does not depend on k) when H is a star.*

It turns out that for some graphs H (e.g., when $H = K_3$), $GR_k(H)$ behaves nicely, while the order of magnitude of $R_k(H)$ seems hopelessly difficult to determine. We will utilize the following important structural result of Gallai [15].

Theorem 1.3 ([15]). *Let (G, τ) be a Gallai k -colored complete graph with $|V(G)| \geq 2$. Then $V(G)$ can be partitioned into nonempty sets V_1, \dots, V_p with $p \geq 2$ so that at most two colors are used on the edges in $E(G) \setminus (E(V_1) \cup \dots \cup E(V_p))$ and only one color is used on the edges between any fixed pair (V_i, V_j) under τ , where $E(V_i)$ denotes the set of edges with both ends in V_i for all $i \in [p]$.*

The partition $\{V_1, \dots, V_p\}$ given in Theorem 1.3 is a *Gallai partition* of (G, τ) . Let (\mathcal{R}, τ) be obtained from (G, τ) by first contracting each V_i into a single vertex v_i and then coloring $v_i v_j$ by the unique color on the edges between V_i and V_j in (G, τ) . We say (\mathcal{R}, τ) is the *reduced graph* of (G, τ) corresponding to the Gallai partition $\{V_1, \dots, V_p\}$. Note that $\mathcal{R} = K_p$. By Theorem 1.3, all edges in \mathcal{R} are colored by at most two colors under τ . One can see that any monochromatic copy of H in (\mathcal{R}, τ) will result in a monochromatic copy of H in (G, τ) . It is not surprising that the 2-color Ramsey number $R_2(H)$ plays an important role in determining the value of $GR_k(H)$ when H is a complete graph. Fox, Grinshpun and Pach [12] posed the following conjecture.

Conjecture 1.4 ([12]). *For all $k \geq 1$ and $t \geq 3$,*

$$GR_k(K_t) = \begin{cases} (R_2(K_t) - 1)^{k/2} + 1 & \text{if } k \text{ is even} \\ (t-1)(R_2(K_t) - 1)^{(k-1)/2} + 1 & \text{if } k \text{ is odd.} \end{cases}$$

The first case of Conjecture 1.4 follows directly from a result of Chung and Graham [8] in 1983. A simpler proof of the first case of Conjecture 1.4 can be found in [18]. The next open case, when $t = 4$, was recently settled in [24].

In this paper, we focus on determining the exact values of $GR_k(\Theta_{2n+1})$ for all $k \geq 1$ and $n \geq 3$, and $GR_k(C_{2n+1})$ for all $k \geq 1$ and $n \geq 2$. It is worth noting that the construction of Bondy and Erdős mentioned earlier for $R_k(C_{2n+1})$ contains no rainbow triangles. Hence $GR_k(\Theta_{2n+1}) \geq n \cdot 2^k + 1$ and $GR_k(C_{2n+1}) \geq n \cdot 2^k + 1$ for all $k \geq 1$ and $n \geq 2$. More recently, the exact value of $GR_k(C_{2n+1})$ for $2 \leq n \leq 7$ has been completely settled, see [2, 3, 4, 14]; the exact value of $GR_k(\Theta_5)$ was recently

settled in [25]. We determine the exact value of $GR_k(\Theta_{2n+1})$ for all $k \geq 1$ and $n \geq 3$. We include C_5 in Theorem 1.5 below in order to provide a unified proof for the Gallai-Ramsey number of all odd cycles on at least five vertices.

Theorem 1.5. *For all $k \geq 1$, $GR_k(C_5) = 2 \cdot 2^k + 1$ and $GR_k(\Theta_{2n+1}) = n \cdot 2^k + 1$ for all $n \geq 3$.*

Note that every monochromatic copy of Θ_{2n+1} contains a monochromatic copy of C_{2n+1} . Theorem 1.5 not only implies the exact value for $GR_k(C_{2n+1})$ but also provides a unified proof for the Gallai-Ramsey number of C_{2n+1} for all $k \geq 1$ and $n \geq 2$.

Theorem 1.6. *For all $k \geq 1$ and $n \geq 2$, $GR_k(C_{2n+1}) = n \cdot 2^k + 1$.*

Theorem 1.6 implies that Conjecture 1.1 is true under Gallai colorings. We want to point out here that Wang et al. have also posted a paper [30] claiming the result of Theorem 1.6. We were thus compelled to share our independent proof in the manuscript [31, Theorem 1.7] which was not submitted for publication. Our proof of Theorem 1.5 given in Section 3 relies on an upper bound result for Gallai-Ramsey number of even cycles in [32, Theorem 1.6]. However, the proof of [32, Theorem 1.6] is rather long. In this paper we prove Theorem 1.5 using a more relaxed upper bound on $GR_k(C_{2n})$ given in Theorem 1.8 below. We include a short proof of Theorem 1.8 here for completeness. We need to introduce more notation before we state the result.

For the remainder of the paper, we use (G, τ) to denote a Gallai k -colored complete graph, where G is a complete graph and $\tau : E(G) \rightarrow [k]$ is a Gallai k -coloring. For each (G, τ) , let G_i^τ denote the spanning subgraph of G with $E(G_i^\tau) := \{e \in E(G) \mid \tau(e) = i\}$ for all $i \in [k]$. We simply write G_r^τ if the color i is red; G_b^τ if the color i is blue. For every $n \in \mathbb{N}$, let $q_\tau(G, n)$ denote the number of colors $i \in [k]$ such that G_i^τ has a component of order at least n . Then $q_\tau(G, n) \leq k$. We begin with Lemma 1.7 (which follows directly from the proof of Lemma 9 given in [20]). A proof of Lemma 1.7 can be found in [32, Lemma 1.5].

Lemma 1.7 ([20]). *Let (G, τ) be a Gallai k -colored complete graph with $|V(G)| \geq n \geq 2$. Let $\{V_1, \dots, V_p\}$ be a Gallai partition of (G, τ) with $p \geq 2$ as small as possible. Then $q_\tau(G, n) \geq 1$, and for any color i on the edges in (\mathcal{R}, τ) , \mathcal{R}_i^τ is connected.*

Theorem 1.8 below establishes an upper bound for $GR_k(C_{2n})$ for all $k \geq 1$ and $n \geq 2$.

Theorem 1.8. *Let (G, τ) be a Gallai k -colored complete graph with $|V(G)| \geq n \geq 2$. If*

$$|V(G)| \geq (n-1) \cdot q_\tau(G, n) + 2n + 2,$$

then (G, τ) has a monochromatic copy of C_{2n} .

We conclude this section by introducing more notation and list several known results that will be used in the proof of Theorem 1.8 and Theorem 1.5. Given a graph G and a set $S \subseteq V(G)$, we use $|G|$ to denote the number of vertices of G , $G \setminus S$ the subgraph obtained from G by deleting all vertices in S , and $G[S]$ the subgraph obtained from G by deleting all vertices in $V(G) \setminus S$. For two disjoint sets $A, B \subseteq V(G)$, A is *complete* to B in G if each vertex in A is adjacent to all vertices in B , and *anti-complete* to B in G if no vertex in A is adjacent to any vertex in B . Let (G, τ) be a Gallai k -colored complete graph. For two disjoint sets $A, B \subseteq V(G)$, A is *mc-complete* to B if all the edges between A and B in (G, τ) are colored the same color. We simply say A is *j -complete* to B if all the edges between A and B in (G, τ) are colored by some color $j \in [k]$, and A is *blue-complete* to B if all the edges between A and B in (G, τ) are colored blue. We say a vertex $x \in V(G)$ is *blue-adjacent* to a vertex $y \in V(G)$ if the edge xy is colored blue in (G, τ) , and x is *blue-complete* to an edge $yz \in E(G)$ if x is blue-complete to $\{y, z\}$ in (G, τ) . Similar definitions hold when blue is replaced by another color. For convenience, we use $A \setminus B$ to denote $A - B$; and $A \setminus b$ to denote $A \setminus \{b\}$ when $B = \{b\}$. We use the convention “ $S :=$ ” to mean that S is defined to be the right-hand side of the relation.

Theorem 1.9 ([1, 21]). *For all $n \geq 2$, $R_2(C_{2n+1}) = R_2(\Theta_{2n+1}) = 4n + 1$.*

For a bipartite graph G with bipartition (M, N) , let $\delta(N) := \min\{d_G(x) : x \in N\}$ and $\Delta(N) := \max\{d_G(x) : x \in M\}$. Lemma 1.10 follows from Lemma 12, Lemma 13, Lemma 14 in [20].

Lemma 1.10 ([20]). *Let G be a bipartite graph with bipartition (M, N) such that $|N| \geq 2$, $|M| \geq 4$ and $\delta(N) \geq (|M| + 1)/2$. Then G has a cycle of length 2ℓ for any ℓ satisfying $2 \leq \ell \leq \min\{|N|, \delta(N) - 1\}$, or $\Delta(N) = (|M| + 1)/2$, $M = M_1 \cup M_2 \cup M_3$ and $N = N_1 \cup N_2$, where M_1, M_2, M_3, N_1, N_2 are non-empty, pairwise disjoint sets in G , $|M_3| = 1$, $|M_1| = |M_2|$, and N_i is complete to $M_i \cup M_3$ but anti-complete to M_{3-i} for all $i \in [2]$.*

Lemma 1.11 ([2]). *Let $\{V_1, \dots, V_p\}$ be a Gallai partition of (G, τ) with $|V_1| \leq \dots \leq |V_p|$ and $|G| \geq 2n + 1$. If $|V_p| \leq n$ and the corresponding reduced graph \mathcal{R} is monochromatic, say blue, then (G, τ) contains a blue C_{2n+1} .*

Lemma 1.12 ([2]). *Let Y, Z be two disjoint sets of vertices in (G, τ) with $|Y| \geq n$ and $|Z| \geq n$. If (G, τ) is C_{2n+1} -free and Y is mc-complete, say blue-complete, to Z , then no vertex in $V(G) \setminus (Y \cup Z)$ is blue-complete to $Y \cup Z$ in (G, τ) . Moreover, if $|Z| \geq n + 1$, then $(G[Z], \tau)$ has no blue edges. Similarly, if $|Y| \geq n + 1$, then $(G[Y], \tau)$ has no blue edges.*

2 Proof of Theorem 1.8

Let (G, τ) and k be as given in the statement. Suppose (G, τ) is C_{2n} -free. Choose (G, τ) so that $q_\tau(G, n)$ is minimum. Let $q := q_\tau(G, n)$. By Lemma 1.7, $q \geq 1$. We may assume that for each color

$i \in [q]$, G_i^τ has a component of order at least n . Let X_1, \dots, X_q be disjoint subsets of $V(G)$ such that for each $i \in [q]$, X_i (possibly empty) is mc-complete in color i to $V(G) \setminus \bigcup_{i=1}^q X_i$. Choose X_1, \dots, X_q so that $|G| - \sum_{i=1}^q |X_i| \geq n$ and $\sum_{i=1}^q |X_i|$ is as large as possible. Denote $X := \bigcup_{i=1}^q X_i$. Then $|G \setminus X| \geq n \geq 2$. Since (G, τ) has no rainbow triangle, we see that for all $i, j \in [q]$ with $i \neq j$, all the edges between X_i and X_j are colored by color i or color j . We next prove a series of claims.

Claim 2.1. *For all $i \in [q]$, $|X_i| \leq n - 1$.*

Proof. Suppose $|X_i| \geq n$ for some color $i \in [q]$, say blue. By the choice of X_1, \dots, X_q , X_i is blue-complete to $V(G) \setminus X$. Thus (G, τ) has a blue C_{2n} using edges between X_i and $V(G) \setminus X$, a contradiction. ■

Let V_1, \dots, V_p be a Gallai partition of $(G \setminus X, \tau)$ with $p \geq 2$ as small as possible. We may assume that $|V_1| \leq \dots \leq |V_p|$. By Theorem 1.3 and Lemma 1.7, all the edges of the reduced graph of $(G \setminus X, \tau)$ are colored by at most two colors in $[q]$, say red or blue. Then for all $i \in [p-1]$, V_i is either red- or blue-complete to V_p in $(G \setminus X, \tau)$. Let

$$\begin{aligned} \mathcal{V}_r &:= \{V_i \in \{V_1, \dots, V_{p-1}\} \mid V_i \text{ is red-complete to } V_p \text{ in } (G \setminus X, \tau)\} \text{ and} \\ \mathcal{V}_b &:= \{V_i \in \{V_1, \dots, V_{p-1}\} \mid V_i \text{ is blue-complete to } V_p \text{ in } (G \setminus X, \tau)\}. \end{aligned}$$

Let $R = \bigcup_{V_j \in \mathcal{V}_r} V_j$ and $B = \bigcup_{V_j \in \mathcal{V}_b} V_j$. Then $R \cup B = V(G) \setminus (X \cup V_p)$, and R and B are disjoint. We may further assume that X_1 is red-complete to $V(G) \setminus X$ and X_2 is blue-complete to $V(G) \setminus X$.

Claim 2.2. $\min\{|B|, |R|\} \geq 1$ and so $q \geq 2$.

Proof. Suppose $\min\{|B|, |R|\} = 0$. We may assume that $B = \emptyset$. Since $p \geq 2$, we see that $|R| \geq 1$ and R is red-complete to V_p . Then $|V_p| \leq n - 1$, else, let $X'_1 := X_1 \cup R$ and $X'_i := X_i$ for all $i \in \{2, \dots, q\}$. But then $|X'_1| + \dots + |X'_q| = |X \cup R| > |X|$, contrary to the choice of X_1, \dots, X_q . Similarly, $|R| \leq n - 1$. By Claim 2.1, $|X_i| \leq n - 1$ for all $i \in [q]$. But then $|G| = |X| + |V_p| + |R| \leq (n-1)q + 2(n-1)$, contrary to the assumption that $|G| \geq (n-1)q + 2n + 2$. This proves that $\min\{|B|, |R|\} \geq 1$. By the choice of p and Lemma 1.7, both $G_r^\tau \setminus X$ and $G_b^\tau \setminus X$ are connected. Thus $q \geq 2$. ■

Claim 2.3. $|V_p| \leq n - 1$ and so $X_i \neq \emptyset$ for any color $i \in [q]$ that is neither red nor blue.

Proof. Suppose $|V_p| \geq n$. Then every vertex in $B \cup R$ is either red- or blue-complete to V_p . Let $X'_1 := X_1 \cup R$, $X'_2 := X_2 \cup B$, and $X'_i := X_i$ for all $i \in \{3, \dots, q\}$. But then $|X'_1| + \dots + |X'_q| = |X \cup B \cup R| > |X|$, contrary to the choice of X_1, \dots, X_q . This proves that $|V_p| \leq n - 1$. Next, suppose there exists a color $j \in [q]$ such that j is neither red nor blue but $X_j = \emptyset$. Then no edges between pairs of X_1, \dots, X_q are colored by color j in (G, τ) . But then G_j^τ has no component of order at least n , because $|V_\ell| \leq |V_p| \leq n - 1$ for all $\ell \in [p-1]$, and $|X_i| \leq n - 1$ for all $i \in [q]$ by Claim 2.1, a contradiction. ■

Claim 2.4. $n \geq 3$ and $q = 2$.

Proof. Suppose $n = 2$. By Claim 2.1 and Claim 2.3, $|V_p| = 1$ and $|X_i| \leq 1$ for all $i \in [q]$. Then $|G \setminus \bigcup_{i=3}^q X_i| \geq q+6-(q-2) = 8$. Thus $(G \setminus \bigcup_{i=3}^q X_i, \tau)$ contains a red or blue C_4 because $R_2(C_4) = 6$ [6], a contradiction. Suppose next $n \geq 3$ and $q \neq 2$. By Claim 2.2, $q \geq 3$ and the color q is neither red nor blue. By Claim 2.3, $X_q \neq \emptyset$ and $|V_p| \leq n-1$. Thus $q_\tau(G \setminus X_q, n) = q_\tau(G, n) - 1 = q-1$. By Claim 2.1, $|X_q| \leq n-1$, and so $|G \setminus X_q| \geq (n-1)q+2n+2-(n-1) = (n-1) \cdot q_\tau(G \setminus X_q, n) + 2n+2$. By the minimality of q , $(G \setminus X_q, \tau)$ has a monochromatic copy of C_{2n} , a contradiction. ■

By Claim 2.4, $q = 2$ and $n \geq 3$ and so $|G| \geq 2(n-1) + 2n+2 = 4n$. We may further assume that $|X_1 \cup R| \geq |X_2 \cup B|$. Then $|X_1 \cup R| \geq \lceil (|G| - |V_p|)/2 \rceil \geq n+2$ and $|V_p \cup X_1 \cup R| \geq |V_p| + \lceil (|G| - |V_p|)/2 \rceil \geq 2n+1$. Let

$$W := \begin{cases} X_1 \cup R & \text{if } |X_1 \cup R| \leq 2n+1 \\ X_1 \cup \left(\bigcup_{V \in \mathcal{V}'_r} V \right) & \text{if } |X_1 \cup R| > 2n+1, \end{cases}$$

where $\mathcal{V}'_r \subseteq \mathcal{V}_r$ is chosen so that $|W| \leq 2n+1$ is as large as possible when $|X_1 \cup R| > 2n+1$. In both cases, we have $|V_p \cup W| \geq 2n+1$ because in the second case $|V_p| \geq |V_i|$ for all $i \in [p-1]$. Let $V_p^* \subseteq V_p$ be such that $|V_p^* \cup W| = 2n+1$. Let $M := V_p^* \cup W$ and $N := V(G) \setminus M$. Then $|N| \geq 4n - (2n+1) = 2n-1$ and $N = (V_p \setminus V_p^*) \cup B \cup (R \setminus W) \cup X_2$. By Claim 2.3, $|W| = |M \setminus V_p| \geq (2n+1) - (n-1) = n+2$. It follows that every vertex in $V_p \setminus V_p^*$ is red-adjacent to every vertex in W , and every vertex in $N \setminus V_p$ is either red- or blue-adjacent to at least $\lceil |M|/2 \rceil = n+1$ vertices in M . Let

$$N_r := \{v \in N \mid v \text{ is red-adjacent to at least } n+1 \text{ vertices in } M\} \text{ and}$$

$$N_b := \{v \in N \mid v \text{ is blue-adjacent to at least } n+1 \text{ vertices in } M\}.$$

Then $|N_r| + |N_b| = |N| \geq 2n-1$. Since every vertex in $V_p \setminus V_p^*$ is red-adjacent to at least $|W| \geq n+2$ vertices in M , we see that $N_b \cap V_p = \emptyset$ and $V_p \setminus V_p^* \subseteq N_r$. For each color $j \in \{r, b\}$, let H_j be the bipartite subgraph of G with bipartition (M, N_j) such that $E(H_j)$ consists of all edges in color j between M and N_j in (G, τ) . Then H_j is C_{2n} -free because (G, τ) is C_{2n} -free.

Suppose $|N_b| \geq n$. Since $N_b \cap V_p = \emptyset$, we see that every edge between M and N_b is colored red or blue in (G, τ) . Note that $\min\{d_{H_b}(v) : v \in N_b\} \geq n+1 = (|M|+1)/2$ and H_b is C_{2n} -free. By Lemma 1.10 applied to H_b with $\ell := n$, we see that $M = M_1 \cup M_2 \cup M_3$ and $N_b = N_1 \cup N_2$, where M_1, M_2, M_3, N_1, N_2 are non-empty, pairwise disjoint sets in H_b , $|M_3| = 1$, $|M_1| = |M_2| = n$, N_i is blue-complete to $M_i \cup M_3$ and red-complete to M_{3-i} for all $i \in [2]$. Since G contains no rainbow triangle, every edge between M_1 and M_2 is colored red or blue. Let $x_1, y_1, z_1 \in M_1$ and $x_2, y_2, z_2 \in M_2$ be all distinct. Since each edge between M_1 and M_2 is colored red or blue, we may

further assume that x_1x_2 and y_1y_2 are colored the same. For each $i \in [2]$, let Q_i^r be an (x_i, y_i) -path between M_i and N_{3-i} such that $|Q_1^r| + |Q_2^r| = 2n$; Q_i^b be an (x_i, y_i) -path between M_i and N_i such that $|Q_1^b| + |Q_2^b| = 2n$. But then (G, τ) contains a red C_{2n} with edge set $E(Q_1^r) \cup E(Q_2^r) \cup \{x_1y_1, x_2y_2\}$ when x_1x_2 and y_1y_2 are colored red, or a blue C_{2n} with edge set $E(Q_1^b) \cup E(Q_2^b) \cup \{x_1y_1, x_2y_2\}$ when x_1x_2 and y_1y_2 are colored blue, a contradiction.

It remains to consider the case $|N_b| \leq n - 1$. Then $|N_r| \geq n$ and $\min\{d_{H_r}(v) : v \in N_r\} \geq n + 1 = (|M| + 1)/2$. Note that H_r is C_{2n} -free. By Lemma 1.10 applied to H_r with $\ell = n$, we see that $M = M_1 \cup M_2 \cup M_3$ and $N_r = N_1 \cup N_2$, where M_1, M_2, M_3, N_1, N_2 are non-empty, pairwise disjoint sets in H_r , $|M_3| = 1$, $|M_1| = |M_2| = n$, N_i is red-complete to $M_i \cup M_3$ and no edges between N_i and M_{3-i} are colored red for each $i \in [2]$. Since every vertex in $V_p \setminus V_p^*$ is red-adjacent to at least $n + 2$ vertices in M , it follows that $N_r \cap V_p = \emptyset$. By the choice of W , N_i is blue-complete to M_{3-i} for each $i \in [2]$. Note that every edge between M_1 and M_2 is colored red or blue because (G, τ) contains no rainbow triangle. Similar to the argument in the previous paragraph, we obtain a monochromatic C_{2n} in (G, τ) , a contradiction.

This completes the proof of Theorem 1.8. ■

3 Proof of Theorem 1.5

Let n, k be as given in the statement. It suffices to show that $GR_k(C_5) \leq 2 \cdot 2^k + 1$ for all $k \geq 1$ and $GR_k(\Theta_{2n+1}) \leq n \cdot 2^k + 1$ for all $k \geq 1$ and $n \geq 3$. The case when $k = 1$ is trivial and the case when $k = 2$ follows from Theorem 1.9. So we may assume that $k \geq 3$. Let $G := K_{n \cdot 2^k + 1}$ and let $\tau : E(G) \rightarrow [k]$ be any Gallai k -coloring of G . Suppose that (G, τ) is Θ_{2n+1} -free when $n \geq 3$ and C_5 -free when $n = 2$. Choose (G, τ) with k minimum. Then τ is an onto mapping. Let X_1, \dots, X_k be disjoint subsets of $V(G)$ such that for each $i \in [k]$, X_i (possibly empty) is mc-complete in color i to $V(G) \setminus \bigcup_{i=1}^k X_i$ under τ . Choose X_1, \dots, X_k so that $\sum_{i=1}^k |X_i| \leq (k+1)n$ is as large as possible. Let $X := \bigcup_{i=1}^k X_i$. Then $|X| \leq (k+1)n$ and $|G \setminus X| \geq n \cdot 2^k + 1 - (k+1)n \geq 4n + 1$. Since (G, τ) has no rainbow triangle, we see that for all $i, j \in [k]$ with $i \neq j$, all the edges between X_i and X_j are colored by color i or color j . We next prove several claims.

Claim 3.1. (G, τ) is C_{2n+1} -free for all $n \geq 2$.

Proof. Suppose (G, τ) contains a monochromatic $C := C_{2n+1}$, say with vertices $x_1, x_2, \dots, x_{2n+1}$ in order. Then $n \geq 3$. We may assume that all edges of C are colored blue. Then no chord of C is colored blue because (G, τ) is Θ_{2n+1} -free. We may further assume that x_1x_3 is colored red. Then x_1x_j is colored red for all $j \in \{3, 4, \dots, 2n\}$ because (G, τ) has a no rainbow triangle. It follows that all chords of C are colored red. Let H be the graph with $V(H) = V(C)$ and $E(H)$ consisting

of all chords of C . Then H is the complement of C_{2n+1} . It can be easily checked that H contains a chorded C_{2n+1} because $n \geq 3$. Thus (G, τ) contains a red copy of Θ_{2n+1} , a contradiction. ■

Claim 3.2. For all $i \in [k]$, $|X_i| \leq n - 1$.

Proof. Suppose $|X_j| \geq n$ for some color $j \in [k]$. Since $|G \setminus X| \geq 4n + 1$, by Lemma 1.12 applied to X_j and $V(G) \setminus X$, we see that $(G \setminus X, \tau)$ has no edge in color j . By minimality of k , $|G \setminus X| \leq n \cdot 2^{k-1}$. But then

$$|G| = |X| + |G \setminus X| \leq (k + 1)n + n \cdot 2^{k-1} < n \cdot 2^k + 1$$

for all $k \geq 3$ and $n \geq 2$, a contradiction. ■

Claim 3.3. $X_i = \emptyset$ for some $i \in [k]$, and so $|X| \leq (k - 1)(n - 1)$.

Proof. Suppose $X_i \neq \emptyset$ for every $i \in [k]$. By Claim 3.2, $|X| \leq k(n - 1)$. Then

$$|G \setminus X| \geq n \cdot 2^k + 1 - k(n - 1) > (n - 1)k + 2n + 2,$$

for all $k \geq 3$ and $n \geq 2$. By Theorem 1.8 and the fact that $q_\tau(G \setminus X, n) \leq k$, we see that $(G \setminus X, \tau)$ contains a monochromatic copy of C_{2n} , say in color 1. Since $X_1 \neq \emptyset$, we see that (G, τ) contains a monochromatic copy of C_{2n+1} in color 1, contrary to Claim 3.1. This proves that $X_i = \emptyset$ for some $i \in [k]$. By Claim 3.2, $|X| \leq (k - 1)(n - 1)$. ■




Figure 1: An overview of (G, τ) .

Let $\{V_1, \dots, V_p\}$ be a Gallai partition of $(G \setminus X, \tau)$ with $p \geq 2$ and $|V_1| \leq \dots \leq |V_p|$. By Theorem 1.3, we may assume that every edge of the corresponding reduced graph (\mathcal{R}, τ) of $(G \setminus X, \tau)$ is colored either red or blue. By the choice of X_1, \dots, X_k , we may further assume that X_1 is red-complete to $V(G) \setminus X$ and X_2 is blue-complete to $V(G) \setminus X$. By Theorem 1.9, $p \leq 4n$. Thus

$|V_p| \geq 2$. Note that for all $i \in [p-1]$, V_i is either red- or blue-complete to V_p in (G, τ) . Let

$$\begin{aligned}\mathcal{V}_r &:= \{V_i \in \{V_1, \dots, V_{p-1}\} \mid V_i \text{ is red-complete to } V_p \text{ in } (G, \tau)\} \text{ and} \\ \mathcal{V}_b &:= \{V_j \in \{V_1, \dots, V_{p-1}\} \mid V_j \text{ is blue-complete to } V_p \text{ in } (G, \tau)\}.\end{aligned}$$

Let $R := \bigcup_{V_j \in \mathcal{V}_r} V_j$ and $B := \bigcup_{V_j \in \mathcal{V}_b} V_j$. Then $R \cup B = V(G) \setminus (X \cup V_p)$. An overview of (G, τ) is depicted in Figure 1.

Claim 3.4. $|R \cup B| \geq 2n + 2$.

Proof. Suppose $|R \cup B| \leq 2n + 1$. Since $p \geq 2$, we see that $|R \cup B| \geq 1$. Let $X'_1 := X_1 \cup R$, $X'_2 := X_2 \cup B$, and $X'_i := X_i$ for all $i \in \{3, \dots, k\}$. Then $V_p = V(G) \setminus \bigcup_{i=1}^k X'_i$, and for all $i \in [k]$, X'_i is mc-complete to V_p in (G, τ) . Thus

$$\sum_{i=1}^k |X_i| < \sum_{i=1}^k |X'_i| = |X \cup R \cup B| \leq (k-1)(n-1) + 2n + 1 = (k+1)n - (k-2) < (k+1)n,$$

contrary to the choice of X_1, \dots, X_k . ■

Claim 3.5. $|V_p| \geq n + 1$.

Proof. Suppose $|V_p| \leq n$. We may assume that $|R| \geq |B|$. Then

$$|R| + |B| = |G| - |V_p| - |X| \geq (n \cdot 2^k + 1) - n - (k-1)(n-1) = n \cdot 2^k - kn + k \geq 5n + 3.$$

Thus $|R| > 2n + 1$. By Lemma 1.11 and Claim 3.1, $(G[R], \tau)$ must have both red and blue edges. Suppose $|X_1 \cup V_p| \geq n$. By Lemma 1.12 applied to (G, τ) with $Y = R$ and $Z = X_1 \cup V_p$, we see that $(G[R], \tau)$ has no red edges, a contradiction. Thus $|X_1 \cup V_p| \leq n - 1$ and so $|V_p| \leq n - 1$. It follows that for any color $i \in [k]$ other than red and blue, $G_i^r \setminus X$ has no component of order at least n . Thus $q_\tau(G \setminus X, n) \leq 2$ and then $|G \setminus X| > (n-1) \cdot q_\tau(G \setminus X, n) + 2n + 2$. By Theorem 1.8, $(G \setminus X, \tau)$ contains a red or blue C_{2n} . This implies that either $X_1 = \emptyset$ or $X_2 = \emptyset$.

Suppose $|R| \geq 4n$. Let $\mathcal{V}'_r \subseteq \mathcal{V}_r$ be chosen so that $|\bigcup_{V_i \in \mathcal{V}'_r} V_i| \geq 4n$ is as small as possible. Let $U := \bigcup_{V_i \in \mathcal{V}'_r} V_i$. Then $|U| \leq 5n - 2$ because $|V_1| \leq \dots \leq |V_p| \leq n - 1$. Since $q_\tau(G \setminus X, n) \leq 2$, we see that $q_\tau(G[U], n) \leq 2$. Then $|U| \geq 4n \geq (n-1) \cdot q_\tau(G[U], n) + 2n + 2$. By Theorem 1.8, $(G[U], \tau)$ contains a monochromatic copy of cycle C of length $2n$, say with vertices u_1, u_2, \dots, u_{2n} in order. Let $U_1 = \{u_1, u_3, \dots, u_{2n-1}\}$ and $U_2 = \{u_2, u_4, \dots, u_{2n}\}$. Since U is red-complete to V_p , we see that the cycle C must be blue. Then $X_2 = \emptyset$. Let $x, y \in V_p$ be distinct. Then

$$|V(G) \setminus (X \cup U \cup \{x, y\})| \geq n \cdot 2^k + 1 - (k-1)(n-1) - (5n-2) - 2 \geq n + 3.$$

Let $W \subseteq V(G) \setminus (X \cup U \cup \{x, y\})$ with $|W| = n$. By the choice of U , every edge between W and $V(C)$ is colored either red or blue in (G, τ) . Note that no vertex in W is blue-adjacent to any edge on the cycle C , else, we obtain a blue C_{2n+1} , contrary to Claim 3.1. It follows that each vertex in W is red-adjacent to at least n vertices on C . Let

$$\begin{aligned} W_0 &:= \{v \in W \mid v \text{ is red-adjacent to at least } n+1 \text{ vertices on } C\}, \\ W_1 &:= \{v \in W \setminus W_0 \mid v \text{ is red-complete to } U_1 \text{ only in } G\} \text{ and} \\ W_2 &:= \{v \in W \setminus W_0 \mid v \text{ is red-complete to } U_2 \text{ only in } G\}. \end{aligned}$$

Then $W = W_0 \cup W_1 \cup W_2$ and W_i is blue-complete to U_{3-i} for each $i \in [2]$. Let H be the bipartite subgraph of G with bipartition $(V(C), W)$ such that $E(H)$ consists of all the red edges between W and $V(C)$ in (G, τ) . Suppose H has a path P on $2n - 3$ vertices with both ends in $V(C)$. We may assume that u_1, u_j are the ends of P for some $j \in [2n]$ with $j \neq 1$. Since $|U \cap V(P)| = n - 1$, we see that $|U \setminus V(P)| \geq 4n - (n - 1) = 3n + 1$. By Lemma 1.11, $(G[U \setminus V(P)], \tau)$ must contain a red edge, say uv . But then (G, τ) has a red C_{2n+1} with edge set $E(P) \cup \{xu_1, xu, uv, vy, yu_j\}$, contrary to Claim 3.1. Thus H has no path P on $2n - 3$ vertices with both ends in $V(C)$. It follows that $|W_1| \leq n - 3$ and $|W_2| \leq n - 3$.

We next show that $|W_0| \geq 1$. Suppose $W_0 = \emptyset$. Then $|W_1| + |W_2| = |W| = n$. We may assume that $|W_1| \leq |W_2|$. Then $|W_1| \geq 3$ because $|W_2| \leq n - 3$. Recall that $|U_1| = n$ and $|V_p| \leq n - 1$, we see that $(G[U_1], \tau)$ contains an edge e that is colored red or blue. We may assume that $e = u_1 u_{2j+1}$ for some $j \in [n - 1]$. Then e must be red, else let $v \in W_1$, we obtain a blue C_{2n+1} with vertices $v, u_{2n}, u_{2n-1}, \dots, u_{2j+1}, u_1, u_2, \dots, u_{2j}$ in order, contrary to Claim 3.1. Let $v_1 \in U_1 \setminus \{u_1, u_{2j+1}\}$ and $v_2 \in U_2 \setminus u_2$; let Q_1 be a (u_1, v_1) -path on $2|W_1| + 1$ vertices using edges between W_1 and $U_1 \setminus u_{2j+1}$ and Q_2 be a (u_2, v_2) -path on $2|W_2| - 3$ vertices using edges between W_2 and U_2 . But then (G, τ) contains a red C_{2n+1} with edge set $E(Q_1) \cup E(Q_2) \cup \{xu_{2j+1}, u_{2j+1}u_1, v_1y, yv_2, u_2x\}$, contrary to Claim 3.1. This proves that $|W_0| \geq 1$, as claimed.

Finally, let H' be obtained from H by adding a new vertex w adjacent to all vertices in W . Then H' is a bipartite graph with bipartition $(V(C) \cup \{w\}, W)$. Note that for each vertex $v \in W$, $d_{H'}(v) \geq (|V(C) \cup \{w\}| + 1)/2$; for each vertex $v \in W_0 \neq \emptyset$, $d_{H'}(v) > (|V(C) \cup \{w\}| + 1)/2$. By Lemma 1.10 applied to H' with $M := V(C) \cup \{w\}$, $N := W$ and $\ell := n$, we see that H' contains a cycle C_{2n} and thus H has a path P on $2n - 3$ vertices with both ends on C , contrary to the fact that H has no path on $2n - 3$ vertices with both ends in $V(C)$.

It remains to consider the case $|R| \leq 4n - 1$. In this case,

$$|B| = |V(G) \setminus (X \cup V_p \cup R)| \geq n \cdot 2^k + 1 - (k - 1)(n - 1) - (n - 1) - (4n - 1) \geq n + 5.$$

By Lemma 1.12, $|X_1 \cup V_p| \leq n$ and $|X_2 \cup V_p| \leq n$. Since $X_1 = \emptyset$ or $X_2 = \emptyset$, we see that $|X_1 \cup X_2 \cup V_p| \leq n$. By Claim 3.3, $|X \cup V_p| = |X_3| + \dots + |X_k| + |X_1 \cup X_2 \cup V_p| \leq (k-2)(n-1) + n$. Thus

$$|B| + |R| = |G| - |X \cup V_p| \geq n \cdot 2^k + 1 - (k-2)(n-1) - n \geq 6n + 2,$$

which implies $|B| \geq 2n + 3$. Then $|R| \geq |B| \geq 2n + 3$. By Lemma 1.11, $(G[R], \tau)$ must have a red edge, say $x_r y_r$; $(G[B], \tau)$ must have a blue edge, say $x_b y_b$. Let $W \subseteq R \setminus \{x_r, y_r\}$ be such that $|W| = 2n + 1$. Then all the edges between B and W are colored red or blue. Let

$$\begin{aligned} B_r &:= \{v \in B \setminus \{x_b, y_b\} \mid v \text{ is red-adjacent to at least } n+1 \text{ vertices in } W\} \text{ and} \\ B_b &:= \{v \in B \setminus \{x_b, y_b\} \mid v \text{ is blue-adjacent to at least } n+1 \text{ vertices in } W\}. \end{aligned}$$

Then $|B_r| \geq n+1$ or $|B_b| \geq n+1$ because $|B \setminus \{x_b, y_b\}| \geq 2n+1$. We may assume that $|B_j| \geq n+1$ for some color $j \in \{r, b\}$. Let H_j be the bipartite subgraph of G with bipartition (W, B_j) such that $E(H_j)$ consists of all edges in color j between W and B_j in (G, τ) . For each vertex $v \in B_j$, $d_{H_j}(v) \geq (|W| + 1)/2$. By Lemma 1.10 applied to H_j with $M := W$, $N := B_j$ and $\ell := n$, we see that H_j has a path P in color j on $2n - 3$ vertices with both ends in W when $j = r$ and both ends in B_j when $j = b$. Let v_1, v_2 be the two ends of P . But then (G, τ) contains a monochromatic copy of C_{2n+1} in color j with edge set $E(P) \cup \{v_1 x, x x_j, x_j y_j, y_j y, y v_2\}$, contrary to Claim 3.1. \blacksquare

Claim 3.6. $|V_{p-2}| \leq n$ if $p \geq 3$.

Proof. Suppose $|V_{p-2}| \geq n + 1$. Then $n + 1 \leq |V_{p-2}| \leq |V_{p-1}| \leq |V_p|$. Let B_1, B_2, B_3 be a permutation of V_{p-2}, V_{p-1}, V_p . Since (G, τ) has no rainbow triangle, we may assume that B_2 is, say blue-complete, to $B_1 \cup B_3$ in (G, τ) . Then B_1 must be red-complete to B_3 in (G, τ) , else we obtain a blue C_{2n+1} , contrary to Claim 3.1. By Lemma 1.12, no vertex in X is either red- or blue-complete to $V(G) \setminus X$. Thus $X_1 = X_2 = \emptyset$ and $|X| \leq (k-2)(n-1)$. Let $A := V(G) \setminus (B_1 \cup B_2 \cup B_3 \cup X)$. By Lemma 1.12, no vertex in A is red-complete to $B_1 \cup B_3$ in (G, τ) , and no vertex in A is blue-complete to $B_1 \cup B_2$ or $B_2 \cup B_3$ in (G, τ) . This implies that A must be red-complete to B_2 in (G, τ) . Suppose that $(G[A], \tau)$ has a blue edge, say, uv . We may assume that u is blue-complete to B_1 . But then we obtain a blue C_{2n+1} , because v is blue-complete to B_1 or B_3 . This proves that $(G[A], \tau)$ has no blue edges. By Lemma 1.12, $(G[B_2], \tau)$ has no blue edges, and neither $(G[B_1], \tau)$ nor $(G[B_3], \tau)$ has red or blue edges. By minimality of k , $|B_1| \leq n \cdot 2^{k-2}$ and $|B_3| \leq n \cdot 2^{k-2}$. Let τ' be obtained from τ by recoloring all the blue edges in $G[X]$ red. Then τ' is a Gallai $(k-1)$ -coloring of $G[B_2 \cup A \cup X]$. Moreover, $(G[B_2 \cup A \cup X], \tau')$ has no blue edges and no monochromatic copy of Θ_{2n+1} when $n \geq 3$ and no monochromatic C_5 when $n = 2$. By minimality of k , $|B_2 \cup A \cup X| \leq n \cdot 2^{k-1}$. But then

$$|G| = |B_1| + |B_2 \cup A \cup X| + |B_3| \leq n \cdot 2^{k-2} + n \cdot 2^{k-1} + n \cdot 2^{k-2} < n \cdot 2^k + 1,$$

for all $k \geq 3$ and $n \geq 2$, a contradiction. ■

Claim 3.7. $|R| \geq n + 1$ and $|B| \geq n + 1$.

Proof. Suppose $|R| \leq n$ or $|B| \leq n$, say the latter. By Claim 3.5 and 3.4, $|V_p| \geq n+1$ and $|R| \geq n+2$. By Lemma 1.12 applied to V_p and R , we see that $X_1 = \emptyset$ and neither $(G[V_p], \tau)$ nor $(G[R], \tau)$ has red edges. Let $i \in [k]$ be a color, say green, that is neither red nor blue. Let τ' be obtained from τ by recoloring all the red edges in $(G[B \cup (X \setminus X_i)], \tau)$ by color green; and recoloring all the red edges in $(G[X_i], \tau)$ by color blue. Then neither $(G[R \cup X_i], \tau')$ nor $(G[B \cup V_p \cup (X \setminus X_i)], \tau')$ has red edges. It follows that τ' is a Gallai $(k-1)$ -coloring of both $G[R \cup X_i]$ and $G[B \cup V_p \cup (X \setminus X_i)]$, and neither $(G[R \cup X_i], \tau')$ nor $(G[B \cup V_p \cup (X \setminus X_i)], \tau')$ has a monochromatic copy of Θ_{2n+1} when $n \geq 3$ and monochromatic C_5 when $n = 2$. By minimality of k , $|R \cup X_i| \leq n \cdot 2^{k-1}$ and $|B \cup V_p \cup (X \setminus X_i)| \leq n \cdot 2^{k-1}$. Then

$$|G| = |R \cup X_i| + |B \cup V_p \cup (X \setminus X_i)| \leq n \cdot 2^{k-1} + n \cdot 2^{k-1} < n \cdot 2^k + 1,$$

for all $k \geq 3$ and $n \geq 2$, a contradiction. ■

By Claim 3.7, $|R| \geq n + 1$ and $|B| \geq n + 1$. By Claim 3.5, $|V_p| \geq n + 1$. By Lemma 1.12 applied to V_p and R , and V_p and B , respectively, we have $|X_1| = |X_2| = 0$; $(G[R], \tau)$ has no red edges; $(G[B], \tau)$ has no blue edges; and $(G[V_p], \tau)$ has neither red nor blue edges. By Claim 3.2, $|X| \leq (k-2)(n-1)$. By minimality of k , $|V_p| \leq n \cdot 2^{k-2}$. We may further assume that $V_{p-1} \subseteq B$. By Claim 3.6, $|V_{p-2}| \leq n$ if $p \geq 3$. By Lemma 1.11, $|B \setminus V_{p-1}| \leq 2n$ and $|R| \leq 2n$. If $k \geq 4$ or $|X| = 0$, then

$$\begin{aligned} |G| &= |V_p| + |V_{p-1}| + |B \setminus V_{p-1}| + |R| + |X| \\ &\leq \begin{cases} n \cdot 2^{k-2} + n \cdot 2^{k-2} + 2n + 2n + (k-2)(n-1) & \text{if } k \geq 4 \\ n \cdot 2^{k-2} + n \cdot 2^{k-2} + 2n + 2n & \text{if } k = 3, |X| = 0 \end{cases} \\ &< n \cdot 2^k + 1, \end{aligned}$$

a contradiction. Thus $k = 3$ and $X \neq \emptyset$. We may assume that X is green-complete to $V(G) \setminus X$. Then all the edges of $G[V_p]$ are colored green. Thus $|X \cup V_p| \leq 2n$, else we obtain a green copy of Θ_{2n+1} . It follows that $|V_{p-1}| \leq |V_p| \leq 2n - 1$. But then

$$|G| = |X \cup V_p| + |V_{p-1}| + |B \setminus V_{p-1}| + |R| \leq 2n + (2n - 1) + 2n + 2n < 8n + 1,$$

for all $n \geq 2$, a contradiction.

This completes the proof of Theorem 1.5. ■

Acknowledgements

The authors would like to thank Christian Bosse and Jingmei Zhang for their helpful discussion.

References

- [1] J. A. Bondy, P. Erdős, Ramsey numbers for cycles in graphs, *J. Combin. Theory Ser. B* 14 (1973) 46–54.
- [2] C. Bosse, Z-X. Song, Multicolor Gallai-Ramsey numbers of C_9 and C_{11} . arXiv:1802.06503.
- [3] C. Bosse, Z-X. Song, J. Zhang, Improved upper bounds for Gallai-Ramsey numbers of odd cycles. arXiv:1808.09963.
- [4] D. Bruce, Z-X. Song, Gallai-Ramsey numbers of C_7 with multiple colors, *Discrete Math.* 342 (2019) 1191–1194.
- [5] K. Cameron, J. Edmonds, L. Lovász, A note on perfect graphs, *Period. Math. Hungar.* 17 (1986) 173–175.
- [6] G. Chartrand, S. Schuster, On the existence of specified cycles in complementary graphs, *Bull. Amer. Math. Soc.* 77 (1971) 995–998.
- [7] M. Chen, Y. Li, C. Pei, Gallai-Ramsey numbers of odd cycles and complete bipartite graphs, *Graphs and Combin.* 34 (2018) 1185–1196.
- [8] F. R. K. Chung, R. Graham, Edge-colored complete graphs with precisely colored subgraphs, *Combinatorica* 3 (1983) 315–324.
- [9] A. N. Day, J. R. Johnson, Multicolour Ramsey numbers of odd cycles, *J. Combin. Theory Ser. B* 124 (2017) 56–63.
- [10] P. Erdős, R. L. Graham, On partition theorems for finite graphs, *Colloq. Math. Soc. János Bolyai* 10 (1973) 515–527.
- [11] P. Erdős, G. Szekeres, A combinatorial problem in geometry, *Compositio Math.* 2 (1935) 463–470.
- [12] J. Fox, A. Grinshpun, J. Pach, The Erdős-Hajnal conjecture for rainbow triangles, *J. Combin. Theory Ser. B* 111 (2015) 75–125.
- [13] S. Fujita, C. Magnant, K. Ozeki, Rainbow generalizations of Ramsey theory: a survey, *Graphs Combin.* 26 (2010) 1–30.
- [14] S. Fujita, C. Magnant, Gallai-Ramsey numbers for cycles, *Discrete Math.* 311 (2011) 1247–1254.
- [15] T. Gallai, Transitiv orientierbare Graphen, *Acta Math. Acad. Sci. Hung.* 18 (1967) 25–66.
- [16] R. L. Graham, B. L. Rothschild, J. H. Spencer, *Ramsey Theory*, Wiley, New York, 1990.

- [17] A. Gyárfás, G. Sárközy, Gallai colorings of non-complete graphs, *Discrete Math.* 310 (2010) 977–980.
- [18] A. Gyárfás, G. Sárközy, A. Sebő, S. Selkow, Ramsey-type results for Gallai colorings, *J. Graph Theory* 64 (2010) 233–243.
- [19] A. Gyárfás, G. Simonyi, Edge colorings of complete graphs without tricolored triangles, *J. Graph Theory* 46 (2004) 211–216.
- [20] M. Hall, C. Magnant, K. Ozeki, M. Tsugaki, Improved upper bounds for Gallai-Ramsey numbers of paths and cycles, *J. Graph Theory* 75 (2014) 59–74.
- [21] M.M.M. Jaradat, M.S. Bataineh, S. Radaideh, Ramsey numbers for theta graphs, *Internat. J. Combin.* (2011) Art. ID 649687, 9pp.
- [22] M. Jenssen, J. Skokan, Exact Ramsey numbers of odd cycles via nonlinear optimisation, [arXiv:1608.05705](https://arxiv.org/abs/1608.05705).
- [23] J. Körner, G. Simonyi, Graph pairs and their entropies: modularity problems, *Combinatorica* 20 (2000) 227–240.
- [24] H. Liu, C. Magnant, A. Saito, I. Schiermeyer, Y. Shi, Gallai-Ramsey number for K_4 , *J. Graph Theory* 94 (2020) 192–205.
- [25] X. Li, L. Wang, Gallai-Ramsey numbers for a class of graphs with five vertices, to appear in *Graphs and Combin.* (2020).
- [26] S. P. Radziszowski, Small Ramsey numbers, *Electron. J. Combin.*, #DS1, revision #15, March 2017.
- [27] Z-X. Song, J. Zhang, A conjecture on Gallai-Ramsey numbers of even cycles and paths, *Australas. J. Combin.* 75 (2019) 296–308.
- [28] Z-X. Song, B. Wei, F. Zhang, Q. Zhao, A note on Gallai-Ramsey number of wheels, *Discrete Math.* 343 (2020) 111725, 5pp.
- [29] F. P. Ramsey, On a problem of formal logic, *Proc. London Math. Soc.* 30 (1930) 264–286.
- [30] Z. Wang, Y. Mao, C. Magnant, I. Schiermeyer, J. Zou, Gallai-Ramsey numbers of odd cycles, [arXiv:1808.09245](https://arxiv.org/abs/1808.09245).
- [31] F. Zhang, Y. Chen, Z-X. Song, Gallai-Ramsey numbers of cycles. [arXiv:1809.00227v1](https://arxiv.org/abs/1809.00227v1).
- [32] F. Zhang, Z-X. Song, Y. Chen, Gallai-Ramsey number of even cycles with chords. [arXiv:1906.05263v3](https://arxiv.org/abs/1906.05263v3).