

The spectrum and automorphism group of the set-inclusion graph

Xueyi Huang^a, Qiongxiang Huang^{b*}, Jianfeng Wang^c

^aSchool of Mathematics and Statistics, Zhengzhou University,
Zhengzhou, Henan 450001, P.R. China

^bCollege of Mathematics and Systems Science, Xinjiang University,
Urumqi, Xinjiang 830046, P.R. China

^cSchool of Mathematics and Statistics, Shandong University of Technology,
Zibo, Shandong 255049, P.R. China

Abstract Let n , k and l be integers with $1 \leq k < l \leq n - 1$. The set-inclusion graph $G(n, k, l)$ is the graph whose vertex set consists of all k - and l -subsets of $[n] = \{1, 2, \dots, n\}$, where two distinct vertices are adjacent if one of them is contained in another. In this paper, we determine the spectrum and automorphism group of $G(n, k, l)$, respectively.

Keywords: set-inclusion graph, spectrum, automorphism group.

AMS Classification: 05C50, 05C25

1 Introduction

Let G be a simple undirected graph with vertex set V and edge set E . As usual, we denote by $\mathcal{L}(G)$ and $A(G)$ the *line graph* and *adjacency matrix* of G , respectively. The *characteristic polynomial* of G is defined as $\phi(G, x) = |x \cdot I - A(G)|$, where I denotes the identity matrix of order $|V|$. The zeros of $\phi(G, x)$ are called the *eigenvalues* of G , and all eigenvalues together with their multiplicities are called the *spectrum* of G . In particular, G is said to be *integral* if its spectrum consists entirely of integers. Also, if G has a bipartition $V = V_1 \cup V_2$ ($|V_i| = n_i$) such that both V_1 and V_2 are independent sets and each vertex of V_1 (resp. V_2) has r_1 (resp. r_2) neighbors in V_2 (resp. V_1), then we say that G is a *semi-regular bipartite graph* with parameters (n_1, n_2, r_1, r_2) .

An *automorphism* of G is a permutation on V that preserves adjacency relations. The *automorphism group* of G , denoted by $\text{Aut}(G)$, is the set of all automorphisms

*Corresponding author.

E-mail address: huangxy@zzu.edu.cn (X. Huang), huangqx@xju.edu.cn (Q. Huang), jfwang@sdut.edu.cn (J. Wang).

of G . The graph G is called *vertex-transitive*, *edge-transitive* and *arc-transitive* if $\text{Aut}(G)$ acts transitively on its vertices, edges and arcs, respectively. As usual, we view each edge of G as a pair of oppositely directed arcs.

For any fixed integers n , k and l with $1 \leq k < l \leq n - 1$, the *set-inclusion graph* $G(n, k, l)$ is defined as the graph with vertex set $V = \{v \mid v \subseteq [n], |v| \in \{k, l\}\}$ and edge set $E = \{\{u, v\} \mid u, v \in V, u \subset v \text{ or } u \supset v\}$ (cf. [1]). It is not difficult to verify that the mapping $\tau : v \mapsto \bar{v} = [n] \setminus v$ builds up an isomorphism between $G(n, k, l)$ and $G(n, n-l, n-k)$. Thus we may always assume that $k+l \leq n$ (thus $k \leq \frac{n-1}{2}$ due to $k < l$). By definition, we see that $G(n, k, l)$ is a connected semi-regular bipartite graph with parameters $((\binom{n}{k}, \binom{n}{l}), (\binom{n-k}{l-k}, \binom{l}{k}))$, where the corresponding bipartition is given by $V = V_1 \cup V_2$ with

$$V_1 = \{v \mid v \subseteq [n], |v| = k\} \text{ and } V_2 = \{v \mid v \subseteq [n], |v| = l\}. \quad (1)$$

Thus the line graph $\mathcal{L}(G(n, k, l))$ is a connected $((\binom{n-k}{l-k} + \binom{l}{k}) - 2)$ -regular graph on $\binom{n-k}{l-k} \binom{n}{k} = \binom{l}{k} \binom{n}{l}$ vertices.

For some special k and l , the set-inclusion graph $G(n, k, l)$ has received particular attention by various researchers. For example, Badakhshian, Katona and Tuza [1] determined the domination number of $G(n, 1, l)$, and gave lower and upper estimates on the domination number of $G(n, 2, l)$. For $l = k + 1$, the graph $G(n, k, k + 1) = B(n, k)$ is just the subgraph of the *hypercube* Q_n induced by all k - and $(k+1)$ -subsets of $[n]$, here Q_n can be viewed as the graph with all subsets of $[n]$ as its vertex set in which two vertices (subsets) are adjacent if their symmetric difference has precisely one element. Particularly, the graph $B(2k + 1, k)$ is known as the *middle layer graph* [17] or the *regular hyper-star graph* [11]. The famous middle levels conjecture, which probably originated with Havel [7] and Buck and Wiedemann [4], and has also been attributed to Dejter, Erdős, Trotter, and various others [9], asserts that $B(2k + 1, k)$ is hamiltonian. In [15], Mütze completely confirmed the conjecture. In addition, the graph $G(n, k, n - k) = H(n, k)$ is known as the *bipartite Kneser graph* [16], which is just the double cover of the Kneser graph (see Section 2 for the definition). It was conjectured independently by Simpson [18] and Roth (see [6, 8]), that $H(n, k)$ (notice that $H(2k + 1, k) = B(2k + 1, k)$) is hamiltonian [18]. Very recently, Mütze and Su [16] confirmed this conjecture.

To study the spectrum and automorphism group of a graph is a basic problem in algebraic graph theory. In this paper, we focus on determining the spectrum and automorphism group of the set-inclusion graph $G(n, k, l)$. First of all, we give the following theorem.

Theorem 1. *Let n , k and l be integers with $1 \leq k < l \leq n - 1$ and $k + l \leq n$. Then the spectrum of $G(n, k, l)$ is given by*

$$\left\{ \left[\pm \sqrt{\beta_s} \right]^{\binom{n}{s} - \binom{n}{s-1}} \mid 0 \leq s \leq k \right\} \cup \left\{ [0]^{\binom{n}{l} - \binom{n}{k}} \right\},$$

where

$$\beta_s = \sum_{i=\max\{2k-l, 0\}}^k \binom{n-2k+i}{l-2k+i} \cdot \left(\sum_{r=0}^s (-1)^{s-r} \binom{s}{r} \binom{k-r}{i-r} \binom{n-k-s+r}{k-i-s+r} \right). \quad (2)$$

As an application, we give the spectra of the line graph $\mathcal{L}(G(n, k, l))$, the graph $B(n, k)$ and its line graph $\mathcal{L}(B(n, k))$, which generalizes a result due to Mirafzal [13] on the eigenvalues of $\mathcal{L}(B(n, 1))$ (see Section 2).

Next we present the automorphism group of $G(n, k, l)$.

Theorem 2. *Let n, k and l be integers with $1 \leq k < l \leq n - 1$ and $k + l \leq n$. Then the automorphism group of $G(n, k, l)$ is*

$$\text{Aut}(G(n, k, l)) \cong \begin{cases} S_n & \text{if } k + l < n, \\ S_n \times \mathbb{Z}_2 & \text{if } k + l = n, \end{cases}$$

where S_n is the symmetric group on $[n]$ and \mathbb{Z}_2 is the cyclic group of order 2.

As corollaries, we obtain the automorphism groups of the graph $B(n, k)$ and the bipartite Kneser graph $H(n, k)$, which have been given by Mirafzal [12, 14] (see Section 3).

2 The spectrum of $G(n, k, l)$

Let n, k and i be integers with $0 \leq i \leq k \leq \frac{n}{2}$. Denote by $J(n, k, i)$ the graph whose vertex set consists of all k -subsets of $[n]$, where two vertices (k -subsets) are adjacent if their intersection has size i (here we regard $J(n, k, k)$ as the graph consisting of $\binom{n}{k}$ isolated vertices with each of them equipping with one loop). In particular, the graphs $J(n, k) = J(n, k, k - 1)$ and $K(n, k) = J(n, k, 0)$ are known as the *Johnson graph* and *Kneser graph*, respectively.

Let A_i be the adjacency matrix of $J(n, k, i)$ for $0 \leq i \leq k$. It is well known that the set of matrices $\{A_i : 0 \leq i \leq k\}$ forms a symmetric association scheme (see Knuth [10], Bannai & Ito [2]), which is called *Johnson scheme*, since these matrices have the properties: $A_k = I$, $\sum_{i=0}^k A_i = J$, $A_i^T = A_i$ and

$$A_i A_j = A_j A_i = \sum_{s=0}^k \left[\sum_{r=0}^s \binom{s}{r} \binom{k-s}{i-r} \binom{k-s}{j-r} \binom{n-2k+s}{k-i-j+r} \right] A_s$$

for $0 \leq i \neq j \leq k$.

The following lemma provides the eigenvalues of $J(n, k, i)$ for each i .

Lemma 1. (Knuth [10]) *Let n, k and i be integers with $0 \leq i \leq k \leq \frac{n}{2}$. Then the spectrum of $J(n, k, i)$ is given by*

$$\left\{ [\alpha_{i,s}] \binom{n}{s} - \binom{n}{s-1} \mid 0 \leq s \leq k \right\},$$

where

$$\alpha_{i,s} = \sum_{r=0}^s (-1)^{s-r} \binom{s}{r} \binom{k-r}{i-r} \binom{n-k-s+r}{k-i-s+r}. \quad (3)$$

Moreover, for any fixed s , there exist $\left(\binom{n}{s} - \binom{n}{s-1} \right)$'s linearly independent vectors which are the common eigenvectors of A_i 's (here A_i is the adjacency matrix of $J(n, k, i)$) with respect to the eigenvalue $\alpha_{i,s}$ for $0 \leq i \leq k$.

Using Lemma 1, we now give the proof of Theorem 1.

Proof of Theorem 1. Since $G(n, k, l)$ is bipartite, its adjacency matrix can be written as

$$A = \begin{bmatrix} 0 & B \\ B^T & 0 \end{bmatrix} \begin{matrix} V_1 \\ V_2 \end{matrix},$$

where V_1 and V_2 are given in (1). Note that $|V_1| = \binom{n}{k}$ and $|V_2| = \binom{n}{l}$. We have

$$|xI - A^2| = |xI - BB^T| \cdot |xI - B^TB| = x^{\binom{n}{l} - \binom{n}{k}} \cdot |xI - BB^T|^2, \quad (4)$$

where $\binom{n}{k} \leq \binom{n}{l}$ due to $k < l$ and $k + l \leq n$. Note that the spectrum of $G(n, k, l)$ is symmetric about 0. If $\xi_1, \xi_2, \dots, \xi_{\binom{n}{k}}$ are all eigenvalues of BB^T , then from (4) we can deduce that the eigenvalues of $G(n, k, l)$ are $\pm\sqrt{\xi_1}, \pm\sqrt{\xi_2}, \dots, \pm\sqrt{\xi_{\binom{n}{k}}}$ and $(\binom{n}{l} - \binom{n}{k})$'s 0. Thus it suffices to determine the eigenvalues of BB^T .

Recall that V_1 and V_2 consist of all k - and l -subsets of $[n]$, respectively. For any two vertices $u, v \in V_1$ (not necessarily distinct), we see that $(BB^T)_{u,v}$ is just the number of common neighbors of u, v in V_2 . Then, according to the definition of $G(n, k, l)$, we obtain

$$(BB^T)_{u,v} = \begin{cases} \binom{n-2k+i}{l-2k+i} & \text{if } |u \cap v| = i \text{ with } \max\{2k-l, 0\} \leq i \leq k, \\ 0 & \text{otherwise,} \end{cases}$$

or equivalently,

$$BB^T = \sum_{i=\max\{2k-l, 0\}}^k \binom{n-2k+i}{l-2k+i} \cdot A_i,$$

where A_i is the adjacency matrix of $J(n, k, i)$. Therefore, by Lemma 1, each eigenvalue of BB^T is of the form

$$\beta_s = \sum_{i=\max\{2k-l, 0\}}^k \binom{n-2k+i}{l-2k+i} \cdot \alpha_{i,s}$$

with multiplicity $\binom{n}{s} - \binom{n}{s-1}$, where $\alpha_{i,s}$ is given in (3) and $0 \leq s \leq k$. Thus we obtain the spectrum of $G(n, k, l)$ immediately. \square

Remark 1. Since $G(n, k, l)$ is a semi-regular bipartite graph with parameters $(\binom{n}{k}, \binom{n}{l}, \binom{n-k}{l-k}, \binom{l}{k})$, its largest eigenvalue must be $\sqrt{\binom{n-k}{l-k} \binom{l}{k}}$. We see that this

eigenvalue is exactly $\sqrt{\beta_0}$ in Theorem 1. In fact, from (2) we obtain

$$\begin{aligned}
\beta_0 &= \sum_{i=\max\{2k-l,0\}}^k \binom{n-2k+i}{l-2k+i} \binom{k}{i} \binom{n-k}{k-i} \\
&= \sum_{i=\max\{2k-l,0\}}^k \frac{(n-k)!}{(l-2k+i)!(n-l)!(k-i)!} \binom{k}{i} \\
&= \sum_{i=\max\{2k-l,0\}}^k \frac{(n-k)!}{(n-l)!(l-k)!} \cdot \frac{(l-k)!}{(l-2k+i)!(k-i)!} \binom{k}{i} \\
&= \binom{n-k}{l-k} \sum_{i=\max\{2k-l,0\}}^k \binom{l-k}{k-i} \binom{k}{i} \\
&= \binom{n-k}{l-k} \binom{l}{k},
\end{aligned}$$

as required.

The following lemma gives the characteristic polynomial of the line graph of a semi-regular bipartite graph.

Lemma 2. (Cvetković, Rowlinson and Simić [5, Corollary 2.4.3]) *If G is a semi-regular bipartite graph with parameters (n_1, n_2, r_1, r_2) ($n_1 \leq n_2$), and if $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_{n_1}$ are the first n_1 largest eigenvalues, then*

$$\begin{aligned}
\phi(\mathcal{L}(G), x) &= (x - r_1 - r_2 + 2)(x - r_2 + 2)^{n_2 - n_1} (x + 2)^{(n_2 r_2 - n_1 - n_2 + 1)} \\
&\quad \cdot \prod_{i=2}^{n_1} ((x - r_1 + 2)(x - r_2 + 2) - \lambda_i^2).
\end{aligned}$$

Corollary 1. *Let n, k and l be integers with $1 \leq k < l \leq n - 1$ and $k + l \leq n$. Then the spectrum of the line graph $\mathcal{L}(G(n, k, l))$ is given by*

$$\begin{aligned}
&\left\{ \left[\binom{n-k}{l-k} + \binom{l}{k} - 2 \right]^1, \left[\binom{l}{k} - 2 \right]^{\binom{n}{l} - \binom{n}{k}}, [-2]^{\binom{n}{l} \binom{l}{k} - \binom{n}{k} - \binom{n}{l} + 1} \right\} \\
&\cup \left\{ [\gamma_{s,1}]^{\binom{n}{s} - \binom{n}{s-1}}, [\gamma_{s,2}]^{\binom{n}{s} - \binom{n}{s-1}} \mid 1 \leq s \leq k \right\},
\end{aligned}$$

where $\gamma_{s,1}, \gamma_{s,2}$ are the two roots of $(x - \binom{n-k}{l-k} + 2) \cdot (x - \binom{l}{k} + 2) - \beta_s = 0$ with β_s shown in (2).

Proof. Recall that $G(n, k, l)$ is a connected semi-regular bipartite graph with parameters $(\binom{n}{k}, \binom{n}{l}, \binom{n-k}{l-k}, \binom{l}{k})$, where $\binom{n}{k} \leq \binom{n}{l}$ due to $k < l$ and $k + l \leq n$. By Theorem 1, the first $\binom{n}{k}$ largest eigenvalues of $G(n, k, l)$ are $\sqrt{\beta_s}$ (with multiplicity $\binom{n}{s} - \binom{n}{s-1}$), $s = 0, 1, \dots, k$. In particular, the largest eigenvalue is $\sqrt{\beta_0} = \sqrt{\binom{n-k}{l-k} \binom{l}{k}}$

according to Remark 1. By Lemma 2, $\mathcal{L}(G(n, k, l))$ has characteristic polynomial

$$\begin{aligned} & \phi(\mathcal{L}(G(n, k, l)), x) \\ &= \left(x - \binom{n-k}{l-k} - \binom{l}{k} + 2 \right) \cdot \left(x - \binom{l}{k} + 2 \right)^{\binom{n}{l} - \binom{n}{k}} \cdot (x+2)^{\binom{n}{l} \binom{l}{k} - \binom{n}{k} - \binom{n}{l} + 1} \\ & \quad \cdot \prod_{s=1}^k \left(\left(x - \binom{n-k}{l-k} + 2 \right) \cdot \left(x - \binom{l}{k} + 2 \right) - \beta_s \right)^{\binom{n}{s} - \binom{n}{s-1}}, \end{aligned}$$

and our result follows. \square

By Theorem 1 and Corollary 1, we can deduce the spectra of the graph $B(n, k)$ and its line graph $\mathcal{L}(B(n, k))$, respectively.

Corollary 2. *Let n and k be integers with $1 \leq k \leq \frac{n-1}{2}$. Then the spectrum of $B(n, k)$ is*

$$\left\{ \left[\pm \sqrt{(n-k-s)(k+1-s)} \right]^{\binom{n}{s} - \binom{n}{s-1}} \mid 0 \leq s \leq k \right\} \cup \left\{ [0]^{\binom{n}{k+1} - \binom{n}{k}} \right\}.$$

Proof. As $B(n, k) = G(n, k, k+1)$, by setting $l = k+1$ in (2), we have

$$\begin{aligned} \beta_s &= \sum_{i=k-1}^k \binom{n-2k+i}{i-k+1} \cdot \left(\sum_{r=0}^s (-1)^{s-r} \binom{s}{r} \binom{k-r}{i-r} \binom{n-k-s+r}{k-i-s+r} \right) \\ &= \sum_{r=0}^s (-1)^{s-r} \binom{s}{r} \binom{k-r}{k-r-1} \binom{n-k-s+r}{1-s+r} + \\ & \quad (n-k) \cdot \sum_{r=0}^s (-1)^{s-r} \binom{s}{r} \binom{k-r}{k-r} \binom{n-k-s+r}{-s+r} \\ &= (n-k-s)(k+1-s), \end{aligned} \tag{5}$$

and the result follows from Theorem 1 immediately. \square

If $l = k+1$, then $(x - \binom{n-k}{l-k} + 2) \cdot (x - \binom{l}{k} + 2) - \beta_s = (x-s+2)(x-n+s+1)$ according to (5). By Theorem 1, we obtain the following result immediately.

Corollary 3. *Let n and k be integers with $1 \leq k \leq \frac{n-1}{2}$. Then the line graph $\mathcal{L}(B(n, k))$ is an integral graph with spectrum*

$$\begin{aligned} & \left\{ [n-1]^1, [k-1]^{\binom{n}{k+1} - \binom{n}{k}}, [-2]^{k \binom{n}{k+1} - \binom{n}{k} + 1} \right\} \\ & \cup \left\{ [s-2]^{\binom{n}{s} - \binom{n}{s-1}}, [n-s-1]^{\binom{n}{s} - \binom{n}{s-1}} \mid 1 \leq s \leq k \right\}. \end{aligned}$$

Putting $k = 1$ in Corollary 3, we obtain the main result of Mirafzal [13].

Corollary 4 (Mirafzal [13]). *Let $n \geq 4$ be an integer. Then the graph $\mathcal{L}(B(n, 1))$ is an integral graph with distinct eigenvalues $-2, -1, 0, n-2, n-1$.*

3 The automorphism group of $G(n, k, l)$

Let S_n be the symmetric group on $[n] = \{1, 2, \dots, n\}$ with $n \geq 3$. For each i with $1 \leq i \leq n-1$, we denote by N_i the set of all i -subsets of $[n]$. Suppose

$$N = \cup_{i \in I} N_i,$$

where I is some nonempty subset of $\{1, 2, \dots, n-1\}$. For any fixed $g \in S_n$, we define σ_g as the mapping

$$\begin{aligned} \sigma_g : N &\rightarrow N \\ v &\mapsto v^g = \{x^g \mid x \in v\}, \text{ for } v \in N. \end{aligned} \quad (6)$$

Clearly, σ_g is one-to-one and onto. Denote by

$$\tilde{S}_n = \{\sigma_g \mid g \in S_n\}. \quad (7)$$

It is not hard to verify that \tilde{S}_n is a permutation group on N which is isomorphic to S_n . If the set $N = \cup_{i \in I} N_i$ satisfies $I = n - I = \{n - i \mid i \in I\}$, we define

$$\begin{aligned} \tau : N &\rightarrow N \\ v &\mapsto \bar{v} = [n] \setminus v, \text{ for } v \in N. \end{aligned} \quad (8)$$

It is clear that τ is a permutation of order 2 on N , and $\tau \notin \tilde{S}_n$ due to $n \geq 3$.

By above arguments, we see that \tilde{S}_n is a permutation group on the vertex set $V = V_1 \cup V_2 = N_k \cup N_l$ of the set-inclusion graph $G(n, k, l)$, where $1 \leq k < l \leq n-1$ and $k+l \leq n$. Let $\sigma_g \in \tilde{S}_n$. For any two vertices $u, v \in V$, we have

$$\begin{aligned} \{u, v\} \in E &\iff u \subset v \text{ or } v \subset u \\ &\iff \sigma_g(u) = u^g \subset v^g = \sigma_g(v) \text{ or } \sigma_g(v) = v^g \subset u^g = \sigma_g(u) \\ &\iff \{\sigma_g(u), \sigma_g(v)\} \in E, \end{aligned}$$

which implies that σ_g is an automorphism of $G(n, k, l)$. Thus we conclude that

$$\tilde{S}_n \leq \text{Aut}(G(n, k, l)).$$

Furthermore, if $l = n - k$, then $G(n, k, n - k)$ has vertex set $V = V_1 \cup V_2 = N_k \cup N_{n-k}$, and so the mapping τ defined in (8) is a permutation on V . For any two vertices $u, v \in V$, we obtain

$$\begin{aligned} \{u, v\} \in E &\iff u \subset v \text{ or } v \subset u \\ &\iff \tau(u) = \bar{u} \supset \bar{v} = \tau(v) \text{ or } \tau(v) = \bar{v} \supset \bar{u} = \tau(u) \\ &\iff \{\tau(u), \tau(v)\} \in E, \end{aligned}$$

implying that τ is also an automorphism of $G(n, k, n - k)$. As $1 \leq k < l = n - k$, we have $n \geq 3$, and $\tau \notin \tilde{S}_n$. Additionally, one can check that $\tau \cdot \sigma_g = \sigma_g \cdot \tau$ for all $\sigma_g \in \tilde{S}_n$. Therefore, we have

$$\tilde{S}_n \times \langle \tau \rangle \leq \text{Aut}(G(n, k, n - k)).$$

The result of Theorem 2 suggests that the two subgroups mentioned above are in fact the full automorphism group of $G(n, k, l)$ for $l < n - k$ and $l = n - k$, respectively. Before giving the proof, we need some powerful lemmas.

Recall that the Johnson graph $J(n, k) = J(n, k, k - 1)$ ($1 \leq k \leq \frac{n}{2}$) is the graph whose vertex set consists of all k -subsets of $[n]$, where two vertices are adjacent if their intersection has size $k - 1$ (see Section 2). As above, we see that $\text{Aut}(J(n, k))$ contains \tilde{S}_n as its subgroup, where \tilde{S}_n is given in (7). Indeed, we have

Lemma 3. (Brouwer, Cohen and Neumaier [3, Theorem 9.1.2]) *Let n and k be integers with $1 \leq k \leq \frac{n}{2}$. Then the automorphism group of the Johnson graph $J(n, k)$ is*

$$\text{Aut}(J(n, k)) = \begin{cases} \tilde{S}_n \times \langle \tau \rangle \cong S_n \times \mathbb{Z}_2 & \text{if } k = \frac{n}{2} \geq 2, \\ \tilde{S}_n \cong S_n & \text{otherwise,} \end{cases}$$

where \tilde{S}_n and τ are given in (7) and (8), respectively.

The next lemma is straightforward.

Lemma 4. *Let G be a connected bipartite graph with bipartition $V = V_1 \cup V_2$. For each $\sigma \in \text{Aut}(G)$, we have either $\sigma(V_1) = V_1$ and $\sigma(V_2) = V_2$, or $\sigma(V_1) = V_2$ and $\sigma(V_2) = V_1$.*

Lemma 5. *Let n , k and l be integers with $1 \leq k < l \leq n - 1$ and $k + l \leq n$, and let $V = V_1 \cup V_2$ be the bipartition of $G(n, k, l)$ with V_1 and V_2 shown in (1). If σ is an automorphism of $G(n, k, l)$ that fixes each vertex of V_1 , then σ is the identity automorphism.*

Proof. Let $v = \{x_1, x_2, \dots, x_l\}$ be an arbitrary vertex of V_2 . Set $u_1 = \{x_1, x_2, \dots, x_k\}$, $u_2 = \{x_2, \dots, x_{k+1}\}, \dots, u_{l-k+1} = \{x_{l-k+1}, \dots, x_l\}$. Clearly, $u_1, u_2, \dots, u_{l-k+1} \in V_1$. By the definition of $G(n, k, l)$, we see that v is the unique common neighbor of $u_1, u_2, \dots, u_{l-k+1}$. Thus we must have $\sigma(v) = v$ because $u_1, u_2, \dots, u_{l-k+1}$ have been fixed by σ according to the assumption. By the arbitrariness of v , the result follows immediately. \square

Lemma 6. *Let n , k and l be integers with $1 \leq k < l \leq n - 1$ and $k + l \leq n$, and let $V = V_1 \cup V_2$ be the bipartition of $G(n, k, l)$ with V_1 and V_2 shown in (1). Then for each i with $\max\{2k - l, 0\} \leq i \leq k - 1$ and for any two vertices $u, v \in V_1$, we have $|u \cap v| = i$ if and only if $|N(u) \cap N(v)| = \binom{n-2k+i}{l-2k+i}$.*

Proof. As $u, v \in V_1$, we have $|u| = |v| = k$. For $\max\{2k - l, 0\} \leq i \leq k - 1$, if $|u \cap v| = i$ then we have $|u \cup v| = 2k - i$, so u, v are simultaneously contained in $\binom{n-2k+i}{l-2k+i}$'s l -subsets of $[n]$, that is, $|N(u) \cap N(v)| = \binom{n-2k+i}{l-2k+i}$. This proves the necessity. Also, since $l \leq n - 1$, for any i, j with $\max\{2k - l, 0\} \leq i \neq j \leq k - 1$, we can verify that $\binom{n-2k+i}{l-2k+i} \neq \binom{n-2k+j}{l-2k+j}$. Thus the sufficiency follows. \square

Now we give the proof of Theorem 2.

Proof of Theorem 2. At the beginning of this section, we have mentioned that \tilde{S}_n and $\tilde{S}_n \times \langle \tau \rangle$ are the subgroups of $\text{Aut}(G(n, k, l))$ for $k + l < n$ and $k + l = n$, respectively.

Let $V = V_1 \cup V_2 = N_k \cup N_l$ be the bipartition of $G(n, k, l)$. Let σ be an arbitrary automorphism of $G(n, k, l)$. By Lemma 4, we only need to consider the following two cases.

Case 1. $\sigma(V_1) = V_1$ and $\sigma(V_2) = V_2$.

In this situation, let $\sigma|_{V_1}$ be the restriction of σ on V_1 . Clearly, $\sigma|_{V_1}$ is a permutation on $V_1 = N_k$. We have the following claim.

Claim 1. $\sigma|_{V_1} \in \text{Aut}(J(n, k)) = \tilde{S}_n$.

Recall that the Johnson graph $J(n, k)$ has $V_1 = N_k = \{v \mid v \subseteq [n], |v| = k\}$ as its vertex set, where $u, v \in V_1$ are adjacent if and only if $|u \cap v| = k - 1$. For any $u, v \in V_1$, by Lemma 6, we obtain

$$\begin{aligned} |u \cap v| = k - 1 &\iff |N(u) \cap N(v)| = \binom{n - k - 1}{l - k - 1} \\ &\iff |N(\sigma|_{V_1}(u)) \cap N(\sigma|_{V_1}(v))| = \binom{n - k - 1}{l - k - 1} \\ &\iff |\sigma|_{V_1}(u) \cap \sigma|_{V_1}(v)| = k - 1, \end{aligned}$$

which implies that $\sigma|_{V_1} \in \text{Aut}(J(n, k))$. Also, by Lemma 3, we have $\text{Aut}(J(n, k)) = \tilde{S}_n$ because $1 \leq k \leq \frac{n-1}{2}$. This proves Claim 1.

By Claim 1, there exists some $g \in S_n$ such that $\sigma|_{V_1} = \sigma_g$, where σ_g is the permutation on $V_1 = N_k$ defined in (6). Using the same g , we can extend σ_g as a permutation on $V = V_1 \cup V_2$ again by (6), and we still denote this permutation by σ_g . Clearly, $\sigma_g \in \tilde{S}_n \leq \text{Aut}(G(n, k, l))$. Then we see that $\sigma_g^{-1}\sigma$ fixes each vertex of V_1 , and so must be the identity automorphism by Lemma 5. Thus $\sigma = \sigma_g \in \tilde{S}_n$.

Case 2. $\sigma(V_1) = V_2$ and $\sigma(V_2) = V_1$.

In this situation, we must have $|V_1| = |V_2|$, and so $k + l = n$. Let τ be the automorphism of $G(n, k, n - k)$ defined in (8). We see that $\sigma \cdot \tau \in \text{Aut}(G(n, k, n - k))$, and $\sigma \cdot \tau(V_i) = V_i$ for $i = 1, 2$. According to what was proved in Case 1, there exists some $g \in \tilde{S}_n$ such that $\sigma \cdot \tau = \sigma_g$, i.e., $\sigma = \sigma_g \cdot \tau$. Therefore, we conclude that $\sigma \in \tilde{S}_n \times \langle \tau \rangle$.

If $k + l < n$, only Case 1 can occur, thus we have $\text{Aut}(G(n, k, l)) \leq \tilde{S}_n \cong S_n$. If $k + l = n$, both Case 1 and Case 2 can occur, and so $\text{Aut}(G(n, k, n - k)) \leq \tilde{S}_n \times \langle \tau \rangle \cong S_n \times \mathbb{Z}_2$, where \mathbb{Z}_2 is the cyclic group of order 2.

We complete the proof. \square

Remark 2. According to Theorem 2, it is easy to see that the set-inclusion graph $G(n, k, l)$ is edge-transitive. Furthermore, for $l = n - k$ we claim that $G(n, k, n - k)$ is arc-transitive, since it is obviously vertex-transitive, and the stabilizer of any given vertex v acts transitively on the set of neighbors.

Recall that $B(n, k) = G(n, k, k + 1)$, and $H(n, k) = G(n, k, n - k)$. Putting $l = k + 1$ and $n - k$ in Theorem 2, we obtain the following two corollaries immediately.

Corollary 5. (Mirafzal [14]) *Let n and k be integers with $1 \leq k \leq \frac{n-1}{2}$. Then the automorphism group of $B(n, k)$ is*

$$\text{Aut}(B(n, k)) \cong \begin{cases} S_n & \text{if } k < \frac{n-1}{2}, \\ S_n \times \mathbb{Z}_2 & \text{if } k = \frac{n-1}{2}, \end{cases}$$

where S_n is the symmetric group on $[n]$ and \mathbb{Z}_2 is the cycle group of order 2.

Corollary 6. (Mirafzal [13]) *Let n and k be integers with $1 \leq k \leq \frac{n-1}{2}$. Then the automorphism group of $H(n, k)$ is*

$$\text{Aut}(H(n, k)) \cong S_n \times \mathbb{Z}_2,$$

where S_n is the symmetric group on $[n]$ and \mathbb{Z}_2 is the cycle group of order 2.

Acknowledgements. The first author is supported by the China Postdoctoral Science Foundation and the Postdoctoral Research Sponsorship in Henan Province (No. 1902011), the second author is supported by the National Natural Science Foundation of China (No. 11671344), and the third author is supported by the National Natural Science Foundation of China (No. 11461054).

References

- [1] L. Badakhshian, G.O.H. Katona, Z. Tuza, The domination number of the graph defined by two levels of the n -cube, *Discrete Appl. Math.*, 2019, <https://doi.org/10.1016/j.dam.2019.02.006>.
- [2] E. Bannai, T. Ito, *Algebraic Combinatorics I: Association Schemes*, Benjamin/Cummings, Menlo Park, California, 1984.
- [3] A.E. Brouwer, A.M. Cohen, A. Neumaier, *Distance-Regular Graphs*, Springer-Verlag, Berlin, 1989.
- [4] M. Buck, D. Wiedemann, Gray codes with restricted density, *Discrete Math.* 48 (2-3) (1984) 163–171.
- [5] D. Cvetković, P. Rowlinson, S.K. Simić, *An Introduction to the Theory of Graph Spectra*, Cambridge University Press, Cambridge, 2010.
- [6] R. Gould, Updating the hamiltonian problem – a survey, *J. Graph Theory* 15 (1991) 121–157.
- [7] I. Havel, Semipaths in directed cubes, in: M. Fiedler (Ed.), *Graphs and other Combinatorial Topics*, Teubner-Texte Math., Teubner, Leipzig, 1983, pp. 101–108.
- [8] G. Hurlbert, The antipodal layers problem, *Discrete Math.* 128 (1994) 237–245.
- [9] H. Kierstead, W. Trotter, Explicit matchings in the middle levels of the Boolean lattice, *Order* 5 (2) (1988) 163–171.
- [10] D.E. Knuth, *Combinatorial Matrices*, in: *Selected Papers on Discrete Mathematics*, CSLI lecture Notes 106, CSLI Publications, Stanford, CA, 2003, pp. 177–186.

- [11] L. Liptak, E. Cheng, J.S. Kim, S.W. Kim, One-to-many node-disjoint paths of hyper-star networks, *Discrete Appl. Math.* 160 (2012) 2006–2014.
- [12] S.M. Mirafzal, Cayley properties of the line graphs induced by consecutive layers of the hypercube, arXiv:1711.02701 [math.GR].
- [13] S.M. Mirafzal, A new class of integral graphs constructed from the hypercube, *Linear Algebra Appl.* 558 (2018) 186–194.
- [14] S.M. Mirafzal, The automorphism group of the bipartite Kneser graph, *Proc. Indian Acad. Sci. (Math. Sci.)* 129 (2019) #34.
- [15] T. Mütze, Proof of the middle levels conjecture, *Proc. Lond. Math. Soc.* 112 (2016) 677–713.
- [16] T. Mütze, P. Su, Bipartite Kneser graphs are Hamiltonian, *Combinatorica* 37 (6) (2017) 1207–1219.
- [17] T. Mütze, F. Weber, Construction of 2-factors in the middle layer of the discrete cube, *J. Combin. Theory Ser. A* 119 (8) (2012) 1832–1855.
- [18] J. Simpson, Hamiltonian bipartite graphs, in: *Proceedings of the Twenty-second Southeastern Conference on Combinatorics, Graph Theory, and Computing* (Baton Rouge, LA, 1991), volume 85, 97–110, 1991.