

Semiparametric model averaging for high dimensional conditional quantile prediction

Jingwen Tu^a, Hu Yang^a, Chaohui Guo^{b*}

^aCollege of Mathematics and Statistics, Chongqing University, Chongqing, 401331, China

^bCollege of Mathematics Science, Chongqing Normal University, Chongqing, 401331, China

Abstract

In this article, we propose a penalized high dimensional semiparametric model average quantile prediction approach that is robust for forecasting the conditional quantile of the response. We consider a two-step estimation procedure. In the first step, we use a local linear regression approach to estimate the individual marginal quantile functions, and approximate the conditional quantile of the response by an affine combination of one-dimensional marginal quantile regression functions. In the second step, based on the nonparametric kernel estimates of the marginal quantile regression functions, we utilize a penalized method to estimate the suitable model weights vector involved in the approximation. The objective of the second step is to select significant variables whose marginal quantile functions make a significant contribution to estimating the joint multivariate conditional quantile function. Under some mild conditions, we have established the asymptotic properties of the proposed robust estimator. Finally, simulations and a real data analysis have been used to illustrate the proposed method.

Keywords: High dimensional data, Kernel estimation, Model averaging, Penalized quantile regression, Prediction accuracy, Semiparametric models

1. Introduction

In many practical situations, especially for economic and medical fields, forecasting and predictive inference are our main goals. In practice, we often face a large number of predictors and uncertain functional forms when making statistical prediction. A popular approach to solve this problem is to consider the model selection tool that can select a optimal model from all candidate models, but we have to recognize that model selection tech-

*Corresponding author (Email:guochaohui2010@126.com)

nique yields only one final model, so useful information may be ignored when significant variables absent from the final model. This may result in misleading predictive outcomes. Instead of depending on only one best model, an alternative method, called model averaging technique, aims to improve the prediction accuracy through giving higher weights to the better marginal models. Thus, model averaging can be regarded as a smoothed extension of model selection and generally leads to a lower risk than model selection. Earlier development for model average was linked closely the Bayesian statistics including Hoeting et al. (1999), Raftery et al. (1997) and Hjort and Claeskens (2003). Recently, various strategies have been developed to construct optimal model averaging weights for frequentist models. For example, Hansen (2007) proposed a frequentist model average approach with weights selected by minimizing a Mallows criterion. Wan et al. (2010) focused on two assumptions of Hansen (2007) and provided a stronger theoretical basis for the use of the Mallows criterion in model averaging. Liang et al. (2011) considered a new procedure of weight choice by minimizing frequentist model average estimators' mean squared errors. To deal with heteroscedastic data, Hansen and Racine (2012) developed a jackknife model averaging approach to choose weights by minimizing a leave-one-out cross-validation criterion and had proved that the proposed approach achieved the lowest possible asymptotic squared error. Zhang et al. (2013) further extended the method of Hansen and Racine (2012) to general models with a non-diagonal error covariance structure or lagged dependent variables. In the framework of linear mixed-effects models, Zhang et al. (2014) constructed an unbiased estimator of the squared risk for the model averaging, which has been demonstrated to be asymptotically optimal in theory under some regularity conditions. Zhang et al. (2016) studied optimal model averaging methods for generalized linear models and generalized linear mixed-effects models, which can be taken as an extension of Zhang et al. (2014)'s. Under the local asymptotic framework, Liu et al. (2015) studied the limiting distributions of least squares averaging estimators and proposed a plug-in averaging estimator by minimizing the sample asymptotic mean squared error. Other related literature can refer to Hansen (2008), Claeskens and Hjort (2008), Zhang et al. (2012), Cheng and Hansen (2015).

Almost all mentioned above research work focus on averaging a set of parametric mod-

els by assuming some parametrically linear or nonlinear relationships between the response and predictors. Although parametric models are easy to understand and widely accepted by scientific researchers, they make strong assumptions in practical applications, which may increase the risk of bias prediction. In contrast, nonparametric models with less structural restriction may provide more flexible predictive inference. Recently, Li et al. (2015) firstly proposed a nonparametric model averaging approach which is more flexible than traditional parametric averaging method. They estimated the multivariate conditional mean regression function by averaging a set of estimated marginal mean regression functions with proper weights obtained by minimizing least squares loss. Motivated by the nonparametric model averaging technique, Chen et al. (2016) studied the semiparametric dynamic portfolio choice and utilized a novel data-driven method to estimate the nonparametric optimal portfolio choice. Huang and Li (2018) extended the method of Li et al. (2015) to panel data and established the asymptotic results of the proposed procedure. Li et al. (2018) approximated the conditional mean regression function by a weighted average of varying coefficient regression functions, which can handle discrete and continuous predictors.

In recent years, we often encounter datasets with a very large number of potential predictors, but only a minority of predictors are truly relevant in prediction. However, most of literature focus on the determination of weights for individual models under a fixed number of covariates. So far, Ando and Li (2014) proposed a two-step model averaging procedure to predict the conditional mean of the response for a ultra-high dimensional linear regression. In order to obtain more accurate prediction of the conditional mean of the response for ultra-high dimensional time series, Chen et al. (2018) introduced a two-step semiparametric procedure that includes the kernel sure independence screening technique and the semiparametric penalized method of model averaging marginal regression. All mentioned above references aim to forecast the conditional mean of the response, but sometimes we are more interested in predicting the conditional quantile of the response. Compared to mean regression, quantile regression not only provides a more complete description of the entire response distribution but also does not require specification of the error distribution, and thus it is more robust.

In this paper, we aim to develop a new semiparametric model averaging procedure for achieving more accurate prediction for the true conditional quantile of the response under the high dimensional setting. This paper may have several innovations as follows: (1) our objective is to predict the conditional quantile of the response rather than its conditional mean. Thus we may encounter more challenge to establish asymptotic theories of model weights since we cannot obtain the closed-form expression of model weights; (2) the proposed approach can offer a complete prediction for the response when different quantiles are adopted; (3) our method produces more accurate in-sample and out-of-sample prediction when non-normal errors are considered.

The rest of the paper is organized as follows. In Section 2, we first give the approximation of the conditional quantile function of the response. Then a two-step semiparametric model averaging approach is applied to estimate the conditional quantile function of the response. In Section 3, we establish the asymptotic theory for the proposed estimator. In Section 4, numerical studies including simulation studies and a real data analysis are carried out to investigate the finite sample performance of the proposed method. Some discussions are reported in Section 5. Finally, all technical proofs are given in the Appendix.

2. Model approximation and estimation method

Let $\{(\mathbf{X}_i, Y_i), 1 \leq i \leq n\}$ be independent and identically distributed observations from (\mathbf{X}, Y) , where $\mathbf{X} = (X_1, \dots, X_{p_n})$ is a p_n -vector of predictors and Y is the response variable. The goal of this paper is to develop a new procedure for forecasting the τ th conditional quantile function of Y given \mathbf{X} , namely, $m_\tau(\mathbf{X}) \equiv Q_\tau(Y|\mathbf{X})$. If the dimension of \mathbf{X} is high, it is not practical to model conditional quantile function $m_\tau(\mathbf{X})$ without any structure assumption due to the curse of dimensionality. Recently, authors approximated the quantile function $m_\tau(\mathbf{X})$ by semiparametric models such as quantile additive models (Horowitz and Lee, 2005, Lv et al., 2017), quantile varying coefficient models (Tang et al., 2013) and among others. However, using a specified model with fixed model structure may increase the risk of model misspecification, which results in poor predictive performance. Therefore, we adopt the model averaging technique to predict $m_\tau(\mathbf{X})$. Specifically, motivated by Li et al. (2015), we model or approximate $m_\tau(\mathbf{X})$ by an affine combination

of one-dimensional nonparametric functions $m_\tau(\mathbf{X}) = w_0 + \sum_{j=1}^{p_n} w_j m_{\tau_j}(X_j)$, where $m_{\tau_j}(X_j) = Q_{\tau_j}(Y|X_j)$ is the τ th conditional quantile of Y given X_j . Here, each marginal regression $m_{\tau_j}(\cdot)$ can be regarded as a candidate model and w_j is the corresponding model weight coefficient. In the rest of the article, we omit τ_j and τ from $m_{\tau_j}(\cdot)$ and $m_\tau(\mathbf{X})$ for notational simplicity, but it is helpful to bear in mind that these quantities are τ_j and τ -specific.

What we are most interested in is to accurately estimate m_j and the model average weight vector $\mathbf{w} = (w_0, w_1, \dots, w_{p_n})^T$. We consider a two-step estimation procedure. In the first step, we employ local linear regression technique to estimate the individual marginal regression functions $m_j(\cdot)$, $j = 1, \dots, p_n$. Specifically, considering a Taylor expansion, we have

$$m_j(X_{ij}) \approx m_j(x) + \dot{m}_j(x)(X_{ij} - x) \equiv a + b(X_{ij} - x), i = 1, \dots, n, j = 1, \dots, p_n,$$

where $\dot{m}_j(x)$ is the first-order derivative of $m_j(x)$. Let $\rho_\tau(u) = \tau u - uI(u < 0)$ be check loss function at τ quantile. Then, we estimate m_j by minimizing the following local weighted quantile loss

$$\sum_{i=1}^n \rho_{\tau_j} \{Y_i - a - b(X_{ij} - x)\} K\left(\frac{X_{ij} - x}{h_j}\right), \quad (1)$$

where $K(\cdot)$ is a kernel function and h_j is a bandwidth. Let (\hat{a}, \hat{b}) be the minimizer of the objective function (1). Then, we have $\hat{m}_j = \hat{a}$.

In the second step, let $\mathbf{w}_o = (w_{o0}, w_{o1}, \dots, w_{op_n})^T$ be the optimal values of the weights in the model averaging defined in Li et al. (2015). To estimate \mathbf{w}_o , we minimize the following function with respect to $\mathbf{w}_n = (w_0, \mathbf{w}^T)^T$ with $\mathbf{w} = (w_1, \dots, w_{p_n})^T$,

$$\mathcal{Q}_n(\mathbf{w}_n) = \sum_{i=1}^n \rho_\tau \left\{ Y_i - w_0 - \sum_{j=1}^{p_n} \hat{m}_j(X_{ij}) w_j \right\} + n \sum_{j=1}^{p_n} p_\lambda(|w_j|), \quad (2)$$

where $p_\lambda(\cdot)$ is a penalty function with a tuning parameter λ , such as SCAD penalty

function, $\dot{p}_\lambda(\cdot)$ is its first order derivative, defined by

$$\dot{p}_\lambda(x) = \lambda \left\{ I(x \leq \lambda) + \frac{(a\lambda - x)_+}{(a-1)\lambda} I(x > \lambda) \right\},$$

where $a > 2$, $p_\lambda(0) = 0$ and λ is a nonnegative penalty parameter which governs sparsity of the model. It is easy to find that $\dot{p}_\lambda(|x|)$ is close to zero if $|x|$ is large.

The estimator of the optimal weights \mathbf{w}_o can be obtained through minimizing the objective function (2), that is, $\hat{\mathbf{w}}_n = \arg \min_{\mathbf{w}_n} \mathcal{Q}_n(\mathbf{w}_n)$. This paper uses the R package ‘‘rqPen’’ to obtain the estimator $\hat{\mathbf{w}}_n$. Finally, for a future observation $\mathbf{x} = (x_1, \dots, x_{p_n})$, we can predict $m(\mathbf{x})$ by $\hat{m}(\mathbf{x}) = \hat{w}_0 + \sum_{j=1}^{p_n} \hat{w}_j \hat{m}_j(x_j)$.

3. The theoretical results

Define $a_n = \max_{1 \leq j \leq p_n} \{|\dot{p}_\lambda(|w_{oj}|)|, |w_{oj}| \neq 0\}$ and $b_n = \max_{1 \leq j \leq p_n} \{|\ddot{p}_\lambda(|w_{oj}|)|, |w_{oj}| \neq 0\}$, where $\ddot{p}_\lambda(x)$ is the second-order derivative of $p_\lambda(x)$, $\eta_{ij} = Y_i - m_j(X_{ij})$, $\eta_i = Y_i - w_{o0} - \sum_{j=1}^{p_n} w_{oj} m_j(X_{ij})$, $\mathcal{M}_i = (m_1(X_{i1}), \dots, m_{p_n}(X_{ip_n}))^T$ and $\hat{\mathcal{M}}_i = (\hat{m}_1(X_{i1}), \dots, \hat{m}_{p_n}(X_{ip_n}))^T$. We assume $p\{\eta_{ij} \leq 0\} = \tau_j$ and $p\{\eta_i \leq 0\} = \tau$. To prove the theoretical results of the proposed estimators, we next present the following technical conditions.

(C1) Let $f_j(\cdot)$ be the marginal density function of the covariates $\{X_{ij}\}$, the j -th element of \mathbf{X}_i . Assume that $f_j(\cdot)$ has continuous derivatives up to the second order and

$$0 < c \leq \inf_j \inf_{x_j \in \mathcal{C}_j} f_j(x_j) \leq \sup_j \sup_{x_j \in \mathcal{C}_j} f_j(x_j) \leq C < \infty$$

where \mathcal{C}_j is the compact support of X_{ij} . For each j , the conditional density functions of Y_i for given X_{ij} exists and satisfies the Lipschitz continuous condition. Furthermore, the length of \mathcal{C}_j is uniformly bounded by a positive constant.

(C2) The kernel function $K(\cdot)$ is a Lipschitz continuous, symmetric and bounded probability density function with a compact support.

(C3) The marginal regression function $m_j(\cdot)$ has continuous derivatives up to the second order and there exists a positive constant c_m such that

$$\sup_j \sup_{x_j \in \mathcal{C}_j} [|m_j(x_j)| + |\dot{m}_j(x_j)| + |\ddot{m}_j(x_j)|] \leq c_m.$$

(C4) Let $f_{\eta_j}(\cdot)$ and $F_{\eta_j}(\cdot)$ be the marginal density and distribution functions of η_{ij} , $f_{\eta}(\cdot)$ and $F_{\eta}(\cdot)$ be the density and distribution functions of η_i . The density functions $f_{\eta_j}(\cdot)$ and $f_{\eta}(\cdot)$ are bounded and bounded away from zero in a neighborhood of zero.

(C5) There exists a sequence of fixed vectors $\{\mathbf{u}\}$ in \mathcal{R}^{p_n} , with $\|\mathbf{u}\|$ bounded, such that $\max_{1 \leq i \leq n} |\hat{\mathcal{M}}_i^T \mathbf{u}| = O(\sqrt{\log n})$, where $\|\cdot\|$ denotes the L_2 norm for any vector.

(C6) The matrix

$$\mathbf{\Lambda}_n = \begin{pmatrix} E[m_1(X_{i1})m_1(X_{i1})] & \cdots & E[m_1(X_{i1})m_{p_n}(X_{ip_n})] \\ \vdots & \vdots & \vdots \\ E[m_{p_n}(X_{ip_n})m_1(X_{i1})] & \cdots & E[m_{p_n}(X_{ip_n})m_{p_n}(X_{ip_n})] \end{pmatrix}$$

is positive definite with the eigenvalues bounded away from zero and infinity. In particular, the smallest eigenvalue of $\mathbf{\Lambda}_n$ is larger than b , a small positive constant.

(C7) $Var(\hat{\mathcal{M}}_i) = \mathbf{G}_n > 0$, $E(\hat{\mathcal{M}}_i^{\otimes 2}) = \mathbf{\Psi}_n$, and $0 < d_1 \leq \lambda_{\min}(\mathbf{\Psi}_n) \leq \lambda_{\max}(\mathbf{\Psi}_n) < d_2 < \infty$ for all n , where $\lambda_{\min}(\mathbf{\Psi}_n)$ and $\lambda_{\max}(\mathbf{\Psi}_n)$ are the smallest and largest eigenvalues of $\mathbf{\Psi}_n$.

(C8) $\liminf_{n \rightarrow \infty} \liminf_{x \rightarrow 0^+} \dot{p}_{\lambda}(x) / \lambda > 0$.

(C9) Let $a_n = O(n^{-1/2})$ and $b_n = o(1)$, and there exist two positive constants C_1 and C_2 such that $|\ddot{p}_{\lambda}(x_1) - \ddot{p}_{\lambda}(x_2)| \leq C_2|x_1 - x_2|$ when $x_1, x_2 > C_1\lambda$.

Without loss of generality, we define the vector of the optimal weights

$$\mathbf{w}_o = (w_{o0}, w_{o1}, \dots, w_{op_n})^T = (w_{o0}, \mathbf{w}_o^T(1), \mathbf{w}_o^T(2))^T,$$

where $\mathbf{w}_o(1) = (w_{o1}, \dots, w_{os_n})^T$ stands for non-zero weights with dimension s_n and $\mathbf{w}_o(2) = (w_{o,s_n+1}, \dots, w_{op_n})^T$ is zero weights with dimension $p_n - s_n$. Let $\hat{\mathbf{w}}_n(1)$ and $\hat{\mathbf{w}}_n(2)$ be the estimators of $\mathbf{w}_o(1)$ and $\mathbf{w}_o(2)$ respectively.

Define $\bar{\eta}_{ij} = m_j(X_{ij})\psi_{\tau}(\eta_i)$, $\boldsymbol{\mu}_n = E(\hat{\mathcal{M}}_i)$, $\boldsymbol{\mu}_{n1}$ is first $s_n \times 1$ vector of $\boldsymbol{\mu}_n$, $\mathbf{\Psi}_{n1}$ and $\mathbf{\Lambda}_{n1}$ are the top-left $s_n \times s_n$ submatrix of $\mathbf{\Psi}_n$ and $\mathbf{\Lambda}_n$, and $\mathbf{G}_{n1} = \mathbf{\Psi}_{n1} - \boldsymbol{\mu}_{n1}\boldsymbol{\mu}_{n1}^T$. Let $\boldsymbol{\xi}_i = \mathbf{m}_i - \boldsymbol{\mu}_{n1}$ with $\mathbf{m}_i = (m_1(X_{i1}), \dots, m_{s_n}(X_{is_n}))^T$, $\tilde{\boldsymbol{\eta}}_i = (\tilde{\eta}_{i1}, \dots, \tilde{\eta}_{is_n})^T$ with $\tilde{\eta}_{ij} = \sum_{k=1}^{s_n} w_{ok}\psi_{\tau_k}(\eta_{ik})\beta_{jk}(X_{ik})$ and $\beta_{jk}(x_k) = E[m_j(X_{ij})|X_{ik} = x_k]$. Obviously, the mean of $(\boldsymbol{\xi}_i\psi_{\tau}(\eta_i) - f_{\eta}(0)\tilde{\boldsymbol{\eta}}_i)$ is zero, and we define $\boldsymbol{\Upsilon}_{n1} = Var\left(n^{-1/2}\sum_{i=1}^n(\boldsymbol{\xi}_i\psi_{\tau}(\eta_i) - f_{\eta}(0)\tilde{\boldsymbol{\eta}}_i)\right)$.

Let $\mathbf{c}_n = (\dot{p}_\lambda(|w_{o1}|) \text{sgn}(w_{o1}), \dots, \dot{p}_\lambda(|w_{os_n}|) \text{sgn}(w_{os_n}))^T$, $\mathbf{\Gamma}_n = \text{diag}\{\ddot{p}_\lambda(|w_{o1}|), \dots, \ddot{p}_\lambda(|w_{os_n}|)\}$ and $\mathcal{X}_j = (X_{1j}, \dots, X_{nj})^T$, where $\text{sgn}(\cdot)$ is the sign function. Define $\mu_k = \int u^k K(u) du$ and $\nu_k = \int u^k K^2(u) du$. In the following theorems, we give the asymptotic theories of $\hat{m}_j(\cdot)$ and $\hat{\mathbf{w}}_n$.

Theorem 1. Suppose that x is an interior point of the support of $f_j(\cdot)$. Under the regularity conditions (C1)–(C4), if $h_j \rightarrow 0$ and $nh_j \rightarrow \infty$ for $j = 1, \dots, p_n$, then the asymptotic conditional bias and variance of the local linear estimator $\hat{m}_j(x)$ are given by

$$\begin{aligned} \text{bias}(\hat{m}_j(x) | \mathcal{X}_j) &= \frac{1}{2} \ddot{m}_j(x) \mu_2 h_j^2 + o_p(h_j^2), \\ \text{Var}(\hat{m}_j(x) | \mathcal{X}_j) &= \frac{1}{nh_j f_j(x)} \frac{\tau_j(1-\tau_j)\nu_0}{(f_{\eta_j}(0))^2} + o_p\left(\frac{1}{nh_j}\right). \end{aligned}$$

Furthermore, conditioning on \mathcal{X}_j , we have

$$\sqrt{nh_j} \left\{ \hat{m}_j(x) - m_j(x) - \frac{1}{2} \ddot{m}_j(x) \mu_2 h_j^2 \right\} \xrightarrow{d} N\left(0, \frac{1}{f_j(x)} \frac{\tau_j(1-\tau_j)\nu_0}{(f_{\eta_j}(0))^2}\right)$$

for $j = 1, \dots, p_n$, where \xrightarrow{d} stands for convergence in distribution.

Remark 1. Theorem 1 shows that the proposed nonparametric estimate $\hat{m}_j(\cdot)$ is $\sqrt{nh_j}$ consistent and enjoys an asymptotically normal distribution.

Theorem 2. Under conditions (C1)–(C9), together with $np_n \left(h_j^2 + (nh_j)^{-1/2}\right) = o(1)$ for $j = 1, \dots, p_n$ and $p_n^2/n \rightarrow 0$, if $\lambda \rightarrow 0$, $\sqrt{n/p_n}\lambda \rightarrow \infty$, then $\forall \mathbf{e}_n \in \mathcal{R}^{s_n}$ such that $\|\mathbf{e}_n\| = 1$, we have

- (i) there exists a local minimizer $\hat{\mathbf{w}}_n$ of the objective function $\mathcal{Q}_n(\mathbf{w}_n)$ defined in (2) such that $\|\hat{\mathbf{w}}_n - \mathbf{w}_o\| = O_p(\sqrt{p_n}(n^{-1/2} + a_n))$;
- (ii) $\hat{\mathbf{w}}_n(2) = \mathbf{0}$ with probability approaching one;
- (iii) $\sqrt{n} \mathbf{e}_n^T \mathbf{\Upsilon}_{n1}^{-1/2} (f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n) [\hat{\mathbf{w}}_n(1) - \mathbf{w}_o(1) + (f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n)^{-1} \mathbf{c}_n] \xrightarrow{d} N(0, 1)$.

Remark 2. Theorem 2 indicates that the estimate of the optimal weight \mathbf{w} is still consistent although the dimension of predictor goes to infinite. Meanwhile, it also shows that the proposed estimate $\hat{\mathbf{w}}_n$ enjoys well-known properties in high dimensional variable selection such as the sparsity and oracle property.

4. Numerical studies

We investigate the performance of the proposed approach by three simulation examples and an empirical application. In our numerical studies, we set the kernel function $K(\cdot)$ as the Epanechnikov kernel, namely, $K(u) = 0.75(1 - u^2)_+$. Bandwidth selection is crucial in local smoothing since it governs the curvature of the fitted function. Similar to Kai et al. (2011), we use the following formula to choose the bandwidth $h = h_{ls} \times \{\tau(1 - \tau)/f(\Phi^{-1}(\tau))\}^{1/5}$, where h_{ls} is the selected optimal bandwidth for least squares, and $f(\cdot)$ and $\Phi(\cdot)$ represent the density function and distribution function of standard normal distribution, respectively. The rule of thumb is used to select the bandwidth h_{ls} . In addition, the tuning parameter λ in the proposed penalized procedure plays an important role. Lian (2012) had proved that the Schwarz information criterion (SIC) is a consistent variable selection criterion under the framework of fixed dimension. In this paper, we select λ by minimizing the following modified SIC criterion (MSIC)

$$\text{MSIC}(\lambda) = \log(\mathcal{Q}_n(\hat{\mathbf{w}}_n)) + df C_n \log(n)/(2n), \quad (3)$$

where $\hat{\mathbf{w}}_n$ is the estimated model weight vector for a given λ , df is the number of nonzero coefficients in $\hat{\mathbf{w}}_n$ and C_n diverges with n . For example, the MSIC criterion reduces to tradition SIC criterion Lian (2012) when $C_n = 1$, and the MSIC criterion is more suitable for high dimensional data if C_n is selected as $\log(p_n)$.

In order to investigate the superiority of the proposed method, we consider the following methods: (1) the proposed semiparametric model average quantile prediction (without SCAD penalty, denoted as SMAQP), (2) the proposed penalized semiparametric model average quantile prediction (with SCAD penalty, denoted as PSMAQP), (3) semiparametric model average mean prediction proposed by Li et al. (2015) (without SCAD penalty, denoted as SMAMP), (4) penalized semiparametric model average mean prediction proposed by Chen et al. (2018) (with SCAD penalty, denoted as PSMAMP). SMAMP and PSMAMP aim to forecast the conditional mean function $E(Y|\mathbf{X})$, and detailed descriptions about the two methods can refer to the section 3 of Li et al. (2015) and subsection 2.1 of Chen et al. (2018). The tuning parameter λ involved in PSMAMP is chosen by the cross-validation according to the advice of Chen et al. (2018), and the R package ‘‘ncvreg’’

can be used to obtain the penalized estimator PSMAMP.

4.1. Simulation studies

In all simulation examples, the sample size n consists of a training set of size n_{tr} and a testing set of size n_{te} , namely, $n = n_{tr} + n_{te}$.

Example 1. For a clear comparison, we adopt similar settings used in Chen et al. (2018) and generate the random samples from the following model

$$Y_i = m_1(X_{i1}) + m_2(X_{i2}) + m_3(X_{i3}) + m_4(X_{i4}) + \varepsilon_i, i = 1, \dots, n, \quad (4)$$

where $m_1(u) = -\sin(2u)$, $m_2(u) = u^2 - 25/12$, $m_3(u) = u$ and $m_4(u) = e^{-u} - \frac{2}{5}\sinh(5/2)$. We fix $n_{te} = 100$ and consider $n_{tr} = 200$ and 400 for example 1. The covariates $\mathbf{X}_i = (X_{i1}, \dots, X_{ip_n})^T$ are independently drawn from $U(-2.5, 2.5)$, and we set the dimension of covariates as $p_n = \lceil n_{tr}^{1/2} \rceil$ which satisfies the theoretical condition $p_n^2/n \rightarrow 0$, where $\lceil s \rceil$ stands for the largest integer not greater than s . Obviously, the first four variables make a significant contribution to estimating the joint multivariate quantile function $m_\tau(\mathbf{X})$, while the rest are not. Therefore, we have reasons to believe that the first four model weights are nonzero and the rest are zero. Please note that the model average component m_j given in section 2 is different from m_j reported in model (4) for $j = 1, \dots, p_n$. Our mission is to achieve the goal of accurately predicting the conditional quantile function $m(\cdot)$, so we are not attempting to estimate m_j in this paper.

In order to examine the robustness of the proposed procedure, we consider the following three different error distributions of ε_i : standard normal distribution (SN), t -distribution with 3 degrees of freedom (t_3), contaminated normal distribution ($MN(\rho, \sigma_1, \sigma_2)$) representing a mixture of $N(0, 1)$ and $N(0, 10^2)$ with weights 0.95 and 0.05 respectively. In addition, four criteria are adopted to evaluate the performance of proposed approach. Firstly, ‘‘C’’, ‘‘IC’’ and ‘‘CF’’ are considered to examine variable selection performance, where ‘‘C’’ represents the average number of zero coefficients in the model weight vector that are correctly estimated to be zero; ‘‘IC’’ represents the average number of nonzero coefficients in the model weight vector that are incorrectly estimated to be zero and ‘‘CF’’ represents the proportion of correctly fitted models (‘‘correctly fit’’ means that the es-

Table 1: Simulation results of C, IC, CF, MPE and their standard deviations (in parenthesis) for $\tau = 0.5$ in example 1.

n_{tr}	Error	method	C	IC	CF	In-Sample Error	Out-of-Sample Error	
200	N(0,1)	SMAMP	–	–	–	0.459 (0.031)	0.577 (0.058)	
		PSMAMP	7.362	0.018	0.258	0.467 (0.032)	0.566 (0.058)	
		SMAQP	–	–	–	0.462 (0.036)	0.580 (0.062)	
	t_3	PSMAQP	9.930	0.108	0.836	0.487 (0.038)	0.557 (0.061)	
		SMAMP	–	–	–	0.601 (0.065)	0.748 (0.085)	
		PSMAMP	6.118	0.030	0.096	0.605 (0.063)	0.730 (0.085)	
	MN	SMAQP	–	–	–	0.592 (0.047)	0.734 (0.081)	
		PSMAQP	9.956	0.206	0.760	0.625 (0.051)	0.705 (0.077)	
		SMAMP	–	–	–	0.702 (0.117)	0.869 (0.141)	
		PSMAMP	4.372	0.048	0.030	0.697 (0.113)	0.846 (0.145)	
		SMAQP	–	–	–	0.637 (0.084)	0.765 (0.122)	
		PSMAQP	9.966	0.222	0.754	0.668 (0.087)	0.738 (0.121)	
	400	N(0,1)	SMAMP	–	–	–	0.445 (0.021)	0.524 (0.046)
			PSMAMP	13.63	0.006	0.414	0.453 (0.022)	0.513 (0.045)
			SMAQP	–	–	–	0.439 (0.023)	0.512 (0.045)
t_3		PSMAQP	15.98	0.006	0.976	0.457 (0.024)	0.494 (0.043)	
		SMAMP	–	–	–	0.597 (0.052)	0.694 (0.081)	
		PSMAMP	11.38	0.002	0.162	0.599 (0.042)	0.673 (0.078)	
MN		SMAQP	–	–	–	0.580 (0.036)	0.664 (0.072)	
		PSMAQP	15.99	0.012	0.978	0.603 (0.037)	0.640 (0.072)	
		SMAMP	–	–	–	0.676 (0.078)	0.789 (0.116)	
		PSMAMP	8.346	0.006	0.040	0.667 (0.075)	0.761 (0.120)	
		SMAQP	–	–	–	0.616 (0.058)	0.693 (0.106)	
		PSMAQP	15.99	0.012	0.984	0.636 (0.058)	0.674 (0.105)	

Notation: To make this a fair comparison, we consider $\tau = 0.5$ for SMAQP and PSMAQP in this table. In addition, the number of zero components of model weight vector is 10 for $n_{tr} = 200$ and 16 for $n_{tr} = 400$.

timination procedure correctly chooses all significant components from the model weight vector). Secondly, the mean prediction error (MPE) is used to measure accuracy of prediction, which is defined as $\sum_{i \in \mathcal{I}} \rho_\tau(Y_i - \hat{Y}_i) / |\mathcal{I}|$, where \mathcal{I} stands for an index set of either the training sample or the testing sample.

Example 2. In this example, similar to Huang and Li (2018), we generate the random samples from the following model

$$Y_i = 3m_1(X_{i1}) + 3m_2(X_{i2}) + 2m_3(X_{i3}) + 2m_4(X_{i4}) + \sqrt{1.74}\varepsilon_i, i = 1, \dots, n, \quad (5)$$

where $m_1(u) = (2u)$, $m_2(u) = (2u - 1)^2$, $m_3(u) = \sin(2\pi u)/(2 - \sin(2\pi u))$ and $m_4(u) = 0.1\sin(2\pi u) + 0.2\cos(2\pi u) + 0.3\sin(2\pi u)^2 + 0.4\cos(2\pi u)^3 + 0.5\sin(2\pi u)^3$. The covariates $\mathbf{X}_i = (X_{i1}, \dots, X_{ip_n})^T$ are simulated by $X_{ij} = (W_{ij} + tU_i) / (1 + t)$ for $t = 1$ and $j = 1, \dots, p_n$, where W_{ij} and U_i are independently drawn from $U(0, 1)$ and $p_n = \lceil n_{tr}^{1/2} \rceil$. We also fix $n_{te} = 100$ and consider $n_{tr} = 400$ and 800 for example 2. Other settings are the same as that in example 1.

It is easy to find that the conditional mean function $E(Y|\mathbf{X})$ is equal to the conditional quantile function $m_\tau(\mathbf{X})$ for $\tau = 0.5$. Thus, we can compare mean prediction approaches (SMAMP and PSMAMP) with quantile prediction approaches (SMAQP and PSMAQP)

Table 2: Simulation results of C, IC, CF, MPE and their standard deviations (in parenthesis) for $\tau = 0.75$ in example 1.

n_{tr}	Error	method	C	IC	CF	In-Sample Error	Out-of-Sample Error
200	N(0,1)	SMAQP	–	–	–	0.917 (0.099)	1.032 (0.163)
		PSMAQP	9.896	0.562	0.468	0.961 (0.108)	1.027 (0.164)
	t_3	SMAQP	–	–	–	1.087 (0.110)	1.239 (0.170)
		PSMAQP	9.866	0.768	0.344	1.139 (0.130)	1.229 (0.177)
	MN	SMAQP	–	–	–	1.108 (0.128)	1.256 (0.213)
		PSMAQP	9.918	0.848	0.338	1.168 (0.148)	1.253 (0.215)
400	N(0,1)	SMAQP	–	–	–	0.850 (0.075)	0.921 (0.119)
		PSMAQP	15.97	0.134	0.844	0.875 (0.077)	0.909 (0.121)
	t_3	SMAQP	–	–	–	1.035 (0.085)	1.129 (0.148)
		PSMAQP	15.95	0.224	0.740	1.066 (0.088)	1.110 (0.147)
	MN	SMAQP	–	–	–	1.054 (0.096)	1.147 (0.181)
		PSMAQP	15.97	0.208	0.776	1.081 (0.100)	1.127 (0.178)

Notation: The number of zero components of model weight vector is 10 for $n_{tr} = 200$ and 16 for $n_{tr} = 400$.

at $\tau = 0.5$. The MPE criterion is reduced to $\sum_{i \in \mathcal{I}} \frac{1}{2} |Y_i - \hat{Y}_i| / |\mathcal{I}|$ for $\tau = 0.5$, and thus this criterion also can be used to assess the prediction performance of mean prediction approaches. The corresponding results of mean prediction approaches (SMAMP and PSMAMP) and quantile prediction approaches (SMAQP and PSMAQP) at $\tau = 0.5$ are reported in Tables 1 and 3. We can obtain the following findings. Firstly, the values in the column labeled ‘‘C’’ gradually tend to the true number of zero components with the training sample size increasing. The CF values are very close to one for a large training sample size (e.g. $n_{tr} = 400$), which shows that the proposed penalized procedure can consistently select significant components in weight vector. However, the existing mean prediction approach PSMAMP performs badly due to lower CF values. Secondly, unpenalized methods always has smaller in-sample MPE than the penalized methods’s, but it does not hold true for out-of-sample MPE. For heavy-tailed distributions t_3 and contaminated distribution MN, it is not hard to find that our proposed penalized method PSMAQP is best in terms of prediction accuracy among all methods. Meanwhile, there is little difference for PSMAMP and PSMAQP under the normal error distribution. Thirdly, Tables 2 and 4 give the simulation results of SMAQP and PSMAQP at $\tau = 0.75$. The results also show that PSMAQP has better prediction performance.

Example 3. The conditional quantile function $Q_\tau(Y|\mathbf{X})$ is considered as

$$m_\tau(\mathbf{X}) = Q_\tau(Y|\mathbf{X}) = 1 + \Phi^{-1}(\tau) + 2X_{i1} + 3X_{i2}^2 - \log(1 - X_{i3}) + \Phi^{-1}(X_{i4}) + (1 + \Phi^{-1}(\tau) - \log(1 - \tau))X_{i5}, i = 1, \dots, n, \quad (6)$$

where Φ is the standard normal distribution function, $\mathbf{X}_i = (X_{i1}, \dots, X_{ip_n})^T$ are inde-

Table 3: Simulation results of C, IC, CF, MPE and their standard deviations (in parenthesis) for $\tau = 0.5$ and $t = 1$ in example 2.

n_{tr}	Error	method	C	IC	CF	In-Sample Error	Out-of-Sample Error
400	N(0,1)	SMAMP	–	–	–	0.507 (0.021)	0.566 (0.045)
		PSMAMP	11.75	0.000	0.212	0.515 (0.022)	0.555 (0.045)
		SMAQP	–	–	–	0.501 (0.020)	0.594 (0.047)
	t_3	PSMAQP	15.97	0.006	0.966	0.530 (0.022)	0.566 (0.045)
		SMAMP	–	–	–	0.724 (0.054)	0.810 (0.105)
		PSMAMP	7.750	0.006	0.026	0.725 (0.052)	0.790 (0.107)
	MN	SMAQP	–	–	–	0.692 (0.042)	0.802 (0.104)
		PSMAQP	15.98	0.048	0.936	0.728 (0.044)	0.769 (0.103)
		SMAMP	–	–	–	0.837 (0.105)	0.932 (0.154)
		PSMAMP	5.740	0.012	0.000	0.826 (0.104)	0.908 (0.156)
		SMAQP	–	–	–	0.740 (0.072)	0.839 (0.151)
		PSMAQP	15.99	0.044	0.946	0.772 (0.073)	0.808 (0.151)
800	N(0,1)	SMAMP	–	–	–	0.511 (0.014)	0.553 (0.044)
		PSMAMP	18.61	0	0.276	0.519 (0.015)	0.544 (0.043)
		SMAQP	–	–	–	0.508 (0.014)	0.567 (0.045)
	t_3	PSMAQP	23.99	0	0.988	0.528 (0.014)	0.546 (0.043)
		SMAMP	–	–	–	0.728 (0.038)	0.795 (0.089)
		PSMAMP	13.89	0	0.036	0.727 (0.035)	0.774 (0.090)
	MN	SMAQP	–	–	–	0.704 (0.030)	0.784 (0.088)
		PSMAQP	23.99	0	0.998	0.730 (0.031)	0.758 (0.087)
		SMAMP	–	–	–	0.817 (0.066)	0.884 (0.140)
		PSMAMP	9.662	0	0.008	0.803 (0.067)	0.856 (0.146)
		SMAQP	–	–	–	0.744 (0.048)	0.810 (0.137)
		PSMAQP	24.00	0	1.000	0.767 (0.048)	0.787 (0.136)

Notation: To make this a fair comparison, we consider $\tau = 0.5$ for RSMAP and PRSMAP in this table. In addition, the number of zero components of model weight vector is 16 for $n_{tr} = 400$ and 24 for $n_{tr} = 800$.

Table 4: Simulation results of C, IC, CF, MPE and their standard deviations (in parenthesis) for $\tau = 0.75$ and $t = 1$ in example 2.

n_{tr}	Error	method	C	IC	CF	In-Sample Error	Out-of-Sample Error
400	N(0,1)	SMAQP	–	–	–	0.848 (0.042)	0.938 (0.095)
		PSMAQP	15.92	0.010	0.912	0.882 (0.043)	0.922 (0.094)
	t_3	SMAQP	–	–	–	1.111 (0.070)	1.220 (0.139)
		PSMAQP	15.91	0.110	0.828	1.147 (0.073)	1.188 (0.133)
	MN	SMAQP	–	–	–	1.121 (0.092)	1.204 (0.195)
		PSMAQP	15.93	0.094	0.852	1.160 (0.094)	1.184 (0.193)
800	N(0,1)	SMAQP	–	–	–	0.853 (0.029)	0.912 (0.083)
		PSMAQP	23.97	0.000	0.968	0.879 (0.030)	0.901 (0.081)
	t_3	SMAQP	–	–	–	1.114 (0.049)	1.179 (0.150)
		PSMAQP	23.97	0.002	0.968	1.136 (0.048)	1.151 (0.143)
	MN	SMAQP	–	–	–	1.116 (0.065)	1.165 (0.179)
		PSMAQP	23.98	0.000	0.984	1.142 (0.065)	1.146 (0.174)

Notation: The number of zero components of model weight vector is 16 for $n_{tr} = 400$ and 24 for $n_{tr} = 800$.

Table 5: Simulation results of C, IC, CF, MEE and their standard deviations (in parenthesis) in example 3.

n_{tr}	τ	method	C	IC	CF	In-Sample Error	Out-of-Sample Error
400	0.5	SMAQP	–	–	–	0.328 (0.038)	0.341 (0.047)
		PSMAQP	14.96	0.350	0.652	0.258 (0.049)	0.266 (0.056)
	0.75	SMAQP	–	–	–	0.383 (0.047)	0.401 (0.060)
800	0.5	PSMAQP	14.90	0.236	0.702	0.301 (0.058)	0.310 (0.066)
		SMAQP	–	–	–	0.266 (0.027)	0.273 (0.036)
	PSMAQP	22.99	0.014	0.974	0.177 (0.029)	0.180 (0.033)	
	0.75	SMAQP	–	–	–	0.316 (0.033)	0.325 (0.046)
	PSMAQP	22.95	0.022	0.932	0.217 (0.037)	0.221 (0.042)	

Notation: The number of zero components of model weight vector is 15 for $n_{tr} = 400$ and 23 for $n_{tr} = 800$.

pendently drawn from $U(0, 1)$ and $1 + \Phi^{-1}(\tau)$ can be regarded as the intercept. We fix $n_{te} = 100$ and consider $n_{tr} = 400$ and 800 for example 3 and $p_n = \lceil n_{tr}^{1/2} \rceil$. Obviously, the fifth covariate's coefficient varies with τ , and only the first five predictors are significant for predicting $Q_\tau(Y|\mathbf{X})$. The first two examples come from the nonparametric additive model, but the proposed approach do not need any model assumption. Thus we aim to confirm that our method is model free in this example. To assess the estimation accuracy of $\hat{m}_\tau(\mathbf{X})$, we consider the mean estimation error (MEE) defined as $MEE = \sum_{i \in \mathcal{I}} \frac{1}{2} |m_\tau(\mathbf{X}) - \hat{m}_\tau(\mathbf{X})| / |\mathcal{I}|$ in this example. Table 5 lists the simulation results which show that the proposed PSMAQP performs well for different quantiles.

Overall, the proposed model free procedure PSMAQP is competitive when compared with the existing methods, and its finite sample performances are satisfactory.

4.2. An application

In this section, we apply our proposed method to analyze the body fat dataset (Johnson, 1996), which is available from <http://lib.stat.cmu.edu/datasets/bodyfat>. This dataset consists of 252 observations without missing values. The purpose of studying this dataset is to predict the percentage of body fat according to various body circumference measurements. Thus, the percentage of body fat is taken as the response variable and other body circumference measurements are regarded as the predictors. Brief descriptions and marginal Pearson correlations of 14 variables are summarized in Table 6. More details can refer to Johnson (1996). Before employing prediction methods, we take the logarithm transformation for all predictors.

To evaluate the predictive performance of various methods, the data is split into two parts. One part including n_{tr} observations is used as a training data set to estimate the weight vector \mathbf{w}_n and the marginal quantile functions $m_j(\cdot), j = 1, \dots, p_n$, and the other part including n_{te} observations is considered as a testing data set to evaluate the predictive ability of various methods. In this real data analysis, we consider $n_{tr} = 150$ and 200 , and $n_{te} = n - n_{tr} = 252 - n_{tr}$.

Table 7 reports the in-sample and out-of-sample mean prediction errors (MPE) and the corresponding sample standard deviations (SD) over 500 random partitions. Firstly, for $\tau = 0.5$ and in-sample performance, it is easy to see that SMAQP performs best among

Table 6: Regressors for the body fat dataset.

Variable	Name	Description	Correlation with Y
X_1	Age	Age (years)	0.2921
X_2	Weight	Weight(lbs)	0.6287
X_3	Height	Height (inches)	-0.0990
X_4	Neck	Neck circumference (cm)	0.4905
X_5	Chest	Chest circumference (cm)	0.7051
X_6	Abdomen	Abdomen 2 circumference (cm)	0.8218
X_7	Hip	Hip circumference (cm)	0.6365
X_8	Thigh	Thigh circumference (cm)	0.5680
X_9	Knee	Knee circumference (cm)	0.5102
X_{10}	Ankle	Ankle circumference (cm)	0.2796
X_{11}	Biceps	Biceps (extended) circumference (cm)	0.5000
X_{12}	Forearm	Forearm circumference (cm)	0.3584
X_{13}	Wrist	Wrist circumference (cm)	0.3447
Y	Body Fat(%)	Percent body fat using Siri's equation: 495/Density - 450	1.0000

Table 7: Prediction results ($\times 10^{-2}$) for analysis of the body fat dataset.

	method	$n_{tr} = 150$				$n_{tr} = 200$			
		In-Sample		Out-of-Sample		In-Sample		Out-of-Sample	
		MPE	SD	MPE	SD	MPE	SD	MPE	SD
$\tau = 0.5$	SMAMP	1.622	0.069	2.116	0.691	1.653	0.043	2.242	1.402
	PSMAMP	1.763	0.136	2.117	0.544	1.829	0.105	2.314	1.357
	SMAQP	1.593	0.067	1.940	0.143	1.634	0.043	1.890	0.174
	PSMAQP	1.683	0.081	1.896	0.125	1.696	0.055	1.858	0.166
$\tau = 0.25$	SMAQP	2.666	0.151	2.955	0.338	2.687	0.098	2.857	0.363
	PSMAQP	2.726	0.165	2.914	0.314	2.742	0.089	2.853	0.348
$\tau = 0.75$	SMAQP	2.768	0.169	3.118	0.383	2.838	0.115	3.079	0.451
	PSMAQP	2.811	0.171	3.080	0.340	2.859	0.122	3.056	0.401

four approaches. For out-of-sample performance, one can see clearly that the proposed penalized approach PSMAQP has smallest MPE and SD for different settings, which shows that our proposed method has better predictive ability. Secondly, for $\tau = 0.25$ and 0.75, we can see that PSMAQP always performs better than SMAQP in terms of out-of-sample performance.

To investigate the estimated weights, we list the estimated weights at $\tau = 0.5$ and their standard deviations (in brackets) calculated by the bootstrap resampling method (Horowitz, 1998). Obviously, the weights for the penalized prediction methods (PSMAMP and PSMAQP) are relatively sparse with much smaller standard deviations than the unpenalized prediction methods (SMAMP and SMAQP). Meanwhile, it is not hard to find that PSMAQP is most efficient among all methods due to the smallest standard deviations. In addition, for PSMAMP, only the sixth predictor (X_6) is chose as the significant variable whose marginal quantile function has significant influence on estimating $m_\tau(\mathbf{X})$. However, PSMAQP selects five predictors (including X_1 , X_4 , X_6 , X_9 and X_{13}) as the significant variables. In summary, our proposed model averaging procedure generally works well and outperforms other existing methods.

Table 8: Estimated weights and their standard deviations (in brackets) for the body fat study at $\tau = 0.5$.

weight	SMAMP	PSMAMP	SMAQP	PSMAQP
w_0	-0.056 (12.091)	-0.044 (15.681)	-0.038 (0.584)	0.063 (0.070)
w_1	0.660 (0.306)	0.000 (0.391)	0.419 (0.189)	0.314 (0.210)
w_2	-0.530 (0.546)	0.000 (0.446)	-0.115 (0.546)	0.000 (0.231)
w_3	1.114 (3.454)	0.000 (2.229)	0.146 (2.976)	0.000 (0.192)
w_4	-0.418 (0.338)	0.000 (0.397)	-0.330 (0.216)	-0.238 (0.225)
w_5	-0.055 (0.210)	0.000 (0.192)	0.005 (0.248)	0.000 (0.128)
w_6	1.643 (0.151)	1.234 (0.185)	1.174 (0.185)	1.187 (0.170)
w_7	-0.188 (0.291)	0.000 (0.259)	-0.164 (0.282)	0.000 (0.190)
w_8	0.515 (0.251)	0.000 (0.291)	0.282 (0.178)	0.000 (0.188)
w_9	-0.185 (0.336)	0.000 (0.264)	-0.195 (0.195)	-0.049 (0.167)
w_{10}	0.330 (0.752)	0.000 (0.590)	0.204 (0.312)	0.000 (0.156)
w_{11}	0.292 (0.209)	0.000 (0.214)	0.124 (0.166)	0.000 (0.167)
w_{12}	0.204 (0.298)	0.000 (0.245)	0.343 (0.205)	0.000 (0.158)
w_{13}	-2.095 (1.273)	0.000 (1.519)	-0.696 (0.303)	-0.561 (0.349)

5. Conclusion

In this paper, we provide a new semiparametric model averaging estimation for forecasting the conditional quantile function $m_\tau(\mathbf{X})$ under the high-dimensional settings. Based on local linear regression, we firstly estimate the individual marginal regression functions $m_j(\cdot)$ by minimizing the local weighted quantile loss function. Then, a penalized quantile regression is developed to select the regressors whose marginal regression functions make significant contribution in estimating the quantile function $m_\tau(\mathbf{X})$. Simulations and empirical example in Section 4 show that the proposed method performs reasonably well in finite samples.

Recently, under the framework of ultra-high dimension setting, Ando and Li (2014) developed a new model averaging approach based on delete-one cross-validation criterion and proved that the proposed method could achieve the lowest possible prediction loss asymptotically. But they only considered high dimensional parametric model averaging, which may increase the risk of model misspecification. Thus, it is interesting to study semiparametric model averaging estimation for ultra-high dimensional data. Research in these aspects is ongoing.

Acknowledgments

This work is supported by the National Social Science Fund of China (Grant No. 17CTJ015).

Appendix

Let C denote a positive constant that may be different at different place throughout this paper. Let $u = \sqrt{nh_j}(a - m_j(x))$, $v = h_j\sqrt{nh_j}(b - \dot{m}_j(x))$, $x_{ij} = (X_{ij} - x)/h_j$,

$K_{ij} = K(x_{ij})$, $d_{ij} = m_j(X_{ij}) - m_j(x) - \dot{m}_j(x)(X_{ij} - x)$ and $\Delta_{ij} = (u + vx_{ij})/\sqrt{nh_j}$. Define $S_{ij}^* = I(\eta_{ij} + d_{ij} \leq 0) - \tau_j$, $\mathbf{W}_n^* = (w_1^*, w_2^*)^T$, $w_1^* = \frac{1}{\sqrt{nh_j}} \sum_{i=1}^n K_{ij} S_{ij}^*$, $w_2^* = \frac{1}{\sqrt{nh_j}} \sum_{i=1}^n K_{ij} S_{ij}^* x_{ij}$.

Lemma 1. Denote $\hat{\boldsymbol{\theta}} = (\hat{u}, \hat{v})^T$ as the minimizer of (1). Then, under the regularity conditions (C1)–(C4), we have

$$\hat{\boldsymbol{\theta}} + \frac{1}{f_j(x)} \mathbf{S}^{-1} E[\mathbf{W}_n^* | \mathcal{X}_j] \xrightarrow{d} N\left(\mathbf{0}, \frac{1}{f_j(x)} \mathbf{S}^{-1} \boldsymbol{\Sigma} \mathbf{S}^{-1}\right),$$

where $\mathbf{S} = \begin{pmatrix} f_{\eta_j}(0) & 0 \\ 0 & f_{\eta_j}(0) \mu_2 \end{pmatrix}$ and $\boldsymbol{\Sigma} = \begin{pmatrix} \tau_j(1 - \tau_j) \nu_0 & 0 \\ 0 & \tau_j(1 - \tau_j) \nu_2 \end{pmatrix}$.

Proof of Lemma 1. To apply the identity (Knight, 1998)

$$\rho_\tau(u - v) - \rho_\tau(u) = v \{I(u \leq 0) - \tau\} + \int_0^v \{I(u \leq t) - I(u \leq 0)\} dt. \quad (\text{A.1})$$

Minimizing expression (1) is equivalent to minimizing

$$L_n(\boldsymbol{\theta}) = \sum_{i=1}^n [\rho_{\tau_j} \{\eta_{ij} + d_{ij} - \Delta_{ij}\} K_{ij} - \rho_{\tau_j} \{\eta_{ij} + d_{ij}\} K_{ij}].$$

Using identity (A.1) and with some straightforward calculations, it follows that

$$\begin{aligned} L_n(\boldsymbol{\theta}) &= \frac{1}{\sqrt{nh_j}} \sum_{i=1}^n K_{ij} S_{ij}^* u + \frac{1}{\sqrt{nh_j}} \sum_{i=1}^n K_{ij} S_{ij}^* x_{ij} v + B_{nj}(\boldsymbol{\theta}) \\ &= \mathbf{W}_n^{*T} \boldsymbol{\theta} + B_{nj}(\boldsymbol{\theta}), \end{aligned} \quad (\text{A.2})$$

where $B_{nj}(\boldsymbol{\theta}) = \sum_{i=1}^n \int_0^{\Delta_{ij}} K_{ij} \{I(\eta_{ij} + d_{ij} \leq t) - I(\eta_{ij} + d_{ij} \leq 0)\} dt$. Then

$$\begin{aligned} E(B_{nj}(\boldsymbol{\theta}) | \mathcal{X}_j) &= \sum_{i=1}^n \int_0^{\Delta_{ij}} K_{ij} \{F_{\eta_j}(t - d_{ij}) - F_{\eta_j}(-d_{ij})\} dt \\ &= \sum_{i=1}^n \int_0^{\Delta_{ij}} K_{ij} \{f_{\eta_j}(-d_{ij}) t (1 + o(1))\} dt \\ &= \frac{1}{2} f_{\eta_j}(0) \sum_{i=1}^n K_{ij} \Delta_{ij}^2 + o_p(1) \\ &= \frac{1}{2} \boldsymbol{\theta}^T \mathbf{S}_n \boldsymbol{\theta} + o_p(1), \end{aligned} \quad (\text{A.3})$$

where

$$\mathbf{S}_n = \begin{pmatrix} \frac{1}{nh_j} f_{\eta_j}(0) \sum_{i=1}^n K_{ij} & \frac{1}{nh_j} f_{\eta_j}(0) \sum_{i=1}^n x_{ij} K_{ij} \\ \frac{1}{nh_j} f_{\eta_j}(0) \sum_{i=1}^n x_{ij} K_{ij} & \frac{1}{nh_j} f_{\eta_j}(0) \sum_{i=1}^n x_{ij}^2 K_{ij} \end{pmatrix},$$

and

$$\begin{aligned} & \text{Var}(B_{nj}(\boldsymbol{\theta}) | \mathcal{X}_j) \\ &= \sum_{i=1}^n \text{Var} \left\{ \left(\int_0^{\Delta_{ij}} K_{ij} \{I(\eta_{ij} + d_{ij} \leq t) - I(\eta_{ij} + d_{ij} \leq 0)\} dt \right) | \mathcal{X}_j \right\} \\ &\leq \sum_{i=1}^n E \left\{ \left(\int_0^{\Delta_{ij}} K_{ij} \{I(\eta_{ij} + d_{ij} \leq t) - I(\eta_{ij} + d_{ij} \leq 0)\} dt \right)^2 | \mathcal{X}_j \right\} \\ &\leq \sum_{i=1}^n K_{ij}^2 \int_0^{|\Delta_{ij}|} \int_0^{|\Delta_{ij}|} \{F_{\eta_j}(|\Delta_{ij}| - d_{ij}) - F_{\eta_j}(-d_{ij})\} dt_1 dt_2 \\ &= o \left(\sum_{i=1}^n K_{ij}^2 \Delta_{ij}^2 \right) = o_p(1). \end{aligned} \tag{A.4}$$

Similar to Parzen (1962), we have

$$\frac{1}{nh_j} \sum_{i=1}^n K_{ij} x_{ij}^k \xrightarrow{p} f_j(x) \mu_k,$$

where \xrightarrow{p} stands for convergence in probability. Thus

$$\mathbf{S}_n \xrightarrow{p} f_j(x) \begin{pmatrix} f_{\eta_j}(0) & 0 \\ 0 & f_{\eta_j}(0) \mu_2 \end{pmatrix} = f_j(x) \mathbf{S}.$$

This together with (A.2)–(A.4), leads to

$$L_n(\boldsymbol{\theta}) = \frac{1}{2} f_j(x) \boldsymbol{\theta}^T \mathbf{S} \boldsymbol{\theta} + \mathbf{W}_n^{*T} \boldsymbol{\theta} + o_p(1)$$

Since the convex function $L_n(\boldsymbol{\theta}) - \mathbf{W}_n^{*T} \boldsymbol{\theta}$ converges in probability to the convex function $\frac{1}{2} f_j(x) \boldsymbol{\theta}^T \mathbf{S} \boldsymbol{\theta}$, it follows from the convexity lemma (Pollard, 1991) that, for any compact set Θ , the quadratic approximation to $L_n(\boldsymbol{\theta})$ holds uniformly for $\boldsymbol{\theta}$ in any compact set, which leads to

$$\hat{\boldsymbol{\theta}} = -\frac{1}{f_j(x)} \mathbf{S}^{-1} \mathbf{W}_n^* + o_p(1).$$

Denote $S_{ij} = I(\eta_{ij} \leq 0) - \tau_j$ and $\mathbf{W}_n = (w_1, w_2)^T$ with $w_1 = \frac{1}{\sqrt{nh_j}} \sum_{i=1}^n K_{ij} S_{ij}$ and $w_2 =$

$\frac{1}{\sqrt{nh_j}} \sum_{i=1}^n K_{ij} S_{ij} x_{ij}$. By the Cramér-Wold theorem and central limit theorem, we have

$$\frac{\mathbf{W}_n | \mathcal{X}_j - E(\mathbf{W}_n | \mathcal{X}_j)}{\sqrt{\text{Var}(\mathbf{W}_n | \mathcal{X}_j)}} \xrightarrow{d} N(\mathbf{0}, \mathbf{I}_{2 \times 2}). \quad (\text{A.5})$$

Note that $E(S_{ij}) = 0$ and $\text{Var}(S_{ij}) = \tau_j(1 - \tau_j)$. Similar to Parzen (1962), we have

$\frac{1}{nh_j} \sum_{i=1}^n K_{ij}^2 x_{ij}^k \xrightarrow{p} f_j(x) \nu_k$. Therefore,

$$\text{Var}(\mathbf{W}_n | \mathcal{X}_j) \xrightarrow{p} f_j(x) \begin{pmatrix} \tau_j(1 - \tau_j) \nu_0 & 0 \\ 0 & \tau_j(1 - \tau_j) \nu_2 \end{pmatrix} = f_j(x) \boldsymbol{\Sigma}.$$

Combined with (A.5), we have $\mathbf{W}_n | \mathcal{X}_j \xrightarrow{d} N(\mathbf{0}, f_j(x) \boldsymbol{\Sigma})$. Moreover, we have

$$\begin{aligned} \text{Var}(w_1^* - w_1 | \mathcal{X}_j) &= \frac{1}{nh_j} \sum_{i=1}^n K_{ij}^2 \text{Var}(S_{ij}^* - S_{ij}) \\ &\leq \frac{1}{nh_j} \sum_{i=1}^n K_{ij}^2 [F_{\eta_j}(|d_{ij}|) - F_{\eta_j}(0)] = o_p(1), \end{aligned}$$

and

$$\begin{aligned} \text{Var}(w_2^* - w_2 | \mathcal{X}_j) &= \frac{1}{nh_j} \sum_{i=1}^n K_{ij}^2 x_{ij}^2 \text{Var}(S_{ij}^* - S_{ij}) \\ &\leq \frac{1}{nh_j} \sum_{i=1}^n K_{ij}^2 x_{ij}^2 [F_{\eta_j}(|d_{ij}|) - F_{\eta_j}(0)] = o_p(1). \end{aligned}$$

Thus, we obtain $\text{Var}(\mathbf{W}_n^* - \mathbf{W}_n | \mathcal{X}_j) = o_p(1)$. So by Slutsky's theorem, conditioning on \mathcal{X}_j , we have

$$\mathbf{W}_n^* | \mathcal{X}_j - E(\mathbf{W}_n^* | \mathcal{X}_j) \xrightarrow{d} N(\mathbf{0}, f_j(x) \boldsymbol{\Sigma}).$$

Therefore,

$$\hat{\boldsymbol{\theta}} + \frac{1}{f_j(x)} \mathbf{S}^{-1} E(\mathbf{W}_n^* | \mathcal{X}_j) \xrightarrow{d} N\left(\mathbf{0}, \frac{1}{f_j(x)} \mathbf{S}^{-1} \boldsymbol{\Sigma} \mathbf{S}^{-1}\right).$$

This completes the proof. \square

Proof of Theorem 1. Let us calculate the conditional bias and variance.

$$\begin{aligned} \text{bias}(\hat{m}_j(x) | \mathcal{X}_j) &= -\frac{1}{\sqrt{nh_j f_j(x)}} \frac{1}{f_{\eta_j}(0)} \frac{1}{\sqrt{nh_j}} E\left(\sum_{i=1}^n K_{ij} S_{ij}^* | \mathcal{X}_j\right) \\ &= -\frac{1}{nh_j} \frac{1}{f_j(x) f_{\eta_j}(0)} \sum_{i=1}^n K_{ij} [F_{\eta_j}(-d_{ij}) - F_{\eta_j}(0)] \\ &= \frac{1}{nh_j} \frac{1}{f_j(x)} \sum_{i=1}^n K_{ij} d_{ij}. \end{aligned}$$

By using the fact that

$$\frac{1}{nh_j} \sum_{i=1}^n K_{ij} d_{ij} = \frac{1}{2} f_j(x) \ddot{m}_j(x) \mu_2 h_j^2 \{1 + o_p(1)\}.$$

We obtain

$$\text{bias}(\hat{m}_j(x) | \mathcal{X}_j) = \frac{1}{2} \ddot{m}_j(x) \mu_2 h_j^2 + o_p(h_j^2).$$

Furthermore, the conditional variance of $\hat{m}_j(x)$ is

$$\text{Var}(\hat{m}_j(x) | \mathcal{X}_j) = \frac{1}{nh_j f_j(x)} \frac{\tau_j (1 - \tau_j) \nu_0}{(f_{\eta_j}(0))^2} + o_p\left(\frac{1}{nh_j}\right).$$

□

Proof of Theorem 2. Let $\alpha_n = \sqrt{p_n} (n^{-1/2} + a_n)$, $\mathbf{w}_o = (w_{o0}, \mathbf{w}_o)^T$, $\mathbf{w}_o = (w_{o1}, \dots, w_{op_n})^T$, $v = \alpha_n^{-1} (\hat{w}_0 - w_{o0})$, $\mathbf{u} = \alpha_n^{-1} (\hat{\mathbf{w}} - \mathbf{w}_o)$ and $\mathcal{C}_n = \left\{ (v, \mathbf{u}) : \left\| (v, \mathbf{u}^T)^T \right\| = C \right\}$. In order to prove the convergence rate in Theorem 2(i), our aim is to show that for any given ϵ there is a large constant C such that, for a large n , we have

$$P \left\{ \inf_{(v, \mathbf{u}) \in \mathcal{C}_n} \mathcal{Q}_n(\mathbf{w}_o + \alpha_n(v, \mathbf{u}^T)^T) \geq \mathcal{Q}_n(\mathbf{w}_o) \right\} \geq 1 - \epsilon. \quad (\text{A.6})$$

Using $p_\lambda(0) = 0$ and identity (A.1) together with some straightforward calculations, it follows that

$$\begin{aligned} L_n &= \mathcal{Q}_n(\mathbf{w}_o + \alpha_n(v, \mathbf{u}^T)^T) - \mathcal{Q}_n(\mathbf{w}_o) \\ &\geq \sum_{i=1}^n \alpha_n (v + \hat{\mathcal{M}}_i^T \mathbf{u}) \left[I \left\{ Y_i - w_{o0} - \hat{\mathcal{M}}_i^T \mathbf{w}_o \leq 0 \right\} - \tau \right] \\ &\quad + \sum_{i=1}^n \int_0^{\alpha_n(v + \hat{\mathcal{M}}_i^T \mathbf{u})} \left[I \left\{ Y_i - w_{o0} - \hat{\mathcal{M}}_i^T \mathbf{w}_o \leq t \right\} - I \left\{ Y_i - w_{o0} - \hat{\mathcal{M}}_i^T \mathbf{w}_o \leq 0 \right\} \right] dt \\ &\quad + n \sum_{j=1}^{s_n} p_\lambda(|w_{oj} + \alpha_n u_j|) - n \sum_{j=1}^{s_n} p_\lambda(|w_{oj}|) \\ &\stackrel{\Delta}{=} \sqrt{n} \alpha_n (\zeta v + \mathbf{z}_n^T \mathbf{u}) + B_n + P_n, \end{aligned} \quad (\text{A.7})$$

where

$$\begin{aligned}\zeta &= n^{-1/2} \sum_{i=1}^n [I\{\eta_i \leq \Delta_i\} - \tau], \\ \mathbf{z}_n &= n^{-1/2} \sum_{i=1}^n \hat{\mathcal{M}}_i [I\{\eta_i \leq \Delta_i\} - \tau], \\ B_n &= \sum_{i=1}^n \int_0^{\alpha_n(v + \hat{\mathcal{M}}_i^T \mathbf{u})} [I\{\eta_i \leq t + \Delta_i\} - I\{\eta_i \leq \Delta_i\}] dt, \\ P_n &= n \sum_{j=1}^{s_n} p_\lambda(|w_{oj} + \alpha_n u_j|) - n \sum_{j=1}^{s_n} p_\lambda(|w_{oj}|),\end{aligned}$$

where $\Delta_i = (\hat{\mathcal{M}}_i - \mathcal{M}_i)^T \mathbf{w}_o$. Note that, by (C7), $E(\mathbf{z}_n^T \mathbf{u}) = 0$ and $E((\mathbf{z}_n^T \mathbf{u})^2) = \mathbf{u}^T E(\mathbf{z}_n \mathbf{z}_n^T) \mathbf{u} = C \mathbf{u}^T \Psi_n \mathbf{u} \leq C \mathbf{u}^T \lambda_{max}(\Psi_n) \mathbf{u} = O(\|\mathbf{u}\|^2)$. Hence, $\mathbf{z}_n^T \mathbf{u} = O(\|\mathbf{u}\|)$.

This combined with (A.7) leads to

$$L_n = B_n + P_n + o_p(n\alpha_n^2) \|\mathbf{u}\|. \quad (\text{A.8})$$

By the definition of $\hat{\mathcal{M}}_i$ and \mathcal{M}_i in section 3 and the consistency result in Theorem 1, we have, for x_j and $j = 1, \dots, p_n$, $\hat{m}_j(x_j) - m_j(x_j) = O(h_j^2 + (nh_j)^{-1/2})$. Observe that

$$\hat{\mathcal{M}}_i \hat{\mathcal{M}}_i^T - \mathcal{M}_i \mathcal{M}_i^T = (\hat{\mathcal{M}}_i - \mathcal{M}_i) \mathcal{M}_i^T + \mathcal{M}_i (\hat{\mathcal{M}}_i - \mathcal{M}_i)^T + (\hat{\mathcal{M}}_i - \mathcal{M}_i) (\hat{\mathcal{M}}_i - \mathcal{M}_i)^T.$$

Then, for any δ , by Chebyshev's inequality and following the proof of Lemma 8 in Fan and Peng (2004), we have

$$P \left\{ \left\| \frac{1}{n} \sum_{i=1}^n \mathcal{M}_i \mathcal{M}_i^T - \Lambda_n \right\|_F > \delta \right\} \leq \frac{1}{n^2 \delta^2} E \left[\left\| \sum_{i=1}^n \mathcal{M}_i \mathcal{M}_i^T - n \Lambda_n \right\|_F^2 \right] = O(p_n^2/n) = o(1),$$

where $\|\mathbf{A}\|_F$ stands for Frobenius norm for a real matrix $\mathbf{A} = (a_{ij})_{i,j}^{m,n}$, namely, $\|\mathbf{A}\|_F = \left(\sum_{i=1}^m \sum_{j=1}^n a_{ij}^2 \right)^{1/2}$. Hence, we have

$$\left\| \frac{1}{n} \sum_{i=1}^n \mathcal{M}_i \mathcal{M}_i^T - \Lambda_n \right\|_F = o_p(1). \quad (\text{A.9})$$

Equation (A.9) and condition (C6) imply that $\mathbf{u}^T \left(\sum_{i=1}^n \mathcal{M}_i \mathcal{M}_i^T / n \right) \mathbf{u}$ is asymptotically dominated by $\mathbf{u}^T \Lambda_n \mathbf{u}$. As $np_n (h_j^2 + (nh_j)^{-1/2}) = o(1)$ for $j = 1, \dots, p_n$, we can easily

prove that

$$\left\| \frac{1}{n} \sum_{i=1}^n (\hat{\mathcal{M}}_i - \mathcal{M}_i) \mathcal{M}_i^T \right\|_F \rightarrow 0, \left\| \frac{1}{n} \sum_{i=1}^n \mathcal{M}_i (\hat{\mathcal{M}}_i - \mathcal{M}_i)^T \right\|_F \rightarrow 0,$$

and

$$\left\| \frac{1}{n} \sum_{i=1}^n (\hat{\mathcal{M}}_i - \mathcal{M}_i) (\hat{\mathcal{M}}_i - \mathcal{M}_i)^T \right\|_F \rightarrow 0.$$

Thus $\frac{1}{n} \sum_{i=1}^n \hat{\mathcal{M}}_i \hat{\mathcal{M}}_i^T - \frac{1}{n} \sum_{i=1}^n \mathcal{M}_i \mathcal{M}_i^T = o_p(1)$. Let $\mathbb{M} = (\hat{\mathcal{M}}_1^T, \dots, \hat{\mathcal{M}}_n^T)^T$. Hence, we have

$$\begin{aligned} E(B_n | \mathbb{M}) &= \sum_{i=1}^n \int_0^{\alpha_n(v + \hat{\mathcal{M}}_i^T \mathbf{u})} [F_\eta\{t + \Delta_i\} - F_\eta\{\Delta_i\}] dt \\ &= \sum_{i=1}^n \int_0^{\alpha_n(v + \hat{\mathcal{M}}_i^T \mathbf{u})} [f_\eta(\Delta_i) t (1 + o(1))] dt \\ &= \frac{1}{2} \sum_{i=1}^n f_\eta(0) \left(\alpha_n (v + \hat{\mathcal{M}}_i^T \mathbf{u}) \right)^2 (1 + o(1)) \\ &= \frac{1}{2} f_\eta(0) n \alpha_n^2 (v^2 + \mathbf{u}^T \Psi_n \mathbf{u} + 2v \boldsymbol{\mu}_n^T \mathbf{u}) (1 + o_p(1)). \end{aligned} \tag{A.10}$$

Because $p_n^2/n \rightarrow 0$, we have $p_n \log n/n \rightarrow 0$, $\sqrt{p_n \log n} a_n \rightarrow 0$ as $n \rightarrow \infty$, with condition (C5) we have $\max_{1 \leq i \leq n} \left| \alpha_n (v + \hat{\mathcal{M}}_i^T \mathbf{u}) \right| \rightarrow 0$. By the Schwarz inequality, it is not difficult to show that

$$\begin{aligned} &Var(B_n | \mathbb{M}) \\ &\leq \sum_{i=1}^n E \left(\int_0^{\alpha_n(v + \hat{\mathcal{M}}_i^T \mathbf{u})} [I\{\eta_i \leq t + \Delta_i\} - I\{\eta_i \leq \Delta_i\}] dt | \mathbb{M} \right)^2 \\ &\leq \sum_{i=1}^n \alpha_n \left| v + \hat{\mathcal{M}}_i^T \mathbf{u} \right| \left| \int_0^{\alpha_n(v + \hat{\mathcal{M}}_i^T \mathbf{u})} E [I\{\eta_i \leq t + \Delta_i\} - I\{\eta_i \leq \Delta_i\}]^2 dt | \mathbb{M} \right| \\ &= o(1) \sum_{i=1}^n \left(\alpha_n (v + \hat{\mathcal{M}}_i^T \mathbf{u}) \right)^2 \\ &= o_p(p_n). \end{aligned} \tag{A.11}$$

From (A.9)–(A.11), we have

$$L_n = \frac{1}{2} f_\eta(0) n \alpha_n^2 (v^2 + \mathbf{u}^T \Psi_n \mathbf{u} + 2v \boldsymbol{\mu}_n^T \mathbf{u}) + o_p(n \alpha_n^2) \|\mathbf{u}\| + P_n. \tag{A.12}$$

For P_n , by the condition (C9) and Taylor's expansion for the penalty function, we have

$$\begin{aligned}
P_n &= n \sum_{j=1}^{s_n} [p_\lambda(|w_{oj} + \alpha_n u_j|) - p_\lambda(|w_{oj}|)] \\
&= n \sum_{j=1}^{s_n} \dot{p}_\lambda(|w_{oj}|) \operatorname{sgn}(w_{oj}) \alpha_n u_j + \frac{1}{2} n \sum_{j=1}^{s_n} \ddot{p}_\lambda(|w_{oj}|) (\alpha_n u_j)^2 \{1 + o(1)\} \\
&\leq n \alpha_n \sqrt{s_n} \max_{1 \leq j \leq s_n} |\dot{p}_\lambda(|w_{oj}|)| \|\mathbf{u}\| + n \alpha_n^2 \max_{1 \leq j \leq s_n} |\ddot{p}_\lambda(|w_{oj}|)| \|\mathbf{u}\|^2 \{1 + o(1)\} \\
&= O(n \alpha_n a_n \sqrt{s_n}) \|\mathbf{u}\| + O(n \alpha_n^2 b_n) \|\mathbf{u}\|^2 \\
&= O(n \alpha_n^2) \|\mathbf{u}\| + o(n \alpha_n^2) \|\mathbf{u}\|^2.
\end{aligned} \tag{A.13}$$

It follows from (A.12) and (A.13) that L_n in (A.7) is dominated by the positive quadratic term $\frac{1}{2} f_\eta(0) n \alpha_n^2 (v^2 + \mathbf{u}^T \boldsymbol{\Psi}_n \mathbf{u} + 2v \boldsymbol{\mu}_n^T \mathbf{u})$ when a sufficiently large C is chosen. Therefore, (A.6) holds and this completes the proof of Theorem 2 (i).

(ii) Let $\hat{\mathbf{w}}_n(1)$ and $\hat{\mathbf{w}}_n(2)$ be the estimators of $\mathbf{w}_n(1)$ and $\mathbf{w}_n(2)$, respectively, where $\mathbf{w}_n(1) = (w_1, \dots, w_{s_n})^T$ and $\mathbf{w}_n(2) = (w_{s_n+1}, \dots, w_{p_n})^T$. To prove Theorem 2(ii), it suffices to show that for any constant C and any given $(w_0, \mathbf{w}_n^T(1))^T$ satisfying $\|(w_0, \mathbf{w}_n^T(1))^T - (w_0, \mathbf{w}_o^T(1))^T\| = O_p(\alpha_n)$, where $\alpha_n = \sqrt{p_n} (n^{-1/2} + a_n)$, we have

$$\mathcal{Q}_n([w_0, \mathbf{w}_n^T(1), \mathbf{0}^T]^T) = \min_{\|\mathbf{w}_n(2)\| \leq C \alpha_n} \mathcal{Q}_n([w_0, \mathbf{w}_n^T(1), \mathbf{w}_n^T(2)]^T). \tag{A.14}$$

By (A.14), it is easy to prove that $\hat{\mathbf{w}}_n(2) = 0$.

To prove (A.14), it is sufficient to show that, with probability approaching one, for any $(p_n + 1)$ -dimensional vector $\mathbf{w}_n = (w_0, \mathbf{w}_n^T(1), \mathbf{w}_n^T(2))^T$ with $(w_0, \mathbf{w}_n^T(1))^T$ satisfying $\|(w_0, \mathbf{w}_n^T(1))^T - (w_0, \mathbf{w}_o^T(1))^T\| = O_p(\alpha_n)$ and for some small $\epsilon_n = C \alpha_n$ and $j = s_n + 1, \dots, p_n$,

$$\frac{\partial \mathcal{Q}_n(\mathbf{w}_n)}{\partial w_j} > 0, \quad 0 < w_j < \epsilon_n, \tag{A.15}$$

and

$$\frac{\partial \mathcal{Q}_n(\mathbf{w}_n)}{\partial w_j} < 0, \quad -\epsilon_n < w_j < 0, \tag{A.16}$$

Taking the first derivative of $\mathcal{Q}_n(\mathbf{w}_n)$ at any differentiable point $\mathbf{w}_n = (w_0, w_1, \dots, w_{p_n})^T$

with respect to $w_j, j = s_n + 1, \dots, p_n$, we have

$$\frac{\partial \mathcal{Q}_n(\mathbf{w}_n)}{\partial w_j} = - \sum_{i=1}^n \psi_\tau \left\{ Y_i - w_0 - \sum_{j=1}^{p_n} \hat{m}_j(X_{ij}) w_j \right\} \hat{m}_j(X_{ij}) + n \dot{p}_\lambda(|w_j|) \operatorname{sgn}(w_j)$$

for $j = 1, \dots, p_n$, where $\psi_\tau(u) = \tau - I(u < 0)$ and

$$\begin{aligned} & - \sum_{i=1}^n \psi_\tau \left\{ Y_i - w_0 - \sum_{j=1}^{p_n} \hat{m}_j(X_{ij}) w_j \right\} \hat{m}_j(X_{ij}) \\ &= - \sum_{i=1}^n \psi_\tau \{ \eta_i - (w_0 - w_{o0}) - \varsigma_i \} \hat{m}_j(X_{ij}) \\ &= \sum_{i=1}^n \{ I(\eta_i \leq 0) - \tau \} \hat{m}_j(X_{ij}) + \sum_{i=1}^n [I\{\eta_i \leq (w_0 - w_{o0}) + \varsigma_i\} - \{I(\eta_i \leq 0)\}] \hat{m}_j(X_{ij}) \\ &\triangleq I + II, \end{aligned}$$

where $\varsigma_i = \sum_{j=1}^{p_n} \{[\hat{m}_j(X_{ij}) - m_j(X_{ij})] w_{oj} + \hat{m}_j(X_{ij}) [w_j - w_{oj}]\}$. As in the proof of Theorem 2 (i) and Theorem 1, it is easy to prove that

$$I = O_p(\sqrt{np_n}) \text{ and } II = O_p(\sqrt{np_n}).$$

Hence, we have

$$\begin{aligned} \frac{\partial \mathcal{Q}_n(\mathbf{w}_n)}{\partial w_j} &= O_p(\sqrt{np_n}) + n \dot{p}_\lambda(|w_j|) \operatorname{sgn}(w_j) \\ &= n \lambda \left\{ O_p\left(\sqrt{p_n/n}/\lambda\right) + \lambda^{-1} \dot{p}_\lambda(|w_j|) \operatorname{sgn}(w_j) \right\}. \end{aligned} \tag{A.17}$$

Whereas $\liminf_{n \rightarrow \infty} \liminf_{|w_j| \rightarrow 0^+} \dot{p}_\lambda(|w_j|) / \lambda > 0$ by condition (C8) and $\sqrt{p_n/n} / \lambda \rightarrow 0$, the sign of the derivative is completely determined by that of w_j . Hence, we can show that (A.15) and (A.16) hold by using (A.17). This completes the proof of Theorem 2 (ii).

(iii) It can be shown easily that there exists a $(\hat{w}_0, \hat{\mathbf{w}}_n^T(1))^T$ in Theorem 2 that is a $\sqrt{n/p_n}$ consistent local minimizer of $\mathcal{Q}_n((w_0, \mathbf{w}_n^T(1), \mathbf{0}^T)^T)$, and satisfies the equations

$$n^{-1} \sum_{i=1}^n \hat{\mathcal{M}}_{1i} \psi_\tau \left\{ Y_i - \hat{w}_0 - \hat{\mathcal{M}}_{1i}^T \hat{\mathbf{w}}_n(1) \right\} = \mathbf{c}_n + \mathbf{\Gamma}_n(\hat{\mathbf{w}}_n(1) - \mathbf{w}_o(1)) \{1 + o_p(1)\}, \tag{A.18}$$

and

$$-n^{-1} \sum_{i=1}^n \psi_\tau \left\{ Y_i - \hat{w}_0 - \hat{\mathcal{M}}_{1i}^T \hat{\mathbf{w}}_n(1) \right\} = 0. \quad (\text{A.19})$$

where $\hat{\mathcal{M}}_{1i} = (\hat{m}_1(X_{i1}), \dots, \hat{m}_{s_n}(X_{is_n}))^T$. We can write

$$n^{-1} \sum_{i=1}^n \hat{\mathcal{M}}_{1i} \psi_\tau \left\{ Y_i - \hat{w}_0 - \hat{\mathcal{M}}_{1i}^T \hat{\mathbf{w}}_n(1) \right\} = -n^{-1/2} \mathbf{z}_{n1} + B_{n1} + B_{n2}, \quad (\text{A.20})$$

where

$$\begin{aligned} \mathbf{z}_{n1} &= n^{-1/2} \sum_{i=1}^n \hat{\mathcal{M}}_{1i} \{I(\eta_i \leq 0) - \tau\}, \\ B_{n1} &= n^{-1} \sum_{i=1}^n \hat{\mathcal{M}}_{1i} \{F_\eta(0) - F_\eta(\zeta_i)\}, \\ B_{n2} &= n^{-1} \sum_{i=1}^n \hat{\mathcal{M}}_{1i} \{[I(\eta_i \leq 0) - I(\eta_i \leq \zeta_i)] - [F_\eta(0) - F_\eta(\zeta_i)]\}, \end{aligned}$$

where $\zeta_i = (\hat{w}_0 - w_{o0}) + \sum_{j=1}^{s_n} [\hat{m}_j(X_{ij})(\hat{w}_j - w_{oj}) + (\hat{m}_j(X_{ij}) - m_j(X_{ij})) w_{oj}]$.

Taking Taylor's explanation for $F_\eta(\zeta_i)$ at 0 gives

$$\begin{aligned} B_{n1} &= n^{-1} \sum_{i=1}^n \hat{\mathcal{M}}_{1i} \{F_\eta(0) - F_\eta(\zeta_i)\} \\ &= -n^{-1} f_\eta(0) \sum_{i=1}^n \hat{\mathcal{M}}_{1i} \zeta_i \{1 + o_p(1)\} \\ &= -n^{-1} f_\eta(0) \left[\sum_{i=1}^n \hat{\mathcal{M}}_{1i} (\hat{w}_0 - w_{o0}) + \sum_{i=1}^n \hat{\mathcal{M}}_{1i} \hat{\mathcal{M}}_{1i}^T (\hat{\mathbf{w}}_n(1) - \mathbf{w}_o(1)) \right. \\ &\quad \left. + \sum_{i=1}^n \hat{\mathcal{M}}_{1i} (\hat{\mathcal{M}}_{1i} - \mathcal{M}_{1i})^T \mathbf{w}_o(1) \right] \{1 + o_p(1)\} \\ &= -n^{-1} f_\eta(0) [n \boldsymbol{\mu}_{n1} (\hat{w}_0 - w_{o0}) + n \boldsymbol{\Psi}_{n1} (\hat{\mathbf{w}}_n(1) - \mathbf{w}_o(1)) \\ &\quad + \sum_{i=1}^n \hat{\mathcal{M}}_{1i} (\hat{\mathcal{M}}_{1i} - \mathcal{M}_{1i})^T \mathbf{w}_o(1)] \{1 + o_p(1)\}. \end{aligned} \quad (\text{A.21})$$

By direct calculation of the mean and variance, we can show, as in Jiang et al. (2001), that $B_{n2} = o_p(\alpha_n)$. This combined with (A.20) and (A.21) leads to

$$\begin{aligned} &[f_\eta(0) \boldsymbol{\Psi}_{n1} + \boldsymbol{\Gamma}_n] (\hat{\mathbf{w}}_n(1) - \mathbf{w}_o(1)) \{1 + o_p(1)\} + n^{-1/2} \mathbf{z}_{n1} + \mathbf{c}_n \\ &= -n^{-1} f_\eta(0) \left[n \boldsymbol{\mu}_{n1} (\hat{w}_0 - w_{o0}) + \sum_{i=1}^n \hat{\mathcal{M}}_{1i} (\hat{\mathcal{M}}_{1i} - \mathcal{M}_{1i})^T \mathbf{w}_o(1) \right] \{1 + o_p(1)\}. \end{aligned} \quad (\text{A.22})$$

Similarly, (A.19) can be simplified as

$$\begin{aligned} n^{-1} \sum_{i=1}^n \{\tau - I(\eta_i \leq 0)\} &= f_\eta(0) (\hat{w}_0 - w_{o0}) \\ &+ n^{-1} f_\eta(0) \sum_{i=1}^n \left[\left(\hat{\mathcal{M}}_{1i} - \mathcal{M}_{1i} \right)^T \mathbf{w}_o(1) + \hat{\mathcal{M}}_{1i}^T (\hat{\mathbf{w}}_n(1) - \mathbf{w}_o(1)) \right] \{1 + o_p(1)\} \end{aligned} \quad (\text{A.23})$$

Solving (A.22) and (A.23), we obtain that

$$\begin{aligned} &(\hat{\mathbf{w}}_n(1) - \mathbf{w}_o(1)) \\ &= -[f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n]^{-1} n^{-1} f_\eta(0) \sum_{i=1}^n \hat{\mathcal{M}}_{1i} \left(\hat{\mathcal{M}}_{1i} - \mathcal{M}_{1i} \right)^T \mathbf{w}_o(1) \\ &\quad - [f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n]^{-1} n^{-1/2} \mathbf{z}_{n1} \\ &\quad + [f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n]^{-1} n^{-1} f_\eta(0) \boldsymbol{\mu}_{n1} \sum_{i=1}^n \left(\hat{\mathcal{M}}_{1i} - \mathcal{M}_{1i} \right)^T \mathbf{w}_o(1) \\ &\quad - [f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n]^{-1} \left(\mathbf{c}_n + n^{-1} \sum_{i=1}^n \{\tau - I(\eta_i \leq 0)\} \boldsymbol{\mu}_{n1} \right) + o_p(n^{-1/2}) \\ &= \Pi_{n1} + \Pi_{n2} + \Pi_{n3} + \Pi_{n4} + o_p(n^{-1/2}). \end{aligned} \quad (\text{A.24})$$

By Theorem 1, we have, uniformly for $x_k \in \mathcal{C}_k$,

$$\hat{m}_k(x_k) - m_k(x_k) = \frac{1}{nh_k} f_k^{-1}(x_k) f_{\eta_k}^{-1}(0) \sum_{t=1}^n K\left(\frac{X_{tk} - x_k}{h_k}\right) \psi_{\tau_k}(\eta_{tk}) \{1 + o_p(1)\}. \quad (\text{A.25})$$

Then, we have

$$\Pi_{n1} \xrightarrow{p} -[f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n]^{-1} n^{-1} f_\eta(0) \Pi_{n5}, \quad (\text{A.26})$$

where

$$\begin{aligned} \Pi_{n5} &= \left\{ \sum_{i=1}^n m_j(X_{ij}) \sum_{k=1}^{s_n} w_{ok} [\hat{m}_k(X_{ik}) - m_k(X_{ik})] \right\}_{j=1, \dots, s_n}^T \\ &= \left\{ \sum_{i=1}^n m_j(X_{ij}) \sum_{k=1}^{s_n} w_{ok} \right. \\ &\quad \times \left. \left[\frac{1}{nh_k} f_k^{-1}(X_{ik}) f_{\eta_k}^{-1}(0) \sum_{t=1}^n K\left(\frac{X_{tk} - X_{ik}}{h_k}\right) \psi_{\tau_k}(\eta_{tk}) \{1 + o_p(1)\} \right] \right\}_{j=1, \dots, s_n}^T \\ &= \left\{ \sum_{t=1}^n \sum_{k=1}^{s_n} w_{ok} \psi_{\tau_k}(\eta_{tk}) \right. \\ &\quad \times \left. \left[\frac{1}{nh_k} \sum_{i=1}^n m_j(X_{ij}) f_k^{-1}(X_{ik}) f_{\eta_k}^{-1}(0) K\left(\frac{X_{tk} - X_{ik}}{h_k}\right) \{1 + o_p(1)\} \right] \right\}_{j=1, \dots, s_n}^T, \end{aligned}$$

where $f_k(\cdot)$ is the marginal density function of X_{ik} . If $k = j$, we have

$$\frac{1}{nh_j} \sum_{i=1}^n m_j(X_{ij}) f_j^{-1}(X_{ij}) f_{\eta_j}^{-1}(0) K\left(\frac{X_{tj} - X_{ij}}{h_j}\right) = m_j(X_{tj}) + o_p(1). \quad (\text{A.27})$$

If $k \neq j$, we have

$$\frac{1}{nh_k} \sum_{i=1}^n m_j(X_{ij}) f_k^{-1}(X_{ik}) f_{\eta_k}^{-1}(0) K\left(\frac{X_{tk} - X_{ik}}{h_k}\right) = \beta_{jk}(X_{tk}) + o_p(1), \quad (\text{A.28})$$

where $\beta_{jk}(X_{tk}) = E(m_j(X_{tj}) | X_{tk})$. Then, by (A.27) and (A.28) and noting that $\beta_{jj}(X_{tj}) = m_j(X_{tj})$, we have

$$\begin{aligned} \Pi_{n5} &= \left(\sum_{t=1}^n \sum_{k=1}^{s_n} w_{ok} \psi_{\tau_k}(\eta_{tk}) \beta_{jk}(X_{tk}) \right)_{j=1, \dots, s_n}^T + O_p\left(ns_n \max_{1 \leq j \leq s_n} (h_j^2 + (nh_j)^{-1/2}) \right) \\ &= \left(\sum_{t=1}^n \tilde{\eta}_{t1}, \dots, \sum_{t=1}^n \tilde{\eta}_{ts_n} \right)^T + O_p\left(ns_n \max_{1 \leq j \leq s_n} (h_j^2 + (nh_j)^{-1/2}) \right), \end{aligned} \quad (\text{A.29})$$

where $\tilde{\eta}_{tj} = \sum_{k=1}^{s_n} w_{ok} \psi_{\tau_k}(\eta_{tk}) \beta_{jk}(X_{tk})$. By (A.26) and (A.29), we have

$$\begin{aligned} \Pi_{n1} &= -[f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n]^{-1} n^{-1} f_\eta(0) \left(\sum_{i=1}^n \tilde{\eta}_{i1}, \dots, \sum_{i=1}^n \tilde{\eta}_{is_n} \right)^T + \\ &\quad O_p\left(s_n \max_{1 \leq j \leq s_n} (h_j^2 + (nh_j)^{-1/2}) \right). \end{aligned} \quad (\text{A.30})$$

We next consider Π_{n2} . Observe that

$$\begin{aligned} \Pi_{n2} &= [f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n]^{-1} n^{-1} \sum_{i=1}^n \hat{\mathcal{M}}_{1i} \psi_\tau(\eta_i) \\ &= [f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n]^{-1} n^{-1} \\ &\quad \times \left[\sum_{i=1}^n \mathcal{M}_{1i} \psi_\tau(\eta_i) + \sum_{i=1}^n \left(\hat{\mathcal{M}}_{1i} - \mathcal{M}_{1i} \right) \psi_\tau(\eta_i) \right] \{1 + o_p(1)\}. \end{aligned}$$

We can show that the leading term of Π_{n2} is $[f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n]^{-1} n^{-1} \sum_{i=1}^n \mathcal{M}_{1i} \psi_\tau(\eta_i)$.

$$\Pi_{n2} = [f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n]^{-1} n^{-1} \left(\sum_{i=1}^n \bar{\eta}_{i1}, \dots, \sum_{i=1}^n \bar{\eta}_{is_n} \right)^T \{1 + o_p(1)\}, \quad (\text{A.31})$$

and $\Pi_{n3} = o_p(n^{-1/2})$. This combined with (A.30) and (A.31) leads to

$$\sqrt{n} [f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n] (\hat{\mathbf{w}}_n(1) - \mathbf{w}_o(1)) + \sqrt{n} \mathbf{c}_n = n^{-1/2} \sum_{i=1}^n (\boldsymbol{\xi}_i \psi_\tau(\eta_i) - f_\eta(0) \tilde{\boldsymbol{\eta}}_i).$$

where $\mathbf{e}_n^T \boldsymbol{\Upsilon}_{n1}^{-1/2} n^{-1/2} \sum_{i=1}^n (\boldsymbol{\xi}_i \psi_\tau(\eta_i) - f_\eta(0) \tilde{\boldsymbol{\eta}}_i) \xrightarrow{d} N(0, 1)$. It follows that

$$\sqrt{n} \mathbf{e}_n^T \boldsymbol{\Upsilon}_{n1}^{-1/2} (f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n) [(\hat{\mathbf{w}}_n(1) - \mathbf{w}_o(1)) + (f_\eta(0) \mathbf{G}_{n1} + \mathbf{\Gamma}_n)^{-1} \mathbf{c}_n] \xrightarrow{d} N(0, 1).$$

□

References

- Ando, T., Li, K.-C., 2014. A model-averaging approach for high-dimensional regression. *Journal of the American Statistical Association*, 109, 254–265.
- Chen, J., Li, D., Linton, O., Lu, Z., 2016. Semiparametric dynamic portfolio choice with multiple conditioning variables. *Journal of Econometrics*, 194, 309–318.
- Chen, J., Li, D., Linton, O., Lu, Z., 2018. Semiparametric ultra-high dimensional model averaging of nonlinear dynamic time series. *Journal of the American Statistical Association*, in press.
- Cheng, X., Hansen, B., 2015. Forecasting with factor-augmented regression: a frequentist model averaging approach. *Journal of Econometrics*, 186, 280–293.
- Claeskens, G., Hjort, N. L., 2008. *Model selection and model averaging*. Cambridge University Press.
- Fan, J., Peng, H., 2004. On non-concave penalized likelihood with diverging number of parameters. *The Annals of Statistics*, 32, 928–961.
- Guo, Y., Zhang, X., Wang, S., Zou, G., 2016. Model averaging based on leave-subject-out cross-validation. *Journal of Econometrics*, 192, 139–151.
- Hansen, B. E., 2007. Least squares model averaging. *Econometrica*, 75, 1175–1189.
- Hansen, B. E., 2008. Least squares forecast averaging. *Journal of Econometrics*, 146, 342–350.

- Hansen, B. E., Racine, J. S., 2012. Jackknife model averaging. *Journal of Econometrics*, 167, 38–46.
- Horowitz, J.L., 1998. Bootstrap methods for median regression models. *Econometrica*, 66, 1327–1351.
- Lian, H., 2012. A note on the consistency of Schwarz's criterion in linear quantile regression with the SCAD penalty. *Statistics and Probability Letters*, 82, 1224–1228.
- Hjort, N., Claeskens, G., 2003. Frequentist model average estimators. *Journal of the American Statistical Association*, 98, 879–899.
- Hoeting, J. A., Madigan, D., Raftery, A. E., Volinsky, C. T., 1999. Bayesian model averaging: a tutorial. *Statistical Science*, 14, 382–417.
- Horowitz, J. L., Lee, S., 2005. Nonparametric estimation of an additive quantile regression model. *Journal of the American Statistical Association*, 100, 1238–1249.
- Huang, T. and Li, J. 2018. Semiparametric model average prediction in panel data analysis. *Journal of Nonparametric Statistics*, 30, 125–144.
- Jiang, J., Zhao, Q., Hui, Y. V., 2001. Robust modelling of ARCH models. *Journal of Forecasting*, 20, 111–133.
- Johnson, R.W., 1996. Fitting percentage of body fat to simple body measurements. *Journal of Statistics Education*, 4, 1–8.
- Kai, B., Li, R., Zou, H., 2011. New efficient estimation and variable selection methods for semiparametric varying-coefficient partially linear models. *The Annals of Statistics*, 39, 305–332.
- Knight, K., 1998. Limiting distributions for L_1 regression estimators under general conditions. *The Annals of Statistics*, 26, 755–770.
- Li, D., Linton, O., Lu, Z., 2015. A flexible semiparametric forecasting model for time series. *Journal of Econometrics*, 187, 345–357.
- Li, J., Xia, X., Wong, W. K., Nott, D., 2018. Varying-coefficient semiparametric model averaging prediction. *Biometrics*, in press.

- Liang, H., Zou, G., Wan, A. T. K., Zhang, X., 2011. Optimal weight choice for frequentist model average estimators. *Journal of the American Statistical Association*, 106, 1053–1066.
- Liu, C.-A., 2015. Distribution theory of the least squares averaging estimator. *Journal of Econometrics*, 186, 142–159.
- Lv, J., Yang, H., Guo, C., Smoothing combined generalized estimating equations quantile partially linear additive models longitudinal data. *Computational Statistics*, 31, 1203–1234.
- Parzen, E., 1962. On estimation of a probability density function and model. *The Annals of Statistics*, 33, 1065–1076.
- Pollard, D., 1991. Asymptotics for least absolute deviation regression estimators. *Econometr. Theory*, 7, 186–199.
- Raftery, A., Madigan, D., Hoeting, J., 1997. Bayesian model averaging for linear regression models. *Journal of the American Statistical Association*, 92, 179–191.
- Tang, Y., Song, X., Wang, H. J., Zhu, Z., 2013. Variable selection in high-dimensional quantile varying coefficient models. *Journal of Multivariate Analysis*, 122, 115–132.
- Wan, T. K., Zhang, X., Zou, G., 2010. Least squares model averaging by Mallows criterion. *Journal of Econometrics*, 156, 277–283.
- Zhang, X., Wan, A. T. K., Zhou, S. Z., 2012. Focused information criteria, model selection, and model averaging in a tobit model with a nonzero threshold. *Journal of Business & Economic Statistics*, 30, 132–142.
- Zhang, X., Wan, A. T. K., Zou, G., 2013. Model averaging by Jackknife criterion in models with dependent data. *Journal of Econometrics*, 174, 82–94.
- Zhang, X., Zou, G., Liang, H., 2014. Model averaging and weight choice in linear mixed-effects models. *Biometrika*, 101, 205–218.
- Zhang, X., Yu, D., Zou, G., Liang, H., 2016. Optimal model averaging estimation for generalized linear models and generalized linear mixed-effects models. *Journal of American Statistical Association*, In Press.