

LINEABILITY AND SPACEABILITY: A NEW APPROACH

V.V. FÁVARO, D. PELLEGRINO, AND D. TOMAZ

ABSTRACT. The area of research called “Lineability” looks for linear structures inside exotic subsets of vector spaces. In the last decade lineability/spaceability has been investigated in rather general settings; for instance, Set Theory, Probability Theory, Functional Analysis, Measure Theory, etc. It is a common feeling that positive results on lineability/spaceability are quite natural (i.e., in general “large” subspaces can be found inside exotic subsets of vector spaces, in quite different settings) and more restrictive approaches have been persecuted. In this paper we introduce and explore a new approach in this direction.

1. INTRODUCTION AND PRELIMINARIES

The notions of lineability and spaceability were introduced by V. Gurariy and L. Quarta in [19] and by Aron, Gurariy, and Seoane-Sepúlveda in [5]. For a comprehensive background on lineability and spaceability we recommend the recent monograph [4]. This line of research investigates the presence of “large” linear subspaces in certain mathematical objects with *a priori* no linear structure. The properties of lineability and spaceability are studied in several contexts with interesting applications in different fields as norm-attaining operators, multilinear forms, homogeneous polynomials, sequence spaces, holomorphic mappings, absolutely summing operators, Peano curves, fractals, among others. See, for instance, [2, 6, 8, 11, 12, 13, 15, 16, 22], and the references therein.

Let E be a vector space and α be a cardinal number. A subset A of E is called α -lineable if $A \cup \{0\}$ contains an α -dimensional linear subspace of E ; if E is a topological vector space, it is called α -spaceable if $A \cup \{0\}$ contains a closed α -dimensional linear subspace of E . It is well known that, in general, positive results of lineability are rather usual and the feeling that “everything is lineable” is somewhat common. So, a more restrictive notion of lineability is in order. Our paper investigates a stronger notion of lineability/spaceability, which is rather more restrictive.

Let us establish some notations that will be carried out along this work. From now on all vector spaces are considered over a fixed scalar field \mathbb{K} which can be either \mathbb{R} or \mathbb{C} . We shall denote by $\mathfrak{c} = \text{card}(\mathbb{R})$ and $\aleph_0 = \text{card}(\mathbb{N})$. The following concepts are more restrictive notions of lineability/spaceability inspired by some ideas from [21]:

Definition 1.1. Let α, β, λ be cardinal numbers and V be a vector space, with $\dim V = \lambda$ and $\alpha < \beta \leq \lambda$. A set $A \subset V$ is:

(i) (α, β) -lineable if it is α -lineable and for every subspace $W_\alpha \subset V$ with $W_\alpha \subset A \cup \{0\}$ and $\dim W_\alpha = \alpha$, there is a subspace $W_\beta \subset V$ with $\dim W_\beta = \beta$ and $W_\alpha \subset W_\beta \subset A \cup \{0\}$.

(ii) (α, β) -spaceable if it is α -lineable and for every subspace $W_\alpha \subset V$ with $W_\alpha \subset A \cup \{0\}$ and $\dim W_\alpha = \alpha$, there is a closed subspace $W_\beta \subset V$ with $\dim W_\beta = \beta$ and $W_\alpha \subset W_\beta \subset A \cup \{0\}$.

The original notion of lineability (or spaceability) is just the case $\alpha = 0$. Let α_1, α_2 be cardinal numbers with $\alpha_1 < \alpha_2 \leq \beta$. We start by showing that β -lineability does not imply

Key words and phrases. Cardinal numbers, lineability, spaceability.

Daniel Tomaz is supported by Capes.

Daniel Pellegrino is supported by CNPq.

Vinicius Fávoro is supported by FAPEMIG Grant APQ-03181-16; and CNPq Grant 310500/2017-6.

2010 Mathematics Subject Classification: 15A03, 46A16; 46A45.

(α, β) -lineability and *vice versa*. To provide examples, let us recall that for $p > 0$, ℓ_p denotes the Banach space (p -Banach space if $p < 1$) of the sequences $(x_j)_{j=1}^\infty$ such that

$$\left\| (x_j)_{j=1}^\infty \right\|_p = \left(\sum_{j=1}^\infty |x_j|^p \right)^{1/p} < \infty.$$

We denote by e_j the canonical vector $(0, \dots, 0, 1, 0, 0, \dots)$ with 1 in the j -th coordinate, for each $j \in \mathbb{N}$.

More generally, let $p \in [1, \infty)$ and Γ be an abstract nonempty set. We denote by $\ell_p(\Gamma)$ the vector space of all functions $f: \Gamma \rightarrow \mathbb{K}$ such that $\sum_{\gamma \in \Gamma} |f(\gamma)|^p < \infty$, which becomes a Banach space with the norm

$$\|f\|_p = \left(\sum_{\gamma \in \Gamma} |f(\gamma)|^p \right)^{1/p},$$

where the sum is defined by

$$\sum_{\gamma \in \Gamma} |f(\gamma)|^p = \sup \left\{ \sum_{\gamma \in F} |f(\gamma)|^p : F \text{ is a finite subset of } \Gamma \right\}.$$

We also denote the elements of the canonical generalized Schauder basis of $\ell_p(\Gamma)$ by e_i , $i \in \Gamma$.

It is clear that, when $\Gamma = \mathbb{N}$, we have $\ell_p(\mathbb{N}) = \ell_p$. For details about the space $\ell_p(\Gamma)$ and related results we refer to [17].

Example 1.2. Let $n \in \mathbb{N}$ and consider the following subset of ℓ_p :

$$A = \text{span} \{e_1, e_2, \dots, e_n\} \cup \left\{ (x_j)_{j=1}^\infty \in \ell_p : x_1 = x_2 = \dots = x_n = 0 \right\}.$$

Thus A is $(n+1, \mathfrak{c})$ -lineable, but it is not (n, \mathfrak{c}) -lineable.

Example 1.3. Let $\mathcal{B} = \{e_i : i \in \Gamma\}$ be the canonical generalized Schauder basis in $\ell_p(\Gamma)$ with $\text{card}(\Gamma) = 2^{\aleph_1}$. Choose $i_0, i_1 \in \Gamma$. We can easily write

$$\mathcal{B} - \{e_{i_0}, e_{i_1}\} = \bigcup_{(\lambda, \mu) \in \mathbb{R}^2} \mathcal{A}_{(\lambda, \mu)}$$

as a pairwise disjoint union, with

$$\text{card}(\mathcal{A}_{(\lambda, \mu)}) = 2^{\aleph_1}$$

for all $(\lambda, \mu) \in \mathbb{R}^2$. Let

$$A = \bigcup_{(\lambda, \mu) \in \mathbb{R}^2} \text{span}(\{\lambda e_{i_0} + \mu e_{i_1}\} \cup \mathcal{A}_{(\lambda, \mu)}).$$

Note that there is no vector space $W \subset \ell_p(\Gamma)$ with $\dim W = 2^{\aleph_1}$ and

$$\text{span}\{e_{i_0}, e_{i_1}\} \subset W \subset A \cup \{0\}.$$

In fact, if such W exists, since

$$e_{i_0} \in W \subset A \cup \{0\},$$

by the very definition of A we conclude that

$$(1.1) \quad W \subset \bigcup_{\substack{\lambda \in \mathbb{R} \\ \lambda \neq 0}} \text{span}(\{e_{i_0}\} \cup \mathcal{A}_{(\lambda, 0)}) \bigcup \text{span}\{e_{i_0}, e_{i_1}\}.$$

Let us prove it: If the inclusion above was false, then there would exist $w \in W$ with

$$w \notin \bigcup_{\substack{\lambda \in \mathbb{R} \\ \lambda \neq 0}} \text{span}(\{e_{i_0}\} \cup \mathcal{A}_{(\lambda, 0)}) \bigcup \text{span}\{e_{i_0}, e_{i_1}\},$$

i.e.,

$$w \in \bigcup_{\substack{(\lambda, \mu) \in \mathbb{R}^2 \\ \mu \neq 0}} \text{span}(\{\lambda e_{i_0} + \mu e_{i_1}\} \cup \mathcal{A}_{(\lambda, \mu)}) \bigcup \text{span}(\mathcal{A}_{(0,0)}) \text{ and } w \notin \text{span}\{e_{i_0}, e_{i_1}\}.$$

Thus

$$w = r(\lambda e_{i_0} + \mu e_{i_1}) + v$$

with $v \in \text{span}(\mathcal{A}_{(\lambda, \mu)})$, $v \neq 0$, and $r \in \mathbb{R}$, where $\mu \neq 0$ or $(\lambda, \mu) = (0, 0)$. First suppose that $\mu \neq 0$. Since W is a vector space containing e_{i_0} and e_{i_1} , we have

$$w + (-r\lambda + \lambda)e_{i_0} + (-r\mu + \mu + s)e_{i_1} \in W,$$

i.e.,

$$\lambda e_{i_0} + (\mu + s)e_{i_1} + v \in W$$

for any choice of $s \in \mathbb{R}$. Let us choose $s \neq -\mu$ such that

$$(\lambda, \mu) \text{ and } (\lambda, \mu + s) \text{ are linearly independent.}$$

Then,

$$\lambda e_{i_0} + (\mu + s)e_{i_1} \neq 0,$$

and since $v \in \text{span}(\mathcal{A}_{(\lambda, \mu)})$ we have that

$$v \notin \bigcup_{(\alpha, \beta) \neq (\lambda, \mu)} \text{span}(\mathcal{A}_{(\alpha, \beta)}).$$

Now it follows from the definition of A that

$$\lambda e_{i_0} + (\mu + s)e_{i_1} + v \notin A,$$

which is a contradiction.

Now, let us consider the case $(\lambda, \mu) = (0, 0)$. In this case

$$w \in \text{span}(\mathcal{A}_{(0,0)}) \text{ and so } w \notin \bigcup_{(\alpha, \beta) \neq (0,0)} \text{span}(\mathcal{A}_{(\alpha, \beta)}).$$

Since $e_{i_0}, e_{i_1} \in W$, for any choice of $(\alpha, \beta) \neq (0, 0)$ we obtain

$$\alpha e_{i_0} + \beta e_{i_1} + w \in W.$$

By the definition of A we have

$$\alpha e_{i_0} + \beta e_{i_1} + w \notin A,$$

a contradiction.

Analogously, we conclude that

$$(1.2) \quad W \subset \bigcup_{\mu \in \mathbb{R} \setminus \{0\}} \text{span}(\{e_{i_1}\} \cup \mathcal{A}_{(0, \mu)}) \bigcup \text{span}\{e_{i_0}, e_{i_1}\}.$$

Thus, from (1.1) and (1.2),

$$W \subset \left(\bigcup_{\lambda \in \mathbb{R} \setminus \{0\}} \text{span}(\{e_{i_0}\} \cup \mathcal{A}_{(\lambda, 0)}) \right) \cap \left(\bigcup_{\mu \in \mathbb{R} \setminus \{0\}} \text{span}(\{e_{i_1}\} \cup \mathcal{A}_{(0, \mu)}) \right) \bigcup \text{span}\{e_{i_0}, e_{i_1}\} = \text{span}\{e_{i_0}, e_{i_1}\},$$

a contradiction. Thus A is not $(2, 2^{\aleph_1})$ -lineable, but it is clear that A is $(1, 2^{\aleph_1})$ -lineable.

Obviously, every known lineability/spaceability result can be investigated in this more restrictive setting. In this paper we develop a technique to characterize the (α, \mathfrak{c}) -spaceability of $\ell_p \setminus \bigcup_{0 < q < p} \ell_q$ for all $\alpha < \mathfrak{c}$ that illustrates the technicalities arisen by this new approach. We also propose open problems in the final section.

2. CHARACTERIZATION OF (α, \mathfrak{c}) -SPACEABILITY IN SEQUENCE SPACES

It is well known that $\ell_p \setminus \bigcup_{0 < q < p} \ell_q$ is \mathfrak{c} -spaceable (see [11]). In this section we characterize

(α, \mathfrak{c}) -spaceability in this setting. For the sake of completeness, we begin by presenting a simple explicit example of a vector inside $\ell_p \setminus \bigcup_{0 < q < p} \ell_q$ that we were not able to find in the literature.

Let

$$(2.1) \quad \mathbb{N} = \bigcup_{j=1}^{\infty} \mathbb{N}_j,$$

with $\mathbb{N}_i \cap \mathbb{N}_j = \emptyset$ whenever $i \neq j$ and $\text{card}(\mathbb{N}_i) = \aleph_0$ for all i . Denote

$$\mathbb{N}_j = \{j_1, j_2, \dots\}$$

with $j_r < j_s$ whenever $r < s$, and define

$$x_j^{(k)} = \begin{cases} 0, & \text{if } j \notin \mathbb{N}_k \\ r^{-1/(p-k-1)}, & \text{if } j = k_r \in \mathbb{N}_k. \end{cases}$$

Note that $(x_j^{(k)})_{j=1}^{\infty} \in \ell_p \setminus \ell_{p-\frac{1}{k}}$. Then, the sequence $(y_j)_{j=1}^{\infty}$ defined by

$$y_j = \frac{x_j^{(k)}}{2^k \left\| (x_l^{(k)})_{l=1}^{\infty} \right\|_p},$$

where $j \in \mathbb{N}_k$, belongs to $\ell_p \setminus \bigcup_{0 < q < p} \ell_q$. In fact,

$$\begin{aligned} \sum_{j=1}^{\infty} |y_j|^p &= \sum_{k=1}^{\infty} \sum_{j \in \mathbb{N}_k} |y_j|^p \\ &= \sum_{k=1}^{\infty} \sum_{j \in \mathbb{N}_k} \left(\frac{|x_j^{(k)}|}{2^k \left\| (x_l^{(k)})_{l=1}^{\infty} \right\|_p} \right)^p \\ &= \sum_{k=1}^{\infty} \frac{1}{2^{kp} \left\| (x_l^{(k)})_{l=1}^{\infty} \right\|_p^p} \sum_{j \in \mathbb{N}_k} |x_j^{(k)}|^p \\ &= \sum_{k=1}^{\infty} \frac{1}{2^{kp}} < \infty, \end{aligned}$$

and a similar estimate shows that

$$\sum_{j=1}^{\infty} |y_j|^q = \infty$$

for all $0 < q < p$.

Lemma 2.1. *Let $p > 0$ and x_1, \dots, x_n be linearly independent vectors of ℓ_p . There exists $n_0 \in \mathbb{N}$ such that the first n_0 coordinates of x_1, \dots, x_n form a linearly independent set of \mathbb{K}^{n_0} .*

Proof. Suppose that, for each $j \in \mathbb{N}$, there exist $a_{1j}, \dots, a_{nj} \in \mathbb{K}$, not all equal to zero, such that

$$(2.2) \quad a_{1j}x_1 + \dots + a_{nj}x_n = (0, \dots, 0, \lambda_{j+1}, \lambda_{j+2}, \dots)$$

It is plain that we may suppose $\|\alpha_j\|_1 = 1$, where $\alpha_j = (a_{1j}, \dots, a_{nj}) \in \mathbb{K}^n$, for all $j \in \mathbb{N}$. Since $(\alpha_j)_{j=1}^\infty$ is bounded, there is a convergent subsequence $\alpha_{j_k} \rightarrow \alpha = (a_1, \dots, a_n) \in \mathbb{K}^n$. Since $\|\alpha_j\|_1 = 1$, it follows that $\alpha \neq 0$. Then

$$(2.3) \quad a_{1j_k}x_1 + \dots + a_{nj_k}x_n \rightarrow a_1x_1 + \dots + a_nx_n, \text{ when } k \rightarrow \infty.$$

On the other hand, if $\pi_m: \ell_p \rightarrow \mathbb{K}$ denotes the m -th canonical projection, then it follows from (2.2) that

$$\pi_m(a_{1j_k}x_1 + \dots + a_{nj_k}x_n) \rightarrow 0, \text{ when } k \rightarrow \infty, \text{ for every } m \in \mathbb{N}.$$

Since convergence in (2.3) implies coordinatewise convergence, it follows that $a_1x_1 + \dots + a_nx_n = (0, 0, \dots)$, which is a contradiction because x_1, \dots, x_n are linearly independent. \square

Theorem 2.2. *For all $p > 0$ the set $\ell_p \setminus \bigcup_{0 < q < p} \ell_q$ is (α, \mathfrak{c}) -spaceable in ℓ_p if, and only if, $\alpha < \aleph_0$.*

Proof. If $\alpha = \aleph_0$, then the following example shows that $\ell_p \setminus \bigcup_{0 < q < p} \ell_q$ is not (\aleph_0, \mathfrak{c}) -spaceable:
Split

$$\mathbb{N} = \bigcup_{j=1}^{\infty} \mathbb{N}_j$$

as in (2.1). Let

$$W := \text{span}\{x^{(j)} : j \in \mathbb{N}\}$$

with

- $x^{(1)} = \left(1, \left(x_j^{(1)}\right)_{j>1}\right)$ where

$$\begin{cases} x_j^{(1)} = 0, & \text{if } j \notin \mathbb{N}_1 \\ \left(x_j^{(1)}\right)_{j \in \mathbb{N}_1(j>1)} \in \ell_p \setminus \bigcup_{0 < q < p} \ell_q \end{cases}$$

and

$$\left\| \left(x_j^{(1)}\right)_{j \in \mathbb{N}_1(j>1)} \right\|_p = 2^{-1}.$$

- $x^{(2)} = \left(1, \left(x_j^{(2)}\right)_{j>1}\right)$ where

$$\begin{cases} x_j^{(2)} = 0, & \text{if } j \notin \mathbb{N}_2 \\ \left(x_j^{(2)}\right)_{j \in \mathbb{N}_2(j>1)} \in \ell_p \setminus \bigcup_{0 < q < p} \ell_q \end{cases}$$

and

$$\left\| \left(x_j^{(2)}\right)_{j \in \mathbb{N}_2(j>1)} \right\|_p = 2^{-2},$$

and so on. It is obvious that $(W \setminus \{0\}) \subset \ell_p \setminus \bigcup_{0 < q < p} \ell_q$ and $\dim(W) = \aleph_0$ and there is no closed subspace W_1 of ℓ_p such that $W \subset W_1$ and $(W_1 \setminus \{0\}) \subset \ell_p \setminus \bigcup_{0 < q < p} \ell_q$, because

$$\lim_{k \rightarrow \infty} \left\| \left(x_j^{(k)}\right)_{j \in \mathbb{N}} - e_1 \right\|_p = \lim_{k \rightarrow \infty} \left\| \left(0, x_j^{(k)}\right)_{j \in \mathbb{N}_k(j>1)} \right\|_p = \lim_{k \rightarrow \infty} \frac{1}{2^k} = 0.$$

Now we prove the theorem for $\alpha = 3$. The general case $\alpha \in \mathbb{N}$ is easily adapted from this case.

Let $x = (x_j)_{j=1}^\infty$, $y = (y_j)_{j=1}^\infty$ and $z = (z_j)_{j=1}^\infty$ be linearly independent and

$$W := \text{span}\{x, y, z\} \subset \left(\ell_p \setminus \bigcup_{0 < q < p} \ell_q \right) \cup \{0\}$$

be a 3-dimensional subspace of ℓ_p . Since $(x_j)_{j=1}^\infty, (y_j)_{j=1}^\infty$ and $(z_j)_{j=1}^\infty$ are linearly independent, Lemma 2.1 assures that there is a natural number n_0 such that the first n_0 coordinates of $(x_j)_{j=1}^\infty, (y_j)_{j=1}^\infty$ and $(z_j)_{j=1}^\infty$ form a linearly independent set of \mathbb{K}^{n_0} .

It is easy to pick an infinite set $\mathbb{N}_1 := \{\alpha_1 < \alpha_2 < \dots\}$ of positive integers such that

$$\left\{ \begin{array}{l} \sum_{j \notin \mathbb{N}_1} x_j e_j \in \ell_p \setminus \bigcup_{0 < q < p} \ell_q, \\ \sum_{j \notin \mathbb{N}_1} y_j e_j \in \ell_p \setminus \bigcup_{0 < q < p} \ell_q, \\ \sum_{j \notin \mathbb{N}_1} z_j e_j \in \ell_p \setminus \bigcup_{0 < q < p} \ell_q \end{array} \right.$$

and such that

$$(2.4) \quad \max\{|x_{\alpha_l}|, |y_{\alpha_l}|, |z_{\alpha_l}|\} < \frac{1}{2^l}$$

for all $l \in \mathbb{N}$. Of course, $\mathbb{N} \setminus \mathbb{N}_1$ is also infinite. Let

$$(2.5) \quad \mathbb{O} := \mathbb{N}_1 \setminus \{1, \dots, n_0\},$$

and note that

$$(2.6) \quad \left\{ \begin{array}{l} \sum_{j \notin \mathbb{O}} x_j e_j \in \ell_p \setminus \bigcup_{0 < q < p} \ell_q, \\ \sum_{j \notin \mathbb{O}} y_j e_j \in \ell_p \setminus \bigcup_{0 < q < p} \ell_q, \\ \sum_{j \notin \mathbb{O}} z_j e_j \in \ell_p \setminus \bigcup_{0 < q < p} \ell_q. \end{array} \right.$$

Split

$$\mathbb{O} = \bigcup_{i=1}^{\infty} \mathbb{O}_i$$

with $\mathbb{O}_i := \{i_1 < i_2 < \dots\}$ and $\mathbb{O}_i \cap \mathbb{O}_j = \emptyset$ for all $i \neq j$. It is plain that $\mathbb{N} \setminus \mathbb{O}$ is infinite and let $f : \mathbb{N} \rightarrow \mathbb{N}$ be injective such that

$$\mathbb{N} \setminus \mathbb{O} := \{f(1), f(2), \dots\}$$

and define, for all i , the vector

$$\varepsilon_i = \sum_{j=1}^{\infty} x_{f(j)} e_{i_j} \in \mathbb{K}^{\mathbb{N}}.$$

It is obvious that $\varepsilon_i \in \ell_p$ for all i . Define $\tilde{p} = 1$ if $p \geq 1$ and $\tilde{p} = p$ if $0 < p < 1$. For $(a_i)_{i=1}^\infty \in \ell_{\tilde{p}}$,

$$\sum_{i=1}^{\infty} \|a_i \varepsilon_i\|_{\tilde{p}}^{\tilde{p}} = \sum_{i=1}^{\infty} |a_i|^{\tilde{p}} \|\varepsilon_i\|_{\tilde{p}}^{\tilde{p}} \leq \|x\|_{\tilde{p}}^{\tilde{p}} \sum_{i=1}^{\infty} |a_i|^{\tilde{p}} = \|x\|_{\tilde{p}}^{\tilde{p}} \|(a_i)_{i=1}^\infty\|_{\tilde{p}}^{\tilde{p}} < \infty.$$

Thus $\sum_{i=1}^{\infty} \|a_i \varepsilon_i\|_{\tilde{p}}^{\tilde{p}} < \infty$ and thus the series $\sum_{i=1}^{\infty} a_i \varepsilon_i$ converges in ℓ_p . Hence, the operator

$$\left\{ \begin{array}{l} T : \ell_{\tilde{p}} \longrightarrow \ell_p, \\ T((a_i)_{i=0}^\infty) = a_0 x + a_1 y + a_2 z + \sum_{i=3}^{\infty} a_i \varepsilon_i \end{array} \right.$$

is well defined. It is easy to see that T is linear and injective. In fact, if

$$T((a_i)_{i=0}^\infty) = 0$$

then

$$a_0x + a_1y + a_2z + \sum_{i=3}^{\infty} a_i \varepsilon_i = 0.$$

In particular, since the first n_0 coordinates of each ε_i are all zero, we have

$$a_0(x_j)_{j=1}^{n_0} + a_1(y_j)_{j=1}^{n_0} + a_2(z_j)_{j=1}^{n_0} = 0$$

and then

$$(2.7) \quad a_0 = a_1 = a_2 = 0.$$

From (2.7) and since the supports of the vectors ε_i are disjoint, we have $a_i = 0$ for all i .

Note that $T((a_i)_{i=0}^\infty) \notin \ell_q$ for all $0 < q < p$ and $(a_i)_{i=0}^\infty \neq 0$. In fact, let i_0 be such that $a_{i_0} \neq 0$. If $a_0 = a_1 = a_2 = 0$ we have

$$\|T((a_i)_{i=0}^\infty)\|_q = \left\| \sum_{i=3}^{\infty} a_i \varepsilon_i \right\|_q \geq |a_{i_0}| \|\varepsilon_{i_0}\|_q = \infty.$$

If $a_i \neq 0$ for some $i = 0, 1, 2$, since the coordinates of ε_i belonging to $\mathbb{N} \setminus \mathbb{O}$ are zero, we have

$$\|T((a_i)_{i=0}^\infty)\|_q = \left\| a_0x + a_1y + a_2z + \sum_{i=3}^{\infty} a_i \varepsilon_i \right\|_q \geq \left\| (a_0x_j + a_1y_j + a_2z_j)_{j \notin \mathbb{O}} \right\|_q.$$

Since

$$\infty = \left\| (a_0x_j + a_1y_j + a_2z_j)_{j \in \mathbb{N}} \right\|_q^q = \left\| (a_0x_j + a_1y_j + a_2z_j)_{j \in \mathbb{O}} \right\|_q^q + \left\| (a_0x_j + a_1y_j + a_2z_j)_{j \notin \mathbb{O}} \right\|_q^q$$

and by (2.4) and (2.5) we have

$$\left\| (a_0x_j + a_1y_j + a_2z_j)_{j \in \mathbb{O}} \right\|_q < \infty,$$

then

$$(2.8) \quad \left\| (a_0x_j + a_1y_j + a_2z_j)_{j \notin \mathbb{O}} \right\|_q = \infty.$$

Thus $T((a_i)_{i=0}^\infty) \notin \ell_q$ for all $0 < q < p$. We have thus just proved the $(3, \mathbf{c})$ -lineability (and of course the (n, \mathbf{c}) -lineability for $n \in \mathbb{N}$ is analogous). Now let us prove the spaceability.

Of course, $\overline{T(\ell_{\bar{p}})}$ is a closed infinite-dimensional subspace of ℓ_p . We just have to show that

$$\overline{T(\ell_{\bar{p}})} \setminus \{0\} \subset \ell_p \setminus \bigcup_{0 < q < p} \ell_q.$$

Let $w = (w_n)_{n=1}^\infty \in \overline{T(\ell_{\bar{p}})}$, $w \neq 0$. There are sequences $(a_i^{(k)})_{i=0}^\infty \in \ell_{\bar{p}}$, $k \in \mathbb{N}$, such that $w = \lim_{k \rightarrow \infty} T\left(\left(a_i^{(k)}\right)_{i=0}^\infty\right)$ in ℓ_p . Note that, for each $k \in \mathbb{N}$,

$$\begin{aligned} T\left(\left(a_i^{(k)}\right)_{i=0}^\infty\right) &= a_0^{(k)}x + a_1^{(k)}y + a_2^{(k)}z + \sum_{i=3}^{\infty} a_i^{(k)}\varepsilon_i \\ &= a_0^{(k)}x + a_1^{(k)}y + a_2^{(k)}z + \sum_{i=3}^{\infty} a_i^{(k)} \sum_{j=1}^{\infty} x_{f(j)}e_{i_j} \\ &= a_0^{(k)}x + a_1^{(k)}y + a_2^{(k)}z + \sum_{i=3}^{\infty} \sum_{j=1}^{\infty} a_i^{(k)} x_{f(j)}e_{i_j}. \end{aligned}$$

Since the coordinates of ε_i belonging to $\mathbb{N} \setminus \mathbb{O}$ are zero, then the coordinates of $\sum_{i=3}^{\infty} a_i^{(k)} \varepsilon_i$ belonging to $\mathbb{N} \setminus \mathbb{O}$ are also zero. Thus, the coordinates of $T\left(\left(a_i^{(k)}\right)_{i=0}^{\infty}\right)$ belonging to $\mathbb{N} \setminus \mathbb{O}$ are the respective coordinates of $a_0^{(k)}x + a_1^{(k)}y + a_2^{(k)}z$ for all k . Hence the limit

$$\lim_{k \rightarrow \infty} \left(a_0^{(k)} x_j + a_1^{(k)} y_j + a_2^{(k)} z_j \right)_{j \in \mathbb{N} \setminus \mathbb{O}}$$

exists in ℓ_p . Since $\{(x_j)_{j \in \mathbb{N} \setminus \mathbb{O}}, (y_j)_{j \in \mathbb{N} \setminus \mathbb{O}}, (z_j)_{j \in \mathbb{N} \setminus \mathbb{O}}\}$ is linearly independent (because $\{1, \dots, n_0\} \subset \mathbb{N} \setminus \mathbb{O}$), it is obvious that this limit can be written in an unique form as

$$(ax_j + by_j + cz_j)_{j \in \mathbb{N} \setminus \mathbb{O}}.$$

Applying a linear functional that sends x_j in 1 and y_j and z_j in 0 we obtain that

$$\lim_{k \rightarrow \infty} \left(a_0^{(k)} \right) = a.$$

Analogously we obtain

$$\lim_{k \rightarrow \infty} \left(a_1^{(k)} \right) = b \text{ and } \lim_{k \rightarrow \infty} \left(a_2^{(k)} \right) = c.$$

Thus

$$w_j = \left(\lim_{k \rightarrow \infty} a_0^{(k)} \right) x_j + \left(\lim_{k \rightarrow \infty} a_1^{(k)} \right) y_j + \left(\lim_{k \rightarrow \infty} a_2^{(k)} \right) z_j$$

for all positive integers j in $\mathbb{N} \setminus \mathbb{O}$. To finish the proof consider the following cases:

- First case: $\lim_{k \rightarrow \infty} a_i^{(k)} \neq 0$, for some $i = 0, 1, 2$.

This case is simple because

$$\|w\|_q^q \geq \sum_{j \in \mathbb{N} \setminus \mathbb{O}} |a_0 x_j + a_1 y_j + a_2 z_j|^q \stackrel{(2.8)}{=} \infty$$

and the proof is done.

- Second case: $\lim_{k \rightarrow \infty} a_i^{(k)} = 0$ for all $i = 0, 1, 2$.

In this case $w_j = 0$ for all j in $\mathbb{N} \setminus \mathbb{O}$. Since $w \neq 0$, for all there is $r \in \mathbb{O}$ such that $w_r \neq 0$.

Since $\mathbb{O} = \bigcup_{j=1}^{\infty} \mathbb{O}_j$, there are (unique) $m, t \in \mathbb{N}$ such that $e_{m_t} = e_r$. Thus, for each $k \in \mathbb{N}$, the r -th coordinate of $T\left(\left(a_i^{(k)}\right)_{i=0}^{\infty}\right)$ is the number $a_0^{(k)}x_r + a_1^{(k)}y_r + a_2^{(k)}z_r + a_m^{(k)}x_{f(t)}$. So

$$0 \neq w_r = \lim_{k \rightarrow \infty} \left(a_0^{(k)} x_r + a_1^{(k)} y_r + a_2^{(k)} z_r + a_m^{(k)} x_{f(t)} \right) = \lim_{k \rightarrow \infty} a_m^{(k)} x_{f(t)} = x_{f(t)} \cdot \lim_{k \rightarrow \infty} a_m^{(k)}.$$

It follows that $x_{f(t)} \neq 0$. Hence $\lim_{k \rightarrow \infty} |a_m^{(k)}| = \frac{|w_r|}{|x_{f(t)}|} \neq 0$. For $j, k \in \mathbb{N}$, the m_j -th coordinate of $T\left(\left(a_i^{(k)}\right)_{i=0}^{\infty}\right)$ is

$$a_0^{(k)} x_{m_j} + a_1^{(k)} y_{m_j} + a_2^{(k)} z_{m_j} + a_m^{(k)} x_{f(j)}.$$

Defining $\alpha_m = \frac{|w_r|}{|x_{f(t)}|}$ we have

$$\lim_{k \rightarrow \infty} \left| a_0^{(k)} x_{m_j} + a_1^{(k)} y_{m_j} + a_2^{(k)} z_{m_j} + a_m^{(k)} x_{f(j)} \right| = \lim_{k \rightarrow \infty} |a_m^{(k)} x_{f(j)}| = |x_{f(j)}| \cdot \lim_{k \rightarrow \infty} |a_m^{(k)}| = \alpha_m |x_{f(j)}|$$

for every $j \in \mathbb{N}$. On the other hand, coordinatewise convergence gives us

$$\lim_{k \rightarrow \infty} \left| a_0^{(k)} x_{m_j} + a_1^{(k)} y_{m_j} + a_2^{(k)} z_{m_j} + a_m^{(k)} x_{f(j)} \right| = |w_{m_j}|,$$

so $|w_{m_j}| = \alpha_m |x_{f(j)}|$ for each $j \in \mathbb{N}$. Hence

$$\|w\|_q^q = \sum_{n=1}^{\infty} |w_n|^q \geq \sum_{j=1}^{\infty} |w_{m_j}|^q = \sum_{j=1}^{\infty} \alpha_m^q \cdot |x_{f(j)}|^q = \alpha_m^q \cdot \left\| (x_{f(j)})_{j=1}^{\infty} \right\|_q^q = \alpha_m^q \cdot \left\| (x_j)_{j \in \mathbb{N} \setminus \mathbb{O}} \right\|_q^q \stackrel{(2.6)}{=} \infty.$$

Therefore $w \notin \bigcup_{0 < q < p} \ell_q$, so $\overline{T(\ell_p)} \setminus \{0\} \subseteq \ell_p \setminus \bigcup_{0 < q < p} \ell_q$. □

3. SOME OPEN PROBLEMS

Of course, any positive result of lineability and/or spaceability is a potential problem to be investigated in our more general framework (for instance, the results of [2, 8, 9, 10, 11, 12, 14, 22]). Our feeling is that, in general, new techniques and tools are needed to deal with these new problems. We finish this paper by illustrating situations in which we were able just to prove (1, \mathfrak{c})-lineability or (1, \mathfrak{c})-spaceability.

3.1. Non injective linear operators. Lineability and spaceability are investigated in the framework of (non) injective and (non) surjective functions/operators in several papers (see, for instance, [1, 3, 7, 18, 20] and the references therein). In this section we prove a related result in the new context initiated in the present paper; our result provides only (1, \mathfrak{c})-lineability; the general case seems to be an interesting open problem, as well as the case of (non) surjective operators.

Theorem 3.1. *Let $A := \{T \in \mathcal{L}(\ell_p; \ell_q) : T \text{ is non injective}\}$. Then, A is (1, \mathfrak{c})-lineable.*

Proof. In fact, let $T \in A \setminus \{0\}$ and consider W_1 the subspace generated by T . By hypothesis, there exist x, y in ℓ_p with $x \neq y$ such that

$$(3.1) \quad T(x) = T(y).$$

Since $T \neq 0$, there is a $z \in \ell_p$ such that $T(z) \neq 0$. Let j_0 be such that $(Tz)_{j_0} \neq 0$, where $(Tz)_{j_0}$ denotes the j_0 -th coordinate of the sequence $T(z)$. Let us choose $(\mathbb{N}_k)_{k=1}^\infty$ a sequence of pairwise disjoint subsets of \mathbb{N} with $\text{card}(\mathbb{N}_k) = \aleph_0$ for all k , and such that $j_0 \notin \bigcup_{k=1}^\infty \mathbb{N}_k$.

Define, for each $k \in \mathbb{N}$, a sequence of linear operators $T_k : \ell_p \rightarrow \ell_q$ of the form

$$T_k(x) = \begin{cases} (T(x))_j, & \text{if } j \in \mathbb{N}_k \\ 0, & \text{otherwise.} \end{cases}$$

Note that $T(x) = T(y) \Rightarrow T_k(x) = T_k(y)$. We conclude that $T_k \in A$ for all $k \in \mathbb{N}$. In addition, the set $\{T, T_k : k \in \mathbb{N}\}$ is linearly independent. In fact, let a, a_1, \dots, a_k be scalars and suppose that

$$aT + a_1T_1 + \dots + a_kT_k = 0.$$

We have

$$aT(z) + a_1T_1(z) + \dots + a_kT_k(z) = 0.$$

In particular

$$a(Tz)_{j_0} + a_1(T_1z)_{j_0} + \dots + a_k(T_kz)_{j_0} = 0.$$

Since $j_0 \notin \bigcup_{k=1}^\infty \mathbb{N}_k$, it follows that $(T_1z)_{j_0} = \dots = (T_kz)_{j_0} = 0$. So, $a(Tz)_{j_0} = 0$ and thus $a = 0$.

Consequently

$$a_1T_1 + \dots + a_kT_k = 0.$$

Since $\{T_1, \dots, T_k\}$ is linearly independent, we conclude that $a_1 = \dots = a_k = 0$.

Now consider the linear operator

$$\Psi : \ell_1 \rightarrow \mathcal{L}(\ell_p; \ell_q)$$

given by

$$\Psi((a_k)_{k=1}^\infty) = a_1T + \sum_{j=2}^\infty a_jT_{j-1}.$$

Note that Ψ is well defined, because

$$\begin{aligned} \|\Psi((a_k)_{k=1}^\infty)\|_{\mathcal{L}(\ell_p; \ell_q)} &= \left\| a_1 T + \sum_{j=2}^\infty a_j T_{j-1} \right\|_{\mathcal{L}(\ell_p; \ell_q)} \\ &\leq \|a_1 T\|_{\mathcal{L}(\ell_p; \ell_q)} + \sum_{j=2}^\infty \|a_j T_{j-1}\|_{\mathcal{L}(\ell_p; \ell_q)} \\ &\leq |a_1| \|T\|_{\mathcal{L}(\ell_p; \ell_q)} + \sum_{j=1}^\infty |a_j| \cdot \|T\|_{\mathcal{L}(\ell_p; \ell_q)} \\ &\leq \sum_{j=1}^\infty |a_j| \cdot \|T\|_{\mathcal{L}(\ell_p; \ell_q)} < \infty. \end{aligned}$$

Moreover, it is not difficult to prove that Ψ is injective. Let $a = (a_k)_{k=1}^\infty \in \ell_1$, with $a_k \neq 0$ for some $k \in \mathbb{N}$. So, from (3.1),

$$\begin{aligned} \Psi((a_k)_{k=1}^\infty)(x) &= a_1 T(x) + a_2 T_1(x) + a_3 T_2(x) + \dots \\ &= a_1 T(y) + a_2 T_1(y) + a_3 T_2(y) + \dots \\ &= \Psi((a_k)_{k=1}^\infty)(y), \end{aligned}$$

and $\Psi((a_k)_{k=1}^\infty)$ is non injective. Hence, $\Psi(\ell_1 \setminus \{0\}) \subseteq A$. Moreover,

$$T \in W_1 \subseteq \Psi(\ell_1 \setminus \{0\}) \subseteq A \cup \{0\}.$$

Therefore, A is $(1, \mathfrak{c})$ -lineable. \square

3.2. (α, \mathfrak{c}) -spaceability in L_p spaces. Recall that, for $1 \leq p < \infty$, $L_p[0, 1]$ denotes the classical space of the (class of equivalence of) measurable functions $f : [0, 1] \rightarrow \mathbb{K}$ equipped with the norm defined by

$$\|f\|_p = \left(\int_0^1 |f(t)|^p dt \right)^{\frac{1}{p}}.$$

By mimicking our constructive example of the beginning of the previous section, we can easily provide a simple construction of a function in $L_p[0, 1] \setminus \bigcup_{q>p} L_q[0, 1]$.

In [14] it was proved that $L_p[0, 1] \setminus \bigcup_{q>p} L_q[0, 1]$ is spaceable, but the proof does not assure that $L_p[0, 1] \setminus \bigcup_{q>p} L_q[0, 1]$ is (α, \mathfrak{c}) -spaceable for some cardinal $\alpha > 0$. The next result shows that this is true for $\alpha = 1$. The question for a cardinal $1 < \alpha < \mathfrak{c}$ remain unanswered. In the next proof, for any $X \subset [0, 1]$, the characteristic function of X on $[0, 1]$ is denoted by χ_X .

Theorem 3.2. $L_p[0, 1] \setminus \bigcup_{q>p} L_q[0, 1]$ is $(1, \mathfrak{c})$ -spaceable in $L_p[0, 1]$.

Proof. Let $f \in L_p[0, 1] \setminus \bigcup_{q>p} L_q[0, 1]$. It is obvious that

$$\tilde{f} = f\chi_{[0, 1/2]} \text{ or } \tilde{f} = f\chi_{[1/2, 1]}$$

belongs to $L_p[0, 1] \setminus \bigcup_{q>p} L_q[0, 1]$. Without loss of generality let us assume that

$$\tilde{f} \in L_p[0, 1] \setminus \bigcup_{q>p} L_q[0, 1].$$

Split $[1/2, 1]$ as an infinite sequence of disjoint intervals $I_n = [c_n, d_n)$. Notice that, for every $n \in \mathbb{N}$ and every $x \in I_n$, there is a unique $t_{x,n} \in [0, 1)$ such that

$$x = (1 - t_{x,n})c_n + t_{x,n}d_n.$$

Define

$$f_n(x) = \begin{cases} \tilde{f}(t_{x,n}) & \text{if } x \in I_n, \\ 0 & \text{if } x \notin I_n. \end{cases}$$

It is simple to verify that our construction provides

$$\|f_n\|_p < \left\| \tilde{f} \right\|_p,$$

for every $n \in \mathbb{N}$ and so $f_n \in L_p[0, 1]$.

Define $\tilde{p} = 1$ if $p \geq 1$ and $\tilde{p} = p$ if $0 < p < 1$. For $(a_i)_{i=1}^\infty \in \ell_{\tilde{p}}$,

$$\sum_{i=1}^{\infty} \|a_i f_i\|_{\tilde{p}}^{\tilde{p}} = \sum_{i=1}^{\infty} |a_i|^{\tilde{p}} \|f_i\|_{\tilde{p}}^{\tilde{p}} \leq \|\tilde{f}\|_{\tilde{p}}^{\tilde{p}} \sum_{i=1}^{\infty} |a_i|^{\tilde{p}} = \|\tilde{f}\|_{\tilde{p}}^{\tilde{p}} \|(a_i)_{i=1}^\infty\|_{\tilde{p}}^{\tilde{p}} < \infty.$$

Thus $\sum_{i=1}^{\infty} \|a_i f_i\|_{\tilde{p}}^{\tilde{p}} < \infty$ and thus the series $\sum_{i=1}^{\infty} a_i f_i$ converges in $L_p[0, 1]$. Hence, the operator

$$T: \ell_{\tilde{p}} \longrightarrow L_p[0, 1], \quad T((a_i)_{i=0}^\infty) = a_0 f + \sum_{i=1}^{\infty} a_i f_i$$

is well defined. It is easy to see that T is linear and injective. In fact, if

$$T((a_i)_{i=0}^\infty) = 0$$

then

$$a_0 f + \sum_{i=1}^{\infty} a_i f_i = 0$$

and choosing $x \in [0, 1/2]$ we have

$$a_0 f(x) = a_0 f(x) + \sum_{i=1}^{\infty} a_i f_i(x) = 0.$$

Since f is non null on $[0, 1/2]$ we conclude that $a_0 = 0$. Since $\{f_i : i \in \mathbb{N}\}$ is linearly independent (they have disjoint supports) we obtain $a_i = 0$ for all i .

Thus $\overline{T(\ell_{\tilde{p}})}$ is a closed infinite-dimensional subspace of $L_p[0, 1]$. We just have to show that

$$\overline{T(\ell_{\tilde{p}})} - \{0\} \subseteq L_p[0, 1] \setminus \bigcup_{q>p} L_q[0, 1].$$

Indeed, let $g \in \overline{T(\ell_{\tilde{p}})} \setminus \{0\}$. Thus $g \neq 0$ a.e., that is, the set $[0, 1] - A$ has null measure, where

$$A = \{x \in [0, 1] : g(x) \neq 0\}.$$

Let us consider sequences $(a_i^{(k)})_{i=0}^\infty \in \ell_{\tilde{p}}$ ($k \in \mathbb{N}$) such that $g = \lim_{k \rightarrow \infty} T\left(\left(a_i^{(k)}\right)_{i=0}^\infty\right)$ in $L_p[0, 1]$. By the definition of T we have

$$T\left(\left(a_i^{(k)}\right)_{i=0}^\infty\right) = a_0^{(k)} f + \sum_{i=1}^{\infty} a_i^{(k)} f_i \xrightarrow{k \rightarrow \infty} g \text{ in } L_p[0, 1].$$

In particular

$$T\left(\left(a_i^{(k)}\right)_{i=0}^\infty\right) \chi_{[0, 1/2]} \xrightarrow{k \rightarrow \infty} g \chi_{[0, 1/2]} \text{ in } L_p[0, 1].$$

Since f_i is null on the interval $[0, 1/2]$ for all i , we have

$$(3.2) \quad a_0^{(k)} \tilde{f} = T\left(\left(a_i^{(k)}\right)_{i=0}^\infty\right) \chi_{[0, 1/2]} \xrightarrow{k \rightarrow \infty} g \chi_{[0, 1/2]} \text{ in } L_p[0, 1].$$

On the other hand, note that

$$(3.3) \quad a_0^{(k)} \tilde{f} \xrightarrow{k \rightarrow \infty} \alpha \tilde{f} \text{ in } L_p[0, 1],$$

where $\alpha = \lim_{k \rightarrow \infty} a_0^{(k)}$. By (3.2) and (3.3) we have

$$g \chi_{[0, 1/2]} = \alpha \tilde{f} \text{ a.e.}$$

So, if $\alpha \neq 0$

$$\|g\|_q^q = \int_0^1 |g(t)|^q dt \geq \int_0^{1/2} |g(t)|^q dt = \alpha^q \|f \chi_{[0, 1/2]}\|_q^q = \infty,$$

proving that $g \notin L_q[0, 1]$.

If $\alpha = 0$, then $g\chi_{[0, 1/2]} = 0$ a.e. Define

$$\tilde{A} = \{x \in [1/2, 1] : g(x) \neq 0\}.$$

Since A has positive measure and $g\chi_{[0, 1/2]} = 0$ a.e., then \tilde{A} has positive measure. Since

$$\left\| \left(a_0^{(k)} f + \sum_{n=1}^{\infty} a_n^{(k)} f_n - g \right) \chi_{[1/2, 1]} \right\|_p \xrightarrow{k \rightarrow \infty} 0,$$

there is a subsequence

$$\left(\left(a_0^{(k_j)} f + \sum_{n=1}^{\infty} a_n^{(k_j)} f_n \right) \chi_{[1/2, 1]} \right)_{j=1}^{\infty}$$

such that

$$\left(a_0^{(k_j)} f(x) + \sum_{n=1}^{\infty} a_n^{(k_j)} f_n(x) \right) \chi_{[1/2, 1]}(x) \xrightarrow{j \rightarrow \infty} g(x) \chi_{[1/2, 1]}(x) \text{ a.e.}$$

Hence the set $[1/2, 1] - B$, where $B = \{x \in [1/2, 1] : \text{the limit above holds}\}$, has measure zero. Since,

$$\tilde{A} = (B \cap \tilde{A}) \cup (([1/2, 1] - B) \cap \tilde{A}),$$

and $([1/2, 1] - B) \cap \tilde{A}$ has measure zero and \tilde{A} has positive measure, it follows that $B \cap \tilde{A}$ has positive measure. Let $C = \{x \in [0, 1/2] : f(x) = 0\}$. Since $f \in L_p[0, 1] \setminus \bigcup_{q > p} L_q[0, 1]$ it follows that C has measure zero. By the fact that each f_n is the reproduction of f on the interval I_n , it follows that the set $C_n = \{x \in I_n : f_n(x) = 0\}$ has measure zero, for all $n \in \mathbb{N}$. Since

$$B \cap \tilde{A} = (B \cap \tilde{A} \cap C_n) \cup (B \cap \tilde{A} \cap (I_n - C_n)),$$

and $B \cap \tilde{A} \cap C_n$ has measure zero, then $B \cap \tilde{A} \cap (I_n - C_n)$ has positive measure, for each $n \in \mathbb{N}$. Fixing $r \in \mathbb{N}$ and choosing $x_0 \in B \cap \tilde{A} \cap (I_r - C_r)$, with $x_0 \neq 1$, we have that $x_0 \in I_r$, $f_r(x_0) \neq 0$, $g(x_0) \neq 0$ and

$$a_0^{(k_j)} f(x_0) + a_r^{(k_j)} f_r(x_0) = a_0^{(k_j)} f(x_0) + \sum_{n=1}^{\infty} a_n^{(k_j)} f_n(x_0) \longrightarrow g(x_0) \text{ when } j \rightarrow \infty.$$

Since $\lim_{k \rightarrow \infty} a_0^{(k)} = \alpha = 0$ we obtain

$$\lim_{j \rightarrow \infty} a_r^{(k_j)} = \frac{g(x_0)}{f_r(x_0)} = \eta \neq 0.$$

Since

$$f_r \chi_{I_r}(x) a_r^{(k_j)} \longrightarrow g \chi_{I_r}(x) \text{ a.e., when } j \rightarrow \infty,$$

by the unicity of the limit we have

$$g \chi_{I_r} = \eta f_r \chi_{I_r} \text{ a.e.,}$$

which implies that $g \chi_{I_r} \notin L_q[0, 1]$ and consequently $g \notin L_q[0, 1]$ (regardless of the $q > p$) finishing the proof. \square

REFERENCES

- [1] N. Albuquerque, Maximal lineability of the set of continuous surjections. *Bull. Belg. Math. Soc. Simon Stevin* 21 (2014), no. 1, 83–87.
- [2] N. Albuquerque, L. Bernal-González, D. Pellegrino, J. B. Seoane-Sepúlveda, *Peano curves on topological vector spaces*, *Linear Algebra Appl.* **460** (2014), 81–96.
- [3] R. M. Aron, L. Bernal-González, P. Jiménez-Rodríguez, G. Muñoz-Fernández, J. B. Seoane-Sepúlveda, *On the size of special families of linear operators*, *Linear Algebra Appl.* **544** (2018), 186–205.
- [4] R. M. Aron, L. Bernal-González, D. Pellegrino, J. B. Seoane-Sepúlveda, *Lineability: The Search for Linearity in Mathematics*, *Monographs and Research Notes in Mathematics*. CRC Press, Boca Raton (2016).

- [5] R. M. Aron, V. I. Gurariy, J. B. Seoane-Sepúlveda, *Lineability and spaceability of sets of functions on \mathbb{R}* , Proc. Amer. Math. Soc. **133** (2005), 795–803.
- [6] C. S. Barroso, G. Botelho, V. V. Fávaro, D. Pellegrino, *Lineability and spaceability for the weak form of Peano's theorem and vector-valued sequence spaces*, Proc. Amer. Math. Soc. **141** (2013), 1913–1923.
- [7] L. Bernal-González, *The algebraic size of the family of injective operators*, Open Math. **15** (2017), 13–20.
- [8] L. Bernal-González, M. O. Cabrera, *Lineability criteria, with applications*, J. Funct. Anal. **266** (2014), 3997–4025.
- [9] G. Botelho, D. Cariello, V. V. Fávaro, D. Pellegrino, *Maximal spaceability in sequence spaces*, Linear Algebra Appl. **437** (2012), 2978–2985.
- [10] G. Botelho, D. Cariello, V. V. Fávaro, D. Pellegrino, J. B. Seoane-Sepúlveda, *On very non-linear subsets of continuous functions*, Quart. J. Math. **65** (2014), 841–850.
- [11] G. Botelho, D. Diniz, V.V. Fávaro, D. Pellegrino, *Spaceability in Banach and quasi-Banach sequence spaces*, Linear Algebra. Appl. **434** (2011), 1255–1260.
- [12] G. Botelho, D. Diniz, D. Pellegrino, *Lineability of the set of bounded linear non-absolutely summing operators*, J. Math. Anal. Appl. **357** (2009), 171–175.
- [13] G. Botelho, V.V. Fávaro, *Constructing Banach spaces of vector-valued sequences with special properties*, Michigan Math. J. **64** (2015), 539–554.
- [14] G. Botelho, V. V. Fávaro, D. Pellegrino, J. B. Seoane-Sepúlveda, *$L_p[0, 1] - \cup_{q>p} L_q[0, 1]$ is spaceable for every $p > 0$* , Linear Algebra Appl. **436** (2012), 2963–2965.
- [15] D. Cariello, V.V. Fávaro, J. B. Seoane-Sepúlveda, *Self-similar functions, fractals and algebraic genericity*, Proc. Amer. Math. Soc. **145** (2017), 4151–4159.
- [16] D. Cariello and J. B. Seoane-Sepúlveda, *Basic sequences and spaceability in ℓ_p spaces*, J. Funct. Anal. **266** (2014), 3797–3814.
- [17] M. Fabian, P. Habala, P. Hájek, V. Montesinos and V. Zizler, *Banach Space Theory: The Basis for Linear and Nonlinear Analysis*, CMS Books in Mathematics (Canadian Mathematical Society/Springer), 2011.
- [18] J. L. Gámez-Merino, *Large algebraic structures inside the set of surjective functions*, Bull. Belg. Math. Soc. Simon Stevin, **18** (2011), 297–300.
- [19] V. I. Gurariy, L. Quarta, *On lineability of sets of continuous functions*, J. Math. Anal. Appl. **294** (2004), 62–72.
- [20] P. Jiménez-Rodríguez, P. S. Maghsoudi, G. Muñoz-Fernández, J. B. Seoane-Sepúlveda, *Injective mappings in $\mathbb{R}^{\mathbb{R}}$ and lineability*, Bull. Belg. Math. Soc. Simon Stevin **23** (2016), 609–623.
- [21] T. Nogueira, D. Pellegrino, *On the size of certain subsets of invariant Banach sequence spaces*, Linear Algebra Appl. **487**, 172–183.
- [22] D. Pellegrino, E. Teixeira, *Norm optimization problem for linear operators in classical Banach spaces*, Bull. Braz. Math. Soc. **40** (2009), 417–431.

FACULDADE DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE UBERLÂNDIA, 38.400-902 - UBERLÂNDIA
E-mail address: vvfavaro@gmail.com

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DA PARAÍBA, 58.051-900 - JOÃO PESSOA, BRAZIL
E-mail address: pellegrino@pq.cnpq.br

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DA PARAÍBA, 58.051-900 - JOÃO PESSOA, BRAZIL
E-mail address: danieltomazmatufpb@gmail.com