

Fano–Mathieu correspondence

Sergey Galkin

Preprint IPMU 10-0150. First version: 2010. Last version: September 11, 2018

Abstract

We show that G -Fano threefolds are *mirror-modular*.

1. *Mirror maps* are inversed reversed *Hauptmoduln* for moonshine subgroups of $SL_2(\mathbb{R})$.
2. *Quantum periods*, shifted by an integer constant (eigenvalue of quantum operator on primitive cohomology) are expansions of weight 2 modular forms (“theta-functions”) in terms of inversed Hauptmoduln.
3. Products of inversed Hauptmoduln with some fractional powers of shifted quantum periods are very nice cuspforms (“eta-quotients”).

The latter cuspforms also appear in work of Mason and others: they are eta-products, related to conjugacy classes of sporadic simple groups, such as Mathieu group M_{24} and Conway’s group of isometries of Leech lattice.

This gives a strange correspondence between deformation classes of G -Fano threefolds and conjugacy classes of Mathieu group M_{24} .

Introduction

The simplest example of Lian–Yau’s *mirror moonshine* for $K3$ surfaces ([16, 17], see also [6, 7, 21, 22]) is the remarkable identity (*modular relation*) of Kachru–Vafa [14], that goes back to Fricke and Klein:

$$\sum_{n \geq 0} \frac{(6n)!}{(3n)!n!^3} j(q)^{-n} = E_4(q)^{\frac{1}{2}} \quad (0.1)$$

expressing modular form $E_4(q)$ of weight 4 as square of hypergeometric series $I_1(t) = \sum \frac{(6n)!}{(3n)!n!^3} t^n$ expanded in terms of inverse modular parameter $t = j(q)^{-1}$. Here $\eta(q) = q^{\frac{1}{24}} \prod_{n \geq 1} (1 - q^n)$ is Dedekind’s eta-function, $\Delta(q) = \eta(q)^{24}$ is modular discriminant, $\sigma_3(n) = \sum_{d|n} d^3$, Eisenstein series $E_4(q) = 1 + 240 \sum_{n \geq 1} \sigma_3(n) q^n$ equals to the theta-series θ_{E_8} for a lattice E_8 , and $j(q) = \frac{E_4^3}{\Delta(q)} = \frac{1}{q} + 744 + 196884q + \dots$ is Klein’s modular j -invariant. The respective weights of these modular forms are: $wt([\eta, \Delta, E_4, j]) = [\frac{1}{2}, 12, 4, 0]$. Multiplying both sides by $j(q)^{-\frac{1}{6}}$, and then taking to the power six implies an identity for $\Delta(q)$:

$$j(q)^{-1} \left(\sum_{n \geq 0} \frac{(6n)!}{(3n)!n!^3} j(q)^{-n} \right)^6 = \left(\sum_{n \geq 0} \frac{(6n)!}{(3n)!n!^3} j(q)^{-n-\frac{1}{6}} \right)^6 = \eta(q)^{24} = \Delta(q) \quad (0.2)$$

One advantage of formula 0.2 over 0.1 is that Δ is a holomorphic cuspform. Moreover, it is an eigenform for Hecke operations, and has no zeroes on upper half-plane. That is, Δ is an *eta-product*.

Smooth anticanonical divisor $S \in |-K_Y|$ in Fano threefold Y is a $K3$ surface endowed with a natural lattice polarization $c_1(Y) \in \text{Pic}(Y) \subset \text{Pic}(S) \subset H^2(S, \mathbb{Z}) = II_{3,19}$. Beauville [2] shows inverse: generic $K3$ surface lattice-polarized by $c_1(Y) \in \text{Pic}(Y)$ is anti-canonically embedded to a Fano threefold Y . So Fano threefolds single out 105 (see [13, 19]) out of countably many families of lattice polarized $K3$ surfaces (and also 105 mirror dual families [6]). In fact, almost all moonshine examples listed in [16, 17, 6, 21, 22] appear in that way, and mirror moonshine makes more sense in the context of Fano threefolds. Golyshev [12] reproduced Iskovskikh’s classification of prime Fano threefolds (see e.g. [13]) by effectively combining three elements: mirror, moonshine and minimality.

G -series G_Y (see 1.4) is a certain invariant of Fano variety Y “counting” rational curves on it.

Mirror conjecture (for variations of Hodge structures) states that Laplace transform of G -series for Fano threefold Y is the solution of Picard–Fuchs differential equation for some 1-parameter family of $K3$ surfaces that is called *mirror dual to Y Ginzburg–Landau model*.

Moonshine (*genus-zero modularity*) is explicitly stated as *miraculous eta-product formula*:

$$I_{N,s}(H_{N,c}^{-1}) = \eta(q)^2 \eta(q^N)^2 H_{N,c}^{\frac{N+1}{12}} \quad (0.3)$$

where G_{Y_N} is G -series of Fano threefold Y_N with invariants $H^2(Y_N, \mathbb{Z}) = \mathbb{Z}c_1(Y_N)$, $c_1(Y_N)^3 = 2N$, $I_{N,s} = \mathbb{R}_s G_{Y_N}(t)$ is Laplace transform of G_{Y_N} multiplied by e^{st} for a particular choice of constant $s = s_N$ (see 1.1.1.7), and H_N is a Hauptmodul on Fricke modular curve $X_0(N)/w_N$ (see [5]) with a particular constant term $c = c_N$: $H_{N,c} = \frac{1}{q} + c_N + O(q)$. In case Y is a sextic double solid (i.e. a smooth sextic hypersurface in weighted projective 4-space $\mathbb{P}(1, 1, 1, 1, 3)$) we have $N = 1$, $H_1 = j(q)$, $s_1 = 120$, $c_1 = 744$ and formula 0.3 specializes to 0.1 (we present other exact equalities in appendix 6). One of the points of this note, is that one also has more natural identity for a cusp-eigenform (eta-product or eta-quotient), that specializes to 0.2, and the latter identity generalizes to a more general class of G -Fano threefolds, with eta-products usually enoded by conjugacy classes of Mathieu group M_{24} .

Minimality is formalized in the notion of *D3 differential equation* (see 4.1). It is a 6-parameter class of differential equations of degree 3, generalizing the construction of regularized quantum differential equations of a Fano threefold from 6 two-point Gromov–Witten invariants.

Modularity conjecture (which is now a theorem) states that function $G = \sum_{n \geq 0} a_i t^n$ is G -series of minimal Fano threefold of index one if and only if for some s function $\mathbb{R}_s G$ is of moonshine type (satisfy 0.3 for some N) and is annihilated by differential equation of type $D3$.

Apart from 17 quantum differential equations of minimal smooth Fano threefolds Golyshev found two more differential equations of type $D3$ and modular origin (4.6 and 4.7) (also these equations were found by Almkvist, van Enckevort, van Straten and Zudilin in [1], and they were already listed in [17]).

It turned out ([9]) that these two examples are quantum differential equations for two deformation classes Y_{28} and Y_{30} of Fano threefolds with $H^2(Y_{28}, \mathbb{Z}) = \mathbb{Z}^2$, $H^2(Y_{30}, \mathbb{Z}) = \mathbb{Z}^3$.

Since Fano threefolds Y_{28} and Y_{30} are not minimal one naively expects their regularized quantum differential equations to be of degree 4 and 5, but these varieties occur to be *quantum minimal* — minimal differential equation vanishing \widehat{G} -series of these varieties has degree 3 (see [10] for the details).

In [9] we made an observation that both Y_{28} and Y_{30} are *G-Fano threefolds* i.e. for some complex structure they admit a finite group action $G : Y$ with $\text{Pic}^G(Y) = \mathbb{Z}c_1(Y)$. In [10] we shown that G -Fano threefolds are quantum minimal. So it is natural to look whether other G -Fano threefolds are mirror-modular.

Families Y_{28} and Y_{30} are two of total eight families of G -Fano threefolds (see [20]). In these article we will show that all 8 families of G -Fano threefolds are mirror-modular.

1 Preliminaries

1.1 Shifts and regularizations.

For a number s and a power series $A = \sum_{n \geq 0} a_i t^n$ define its regularization (Laplace transform \mathcal{L}), inverse Laplace transform \mathcal{L}^{-1} , shifted regularization \mathcal{L}_s , regular shift \mathcal{S}_g and normalization \mathcal{N} by the formulas

$$\begin{aligned} \widehat{A} &= \mathcal{L}A = \sum (a_i \cdot n!) t^n, \\ \mathcal{L}^{-1}A &= \sum \frac{a_i}{n!} t^n, \\ \mathcal{L}_s A &= \mathcal{L}(e^{s \cdot t} \cdot A) \\ \mathcal{S}_s A &= \mathcal{L}_s \mathcal{L}^{-1}A \\ \mathcal{N}A &= S_{-a_1} A \end{aligned}$$

1.2 Fano varieties

[see e.g. [13]]

Let Y be a Fano variety — smooth variety with ample anticanonical class ω_Y^{-1} .

Y is simply-connected, by Kodaira vanishing $H^i(Y, \mathcal{O}_Y) = 0$ for $i > 0$, so $c_1 : \text{Pic}(Y) \rightarrow H^2(Y, \mathbb{Z})$ is an isomorphism and both are isomorphic to \mathbb{Z}^ρ , where ρ is called Picard number. Lefschetz pairing on $H^2(Y, \mathbb{Z})$ defined by $(A, B) = \int_{[Y]} A \cup B \cup c_1(Y)^{\dim Y - 2}$ is nondegenerate (by Hard Lefschetz theorem), so $H^2(Y, \mathbb{Z})$ is a lattice and we denote its discriminant by $d(Y)$.

Anticanonical *degree* of Fano variety Y is $\text{deg}(Y) = (c_1(Y), c_1(Y)) = \int_{[Y]} c_1(Y)^{\dim Y}$. Euler number is a topological Euler characteristic $\chi(Y) = \int_{[Y]} c_{\dim Y}$.

Fano index $r(Y)$ is divisibility of $c_1(Y)$ in the lattice $H^2(Y, \mathbb{Z})$ i.e. $r(Y) = \max\{r \in \mathbb{Z} | c_1(Y) = rH, H \in H^2(Y, \mathbb{Z})\}$.

Definition 1.1 ([20]). Fano variety Y with group action $G : Y$ is called G -Fano if $H^2(Y, \mathbb{Z})^G = \mathbb{Z}$.

1.3 Quantum differential equations, G-series and Givental's constant

Let \star be quantum multiplication on cohomologies of Y defined by

$$\int_{[Y]} (\gamma_1 \star \gamma_2) \cup \gamma_3 = \sum_{d \geq 0} \langle \gamma_1, \gamma_2, \gamma_3 \rangle_d t^d \quad (1.2)$$

where closed genus 0 3-point correlator $\langle \gamma_1, \gamma_2, \gamma_3 \rangle_d = \int_{\overline{\mathcal{M}}_{0,3}(Y,d)} \prod ev_i^*(\gamma_i)$ is a Gromov-Witten invariant “counting” maps $f : \mathbb{P}^1 \rightarrow Y$ of degree $d = \int_{[\mathbb{P}^1]} f^* c_1(Y)$ passing through homology classes Poincare-dual to γ_i .

Take $D = t \frac{d}{dt}$ and define *quantum differential equation* as a trivial vector bundle over $\text{Spec } \mathbb{C}[t, t^{-1}]$ with fibre $H^*(Y)$ and connection

$$D\Phi = c_1(Y) \star \Phi \quad (1.3)$$

where $\Phi \in H^*(Y)[[t]]$.

Let $\mathcal{G}_Y(t) = [pt] + \sum_{n \geq 1} \mathcal{G}^{(n)} t^n$ be the unique analytic solution of 1.3 starting with class Poincare-dual to the the class of the point, and define G -series as

$$G_Y(t) = \int_{[Y]} \mathcal{G}_Y(t) = 1 + \sum_{n \geq 1} G^{(n)} t^n \quad (1.4)$$

We note that the first coefficient $G^{(1)} = \langle [pt], c_1(Y), [Y] \rangle_1 = \int_{|t|=\epsilon} (\int_{[Y]} c_1(Y) \star [pt]) \frac{dt}{t^2}$ is zero according to the String equation, and we name $G^{(2)} = \langle [pt] \rangle_2$ (the expected number of anticanonical conics passing through a point) as *Givental's constant* $G(Y)$, so

$$G_Y = 1 + G(Y) \cdot t^2 + O(t^3). \quad (1.5)$$

Define \widehat{G} -series (also known as *the quantum period* of Y) as

$$\widehat{G}_Y = \widehat{G}_Y = 1 + \sum_{n \geq 1} n! \cdot G^{(n)} t^n = 1 + 2G(Y) \cdot t^2 + \dots \quad (1.6)$$

Conjecturally \widehat{G}_Y should have integer coefficients.

For convenience we define I -series to be a regularized shifted G -series:

Definition 1.7. For a given number s define $I_{Y,s} = \mathcal{L}_s G_Y$, in particular $I_{Y,0} = \widehat{G}_Y$. By a slight abuse of notation we will say that power series $I(t) = 1 + \sum_{n \geq 1} i^{(n)} t^n$ is a regular I -series of smooth Fano variety Y if $\mathcal{N}I = \widehat{G}_Y$ i.e. $I = I_{Y,i(1)}$.

2 G-Fano threefolds and A-model G-series

There are 8 deformation classes of G -Fano threefolds Y with $\text{rk } H^2(Y, \mathbb{Z}) > 1$ (see e.g. [20]). Two of them has Fano index two, these are $Y_{48}^{(3)} = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ and $Y_{48}^{(2)} = W \subset \mathbb{P}^2 \times \mathbb{P}^2$ of degree 48. Six other have Fano index one: Y_{30} of degree 30, Y_{28} of degree 28, Y_{24} of degree 24, Y_{20} of degree 20, $Y_{12}^{(2)}$ and $Y_{12}^{(3)}$ of degree 12.

In this section we are going to describe all of them geometrically. The details regarding the computation of the respective quantum periods can be found in [3].

Definition 2.1. Fano threefold $Y_{48}^{(2)}$ is the variety $Fl(1, 2, 3)$ of complete flags in \mathbb{P}^2 i.e. a hyperplane section of Segre fourfold $\mathbb{P}^2 \times \mathbb{P}^2 \subset \mathbb{P}^8$. This variety is unique in its deformation class number 32 in table 2 of [19], it has Fano index 2, degree 48, $\chi = 6$ and $\rho = 2$.

This variety can also be described as a projectivization of tangent bundle on \mathbb{P}^2 or variety of complete flags in \mathbb{P}^2 .

Corollary 2.2. I -series $I_{6,2;2} = \mathbb{R}G_{Fl(1,2,3)}$ is given by the pullback of hypergeometric series from two-dimensional torus

$$I_{6,2;2} = \sum_{a,b \geq 0} \frac{(a+b)!(2a+2b)!}{a!^3 b!^3} t^{2(a+b)} = 1 + 4t^2 + 60t^4 + 1120t^6 + 24220t^8 + 567504t^{10} + \dots \quad (2.3)$$

Proof. Combine the computation of I -series of toric variety $\mathbb{P}^2 \times \mathbb{P}^2$ in [11] and quantum Lefschetz principle in [4]. \square

Definition 2.4. Fano threefold $Y_{48}^{(3)}$ is just a Cartesian cube of a line i.e. Segre threefold $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \subset \mathbb{P}^7$. This variety is unique in its deformation class number 27 in table 3 of [19], it has Fano index 2, degree 48, $\chi = 8$ and $\rho = 3$.

Corollary 2.5. I -series $I_{6,3;2} = \mathbb{R}G_{\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1}$ is given by the pullback of hypergeometric series from three-dimensional torus

$$I_{6,3;2} = \sum_{a,b,c \geq 0} \frac{(2a+2b+2c)!}{a!^2 b!^2 c!^2} t^{2(a+b+c)} = 1 + 6t^2 + 90t^4 + 1860t^6 + 44730t^8 + 1172556t^{10} + \dots \quad (2.6)$$

Remark 2.7. Threefolds $Y_{48}^{(2)}$ and $Y_{48}^{(3)}$ has isomorphic hyperplane sections — del Pezzo surface of degree 6. This implies the relation $G_{\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1}(\sqrt{t}) = G_{Fl(1,2,3)}(\sqrt{t}) \cdot e^t$

Definition 2.8. Fano threefold Y_{30} is the blowup of a curve of bidegree $(2, 2)$ on $Fl(1, 2, 3) \subset \mathbb{P}^2 \times \mathbb{P}^2$. This deformation class of varieties has number 13 in table 3 of [19], it has degree 30, $\chi = 8$ and $\rho = 3$.

Proposition 2.9. Y_{30} is a complete intersection of three numerically effective divisors of tridegrees $(1, 1, 0)$, $(1, 0, 1)$ and $(0, 1, 1)$ on $\mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^2$.

Corollary 2.10. I -series $I_{15} = \mathcal{L}_3 G_{Y_{30}}$ is given by the pullback of hypergeometric series from three-dimensional torus

$$I_{15} = \sum_{a,b,c \geq 0} \frac{(a+b)!(a+c)!(b+c)!(a+b+c)!}{a!^3 b!^3 c!^3} t^{a+b+c} = 1 + 3t + 15t^2 + 105t^3 + 855t^4 + 7533t^5 + \dots \quad (2.11)$$

Let Q be 3-dimensional quadric.

Definition 2.12. Let Y_{28} be the blowup of a twisted quartic on Q . This deformation class of varieties has number 21 in table 2 of [19], it has degree 28, $\chi = 6$ and $\rho = 2$.

Denote the I -series of Y_{28} as $I_{14} = \widehat{G}_{Y_{28}}$.

Definition 2.13. Fano threefold Y_{24} is a hyperplane section of Segre embedding $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \subset \mathbb{P}^{15}$. This deformation class of varieties has number 1 in table 4 of [19], it has degree 24, $\chi = 6$ and $\rho = 4$.

Corollary 2.14. I -series $I_{12} = \mathcal{L}_4 G_{Y_{24}}$ is given by the pullback of hypergeometric series from four-dimensional torus

$$I_{12} = \sum_{a,b,c,d \geq 0} \frac{(a+b+c+d)!^2}{a!^2 b!^2 c!^2 d!^2} t^{a+b+c+d} = 1 + 4t + 28t^2 + 256t^3 + 2716t^4 + 31504t^5 + \dots \quad (2.15)$$

Definition 2.16. Let Y_{20} be the blowup of projective space \mathbb{P}^3 with center a curve of degree 6 and genus 3 which is an intersection of cubics. This deformation class of varieties has number 12 in table 2 of [19], it has degree 20, $\chi = 0$ and $\rho = 2$.

Proposition 2.17. Y_{20} is an intersection of Segre variety $\mathbb{P}^3 \times \mathbb{P}^3$ by linear subspace of codimension 3.

Corollary 2.18. I -series $I_{10} = \mathcal{L}_2 G_{Y_{20}}$ is given by the pullback of hypergeometric series from two-dimensional torus

$$I_{10} = \sum_{a,b \geq 0} \frac{(a+b)!^4}{a!^4 b!^4} t^{a+b} = 1 + 2t + 18t^2 + 164t^3 + 1810t^4 + 21252t^5 + 263844t^6 + 3395016t^7 + \dots \quad (2.19)$$

Definition 2.20. Fano threefold $Y_{12}^{(2)}$ can be described either as a section of Segre fourfold $\mathbb{P}^2 \times \mathbb{P}^2 \subset \mathbb{P}^8$ by quadric or as double cover of W with branch locus in anticanonical divisor. This deformation class of varieties has number 6 in table 2 of [19], it has degree 12, $\chi = -12$ and $\rho = 2$.

Corollary 2.21. I -series $I_{6,2} = \mathcal{L}_4 G_{Y_{12}^{(2)}}$ is given by the pullback of hypergeometric series from two-dimensional torus

$$I_{6,2} = \sum_{a,b \geq 0} \frac{(a+b)!(2a+2b)!}{a!^3 b!^3} t^{a+b} = 1 + 4t + 60t^2 + 1120t^3 + 24220t^4 + 567504t^5 + \dots \quad (2.22)$$

Remark 2.23. Series $I_{6,2}$ and $I_{6,2;2}$ are related by change of coordinates $I_{6,2;2}(t) = I_{6,2}(t^2)$.

Definition 2.24. Fano threefold $Y_{12}^{(3)}$ is a double cover of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ with branch locus in anticanonical divisor. This deformation class of varieties has number 1 in table 3 of [19], it has degree 12, $\chi = -8$ and $\rho = 3$.

Corollary 2.25. I -series $I_{6,3} = \mathcal{L}_6 G_{Y_{12}^{(3)}}$ is given by the pullback of hypergeometric series from two-dimensional torus

$$I_{6,3} = \sum_{a,b,c \geq 0} \frac{(2a+2b+2c)!}{a!^2 b!^2 c!^2} t^{a+b+c} = 1 + 6t + 90t^2 + 1860t^3 + 44730t^4 + 1172556t^5 + \dots \quad (2.26)$$

Remark 2.27. Series $I_{6,3}$ and $I_{6,3;2}$ are related by change of coordinates $I_{6,3;2}(t) = I_{6,3}(t^2)$.

3 Eta-products, Hauptmoduln and their M-series

Let $\eta(q)$ be Dedekind's eta-function: $\eta(q) = q^{\frac{1}{24}} \prod_{n \geq 1} (1 - q^n)$.

Consider 4 eta-products and 1 eta-quotient:

$$\eta_{6+} = \eta(q)\eta(q^2)\eta(q^3)\eta(q^6) \quad (3.1)$$

$$\eta_{10+} = \eta(q)\eta(q^2)\eta(q^5)\eta(q^{10}) \quad (3.2)$$

$$\eta_{12+} = \frac{\eta(q^2)^4 \eta(q^6)^4}{\eta(q)\eta(q^3)\eta(q^4)\eta(q^{12})} \quad (3.3)$$

$$\eta_{14+} = \eta(q)\eta(q^2)\eta(q^7)\eta(q^{14}) \quad (3.4)$$

$$\eta_{15+} = \eta(q)\eta(q^3)\eta(q^5)\eta(q^{15}) \quad (3.5)$$

Let $\sigma_1(n)$ be -24 times valuation of η_{n+} .

For a constant c and a conjugacy class g of Monster simple group denote by $T_{g,c}$ its McKay-Thompson series (see [5]) with constant term normalized to be c : $T_{g,c} = \frac{1}{q} + c + \sum_{n \geq 1} a_i(g)q^n$.

Take $H_{6A,2} = T_{6A,10}$, $H_{6A,3} = T_{6A,14}$, $H_{10A} = T_{10A,4}$, $H_{12A} = T_{12A,6}$, $H_{14A,s} = T_{14A,s}$, $H_{15A,s} = T_{15A,s}$:

$$T_{6A,0} = \frac{1}{q} + 79q + 352q^2 + 1431q^3 + 4160q^4 + 13015q^5 + 31968q^6 + \dots \quad (3.6)$$

$$H_{10A} = 8 + \frac{\eta^4(q)\eta^4(q^5)}{\eta^4(q^2)\eta^4(q^{10})} + 16 \frac{\eta^4(q^2)\eta^4(q^{10})}{\eta^4(q)\eta^4(q^5)} = \frac{1}{q} + 4 + 22q + 56q^2 + 177q^3 + 352q^4 + \dots \quad (3.7)$$

$$H_{12A} = \left(\frac{\eta(q^2)^2 \eta(q^6)^2}{\eta(q)\eta(q^3)\eta(q^4)\eta(q^{12})} \right)^6 = \frac{1}{q} + 6 + 15q + 32q^2 + 87q^3 + 192q^4 + \dots \quad (3.8)$$

$$H_{14A} = 4 + \frac{\eta^3(q)\eta^3(q^7)}{\eta^3(q^2)\eta^3(q^{14})} + 8 \frac{\eta^3(q^2)\eta^3(q^{14})}{\eta^3(q)\eta^3(q^7)} = \frac{1}{q} + 1 + 11q + 20q^2 + 57q^3 + 92q^4 + \dots \quad (3.9)$$

$$H_{15A} = 3 + \frac{\eta^2(q)\eta^2(q^5)}{\eta^2(q^3)\eta^2(q^{15})} + 9 \frac{\eta^2(q^3)\eta^2(q^{15})}{\eta^2(q)\eta^2(q^5)} = \frac{1}{q} + 1 + 8q + 22q^2 + 42q^3 + 70q^4 + \dots \quad (3.10)$$

Define $M_n(t)$ as power-series satisfying the functional equation $M_n\left(\frac{1}{H_n(q)}\right) = \eta_{n+} \cdot H_n^{\frac{\sigma_1(n)}{24}}$:

$$M_{6,2}(H_{6A,2}^{-1}(q)) = \eta_{6+} \cdot H_{6A,2}^{\frac{1}{2}} \quad (3.11)$$

$$M_{6,3}(H_{6A,3}^{-1}(q)) = \eta_{6+} \cdot H_{6A,3}^{\frac{1}{2}} \quad (3.12)$$

$$M_{10}(H_{10}^{-1}(q)) = \eta_{10+} \cdot H_{10A}^{\frac{3}{4}} \quad (3.13)$$

$$M_{12}(H_{12}^{-1}(q)) = \eta_{12+} \cdot H_{12A}^{\frac{1}{2}} \quad (3.14)$$

$$M_{14,s}(H_{14,s}^{-1}(q)) = \eta_{14+} \cdot H_{14A,s} \quad (3.15)$$

$$M_{15,s}(H_{15,s}^{-1}(q)) = \eta_{15+} \cdot H_{15A,s} \quad (3.16)$$

4 Differential equations and their solutions

Let t be a coordinate on $G_m = \text{Spec } \mathbb{C}[t, t^{-1}]$ and $D = t \frac{d}{dt}$.

Definition 4.1 ([12]). *Normalized operators of type D3* is the following 5-dimensional family of differential operators depending on parameters b_1, b_2, b_3, b_4, b_5 ¹:

$$L(b_1, b_2, b_3, b_4, b_5) = D^3 - t \cdot b_1 D(D+1)(2D+1) - t^2 \cdot (D+1)(b_2 D(D+2) + 4b_3) - t^3 \cdot b_4(D+1)(D+2)(2D+3) - t^4 \cdot b_5(D+1)(D+2)(D+3)$$

Define $L_{6,2}, L_{6,3}, L_{10}, L_{12}, L_{14}, L_{15}$ as follows:

$$L_{6,2} = L(6, 368, 88, 1056, 3584) \quad (4.2)$$

$$L_{6,3} = L(8, 360, 108, 864, 2160) \quad (4.3)$$

$$L_{10} = L(2, 112, 28, 184, 336) \quad (4.4)$$

$$L_{12} = L(2, 80, 24, 96, 0) \quad (4.5)$$

$$L_{14} = L(1, 59, 16, 68, 80) \quad (4.6)$$

$$L_{15} = L(1, 43, 12, 78, 216) \quad (4.7)$$

Locally equation L_n has 1-dimensional space of analytic solutions spanned by F_n :

$$F_{6,2}(t) = 1 + 44t^2 + 528t^3 + 11292t^4 + 228000t^5 + 4999040t^6 + 112654080t^7 + \dots \quad (4.8)$$

$$F_{6,3}(t) = 1 + 54t^2 + 672t^3 + 15642t^4 + 336960t^5 + 7919460t^6 + 191177280t^7 + \dots \quad (4.9)$$

$$F_{10}(t) = 1 + 14t^2 + 72t^3 + 882t^4 + 8400t^5 + 95180t^6 + 1060080t^7 + \dots \quad (4.10)$$

$$F_{12}(t) = 1 + 12t^2 + 48t^3 + 540t^4 + 4320t^5 + 42240t^6 + 403200t^7 + \dots \quad (4.11)$$

$$F_{14}(t) = 1 + 8t^2 + 24t^3 + 240t^4 + 1440t^5 + 11960t^6 + 89040t^7 + \dots \quad (4.12)$$

$$F_{15}(t) = 1 + 6t^2 + 24t^3 + 162t^4 + 1080t^5 + 7620t^6 + 55440t^7 + \dots \quad (4.13)$$

5 Equivalence of realizations

Lemma 5.1. *I-series I_n are solutions to differential equations L_n listed in 4, i.e. $\mathcal{N}I_n = F_n$.*

Proof. By [10] G -function G_V of G -Fano threefold V satisfy ODE of order 4 and its Fourier–Laplace transform \widehat{G}_V satisfy a Fuchsian ODE of order 3. \square

Lemma 5.2. *M -series M_n are solutions to differential equations L_n listed in 4, i.e. $\mathcal{N}M_n = F_n$.*

Proof. By Proposition 21 of [23] function M_n satisfy some differential equation of order 3 in $D = t \frac{d}{dt}$. \square

We have an immediate

Corollary 5.3. *For every n series $F_n, \mathcal{N}I_n$ and $\mathcal{N}M_n$ coincide.*

and it implies the main

Theorem 5.4. *For every n there are constants s_n and c_n such that I-series of G -Fano threefolds satisfy generalized miraculous eta-product formula 0.3: $I_{n,s}(H_{n,c}^{-1}) = \eta_{n+} \cdot H_{n,c}^{\frac{\sigma_1(n)}{2^4}}$.*

6 Golyshev’s modularity of minimal Fano threefolds, and beyond

Let Y_N be a Fano threefold with $H^2(Y, \mathbb{Z}) = \mathbb{Z}K_Y$ and $(-K_Y)^3 = 2N$. Let $G_Y = 1 + \sum_{n \geq 2} G^{(n)}(Y)t^n$ be its G -series. Golyshev’s modularity conjecture states that for every $N = 1, \dots, 9, 11$ there exists such a constant s_N , and a Monster conjugacy class g_N ($N + N$ in notations of [5]), and a constant c_N such that

$$\eta^2(q)\eta^2(q^N) \cdot T_{g_N, c_N}^{\frac{N+1}{12}}(q) = I_{Y_N, s_N} \left(\frac{1}{T_{g_N, c_N}(q)} \right)$$

where $T_{g_N, c_N} = \frac{1}{q} + c_N + O(q)$ is McKay–Thompson series for conjugacy class g_N with constant term c_N .

For $N = 6$ the conjugacy class $6 + 6$ is $6B$, for other values of N it is NA .

¹Original definition has the other basis $a_{01}, a_{02}, a_{03}, a_{11}, a_{12}$ for parameter space \mathbb{A}^5 . Bases a and b are equivalent over \mathbb{Z} : $b_1 = a_{11}$, $b_2 = a_{12} + 2a_{01} - a_{11}^2$, $b_3 = a_{01}$, $b_4 = a_{02} - a_{01}a_{11}$, $b_5 = a_{03} - a_{01}^2$; $a_{01} = b_3$, $a_{02} = b_4 + b_1b_3$, $a_{03} = b_5 + b_3^2$, $a_{11} = b_1$, $a_{12} = b_2 - 2b_3 + b_1^2$.

In the table we specify values N , $g = g_N$, $c = c_N$ and $s = s_N$ for 16 G -Fano threefolds of index 1 – 10 Golyshev's cases and 6 other G -Fano threefolds with $N = 6, 6, 10, 12, 14, 15$ that are explained in this paper.

For integer N we define numbers:

$$\phi(N) = N \prod_{p|N} (1 - p^{-1}), \psi(N) = N \prod_{p|N} (1 + p^{-1}), \epsilon(N) = \frac{24}{\psi(N)}, \iota(N) = \sum_{M|N} \phi(M) \epsilon(M).$$

N	1	2	3	4	5	6	6	6	7	8	9	10	11	12	14	15
$\epsilon(N)$	24	8	6	4	4	2	2	2	3	2	2	4/3	2	1	1	1
$\iota(N)$	24	16	12	10	8	8	8	8	6	6	4	8*	4	5*	4	4
s	120	24	12	8	6	6	5	4	4	4	3	2	*	4	*	*
c	744	104	42	24	16	14	12	10	9	8	6	4	s+2	6	s+1	s+1
g	1A	2A	3A	4A	5A	6A	6B	6A	7A	8A	9A	10A	11A	12A	14A	15A
ρ	1	1	1	1	1	3	1	2	1	1	1	2	1	4	2	3

Proposition 6.1. Number $\psi(N)$ equals to index of $\Gamma_0(N)$ in $SL(2, \mathbb{Z})$. Number $\phi(N)$ equals to index of $\Gamma_1(N)$ in $\Gamma_0(N)$.

Number $\epsilon(N)$ is integer only for 15 values of N .

Proposition 6.2. Let N be one of $1, \dots, 8, 11, 14, 15, 23$. Denote by M_{23} the Mathieu group and by V its natural 24-dimensional representation induced from Mathieu group M_{24} (which in turn induced from Conway group Co_0). Let $g \in M_{23}$ be an element of Mathieu group M_{23} of order N . Then number $\epsilon(N)$ equals to the trace $\text{Tr}g : V$ and number $\iota(N)$ equals to the dimension of invariants $\dim V^g$.

Remark 6.3. Note that for $N = 11, 14, 15$ the respective space of modular forms is 2-dimensional and any choice of s produces a modular relation. There is a particular choice of c depending on s : the difference $(c - s)$ is an invariant of Fano threefold. For $N \neq 11, 14, 15$ the choice of s is unique: s is a natural number such that I -function has singularity at 0^2 .

7 Mathieu groups.

Definition 7.1. Let S be the set of 24 points and $S_{24} = \text{Aut}(S)$ is its group of automorphisms. Let $M = S^{\mathcal{Q}}$ be a vector space of the tautological 24-dimensional representation of S_{24} . Mathieu group M_{24} is a particular simple subgroup of S_{24} of order $244823040 = 23 \cdot 11 \cdot 7 \cdot 5 \cdot 3^3 \cdot 2^{10} = 24 \cdot 23 \cdot 22 \cdot 21 \cdot 20 \cdot (16 \cdot 3)$.

Natural action $M_{24} : S$ is 5-transitive.

Definition 7.2. Stabilizer of one point in this action is simple Mathieu group M_{23} of order $23 \cdot 11 \cdot 7 \cdot 5 \cdot 3^2 \cdot 2^7$.

Definition 7.3. Any transposition $h \in S_{24}$ of symmetric group decomposes into the product of cycles, so we will say *Frame shape* of h is $\prod_{n \geq 1} i^{a_i}$ where $a_i(h)$ is number of cycles of length n . Frame shape is a complete invariant for conjugacy classes of S_{24} (and complete up to power-equivalence for M_{24} and M_{23}).

Example 7.4. 1. S_{24} has 1575 Frame shapes and conjugacy classes (equal to number of partitions of 24),

2. M_{24} has 21 Frame shapes and 26 conjugacy classes,

3. M_{23} has 12 Frame shapes and 17 conjugacy classes.

Definition 7.5. Order of element (conjugacy class, Frame shape) h equals to the least common multiple of cycle lengths in its cycle decomposition: $n(h) = \text{lcm}(a_1, \dots, a_{24})$. Denote by $G(h) = \text{Tr}h : \mathbb{Q}^{24} = H^*(24\text{points}, \mathbb{Q})$ the number a_1 of cycles of length one (it can be computed via Lefschetz fixed point formula on finite set of 24 elements).

Obviously $G(h)$ is integer non-negative and if $h \in M_{23}$ then $G(h) \geq 1$.

Proposition 7.6 (Frobenius, Mukai). 1. $h \in M_{24}$ comes from $M_{23} \iff G(h) \geq 1$.

2. For element $h \in M_{23}$ number $G(h)$ depends only on $n(h)$: $G(h) = \epsilon(n)$.

3. There are 12 orders of elements in M_{23} : from 1 to 8, 11, 14, 15 and 23. Frame shapes of M_{23} are determined by the orders.

Proposition 7.7. M_{23} has the following 12 Frame shapes:

g	1^{24}	$1^8 2^8$	$1^6 3^6$	$1^4 2^2 4^4$	$1^4 5^4$	$1^2 2^2 3^2 6^2$	$1^3 7^3$	$1^2 2^4 4^8$	$1^2 11^2$	$1^1 2^7 14^1$	$1^1 3^1 5^1 15^1$	$1^1 23^1$
N	1	2	3	4	5	6	7	8	11	14	15	23
w	12	8	6	5	4	4	3	3	2	2	2	1

g	2^{12}	3^8	$2^4 4^4$	4^6	6^4	$2^2 10^2$	$2^1 4^1 6^1 12^1$	12^2	$3^1 21^1$
n	2	3	4	4	6	10	12	12	21
N	4	9	8	16	36	20	24	144	63
w	6	4	4	3	2	2	2	1	1

M_{24} has the following 9 extra Frame shapes:

²There is a single ambiguity in case $N = 7$: one have to choose $s = 4$ but not $s = 5$.

8 Mason's cusp-forms.

Given a Frame shape $g = \prod i^{a_i}$ consider a function $\eta_g = \prod \eta(q^n)^{a_i}$, where $\eta(q) = q^{\frac{1}{24}} \prod_{m \geq 1} (1 - q^m)$ is Dedekind's eta-function.

Define *weight* of Frame shape as $w(g) = \frac{\sum a_i}{2}$ and *level* as $N(g) = \gcd(a_1, \dots, a_{24}) \cdot \text{lcm}(a_1, \dots, a_{24})$.³

Theorem 8.1 (Mason [18]). *Let g be one of 21 Frame shapes of M_{24} . Then η_g is a cusp-form and Hecke-eigenform of weight $w(g)$ and level $N(g)$ with quadratic nebentypus character (if weight $w(g)$ is even then the character is trivial). Moreover, all these functions η_g form a character of a particular graded representation of M_{24} functorially constructed from M .*

Theorem 8.2 (Dummit, Kisilevsky, McKay [8]; Koike [15]; Mason [18]). *Only for 30 out of 1575 Frame shapes g of S_{24} the respective eta-product η_g is a Hecke eigen-cuspform. There are 2 extra Frame shapes with non-integer weight (24^1 and 8^3) and 7 extra Frame shapes with integer weight:*

g	$3^2 9^2$	$4^2 8^2$	$2^3 6^3$	$2^1 22^1$	$4^1 20^1$	$6^1 18^1$	$8^1 16^1$
n	9	8	6	22	20	18	16
N	27	32	12	44	80	108	128
w	2	2	3	1	1	1	1

It is known all of them come as characters of ex-

tension $2^{11} M_{24}$.

9 Symplectic automorphisms of K3 surfaces.

Theorem 9.1 (Nikulin). *Let g be an automorphism of finite order N on K3 surface S preserving the holomorphic volume form $\omega: g^N = 1, g^* \omega = \omega$. Denote by $F(g)$ the number of its fixed points: $F(g) = \text{Tr} g: H^*(S, \mathbb{Q})$, number $F(g)$ can be computed by combining topological and holomorphic Lefschetz fixed point formulae on K3 surface. Then*

1. *Order of symplectic automorphism is bounded by $N \leq 8$.*
2. *$F(g)$ depends only on the order and $F(g) = \epsilon(N)$.*

Theorem 9.2 (Mukai). *Finite group G acts on K3 surface S preserving the holomorphic volume form $\omega \iff$ the following two conditions are satisfied:*

1. *G can be embedded in M_{23}*
2. *tautological action of G on set S has at least 5 orbits, or in other words – the dimension of invariants of G -representation $H^*(S, \mathbb{Q})$ is at least 5.*

Problem 9.3. *Is there any direct geometric relation between G -Fano threefolds and symplectic automorphisms of K3 surfaces?*

10 Measuring rationality

Definition 10.1. We say that deformation class of smooth Fano threefolds is of *irrational* type if there is at least one irrational (over \mathbb{C}) variety in this family. Otherwise we say it is of *rational* type.

Proposition 10.2. *All G -Fano threefolds of higher rank are rational.*

Proof. One can check it case by case. Two varieties of index Fano index 2 ($\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ and variety of complete flags W) are obviously rational. For six varieties of index one:

1. Projection from $Y_{15} \subset \mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^2$ to $\mathbb{P}^1 \times \mathbb{P}^2$ (contract the first \mathbb{P}^2 and project from a point on the second \mathbb{P}^2 factor) is birational. Also Y_{15} is known to be blowup of W .
2. Projection from $Y_{14} \subset Q \times Q$ to one of the factors is birational (inverse to the blowup of a twisted quartic).
3. Projection from $Y_{12} = X_{(1,1,1,1)} \subset \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ to $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ is birational. In fact it is a blowup with center in elliptic curve, which is the base locus of a pencil in linear system $\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1}(1, 1, 1)$.
4. Projection from $Y_{10} = X_{(1,1),(1,1),(1,1)} \subset \mathbb{P}^3 \times \mathbb{P}^3$ to one of the factors is birational.
5. Varieties $Y_{6,2}$ are divisors of type $(2, 2)$ in $\mathbb{P}^2 \times \mathbb{P}^2$. Projection to the first factor is a conic bundle over \mathbb{P}^2 with degeneracy locus of degree six. By Iskovskikh criterion they are rational.

³In our examples it will be $\min\{i|a_i \neq 0\} \cdot \max\{i|a_i \neq 0\}$.

6. Varieties $Y_{6,3}$ are covers of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ branched in anticanonical divisor. Composition of the double cover and projection to the product of first two factors gives a conic bundle over $\mathbb{P}^1 \times \mathbb{P}^1$.

□

Proposition 10.3. *Let X_N be a deformation class G -Fano threefolds of index $r = 1$ and degree $2N$. Then X_N is of rational type $\iff \epsilon(N) \geq 2$.*

Proof. Case by case. Combine proposition 10.2 and tables in the end of [13]. □

This work was done in 2010 and I reported it in some talks:

- “(Mirror) modularity of Fano threefolds” on 2010.08.24 at Tokyo Metropolitan University,
- “G-Fano threefolds and Mathieu group” on 2011.01.31 at University of Miami,
- “Fano and Mathieu” on 2011.03.28 at the joint seminar of the Sector 4.1 (IITP RAS) and Poncelet French-Russian laboratory “Arithmetic Geometry and Coding Theory” in Moscow,
- “Fano and Mathieu” on 2011.05.12 at *Number Theory and Physics at the Crossroads* workshop in Banff International Research Station,
- “New ways to count up to 15” on 2011.06.03 at Algebraic Geometry Seminar in Kyoto University.

Acknowledgements. Author thanks Tom Coates, Alessio Corti, Vasily Golyshev, Shinobu Hosono, Ludmil Katzarkov, Bumsig Kim, Yuri Prokhorov, Duco van Straten, Todor Milanov, Kyoji Saito, Maria Vlasenko, Noriko Yui for useful discussions.

Funding. This work was supported by World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan, Grant-in-Aid for Scientific Research (10554503) from Japan Society for Promotion of Science, Grant of Leading Scientific Schools (N.Sh. 1987.2008.1 and 4713.2010.1) This is an update for the second part of my earlier SFB45 preprint [9], supported by INTAS 05-100000-8118.

References

- [1] G. Almkvist, C. van Enckevort, D. van Straten, W. Zudilin: *Tables of Calabi–Yau equations*, [arXiv:math/0507430](#).
- [2] A. Beauville: *Fano threefolds and K3 surfaces*, [arXiv:math/0211313](#)
- [3] T. Coates, A. Corti, S. Galkin, A. Kasprzyk: *Quantum periods for 3-dimensional Fano manifolds*, [arXiv:1303.3288](#)
- [4] T. Coates, A. Givental: *Quantum Riemann-Roch, Lefschetz and Serre*, [arXiv:math/0110142](#) Annals of Mathematics, 165 (2007), no. 1, pages 15–53.
- [5] J. H. Conway, S. P. Norton: *Monstrous Moonshine*, Bull. London Math. Soc. 11 (1979), no. 3, 308–339.
- [6] I. V. Dolgachev: *Mirror symmetry for lattice polarized K3 surfaces*, [arXiv:alg-geom/9502005](#)
- [7] C. F. Doran: *Picard-Fuchs uniformization: Modularity of the mirror map and mirror-moonshine*, [arXiv:math/9812162](#).
- [8] D. Dummit, H. Kisilevsky, J. McKay: *Multiplicative products of η -functions*, Contemp. Math, 45, 89–98 (1985).
- [9] S. S. Galkin: *Two instances of fake minimal Fano threefolds II*, 2008, preprint IPMU 11-0099
- [10] S. S. Galkin: *G-minimal varieties are quantum minimal*, preprint IPMU 10-0101
- [11] A. B. Givental: *A mirror theorem for toric complete intersections*, [arXiv:alg-geom/9701016](#), Kashiwara, Masaki (ed.) et al., Topological field theory, primitive forms and related topics. Proceedings of the 38th Taniguchi symposium, Kyoto, Japan, December 9–13, 1996 Boston, MA: Birkhauser. Prog. Math. 160, 141–175.
- [12] V. V. Golyshev: *Classification problems and mirror duality*, [arXiv:math.AG/0510287](#), LMS Lecture Note, ed. N. Young, 338 (2007).
- [13] V. A. Iskovskikh, Y. G. Prokhorov: *Fano varieties. Algebraic geometry, V*, Encyclopaedia Math. Sci, 47, 1999.
- [14] S. Kachru, C. Vafa, *Exact Results for $N = 2$ Compactifications of Heterotic Strings*, [arXiv:hep-th/9505105](#) Nucl.Phys.B450:69–89,1995.

- [15] M. Koike: *On McKay's conjecture*, Nagoya Math. J. Volume 95 (1984), 85–89.
- [16] B. H. Lian, S.-T. Yau, *Arithmetic properties of mirror map and quantum coupling*, [arXiv:hep-th/9411234](#).
- [17] B. H. Lian, S.-T. Yau, *Mirror maps, modular relations and hypergeometric series I*, [arXiv:hep-th/9507151](#).
- [18] G. Mason: *M_{24} and Certain Automorphic Forms*, Contemporary Mathematics 45 (1985), 223–244.
- [19] S. Mori, S. Mukai, *Classification of Fano 3-folds with $b_2 \geq 2$* , *Manuscr. Math.*, **36**:147–162 (1981). Erratum **110**: 407 (2003).
- [20] Yu. G. Prokhorov, *G-Fano threefolds I, II*, [arXiv:1012.4959](#), [arXiv:1101.3854](#).
- [21] H. Verril, N. Yui, *Thompson series, and the mirror maps of pencils of K3 surfaces*, The arithmetic and geometry of algebraic cycles: proceedings of the CRM .
- [22] N. Yui, *Arithmetic of certain Calabi–Yau varieties and mirror symmetry: K3 surfaces and mirror moonshine phenomenon*, Arithmetic Algebraic Geometry, Vol.9, p. 549–564
- [23] D. Zagier: *Elliptic modular forms and their applications*, The 1 – 2 – 3 of Modular Forms: lectures at a summer school in Nordfjordeid, pages 1–103, 2008.