

Alon-Tarsi number of signed planar graphs*

Wei Wang^{a,b}, Jianguo Qian^{a†}

^aSchool of Mathematical Sciences, Xiamen University, Xiamen 361005, P. R. China

^bCollege of Information Engineering, Tarim University, Alar 843300, P. R. China

Abstract

Let (G, σ) be any signed planar graph. We show that the Alon-Tarsi number of (G, σ) is at most 5, generalizing a recent result of Zhu for unsigned case. In addition, if (G, σ) is 2-colorable then (G, σ) has the Alon-Tarsi number at most 4. We also construct a signed planar graph which is 2-colorable but not 3-choosable.

Key words. signed graph; planar graph; list coloring; Alon-Tarsi number

AMS subject classification. 05C15; 05C22; 05C10

1 Introduction

Let G be a simple graph with vertex set $V(G)$ and edge set $E(G)$. A *signed graph* with underling graph G is a pair (G, σ) , where σ is a mapping from $E(G)$ to $\{+1, -1\}$. An edge e is *positive* (resp. *negative*) if $\sigma(e) = +1$ (resp. $\sigma(e) = -1$). In particular, we denote by $(G, +)$ (resp. $(G, -)$) the signed graph (G, σ) if every edge is positive (resp. negative). We often identify $(G, +)$ with the (unsigned) underling graph G .

Recently, based on the work of Zaslavsky [12], Máčajová et al. [7] generalized the concept of chromatic number of an unsigned graph to a signed

*Supported by the National Natural Science Foundation of China under Grant No. 11471273 and 11561058.

†Corresponding author: jgqian@xmu.edu.cn.

graph. For a signed graph (G, σ) and a color set $C \subset \mathbb{Z}$, a *proper coloring* [12] with color set C is a mapping $\phi: V(G) \mapsto C$ such that

$$\phi(u) \neq \sigma(uv)\phi(v) \quad (1)$$

for each edge $uv \in E(G)$. For $k \geq 1$, set $M_k = \{\pm 1, \pm 2, \dots, \pm k/2\}$ if k is even and $M_k = \{0, \pm 1, \pm 2, \dots, \pm(k-1)/2\}$ if k is odd. A (proper) k -coloring of a signed graph (G, σ) is a proper coloring with color set M_k . A signed graph (G, σ) is k -colorable if it admits a k -coloring. The *chromatic number* of (G, σ) , denoted $\chi(G, \sigma)$, is the minimum k for which (G, σ) is k -colorable.

Jin et al. [6] and Schweser et al. [9] further considered the list coloring of signed graphs. For a positive integer k , a k -list assignment of (G, σ) is a mapping L which assigns to each vertex v a set $L(v) \subset \mathbb{Z}$ of k permissible colors. For a k -list assignment L of (G, σ) , an L -coloring is a proper coloring $\phi: V(G) \mapsto \cup_{v \in V(G)} L(v)$ such that $\phi(v) \in L(v)$ for every vertex $v \in V(G)$. We say that (G, σ) is L -colorable if G has an L -coloring. A signed graph (G, σ) is called k -choosable if G is L -colorable for any k -list assignment L . The *list chromatic number* (or *choice number*) $\chi_l(G, \sigma)$ is the minimum k for which G is k -choosable. Clearly, $\chi_l(G, \sigma) \geq \chi(G, \sigma)$. We note that when we restrict the signed graphs (G, σ) to $(G, +)$, both the chromatic number and list chromatic number agree with the ordinary chromatic number and list chromatic number of its underlying graph G . This explains why we can identify $(G, +)$ with G .

Let ' $<$ ' be an arbitrary fixed ordering of the vertices of (G, σ) . In view of (1), we define the *graph polynomial* of (G, σ) as

$$P_{G, \sigma}(\mathbf{x}) = \prod_{u \sim v, u < v} (x_u - \sigma(uv)x_v),$$

where $u \sim v$ means that u and v are adjacent, and $\mathbf{x} = (x_v)_{v \in V(G)}$ is a vector of $|V(G)|$ variables indexed by the vertices of G . It is easy to see that a mapping $\phi: V(G) \mapsto \mathbb{Z}$ is a proper coloring of (G, σ) if and only if $P_{G, \sigma}((\phi(v))_{v \in V(G)}) \neq 0$.

Lemma 1.1. [1](Combinatorial Nullstellensatz) *Let \mathbb{F} be an arbitrary field and let $f = f(x_1, x_2, \dots, x_n)$ be a polynomial in $\mathbb{F}[x_1, x_2, \dots, x_n]$. Suppose that the degree $\deg(f)$ of f is $\sum_{i=1}^n t_i$ where each t_i is a nonnegative integer, and suppose that the coefficient of $\prod_{i=1}^n x_i^{t_i}$ of f is nonzero. Then if S_1, S_2, \dots, S_n are subsets of \mathbb{F} with $|S_i| \geq t_i + 1$, then there are $s_1 \in S_1, s_2 \in S_2, \dots, s_n \in S_n$ so that $f(x_1, x_2, \dots, x_n) \neq 0$.*

Note that $P_{G, \sigma}(\mathbf{x})$ is a homogeneous polynomial. It follows from Lemma 1.1 that if there exists a monomial $c \prod_{v \in V(G)} x_v^{t_v}$ in the expansion of $P_{G, \sigma}(\mathbf{x})$

such that $c \neq 0$ and $t_v < k$ for all $v \in V(G)$, then (G, σ) is k -choosable. Thus, the notion of Alon-Tarsi number of unsigned graphs defined by Jensen and Toft [5] can be naturally extended to signed graphs.

Definition 1.2. The *Alon-Tarsi number* of (G, σ) , denoted $AT(G, \sigma)$, is the minimum k for which there exists a monomial $c \prod_{v \in V(G)} x_v^{t_v}$ in the expansion of $P_{G, \sigma}(\mathbf{x})$ such that $c \neq 0$ and $t_v < k$ for all $v \in V(G)$.

Parallel to the unsigned case, we have

$$AT(G, \sigma) \geq \chi_l(G, \sigma) \geq \chi(G, \sigma).$$

For a subgraph H of G , we use (H, σ) to denote the signed subgraph of (G, σ) restricted on H , i.e., $(H, \sigma) = (H, \sigma|_{E(H)})$. Note that $P_{H, \sigma}(\mathbf{x})$ is a factor of $P_{G, \sigma}(\mathbf{x})$. From Definition 1.2, it is clear that $AT(H, \sigma) \leq AT(G, \sigma)$.

For a vertex v in a signed graph (G, σ) , a *switching* at v means changing the sign of each edge incident to v . For $X \subseteq V(G)$, a switching at X means switching at every vertex in X one by one. Equivalently, a switching at X means changing the sign of every edge with exactly one end in X . We denote the switched graph by (G, σ^X) . In particular, when $X = \{v\}$ we use (G, σ^v) to denote $(G, \sigma^{\{v\}})$. Two signed graphs (G, σ) and (G, σ') are *switching equivalent* if $\sigma' = \sigma^X$ for some $X \subseteq V(G)$.

It is easy to show that two switching equivalent signed graphs have the same chromatic number [7] as well as the same list chromatic number [6, 9]. For the Alon-Tarsi numbers, we have the following similar result.

Proposition 1.3. *If two signed graphs (G, σ) and (G, σ') are switching equivalent then $AT(G, \sigma) = AT(G, \sigma')$.*

Proof. It clearly suffices to consider the case that $\sigma' = \sigma^v$, where $v \in V(G)$. For any edge incident with v , say uv , we have $\sigma^v(uv) = -\sigma(uv)$. We use $T(x_u, x_v)$ and $T^v(x_u, x_v)$ to denote the factors corresponding to this edge in $P_{G, \sigma}(\mathbf{x})$ and $P_{G, \sigma^v}(\mathbf{x})$, respectively. If $u < v$ then $T(x_u, x_v) = x_u - \sigma(uv)x_v$, $T^v(x_u, x_v) = x_u - \sigma^v(uv)x_v$ and hence $T(x_u, x_v) = T^v(x_u, -x_v)$. If $v < u$ then $T(x_u, x_v) = x_v - \sigma(uv)x_u$ and $T^v(x_u, x_v) = x_v - \sigma^v(uv)x_u$ and hence $T(x_u, x_v) = -T^v(x_u, -x_v)$. In either case we have $T(x_u, x_v) = \pm T^v(x_u, -x_v)$. Letting \mathbf{x}^v be obtained from \mathbf{x} by changing x_v to $-x_v$, we have $P_{G, \sigma}(\mathbf{x}) = \pm P_{G, \sigma^v}(\mathbf{x}^v)$. Therefore, for each monomial $\prod_{v \in V(G)} x_v^{t_v}$, the coefficients of this monomial in $P_{G, \sigma}(\mathbf{x})$ and $P_{G, \sigma^v}(\mathbf{x}^v)$ and hence in $P_{G, \sigma^v}(\mathbf{x})$ have the same absolute value. This implies that $AT(G, \sigma) = AT(G, \sigma^v)$. \square

Recently, a few classical results on colorability [4] and choosability [6] of planar graphs were generalized to signed planar graphs. In particular, Jin et al. [6] showed that every signed planar graph is 5-choosable, generalizing the well-known result of Thomassen [10] which states that every (unsigned) planar graph is 5-choosable. Another generalization of Thomassen's result was given by Zhu [11], who showed that $AT(G) \leq 5$ for any planar graph G , which solved an open problem proposed by Hefetz [3]. Considering the above results of Jin et. al [6] and Zhu [11], it is natural to ask whether the Alon-Tarsi number of each signed planar graph is at most 5. The main aim of this paper is to give an affirmative answer to this question.

Theorem 1.4. *For any signed graph (G, σ) , if G is a planar graph then $AT(G, \sigma) \leq 5$.*

In [2], Alon and Tarsi showed that every bipartite planar graph is 3-choosable. The result is sharp as $K_{2,4}$ is a bipartite planar graph and $\chi_l(K_{2,4}) = 3$. The following result is a natural extension of this result for signed planar graphs.

Theorem 1.5. *For any signed graph (G, σ) , if G is planar and 2-colorable then $AT(G, \sigma) \leq 4$. Moreover, there is a signed planar graph which is 2-colorable but not 3-choosable.*

2 Orientation and Alon-Tarsi number for signed graphs

For an unsigned graph G , Alon and Tarsi [2] found a useful combinatorial interpretation of the coefficient for each monomial in the graph polynomial $P_G(\mathbf{x})$ in terms of orientations and Eulerian subgraphs. By defining hypergraph polynomial and hypergraph orientation, Ramamurthi and West [8] generalized the result of Alon and Tarsi to k -uniform hypergraph for prime k . In this section we consider the signed graphs. Instead of using orientations of signed graphs, we use orientations of the underlying graphs and find that the result of Alon and Tarsi has a very natural extension for signed graphs.

Let (G, σ) be a signed graph and ' $<$ ' be an arbitrary fixed ordering of $V(G)$. For an orientation D of the underlying graph G , we denote by (v, u) the oriented edge of D with direction from v to u . We call an oriented edge (v, u) σ -decreasing if $v > u$ and $\sigma(uv) = +1$, that is, (v, u) is positive and oriented from the larger vertex to the smaller vertex. We note that a negative edge

will never be σ -decreasing, no matter how it is oriented. An orientation D of G is called σ -even if it has an even number of σ -decreasing edges and called σ -odd otherwise. For a nonnegative sequence $\mathbf{d} = (d_v)_{v \in V(G)}$, let $\sigma EO(\mathbf{d})$ and $\sigma OO(\mathbf{d})$ denote the sets of all σ -even and σ -odd orientations of G having outdegree sequence \mathbf{d} , respectively.

Lemma 2.1. $P_{G,\sigma}(\mathbf{x}) = \sum (|\sigma EO(\mathbf{d})| - |\sigma OO(\mathbf{d})|) \prod_{v \in V(G)} x_v^{d_v}$, where $\mathbf{d} = (d_v)_{v \in V(G)}$ and the summation is taken over all \mathbf{d} such that $d_v \geq 0$ and $\sum_{v \in V(G)} d_v = |E(G)|$.

Proof. Let D be an arbitrary orientation of G . For each oriented edge $e = (v, u)$, define

$$w(e) = \begin{cases} -x_v, & \text{if } e \text{ is } \sigma\text{-decreasing} \\ x_v, & \text{otherwise.} \end{cases} \quad (2)$$

and $w(D) = \prod_{e \in E(D)} w(e)$. Let d_v be the outdegree of v in D for each $v \in V(G)$ and let t be the number of σ -decreasing edges in D . It is easy to see that

$$w(D) = (-1)^t \prod_{v \in V(G)} x_v^{d_v}. \quad (3)$$

Recall that

$$P_{G,\sigma}(\mathbf{x}) = \prod_{u \sim v, u < v} (x_u - \sigma(uv)x_v).$$

By selecting x_u or $-\sigma(uv)x_v$ from each factor $(x_u - \sigma(uv)x_v)$, we expand $P_{G,\sigma}(\mathbf{x})$ and obtain $2^{|E(G)|}$ monomials, each of which has coefficient ± 1 . For each monomial, we orient the edge uv of G with direction from u to v if, in the factor $(x_u - \sigma(uv)x_v)$, x_u is selected; or from v to u if $-\sigma(uv)x_v$ is selected. This is clearly a bijection between the $2^{|E(G)|}$ monomials and the $2^{|E(G)|}$ orientations of G . Therefore,

$$P_{G,\sigma}(\mathbf{x}) = \sum w(D), \quad (4)$$

where D ranges over all orientations of G .

Let $\mathbf{d} = (d_v)_{v \in V(G)}$ be the sequence of outdegrees of some orientation D . Clearly, $d_v \geq 0$ and $\sum_{v \in V(G)} d_v = |E(G)|$. Note that there are exactly $|\sigma EO(\mathbf{d})|$ (resp. $|\sigma OO(\mathbf{d})|$) σ -even (resp. σ -odd) orientations of G . It follows from (3) and (4) that the coefficient of $\prod_{v \in V(G)} x_v^{d_v}$ in the expansion of $P_{G,\sigma}(\mathbf{x})$ is $|\sigma EO(\mathbf{d})| - |\sigma OO(\mathbf{d})|$. This proves the lemma. \square

For an orientation D of G , a subdigraph H of D is called *Eulerian* if $V(H) = V(D)$ and the indegree of every vertex equals its outdegree. We note

that an Eulerian subdigraph H defined here is not necessarily connected. In particular, a vertex is called *isolated* in H if it has indegree 0 (and therefore, has outdegree 0) in H . Further, H is called σ -even (resp. σ -odd) if H has an even (resp. odd) number of positive edges. Let $\sigma EE(D)$ (resp. $\sigma OE(D)$) denote the set of all σ -even (resp. σ -odd) Eulerian subdigraphs of D .

Lemma 2.2. *Let (G, σ) be a signed graph and D be an orientation of G with outdegree sequence $\mathbf{d} = (d_v)_{v \in V(G)}$. Then the coefficient of $\prod_{v \in V(G)} x_v^{d_v}$ in the expansion of $P_{G, \sigma}(\mathbf{x})$ is equal to $\pm(|\sigma EE(D)| - |\sigma OE(D)|)$.*

Proof. For any orientation $D' \in \sigma EO(\mathbf{d}) \cup \sigma OO(\mathbf{d})$, let $D \oplus D'$ denote the set of all oriented edges of D whose orientation in D' is in the opposite direction. Since D and D' have the same outdegree sequence, $D \oplus D'$ is Eulerian. Moreover, $D \oplus D'$ contains an even number of positive edges if and only if D and D' are both σ -even or both σ -odd.

Now, the map $\tau: D' \mapsto D \oplus D'$ is clearly a bijection between $\sigma EO(\mathbf{d}) \cup \sigma OO(\mathbf{d})$ and $\sigma EE(D) \cup \sigma OE(D)$. If D is σ -even, then τ maps $\sigma EO(\mathbf{d})$ to $\sigma EE(D)$ and maps $\sigma OO(\mathbf{d})$ to $\sigma OE(D)$. In this case $|\sigma EO(\mathbf{d})| = |\sigma EE(D)|$ and $|\sigma OO(\mathbf{d})| = |\sigma OE(D)|$. Thus, $|\sigma EO(\mathbf{d})| - |\sigma OO(\mathbf{d})| = |\sigma EE(D)| - |\sigma OE(D)|$. It follows from Lemma 2.1 that the coefficient of $\prod_{v \in V(G)} x_v^{d_v}$ in the expansion of $P_{G, \sigma}(\mathbf{x})$ is equal to $|\sigma EE(D)| - |\sigma OE(D)|$. Similarly, if D is σ -odd, then the coefficient of $\prod_{v \in V(G)} x_v^{d_v}$ in the expansion of $P_{G, \sigma}(\mathbf{x})$ is equal to $|\sigma OE(D)| - |\sigma EE(D)|$. This proves the lemma. \square

By Lemma 2.2 and Definition 1.2, we have the following characterization of the Alon-Tarsi number $AT(G, \sigma)$.

Corollary 2.3. *For any signed graph (G, σ) , $AT(G, \sigma)$ equals the minimum k for which there exists an orientation D of G such that $|\sigma EE(D)| \neq |\sigma OE(D)|$ and every vertex has outdegree less than k .*

3 Proof of Theorem 1.4

We call a plane graph (a planar graph embedded on the plane) a *near triangulation* if the boundary of the outer face is a cycle, called the outer facial cycle, and the boundaries of all inner faces are triangles.

Definition 3.1. Let (G, σ) be a signed graph where G is a near triangulation with outer facial cycle $v_1 v_2 \cdots v_k$ and let $e = v_1 v_2$. An orientation D of $G - e$ is σ -nice for $G - e$ if the following hold:

- $|\sigma EE(D)| \neq |\sigma OE(D)|$.
- v_1 and v_2 have outdegree 0, v_i has outdegree at most 2 for $i \in \{3, 4, \dots, k\}$, and every interior vertex has outdegree at most 4.

We use the method presented in [11] to prove the following theorem.

Theorem 3.2. *Let (G, σ) be a signed graph where G is a near triangulation with outer facial cycle $C = v_1v_2 \cdots v_k$ and let $e = v_1v_2$. Then $G - e$ has a σ -nice orientation.*

Proof. We prove the theorem by induction on $|V(G)|$. If $|V(G)| = 3$ then $G - e$ is a path $v_2v_3v_1$. Let D be the orientation of $G - e$ such that $E(D) = \{(v_3, v_2), (v_3, v_1)\}$. Clearly, D is σ -nice. Now assume that $|V(G)| > 3$ and the assertion holds for graphs of order less than $|V(G)|$. We shall distinguish two cases, according to whether the outer facial cycle C contains a chord incident with v_k .

First we consider the case that C has a chord $e' = v_kv_j$ where $2 \leq j \leq k-2$ (see Figure 1(a)). In this case $C_1 = v_1v_2 \cdots v_jv_k$ and $C_2 = v_kv_jv_{j+1} \cdots v_{k-1}$ are two cycles of G . For $i \in \{1, 2\}$, let G_i be the subgraph of G formed by C_i and its interior part. By the induction hypothesis, $G_1 - e$ has a σ -nice orientation D_1 , and $G_2 - e'$ has a σ -nice orientation D_2 . We notice that D_1 and D_2 are edge disjoint. Let $D = D_1 \cup D_2$. It is clear that D is an orientation of $G - e$. We will show that D is σ -nice for $G - e$. It can be checked that D satisfies the outdegree condition in Definition 3.1. It remains to check that $|\sigma EE(D)| \neq |\sigma OE(D)|$.

Note that both v_k and v_j have outdegree 0 in D_2 . This implies that v_k and v_j are both isolated in any Eulerian subdigraph of D . Therefore, any Eulerian subdigraph H of D has an edge-disjoint decomposition $H = H_1 \cup H_2$, where H_1 and H_2 are Eulerian subdigraphs in D_1 and D_2 , respectively. Thus, the map $\tau: H \mapsto (H_1, H_2)$ is a bijection between $\sigma EE(D) \cup \sigma OE(D)$ and $(\sigma EE(D_1) \cup \sigma OE(D_1)) \times (\sigma EE(D_2) \cup \sigma OE(D_2))$. Moreover, H is σ -even if and only if both H_1 and H_2 are σ -even, or both are σ -odd. Thus, we have

$$\begin{aligned}
& |\sigma EE(D)| - |\sigma OE(D)| \\
&= (|\sigma EE(D_1) \times \sigma EE(D_2)| + |\sigma OE(D_1) \times \sigma OE(D_2)|) \\
&\quad - (|\sigma EE(D_1) \times \sigma OE(D_2)| + |\sigma OE(D_1) \times \sigma EE(D_2)|) \\
&= (|\sigma EE(D_1)| - |\sigma OE(D_1)|) \cdot (|\sigma EE(D_2)| - |\sigma OE(D_2)|) \\
&\neq 0,
\end{aligned}$$

where the last inequality holds since D_1 and D_2 are σ -nice. This proves that D is a σ -nice orientation of $G - e$.

Next assume that C contains no chord of the form $v_k v_j$ for $j \in \{2, 3, \dots, k-2\}$. Let $v_{k-1}, u_1, u_2, \dots, u_s, v_1$ be the neighbors of v_k and be ordered so that $v_k v_{k-1} u_1, v_k u_1 u_2, \dots, v_k u_s v_1$ are inner facial cycles of G (see Figure 1(b) when $k = 3$ and Figure 1(c) when $k > 3$). Let $G' = G - v_k$. It is clear that G' is a near triangulation with outer facial cycle $v_1 v_2 \dots v_{k-1} u_1 u_2 \dots u_s$. Therefore, by the induction hypothesis, $G' - e$ has a σ -nice orientation D' .




Figure 1. Proof of Theorem 1.4.

If $k = 3$ (i.e., C is a triangle), then let D be the orientation of $G - e$ obtained from D' by adding the vertex v_3 and oriented edges (v_3, v_1) , (v_3, v_2) and (u_i, v_3) for $i \in \{1, 2, \dots, s\}$, as shown in Figure 1(b). It is easy to verify that D satisfies the outdegree condition in Definition 3.1. In particular, both v_1 and v_2 have outdegree 0. Thus, v_1 and v_2 are both isolated in any Eulerian subdigraph of D and therefore, by the definition of D , v_3 is also isolated in any Eulerian subdigraph of D . This means that each Eulerian subdigraph of D is an Eulerian subdigraph of D' by ignoring the isolated vertex v_k . Thus, $\sigma EE(D) = \sigma EE(D')$ and $\sigma OE(D) = \sigma EE(D')$. As D' is σ -nice,

$|\sigma EE(D')| \neq |\sigma OE(D')|$ and hence $|\sigma EE(D)| \neq |\sigma OE(D)|$. This proves that D is a σ -nice orientation of $G - e$.

Now assume that $k \geq 4$. We call an orientation D of $G' - e$ *special* if the following hold:

- v_1 and v_2 have outdegree 0, v_{k-1} has outdegree at most 1, each of v_3, v_4, \dots, v_{k-1} has outdegree at most 2, and each of u_1, u_2, \dots, u_s has outdegree at most 3.
- Every interior vertex has outdegree at most 4.

To show that $G - e$ has a σ -nice orientation, we consider two cases:

Case 1. $G' - e$ has a special orientation D'' with $|\sigma EE(D'')| \neq |\sigma OE(D'')|$.

Let D be the orientation of $G - e$ obtained from D'' by adding the vertex v_k and $s + 2$ oriented edges $(v_k, v_1), (v_{k-1}, v_k)$ and (u_i, v_k) for $i \in \{1, 2, \dots, s\}$, see Figure 1(c). Then D satisfies the outdegree condition of a σ -nice orientation. Since v_1 has outdegree 0 in D , by a similar discussion as above, v_k is isolated in any Eulerian subdigraph of D . Therefore, each Eulerian subdigraph of D is an Eulerian subdigraph of D'' by ignoring the isolated vertex v_k , i.e., $\sigma EE(D) = \sigma EE(D'')$ and $\sigma OE(D) = \sigma OE(D'')$. This yields that $|\sigma EE(D)| \neq |\sigma OE(D)|$ by the condition of this case. Thus, D is a σ -nice orientation of $G - e$, as desired.

Case 2. For any special orientation D'' (if exists), $|\sigma EE(D'')| = |\sigma OE(D'')|$.

Recall that D' is a σ -nice orientation of $G' - e$. Let D be the orientation of $G - e$ obtained from D' by adding the vertex v_k and $s + 2$ oriented edges $(v_k, v_1), (v_k, v_{k-1})$ and (u_i, v_k) for $i \in \{1, 2, \dots, s\}$, as shown in Figure 1(d). Clearly, D satisfies the outdegree condition of a σ -nice orientation. To show that D is σ -nice for $G - e$, it remains to show that $|\sigma EE(D)| \neq |\sigma OE(D)|$.

Notice that v_1 has outdegree 0 in D and therefore, is isolated in any Eulerian subdigraph of D . Thus, if H is an Eulerian subdigraph of D and v_k is non-isolated in H then H contains the oriented edge (v_k, v_{k-1}) and exactly one of the s oriented edges $(u_1, v_k), (u_2, v_k), \dots, (u_s, v_k)$. For $i \in \{1, 2, \dots, s\}$, let

$$\sigma EE_i(D) = \{H \in \sigma EE(D) : (u_i, v_k) \in H\},$$

and similarly,

$$\sigma OE_i(D) = \{H \in \sigma OE(D) : (u_i, v_k) \in H\}.$$

For an Eulerian subdigraph of D' , we regard it as an Eulerian subdigraph of D by adding v_k as an isolated vertex. Then we have

$$\sigma EE(D) = \sigma EE(D') \cup \bigcup_{i=1}^s \sigma EE_i(D), \sigma OE(D) = \sigma OE(D') \cup \bigcup_{i=1}^s \sigma OE_i(D).$$

Since D' is σ -nice, $|\sigma EE(D')| \neq |\sigma OE(D')|$. If we can show that $|\sigma EE_i(D)| = |\sigma OE_i(D)|$ for each $i \in \{1, 2, \dots, s\}$, then $|\sigma EE(D)| \neq |\sigma OE(D)|$ and we are done.

Let i be any integer in $\{1, 2, \dots, s\}$. If $\sigma EE_i(D) \cup \sigma OE_i(D) = \emptyset$ then $|\sigma EE_i(D)| = |\sigma OE_i(D)| = 0$, as desired. Thus, we may assume that $\sigma EE_i(D) \cup \sigma OE_i(D) \neq \emptyset$. Therefore, D has an Eulerian subdigraph and hence a directed cycle containing (u_i, v_k) . Let $C_i = u_i v_k v_{k-1} w_1 w_2 \cdots w_p$ be such a directed cycle and let D'_i be the orientation of $G' - e$ obtained from D' by reversing the direction of edges in the path $v_{k-1} w_1 w_2 \cdots w_p u_i$. The reversing operation decreases the outdegree of v_{k-1} by 1, increases the outdegree of u_i by 1, and leaves the outdegrees of other vertices in $G' - e$ unchanged. Since D' is σ -nice for $G' - e$, the outdegree condition of D' implies that D'_i is special. Hence, $|\sigma EE(D'_i)| = |\sigma OE(D'_i)|$ by the condition of this case.

Let C_i^{-1} be the reverse of C_i , i.e., $C_i^{-1} = w_p w_{p-1} \cdots w_1 v_{k-1} v_k u_i$. For each Eulerian subdigraph $H \in \sigma EE_i(D) \cup \sigma OE_i(D)$, let $H \triangle C_i^{-1}$ be the symmetry difference of the edge sets of H and C_i^{-1} , that is, the set obtained from the edge union $H \cup C_i^{-1}$ of H and C_i^{-1} by deleting the directed 2-cycles. One may verify that $H \triangle C_i^{-1}$ is an Eulerian subdigraph of D'_i and the map $\tau: H \mapsto H \triangle C_i^{-1}$ is a bijection between $\sigma EE_i(D) \cup \sigma OE_i(D)$ and $\sigma EE(D'_i) \cup \sigma OE(D'_i)$.

For a set S of some oriented edges in an orientation of (G, σ) , we use $N^\sigma(S)$ to denote the number of positive edges in S . If S is a directed 2-cycle, then either $N^\sigma(S) = 2$ or $N^\sigma(S) = 0$. Thus, $N^\sigma(H \triangle C_i^{-1})$ and $N^\sigma(H \cup C_i^{-1})$ have the same parity. Of course, $N^\sigma(H \cup C_i^{-1}) = N^\sigma(H) + N^\sigma(C_i^{-1}) = N^\sigma(H) + N^\sigma(C_i)$. Therefore, if $N^\sigma(C_i)$ is even, then $\tau: H \mapsto H \triangle C_i^{-1}$ maps $\sigma EE_i(D)$ to $\sigma EE(D'_i)$ and $\sigma OE_i(D)$ to $\sigma OE(D'_i)$. Similarly, if $N^\sigma(C_i)$ is odd, then it maps $\sigma EE_i(D)$ to $\sigma OE(D'_i)$ and $\sigma OE_i(D)$ to $\sigma EE(D'_i)$. Therefore, we have $|\sigma EE_i(D)| - |\sigma OE_i(D)| = \pm(|\sigma EE(D'_i)| - |\sigma OE(D'_i)|)$. Note that D'_i is special. It follows from the condition of this case that $|\sigma EE_i(D)| = |\sigma OE_i(D)|$. This completes the proof of this theorem. \square

Proof of Theorem 1.4 Note that $AT(H, \sigma) \leq AT(G, \sigma)$ for any subgraph H of G . To show that $AT(G, \sigma) \leq 5$ for any planar graph, it suffices to consider the case when G is a near triangulation. Let $v_1 v_2 \cdots v_k$ be the outer facial

cycle of G and $e = v_1v_2$. By Theorem 3.2, $G - e$ has a σ -nice orientation D . Let D' be obtained from D by adding the oriented edge (v_1, v_2) . Clearly, each vertex has outdegree at most 4 in D' . Moreover, as v_2 has outdegree 0 in D , the orientated edge (v_1, v_2) will never appears in any Eulerian subgraph of D' . Thus, $|\sigma EE(D')| = |\sigma EE(D)|$ and $|\sigma OE(D')| = |\sigma OE(D)|$. As D is σ -nice, we have $|\sigma EE(D)| \neq |\sigma OE(D)|$. Therefore, $|\sigma EE(D')| \neq |\sigma OE(D')|$ and hence $AT(G, \sigma) \leq 5$ by Corollary 2.3.

4 Proof of Theorem 1.5

For a graph G , the maximum average degree of G , denoted $\text{mad}(G)$, is the maximum of $2|E(H)|/|V(H)|$, where H ranges over all subgraphs of G . The following useful criterion on the existence of an orientation with bounded outdegree appeared in [2].

Lemma 4.1. *A graph G has an orientation D such that every vertex has outdegree at most p if and only if $\text{mad}(G) \leq 2p$.*

Corollary 4.2. *For any graph G ,*

$$AT(G, -) = \left\lceil \frac{\text{mad}(G)}{2} \right\rceil + 1. \quad (5)$$

Proof. Let $p = \lceil \frac{\text{mad}(G)}{2} \rceil$. Then $\text{mad}(G) \leq 2p$ and hence, by Lemma 4.1, G has an orientation \bar{D} in which every outdegree is at most p . As all edge in $(G, -)$ is negative, each Eulerian subdigraph of \bar{D} contains no positive edge and hence is σ -even. Thus $|\sigma OE(\bar{D})| = 0$. Since the empty subdigraph is a σ -even Eulerian subdigraph, we have $|\sigma EE(\bar{D})| \geq 1$ and hence $|\sigma EE(\bar{D})| \neq |\sigma OE(\bar{D})|$. Thus by Corollary 2.3, $AT(G, -) \leq p + 1$.

On the other hand, by Corollary 2.3, G has an orientation D such that each outdegree is at most $AT(G, -) - 1$. Thus, by Lemma 4.1, $\text{mad}(G) \leq 2(AT(G, -) - 1)$, i.e., $AT(G, -) \geq \frac{\text{mad}(G)}{2} + 1$. Therefore, $AT(G, -) \geq p + 1$ since $AT(G, -)$ is an integer. This proves the corollary. \square

Proof of Theorem 1.5. For a signed graph (G, σ) , Schweser and Stiebitz [9] showed that $\chi(G, \sigma) \leq 2$ if and only if (G, σ) is switching equivalent to $(G, -)$. Thus, by Proposition 1.3, it suffices to consider the case when $(G, \sigma) = (G, -)$, i.e., $\sigma(uv) = -1$ for each $uv \in E(G)$. Let H be any subgraph of a planar graph G . Then by Euler's formula for planar graph we

have $2|E(H)|/|V(H)| \leq 6$, i.e., $\text{mad}(G) \leq 6$. By Corollary 4.2, $AT(G, -) \leq 4$. This proves the first part of Theorem 1.5.

Let $(G, -)$ be the negative planar graph as shown in Figure 2. We show that $(G, -)$ is not 3-choosable.




Figure 2. A non-3-choosable negative planar graph $(G, -)$.

Define a 3-list assignment L as follows:

- $L(a) = L(a') = \{0, -1, -2\}$.
- $L(b) = L(b') = \{0, -1, 2\}$.
- $L(c) = L(c') = \{0, 1, -2\}$.
- $L(d) = L(d') = \{0, 1, 2\}$.

It suffices to show that $(G, -)$ is not L -colorable. Suppose to the contrary that ϕ is an L -coloring of $(G, -)$. Let $V = \{a, b, c, d\}$.

Claim 1: There exists some $x \in V$ such that $\phi(x) = 0$.

Suppose to the contrary that $\phi(x) \neq 0$ for each $x \in V$. Then $\phi(a) \in \{-1, -2\}$, $\phi(b) \in \{-1, 2\}$, $\phi(c) \in \{1, -2\}$ and $\phi(d) \in \{1, 2\}$. Note that $(G[V], -)$ is a negative complete graph. Thus $\phi(x) \neq -\phi(y)$ for two distinct x, y in V . If $\phi(a) = -1$ then $\phi(c) = -2$ and $\phi(d) = 2$. Now, $\phi(c) = -\phi(d)$, a contradiction. Similarly, if $\phi(a) = -2$ then $\phi(b) = -1$ and $\phi(d) = 1$ and hence $\phi(b) = -\phi(d)$. This is also a contradiction. Thus, Claim 1 follows.

Claim 2: Let $x \in V$. If $\phi(x) = 0$ then $\phi(N(x')) = -L(x')$.

We only prove the case that $x = a$ and the other three cases can be settled in the same way. Since $\phi(a) = 0$, we have $\phi(b) \in \{-1, 2\}$, $\phi(c) \in \{1, -2\}$ and $\phi(d) \in \{1, 2\}$. If $\phi(b) = -1$ then $\phi(c) = -2$ and $\phi(d) = 2$. Thus,

$\phi(c) = -\phi(d)$, a contradiction. Therefore, $\phi(b) = 2$. Similarly, if $\phi(c) = -2$ then $\phi(b) = -1$ and $\phi(d) = 1$. We also have a contradiction as $\phi(b) = -\phi(d)$. Therefore, $\phi(c) = 1$. Finally, as $N(a') = \{a, b, c\}$ and $L(a') = \{0, -1, -2\}$, we have $\phi(N(a')) = \{\phi(a), \phi(b), \phi(c)\} = \{0, 2, 1\} = -L(a')$. This proves Claim 2.

Now, by Claim 1, let $x \in V$ satisfy $\phi(x) = 0$. Then, $\phi(N(x')) = -L(x')$ by Claim 2. As $\phi(x') \in L(x')$ we have $-\phi(x') \in \phi(N(x'))$, that is, $-\phi(x') = \phi(y)$ for some $y \in N(x')$. Thus, ϕ is not proper since $x'y$ is a negative edge. This is a contradiction and hence completes the proof of Theorem 1.5.

References

- [1] N. Alon, Combinatorial Nullstellensatz, *Combin. Probab. Comput.* 8 (1999) 7-29. <http://dx.doi.org/10.1017/S0963548398003411>.
- [2] N. Alon, M. Tarsi, Colorings and orientations of graphs, *Combinatorica* 12(2)(1992)125-134. <http://dx.doi.org/10.1007/BF01204715>.
- [3] D. Hefetz, On two generalizations of the Alon-Tarsi polynomial method, *J. Combina. Theory Ser. B* 101(2011) 403-414. <http://dx.doi.org/10.1016/j.jctb.2010.12.007>.
- [4] L. Hu, X. Li, Every signed planar graph without cycles of length from 4 to 8 is 3-colorable, *Discrete Math.* 341 (2018) 513-519. <http://dx.doi.org/10.1016/j.disc.2017.09.019>.
- [5] T. Jensen, B. Toft, *Graph Coloring Problems*, Wiley, New York, 1995.
- [6] L. Jin, Y. Kang, E. Steffen, Choosability in signed planar graphs, *Europ. J. Combin.* 52(2016)234-243. <http://dx.doi.org/10.1016/j.ejc.2015.10.001>.
- [7] E. Máčajová, A. Raspaud, M. Škoviera, The chromatic number of a signed graph, *Electro. J. Combin.* 23(2016)#P1.14.
- [8] R. Ramamurthi, D. B. West, Hypergraph extension of the Alon-Tarsi list coloring theorem, *Combinatorica* 25(2005)355-366. <http://dx.doi.org/10.1007/s00493-005-0020-8>.
- [9] T. Schweser, M. Stiebitz, Degree choosable signed graphs, *Discrete Math.* 340(2017)882-891. <http://dx.doi.org/10.1016/j.disc.2017.01.007>.
- [10] C. Thomassen, Every planar graph is 5-choosable, *J. Combin. Theory, Ser. B* 62(1) (1994) 180-181. <http://dx.doi.org/10.1006/jctb.1994.1062>.

- [11] X. Zhu, The Alon-Tarsi number of planar graphs, J. Combin. Theory, Ser. B., available online. <http://dx.doi.org/10.1016/j.jctb.2018.06.004>.
- [12] T. Zaslavsky, Signed graph coloring, Discrete Math. 39 (1982) 215-228. [http://dx.doi.org/10.1016/0012-365X\(82\)90144-3](http://dx.doi.org/10.1016/0012-365X(82)90144-3).