

WHEN IS A GIVEN REAL GENERIC OVER L ?

FABIANA CASTIBLANCO AND RALF SCHINDLER

ABSTRACT. In this paper we isolate a new criterion for when a given real x is generic over L in terms of x 's capability of lifting elementary embeddings of initial segments of L .

1. INTRODUCTION

The results established in [2] show that the property of closure under sharps for reals is preserved by certain tree forcing notions such as Sacks, Silver, Mathias, Miller and Laver by proving that enough ground-model embeddings $j : L[x] \rightarrow L[x]$, $x \in {}^\omega\omega$, lift to the generic extension. Here we turn our attention to a related problem:

Question 1.1. *Suppose that $0^\#$ exists. Assume that $x \in {}^\omega\omega$ is such that every elementary embedding $j : L \rightarrow L$ lifts to $j^* : L[x] \rightarrow L[x]$. Is x set generic over L ?*

In this paper we will characterize the reals $x \in V$ which are generic by set forcing over L by means of their capability to lift partial elementary embeddings of L (see theorem 3.2). In order to prove our result, we present a version of Woodin's extender algebra for (partial) extenders which exist in L , and also we introduce the notion of "weak Woodiness," which turns out not to be a large cardinal concept at all.

2. WOODIN'S EXTENDER ALGEBRA, MODIFIED, AND BUKOWSKÝ'S THEOREM

Definition 2.1. For a regular cardinal $\delta \geq \omega_1$ and an ordinal $\mu \leq \delta$ let $\mathcal{L}_{\mu, \delta}$ be the least infinitary language which has constants $\check{\xi}$, all $\xi < \mu$, as well as $\dot{a}$, and which has atomic formulae " $\check{\xi} \in \dot{a}$ ", $\xi < \mu$, and is closed under negation and disjunction of length $< \delta$, i.e.,

- (i) if $\phi \in \mathcal{L}_{\mu, \delta}$, then $\neg\phi \in \mathcal{L}_{\mu, \delta}$, and
- (ii) if $\theta < \delta$ and $\phi_\alpha \in \mathcal{L}_{\mu, \delta}$ for all $\alpha < \theta$, then $\bigvee_{\alpha < \theta} \phi_\alpha \in \mathcal{L}_{\mu, \delta}$.

Each $x \subseteq \mu$, x not necessarily in V , defines a model for the logic $\mathcal{L}_{\mu, \delta}$. Given $\varphi \in \mathcal{L}_{\mu, \delta}$ we may define the meaning of $x \models \varphi$ recursively:

- (1) $x \models \check{\xi} \in \dot{x}$ if and only if $\xi \in x$,
- (2) $x \models \neg\varphi$ if and only if $x \not\models \varphi$, and
- (3) $x \models \bigvee \Gamma$ if and only if $x \models \varphi$ for some $\varphi \in \Gamma$, where Γ is an enumeration of formulas in $\mathcal{L}_{\mu, \delta}$ of length $< \delta$.

In this setting, notice that the statement " $x \models \varphi$ " is absolute between transitive models of ZFC containing x and φ .

Definition 2.2. Let $\mathbb{P} \in V$ be a forcing notion and suppose that g is $\mathbb{P}$ -generic over V . For $\varphi \in \mathcal{L}_{\mu, \delta}$, let

$$A_\varphi^g = \{x \in \wp(\mu) \cap V[g] : x \models \varphi\}$$

Further, if $\Gamma \subseteq \mathcal{L}_{\mu, \delta}$ is a theory we set $A_\Gamma^g = \{x \in \wp(\mu) \cap V[g] : \text{for all } \varphi \in \Gamma, x \models \varphi\}$.

Definition 2.3. Let $\mathbb{P} \in V$ be a forcing notion. We say that a theory Γ in $\mathcal{L}_{\mu, \delta}$ is *consistent* if and only if $A_\Gamma^g \neq \emptyset$ for some $\mathbb{P}$ -generic filter g . If $\Gamma \cup \{\phi\} \subseteq \mathcal{L}_{\mu, \delta}$, then we write $\Gamma \vdash \phi$ iff $\Gamma \cup \{\neg\phi\}$ is inconsistent.

Lemma 2.4. *For every theory $\Gamma \subseteq \mathcal{L}_{\mu, \delta}$ and every $\varphi \in \mathcal{L}_{\mu, \delta}$ the following are equivalent:*

- (1) $\Gamma \vdash \varphi$

Both authors gratefully acknowledge support from the SFB 878 program "Groups, Geometry & Actions" financed by the DFG (Deutsche Forschungsgemeinschaft).

- (2) $A_{\Gamma \cup \{\varphi\}}^g = A_\Gamma^g$ for any g which is $\mathbb{P}$ -generic for a forcing notion $\mathbb{P}$ which makes $\delta^{<\delta}$ countable.
- (3) $A_{\Gamma \cup \{\varphi\}}^g = A_\Gamma^g$ for any g which is $\text{Coll}(\omega, \delta^{<\delta})$ -generic over V .

Proof. Easy. See [3, Lemma 2.2] or [4, Lemma 1.2]. $\square$

Definition 2.5. Let $\delta \geq \omega_1$ be a regular cardinal such that $\delta = \delta^{<\delta}$ and let $\mu \leq \delta$. Let $\mathcal{E}$ be a collection of elementary embeddings $j : M \rightarrow N$ with critical point $\kappa = \kappa_j$ such that M, N are transitive, and $M \models \text{ZFC}^-$, $|N| < \delta$, together with a map $\nu_{\mathcal{E}} : \mathcal{E} \rightarrow \text{OR}$ satisfying $\kappa_j + 1 \leq \nu_{\mathcal{E}}(j) \leq j(\kappa_j)$ for each j . We associate to $\mathcal{E}$, in addition to the axioms and rules for the infinitary logic $\mathcal{L}_{\mu, \delta}$, a set of axioms $A_{\mathcal{E}}$ as follows:

$$A_{\mathcal{E}} : \text{ Whenever } j \in \mathcal{E}, j : M \rightarrow N \text{ and } \vec{\phi} = (\phi_i : i < \kappa_j) \in M \text{ then} \\ \mathbb{W} j(\vec{\phi}) \upharpoonright \nu_{\mathcal{E}}(j) \rightarrow \mathbb{W} \vec{\phi}.$$

Definition 2.6. Given $\mathcal{E}$ as above and $\phi, \psi \in \mathcal{L}_{\mu, \delta}$, we define

$$\phi \sim_{\mathcal{E}} \psi \iff A_{\mathcal{E}} \vdash \psi \leftrightarrow \phi,$$

and we let $[\phi] = \{\psi : \psi \sim_{\mathcal{E}} \phi\}$. We write $\mathbb{P}_{\mathcal{E}} = \{[\phi] : \phi \in \mathcal{L}_{\mu, \delta}, \phi \text{ consistent with } A_{\mathcal{E}}\}$. In $\mathbb{P}_{\mathcal{E}}$ we stipulate

$$[\phi] \leq [\psi] \iff A_{\mathcal{E}} \vdash \phi \rightarrow \psi.$$

Definition 2.7. We say that δ is *weakly Woodin* as being witnessed by $\mathcal{E}$ iff for all $A \subset \delta$ there is some $\kappa < \delta$ such that for all $\alpha < \delta$ there is some $j : M \rightarrow N$, $j \in \mathcal{E}$ with $\kappa_j = \kappa$, $A \cap \kappa \in M$ and $j(A \cap \kappa) \cap \alpha = A \cap \alpha$.

By taking hulls, it is easy to see that every regular cardinal $\delta \geq \omega_1$ is weakly Woodin, so “weak Woodinness” is not a large cardinal concept.

Lemma 2.8. Let $\delta \geq \omega_1$ be regular. Let $\theta > \delta$, $p \in H_{\delta}$, and let $\mathcal{E}$ be the collection of all $j : M \rightarrow N$ such that M and N are transitive, $M \models \text{ZFC}^-$, $|N| < \delta$ and $p \in H_{\kappa_j}^M$. Then $\mathcal{E}$ witness that δ is weakly Woodin.

Proof. Let $A \subset \delta$, and let $X \prec H_{\delta^+}$ with $X \cap \delta \in \delta$, $p \in X$, $|X| < \delta$ and $A \in X$. Let $i : M \cong X$ where M is transitive. Write $\kappa = \kappa(i) = X \cap \delta < \delta$. Of course, $A \cap \kappa = i^{-1}(A)$.

Let $\alpha < \delta$ and let $Y = \text{Hull}^{H_{\delta^+}}(X \cup \{\kappa\} \cup (\alpha + 1)) \prec H_{\delta^+}$. Suppose $k : N \cong Y$, where N is transitive. Setting $j = k^{-1} \circ i$, $j : M \rightarrow N$ has critical point κ , and $j(A \cap \alpha) = k^{-1}(A)$ and $k^{-1}(A) \cap \alpha = A \cap \alpha$ as $k \upharpoonright \alpha + 1 = \text{id}$. So $j(A \cap \kappa) \cap \alpha = A \cap \alpha$. $\square$

Lemma 2.9. Suppose $\delta \geq \omega_1$ is a regular cardinal with $\delta^{<\delta} = \delta$. Let $\mathcal{E}$ witness that δ is weakly Woodin, and let $\nu : \mathcal{E} \rightarrow \text{OR}$ be a map with $\kappa_j + 1 \leq \nu_{\mathcal{E}}(j) \leq j(\kappa_j)$ for each $j \in \mathcal{E}$. Then $\mathbb{P}_{\mathcal{E}}$ as being defined in 2.6 with the theory $A_{\mathcal{E}}$, has the δ -c.c.

Proof. Let $\vec{\phi} = \langle \phi_i : i < \delta \rangle$ be such that $\langle [\phi_i] : i < \delta \rangle$ is an antichain in $\mathbb{P}_{\mathcal{E}}$. As $\delta^{<\delta} = \delta$ and every ϕ_i is a formula in $\mathcal{L}_{\mu, \delta}$, we may code $\vec{\phi}$ by a subset A of δ (so $\vec{\phi}$ may be identified with A). As δ is weakly Woodin, we may pick κ as in 2.7. We have $\kappa + 1 < \delta$, and we may pick $j : M \rightarrow N$ in $\mathcal{E}$ such that if $\kappa = \kappa_j$, then

$$j(\vec{\phi} \upharpoonright \kappa)(\kappa) = \phi_{\kappa}$$

Since $\nu(j) \geq \kappa + 1$, the axioms of $A_{\mathcal{E}}$ tell us that:

$$A_{\mathcal{E}} \vdash \phi_{\kappa} \rightarrow \mathbb{W} j(\vec{\phi} \upharpoonright \kappa) \upharpoonright \nu(j) \rightarrow \mathbb{W} \vec{\phi} \upharpoonright \kappa$$

Therefore, $\{[\phi_i] : i \leq \kappa\}$ is not an antichain, which leads to a contradiction. $\square$

For $x \subset \mu$ such that $x \models A_{\mathcal{E}}$, let $G_x = \{[\phi] \in \mathbb{P}_{\mathcal{E}} : x \models \phi\}$. It is easy to see that $G_x \subset \mathbb{P}_{\mathcal{E}}$ is an ultrafilter. Now, given a $\mathbb{P}_{\mathcal{E}}$ -generic filter G over V , notice that for $x := \{\xi < \mu : “\dot{\xi} \in \dot{x}” \in G\}$ we have $G_x = G$. In such case, we say that x is $\mathbb{P}_{\mathcal{E}}$ -generic over V .

Lemma 2.10. Let $\mathcal{E}$ witness that δ is weakly Woodin, where $\delta^{<\delta} = \delta$. Let $x \subset \mu$, x not necessarily in V but in a transitive outer model $V[x]$ of ZFC , and assume that $x \models A_{\mathcal{E}}$. Then x is $\mathbb{P}_{\mathcal{E}}$ -generic over V .

Proof. We show that $G_x = \{[\varphi] \in \mathbb{P}_\mathcal{E} : x \models \varphi\}$ is $\mathbb{P}_\mathcal{E}$ -generic over V . Let $A = \{[\phi_i] : i < \theta\}$ be a maximal antichain of $\mathbb{P}_\mathcal{E}$ in V . By 2.9 we have that $\mathbb{P}_\mathcal{E}$ is δ -c.c. so $\theta < \delta$ and hence $\bigvee_{i < \theta} \phi_i$ is in $\mathcal{L}_{\mu, \delta}$ and in fact $[\bigvee_{i < \theta} \phi_i] \in \mathbb{P}_\mathcal{E}$. Since A is maximal, we have that $A_\mathcal{E} \vdash \bigvee_{i < \theta} \phi_i$.

We claim that $x \models A_\mathcal{E}$ yields that

$$x \models \bigvee_{i < \theta} \phi_i.$$

Suppose that $x \models A_\mathcal{E} \cup \{\neg \bigvee_{i < \theta} \phi_i\}$. Then in $V[x]$ and hence in $V[x]^{\text{Coll}(\omega, \delta)}$ there is some x' with $x' \models A_\mathcal{E} \cup \{\neg \bigvee_{i < \theta} \phi_i\}$, and by Shoenfield absoluteness there will be an x' in $V^{\text{Coll}(\omega, \delta)}$ with $x' \models A_\mathcal{E} \cup \{\neg \bigvee_{i < \theta} \phi_i\}$. But this contradicts $A_\mathcal{E} \vdash \bigvee_{i < \theta} \phi_i$.

Thus $x \models \phi_i$ for some $i < \theta$, so $G_x \cap A \neq \emptyset$ and hence G_x is $\mathbb{P}_\mathcal{E}$ -generic. $\square$

Lemma 2.11. *Let $\delta, p, \mathcal{E}$ be as in the statement of lemma 2.8. Let $x \subset \mu$, x not necessarily in V , be such that for every $j : M \rightarrow N$ in $\mathcal{E}$, there is some elementary $\hat{j} : M[x] \rightarrow N[x]$ with $\hat{j} \supset j$ and $M[x] \models \text{ZFC}^-$. Then x is $\mathbb{P}_\mathcal{E}$ -generic over V .*

Proof. By the previous lemmas, it suffices to show that $x \models A_\mathcal{E}$. So let $j : M \rightarrow N$ in $\mathcal{E}$ and let $\vec{\phi} = (\phi_i : i < \kappa(j)) \in M$. Let us assume that $x \models \bigvee j(\vec{\phi}) \upharpoonright \nu_\mathcal{E}(j)$. Then

$$N[x] \models "x \models \bigvee j(\vec{\phi}) \upharpoonright \nu_\mathcal{E}(j)"$$

hence by elementarity of $\hat{j}$, $M[x] \models "x \models \bigvee \vec{\phi} \upharpoonright \kappa"$ and so $x \models \bigvee \vec{\phi}$. $\square$

The arguments given so far allow us to reprove a theorem of Bukowský's, see [1], which we state as follows:

Theorem 2.12. *Let $X \subset \mu$, X not necessarily in V but in a transitive outer model $V[X]$ of ZFC, and let δ be a regular uncountable cardinal. The following are equivalent:*

- (1) *There is some $\mathbb{P}$ such that $\mathbb{P}$ has the δ -c.c. and X is $\mathbb{P}$ -generic over V .*
- (2) *There is some $\theta > \mu$ and some club C of $Y \prec H(\theta)$ with $Y \cap \delta \in \delta$, $|Y| < \delta$ such that if $j : M \cong Y$, where M is transitive, then there is some elementary $\hat{j} : M[\bar{X}] \rightarrow H(\theta)[X]$ for some $\bar{X}$ and $H(\theta)[X] \models \text{ZFC}^-$.*
- (3) *If $f : \theta \rightarrow \text{Ord}$, some $\theta, f \in V[x]$, then there is $g : \theta \rightarrow \wp(\text{Ord})$, $g \in V$ such that $|g(\xi)| < \delta$ in V and $f(\xi) \in g(\xi)$ for all $\xi < \theta$.*

Proof. “(2) $\Rightarrow$ (1)”: This is by the proof of Lemma 2.11.

“(3) $\Rightarrow$ (2)”: Let $\theta > \mu$ be sufficiently big. Notice that for all X , $H(\theta) \subset H(\theta)^{V[X]} = H(\theta)[X]$. Doing a suitable book-keeping, let $\tilde{h} : \omega \times (H(\theta)^{V[G]})^{<\omega} \rightarrow H(\theta)^{V[G]} \in V[G]$ be a Skolem function. Thus, for all $A \subset H(\theta)$, $\tilde{h}'' A \subset H(\theta)^{V[G]}$.

Let us define $h : \theta \rightarrow \text{Ord}$ by using $\tilde{h}$ as follows:

$$h(x) = \begin{cases} \tilde{h}(x) & \text{if } \tilde{h}(x) \in H(\theta) \\ \emptyset & \text{otherwise} \end{cases}$$

By using (3), we may find some $g \in V$ such that for all $\xi < \theta$, $|g(\xi)| < \delta$ and $h(\xi) \in g(\xi)$. In particular, for every $A \subset H(\theta)$ of size $< \delta$ in V , $g'' A \cap \delta \in \delta$ and $h'' A \subset g'' A$.

Notice that if $\tilde{h}(x) \in V$, $x \in A$, then $\tilde{h}(x) = h(x) \in g(x) \subset A$, so $\tilde{h}'' A \cap V \subset A$. On the other side, as $\tilde{h}$ is a Skolem function we have that $A \subset \tilde{h}'' A \cap V$ for $A \in V$. Thus if $A \in V$ then $\tilde{h}'' A \cap V = A$. This shows that for every $A \in V$, $A \subset H(\theta)$, $A = g'' A$ holds. Therefore, as $\tilde{h}'' A \prec H(\theta)^{V[G]}$ and $\tilde{h}'' A \cap V = A$, we have that $A \prec H(\theta)^V$.

According to this, let us define C as the collection of all g -closed $Y \prec H(\theta)$ with $Y \cap \delta \in \delta$ and $|Y| < \delta$. By construction, we have that the collection C satisfy (2).

“(1) $\Rightarrow$ (3)”: This is a standard argument. $\square$

Question 2.13. *Suppose that $\mathcal{E} \subset V_\delta$ is a collection of extenders. Define $A_\mathcal{E}$ as in section 2, that is for each extender $E \in \mathcal{E}$ and for every sequence of formulae $\vec{\varphi} = (\varphi_i : i < \text{crit}(E))$ we associate the axiom*

$$\bigvee i_E(\vec{\varphi}) \upharpoonright \nu(E) \rightarrow \bigvee \vec{\varphi}$$

where $i_E : M \rightarrow \text{Ult}(M, E)$ and $\nu(E)$ is the strength of the extender. Assume that $\mathbb{P}_\mathcal{E}$ has the δ -c.c. Is δ Woodin?

3. A CRITERION FOR SET-GENERICITY OVER L

Proposition 3.1 (Kunen). *Let $j : L \rightarrow L$ be an elementary embedding with critical point κ . Then, for all $\alpha < \kappa^{+L}$, $j \restriction L_\alpha \in L$.*

Proof. This is by the “ancient Kunen argument.” Let $\alpha < \kappa^{+L}$ and pick $f : \kappa \rightarrow L_\alpha$ onto, $f \in L$. Note that for every $x \in L_\alpha$, $j(x) = y$ if and only if

$$\exists \xi < \kappa (x = f(\xi) \wedge y = j(f)(\xi))$$

Since $j(f) \in L$, for every $x \in L_\alpha$ we can compute $j(x)$ in L from f and $j(f)$. Therefore, $j \restriction L_\alpha \in L$ as required. $\square$

Theorem 3.2. *The following are equivalent for a given real $x \in V$:*

- (1) *x is set-generic over L*
- (2) *There is some $p \in L$ such that for all elementary $j : L_\alpha \rightarrow L_\beta$ with critical point κ , where $j \in L$, $L_\alpha \models \text{ZFC}^-$ and $p \in L_\kappa (\subsetneq L_\alpha)$, there is some $\hat{j} : L_\alpha[x] \rightarrow L_\beta[x]$ with $\hat{j} \supset j$ and $L_\alpha[x] \models \text{ZFC}^-$.*

Proof. “ $\Leftarrow$ ”: Given p , let $\delta \geq \omega_1$ be regular with $p \in L_\delta$. Let $\mathcal{E}$ be defined in L as in the statement of lemma 2.8. Then if $\mathbb{P}_\mathcal{E}$ is defined as in definition 2.6 inside L , x is $\mathbb{P}_\mathcal{E}$ -generic over L .

“ $\Rightarrow$ ”: Let $\mathbb{P} \in L$ be such that x is $\mathbb{P}$ -generic over L . Write $\mu = \text{Card}^L(\mathbb{P})$ and let $p \in L_{\mu^+}$, where $\mu^+ = \mu^{+L}$. Let $j : L_\alpha \rightarrow L_\beta$ with $\text{crit}(j) = \kappa$, $\mu^+ < \kappa$, $L_\alpha \models \text{ZFC}^-$.

Without loss of generality, let us assume that $\mathbb{P} \in p \in L_\kappa$, so that x is $\mathbb{P}$ -generic over L_α . This gives $L_\alpha[x] \models \text{ZFC}^-$. The real x is also $\mathbb{P}$ -generic over L_β , and by writing $X = \text{rng}(j)$, $\mathbb{P} \in p \in X$. In order to see that there is $\hat{j} : L_\alpha[x] \rightarrow L_\beta[x]$ with $\hat{j} \supset j$ it suffices to verify that $X[x] \cap \beta = X \cap \beta^1$. Let $\tau \in X \cap L^\mathbb{P}$ be a name for an ordinal. Note that $B = \{\gamma : \exists p \in \mathbb{P} (p \Vdash_{L_\beta}^\mathbb{P} \tau = \hat{\gamma})\} \in X$ and $\text{otp}(B) < \mu^+$. Thus, the order isomorphism $\pi : \text{otp}(B) \cong B$ is in X , and since $\mu^+ < \kappa \subset X$, we have $\text{otp}(B) \cup \{\text{otp}(B)\} \subseteq X$. Hence $B \subseteq X$ and, in particular, $\tau^x \in X$. $\square$

The paramater p as in the statement (2) above is necessary. Suppose that $j : L_\alpha \rightarrow L_\beta$ is an elementary embedding with critical point κ , and let x be $\text{Coll}(\omega, \kappa)$ -generic over L . In this case, such an embedding cannot be lifted to a $\hat{j} : L_\alpha[x] \rightarrow L_\beta[x]$ because $\text{crit}(j)^{L_\alpha[x]}$ is countable.

By Theorem 3.2, if $x \in \mathbb{R} \cap V$ is such that (2) of the statement of Theorem 3.2 holds true and $0^\#$ exists, then $x^\#$ exists.

REFERENCES

- [1] BUKOWSKÝ, L. Characterization of generic extensions of models of set theory. *Fundamenta Mathematica* 83 (1973), 35–46.
- [2] CASTIBLANCO, F., AND SCHLICHT, P. Preserving levels of projective determinacy by tree forcings. Preprint, 2018.
- [3] DOEBLER, P., AND SCHINDLER, R. The extender Algebra and Vagaries of Σ_1^2 absoluteness. *Münster J. of Math.* (2008), 99999–100048.
- [4] FARAH, I. The extender algebra and Σ_1^2 -absoluteness. to appear in The Cabal Seminar, vol IV.

FABIANA CASTIBLANCO, INSTITUT FÜR MATEMATISCHE LOGIK UND GRUNDLAGENFORSCHUNG, UNIVERSITÄT MÜNSTER,
EINSTEINSTRASSE 62, 48149 MÜNSTER, GERMANY
E-mail address: fabi.cast@wwu.de

RALF SCHINDLER, INSTITUT FÜR MATEMATISCHE LOGIK UND GRUNDLAGENFORSCHUNG, UNIVERSITÄT MÜNSTER,
EINSTEINSTRASSE 62, 48149 MÜNSTER, GERMANY
E-mail address: rds@wwu.de

¹We pretend x is the $\mathbb{P}$ -generic filter.