UNIVERSAL MIXED SUMS OF GENERALIZED 4- AND 8-GONAL NUMBERS

JANGWON JU AND BYEONG-KWEON OH

ABSTRACT. An integer of the form $P_m(x) = \frac{(m-2)x^2 - (m-4)x}{2}$ for an integer x, is called a generalized m-gonal number. For positive integers $\alpha_1, \ldots, \alpha_u$ and β_1, \ldots, β_v , a mixed sum $\Phi = \alpha_1 P_4(x_1) + \cdots + \alpha_u P_4(x_u) + \beta_1 P_8(y_1) + \cdots + \beta_v P_8(y_v)$ of generalized 4- and 8-gonal numbers is called universal if $\Phi = N$ has an integer solution for any nonnegative integer N. In this article, we prove that there are exactly 1271 proper universal mixed sums of generalized 4- and 8-gonal numbers. Furthermore, the "61-theorem" is proved, which states that an arbitrary mixed sum of generalized 4- and 8-gonal numbers is universal if and only if it represents the integers 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 15, 18, 20, 30, 60, and 61.

1. INTRODUCTION

The famous Lagrange's four-square theorem says that every nonnegative integer can be written as a sum of at most four squares of integers. Motivated by this celebrated four-square theorem, Ramanujan provided a list of 55 candidates of diagonal quaternary integral quadratic forms that represent all nonnegative integers. Dickson pointed out that the diagonal quaternary quadratic form $x^2 + 2y^2 + 5z^2 + 5t^2$ in the Ramanujan's list does not represent the integer 15, and confirmed that Ramanujan's assertion is correct for all the other 54 quaternary forms (for details, see [3]). Recently, Conway, Miller, and Schneeberger proved the so called "15-theorem", which states that a positive definite integral quadratic form represents all nonnegative integers if and only if it represents the integers 1, 2, 3, 5, 6, 7, 10, 14, and 15, irrespective of its rank (for details, see [1]). As a natural generalization of the 15theorem, Bhargava and Hanke [2] proved the so-called "290-theorem", which states that every positive-definite integer-valued quadratic form represents all nonnegative integers if and only if it represents

1, 2, 3, 5, 6, 7, 10, 13, 14, 15, 17, 19, 21, 22, 23, 26, 29, 30, 31, 34, 35, 37, 42, 58, 93, 110, 145, 203, and 290.

Key words and phrases. Generalized polygonal numbers, Universal mixed sums.

²⁰¹⁰ Mathematics Subject Classification. Primary 11E12, 11E20.

This work of the first author was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education(NRF-2018R1A6A3A01012662).

This work of the second author was supported by the National Research Foundation of Korea (NRF-2017R1A2B4003758).

Here a quadratic form $f(x_1, x_2, ..., x_n) = \sum_{1 \leq i,j \leq n} a_{ij} x_i x_j$ $(a_{ij} = a_{ji})$ is called *integral* if $a_{ij} \in \mathbb{Z}$ for any i, j, and is called *integer-valued* if $a_{ii} \in \mathbb{Z}$ and $a_{ij} + a_{ji} \in \mathbb{Z}$ for any i, j.

For an integer $m \ge 3$, a polygonal number of order m (or an m-gonal number) is defined by

$$P_m(x) = \frac{(m-2)x^2 - (m-4)x}{2},$$

for some nonnegative integer x. If we admit that x is a negative integer, then $P_m(x)$ is called a generalized polygonal number of order m (or a generalized mgonal number). Lagrange's four-square theorem implies that the equation

$$P_4(x) + P_4(y) + P_4(z) + P_4(t) = N$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$, for any nonnegative integer N.

As a natural generalization of Ramanujan's result on diagonal quaternary quadratic forms representing all nonnegative integers, it was proved in [5] and [13] that there are exactly 40 quaternary sums of generalized octagonal numbers which represent all nonnegative integers. Furthermore, the "octagonal theorem of sixty" was proved in [5], which states that a sum $a_1P_8(x_1) + a_2P_8(x_2) + \cdots + a_kP_8(x_k)$ of generalized octagonal numbers represents all nonnegative integers if and only if it represents

1, 2, 3, 4, 6, 7, 9, 12, 13, 14, 18, and 60,

which might be considered this as a natural generalization of the "15-theorem".

In this article, we generalize the above theorem to the mixed sum of generalized 4- and 8-gonal numbers. Let $\alpha_1, \ldots, \alpha_u$, and β_1, \ldots, β_v be positive integers. The mixed sum

$$\Phi = \alpha_1 P_4(x_1) + \dots + \alpha_u P_4(x_u) + \beta_1 P_8(y_1) + \dots + \beta_v P_8(y_v)$$

of generalized 4- and 8-gonal numbers is said to be *universal* if the diophantine equation $\Phi = N$ has an integer solution for any nonnegative integer N. The aim of this article is to prove that there are exactly 1271 *proper universal* mixed sums of generalized 4- and 8-gonal numbers. Here, a universal sum Φ of generalized 4- and 8-gonal numbers is called *proper* if there does not exist a proper sum of Φ that is universal. Furthermore, in Section 5, we prove that a mixed sum of generalized 4and 8-gonal numbers is universal if and only if it represents the integers

 $1, \ 2, \ 3, \ 4, \ 5, \ 6, \ 7, \ 8, \ 9, \ 10, \ 12, \ 13, \ 14, \ 15, \ 18, \ 20, \ 30, \ 60, \ and \ 61,$

which might also be considered as a generalization of the "15-theorem". Let

$$f(x_1, x_2, \dots, x_n) = \sum_{1 \le i, j \le n} a_{ij} x_i x_j \ (a_{ij} = a_{ji} \in \mathbb{Z})$$

be a positive definite integral quadratic form. The corresponding symmetric matrix of f is defined by $M_f = (a_{ij})$. For a diagonal quadratic form $f(x_1, x_2, \ldots, x_n) = a_1 x_1^2 + a_2 x_2^2 + \cdots + a_n x_n^2$, we simply write

$$f = \langle a_1, a_2, \dots, a_n \rangle.$$

For an integer k, we say k is represented by f if the equation $f(x_1, x_2, \ldots, x_n) = k$ has an integer solution. The genus of f, denoted by gen(f), is the set of all quadratic forms that are isometric to f over the p-adic integer ring \mathbb{Z}_p for any prime p. The number of isometry classes in gen(f) is called the class number of f, and denoted by h(f).

Any unexplained notations and terminologies can be found in [8] or [12].

2. General tools

Let $\alpha_1, \ldots, \alpha_u$ and β_1, \ldots, β_v be positive integers such that $\alpha_1 \leq \cdots \leq \alpha_u$ and $\beta_1 \leq \cdots \leq \beta_v$. Let

(2.1)
$$\Phi = \alpha_1 P_4(x_1) + \dots + \alpha_u P_4(x_u) + \beta_1 P_8(y_1) + \dots + \beta_v P_8(y_v)$$

be a mixed sum of generalized 4- and 8-gonal numbers. For simplicity, we will use the notation

$$\Phi = \llbracket \alpha_1, \ldots, \alpha_u, \beta_1, \ldots, \beta_v \rrbracket$$

Recall that Φ is universal if the diophantine equation $\Phi = N$ has an integer solution $(x_1, \ldots, x_u, y_1, \ldots, y_v) \in \mathbb{Z}^{u+v}$ for any nonnegative integer N. One may directly check that Φ is universal if and only if the equation

$$3\alpha_1 x_1^2 + \dots + 3\alpha_u x_u^2 + \beta_1 (3y_1 - 1)^2 + \dots + \beta_v (3y_v - 1)^2 = 3N + \beta_1 + \dots + \beta_v$$

has an integer solution for any nonnegative integer N. This is equivalent to the existence of an integer solution $(x_1, \ldots, x_u, y_1, \ldots, y_v) \in \mathbb{Z}^{u+v}$ of

(2.2)
$$\begin{aligned} 3\alpha_1 x_1^2 + \dots + 3\alpha_u x_u^2 + \beta_1 y_1^2 + \dots + \beta_v y_v^2 \\ &= 3N + \beta_1 + \dots + \beta_v \text{ with } y_1 \dots y_v \not\equiv 0 \pmod{3}. \end{aligned}$$

Note that Equation (2.2) corresponds to the representation of a quadratic form with congruence condition. Since there are some method on determineing the existence of the representations of integers by a quadratic form, we try to find a suitable method on reducing Equation (2.2) to the representation of a quadratic form without congruence condition. To explain our method, for example, assume that $\beta_1 + \cdots + \beta_v \neq 0 \pmod{3}$. Since at least one of β_i is not divisible by 3, we may assume that, without loss of generality, $\beta_v \neq 0 \pmod{3}$. Then (2.2) has an integer solution if and only if

$$3\alpha_1 x_1^2 + \dots + 3\alpha_u x_u^2 + \beta_1 (y_v - 3y_1)^2 + \dots + \beta_{v-1} (y_v - 3y_{v-1})^2 + \beta_v y_v^2 = 3N + \beta_1 + \dots + \beta_v$$

has an integer solution. Hence, in this case, the problem can be reduced to the representations of a quadratic form. Now, assume that $\beta_1 + \cdots + \beta_v \equiv 0 \pmod{3}$. As a representative case, suppose that $u \ge 3$, and the set of all integers that are represented by $3\alpha_1x_1^2 + \cdots + 3\alpha_ux_u^2$ is known. In this case, we try to find sets S_i of integers not divisible by 3 such that for each $i = 1, 2, \ldots, v$, there is a $y_i \in S_i$ depending on N such that

(2.3)
$$3\alpha_1 x_1^2 + \dots + 3\alpha_u x_u^2 = 3N + \beta_1 + \dots + \beta_v - (\beta_1 y_1^2 + \dots + \beta_v y_v^2)$$

has an integer solution $(x_1, \ldots, x_u) \in \mathbb{Z}^u$.

In [6], [9], and [11], we developed a method that determines whether or not integers in an arithmetic progression are represented by some particular ternary quadratic form. We briefly introduce this method for those who are unfamiliar with it.

Let d be a positive integer and let a be a nonnegative integer $(a \leq d)$. We define

$$S_{d,a} = \{dn + a \mid n \in \mathbb{N} \cup \{0\}\}$$

For integral ternary quadratic forms f, g, we define

$$R(g,d,a) = \{ v \in (\mathbb{Z}/d\mathbb{Z})^3 \mid vM_g v^t \equiv a \pmod{d} \}$$

and

$$R(f, g, d) = \{ T \in M_3(\mathbb{Z}) \mid T^t M_f T = d^2 M_g \}.$$

A coset (or, a vector in the coset) $v \in R(g, d, a)$ is said to be *good* with respect to f, g, d, and a if there is a $T \in R(f, g, d)$ such that $\frac{1}{d} \cdot vT^t \in \mathbb{Z}^3$. The set of all good vectors in R(g, d, a) is denoted by $R_f(g, d, a)$. If $R(g, d, a) = R_f(g, d, a)$, we write

$$g \prec_{d,a} f.$$

If $g <_{d,a} f$, then by Lemma 2.2 of [9], we have

$$(2.4) S_{d,a} \cap Q(g) \subset Q(f).$$

We denote the set $R(g, d, a) \setminus R_f(g, d, a)$ by $B_f(g, d, a)$. In general, if d is large, then the set $B_f(g, d, a)$ has too many vectors to compute it exactly by hand. A MAPLE based computer program for computing this set is available upon request to the authors.

Theorem 2.1. Under the same notation given above, assume that $T \in M_3(\mathbb{Z})$ satisfies the following conditions:

(i) $\frac{1}{d}T$ has an infinite order;

(ii) $\overset{a}{T}{}^{t}M_{g}T = d^{2}M_{g};$

(iii) for any vector $v \in \mathbb{Z}^3$ such that $v \pmod{d} \in B_f(g, d, a), \frac{1}{d} \cdot vT^t \in \mathbb{Z}^3$.

Then we have

$$S_{d,a} \cap Q(g) \setminus \{g(z) \cdot s^2 \mid s \in \mathbb{Z}\} \subset Q(f)$$

where the vector z is any integral primitive eigenvector of T.

Proof. See Theorem 2.3 of [6].

3. Ternary mixed sums of generalized 4- and 8-gonal numbers

In this section, we determine all ternary universal mixed sums of generalized 4and 8-gonal numbers.

Let Φ be a mixed sum of generalized 4- and 8-gonal numbers defined in (2.1). For an integer N, if the diophantine equation $\Phi = N$ has an integer solution, then we say the mixed sum Φ represents N, and we write $N \to \Phi$. When the mixed sum Φ is not universal, the least positive integer that is not represented by Φ is called the *truant* of Φ , and is denoted by $t(\Phi)$. One may easily show that there does not exist a binary universal sum of generalized polygonal numbers. For the ternary case, we have the following:

Theorem 3.1. There are exactly 6 ternary universal mixed sums of generalized 4and 8-gonal numbers. In fact, they are

$$(3.1) [[1,1,1]], [[1,1,2]], [[1,3,1]], [[1,1,1]], [[2,3,1]], and [[2,1,1]].$$

Proof. Let u, v be nonnegative integers such that u + v = 3. For positive integers $\alpha_1, \ldots, \alpha_u$ and β_1, \ldots, β_v , assume that a ternary mixed sum

 $\Phi = \llbracket \alpha_1, \dots, \alpha_u, \beta_1, \dots, \beta_v \rrbracket$

of generalized 4- and 8-gonal numbers is universal. At first, we find all candidates of ternary mixed sum Φ of generalized 4-and 8-goanl numbers by using, so called, the escalation method. Since $1 \to \Phi$, $\alpha_1 = 1$ or $\beta_1 = 1$. If $\alpha_1 = 1$, then $1 \leq \alpha_2 \leq 2$ or $1 \leq \beta_1 \leq 2$, for $2 \to \Phi$ and $t(\llbracket 1 \rrbracket) = 2$. If $\alpha_1 \neq 1$ and $\beta_1 = 1$, then $\alpha_1 = 2$ or $1 \leq \beta_2 \leq 2$, for $2 \to \Phi$ and $t(\llbracket 1 \rrbracket) = 2$. Let u_1, v_1 be nonnegative integers such that $u_1 + v_1 = 2$ and

$$\Phi' = \llbracket \alpha_1, \ldots, \alpha_{u_1}, \beta_1, \ldots, \beta_{v_1} \rrbracket$$

is one of the above proper sums of Φ . Since $t(\Phi')$ is represented by Φ for each possible case, at least one of α_{u_1+1} or β_{v_1+1} is less than equal to $t(\Phi')$ given in Table 3.1. From this, one may easily show that there are 42 candidates of ternary universal mixed sums of generalized 4- and 8-gonal numbers (see Table 3.1). Among them, 36 ternary sums are, in fact, not universal (for truants of these sums, see Table (4.1)). Since the proofs of the universalities of the remaining 6 candidates are quite similar to each other, we only provide the proof of the case (3-1).

Table 3.1. Ternary universal mixed sums

Case	Φ'	$t(\Phi')$	C. A.	Possible Candidates		Universal Cases	
(3-1)	$[\![1,1]\!]$	3		$1 \leq \alpha_3 \leq 3,$	$1\leqslant\beta_1\leqslant3$		$\beta_1 = 1, 2$
(3-2)	$[\![1,2]\!]$	5		$2 \leqslant \alpha_3 \leqslant 5,$	$1\leqslant\beta_1\leqslant5$		
(3-3)	$[\![1, 1]\!]$	3	$\alpha_2 = 1, 2$	$\alpha_2 = 3,$	$1\leqslant\beta_2\leqslant 3$	$\alpha_2 = 3,$	$\beta_2 = 1$
(3-4)	$[\![1, 2]\!]$	5	$\alpha_2 = 1, 2$	$3 \leqslant \alpha_2 \leqslant 5,$	$2 \leqslant \beta_2 \leqslant 5$		
(3-5)	$\llbracket 2, 1 \rrbracket$	4		$2 \leqslant \alpha_2 \leqslant 4,$	$1\leqslant\beta_2\leqslant4$	$\alpha_2 = 3,$	$\beta_2 = 1$
(3-6)	$[\![1,1]\!]$	3	$\alpha_1 = 1, 2$	$\alpha_1 = 3,$	$1\leqslant\beta_3\leqslant3$		
(3-7)	$[\![1,2]\!]$	4	$\alpha_1 = 1, 2$	$3 \leqslant \alpha_1 \leqslant 4,$	$2 \leqslant \beta_4 \leqslant 4$		

C. A.= Considered Already

Case (3-1) $(\alpha_1, \alpha_2) = (1, 1)$. It suffices to show that for $\beta_1 \in \{1, 2\}$, the equation $3x^2 + 3y^2 + \beta_1 z^2 = 3N + \beta_1$

has an integer solution $(x, y, z) \in \mathbb{Z}^3$ such that $z \neq 0 \pmod{3}$. Since $h(\langle \beta_1, 3, 3 \rangle) = 1$, one may easily show that a positive integer m congruent to β_1 modulo 3 is represented by $\langle \beta_1, 3, 3 \rangle$. This competes the proof.

4. QUATERNARY MIXED SUMS OF GENERALIZED 4- AND 8-GONAL NUMBERS

In [3], Dickson proved that there are exactly 54 quaternary universal sums of squares. Note that any square of an integer is a generalized 4-gonal numbers. Recently, it was proved in [5] and [13] that there are exactly 40 quaternary universal sum of generalized 8-gonal numbers. In this section, we determine all quaternary

proper universal mixed sums of generalized 4- and 8-gonal numbers including the above 94 quaternary universal sums.

Recall that a mixed sum $\Phi = [\![\alpha_1, \ldots, \alpha_u, \beta_1, \ldots, \beta_v]\!]$ of generalized 4-and 8goanl numbers is universal if and only if the equation

(4.1)
$$\begin{array}{c} 3\alpha_1 x_1^2 + \dots + 3\alpha_u x_u^2 + \beta_1 y_1^2 + \dots + \beta_v y_v^2 \\ = 3N + \beta_1 + \dots + \beta_v \text{ with } y_1 \dots y_v \not\equiv 0 \pmod{3} \end{array}$$

has an integer solution $(x_1, \ldots, x_u, y_1, \ldots, y_v) \in \mathbb{Z}^{u+v}$ for any nonnegative integer N. Let

$$E = \{1 \leq n < \beta_1 + \dots + \beta_v \mid n \equiv \beta_1 + \dots + \beta_v \pmod{3}\}.$$

For each $n \in E$, let $\nu(n)$ be the positive integer such that

$$4^{\nu(n)-1} \cdot n < \beta_1 + \dots + \beta_v \leq 4^{\nu(n)} \cdot n$$

Lemma 4.1. Under the same notations given above, assume that

(4.2)
$$\begin{aligned} 3\alpha_1 x_1^2 + \dots + 3\alpha_u x_u^2 + \beta_1 y_1^2 + \dots + \beta_v y_v^2 \\ = 4^{\nu(n)} \cdot n \quad with \quad y_1 \cdots y_v \not\equiv 0 \pmod{3} \end{aligned}$$

has an integer solution for any $n \in E$. Then the mixed sum Φ is universal if and only if the equation

$$3\alpha_1 x_1^2 + \dots + 3\alpha_u x_u^2 + \beta_1 y_1^2 + \dots + \beta_v y_v^2$$

= $3N + \beta_1 + \dots + \beta_v$ with $y_1 \dots y_v \neq 0 \pmod{3}$

has an integer solution for any nonnegative integer N such that $3N + \beta_1 + \cdots + \beta_v \not\equiv 0 \pmod{4}$.

Proof. Note that "only if" part is trivial. To prove "if" part, let N be any nonnegative integer. Let s be a positive integer and n be a positive integer not divisible by 4 such that $3N + \beta_1 + \cdots + \beta_v = 4^s \cdot n$. If $n \ge \beta_1 + \cdots + \beta_v$, then there is a nonnegative integer N_0 such that $3N_0 + \beta_1 + \cdots + \beta_v = n$. Then Equation (4.1) has an integer solution for this N_0 by assumption, and therefore Equation (4.1) has also an integer solution for N. If $n < \beta_1 + \cdots + \beta_v$, then $n \in E$ and $\nu(n) \le s$ from the choice of $\nu(n)$. Since Equation (4.2) has an integer solution for $4^{\nu(n)} \cdot n$, Equation (4.1) has an integer solution for N.

Theorem 4.2. There are exactly 547 proper quaternary universal mixed sums of generalized 4- and 8-gonal numbers (for the list of them, see Table 4.1).

Proof. Let u, v be nonnegative integers such that u + v = 4. For positive integers $\alpha_1, \ldots, \alpha_u$ and β_1, \ldots, β_v , assume that a quaternary mixed sum

$$\Phi = \llbracket \alpha_1, \ldots, \alpha_u, \beta_1, \ldots, \beta_v \rrbracket$$

is proper universal. Then by theorem 3.1, there are nonnegative integers u_1, v_1 such that $u_1 + v_1 = 3$ and the proper sum

$$\Phi' = \llbracket \alpha_1, \ldots, \alpha_{u_1}, \beta_1, \ldots, \beta_{v_1} \rrbracket$$

of Φ is one of the sums in Table 4.1 which is not universal (see also Table 3.1). For each case, from the assumption that Φ is universal, we know that at least one of α_{u_1+1} or β_{v_1+1} is less than or equal to $t(\Phi')$ given in Table 4.1. Therefore, we have exactly 564 candidates of proper quaternary universal mixed sums as in Table 4.1.

Now, we show that there are exactly 547 proper quaternary universal mixed sums among those candidates. The truants of the remaining 17 quaternary sums are given in Table 5.1. If u = 0 (v = 0), then the universality of Φ can directly be proved by the "60-theorem" in [5] ("15-theorem" of [1], respectively). Therefore, it suffices to consider the cases when $uv \neq 0$. Since the proofs are quite similar to each other, we only provide, as representative cases, the proofs of Cases

$$(4-1), (4-4), (4-11), (4-15), (4-18), (4-31), and (4-35).$$

In fact, the proof of Case (4-15) is one of the most complicate cases.

To explain how to read Table 4.1, let us consider the case (4-11). Note that $t(\llbracket 1, 2, 3 \rrbracket) = 10$. Hence all possible candidates of quaternary universal sums containing $\llbracket 1, 2, 3 \rrbracket$ are

$$\llbracket 1, 2, \alpha_3, \mathbf{3} \rrbracket$$
 for $2 \leq \alpha_3 \leq 10$ or $\llbracket 1, 2, \mathbf{3}, \boldsymbol{\beta_2} \rrbracket$ for $3 \leq \beta_2 \leq 10$.

Among these quaternary sums, the cases when $2 \le \alpha_3 \le 5$ are already considered. For example, the case when $\alpha_3 = 2$ is considered in Case (4-5). All the other quaternary sums except $\beta_2 = 3$ is, in fact, universal. For the exceptional case, note that t([1, 2, 3, 3]) = 13, which is considered in Section 5 in more detail.

Case (4-1) $(\alpha_1, \alpha_2, \alpha_3) = (1, 1, 1)$. It is enough to show that for any β_1 such that $3 \leq \beta_1 \leq 7$, the equation

$$3x^2 + 3y^2 + 3z^2 + \beta_1 t^2 = 3N + \beta_1$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $t \not\equiv 0 \pmod{3}$ for any nonnegative integer N.

Since the proofs are quite similar to each other, we only provide the proof of the case $\beta_1 = 3$. By Lemma 4.1, we may assume that $3N + 3 \neq 0 \pmod{4}$. If $1 \leq N \leq 15$, then one may directly check that the equation

$$x^2 + y^2 + z^2 + t^2 = N + 1$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $t \neq 0 \pmod{3}$. Therefore, we assume that $N \geq 16$. Note that an integer is represented by $\langle 1, 1, 1 \rangle$ if and only if it is not of the form $4^l(8k+7)$ for any nonnegative integers l, k. Hence one may easily show that there is an integer $d \in \{1, 4\}$ such that $N + 1 - d^2 > 0$, and furthermore $N + 1 - d^2$ is represented by $\langle 1, 1, 1 \rangle$. Therefore, the equation

$$x^2 + y^2 + z^2 + t^2 = N + 1$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $t \not\equiv 0 \pmod{3}$ for any nonnegative integer N. This completes the proof.

Case (4-4) $(\alpha_1, \alpha_2, \beta_1) = (1, 1, 3)$. It is enough to show that for any $\alpha_3(\beta_2)$ such that $4 \leq \alpha_3 \leq 6$ $(3 \leq \beta_2 \leq 6)$, the equation

$$x^{2} + y^{2} + \alpha_{3}z^{2} + t^{2} = N + 1 (3x^{2} + 3y^{2} + 3z^{2} + \beta_{2}t^{2} = 3N + 3 + \beta_{2})$$

Case	Φ'	$t(\Phi')$	Considered	Already		Universal (Cases	
(4-1)	$[\![1,1,1]\!]$	7		$\beta_1 = 1, 2$	$1 \leq \alpha_4 \leq 7,$	$3\leqslant\beta_1\leqslant7$		
(4-2)	$[\![1,1,2]\!]$	14		$\beta_1 = 1, 2$	$2 \leqslant \alpha_4 \leqslant 14,$	$3\leqslant\beta_1\leqslant14$		
(4-3)	$[\![1, 1, 3]\!]$	6		$\beta_1 = 1, 2$	$3 \leq \alpha_4 \leq 6,$	$3\leqslant\beta_1\leqslant6$		
(4-4)	$[\![1, 1, 3]\!]$	6	$1 \leq \alpha_3 \leq 3$		$4 \leqslant \alpha_3 \leqslant 6,$	$3 \leqslant \beta_2 \leqslant 6$		
(4-5)	$[\![1,2,2]\!]$	7			$2 \leqslant \alpha_4 \leqslant 7,$	$1\leqslant\beta_1\leqslant7$		
(4-6)	$[\![1,2,3]\!]$	10		$\beta_1 = 1$	$3 \leq \alpha_4 \leq 10,$	$2\leqslant\beta_1\leqslant10$		
(4-7)	$[\![1,2,4]\!]$	14			$4 \leqslant \alpha_4 \leqslant 14,$	$1 \leqslant \beta_1 \leqslant 14$		
(4-8)	$[\![1,2,5]\!]$	10			$5 \leq \alpha_4 \leq 10,$	$1\leqslant\beta_1\leqslant10$	$\alpha_4 \neq 5,$	$\beta_1 \neq 5$
(4-9)	$[\![1, 2, 1]\!]$	15	$2 \leqslant \alpha_3 \leqslant 6,$	$\beta_2 = 1$	$6 \leq \alpha_3 \leq 15,$	$2 \leqslant \beta_2 \leqslant 15$		$\beta_2 \neq 14$
(4-10)	$[\![1, 2, 2]\!]$	7	$2 \leqslant \alpha_3 \leqslant 5$		$6 \leq \alpha_3 \leq 7,$	$2 \leqslant \beta_2 \leqslant 7$		$\beta_2 \neq 7$
(4-11)	$[\![1, 2, 3]\!]$	10	$2 \leqslant \alpha_3 \leqslant 5$		$6 \leq \alpha_3 \leq 10,$	$3 \leqslant \beta_2 \leqslant 10$		$\beta_2 \neq 3$
(4-12)	$[\![1, 2, 4]\!]$	14	$2 \leqslant \alpha_3 \leqslant 5$		$6 \leq \alpha_3 \leq 14,$	$4 \leqslant \beta_2 \leqslant 14$		$\beta_2 \neq 14$
(4-13)	$[\![1, 2, 5]\!]$	10	$2 \leqslant \alpha_3 \leqslant 5$		$6 \leq \alpha_3 \leq 10,$	$5 \leqslant \beta_2 \leqslant 10$	$\alpha_3 \neq 10,$	$\beta_2 \neq 5, 10$
(4-14)	$[\![1, 1, 2]\!]$	13	$1 \leq \alpha_2 \leq 3$		$4 \leqslant \alpha_2 \leqslant 13,$	$2 \leqslant \beta_3 \leqslant 13$		
(4-15)	$[\![1, 1, 3]\!]$	18	$1 \leq \alpha_2 \leq 3$		$4 \leqslant \alpha_2 \leqslant 18,$	$3\leqslant\beta_3\leqslant18$		
(4-16)	$[\![1,3,2]\!]$	8			$3 \leq \alpha_3 \leq 8,$	$2 \leqslant \beta_2 \leqslant 8$		
(4-17)	$[\![1, 4, 2]\!]$	12			$4 \leqslant \alpha_3 \leqslant 12,$	$2 \leqslant \beta_2 \leqslant 12$		
(4-18)	$[\![1, 5, 2]\!]$	12			$5 \leq \alpha_3 \leq 12,$	$2 \leqslant \beta_2 \leqslant 12$		
(4-19)	$\llbracket 1, 2, 2 rbracket$	7	$1 \leq \alpha_2 \leq 5$		$6 \leq \alpha_2 \leq 7,$	$2 \leqslant \beta_3 \leqslant 7$		
(4-20)	$\llbracket 1, 2, 3 \rrbracket$	8	$1 \leq \alpha_2 \leq 5$		$6 \leq \alpha_2 \leq 8,$	$3 \leqslant \beta_3 \leqslant 8$		
(4-21)	$\llbracket 1, 2, 4 \rrbracket$	12	$1 \leq \alpha_2 \leq 5$		$6 \leq \alpha_2 \leq 12,$	$4 \leqslant \beta_3 \leqslant 12$		
(4-22)	$[\![1, 2, 5]\!]$	12	$1 \leq \alpha_2 \leq 5$		$6 \leqslant \alpha_2 \leqslant 12,$	$5 \leqslant \beta_3 \leqslant 12$		
(4-23)	$[\![2, 2, 1]\!]$	6	$\alpha_3 = 3,$	$\beta_2 = 1$	$2 \leqslant \alpha_3 \leqslant 6,$	$2 \leqslant \beta_2 \leqslant 6$		
(4-24)	$[\![2, 4, 1]\!]$	15		$\beta_2 = 1$	$4 \leqslant \alpha_3 \leqslant 15,$	$2 \leqslant \beta_2 \leqslant 15$		$\beta_2 \neq 14$
(4-25)	$[\![2, 1, 2]\!]$	6	$2 \leqslant \alpha_2 \leqslant 4$		$5 \leq \alpha_2 \leq 6,$	$2 \leqslant \beta_3 \leqslant 6$		
(4-26)	$[\![2, 1, 3]\!]$	14	$2 \leqslant \alpha_2 \leqslant 4$		$5 \leq \alpha_2 \leq 14,$	$3 \leq \beta_3 \leq 14$		$\beta_3 \neq 14$
(4-27)	$[\![2, 1, 4]\!]$	15	$2 \leqslant \alpha_2 \leqslant 4$		$5 \leq \alpha_2 \leq 15,$	$4 \leqslant \beta_3 \leqslant 15$		
(4-28)	$[\![3, 1, 1]\!]$	7			$3 \leq \alpha_2 \leq 7,$	$1 \leqslant \beta_3 \leqslant 7$		$\beta_3 \neq 7$
(4-29)	$\llbracket 1, 1, 1 rbracket$	4	$1 \leq \alpha_1 \leq 3$		$\alpha_1 = 4,$	$1 \leqslant \beta_4 \leqslant 4$		
(4-30)	$\llbracket 1, 1, 2 rbracket$	14	$1 \leq \alpha_1 \leq 3$		$4 \leqslant \alpha_1 \leqslant 14,$	$2 \leqslant \beta_4 \leqslant 14$		$\beta_4 \neq 14$
(4-31)	$\llbracket 1, 1, 3 rbracket$	7	$1 \leq \alpha_1 \leq 3$		$4 \leqslant \alpha_1 \leqslant 7,$	$3 \leqslant \beta_4 \leqslant 7$	$\alpha_1 \neq 7,$	$\beta_4 \neq 4,7$
(4-32)	[3, 1, 2]	9			$3 \leqslant \alpha_2 \leqslant 9,$	$2 \leqslant \beta_3 \leqslant 9$		
(4-33)	$[\![4, 1, 2]\!]$	13			$4 \leqslant \alpha_2 \leqslant 13,$	$2 \leqslant \beta_3 \leqslant 13$		
(4-34)	$\llbracket 1, 2, 2 rbracket$	6	$1 \leqslant \alpha_1 \leqslant 4$		$5 \leqslant \alpha_1 \leqslant 6,$	$2 \leqslant \beta_4 \leqslant 6$		_
(4-35)	$\llbracket 1, 2, 3 rbracket$	9	$1 \leqslant \alpha_1 \leqslant 4$		$5 \leqslant \alpha_1 \leqslant 9,$	$3 \leqslant \beta_4 \leqslant 9$		$\beta_4 \neq 3$
(4-36)	$\llbracket 1, 2, 4 \rrbracket$	13	$1 \leq \alpha_1 \leq 4$		$5 \leq \alpha_1 \leq 13,$	$4 \leqslant \beta_4 \leqslant 13$		

Table 4.1. Quaternary universal mixed sums

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $t \neq 0 \pmod{3}$ $(zt \neq 0 \pmod{3})$, respectively) for any nonnegative integer N.

Since the proofs are quite similar to each other, we only provide the proof of the case $\alpha_3 = 4$. By Lemma 4.1, we may assume that $N + 1 \neq 0 \pmod{4}$. If $N \leq 3$, then one may directly check that

$$x^2 + y^2 + 4z^2 + t^2 = N + 1$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $t \neq 0 \pmod{3}$. Therefore, we assume that $N \geq 4$. Note that a positive integer m is represented by $\langle 1, 1, 1 \rangle$ if and only if it is not of the form $4^l(8k+7)$ for any nonnegative integers l, k. Furthermore, if m is represented by $\langle 1, 1, 1 \rangle$, then there is an integer solution $(x, y, t) \in \mathbb{Z}^3$ such that $x^2 + y^2 + t^2 = m$ and gcd(x, y, t, p) = 1 for any odd prime p. This can be proved by using a slightly modified version of 102:5 of [12]. Hence, there is an

integer $c \in \{0, 1\}$ such that the diophantine equation

$$x^{2} + y^{2} + t^{2} = N + 1 - 4c^{2}$$
 with $t \neq 0 \pmod{3}$

has an integer solution $(x, y, t) \in \mathbb{Z}^3$. This completes the proof of Case (4-4).

Case (4-11) $(\alpha_1, \alpha_2, \beta_1) = (1, 2, 3)$. It suffices to show that for any $\alpha_3(\beta_2)$ such that $6 \leq \alpha_3 \leq 10$ $(4 \leq \beta_2 \leq 10)$, the equation

$$x^{2} + 2y^{2} + \alpha_{3}z^{2} + t^{2} = N + 1 \left(3x^{2} + 6y^{2} + 3z^{2} + \beta_{2}t^{2} = 3N + 3 + \beta_{2}\right)$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $t \neq 0 \pmod{3}(zt \neq 0 \pmod{3})$, respectively) for any nonnegative integer N.

Assume $\alpha_3 = 7$. By Lemma 4.1, we may assume that $N + 1 \not\equiv 0 \pmod{4}$. If $1 \leq N \leq 111$ then one may directly check that such an integer solution always exists. Therefore, we may assume that $N \geq 112$. Note that an integer is represented by $\langle 1, 1, 2 \rangle$ if and only if it is not of the form $2^{2l+1}(8k+7)$ for any nonnegative integers l, k. Hence, one may easily show that there is an integer $c \in \{0, 1, 2, 3, 4\}$ such that $N + 1 - 7c^2 > 0$, $N + 1 - 7c^2 \equiv 0, 1 \pmod{3}$, and furthermore, $N + 1 - 7c^2$ is represented by $\langle 1, 1, 2 \rangle$. Therefore, the equation

$$x^2 + 2y^2 + t^2 = N + 1 - 7c^2$$

has an integer solution $(x, y, t) = (a, b, d) \in \mathbb{Z}^3$ such that $a \neq 0 \pmod{3}$ or $d \neq 0 \pmod{3}$ or $a \equiv b \equiv d \equiv 0 \pmod{3}$. Assume that $a \equiv b \equiv d \equiv 0 \pmod{3}$. We may assume that $a^2 + 2b^2 \neq 0$. By Theorem 9 of [7] (see also [4], and for more generalization, see [10]), there are integers e, f such that $e^2 + 2f^2 = a^2 + 2b^2$ and $ef \neq 0 \pmod{3}$. Therefore, the equation

$$x^2 + 2y^2 + 7z^2 + t^2 = N + 1$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $t \not\equiv 0 \pmod{3}$ for any nonnegative integer N.

Assume $\beta_2 = 9$. It suffices to show that the equation

$$x^2 + 2y^2 + z^2 + 3t^2 = N + 4$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $zt \neq 0 \pmod{3}$. By Lemma 4.1, we may assume that $N + 4 \neq 0 \pmod{4}$. If $N \leq 165$, then one may directly check that

$$x^2 + 2y^2 + z^2 + 3t^2 = N + 4$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $zt \neq 0 \pmod{3}$. Therefore, we assume $N \geq 166$. If $N + 4 \equiv 0, 1 \pmod{3}$, then the proof is quite similar to the proof of the case $\alpha_3 = 7$. Assume $N + 4 \equiv 2 \pmod{3}$. Since $h(\langle 1, 2, 3 \rangle) = 1$, one may easily show that a positive integer which is not of the form $2^{2l+1}(8k+5)$ for any nonnegative integers l, k is represented by $\langle 1, 2, 3 \rangle$. Hence, there is an integer $c \in \{1, 2, 4, 5, 7, 13\}$ such that $N + 4 - c^2 > 0$, $N + 4 - c^2 \equiv 4 \pmod{6}$, and furthermore, $N + 4 - c^2$ is represented by $\langle 1, 2, 3 \rangle$. Therefore, there are integers a, b, d such that $a^2 + 2b^2 + 3d^2 = N + 4 - c^2$ with $a \equiv d \pmod{2}$. If $d \neq 0 \pmod{3}$, then we are done. Assume $d \equiv 0 \pmod{3}$. Note that $\left(\frac{a+3d}{2}\right)^2 + 3\left(\frac{a-d}{2}\right)^2 = a^2 + 3d^2$ and $\left(\frac{a+3d}{2}\right) \cdot \left(\frac{a-d}{2}\right) \neq 0 \pmod{3}$. Therefore, the equation

$$x^2 + 2y^2 + z^2 + 3t^2 = N + 4$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $zt \neq 0 \pmod{3}$.

The proofs of all the other cases are quite similar to those of the case $\alpha_3 = 7$ or Case (4-1).

Assume that the proper sum Φ' of Φ is one of the sums in Table 4.2. Then the proofs are quite similar to those of Cases (4-1) or (4-11).

Case	Φ'	Case	Φ'	Case	Φ'
(4-2)	$[\![1,1,2]\!]$	(4-10)	$\llbracket 1, 2, 2 \rrbracket$	(4-27)	$\llbracket 2, 1, 4 \rrbracket$
(4-3)	$[\![1,1,3]\!]$	(4-14)	$[\![1,1,2]\!]$	(4-28)	$\llbracket 3, 1, 1 rbracket$
(4-5)	$[\![1,2,2]\!]$	(4-16)	$[\![1,3,2]\!]$	(4-29)	$\llbracket 1, 1, 1 rbracket$
(4-6)	$[\![1,2,3]\!]$	(4-17)	$[\![1,4,2]\!]$	(4-30)	$[\![1,1,2]\!]$
(4-7)	$[\![1,2,4]\!]$	(4-19)	$[\![1,\boldsymbol{2},\boldsymbol{2}]\!]$	(4-34)	$[\![1,2,2]\!]$
(4-8)	$[\![1,2,5]\!]$	(4-23)	$\llbracket 2, 2, 1 \rrbracket$	(4-36)	$[\![1,2,4]\!]$
(4-9)	$[\![1,2,1]\!]$	(4-25)	$\llbracket 2, 1, 2 \rrbracket$		

Table 4.2. All cases whose proofs are similar to (4-1) or (4-11)

Note that for each case, the proof uses the set of integers that are represented by a particular ternary quadratic form having class number one, which are easily computable by Hasse principle.

One of most important ingredients in the proofs of Cases (4-1), (4-4) and (4-11) is the existence of a ternary quadratic form having class number 1. We crucially used the fact that the set of integers represented by such a ternary quadratic form is completely known. In the most of remaining cases, the quaternary quadratic form induced by a quaternary mixed sum of 4- and 8-gonal numbers under consideration does not have any ternary quadratic subform having class number one. Hence we use Theorem 2.1 to compute the set of integers in some arithmetic progression that are represented by a ternary quadratic subform.

Case (4-15) $(\alpha_1, \beta_1, \beta_2) = (1, 1, 3)$. It suffices to show that for any $\alpha_2(\beta_3)$ such that $4 \leq \alpha_2 \leq 18$ $(3 \leq \beta_3 \leq 18)$, the equation

$$3x^{2} + 3\alpha_{2}y^{2} + z^{2} + 3t^{2} = 3N + 4 (3x^{2} + y^{2} + 3z^{2} + \beta_{3}t^{2} = 3N + 4 + \beta_{3})$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $zt \neq 0 \pmod{3}(yzt \neq 0 \pmod{3})$, respectively) for any nonnegative integer N.

First, assume $\alpha_2 = 4$. If $N \leq 34$, then one may directly check that such an integer solution always exists. Therefore, we may assume $N \geq 35$. Define

$$f(x, z, t) = 3x^{2} + (3z + t)^{2} + 3t^{2}$$
 and $g(x, z, t) = 27x^{2} + z^{2} + 27t^{2}$

Then one may directly compute that

for any remainder r modulo 11, where r is 0 or a quadratic non-residue modulo 11.

Now, choose an integer $d \in \{0, 1, 2, 3\}$ such that $3N + 4 - 12b^2 > 0$ and $3N + 4 - 12b^2$ is divisible by 11 or a quadratic non-residue modulo 11. Since $h(\langle 1, 3, 3 \rangle) = 1$, one may easily check that

$$3x^2 + z^2 + 3t^2 = 3N + 4 - 12b^2$$

has an integer solution $(x, z, t) \in \mathbb{Z}^3$. Note that z is not divisible by 3. If either x or t is not divisible by 3, then we are done. If both x and t are divisible by 3, that is, $3N + 4 - 12b^2$ is represented by g, then by (4.3), it is represented by f. Therefore, the equation

$$3x^2 + 12y^2 + z^2 + 3t^2 = 3N + 4$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $zt \neq 0 \pmod{3}$.

In fact, if $\alpha_2 \neq 11$ or $\beta_3 \neq 11$, then the proofs of the cases are quite similar to the above.

Assume $\alpha_2 = 11$. If $N \leq 9$, then one may directly check that the equation

$$3x^2 + 33y^2 + z^2 + 3t^2 = 3N + 4$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $zt \neq 0 \pmod{3}$. Therefore, we may assume $N \geq 10$. Note that the genus of $g(x, z, t) = 27x^2 + z^2 + 27t^2$ consists of

$$M_g = \langle 1, 27, 27, \rangle, \quad M_2 = \begin{pmatrix} 4 & 1 & 0 \\ 1 & 7 & 0 \\ 0 & 0 & 27 \end{pmatrix}, \text{ and } M_3 = \begin{pmatrix} 7 & -3 & 2 \\ -3 & 9 & 3 \\ 2 & 3 & 16 \end{pmatrix}.$$

By Minkowski-Siegel formula (for details, see [14]), we have

$$(4.4) \ r(3N+4,\langle 1,3,3\rangle) - r(3N+4,\langle 1,27,27\rangle) = 4 \cdot r(3N+4,M_2) + 4 \cdot r(3N+4,M_3).$$

One may easily check that

$$M_g <_{4,0} M_2$$
 and $M_g <_{4,3} M_2$.

This implies that any integer congruent to 4 or 7 modulo 12 is represented by M_2 or M_3 . First, assume $3N + 4 \equiv 0, 1, 3 \pmod{4}$. Then there is an integer $b \in \{0, 1\}$ such that $3N + 4 - 33b^2 > 0$ and $3N + 4 - 33b^2$ is congruent to 4 or 7 modulo 12. Therefore by (4.4), we have

$$r(3N+4-33b^2,\langle 1,3,3\rangle) - r(3N+4-33b^2,\langle 1,27,27\rangle) > 0.$$

Consequently, the equation

$$3x^2 + 33y^2 + z^2 + 3t^2 = 3N + 4$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $zt \neq 0 \pmod{3}$. Now, assume $3N + 4 \equiv 2 \pmod{4}$. Define $h(x, y, z) = 2x^2 + 2xy + 17y^2 + 3z^2$. The genus of h consists of

$$M_h = \begin{pmatrix} 2 & 1 \\ 1 & 17 \end{pmatrix} \perp \langle 3 \rangle$$
 and $M_4 = \begin{pmatrix} 5 & 2 & 1 \\ 2 & 5 & 1 \\ 1 & 1 & 5 \end{pmatrix}$.

One may easily check that any positive integer m congruent to 2 modulo 3 is represented by M_h or M_4 by 102:5 of [12], for it is represented by M_h over \mathbb{Z}_p for any prime p. One may easily check that

$$M_4 \prec_{4,0} M_h$$
 and $M_4 \prec_{4,1} M_h$

Now, choose an integer $d \in \{1, 2\}$ such that $\frac{1}{2}(3N+4) - 3d^2 > 0$ and $\frac{1}{2}(3N+4) - 3d^2$ is congruent to 5 or 8 modulo 12. Then $\frac{1}{2}(3N+4) - 3d^2$ is represented by M_h . Therefore there are integers a, b, c such that

$$(2a+b)^{2} + 33b^{2} + 3(c+d)^{2} + 3(c-d)^{2} = 3N+4.$$

Since $d \neq 0 \pmod{3}$, $c + d \neq 0 \pmod{3}$ or $c - d \neq 0 \pmod{3}$. Consequently, the equation

$$3x^2 + 33y^2 + z^2 + 3t^2 = 3N + 4$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $zt \neq 0 \pmod{3}$.

Assume $\beta_3 = 11$. First, we assume that $N \ge 88$. Under the same notations as above, one may directly compute that

$$B_{M_3}(M_g, 5, 1) = \{\pm (1, 0, 0)\}$$
 and $B_{M_3}(M_g, 5, 4) = \{\pm (2, 0, 0)\}.$

In each case, if we define $T = \begin{pmatrix} 0 & 4 & -3 \\ 0 & 3 & 4 \end{pmatrix}$, then one may easily show that it

satisfies all conditions in Theorem 2.1. Note that $\pm(1,0,0)$ are the only integral primitive eigenvectors of T. Therefore, we have

(4.5)
$$S_{5,r} \cap Q(M_g) \setminus \{s^2 \mid s \in \mathbb{Z}\} \subset Q(M_3),$$

for any remainder r modulo 5, where r is a quadratic residue modulo 5. Now, choose an integer $d \in \{1, 2, 5\}$ such that $I(N, d) := 3N + 15 - 11d^2 \ge 2$ and $I(N, d) \equiv 1, 4 \pmod{5}$. We show that I(N, d) is also represented by M_2 or M_3 . If I(N, d) is not a square of an integer, then it is represented by M_2 or M_3 by (4.5). Assume that I(N, d) is a square of an integer. Since 4 is represented by M_2 , any square divisible by 4 is represented by M_2 . Since 9 is represented by M_3 , we may assume that I(N, d) is an odd integer not divisible by 3. For any prime p dividing $I(N, d), p^2$ is represented by M_2 or M_3 by Lemma 2.4 of [6], for both M_2 and M_3 are contained in the spinor genus of M_g . Therefore

$$r(I(N,d),\langle 1,3,3\rangle) - r(I(N,d),\langle 1,27,27\rangle) > 0,$$

by (4.4). Hence, the equation

$$3x^2 + y^2 + 3z^2 + 11t^2 = 3N + 15$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $yzt \neq 0 \pmod{3}$. If $N \leq 87$, then one may directly check that such an integer solution exists.

Recall that we use (2.4) and Theorem 2.1 to show that integers in an arithmetic progressions are represented by the ternary quadratic forms. In Case (4-18), as a representative case, we explain how our method works in detail. Since everything is quite similar to this for all the other cases, we only provide all parameters needed for computations in Table 4.3.

Case (4-18) $(\alpha_1, \alpha_2, \beta_1) = (1, 5, 2)$. It suffices to show that for any $\alpha_3(\beta_2)$ such that $5 \leq \alpha_3 \leq 12$ $(2 \leq \beta_2 \leq 12)$, the equation

$$3x^{2} + 15y^{2} + 3\alpha_{3}z^{2} + 2t^{2} = 3N + 2 (3x^{2} + 15y^{2} + 2z^{2} + \beta_{2}t^{2} = 3N + 2 + \beta_{2})$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $t \neq 0 \pmod{3}(zt \neq 0 \pmod{3})$, respectively) for any nonnegative integer N.

First, assume that $\alpha_3 = 5$. By Lemma 4.1, we may assume that $3N + 2 \neq 0 \pmod{4}$. If $N \leq 124$, then one may directly check that

$$3x^2 + 15y^2 + 15z^2 + 2t^2 = 3N + 2$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $t \neq 0 \pmod{3}$. Therefore we may assume that $N \ge 125$. Note that the genus of $f(x, y, t) = 3x^2 + 15y^2 + 2t^2$ consists of

$$M_f = \langle 2, 3, 15 \rangle$$
 and $M_2 = \begin{pmatrix} 2 & 1 & -1 \\ 1 & 5 & 1 \\ -1 & 1 & 11 \end{pmatrix}$.

One may easily show that any positive integer m congruent to 2 modulo 3 that is not of the form $2^{2l+1}(8k+3)$ for any nonnegative integers l, k is represented by M_f or M_2 by 102:5 of [12]. Now, one may easily check that

$$M_2 \prec_{2,0} M_f.$$

Therefore, any positive integer m congruent to 2 modulo 6 that is not of the form $2^{2l+1}(8k+3)$ is represented by M_f . Hence we may choose an integer $c \in \{1, 2, 3, 4, 5\}$ such that $3N + 2 - 15c^2 > 0$, $3N + 2 - 15c^2 \equiv 2 \pmod{6}$, and furthermore, $3N + 2 - 15c^2$ is represented by M_f . Therefore, the equation

$$3x^2 + 15y^2 + 15z^2 + 2t^2 = 3N + 2$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $t \neq 0 \pmod{3}$.

If either α_3 or β_2 is odd, then each proof is quite similar to that of the case $\alpha_3 = 5$.

Assume $\alpha_3 = 8$. Note that the genus of $f(x, z, t) = 3x^2 + 24z^2 + 2t^2$ consists of

$$M_f = \langle 2, 3, 24 \rangle$$
 and $M_2 = \begin{pmatrix} 5 & 1 \\ 1 & 5 \end{pmatrix} \perp \langle 6 \rangle$

One may easily show that a positive integer m which is congruent to 2 modulo 3 and not of the form $4^{l}(8k + 1)$ and $4^{l}(8k + 7)$ for any nonnegative integers l, k is represented by M_{f} or M_{2} by 102:5 in [12] for it is represented by M_{f} over \mathbb{Z}_{p} for any prime p. Note that

$$B_f(M_2, 3, 2) = \{\pm(1, 2, 0)\}$$

If we define $T = \begin{pmatrix} 2 & -1 & -2 \\ -1 & 2 & -2 \\ 2 & 2 & 1 \end{pmatrix}$, then one may easily show that it satisfies all

conditions in Theorem 2.1. Note that $\pm(1, -1, 0)$ are the only integral primitive eigenvectors of T. Therefore we have

$$S_{3,2} \cap Q(M_2) \setminus \{8s^2 \mid s \in \mathbb{Z}\} \subset Q(M_f).$$

Since 8 is represented by M_f , $8s^2$ is represented by M_f for any $s \in \mathbb{Z}$. Therefore, a positive integer m which is congruent to 2 modulo 3 and not of the form $4^l(8k+1)$ and $4^l(8k+7)$ is represented by M_f . The rest of the proof is quite similar to that of $\alpha_3 = 5$.

Assume $\alpha_3 = 12$. Note that the genus of $f(x, z, t) = 3x^2 + 36z^2 + 2t^2$ consists of

$$M_f = \langle 2, 3, 36 \rangle$$
 and $M_2 = \langle 2, 9, 12 \rangle$.

Note that

$$M_2 \prec_{2,0} M_f.$$

One may easily show that every positive integer congruent to 2 modulo 6 that is not of the form $2^{2l+1}(8k+7)$ for any nonnegative integers l, k is represented by M_f . The rest of the proof is quite similar when $\alpha_3 = 5$.

Assume $\beta_2 = 10$. By Lemma 4.1, we may assume that $3N + 12 \neq 0 \pmod{4}$. If $N \leq 1276$, then one may directly check that the equation

$$3x^2 + 15y^2 + 2z^2 + 10t^2 = 3N + 12$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $zt \neq 0 \pmod{3}$. Therefore, assume $N \geq 1277$. Note that the genus of $f(x, z, t) = \frac{1}{3}(3x^2 + 2(3z + t)^2 + 10t^2) = x^2 + 6z^2 + 4t^2 + 4zt$ consists of

$$M_f = \langle 1 \rangle \perp \begin{pmatrix} 4 & 2 \\ 2 & 6 \end{pmatrix}$$
 and $M_2 = \langle 2, 2, 5 \rangle$.

One may easily show that a positive integer m which is not of the form $4^{l}(8k + 3)$ for any nonnegative integers l, k is represented by M_{f} or M_{2} by 102:5 in [12] for it is represented by M_{f} over \mathbb{Z}_{p} for any prime p. Note that

$$M_2 \prec_{3,0} M_f$$
 and $M_2 \prec_{3,1} M_f$.

Therefore, a positive integer m that is congruent to 0 or 1 modulo 3 and not of the form $4^{l}(8k+3)$ is represented by M_{f} . Hence there is an integer $b \in \{0, 1, 3, 4\}$ such that both $N + 4 - 5b^{2}$ and $N + 4 - 5(b+12)^{2}$ are positive and represented by M_{f} . For all possible cases when both $N + 4 - 5b^{2}$ and $N + 4 - 5(b+12)^{2}$ are squares of integers, one may easily check that the equation

$$3x^2 + 15y^2 + 2z^2 + 10t^2 = 3N + 12$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $zt \neq 0 \pmod{3}$. Therefore we may assume that $3x^2 + 2z^2 + 10t^2 = 3N + 12 - 15b^2$ has an integer solution (x, z, t) = $(a, c, d) \in \mathbb{Z}^3$ such that $2c^2 + 10d^2 \neq 0$. If $c \equiv d \equiv 0 \pmod{3}$, then there are integers e, f such that $e^2 + 5f^2 = c^2 + 5d^2$ and $ef \neq 0 \pmod{3}$, by Theorem 9 of [7]. This completes the proof.

Assume $\beta_2 = 12$. By Lemma 4.1, we may assume that $3N + 14 \neq 0 \pmod{4}$. If $N \leq 1800$, then one may directly check that the equation

$$3x^2 + 15y^2 + 2z^2 + 12t^2 = 3N + 14$$

14

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $zt \neq 0 \pmod{3}$. Therefore, assume $N \geq 1801$. Define

$$f(x, z, t) = 12x^{2} + 2(3z + t)^{2} + 12t^{2}$$
 and $g(x, z, t) = 108x^{2} + 2z^{2} + 108t^{2}$.

Note that

(4.6)
$$g \prec_{7,r} f_4,$$

for any remainder r modulo 7, where r is 0 or a quadratic non-residue modulo 7. Since $h(\langle 1, 6, 6 \rangle) = 1$, one may easily show that a positive integer m which is congruent to 1 modulo 3 and not of the form $4^l(8k + 3)$ for any nonnegative integers l, k is represented by $\langle 1, 6, 6 \rangle$. Hence there is an integer $0 \leq b \leq 19$ such that $3N + 14 - 15b^2$ is positive, $3N + 14 - 15b^2$ is divisible by 7 or a quadratic nonresidue modulo 7, and furthermore, $3N + 14 - 15b^2$ is represented by $\langle 2, 12, 12 \rangle$. If $3N + 14 - 15b^2$ is not represented by g, then we are done. Assume that $3N + 14 - 15b^2$ is represented by g. By Equation (4.6), $3N + 14 - 15b^2$ is represented by f. Therefore the equation

$$3(2x)^2 + 15y^2 + 2(3z+t)^2 + 12t^2 = 3N + 14$$

has an integer solution. This completes the proof.

The proofs of all remaining cases are quite similar to those of Cases (4-1) or (4-11). This completes the proof of Case (4-18).

As noted earlier, since every proof of the case in the Table 4.3 is quite similar to that of Case (4-18), we only provide all parameters needed for the computations.

Case (4-31) $(\beta_1, \beta_2, \beta_3) = (1, 1, 3)$. It is enough to show that for any α_1 such that $4 \leq \alpha_1 \leq 6$, the equation

$$3\alpha_1 x^2 + y^2 + z^2 + 3t^2 = 3N + 5$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $yzt \neq 0 \pmod{3}$ for any nonnegative integer N.

Since the proof of the case $\alpha_1 = 4$ is quite similar to that of Case (4-1), we only provide proofs of the cases when $\alpha_1 \in \{5, 6\}$.

Assume $\alpha_1 = 5$. Note that $h(\langle 1, 1, 3 \rangle) = 1$. Every positive integer congruent to 2 modulo 3 is represented by $\langle 1, 1, 3 \rangle$. Therefore, the equation

$$y^2 + z^2 + 3t^2 = 3N + 5$$

has an integer solution $(y, z, t) = (b, c, d) \in \mathbb{Z}^3$. If $d \neq 0 \pmod{3}$, then we are done. Assume that $d \equiv 0 \pmod{3}$. First, we consider the case when $d \neq 0$. Then there are a positive integer s and a positive integer d_1 not divisible by 3 such that $d = 3^s \cdot d_1$. Now, by Theorem 9 of [7], there are integers d_2 and a such that $3^{2s+1} - 15a^2 = 3d_2^2$ and $d_2 \neq 0 \pmod{3}$. Therefore, we have

$$b^{2} + c^{2} + 3(d_{2} \cdot d_{1})^{2} + 15(a \cdot d_{1})^{2} = 3N + 5.$$

Now, assume d = 0. If $c \equiv 0 \pmod{2}$, then $b^2 + c^2 = b^2 + 4(\frac{c}{2})^2 = b^2 + (\frac{c}{2})^2 + 3(\frac{c}{2})^2$. Therefore, we may assume that $b \equiv c \equiv 1 \pmod{2}$. Note that $b^2 + c^2 = 2\left(\frac{b+c}{2}\right)^2 + 2\left(\frac{b-c}{2}\right)^2$ and $b \equiv c \pmod{3}$. If $b \neq c$, then there are integers b_1, c_1 such that $b^2 + c^2 = 2b_1^2 + 18c_1^2$ and $c_1 \neq 0$. By theorem 9 of [7], there are integers d_3, a_2 such that

Case	M_{f}	M_2	(d,a)	$B_f(M_2, d, a)$	Т	1	f represents m
(4-12)	$\left< 3,4,6 \right>$	$\left< 1, 6, 12 \right>$	(3, 1)	$\{\pm(1,0,0)\}$	$ \begin{pmatrix} 3 & 0 \\ 0 & 1 \\ 0 & 2 \end{pmatrix} $	$\begin{pmatrix} 0\\ -4\\ 1 \end{pmatrix}$	$m \ge 2,$ $m \equiv 1 \pmod{3},$ $m \ne 2^{2l+1}(8k+7)$
(4-13)	$\left< 3,5,6 \right>$	$\begin{pmatrix} 2 & 0 & 0 \\ 0 & 6 & 3 \\ 0 & 3 & 9 \end{pmatrix}$	(3, 2)	$\{\pm(1,0,0)\}$	$\begin{pmatrix} 3 & 0 \\ 0 & 3 \\ 0 & -3 \end{pmatrix}$	$\begin{pmatrix} 0 \\ 3 \\ 2 & 1 \end{pmatrix}$	$ \begin{split} m &\equiv 3 \pmod{2}, \\ m &\neq 5^{2l+1}(5k+2), \\ m &\neq 5^{2l+1}(5k+3), \\ m &\neq 2 \cdot 25^l \end{split} $
(4-20) $\beta_3 \neq 3, 4$	$\begin{pmatrix} 3 & 0 & 0 \\ 0 & 5 & 1 \\ 0 & 1 & 11 \end{pmatrix}$	$\langle 2,3,27 angle$	$\begin{array}{c} (13,2) \\ (13,5) \\ (13,6) \\ (13,7) \\ (13,8) \\ (13,11) \end{array}$	$ \{ \pm (1, 0, 0) \} \\ \{ \pm (3, 0, 0) \} \\ \{ \pm (4, 0, 0) \} \\ \{ \pm (6, 0, 0) \} \\ \{ \pm (2, 0, 0) \} \\ \{ \pm (5, 0, 0) \} $	$\begin{pmatrix} 13 & 0 \\ 0 & 5 \\ 0 & 4 \end{pmatrix}$	$\begin{pmatrix} 0\\ -36\\ 5 \end{pmatrix}$	$m \ge 3,$ $m \equiv 2 \pmod{3},$ $m \equiv r_1 \pmod{13}$
(4-21)	$\langle 2,3,4 angle$	$\left<1,2,12\right>$	(3, 0)	Ø			$m \equiv 0 \pmod{3},$ $m \neq 2^{2l+1}(8k+5)$
(4-22) $\alpha_2 \neq 7$ $\beta_3 \neq 7$	$\langle 2,3,5 angle$	$\langle 1, 1, 30 angle$	$(7, r_2)$	Ø			$m \equiv 1 \pmod{3}, m \equiv r_2 \pmod{7}$
(4-22)	$\left<3,5,21\right>$	$ \begin{pmatrix} 5 & 0 & 0 \\ 0 & 6 & 3 \\ 0 & 3 & 12 \end{pmatrix} $	$(8, r_3)$	Ø			$m \equiv 2 \pmod{3},$ $m \not\equiv 0 \pmod{5},$
$\alpha_2 = 7$	$\left<3,5,21\right>$	$\begin{pmatrix} 2 & 1 & 0 \\ 1 & 8 & 3 \\ 0 & 0 & 21 \end{pmatrix}$	$(8, r_3)$	Ø			$m \not\equiv 0 \pmod{7}, m \equiv r_3 \pmod{8}$
(4-22) $\beta_3 = 7$	$\langle 2, 3, 7 \rangle$	$ \begin{pmatrix} 2 & 1 & 1 \\ 1 & 3 & 0 \\ 1 & 0 & 9 \end{pmatrix} $	(3,0)	Ø			$m \equiv 6 \pmod{9},$ $m \neq 2^{2l+1}(8k+3)$
(4-24)	$\langle 1, 6, 12 \rangle$	$\langle 3, 4, 6 \rangle$	(3,0) (3,1)	$\{\pm(0,1,0)\}$	$ \begin{pmatrix} 1 & 0 \\ 0 & 3 \\ 2 & 0 \end{pmatrix} $	$\begin{pmatrix} -4\\ 0\\ 1 \end{pmatrix}$	$m \equiv 1 \pmod{3},$ $m \neq 2^{2l+1}(8k+7)$
$(4-26)$ $\alpha_2 \neq 13$ $\beta_3 \neq 13$	$\begin{pmatrix} 4 & 1 & 0 \\ 1 & 7 & 0 \\ 0 & 0 & 6 \end{pmatrix}$	$\left< 1, 6, 27 \right>$	$(13, r_4)$	Ø			$m \equiv 1 \pmod{3},$ $m \equiv r_4 \pmod{13},$ $m \neq 2^{2l+1}(8k+7)$
(4-32)	$\langle 1,2,9 angle$	$\begin{pmatrix} 2 & 1 & 0 \\ 1 & 3 & 1 \\ 0 & 1 & 4 \end{pmatrix}$	(3, 0)	Ø			$m \equiv 0 \pmod{3},$ $m \neq 2^{2l+1}(8k+7)$
(4-33)	$\left< 1, 2, 12 \right>$	$\langle 2,3,4 angle$	(3, 0)	Ø			$m \equiv 0 \pmod{3},$ $m \neq 2^{2l+1}(8k+5)$
				$r_1 \in \{\overline{2, 5}$	$, 6, 7, 8, \overline{11}$	$\}, r_2$	$\in \{0, 3, 5, 6\},$

Table 4.3. Some data for quaternary universal mixed sums

 $r_3 \in \{0, 1, 4, 5\},$ $r_4 \in \{0, 2, 5, 6, 7, 8, 11\}$

 $18c_1^2 = 3d_3^2 + 15a_2^2$ and $a_2d_3 \not\equiv 0 \pmod{3}$. Then $b_1^2 + b_1^2 + 3d_3^2 + 15a_2^2 = 3N + 5$. Finally, assume that $3N + 5 = 2b^2$. Note that the genus of $f(y, z, t) = (3y+t)^2 + (3z+t)^2 + 3t^2$ consists of

 $M_f = \begin{pmatrix} 5 & -2 & 2 \\ -2 & 8 & 1 \\ 2 & 1 & 8 \end{pmatrix} \text{ and } M_2 = \begin{pmatrix} 2 & 1 \\ 1 & 5 \end{pmatrix} \perp \langle 27 \rangle.$

Note that M_2 is in the spinor genus of M_f . Clearly, it suffices to show that 3N+5 = $2b^2$ is represented by M_f . Assume that b is even. Since 8 is represented by M_f , $2b^2$ is represented by M_f . Since 18 is also represented by M_f , we may assume that b is relatively prime to 6 by a similar reasoning to the above. Since b > 1, there is a prime p > 3 dividing b. Since 2 is represented by M_2 , $2p^2$ and therefore $2b^2$, is represented by M_f by Lemma 2.4 of [6]. This completes the proof.

Assume $\alpha_1 = 6$. Note that the genus of $f_1(x, y, z) = 18x^2 + y^2 + z^2$ consists of

$$M_{f_1} = \langle 1, 1, 18 \rangle$$
 and $M_3 = \langle 2 \rangle \perp \begin{pmatrix} 2 & 1 \\ 1 & 5 \end{pmatrix}$.

Note that $M_3 <_{2,0} M_{f_1}$. Then one may easily show that for a positive integer which is congruent to 2 modulo 6 and not of the form $2^{2l+1}(8k+7)$ for any nonnegative integers l, k is represented by M_f . The rest of the proof is quite similar to that of Case (4-18).

Case (4-35) $(\beta_1, \beta_2, \beta_3) = (1, 2, 3)$. It is enough to show that for any α_1 such that $5 \leq \alpha_1 \leq 9$, the equation

$$3\alpha_1 x^2 + y^2 + 2z^2 + 3t^2 = 3N + 6$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $yzt \neq 0 \pmod{3}$ for any nonnegative integer N.

First, assume $\alpha_1 = 5$. By Lemma 4.1, we may assume that $3N + 6 \neq 0 \pmod{4}$. If $N \leq 2643$, the one may directly check that such an integer solution exists. Therefore, we may assume that $n \geq 2644$. Note that the genus of $f(y, z, t) = \frac{1}{3}((t+3y)^2 + 2(t+3z)^2 + 3t^2)$ consists of

$$M_f = \begin{pmatrix} 2 & 1 & 0 \\ 1 & 3 & 1 \\ 0 & 1 & 4 \end{pmatrix}$$
 and $M_2 = \langle 1, 2, 9 \rangle$.

One may easily check that any positive integer m which is not of the form $2^{2l+1}(8k+7)$ for any nonnegative integers l, k is represented by M_f or M_2 . Furthermore, we may check by a direct computation that

$$M_2 \prec_{2,0} M_f.$$

Then there is an integer $a \in \{0, 1, 2, 3, 5, 7\}$ such that both $N + 2 - 5a^2$ and $N + 2 - 5(a + 16)^2$ are positive and represented by M_f . For all possible cases when both $N + 2 - 5a^2$ and $N + 2 - 5(a + 16)^2$ are of the form $10s^2$ for some $s \in \mathbb{Z}$, one may directly check that the equation

$$15x^2 + y^2 + 2z^2 + 3t^2 = 3N + 6$$

has an integer solution $(x, y, z, t) \in \mathbb{Z}^4$ such that $yzt \neq 0 \pmod{3}$. We may assume that at least one of $N + 2 - 5a^2$ and $N + 2 - 5(a + 16)^2$ is not of the form $10s^2$ for any $s \in \mathbb{Z}$. Therefore, the equation

$$15x^2 + y^2 + 2z^2 + 3t^2 = 3N + 6$$

has an integer solution $(x, y, z, t) = (a, b, c, d) \in \mathbb{Z}^4$ such that $b \equiv c \equiv d \pmod{3}$ and $3N + 6 - 15a^2$ is not of the form $30s^2$. If $b \equiv c \equiv d \neq 0 \pmod{3}$, then we

are done. Assume
$$b \equiv c \equiv d \equiv 0 \pmod{3}$$
. If $\tau = \frac{1}{3} \begin{pmatrix} -1 & -1 & 3 \\ -1 & 2 & 0 \end{pmatrix}$, then one may

easily check that $\tau(b, c, d)^t = (b_1, c_1, d_1)^t$ is also an integer solution of

(4.7)
$$y^2 + 2z^2 + 3t^2 = 3N + 6 - 15a^2$$

Such that $b_1 \equiv c_1 \equiv d_1 \pmod{3}$. Therefore, there is a positive integer m such that $\tau^m(a, b, c)^t = (a_m, b_m, c_m)^t$ is an integer solution of Equation (4.7) such that each of whose component is not divisible by 3 or for any positive integer $m, \tau^m(a, b, c)^t = (a_m, b_m, c_m)^t$ is an integer solution of Equation (4.7) each of whose component is divisible by 3. Since there are only finitely many integer solution of Equation (4.7) and τ has an infinite order, the latter is impossible unless (a, b, c) is an eigenvector of τ . Note that $\pm(0, -3, 2)$ are the only integer primitive eigenvectors of τ . Since $3N + 6 - 15a^2$ is not of the form $30s^2$, the equation

$$y^2 + 2z^2 + 3t^2 = 3N + 6 - 15a^2$$

has an integer solution $(y, z, t) \in \mathbb{Z}^3$ such that $yzt \neq 0 \pmod{3}$.

Assume that α_1 is odd. Then every proof is quite similar to that of the case $\alpha_1 = 5$.

Assume $\alpha_1 = 6$. Note that the genus of $f_1(x, y, z) = 18x^2 + y^2 + 2z^2$ consists of

$$M_f = \langle 1, 2, 18 \rangle$$
 and $M_2 = \begin{pmatrix} 3 & -1 & 1 \\ -1 & 3 & -1 \\ 1 & -1 & 5 \end{pmatrix}$.

Note that

$$M_3 \prec_{3,0} M_{f_1}$$

Therefore, a positive integer which is divisible by 3 and not of the form $4^{l}(8k + 7)$ for any nonnegative integers l, k is represented by M_{f} . The rest of the proof is quite similar to that of Case (4-18).

Assume $\alpha_1 = 8$. By Lemma 4.1, we may assume that $3N + 6 \neq 0 \pmod{4}$. If $1 \leq N \leq 286$, the one may directly check that the equation

$$24x^2 + y^2 + 2z^2 + 3t^2 = 3N + 6$$

has an integer solution such that $yzt \neq 0 \pmod{3}$. Therefore, we assume that $N \geq 277$. There is an integer $a \in \{3, 6\}$ such that $3N+6-24a^2 > 0$ and furthermore, $3N+6-24a^2$ is represented by $\langle 1, 2, 3 \rangle$. Therefore, the equation

$$y^2 + 2z^2 + 3t^2 = 3N + 6 - 24a^2$$

has an integer solution $(y, z, t) = (b, c, d) \in \mathbb{Z}^3$. We may assume that $b^2 + 2c^2 \neq 0$ and further assume that $bc \neq 0 \pmod{3}$ by theorem 9 of [7]. If $d \neq 0 \pmod{3}$, then we are done. Assume $d \equiv 0 \pmod{3}$. Since $d^2 + 8a^2 \neq 0$ and $d^2 + 8a^2 \equiv 0 \pmod{3}$, there are integers e, f such that $d^2 + 8a^2 = e^2 + 8f^2$ and $ef \neq 0 \pmod{3}$ by Theorem 9 of [7]. This completes the proof of Case (4-35).

5. The 61-theorem of generalized 4- and 8-gonal numbers

In this section, we determine all *n*-ary proper universal mixed sums of generalized 4- and 8-gonal numbers for any $n \ge 5$. Furthermore, we give a simple criterion on the universality of an arbitrary mixed sum of generalized 4- and 8-gonal numbers, which is a generalization of the "15-theorem".

Theorem 5.1. There are exactly 707 quinary proper universal mixed sums of generalized 4- and 8-gonal numbers (for the list of them, see Table 5.1).

Proof. Let u, v be nonnegative integers such that u + v = 5. For positive integers $\alpha_1, \ldots, \alpha_u$ and β_1, \ldots, β_v , assume that a quinary mixed sum

$$\Phi = \llbracket \alpha_1, \dots, \alpha_u, \beta_1, \dots, \beta_v \rrbracket$$

is proper universal. Then by Theorems 3.1 and 4.2, there are nonnegative integers u_1, v_1 such that $u_1 + v_1 = 4$, and the proper sum

$$\Phi' = \llbracket \alpha_1, \dots, \alpha_{u_1}, \beta_1, \dots, \beta_{v_1} \rrbracket$$

of Φ is one of the sums in Table 5.1 which is not universal(see also Table 4.1). For each case, from the universality of Φ we know that at least one of α_{u_1+1} or β_{v_1+1} is less than equal to the truant of the mixed sum Φ' given in Table 5.1. Therefore, we have exactly 708 candidates of proper quinary universal mixed sums.

Now, we show that all candidates are, in fact, universal except [[1, 2, 5, 5, 5]]. The truant of the remaining quinary sum is 20. As in Theorem 4.2, we may assume that $uv \neq 0$. Since the proofs are quite similar to each other, we only provide, as representative cases, the proofs of Cases

$$(5-2)$$
, $(5-14)$, and $(5-16)$.

Table 5.1. Quinary and senary universal mixed sums

		Tab	le J.I. Quinary and senar	y universai mixeu sums	
Case	Φ'	$t(\Phi')$	Considered already	Universal case	
(5-1)	$\llbracket 1, 2, 5, 5 \rrbracket$	15	$6 \leq \alpha_5 \leq 10, 6 \leq \beta_1 \leq 10$	$\alpha_5, \beta_1 = 5, \ 11 \leqslant \alpha_5, \beta_1 \leqslant 15$	
(5-2)	$\llbracket 1, 2, 5, 5 \rrbracket$	15	$5 \leq \alpha_4 \leq 10, 6 \leq \beta_2 \leq 10$	$11 \leqslant \alpha_4 \leqslant 15, \beta_2 = 5, 11 \leqslant \beta_2 \leqslant 15$	
(5-3)	$\llbracket 1,2,1,14 \rrbracket$	61	$2 \leq \alpha_3 \leq 15, \ \beta_3 = 15$	$16 \leqslant \alpha_3 \leqslant 61, \beta_3 = 14, 16 \leqslant \beta_3 \leqslant 61$	
(5-4)	$[\![1, 2, 2, 7]\!]$	30	$2 \leqslant \alpha_3 \leqslant 7$	$8 \leqslant \alpha_3 \leqslant 30, 7 \leqslant \beta_3 \leqslant 30$	
(5-5)	$\llbracket 1,2,\boldsymbol{3},\boldsymbol{3} \rrbracket$	13	$2 \leq \alpha_3 \leq 10, \ 4 \leq \beta_3 \leq 10$	$11 \leqslant \alpha_3 \leqslant 13, \beta_3 = 3, 11 \leqslant \beta_3 \leqslant 13$	
(5-6)	$[\![1, 2, 4, 14]\!]$	60	$2 \leq \alpha_3 \leq 14$	$15 \leqslant \alpha_3 \leqslant 60, \ 14 \leqslant \beta_3 \leqslant 60$	
(5-7)	$\llbracket 1, 2, 10, 5 \rrbracket$	20	$6 \leqslant \beta_2 \leqslant 9$	$10 \leqslant \alpha_4 \leqslant 20, \beta_2 = 5, 10 \leqslant \beta_2 \leqslant 20$	
(5-8)	$[\![1, 2, 5, 5]\!]$	15	$2 \leq \alpha_3 \leq 10, 6 \leq \beta_3 \leq 9$	$11 \leq \alpha_3 \leq 15, \ \beta_3 = 5, 10 \leq \beta_3 \leq 15 \ \beta_3 = 5$	$\neq 5$
(5-9)	$[\![1, 2, 5, 10]\!]$	20	$2 \leq \alpha_3 \leq 10$	$11 \leqslant \alpha_3 \leqslant 20, \ 10 \leqslant \beta_3 \leqslant 20$	
(5-10)	$[\![2,4,1,14]\!]$	61	$4 \leqslant \alpha_3 \leqslant 15, \beta_3 = 15$	$16 \leqslant \alpha_3 \leqslant 61, \ \beta_3 = 14, 16 \leqslant \beta_3 \leqslant 61$	
(5-11)	$[\![2, 1, 3, 14]\!]$	60	$2 \leqslant \alpha_2 \leqslant 14$	$15 \leqslant \alpha_2 \leqslant 60, \ 14 \leqslant \beta_4 \leqslant 60$	
(5-12)	$[\![3, 1, 1, 7]\!]$	30	$3 \leq \alpha_2 \leq 7$	$8 \leqslant \alpha_2 \leqslant 30, 7 \leqslant \beta_4 \leqslant 30$	
(5-13)	$[\![1,1,2,14]\!]$	60	$1 \leq \alpha_1 \leq 14$	$15 \leq \alpha_1 \leq 60, \ 14 \leq \beta_5 \leq 60$	
(5-14)	$\llbracket 7, 1, 1, 3 \rrbracket$	14	$\beta_4 = 3, 5, 6$	$7 \leqslant \alpha_2 \leqslant 14, \beta_4 = 4, 7 \leqslant \beta_4 \leqslant 14$	
(5-15)	$[\![1,1,3,4]\!]$	18	$1 \leqslant \alpha_1 \leqslant 7, \beta_5 = 5, 6$	$8 \leqslant \alpha_1 \leqslant 18, \beta_5 = 4, 7 \leqslant \beta_5 \leqslant 18$	
(5-16)	$[\![1,1,3,7]\!]$	14	$1 \leqslant \alpha_1 \leqslant 7$	$8 \leqslant \alpha_1 \leqslant 14, 7 \leqslant \beta_5 \leqslant 14$	
(5-17)	$[\![\boldsymbol{1},\boldsymbol{2},\boldsymbol{3},\boldsymbol{3}]\!]$	12	$1 \leqslant \alpha_1 \leqslant 9, \ 4 \leqslant \beta_5 \leqslant 9$	$10 \leqslant \alpha_1 \leqslant 12, \beta_5 = 3, 10 \leqslant \beta_5 \leqslant 12$	
(6-1)	$[\![1, 2, 5, 5, 5]\!]$	20	$1 \leq \alpha_3 \leq 15, \ 6 \leq \beta_4 \leq 15$	$16 \leq \alpha_3 \leq 20$ $\beta_4 = 5, 16 \leq \beta_4 \leq 20$	

Case (5-2) $(\alpha_1, \alpha_2, \alpha_3, \beta_1) = (1, 2, 5, 5)$. It is enough to show that for any $\alpha_4(\beta_2)$ such that $11 \leq \alpha_4 \leq 15$ $(\beta_2 = 5 \text{ or } 11 \leq \beta_2 \leq 15)$, the equation

$$x^{2} + 2y^{2} + 5z^{2} + \alpha_{4}t^{2} + 5(3s^{2} - 2s) = N$$

(x² + 2y² + 5z² + 5(3t² - 2t) + \beta_{2}(3s^{2} - 2s) = N, respectively)

has an integer solution $(x, y, z, t, s) \in \mathbb{Z}^5$ for any nonnegative integer N.

Assume $\alpha_4 = 11$. If $N \leq 10$, then one may directly check that the equation

$$x^{2} + 2y^{2} + 5z^{2} + 11t^{2} + 5(3s^{2} - 2s) = N$$

has an integer solution $(x, y, z, t, s) \in \mathbb{Z}^5$. Since $h(\langle 1, 2, 5 \rangle) = 1$, one may easily show that every positive integer not of the form $5^{2l+1}(5k+2)$ and $5^{2l+1}(5k+3)$ for any nonnegative integer l, k is represented by $\langle 1, 2, 5 \rangle$. Therefore, we may assume $N \equiv 0 \pmod{5}$. Then N - 11 is represented by $\langle 1, 2, 5 \rangle$. If α_4 or β_2 is not divisible by 5, then the proofs are quite similar to this.

Assume $\alpha_4 = 15$. If $N \leq 20$, then one may directly check that the equation

$$x^{2} + 2y^{2} + 5z^{2} + 15t^{2} + 5(3s^{2} - 2s) = N$$

has an integer solution $(x, y, z, t, s) \in \mathbb{Z}^5$. Assume $N \ge 21$. We may assume that $N \equiv 0 \pmod{5}$. Then there are integers $d, e \in \{0, 1\}$ such that $N - 15d^2 - 5(3e^2 - 2e)$ is represented by $\langle 1, 2, 5 \rangle$.

If $\beta_2 \in \{5, 15\}$, then the proof is quite similar to the above. This completes the proof of Case (5-2).

Case (5-14) $(\alpha_1, \beta_1, \beta_2, \beta_3) = (7, 1, 1, 3)$. It is enough to show that for any $\alpha_2(\beta_4)$ such that $7 \leq \alpha_2 \leq 14$ $(\beta_4 = 4 \text{ or } 7 \leq \beta_4 \leq 14)$, the equation

$$21x^{2} + 3\alpha_{2}y^{2} + z^{2} + t^{2} + 3s^{2} = 3N + 5(21x^{2} + y^{2} + z^{2} + 3t^{2} + \beta_{4}s^{2} = 3N + 5 + \beta_{4})$$

has an integer solution $(x, y, z, t, s) \in \mathbb{Z}^5$ such that $zts \neq 0 \pmod{3}$ $(yzts \neq 0 \pmod{3})$, respectively) for any nonnegative integer N.

We show that

(5.1)
$$x_1^2 + y_1^2 + 3z_1^2 + 21t_1^2 = 3N + 5 \text{ with } x_1y_1z_1 \neq 0 \pmod{3}$$

has an integer solution $(x_1, y_1, z_1, t_1) \in \mathbb{Z}^4$ for any nonnegative integer N except 14. If $N \leq 782$ and $N \neq 14$, then one may directly check that Equation (5.1) has an integer solution. Note that Equation (5.1) does not have an integer solution for N = 14, whereas it has an integer solution for $N = 4 \cdot 14 = 56$. Therefore, we may assume that $N \geq 783$ and $3N + 5 \neq 0 \pmod{4}$ by similar reasoning of Lemma 4.1. Since $h(\langle 1, 1, 21 \rangle) = 1$, one may easily show that a positive integer which is congruent to 2 modulo 3, not divisible by 7, and not of the form $4^l(8k + 3)$ for any nonnegative integers l, k is represented by $\langle 1, 1, 21 \rangle$. Then there is an integer $c \in \{2, 4, 14, 28\}$ such that $3N + 5 - 3c^2 > 0$, and furthermore, $3N + 5 - 3c^2$ is represented by $\langle 1, 1, 21 \rangle$. Therefore, the equation

$$x_1^2 + y_1^2 + 3z_1^2 + 21t_1^2 = 3N + 5$$

has an integer solution $(x_1, y_1, z_1, t_1) \in \mathbb{Z}^4$ such that $x_1y_1z_1 \not\equiv 0 \pmod{3}$ for any nonnegative integer N except 14. The proof of Case (5-14) follows immediately.

In most of the cases, we can exactly compute the set of all positive integers not represented by Φ' , denoted by $E(\Phi')$. If a proper sum Φ' of Φ is a one of the sums in Table 5.2, then the proof is quite similar to that of Case (5-14).

				,	,
Case	Φ'	$E(\phi')$	Case	Φ'	$E(\phi')$
(5-1)	$\llbracket 1, 2, 5, 5 \rrbracket$	$\{15\}$	(5-9)	$[\![1, 2, 5, 10]\!]$	E^*
(5-3)	$[\![1,2,1,14]\!]$	$\{61\}$	(5-10)	$[\![2,4,1,14]\!]$	$\{61\}$
(5-4)	$[\![1,2,{f 2},{f 7}]\!]$	{30}	(5-11)	$[\![2, 1, 3, 14]\!]$	$\{60\}$
(5-5)	$[\![1,2,3,3]\!]$	$\{13\}$	(5-12)	$[\![3, 1, 1, 7]\!]$	$\{30\}$
(5-6)	$[\![1, 2, 4, 14]\!]$	{60}	(5-13)	$[\![\boldsymbol{1},\boldsymbol{1},\boldsymbol{2},\boldsymbol{14}]\!]$	$\{60\}$
(5-7)	$[\![1, 2, 10, 5]\!]$	$\{20\}$	(5-15)	$[\![1,1,3,4]\!]$	$\{18\}$
(5-8)	$[\![1,2,{\bf 5},{\bf 5}]\!]$	$\{15, 20\}$	(5-17)	$[\![1,2,3,3]\!]$	$\{12\}$
		E	$C^* = \{r \cdot 25\}$	$s - 5 \mid 1 \leq r \leq 4$	$, s \ge 1$

Table 5.2. All cases whose proofs are simair to that of (5-14)

Remark 5.2. (i) In [3], Dickson proved that the sum [[1, 2, 5, 5]] represents all positive integers, except 15.

(ii) In [5], it was proved that the sums [[1, 1, 2, 14]], [[1, 1, 3, 4]], and [[1, 2, 3, 3]] represent all positive integers, except 60, 18, and 12, respectively.

Case (5-16) $(\beta_1, \beta_2, \beta_3, \beta_4) = (1, 1, 3, 7)$. It is enough to show that for any α_1 such that $8 \leq \alpha_1 \leq 14$, the equation

$$3\alpha_1 x^2 + y^2 + z^2 + 3t^2 + 7s^2 = 3N + 12$$

has an integer solution $(x, y, z, t, s) \in \mathbb{Z}^5$ such that $yzts \neq 0 \pmod{3}$ for any nonnegative integer N.

Assume $\alpha_1 = 8$. If $N \leq 1742$, then one may directly check that the equation

$$24x^2 + y^2 + z^2 + 3t^2 + 7s^2 = 3N + 12$$

has an integer solution $(x, y, z, t, s) \in \mathbb{Z}^5$ such that $yzts \neq 0 \pmod{3}$. Therefore, assume $N \geq 1743$. Note that the genus of $f(y, z, t) = (3y + t)^2 + (3z + t)^2 + 3t^2$ consists of

$$M_f = \begin{pmatrix} 5 & -2 & 2\\ -2 & 8 & 1\\ 2 & 1 & 8 \end{pmatrix} \text{ and } M_2 = \begin{pmatrix} 2 & 1\\ 1 & 5 \end{pmatrix} \perp \langle 27 \rangle.$$

One may easily show that a positive integer congruent to 2 modulo 3 is represented by M_f or M_2 by 102:5 of [12] for it is represented by M_f over \mathbb{Z}_p for any prime p. Note that

$$B_f(M_2, 13, 1) = \{\pm(0, 0, 1)\}, \qquad B_f(M_2, 13, 3) = \{\pm(0, 0, 4)\}, \\B_f(M_2, 13, 4) = \{\pm(0, 0, 2)\}, \qquad B_f(M_2, 13, 9) = \{\pm(0, 0, 3)\}, \\B_f(M_2, 13, 10) = \{\pm(0, 0, 6)\}, \qquad B_f(M_2, 13, 12) = \{\pm(0, 0, 5)\}.$$

In each case, if we define $T = \begin{pmatrix} 9 & 20 & 0 \\ -8 & 1 & 0 \\ 0 & 0 & 13 \end{pmatrix}$, then one may easily show that it

satisfies all conditions in Theorem 2.1. Note that $\pm(0,0,1)$ are the only integral primitive eigenvectors of T. Then we have

$$S_{13,r} \cap Q(M_2) \setminus \{27s^2 \mid s \in \mathbb{Z}\} \subset Q(M_f),$$

for any remainder r modulo 13, where r is a quadratic residue modulo 13. Therefore, a positive integer which is congruent to 2 modulo 3 and quadratic residue modulo 13 is represented by M_f . There are integers $a \in \{1, 2, 3, 13\}$ and $e \in \{1, 2, 13\}$ such that $3N + 12 - 7e^2 - 24a^2 > 0$ and furthermore, $3N + 12 - 7e^2 - 24a^2$ is represented by M_f . Therefore, the equation

$$y^2 + z^2 + 3t^2 = 3N + 12 - 7e^2 - 24a^2$$

has an integer solution $(y, z, t) = (b, c, d) \in \mathbb{Z}^3$ such that $b \equiv c \equiv d \pmod{3}$. This competes the proof.

Assume that $\alpha_1 \neq 13$. Then every proof is quite similar to that of the case $\alpha_1 = 8$.

Assume $\alpha_1 = 13$. If $N \leq 18$, then one may directly check that the equation

$$39x^2 + y^2 + z^2 + 3t^2 + 7s^2 = 3N + 12$$

has an integer solution $(x, y, z, t, s) \in \mathbb{Z}^5$ such that $yzts \neq 0 \pmod{3}$. Therefore, assume $N \geq 19$. Note that $h(\langle 1, 1, 3 \rangle) = 1$. There are integers $e \in \{1, 2\}$ and $a \in \{0, 1\}$ such that $3N + 12 - 7e^2 - 39a^2 > 0$ and $3N + 12 - 7e^2 - 39a^2$ is congruent to 5 or 8 modulo 12. Therefore, the equation

$$y^2 + z^2 + 3t^2 = 3N + 12 - 7e^2 - 39a^2$$

has an integer solution $(y, z, t) = (b, c, d) \in \mathbb{Z}^3$ such that $bc \neq 0 \pmod{3}$ and $c \equiv d \pmod{2}$. If $d \neq 0 \pmod{3}$, then we are done. Assume $d \equiv 0 \pmod{3}$. Then $c^2 + 3d^2 = \left(\frac{c+3d}{2}\right)^2 + 3 \cdot \left(\frac{c-d}{2}\right)^2$ and $\left(\frac{c+3d}{2}\right) \cdot \left(\frac{c-d}{2}\right) \neq 0 \pmod{3}$. This completes the proof of Case (5-16).

Theorem 5.3. There are exactly 11 senary proper universal mixed sums of generalized 4- and 8-gonal numbers (for the list of them, see Table 5.1).

Proof. Let u, v be nonnegative integers such that u + v = 6. For positive integers $\alpha_1, \ldots, \alpha_u$ and β_1, \ldots, β_v , assume that a senary mixed sum

$$\Phi = \llbracket \alpha_1, \ldots, \alpha_u, \beta_1, \ldots, \beta_v \rrbracket$$

of generalized 4- and 8-gonal numbers is proper universal. Then by Theorems 3.1, 4.2, and 5.1, Φ has a proper sum $[\![1, 2, 5, 5, 5]\!]$ (see Table 5.1). From the universality of Φ , we know that at least one of α_3 or β_4 is less than equal to 20 that is the truant of $[\![1, 2, 5, 5, 5]\!]$.

Case (6-1) $(\alpha_1, \alpha_2, \beta_1, \beta_2, \beta_3) = (1, 2, 5, 5, 5)$. It is enough to show that for any $\alpha_3(\beta_4)$ such that $16 \leq \alpha_3 \leq 20$ $(\beta_4 = 5 \text{ or } 16 \leq \beta_4 \leq 20)$, the equation

$$x^{2} + 2y^{2} + \alpha_{3}z^{2} + 5(3t^{2} - 2t) + 5(3s^{2} - 2s) + 5(3u^{2} - 2u) = N$$

(x² + 2y² + 5(3z² - 2z) + 5(3t² - 2t) + 5(3s^{2} - 2s) + \beta_{4}(3u^{2} - 2u) = N, \text{ respectively})

has an integer solution $(x, y, z, t, s, u) \in \mathbb{Z}^6$ for any nonnegative integer N. Since the mixed sum $[\![1, 2, 5, 5]\!]$ of generalized 4- and 8-gonal numbers represents all nonnegative integers except 15 and 20 (see Table 5.2), the quinary mixed sum $[\![1, 2, 5, 5, 5]\!]$ of generalized 4- and 8-gonal numbers represents all positive integers except 20. The proof follows immediately. Finally, we give an effective criterion on the universality of an arbitrary mixed sum of generalized 4- and 8-gonal numbers.

Theorem 5.4. Let $\alpha_1, \ldots, \alpha_u$ and β_1, \ldots, β_v be positive integers. The mixed sum

$$\Phi = \alpha_1 P_4(x_1) + \dots + \alpha_u P_4(x_u) + \beta_1 P_8(y_1) + \dots + \beta_v P_8(y_v)$$

of generalized 4- and 8-gonal numbers is universal if and only if it represents the integers

1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 15, 18, 20, 30, 60, and 61.

Proof. Assume that $\Phi = [\![\alpha_1, \ldots, \alpha_u, \beta_1, \ldots, \beta_v]\!]$ represents above 19 integers. Then one may easily show that there are nonnegative integers u_1, v_1 such that $u_1 \leq u, v_1 \leq v$, and furthermore, the proper sum

$$\Phi = \llbracket \alpha_1, \ldots, \alpha_{u_1}, \beta_1, \ldots, \beta_{v_1} \rrbracket$$

of Φ is one of the mixed sums of Tables 3.1, 4.1, and 5.1. Then the the proper sum $\tilde{\Phi}$ of Φ is universal by Theorems 3.1, 4.2, 5.1, and 5.3. Therefore, the mixed sum Φ of generalized 4- and 8-gonal numbers is universal.

References

- M. Bhargava, On the Conway-Schneeberger fifteen theorem, Contem. Math. 272(2000), 27-38.
- [2] M. Bhargava and J. Hanke, Universal quadratic forms and the 290 theorem, to appear.
- [3] L. E. Dickson, Quaternary quadratic forms representing all integers, Amer. J. Math. 49(1927), 39-56.
- W. C. Jagy, Five regular or nearly-regular ternary quadratic forms, Acta Arith. 77(1996), 361-367.
- [5] J. Ju and B.-K. Oh. Universal sums of generalized octagonal numbers, J. Number Theory 190(2018), 292-302.
- [6] J. Ju, B.-K. Oh, and B. Seo, Ternary universal sums of generalized polygonal numbers, submitted.
- [7] B. W. Jones, Representation by Positive Ternary Quadratic Forms, unpublished Ph.D. dissertation, University of Chicago, 1928.
- [8] Y. Kitaoka, Arithmetic of quadratic forms, Cambridge University Press, 1993.
- [9] B.-K. Oh, Regular positive ternary quadratic forms, Acta Arith. 147(2011), 233-243.
- [10] B.-K. Oh and H. Yu, Completely p-primitive binary quadratic forms, J. Number Theory 193(2018), 373-385.
- [11] B.-K. Oh, Ternary universal sums of generalized pentagonal numbers, J. Korean Math. Soc. 48(2011), 837-847.
- [12] O. T. O'Meara, Introduction to quadratic forms, Springer Verlag, New York, 1963.
- [13] Z.-W. Sun, A result similar to Lagrange's theorem, J. Number Theory 162(2016), 190-211.
- [14] T. Yang, An explicit formula for local densities of quadratic forms, J. Number Theory 72(1998), 309-356.

RESEARCH INSTITUTE OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL 08826, KOREA *E-mail address*: jjw@snu.ac.kr

DEPARTMENT OF MATHEMATICAL SCIENCES AND RESEARCH INSTITUTE OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL 08826, KOREA

E-mail address: bkoh@snu.ac.kr