# A CLASSIFICATION OF PROPER HOLOMORPHIC MAPPINGS BETWEEN GENERALIZED PSEUDOELLIPSOIDS OF DIFFERENT DIMENSIONS

#### ATSUSHI HAYASHIMOTO

ABSTRACT. We classify proper holomorphic mappings between generalized pseudoellipsoids of different dimensions. Those domains are parametrized by the exponents. The relations among them are also obtained. Main tool is the orthogonal decomposition of a CR bundle. Such a decomposition derives the "variable-splitting" of the mapping.

#### 1. Introduction

Let  $\alpha_1, \ldots, \alpha_{N-1}$  be positive integers with  $\alpha_1, \ldots, \alpha_{N-1} \geq 2$  and  $w_j = (w_j^1, \ldots, w_j^{m_j}) \in \mathbb{C}^{m_j}, w = (w_1, \ldots, w_N) \in \mathbb{C}^{m_1} \times \cdots \times \mathbb{C}^{m_N}, m_1 + \cdots + m_N = m$  and  $z \in \mathbb{C}$ . Let R(z, w) be a real analytic function defined by

$$R(z, w) = |z|^2 + \sum_{j=1}^{N-1} ||w_j||^{2\alpha_j} + ||w_N||^2 - 1,$$

where  $||w_j||^2 = |w_j^1|^2 + \cdots + |w_j^{m_j}|^2$  denotes the squared Euclidean norm.

Let  $E(m; m_1, \ldots, m_N; \alpha_1, \ldots, \alpha_{N-1}, 1)$  be a bounded domain in  $\mathbb{C}^{m+1}$  with real analytic boundary defined by the following:

$$E(m; m_1, \dots, m_N; \alpha_1, \dots, \alpha_{N-1}, 1)$$

$$= \{ (z, w_1, \dots, w_N) \in \mathbb{C} \times \mathbb{C}^{m_1} \times \dots \times \mathbb{C}^{m_N} : R(z, w) < 0 \}.$$

We call this domain a generalized pseudoellipsoid with N blocks. We abbreviate  $E(m; m_1, \ldots, m_N; \alpha_1, \ldots, \alpha_{N-1}, 1)$  as  $E(m; (m); (\alpha))$ . We sometimes write  $\alpha_N = 1$ . Let  $H(m; (m); (\alpha))$  be a pseudoconvex domain defined by

$$H(m;(m);(\alpha)) = \{(z, w_1, \dots, w_N) \in \mathbb{C} \times \mathbb{C}^{m_1} \times \dots \times \mathbb{C}^{m_N} : \\ \operatorname{Im} z > ||w_1||^{2\alpha_1} + \dots + ||w_{N-1}||^{2\alpha_{N-1}} + ||w_N||^2 \}.$$

The generalized pseudoellipsoid  $E(m;(m);(\alpha))$  is biholomorphically equivalent to its unbounded representation  $H(m;(m);(\alpha))$  via the mapping

(1) 
$$\Psi: H(m; (m); (\alpha)) \ni (z, w_1, \dots, w_N)$$

$$\mapsto (\frac{i-z}{i+z}, \frac{2^{1/\alpha_1} w_1}{(i+z)^{1/\alpha_1}}, \dots, \frac{2^{1/\alpha_{N-1}} w_{N-1}}{(i+z)^{1/\alpha_{N-1}}}, \frac{2w_N}{i+z}) \in E(m; (m); (\alpha)).$$

We study a proper holomorphic mapping

$$(\mathcal{F},\mathcal{G}) = (\mathcal{F},\mathcal{G}_1,\ldots,\mathcal{G}_N) : E(m;(m);(\alpha)) \to E(n;(n);(\beta))$$

This work was supported by Grant-in-Aid for Scientific Research (C) 17K05308

or its unbounded representation

$$\Phi = \widetilde{\Psi}^{-1} \circ (\mathcal{F}, \mathcal{G}) \circ \Psi : H(m; (m); (\alpha)) \to H(n; (n); (\beta)),$$

here  $\widetilde{\Psi}$  is a mapping between  $H(n;(n);(\beta))$  and  $E(n;(n);(\beta))$  defined analogously by (1). Let  $\Phi = (F,G) = (F_1,G_1,\ldots,G_N)$ . If  $(\mathcal{F},\mathcal{G})$  extends holomorphically past the boundary, so does (F,G). In this case, we use the extended mapping (F,G) in two meanings without any mention: one is the mapping between their closures:  $\overline{H(m;(m);(\alpha))} \to \overline{H(n;(n);(\beta))}$  and the other is its restriction to the boundaries:  $\partial H(m;(m);(\alpha)) \to \partial H(n;(n);(\beta))$ .

Let  $f, g: D_1 \to D_2$  be mappings with the same source and target domains. We say that f is equivalent to g if there exist automorphisms  $\phi_1 \in \operatorname{Aut}(D_1)$  and  $\phi_2 \in \operatorname{Aut}(D_2)$  such that  $f = \phi_2 \circ g \circ \phi_1$  holds.

Now we introduce a homogeneous proper holomorphic mapping  $H_M(z)$  between balls of different dimensions. For a variable  $z = (z_1, \ldots, z_n) \in \mathbb{C}^n$ , we define the tensor product by

$$z \otimes z = (z_1 z_1, z_2 z_1, \dots, z_n z_1, z_1 z_2, \dots, z_n z_n).$$

Namely, the tensor product of z and z is the mapping whose components are all possible products of the components of z. Repeat tensor products M times and after applying a unitary transformation that collects the same monomials together and ignoring zero components, we denote by the resulted mapping  $H_M(z)$ , which has an expression:

(2) 
$$H_M(z) = (\ldots, \sqrt{\frac{M!}{p_1! \ldots p_n!}} z^p, \ldots)$$

for  $p = (p_1, \ldots, p_n) \in (\mathbb{Z}_{\geq 0})^n$ ,  $p_1 + \cdots + p_n = M$ . D'Angelo introduced this mapping to classify proper holomorphic mappings between balls of different dimensions, in fact,

$$H_M(z): B^n \to B^N, N = \binom{n+M-1}{M}$$

is a proper holomorphic mapping and satisfies the equality  $||z||^{2M} = ||H_M(z)||^2$ . For this mapping, refer the book [D].

With the above notations and terminology, we can state our result as follows:

**Main Theorem 1.1.** Let  $E(m;(m);(\alpha))$  and  $E(n;(n);(\beta))$  be generalized pseudoellipsoids with N blocks. Let  $(\mathcal{F}, \mathcal{G}_1, \ldots, \mathcal{G}_N) : E(m;(m);(\alpha)) \to E(n;(n);(\beta))$  be a proper holomorphic mapping that is holomorphic up to the boundary. Assume that all components  $\mathcal{F}, \mathcal{G}_1, \ldots, \mathcal{G}_N$  are not constant and that  $m_j \geq 2, j = 1, \ldots, N-1$ . Then we have the following:

(1) There exists a permutation  $\sigma$  of  $\{1, \ldots, N-1\}$  such that

$$\mathcal{G}_j|_{w_{\sigma(j)}=0}=0, \quad j=1,\ldots,N-1.$$

(2) There exist integers  $M_1, \ldots, M_{N-1}$  such that  $\alpha_{\sigma(j)} = M_j \beta_j, j = 1, \ldots, N-1$ .

(3) 
$$n_j \ge \frac{(M_j + m_{\sigma(j)} - 1)!}{M_j!(m_{\sigma(j)} - 1)!}, \quad j = 1, \dots, N - 1.$$

(4) 
$$(\mathcal{F}, \mathcal{G}_1, \dots, \mathcal{G}_N)$$
 is equivalent to  $(\widetilde{\mathcal{F}}, \widetilde{\mathcal{G}}_1, \dots, \widetilde{\mathcal{G}}_N)$  of the form  $\widetilde{\mathcal{F}}(z, w_1, \dots, w_N) = z,$   $\widetilde{\mathcal{G}}_j(z, w_1, \dots, w_N) = (H_{M_j}(w_{\sigma(j)}), 0), \quad j = 1, \dots, N-1,$   $\widetilde{\mathcal{G}}_N(z, w_1, \dots, w_N) = (w_N, 0).$ 

Remark 1.1. Outline of the proof is the following.

- (1) Decompose a CR vector bundle into a special kind of direct sum and the decomposition derives expansions of the mapping. See Sections 2 and 3.
- (2) Restricting our situation to a certain variety, we get the relations of exponents  $\alpha_i$  and  $\beta_i$ . See Section 4.
- (3) Again we restrict to a certain variety to reduce an N blocks case to a one block case. Then we obtain the normalization of a certain component of the mapping. Varying the variety to restrict, we obtain the whole normalization of the mapping. See Sections 5 and 6.

If  $m_j = 1$ , we call the domain  $E(m; (1), \alpha)$  a pseudoellipsoid. For pseudoellipsoids and generalized pseudoellipsoids, following topics have been studied.

- (1) Given two (generalized) pseudoellipsoids, find necessary and sufficient conditions to exist proper holomorphic mappings between them. Classify the mappings between them. [DP1], [DP2], [Ha1], [Ha2], [K4], [L], [MM]
- (2) Given any local biholomorphic mapping on a neighborhood of a boundary point such that the point is mapped to a boundary point, prove that it can be extended to a global biholomorphic mapping. [DP2], [K3]
- (3) Characterize (generalized) pseudoellipsoids by mean of their holomorphic automorphism groups. [GK], [K1], [K2], [K3], [KKM]

Present article is related to (1). The first result on this line is done by M. Landucci [L]. He studied pseudoellipsoids

$$E(\alpha) = \{ z \in \mathbb{C}^n : \sum_{k=1}^n |z_k|^{2\alpha_k} < 1 \}$$

for  $\alpha_k \in \mathbb{N}$ . If there exists a proper holomorphic mapping  $f: E(\alpha) \to E(\beta)$ , then, after reordering the variables,  $\alpha_j/\beta_j \in \mathbb{N}$  holds, and the mapping is equivalent to  $(z_1,\ldots,z_n) \mapsto (z_1^{\alpha_1/\beta_1},\ldots,z_n^{\alpha_n/\beta_n})$ . Another proof of the relations between  $\alpha_j$  and  $\beta_j$  was given by the author [Ha1] using the tangential Cauchy-Riemann equation. After that, G. Dini-A. S. Primicerio [DP1] proved the case of  $\alpha_j \in \mathbb{R}_{>0}$  in Landucci's theorem. The determination of the mappings between generalized pseudoellipsoids was given by R. Monti-D. Morbidelli [MM]. They studied local CR mappings, which are viewed as boundary values of proper holomorphic mappings, between boundaries of generalized pseudoellipsoids and decomposed them to elementary mappings, namely, any such local CR mappings are composite mappings of inversions, one-parameter group of dilations, a kind of unitary transformations, and shifts. G. Dini and A. S. Primicerio studied more general domain, which is denoted by  $\Sigma(m,\beta,d)$ , and obtained the relations of indices. Let  $\Sigma(m,\beta,d)$  be a domain defined by

$$\Sigma(m,\beta,d) = \{ z \in \mathbb{C}^N : \sum_{k=1}^n \left( \sum_{j=s_{k-1}}^{s_k-1} |z_j|^{2d_j} \right)^{\beta_k} < 1 \},$$

where  $d = (d_1, \ldots, d_N) \in \mathbb{N}^N$ ,  $\beta = (\beta_1, \ldots, \beta_n) \in \mathbb{N}^n$ , and  $N = m_1 + \cdots + m_n$ ,  $s_0 = 1$ ,  $s_k = s_{k-1} + m_k$  and  $m = (m_1, \ldots, m_n)$ . If there exists a proper holomorphic mapping between  $\Sigma(p, \alpha, c)$  and  $\Sigma(m, \beta, d)$ , then, up to reordering the variables, p = m,  $\alpha = \beta$  and  $c_j/d_j \in \mathbb{N}$ .

All results mentioned above are for the equi-dimensional case, namely, source and target domains have the same dimensions. On the other hand, our present research is to study the higher codimensional case. By codimension, we mean the difference of the source and the target dimensions. When we study such a case, we find some strange phenomena. For instance, by [Do], there exists a proper holomorphic mapping  $f:B^N\to B^{N+1}$  that can be extended continuously to the closure  $\overline{B^N}$  but can not be extended in a  $C^2$  way to any open subset of the boundary of  $B^N$ . The determination of proper holomorphic mappings between pseudoellipsoids for small codimensional case was studied by P. Ebenfelt-D. N. Son [ES] and they obtained the explicit parametrization of the mappings. In the case of generalized pseudoellipsoids, under some assumptions on dimensions and nondegeneracy condition of a certain kind of Jacobian matrix, the author [Ha2] obtained the conditions to exist mappings between those domains and gave equivalence classes of the mappings. The purpose of this article is to weaken those assumptions. The relations between author's previous results and the present results are discussed in the final section.

We list some notation. Let  $w_j=(w_j^1,\ldots,w_j^{m_j})\in\mathbb{C}^{m_j}, w=(w_1,\ldots,w_N)\in\mathbb{C}^{m_1}\times\cdots\times\mathbb{C}^{m_N},$  and  $p_j=(p_j^1,\ldots,p_j^{m_j})\in(\mathbb{Z}_{\geq 0})^{m_j},\ p=(p_1,\ldots,p_N)\in(\mathbb{Z}_{\geq 0})^{m_1}\times\cdots\times(\mathbb{Z}_{\geq 0})^{m_N},$  and  $\alpha=(\alpha_1,\ldots,\alpha_{N-1},1)\in\mathbb{N}^N.$ 

- The multi-index notations;  $(w_j)^{p_j} = (w_j^1)^{p_j^1} \cdots (w_j^{m_j})^{p_j^{m_j}}$  and  $(w)^p = (w_1)^{p_1} \times \cdots \times (w_N)^{p_N}$ .
- $|||w|||^{2\alpha} = ||w_1||^{2\alpha_1} + \dots + ||w_{N-1}||^{2\alpha_{N-1}} + ||w_N||^2$ .  $|p_j| = p_j^1 + \dots + p_j^{m_j}$  and  $|p| = |p_1| + \dots + |p_N|$ .
- The total degree of  $w_j$  and  $\overline{w_j}$  in a monomial  $(w_j)^{p_j} \times (\overline{w_j})^{q_j}$  is the sum  $|p_j| + |q_j|$ .
- $\partial\Omega$  is the boundary of the domain  $\Omega$ .  $\partial^*\Omega$  is the strongly pseudoconvex part of  $\partial\Omega$ .
- Re z = x.

When we need to distinguish the notation of source and target objects, we add 'tilde' on the target objects.

The author would like to express his sincere thanks to Professor Akio Kodama for reading this article and pointing out some mistakes. His advice was very valuable and made some arguments clear.

## 2. DECOMPOSITION OF A CR VECTOR BUNDLE

Following [MM], we decompose a CR vector bundle. We employ the usual coordinate  $(x, w) \in \mathbb{R} \times \mathbb{C}^m$  on the boundary of  $H(m; (m); (\alpha))$ , that is, we identify naturally (x, w) with  $(x + i|||w|||^{2\alpha}, w)$  throughout this paper.

We define CR vector fields  $L_i^{\lambda}$  by

$$L_j^{\lambda} = \frac{\partial}{\partial w_j^{\lambda}} + i\alpha_j ||w_j||^{2(\alpha_j - 1)} \overline{w_j^{\lambda}} \frac{\partial}{\partial x}, \qquad j = 1, \dots, N, \quad \lambda = 1, \dots, m_j.$$

Define the pseudohermitian structure  $\vartheta$  as

$$\vartheta = dx - i \sum_{j=1}^{N} \sum_{\mu=1}^{m_j} \alpha_j ||w_j||^{2(\alpha_j - 1)} \left( \overline{w_j^{\mu}} dw_j^{\mu} - w_j^{\mu} d\overline{w_j^{\mu}} \right).$$

We define the Levi form on  $\partial H(m;(m);(\alpha))$  to be a 2-form: Levi =  $-id\vartheta$ . We introduce the holomorphic vector fields on  $\partial^* H(m;(m);(\alpha))$ :

$$E_j = \frac{1}{\alpha_j} \sum_{\mu=1}^{m_j} w_j^{\mu} L_j^{\mu}, \ j = 1, \dots, N,$$

which is called a vector field of radial type. The vector fields  $E_1, \ldots, E_N$  span an N-dimensional subbundle  $\mathcal{E} \subset T^{1,0}\partial^*H(m;(m);(\alpha))$ . We denote by  $\mathcal{E}^{\perp}$  the orthogonal component of  $\mathcal{E}$  in  $T^{1,0}\partial^*H(m;(m);(\alpha))$  with respect to the Levi form.

Let  $Q: T^{1,0}\partial^*H(m;(m);(\alpha)) \to \mathcal{E}^{\perp}$  be the projection

$$Q(X) = X - \sum_{j=1}^{N} \frac{\text{Levi}(X, \overline{E_j})}{\text{Levi}(E_j, \overline{E_j})} E_j.$$

Let  $W_i^{\lambda} = Q(L_i^{\lambda})$ . Then, by calculation, it is written as

$$W_{j}^{\lambda} = \frac{\partial}{\partial w_{j}^{\lambda}} - \sum_{\mu=1}^{m_{j}} \frac{\overline{w_{j}^{\lambda}} w_{j}^{\mu}}{||w_{j}||^{2}} \frac{\partial}{\partial w_{j}^{\mu}}$$

$$= \sum_{\mu=1}^{m_{j}} \left\{ \delta_{\mu}^{\lambda} - \frac{\overline{w_{j}^{\lambda}} w_{j}^{\mu}}{||w_{j}||^{2}} \right\} L_{j}^{\mu}.$$
(3)

Here  $\delta_{\mu}^{\lambda}$  is the Kronecker's delta. Then the bundle  $\mathcal{E}^{\perp}$  is an (m-N)-dimensional subbundle of  $T^{1,0}\partial^*H(m;(m);(\alpha))$  and is generated by the set of vector fields  $\{W_i^{\lambda}\}$ .

Let  $W_j$  be the subbundle of  $T^{1,0}\partial^*H(m;(m);(\alpha))$  generated by  $W_j^1,\ldots,W_j^{m_j}$ . Since the relation  $w_j^1W_j^1+\cdots+w_j^{m_j}W_j^{m_j}=0$  holds, the subbundle  $\mathcal{W}_j$  is, in fact, generated by, say,  $W_j^1,\ldots,W_j^{m_j-1}$ . Then we have  $\mathcal{E}^{\perp}=\mathcal{W}_1\oplus\cdots\oplus\mathcal{W}_N$  and  $\mathcal{W}_j\perp\mathcal{W}_k$  if  $j\neq k$ . Hence we obtain the orthogonal decomposition

(4) 
$$T^{1,0}\partial^*H(m;(m);(\alpha)) = \mathcal{W}_1 \oplus \cdots \oplus \mathcal{W}_N \oplus \mathcal{E}.$$

We sometimes use the notation  $\mathcal{E} = \mathcal{W}_{N+1}$ . For later use, we here list the values of the Levi form as follows: Using the fact that  $\theta(L_i^{\lambda}) = 0$  and

$$\theta([L_j^{\lambda}, \overline{L_j^{\mu}}]) = -2i\alpha_j ||w_j||^{2(\alpha_j - 1)} \Big\{ \delta_{\lambda}^{\mu} + (\alpha_j - 1) \frac{\overline{w_j^{\lambda}} w_j^{\mu}}{||w_j||^2} \Big\},$$

we obtain that

(5) 
$$\operatorname{Levi}(W_j^{\lambda}, \overline{W_j^{\mu}}) = -2\alpha_j ||w_j||^{2(\alpha_j - 1)} \frac{\overline{w_j^{\lambda}} w_j^{\mu}}{||w_j||^2}, \quad \operatorname{Levi}(W_j^{\lambda}, \overline{W_k^{\mu}}) = 0, \ j \neq k,$$

$$\operatorname{Levi}(W_j^{\lambda}, \overline{E_k}) = 0, \quad \operatorname{Levi}(E_j, \overline{E_j}) = 2||w_j||^{2\alpha_j}.$$

Rest of this section is devoted to obtain relations among  $W_j$ 's,  $\widetilde{W}_i$ 's and  $\Phi_*$ . In the following, for  $j, k = 1, \ldots, N$ , we denote by  $\widetilde{W}_{j,k}^{\lambda}$  (resp.  $\widetilde{W}_{j,N+1}^{\lambda}$ ) the  $\widetilde{W}_k$ -component (resp. the  $\widetilde{\mathcal{E}}$ -component) of  $\Phi_*W_j^{\lambda}$ . For a fixed  $j = 1, \ldots, N$ , also denote by  $\widetilde{W}_{N+1,k}$ ,

k = 1, ..., N (resp.  $\widetilde{W}_{N+1,N+1}$ ) the  $\widetilde{W}_k$ -component (resp. the  $\widetilde{\mathcal{E}}$ -component) of  $\Phi_* E_j$ . Let  $J_i$  be a subset of  $\{1, ..., N\}$  defined by  $J_i = \{\ell \in \{1, ..., N\} : \widetilde{W}_{\ell}$ -component of  $\Phi_* W_i$  is equal to zero $\}$ .

**Lemma 2.1.** For any j, k = 1, ..., N with  $j \neq k$ , we have  $J_i \cap J_k = \emptyset$ .

*Proof.* Take any  $W_j^{\lambda} \in \mathcal{W}_j$  and  $W_k^{\mu} \in \mathcal{W}_k$  for j, k = 1, ..., N with  $j \neq k$ . Let us denote by

(6) 
$$\Phi_* W_i^{\lambda} = \widetilde{W}_{i,1}^{\lambda} + \dots + \widetilde{W}_{i,N}^{\lambda} + \widetilde{W}_{i,N+1}^{\lambda},$$

(7) 
$$\Phi_* W_k^{\mu} = \widetilde{W}_{k,1}^{\mu} + \dots + \widetilde{W}_{k,N}^{\mu} + \widetilde{W}_{k,N+1}^{\mu}.$$

We calculate the Levi form  $\widetilde{\text{Levi}}(\Phi_*W_j^{\lambda}, \overline{\Phi_*W_k^{\mu}})$  in two ways. Since the mapping  $\Phi$  is CR, there exists a positive function f such that  $\Phi^*\widetilde{\vartheta} = f\vartheta$ . Hence we have

$$\widetilde{\operatorname{Levi}}(\Phi_*W_j^{\lambda},\overline{\Phi_*W_k^{\mu}}) = f\operatorname{Levi}(W_j^{\lambda},\overline{W_k^{\mu}}) = 0.$$

On the other hand, by using (6) and (7), it is expanded as follows:

$$(8) \quad \widetilde{\operatorname{Levi}}(\Phi_{*}W_{j}^{\lambda}, \overline{\Phi_{*}W_{k}^{\mu}}) \\ = \widetilde{\operatorname{Levi}}(\widetilde{W}_{j,1}^{\lambda}, \overline{\widetilde{W}_{k,1}^{\mu}}) + \cdots + \widetilde{\operatorname{Levi}}(\widetilde{W}_{j,N}^{\lambda}, \overline{\widetilde{W}_{k,N}^{\mu}}) + \widetilde{\operatorname{Levi}}(\widetilde{W}_{j,N+1}^{\lambda}, \overline{\widetilde{W}_{k,N+1}^{\mu}}) \\ + \widetilde{\operatorname{Levi}}(\widetilde{W}_{j,1}^{\lambda}, \overline{\widetilde{W}_{k,N+1}^{\mu}}) + \cdots + \widetilde{\operatorname{Levi}}(\widetilde{W}_{j,N}^{\lambda}, \overline{\widetilde{W}_{k,N+1}^{\mu}}) \\ + \widetilde{\operatorname{Levi}}(\widetilde{W}_{i,N+1}^{\lambda}, \overline{\widetilde{W}_{k,1}^{\mu}}) + \cdots + \widetilde{\operatorname{Levi}}(\widetilde{W}_{i,N+1}^{\lambda}, \overline{\widetilde{W}_{k,N}^{\mu}}).$$

For i = 1, ..., N, let us denote by

$$\begin{split} \widetilde{W}_{j,i}^{\lambda} &= a_i^1 \widetilde{W}_i^1 + \dots + a_i^{n_i - 1} \widetilde{W}_i^{n_i - 1}, \quad \widetilde{W}_{k,i}^{\mu} = b_i^1 \widetilde{W}_i^1 + \dots + b_i^{n_i - 1} \widetilde{W}_i^{n_i - 1}, \\ \widetilde{W}_{j,N+1}^{\lambda} &= A_1 \widetilde{E}_1 + \dots + A_N \widetilde{E}_N, \quad \widetilde{W}_{k,N+1}^{\mu} = B_1 \widetilde{E}_1 + \dots + B_N \widetilde{E}_N, \end{split}$$

then it follows from (5) that each term on the right hand side of (8) is calculated as follows:

$$\widetilde{\operatorname{Levi}}(\widetilde{W}_{j,i}^{\lambda}, \overline{\widetilde{W}_{k,i}^{\mu}}) = -2\beta_{i}||\widetilde{w}_{i}||^{2(\beta_{i}-1)} \sum_{\nu_{1},\nu_{2}=1}^{n_{i}-1} a_{i}^{\nu_{1}} \overline{b_{i}^{\nu_{2}}} \overline{\widetilde{w}_{i}^{\nu_{1}}} \widetilde{w}_{i}^{\nu_{2}}, \ i = 1, \dots, N,$$

$$\widetilde{\operatorname{Levi}}(\widetilde{W}_{j,N+1}^{\lambda}, \overline{\widetilde{W}_{k,N+1}^{\mu}}) = 2(A_{1}\overline{B_{1}}||\widetilde{w}_{1}||^{2\beta_{1}} + \dots + A_{N}\overline{B_{N}}||\widetilde{w}_{N}||^{2}),$$

$$\widetilde{\operatorname{Levi}}(\widetilde{W}_{j,i}^{\lambda}, \overline{\widetilde{W}_{k,N+1}^{\mu}}) = \widetilde{\operatorname{Levi}}(\widetilde{W}_{j,N+1}^{\lambda}, \overline{\widetilde{W}_{k,i}^{\mu}}) = 0, \ i = 1, \dots, N.$$

Hence we obtain

$$-\sum_{i=1}^{N} 2\beta_{i}||\widetilde{w}_{i}||^{2(\beta_{i}-1)} \sum_{\nu_{1},\nu_{2}=1}^{n_{i}-1} a_{i}^{\nu_{1}} \overline{b_{i}^{\nu_{2}}} \frac{\overline{\widetilde{w}_{i}^{\nu_{1}}} \widetilde{w}_{i}^{\nu_{2}}}{||\widetilde{w}_{i}||^{2}} + 2\sum_{i=1}^{N} A_{i} \overline{B_{i}}||\widetilde{w}_{i}||^{2\beta_{i}} = 0,$$

which means that for each i = 1, ..., N,

$$a_i^1 \overline{b_i^1} = \dots = a_i^1 \overline{b_i^{n_i - 1}} = 0,$$

$$\vdots$$

$$a_i^{n_i - 1} \overline{b_i^1} = \dots = a_i^{n_i - 1} \overline{b_i^{n_i - 1}} = 0.$$

Take any  $i=1,\ldots,N$  and fix it. If  $a_i^{\lambda}=0$  for any  $\lambda=1,\ldots,n_i-1$ , then  $\widetilde{W}_{j,i}^{\lambda}=0$ ; and hence, the  $\widetilde{W}_i$ -component of  $\Phi_*W_j^{\lambda}$  is equal to zero. If there exists

 $\lambda = 1, \dots, n_i - 1$  such that  $a_i^{\lambda} \neq 0$ , then  $b_i^1 = \dots = b_i^{n_i - 1} = 0$  and this means that the  $\widetilde{\mathcal{W}}_i$ -component of  $\Phi_* W_k^{\mu}$  is equal to zero. Hence both  $\Phi_* W_j^{\lambda}$  and  $\Phi_* W_k^{\mu}$  do not have the  $\widetilde{\mathcal{W}}_i$ -component simultaneously.

The same conclusion holds for  $\Phi_*\mathcal{E}$  and  $\Phi_*\mathcal{W}_j$ .

**Lemma 2.2.** For any j = 1, ..., N, we have  $J_j \cap J_{N+1} = \emptyset$ .

*Proof.* Take any k = 1, ..., N and fix it. Let us denote by

$$\Phi_* E_k = \widetilde{W}_{N+1,1} + \dots + \widetilde{W}_{N+1,N} + \widetilde{W}_{N+1,N+1},$$
  
$$\Phi_* W_j^{\lambda} = \widetilde{W}_{j,1}^{\lambda} + \dots + \widetilde{W}_{j,N}^{\lambda} + \widetilde{W}_{j,N+1}^{\lambda}.$$

Lemma follows by the same argument as in Lemma 2.1.

**Lemma 2.3.** There does not exist  $W_j^{\lambda} \in \mathcal{W}_j$ , j = 1, ..., N such that  $\Phi_* W_j^{\lambda} \in \widetilde{\mathcal{E}}$ .

*Proof.* We use the expansion of  $G_{\ell}^{\mu}$ :

$$G_{\ell}^{\mu}(x,w) = \sum_{|p|+q>0} b_{\ell,p,q}^{\mu}(w)^{p} (x+i|||w|||^{2\alpha})^{q}.$$

Suppose that there exists  $W_j^{\lambda} \in \mathcal{W}_j$  such that  $\Phi_* W_j^{\lambda} \in \widetilde{\mathcal{E}}$  and express it as  $\Phi_* W_j^{\lambda} = A_1^{\lambda} \widetilde{E}_1 + \dots + A_N^{\lambda} \widetilde{E}_N$ . Take  $\ell = 1, \dots, N$  such that  $A_\ell^{\lambda} \neq 0$ . Apply  $\widetilde{w}_\ell^{\mu} = G_\ell^{\mu}$  to this relation to get the following:

$$(9) \qquad \frac{A_{\ell}^{\lambda}}{\beta_{\ell}} G_{\ell}^{\mu} = W_{j}^{\lambda} G_{\ell}^{\mu} = \sum_{\nu=1}^{m_{j}} \left( \delta_{\lambda}^{\nu} - \frac{\overline{w_{j}^{\lambda}} w_{j}^{\nu}}{||w_{j}||^{2}} \right) L_{j}^{\nu} G_{\ell}^{\mu}$$

$$= \sum_{\substack{|p|+q \geq 0 \\ p_{j}^{\lambda} \geq 1}} b_{\ell,p,q}^{\mu} \frac{\partial(w)^{p}}{\partial w_{j}^{\lambda}} (x+i|||w|||^{2\alpha})^{q}$$

$$- \frac{\overline{w_{j}^{\lambda}}}{||w_{j}||^{2}} \sum_{\nu=1}^{m_{j}} \sum_{\substack{|p|+q \geq 0 \\ p_{j}^{\nu} \geq 1}} b_{\ell,p,q}^{\mu} p_{j}^{\nu}(w)^{p} (x+i|||w|||^{2\alpha})^{q}.$$

For convenience, we want to rewrite the last summation in (9) as follows. To this end, let us introduce a subset  $P_k$  of  $\mathbb{Z}^{m_j}$  defined by

$$P_k = \{(p_j^1, \dots, p_j^{m_j}) \in \mathbb{Z}^{m_j} : k \text{ components among } p_j^1, \dots, p_j^{m_j} \text{ are positive and the rest are zero}\}.$$

Take any  $p_j^0 \in P_k$  such that its  $\lambda_1$ -th, ...,  $\lambda_k$ -th components are positive and the others are zero. Then, among  $m_j$  summations in the second term on the far right hand side of (9).

$$\sum_{\substack{|p|+q\geq 0\\ p_j^1\geq 1}} b_{\ell,p,q}^{\mu} p_j^1(w)^p (x+i|||w|||^{2\alpha})^q, \ldots, \sum_{\substack{|p|+q\geq 0\\ p_j^{m_j}\geq 1}} b_{\ell,p,q}^{\mu} p_j^{m_j}(w)^p (x+i|||w|||^{2\alpha})^q,$$

the term of the form:

$$\sum_{\substack{|p|+q \ge 0 \\ p_j = p_j^0}} b_{\ell,p,q}^{\mu} p_j^{\nu}(w)^p (x+i|||w|||^{2\alpha})^q$$

comes only from the  $\lambda_1$ -th, ..., the  $\lambda_k$ -th summations. Thus (9) can be rewritten as

$$\frac{A_{\ell}^{\lambda}}{\beta_{\ell}}G_{\ell}^{\mu} = \sum_{\substack{|p|+q \geq 0 \\ p_{j}^{\lambda} \geq 1}} b_{\ell,p,q}^{\mu} \frac{\partial(w)^{p}}{\partial w_{j}^{\lambda}} (x+i||w|||^{2\alpha})^{q} 
- \frac{\overline{w_{j}^{\lambda}}}{||w_{j}||^{2}} \Big\{ m_{j} \sum_{\substack{|p|+q \geq 0 \\ p_{j} \in P_{m_{j}}}} b_{\ell,p,q}^{\mu} |p_{j}| (w)^{p} (x+i|||w|||^{2\alpha})^{q} 
+ \dots + (m_{j} - k) \sum_{\substack{|p|+q \geq 0 \\ p_{j} \in P_{m_{j}-k}}} b_{\ell,p,q}^{\mu} |p_{j}| (w)^{p} (x+i|||w|||^{2\alpha})^{q} \Big\} 
+ \dots + \sum_{\substack{|p|+q \geq 0 \\ p_{j} \in P_{1}}} b_{\ell,p,q}^{\mu} |p_{j}| (w)^{p} (x+i|||w|||^{2\alpha})^{q} \Big\}.$$

Hence comparing the  $\overline{w_j^{\lambda}}$  terms and the  $(w)^p$  terms with  $p_j \in P_{m_j-k}$ , we have

$$\sum_{\substack{|p|+q\geq 0\\ p_j\in P_{m_j-k}}} b_{\ell,p,q}^{\mu} |p_j|(w)^p (x+i|||w|||^{2\alpha})^q = 0, \quad k = 0, \dots, m_j - 1$$

and thus  $G_{\ell}^{\mu}$  has an expansion

$$G_{\ell}^{\mu}(x,w) = \sum_{\substack{|p|+q \ge 0 \\ p_i = 0}} b_{\ell,p,q}^{\mu}(w)^p (x+i|||w|||^{2\alpha})^q.$$

This means that the variables  $w_j$  do not appear in  $(w)^p$ . Therefore again by the relation (9) and by  $A_\ell^{\lambda} \neq 0$ , we conclude that  $G_\ell^{\mu} = 0$  for any  $\mu = 1, \dots n_\ell$ . This contradicts the assumption on the main theorem.

From the proof of this lemma, the following corollary holds, which will be used in the next argument.

**Corollary 2.1.** Let f be a CR function on  $\partial H(m;(m);(\alpha))$ . Assume that there exist  $W_j^{\lambda} \in W_j$  and a constant A such that  $W_j^{\lambda} f = Af$ . If  $A \neq 0$ , then  $f \equiv 0$  and if A = 0, then f has an expansion:

$$f = \sum_{\substack{|p|+q \ge 0 \\ p_s = 0}} a_{p,q}(w)^p (x+i|||w|||^{2\alpha})^q.$$

From Lemma 2.1 to Lemma 2.3, we have proved that there exists a permutation  $\sigma$  of  $\{1,\ldots,N\}$  such that  $\Phi_*\mathcal{W}_{\sigma(j)}\subset\widetilde{\mathcal{W}}_j\oplus\widetilde{\mathcal{E}}$ . Here we assert that  $\Phi_*\mathcal{W}_{\sigma(j)}\subset\widetilde{\mathcal{W}}_j$  for all  $j=1,\ldots,N$ . We now verify this only in the case where j=1, since the verification in the general case is almost identical. So, take any  $W^{\lambda}_{\sigma(1)}\in\mathcal{W}_{\sigma(1)}$  and write

(10) 
$$\Phi_* W_{\sigma(1)}^{\lambda} = a_1^{\lambda} \widetilde{W}_1^1 + \dots + a_{n_1 - 1}^{\lambda} \widetilde{W}_1^{n_1 - 1} + A_1^{\lambda} \widetilde{E}_1 + \dots + A_N^{\lambda} \widetilde{E}_N.$$

Assume that there exist  $\lambda=1,\ldots,m_{\sigma(1)}$  and  $\ell=2,\ldots,N$  such that  $A_\ell^\lambda\neq 0$ . Apply  $\widetilde{w}_\ell^\mu=G_\ell^\mu$  to (10) to get  $W_{\sigma(1)}^\lambda G_\ell^\mu=(A_\ell^\lambda/\beta_\ell)G_\ell^\mu$ . Then by Corollary 2.1,  $G_\ell^\mu=0$  for any  $\mu=1,\ldots,n_\ell$ , which is a contradiction. Thus  $A_\ell^\lambda=0$  for any  $\lambda=1,\ldots,m_{\sigma(1)}$ 

and  $\ell=2,\ldots,N$ . Among  $A_1^1,\ldots,A_1^{m_{\sigma(1)}}$ , assume that  $A_1^{\lambda_1},\ldots,A_1^{\lambda_i}\neq 0$  and the others are zero. Apply  $\widetilde{x}=(F+\overline{F})/2$  to (10) to get  $(1/2)W_{\sigma(1)}^{\lambda}F=iA_1^{\lambda}||G_1||^{2\beta_1}$ . Since F has an expansion

(11) 
$$F = \sum_{|p|+q>0} a_{p,q}(w)^p (x+i|||w|||^{2\alpha})^q,$$

we get

(12) 
$$\frac{1}{2} \sum_{\substack{|p|+q \geq 0 \\ p_{\sigma(1)}^{\lambda} \geq 1}} a_{p,q} \frac{\partial(w)^{p}}{\partial w_{\sigma(1)}^{\lambda}} (x+i|||w|||^{2\alpha})^{q} \\ -\frac{1}{2} \frac{\overline{w_{\sigma(1)}^{\lambda}}}{||w_{\sigma(1)}||^{2}} \sum_{\nu=1}^{m_{\sigma(1)}} \sum_{\substack{|p|+q \geq 0 \\ p_{\sigma(1)}^{\nu} \geq 1}} a_{p,q} p_{\sigma(1)}^{\nu} (w)^{p} (x+i|||w|||^{2\alpha})^{q} \\ = iA_{1}^{\lambda} ||G_{1}||^{2\beta_{1}}$$

for any  $\lambda=1,\ldots,m_{\sigma(1)}$ . Multiply (12) by  $w_{\sigma(1)}^{\lambda}$  and add them from  $\lambda=1$  to  $\lambda=m_{\sigma(1)}$ . Then we have

$$0 = i||G_1||^{2\beta_1} \{ A_1^{\lambda_1} w_{\sigma(1)}^{\lambda_1} + \dots + A_1^{\lambda_i} w_{\sigma(1)}^{\lambda_i} \}$$

and this means that  $G_1 = 0$ . This is a contradiction. Thus  $A_1^{\lambda} = 0$  for all  $\lambda$ . Hence, the  $\widetilde{\mathcal{E}}$ -component of  $\Phi_*W_{\sigma(1)}^{\lambda}$  is equal to zero and  $\Phi_*W_{\sigma(1)} \subset \widetilde{W}_1$ , as asserted. By the same argument as above, it can be seen that  $\Phi_*\mathcal{E} \subset \widetilde{\mathcal{E}}$ . Summarizing the above, we obtain the following:

**Proposition 2.1.** There exists a permutation  $\sigma$  of  $\{1, ..., N\}$  such that  $\Phi_* W_{\sigma(j)} \subset \widetilde{W}_j$  for all j = 1, ..., N; and  $\Phi_* \mathcal{E} \subset \widetilde{\mathcal{E}}$ .

### 3. Expansions of F and G

Using Proposition 2.1, we can characterize the expansions of F and  $G_i^{\lambda}$ .

**Lemma 3.1.** Up to an automorphism of  $H(n;(n);(\beta))$ , the mapping (F,G) has the following expansions on  $\partial H(m;(m);(\alpha))$ :

(13) 
$$F(x,w) = x + i|||w|||^{2\alpha},$$

(14) 
$$G_j^{\mu}(w) = \sum_{|p_{\sigma(j)}| \ge 1} b_{j,p_{\sigma(j)}}^{\mu}(w_{\sigma(j)})^{p_{\sigma(j)}}, \ j = 1,\dots, N.$$

Proof.

Claim 3.1.  $G_j^{\mu}$  is expanded as

(15) 
$$G_j^{\mu}(w) = \sum_{|p_{\sigma(j)}| \ge 0} b_{j,p_{\sigma(j)}}^{\mu} (w_{\sigma(j)})^{p_{\sigma(j)}}.$$

Take any j = 1, ..., N and fix it. Since  $\Phi_* W_{\sigma(i)}^{\lambda} \in \widetilde{W}_i$ , by applying  $\widetilde{w}_j^{\mu} = G_j^{\mu}$  to  $\Phi_* W_{\sigma(i)}^{\lambda}$ ,  $i = 1, ..., N, \lambda = 1, ..., m_{\sigma(i)}$  with  $i \neq j$ , we get the relations:

$$\begin{split} W^1_{\sigma(1)}G^{\mu}_j &= 0, \dots, W^{m_{\sigma(1)}}_{\sigma(1)}G^{\mu}_j = 0, \\ &\vdots \\ W^1_{\sigma(N)}G^{\mu}_j &= 0, \dots, W^{m_{\sigma(N)}}_{\sigma(N)}G^{\mu}_j = 0. \end{split}$$

Note that  $W^1_{\sigma(j)}G^\mu_j=0,\ldots,W^{m_{\sigma(j)}}_{\sigma(j)}G^\mu_j=0$  are not contained in these relations. It follows from Corollary 2.1 that we obtain the expansion:

$$G_j^{\mu}(x,w) = \sum_{|p_{\sigma(j)}|+q \ge 0} b_{j,p_{\sigma(j)},q}^{\mu}(w_{\sigma(j)})^{p_{\sigma(j)}} (x+i|||w|||^{2\alpha})^q.$$

We apply  $\widetilde{w}_j^{\mu} = G_j^{\mu}$  to  $\Phi_* E_k = A_1^k \widetilde{E}_1 + \dots + A_N^k \widetilde{E}_N$  with  $k \neq \sigma(j)$ . Assume that  $A_j^k \neq 0$  for all  $k = 1, \dots, N$  and  $k \neq \sigma(j)$ . Then we get the relation  $E_k G_j^{\mu} = A_j^k \widetilde{E}_j \widetilde{w}_j^{\mu} = (A_j^k / \beta_j) G_j^{\mu}$ . This relation is rewritten as

$$2i||w_k||^{2\alpha_k} \sum_{\substack{|p_{\sigma(j)}|+q \ge 0 \\ a > 1}} b_{j,p_{\sigma(j),q}}^{\mu} (w_{\sigma(j)})^{p_{\sigma(j)}} q(x+i|||w|||^{2\alpha})^{q-1} = \frac{A_j^k}{\beta_j} G_j^{\mu}.$$

By comparing the terms with  $||w_k||^{2\alpha_k}$ , the both sides must be zero. Since  $G_j^{\mu}$  does not vanish identically,  $A_j^k = 0$ . This is a contradiction. Thus there exists  $k = 1, \ldots, N$  such that  $k \neq \sigma(j)$  and  $A_j^k = 0$ . Take such k and apply  $\widetilde{w}_j^{\mu} = G_j^{\mu}$  to  $\Phi_* E_k = A_1^k \widetilde{E}_1 + \cdots + A_N^k \widetilde{E}_N$  to get the relation  $E_k G_j^{\mu} = 0$ . This is rewritten as

$$2i||w_k||^{2\alpha_k} \sum_{\substack{|p_{\sigma(j)}|+q \ge 0 \\ q > 1}} b_{j,p_{\sigma(j),q}}^{\mu} (w_{\sigma(j)})^{p_{\sigma(j)}} q(x+i|||w|||^{2\alpha})^{q-1} = 0.$$

This means that the coefficients  $b_{j,p_{\sigma(j)},q}^{\mu}$  vanish for  $|p_{\sigma(j)}| + q \ge 0, q \ge 1$ . Now we conclude that  $G_j^{\mu}$  has an expansion of the form (15).

Let us denote by  $\Phi_* E_{\sigma(1)} = A_1^{\sigma(1)} \widetilde{E}_1 + \dots + A_N^{\sigma(1)} \widetilde{E}_N$ . Apply  $\widetilde{w}_j^{\mu} = G_j^{\mu}$  with  $j \neq 1$  to get  $(A_j^{\sigma(1)}/\beta_j)G_j^{\mu} = 0$  and this means that  $A_j^{\sigma(1)} = 0$  for  $j \neq 1$ . By the same reason, we have  $A_j^{\sigma(k)} = 0$  for  $j \neq k$ , namely, we have  $\Phi_* E_{\sigma(j)} = A_j^{\sigma(j)} \widetilde{E}_j$ , which will be used in the next claim.

### Claim 3.2. F is expanded as

(16) 
$$F(x,w) = a_0 + a_1(x+i|||w|||^{2\alpha}).$$

First of all we apply  $\widetilde{x} = (F + \overline{F})/2$  to  $\Phi_* W_{\sigma(1)}^{\lambda} \in \widetilde{\mathcal{W}}_1, \dots, \Phi_* W_{\sigma(N)}^{\lambda} \in \widetilde{\mathcal{W}}_N$  to get

$$W^{1}_{\sigma(1)}F = 0, \dots, W^{m_{\sigma(1)}}_{\sigma(1)}F = 0,$$
  
 $\vdots$   
 $W^{1}_{\sigma(N)}F = 0, \dots, W^{m_{\sigma(N)}}_{\sigma(N)}F = 0.$ 

By the same argument as in Claim 3.1, F is a function of variable  $x + i|||w|||^{2\alpha}$ . Next applying  $\tilde{x} = (F + \overline{F})/2$  to  $\Phi_* E_{\sigma(j)} = A_j^{\sigma(j)} \tilde{E}_j$  and using the expansion (11), we get

(17) 
$$||w_{\sigma(j)}||^{2\alpha_{\sigma(j)}} \sum_{q>1} a_{p,q} q(x+i|||w|||^{2\alpha})^{q-1} = A_j^{\sigma(j)} ||G_j||^{2\beta_j}.$$

Since the right hand side does not contain the variable x, we obtain  $a_{p,q} = 0$  for  $q \ge 2$ . This leads to an expansion of the form (16).

Go back to the proof of lemma. The equation (17) is reduced to  $||w_{\sigma(j)}||^{2\alpha_{\sigma(j)}}a_1 = A_j^{\sigma(j)}||G_j||^{2\beta_j}$ . This means that  $G_j^{\mu}$  does not have a constant term, which implies the desired expansion of (14). On the boundary  $\partial H(m;(m);(\alpha))$ , we have

(18) 
$$\operatorname{Im} (a_0 + a_1(x+i|||w|||^{2\alpha})) = |||G|||^{2\beta}.$$

Since the right hand side does not have the constant term and the terms containing x, we obtain that  $a_0$  and  $a_1$  are real and also obtain that  $a_1 > 0$ . Since the mapping

$$(19) \qquad (\widetilde{z}, \widetilde{w}) \mapsto (\frac{\widetilde{z} - a_0}{a_1}, \frac{\widetilde{w}_1}{a_1^{1/2\beta_{\sigma(1)}}}, \dots, \frac{\widetilde{w}_{N-1}}{a_1^{1/2\beta_{\sigma(N-1)}}}, \frac{\widetilde{w}_N}{a_1^{1/2}})$$

is an automorphism of  $H(n;(n);(\beta))$ . It follows that F has an desired expansion (13). This automorphism does not change the form of (14).

### 4. The relations between exponents

Making use of the expansions of F and  $G_j^{\lambda}$ , we will get the relations between exponents  $\alpha_j$  and  $\beta_j$ . It follows from Lemma 3.1 that we have  $|||w|||^{2\alpha} = |||G|||^{2\beta}$  on the boundary of  $H(m;(m);(\alpha))$ . Take j with  $\sigma(j) = N$ . We restrict  $|||w|||^{2\alpha} = |||G||^{2\beta}$  to the variety  $w_1 = \cdots = w_{N-1} = 0$ . Then it is reduced to  $||w_N||^2 = ||G_j(w_N)||^{2\beta_j}$ . Substitute the expansion of  $G_j^{\lambda}(w_N)$  into this and compare the degree of  $w_N$ . Then we obtain  $\beta_j = 1$ , which implies that j = N. Hence we get  $\sigma(N) = N$ . Thus, from now on, we consider  $\sigma$  as a permutation of  $\{1, \ldots, N-1\}$ . Now we obtain the relations between the exponents  $\alpha$  and  $\beta$ .

**Lemma 4.1.** There exist  $M_1, \ldots, M_{N-1} \in \mathbb{N}$  such that  $\alpha_{\sigma(j)} = M_j \beta_j$  for  $j = 1, \ldots, N-1$ .

*Proof.* Once we get the relation between  $\alpha_{\sigma(1)}$  and  $\beta_1$ , the relations between  $\alpha_{\sigma(j)}$  and  $\beta_j$  for  $2 \leq j \leq N-1$  are obtained by the same argument. So, it suffices to show the case j=1. First we restrict the expansions (13) and (14) to the variety  $w_{\sigma(2)} = \cdots = w_{\sigma(N-1)} = w_N = 0$  and obtain the relation on the boundary  $\partial H(m; (m); (\alpha))$ :

(20) 
$$||w_{\sigma(1)}||^{2\alpha_{\sigma(1)}} = ||\sum_{|p_{\sigma(1)}| \ge 1} b_{1,p_{\sigma(1)}} (w_{\sigma(1)})^{p_{\sigma(1)}}||^{2\beta_1}.$$

Case 1:  $\alpha_{\sigma(1)} < \beta_1$ . Comparing the minimal total degree of  $w_{\sigma(1)}$  and  $\overline{w_{\sigma(1)}}$  in the both sides, we reach a contradiction.

Case 2:  $\alpha_{\sigma(1)} \geq \beta_1$ . Assume that there does not exist  $M_1 \in \mathbb{N}$  as in the lemma. Take  $M \in \mathbb{N}$  such that  $M\beta_1 < \alpha_{\sigma(1)} < (M+1)\beta_1$ . The terms of  $w_{\sigma(1)}$  and  $\overline{w_{\sigma(1)}}$ 

which have total degree less than or equal to  $2M\beta_1$  on the right hand side are zero. Hence the equation (20) becomes

$$||w_{\sigma(1)}||^{2\alpha_{\sigma(1)}} = ||\sum_{|p_{\sigma(1)}| \geq M+1} b_{1;p_{\sigma(1)}} (w_{\sigma(1)})^{p_{\sigma(1)}}||^{2\beta_1}.$$

Comparing the minimal total degree of  $w_{\sigma(1)}$  and  $\overline{w_{\sigma(1)}}$  in the both sides, we reach a contradiction. Hence we obtain  $\alpha_{\sigma(1)} = M_1\beta_1$  for some  $M_1 \in \mathbb{N}$ . This completes the proof.

### 5. Normalization of the mapping

Substituting  $\alpha_{\sigma(1)} = M_1 \beta_1$  to (20) and comparing the total degree of  $w_{\sigma(1)}$  and  $\overline{w_{\sigma(1)}}$ , we conclude that  $G_1^{\mu}$  is of the form:

$$G_1^{\mu}(w_{\sigma(1)}) = \sum_{|p_{\sigma(1)}|=M_1} b_{1;p_{\sigma(1)}}^{\mu} (w_{\sigma(1)})^{p_{\sigma(1)}}.$$

Analogously, we conclude that  $G_j^{\mu}$ ,  $j=1,\ldots,N$ , have the following expansions:

(21) 
$$G_{j}^{\mu}(w_{\sigma(j)}) = \sum_{|p_{\sigma(j)}|=M_{j}} b_{j,p_{\sigma(j)}}^{\mu}(w_{\sigma(j)})^{p_{\sigma(j)}}, \ j = 1,\dots, N-1, \ \mu = 1,\dots, n_{j},$$
$$G_{N}^{\mu}(w_{N}) = \sum_{|p_{N}|=1} b_{N,p_{N}}^{\mu}(w_{N})^{p_{N}}, \ \mu = 1,\dots, n_{N}.$$

We give a normalization of one block case by restricting  $\Phi = (F, G)$  to the variety  $w_{\sigma(2)} = \cdots = w_{\sigma(N-1)} = w_N = 0$ . This restriction does not influence to  $G_1$ . The reduced mapping is now

(22) 
$$(F, G_1) : \{ \operatorname{Im} z = ||w_{\sigma(1)}||^{2\alpha_{\sigma(1)}} \} \to \{ \operatorname{Im} \widetilde{z} = ||\widetilde{w}_1||^{2\beta_1} \}.$$

**Lemma 5.1.** Let  $\alpha_{\sigma(1)} = M_1\beta_1$ . Then we have the following:

(a) 
$$n_1 \ge \frac{(M_1 + m_{\sigma(1)} - 1)!}{M_1!(m_{\sigma(1)} - 1)!}$$
; and

(b)  $(F, G_1)$  is equivalent to  $(z, H_{M_1}(w_{\sigma(1)}), 0)$ .

*Proof.* We introduce two notations:

$$T_{M,m} = \frac{(M+m-1)!}{M!(m-1)!} = {M+m-1 \choose m-1}, \quad C_{M,p} = \sqrt{\frac{M!}{p^1!\dots p^m!}}$$

for  $M, m \in \mathbb{N}$  and  $p = (p^1, \dots, p^m) \in \mathbb{N}^m$ . Note that  $T_{M,m}$  is the number of solutions of  $x_1 + \dots + x_m = M$  for  $x_1, \dots, x_m \in \mathbb{Z}_{\geq 0}$ . Substituting (21) and  $a_{\sigma(1)} = M_1\beta_1$  into (20) and taking the  $\beta_1$ -th root, we get the equation:

(23) 
$$||w_{\sigma(1)}||^{2M_1} = ||\sum_{|p_{\sigma(1)}|=M_1} b_{1;p_{\sigma(1)}} (w_{\sigma(1)})^{p_{\sigma(1)}}||^2.$$

Since the coefficients of  $|(w_{\sigma(j)})^{p_{\sigma(j)}}|^2$  with  $|p_{\sigma(j)}| = M_1$  on the left hand side is  $C^2_{M_1,p_{\sigma(1)}}$  and the one on the right hand side is  $|b^1_{1;p_{\sigma(1)}}|^2 + \cdots + |b^{n_1}_{1;p_{\sigma(1)}}|^2$ , the vector  $(1/C_{M_1,p_{\sigma(1)}})b_{1;p_{\sigma(1)}} \in \mathbb{C}^{n_1}$  is a unit vector. On the other hand, picking up the terms  $(w_1)^{p_{\sigma(1)}}(\overline{w_1})^{\hat{p}_{\sigma(1)}}$  with  $p_{\sigma(1)} \neq \hat{p}_{\sigma(1)}$  from the both sides of (23), we have  $\langle b_{1;p_{\sigma(1)}}, b_{1;\hat{p}_{\sigma(1)}} \rangle = 0$  for  $p_{\sigma(1)} \neq \hat{p}_{\sigma(1)}$ . Thus the set  $\{(1/C_{M_1,p_{\sigma(1)}})b_{1;p_{\sigma(1)}}\}$  consisting of  $T_{M_1,m_{\sigma(1)}}$  vectors is an orthogonal systems in  $\mathbb{C}^{n_1}$ ; and hence,  $T_{M_1,m_{\sigma(1)}} \leq n_1$ . This is the inequality required in (a). For simplicity, we now denote by

 $\{U_1,\ldots,U_{T_{M_1,m_{\sigma(1)}}}\}$  this orthogonal system. Then one can find unit vectors  $D_1,\ldots,D_{n_1-T_{M_1,m_{\sigma(1)}}}\in\mathbb{C}^{n_1}$  in such a way that

$$U = \begin{pmatrix} U_1 \\ \vdots \\ U_{T_{M_1, m_{\sigma(1)}}} \\ D_1 \\ \vdots \\ D_{n_1 - T_{M_1, m_{\sigma(1)}}} \end{pmatrix}$$

is an  $n_1 \times n_1$  unitary matrix. Note that the transformation

$$H(n;(n);(\beta)) \ni (\widetilde{z},\widetilde{w}_1,\ldots,\widetilde{w}_N) \mapsto (\widetilde{z},\widetilde{w}_1\overline{U}^t,\widetilde{w}_2,\ldots,\widetilde{w}_N) \in H(n;(n);(\beta))$$

is an automorphism of  $H(n; (n); (\beta))$ . Applying U to  $G_1$  as follows, we obtain the second part of the lemma.

$$G_{1}\overline{U}^{t} = \left(\sum_{|p_{\sigma(1)}|=M_{1}} b_{1;p_{\sigma(1)}}^{1}(w_{\sigma(1)})^{p_{\sigma(1)}}, \dots, \sum_{|p_{\sigma(1)}|=M_{1}} b_{1;p_{\sigma(1)}}^{n_{1}}(w_{\sigma(1)})^{p_{\sigma(1)}}\right) \overline{U}^{t}$$

$$= \left(\dots, \frac{C_{M_{1},m_{\sigma(1)}}^{2}}{C_{M_{1},p_{\sigma(1)}}}(w_{\sigma(1)})^{p_{\sigma(1)}}, \dots, 0, \dots\right)$$

$$= \left(\dots, C_{M_{1},p_{\sigma(1)}}(w_{\sigma(1)})^{p_{\sigma(1)}}, \dots, 0, \dots\right).$$

The normalization of  $G_1$  in Lemma 5.1 is obtained by restricting the situation to  $w_{\sigma(2)} = \cdots = w_{\sigma(N-1)} = w_N = 0$ . By the same reason, restricting the situation to the variety

$$\left(\bigcap_{\substack{j=1,\dots,N-1\\ j\neq k}} \{w_{\sigma(j)} = 0\}\right) \cap \{w_N = 0\},\,$$

we can normalize  $G_k$  as

$$G_k(w_{\sigma(k)}) = (H_{M_k}(w_{\sigma(k)}), 0)$$

Restricting the mapping to  $w_{\sigma(1)} = \cdots = w_{\sigma(N-1)} = 0$  and tracing the same argument as above, we can normalize  $G_N^{\lambda}$  by using  $T_{1,m_N}$  and  $C_{1,p_N}$ . Then we conclude that  $G_N$  is equivalent to  $(w_N, 0)$ .

#### 6. PROOF OF THE MAIN THEOREM

Now we have proved that (F, G) is equivalent to

$$(24) (z, H_{M_1}(w_{\sigma(1)}), 0, \dots, H_{M_{N-1}}(w_{\sigma(N-1)}), 0, w_N, 0)$$

as a mapping between  $H(m;(m);(\alpha))$  and  $H(n;(n);(\beta))$ . We pull back this normalization to the mapping between  $E(m;(m);(\alpha))$  and  $E(n;(n);(\beta))$  via  $\Psi$  and  $\widetilde{\Psi}$ . By calculation, we obtain that  $(\mathcal{F},\mathcal{G})$  is equivalent to

$$(z, H_{M_1}(w_{\sigma(1)}), 0, \dots, H_{M_{N-1}}(w_{\sigma(N-1)}), 0, w_N, 0).$$

This finishes the proof of the Main Theorem.

#### 7. Comparison with the previous result

In this section, we compare our result with author's previous result in [Ha2]. Notation is the same as in the present paper.

**Main Theorem in [Ha2].** Let  $E(m;(m);(\alpha))$  and  $E(n;(n);(\beta))$  be generalized pseudoellipsoids with N blocks. Suppose that  $2 < m_j, 3 < n_j$  and that  $n - m < \min\{n_1, \ldots, n_N\}$ . Let  $(\mathcal{F}, \mathcal{G}_1, \ldots, \mathcal{G}_N) : E(m;(m);(\alpha)) \to E(n;(n);(\beta))$  be a proper holomorphic mapping that is holomorphic up to the boundary. Assume that non-zero columns of any block row in the Jacobian matrix of the unbounded representation of  $(\mathcal{F}, \mathcal{G})$  are linearly independent. Then we have the following:

- (1) There exists a permutation  $\sigma$  of  $\{1, ..., N\}$  such that  $\mathcal{G}_j|_{w_{\sigma(j)}=0}=0$  for every j.
- (2) If the permutation  $\sigma$  in (1) satisfies the inequality  $m_{\sigma(j)} \leq n_j < 2m_{\sigma(j)} 1$  for every j, then  $\alpha_{\sigma(j)} = \beta_j$  for all j and the proper holomorphic mapping  $(\mathcal{F}, \mathcal{G}_1, \ldots, \mathcal{G}_N)$  is equivalent to  $(\widetilde{\mathcal{F}}, \widetilde{\mathcal{G}}_1, \ldots, \widetilde{\mathcal{G}}_N)$  of the form

$$\widetilde{\mathcal{F}}(z, w_1, \dots, w_N) = z, \ \widetilde{\mathcal{G}}_j(z, w_1, \dots, w_N) = (w_{\sigma(j)}, 0), \ 1 \le j \le N.$$

In [Ha2], we assume that  $m_{\sigma(j)} \leq n_j < 2m_{\sigma(j)} - 1$  for every j. If we add this inequality to Main Theorem 1.1 as an assumption, then, combining with (b) in Main Theorem 1.1, we have an inequality:

(25) 
$$\frac{(M_j + m_{\sigma(j)} - 1)!}{M_j!(m_{\sigma(j)} - 1)!} \le n_j < 2m_{\sigma(j)} - 1.$$

Note that  $\frac{(M_j+m_{\sigma(j)}-1)!}{M_j!(m_{\sigma(j)}-1)!}$  is an increasing function with respect to  $M_j$  and inequality (25) does not hold for  $M_j=2$ . Hence inequality (25) implies that  $M_j=1$ . This implies  $\alpha_{\sigma(j)}=\beta_j$  in Main Theorem 1.1, which is the same conclusion as in Main Theorem in [Ha2]. Since the function defined by (2) satisfies  $H_1(z)=z$ , under the assumption on dimensions  $m_{\sigma(j)}\leq n_j<2m_{\sigma(j)}-1$  as in [Ha2], our mapping  $(\mathcal{F},\mathcal{G})$  is equivalent to the mapping

(26) 
$$(z, w) \mapsto (z, w_{\sigma(1)}, 0, \dots, w_{\sigma(N-1)}, 0, w_N, 0),$$

which is also the same conclusion in [Ha2]. As a result, we conclude the following.

**Theorem 7.1.** We add the assumption  $m_{\sigma(j)} \leq n_j < 2m_{\sigma(j)} - 1$  to Main Theorem 1.1. Then it implies Main Theorem in [Ha2].

Since  $M_j$  is invariant under the composition of automorphisms of  $E(m; (m); (\alpha))$  and  $E(n; (n); (\beta))$ , the mapping

$$(z,w) \mapsto (z, H_{M_1}(w_{\sigma(1)}), 0, \dots, H_{M_{N-1}}(w_{\sigma(N-1)}), 0, w_N, 0)$$

with  $M_j \neq 1$  is not equivalent to the mapping (26). Finally we would like to refer to the assumption on the Jacobian matrix in Main Theorem in [Ha2]. Let  $\mathcal{G}_j$  be a component of a proper holomorphic mapping as in Main Theorem 1.1. The assumption on the Jacobian matrix in Main Theorem in [Ha2] means that the non-zero columns of the Jacobian matrix of  $\mathcal{G}_j$  are linearly independent for any  $j = 1, \ldots, N-1$ . This condition is satisfied for  $\mathcal{G}_j(w_{\sigma(j)}) = (H_{M_j}(w_{\sigma(j)}), 0)$ .

#### References

- [D] J. D'Angelo, Several Complex Variables and the Geometry of Real Hypersurfaces, Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1993
- [Do] A. Dor, Proper holomorphic maps between balls in one co-dimension, Arkiv for Mathematik vol. 28, no. 1, (1990) 49–100
- [DP1] G. Dini, A. S. Primicerio, Propr holomorphic mappings between generalized pseudoellipsoids, Annali di matematica pura e applicata (IV) vol. CLVIII, (1991) 219–229
- [DP2] G. Dini, A. S. Primicerio, Localization principle of automorphisms on generalized pseudoellipsoids, J. Geom. Anal. vol. 7, no. 4, (1997) 575–584
- [ES] P. Ebenfelt, D. N. Son, Holomorphic mappings between pseudoellipsoids in different dimension, arXiv:1210.4434v1[math.CV]
- [GK] R. E. Green, S. G. Krantz, Characterizations of certain weakly pseudoconvex domains with non-compact automorphism groups, Lecture Notes in Math. 1268, Springer-Verlag, Berlin, Heidelberg, New York, London, Paris and Tokyo, 1987 121–157
- [Ha1] A. Hayashimoto, Application of the  $C^{\omega}$ -extendability theorem for proper holomorphic mappings to the classification of their mappings between two pseudoellipsoids, Math. Japonica, 44, no.2 (1996) 221–237
- [Ha2] A. Hayashimoto, A rigidity theorem for proper holomorphic mappings between generalized pseudoellipsoids, Tokyo J. Math. vol. 39, no. 2, (2016) 389–421
- [K1] A. Kodama, A characterization of certain domains with good boundary points in the sense of Greene-Krantz, Kodai Math. J. 12 (1989) 257–269
- [K2] A. Kodama, A characterization of certain domains with good boundary points in the sense of Greene-Krantz, II, Tohoku J. Math. 43 (1991) 9–25
- [K3] A. Kodama, A characterization of certain weakly pseudoconvex domains, Tohoku Math. J. 51 (1999), 55-64
- [K4] A. Kodama, On the holomorphic automorphism group of a generalized complex ellipsoid, Complex Var. Elliptic Equ. 59, no.9, (2014) 1342–1349
- [KKM] A. Kodama, S. G. Krantz, D. Ma, A characterization of generalized complex ellipsoids in  $\mathbb{C}^n$  and related results, Indiana Univ. Math. J. 41 (1992) 173–19
- [L] M. Landucci, On the proper holomorphic equivalence for a class of pseudoconvex domains, Trans. Amer. Math. Soc. 282, (1984) 807–811
- [MM] R. Monti, D. Morbidelli, D., Pseudohermitian invariants and classification of CR mappings in generalized ellipsoids,

ATSUSHI HAYASHIMOTO: 716 TOKUMA, NAGANO 381-8550, JAPAN E-mail address: atsushi@nagano-nct.ac.jp