# On multiple translative tiling in the plane Qi Yang

Abstract. This paper shows that in the plane, a multiple translative tile is a multiple lattice tile.

### 1. Introduction

Let K be a convex body with non-empty interior  $\operatorname{int}(K)$  and boundary  $\partial K$ , and let X be a discrete multiset in  $\mathbb{E}^d$ . We call K + X a translative tiling of  $\mathbb{E}^d$  and call K a translative tile if  $K + X = \mathbb{E}^d$  and any pair of translates  $\operatorname{int}(K) + \mathbf{x}_i$  are disjoint. In other words, K + X is both a packing and a covering in  $\mathbb{E}^d$ . Particularly, if X is a lattice in  $\mathbb{E}^d$ , we call K + X a lattice tiling of  $\mathbb{E}^d$  and call K a lattice tile.

Apparently, a translative tile is a convex polytope. It was shown by Minskowski [8] in 1897 that every translative tile must be centrally symmetric. In 1954, Venkov [11] proved that every translative tile must be a lattice tile. Later, a new proof for this beautiful result was independently discovered by McMullen [9].

Let X be a discrete multiset in  $\mathbb{E}^d$  and let k be a positive integer. We call K + X a k-fold translative tiling of  $\mathbb{E}^d$  and call K a translative k-tile if any point  $x \in \mathbb{E}^d$  belongs to at least k translates of K in K + X and every point  $x \in \mathbb{E}^d$  belongs to at most k translates of int(K)in int(K) + X. In other words, if K + X is both a k-fold packing and a k-fold covering in  $\mathbb{E}^d$ . Particularly, if X is a lattice in  $\mathbb{E}^d$ , we call K + X a k-fold lattice tiling of  $\mathbb{E}^d$  and call K a lattice k-tile. We call K a multiple translative (lattice) tile if K is a translative (lattice) k-tile for some positive integer k.

In 1936, multiple tiling was first investigated by FurtwÄd'ngler [2] as a generalization of MinkowskiâĂŹs conjecture on cube tiling. For more information, see [10], [5] and [14]. Similar to MinkowskiâĂŹs characterization, it was shown by Gravin, Robins and Shiryaev [4] that a translational k-tile must be a centrally symmetric polytope with centrally symmetric facets. As an analogy to the beautiful results of Venkov [11] and McMullen [9], it is natural to ask if a multiple translative tile is a multiple lattice tile.

In 2000, Kolountzakis [6] studied the structure of a multiple translative tiling by proving that, if D is a two dimensional convex domain which is not a parallelogram and D + X is a multiple tiling in  $\mathbb{E}^2$ , then X must be a finite union of 2-dimensional translated lattice. In 2013, Gravin, Kolountzakis, Robins and Shiryae [3] discovered a similar result in  $\mathbb{E}^3$ .

Let  $\tau(K)$  denote the smallest integer k such that K can form a k-fold translative tiling in  $\mathbb{E}^d$ , and let  $\tau^*(K)$  denote the smallest integer k such that K can form a k-fold lattice tiling in  $\mathbb{E}^d$ . For convenience, we define  $\tau(K) = \infty$  if K cannot form translative tiling of any multiplicity. Clearly, for every centrally symmetric convex polytope we have  $\tau(K) \leq \tau^*(K)$ .

At the end of [3], several open problems were proposed. One of them is: Prove or disprove that if any polytope k tile  $\mathbb{E}^d$  by translations, then it is also m tile  $\mathbb{E}^d$  by lattice, for a possibly different m. i.e. Prove or disprove that  $\tau(K) < \infty$  imply that  $\tau^*(K) < \infty$ . This paper confirms the two-dimensional case of this problem. We acknowledge that this result is independently discovered by Liu [7].

In 2017, Yang and Zong [12] studied the multiple translative tiling with given multiplicity. Later, Zong [15] characterized all the two-dimensional five-fold lattice tile. Afterwards, Yang and Zong [13] showed that in the plane  $\tau(P) = 5$  imply that  $\tau^*(P) = 5$  and thus characterized all the two-dimensional translative tile. There is very little known in relations between  $\tau(P)$  and  $\tau^*(P)$ , even in the plane. The main result of this paper is the following theorem :

Theorem 1. In the plane, a multiple translative tile is a multiple lattice tile.

#### 2. Preparation

To prove theorem 1 we need the following known results.

#### The structure of a multiple translative tiling in the plane

**Theorem 2 (Mihail N.Kolountzakis [6]).** Suppose that P + X is a multiple translative tiling, where X is a multiset in the plane. If P is not a parallelogram, then X is a finite union of two-dimensional lattices.

By slightly modifying the method used in U.Bolle [1], we can get more information about the structure of a multiple translative tiling. Without specific statement, assume that P+X is a k-fold translative tiling in the plane for some positive integer k.

**Definition 1 (U.Bolle [1]).** Let  $L(\mathbf{e})$  be the straight line containing  $\mathbf{e}$ , where  $\mathbf{e}$  is an edge of  $P + \mathbf{x}$  and  $\mathbf{x} \in X$ .

A point  $\mathbf{p} \in L(\mathbf{e})$  is called a normal point if there is an  $\varepsilon > 0$  with

$$(B_{\varepsilon}(\mathbf{p}) \setminus L(\mathbf{e})) \cap (\bigcup_{\mathbf{x} \in X} (\mathbf{x} + \partial P)) = \emptyset,$$

where  $B_{\varepsilon}(\mathbf{p})$  denotes the open circular disc with center  $\mathbf{p}$  and radius  $\varepsilon$ . Since X is a discrete multiset in  $\mathbb{E}^2$ , one can deduce that almost all points of  $L(\mathbf{e})$  are normal and the non-normal points form a discrete set.

For normal points we define two functions  $n_i (i = 1, 2)$  by

$$N_i(\mathbf{p}) = \{ \mathbf{x} \in X \mid \mathbf{x} + P \subseteq cl\{H_i\} \text{ and } \mathbf{p} \in \mathbf{x} + \partial P \}$$
(1)

$$n_i(\mathbf{p}) = |N_i(\mathbf{p})| \tag{2}$$

where  $H_i(i = 1, 2)$  are the two half-planes defined by  $L(\mathbf{e})$  and  $d\{H_i\}$  are the closure of  $\{H_i\}$ .

In fact,  $n_1(\mathbf{p}) = n_2(\mathbf{p})$  for all normal points  $\mathbf{p}$ . For, if  $B_i = B_{\varepsilon}(\mathbf{p}) \cap H_i$ , each point in  $B_i$  is cover exactly k times, and if we cross L(e) in  $\mathbf{p}$  from  $B_1$  to  $B_2$ , then we leave  $n_1$  translates of Pand enter  $n_2$  translates of P.

Assume that P is a centrally symmetric polygon with center **o** and 2m edges for some positive integer  $m \ge 4$ . Let  $\mathbf{v}_1, \mathbf{v}_2, ..., \mathbf{v}_{2m}$  be the 2m vertices of P enumerated in the counterclockwise order. Define  $\mathbf{e}_i = \mathbf{v}_{i+1} - \mathbf{v}_i$   $(1 \le i \le 2m, \mathbf{v}_1 = \mathbf{v}_{2m+1}), \mathbf{e}_i^* = \mathbf{v}_{i+m} - \mathbf{v}_{i+1}, 1 \le i \le m$ .

**Lemma 3.** Suppose that  $\mathbf{x} \in X$ . For each  $i \ (1 \le i \le m)$ , either  $\mathbf{x} - \mathbf{e}_i$  or  $\mathbf{x} - \mathbf{e}_i^*$  belongs to X.

**Proof.** Assume that  $\mathbf{x} - \mathbf{e}_i^* \notin X$  and  $\mathbf{x} - \mathbf{e}_i \notin X$ . Let *L* be the line determined by  $\mathbf{e}_i + \mathbf{x}$ . Since the value of  $n_1$  of the normal points on the line *L* can only change at endpoints of translates of  $\mathbf{e}_i$ . Let *R* be the endpoint  $\mathbf{v}_i + \mathbf{x}$  of  $\mathbf{e}_i + \mathbf{x}$ , let  $Q_1$  and  $Q_2$  be normal points of *L* separated by *R* such that there are only normal points between *R* and  $Q_i$ .

Since  $\mathbf{x} - \mathbf{e}_i \notin X$ , then we have  $n_1(Q_1) > n_1(Q_2)$  and so  $n_2(Q_1) \neq n_2(Q_2)$ . It is easy to see that R is an endpoint of a translates of  $\mathbf{e}_{i+m}$ , as shown by Figure 1. By assumption, the other endpoint  $R^*$  of  $\mathbf{e}_i + \mathbf{x}$  cannot belong to the same translate of  $\mathbf{e}_{i+m}$ , then we have  $n_2(Q_1) < n_2(Q_2)$ , a contradiction. As a conclusion, we've proved the lemma 3.



**Theorem 4 (Bolle[1]).** A convex polygon is a k-fold lattice tile for a lattice  $\Lambda$  and some positive integer k if and only if the following conditions are satisfied:

**1.** It is centrally symmetric.

2. When it is centered at the origin, in the relative interior of each edge g there is a point of  $\frac{1}{2}\Lambda$ .

**3.** If the midpoint of **g** is not in  $\frac{1}{2}\Lambda$  then **g** is a lattice vector of  $\Lambda$ .

## 3. Proof of Theorem 1

By Theorem 2, we can easily deduce that X can be represented as

$$X = \bigcup_{j=1}^{l} \bigcup_{k=1}^{l_j} (\Lambda_j + \mathbf{x}_j^k) (\mathbf{x}_j^k \in \mathbb{E}^2)$$
(3)

where  $\Lambda_j \cap \Lambda_{j'}$  is at most one-dimensional sublattice, for any  $1 \leq j \neq j' \leq l$ .

**Lemma 5.** For each  $1 \leq j \leq l$  there exists some positive integer  $\beta_j$  such that either  $\mathbf{e}_i$  or  $\mathbf{e}_i^*$ belongs to  $\frac{1}{\beta_i} \Lambda_j$  for all i = 1, 2, ..., m.

**Proof.** Without loss of generality, we are going to show that there exists some positive integer  $\beta_1$ such that either  $\mathbf{e}_i$  or  $\mathbf{e}_i^*$  belongs to  $\frac{1}{\beta_1}\Lambda_1$  for all i = 1, 2, ..., m.

First, the following will be shown:

(i) For each  $1 \leq i \leq m$ ,

 $p_i \mathbf{e}_i + q_i \mathbf{e}_i^* \in \Lambda_1$ , where  $p_i, q_i (1 \le i \le m)$  are some non-negative integers and can't be zero at the same time.

Given  $\Lambda_1 + \mathbf{x}_1^k$ , the lattice  $\Lambda_1 + \mathbf{x}_1^k$  can be divided into two parts A and B,

$$A = \{ \mathbf{x} \in \Lambda_1 + \mathbf{x}_1^k | \mathbf{x} - \mathbf{e}_i \in X \}$$
$$B = \{ \mathbf{x} \in \Lambda_1 + \mathbf{x}_1^k | \mathbf{x} - \mathbf{e}_i^* \in X \}$$

and obviously,  $A \cup B = \Lambda_1 + \mathbf{x}_1^k$  by lemma 3.

Since  $\Lambda_1 \cap \Lambda_j$   $(j \neq 1)$  is at most one-dimensional sublattice, so  $((A - \mathbf{e}_1) \cap \Lambda_j) \cup ((B - \mathbf{e}_1^*) \cap \Lambda_j)$ for  $j \neq 1$  is at most one-dimensional sublattice. Therefore there must be a lattice  $\Lambda_1 + \mathbf{x}_1^{\psi(k,1)} (1 \leq 1)$  $\psi(k,1) \leq l_1$ ) such that

$$(A - \mathbf{e}_1) \cap (\Lambda_1 + \mathbf{x}_1^{\psi(k,1)}) \text{ or } (B - \mathbf{e}_1^*) \cap (\Lambda_1 + \mathbf{x}_1^{\psi(k,1)})$$

contains infinite elements and is two-dimensional. So we have either  $\mathbf{x}_1^k - \mathbf{e}_i = \mathbf{x}_1^{\psi(k,1)}$  or  $\mathbf{x}_1^k - \mathbf{e}_i^* = \mathbf{x}_1^{\psi(k,1)}$ . Then we can define a morphism  $\psi_1$  from  $\{1, 2, ..., l_1\}$  to itself:

$$\psi_1(k) = \psi(k, 1)$$

such that  $\mathbf{x}_1^k - \mathbf{e}_i = \mathbf{x}_1^{\psi_1(k)}$  or  $\mathbf{x}_1^k - \mathbf{e}_i^* = \mathbf{x}_1^{\psi_1(k)}$ . In this way we get a sequence  $\{a_i\}(a_i \in \{1, ..., l_1\})$  with

$$a_{i+1} = \psi_1(a_i)$$

Clearly, there exist two elements  $a_{n_1}, a_{n_2}$  of this sequence that  $a_{n_1} = a_{n_2}$ , which means there exist non-negative integers  $p_i, q_i$  such that

$$-p_i \mathbf{e}_i - q_i \mathbf{e}_i^* \in \Lambda_1 (1 \le i \le m)$$

where  $p_i, q_i$  are not zero at same time.

Suppose that  $\mathbf{u}, \mathbf{v}$  are the basis vectors of  $\Lambda_1$ , then we have that

$$p_i \mathbf{e}_i + q_i \mathbf{e}_i^* = a_i \mathbf{u} + b_i \mathbf{v} \tag{4}$$

where  $a_i, b_i \in \mathbb{Z}$ .

It's easy to see that

$$\mathbf{e}_1^* = \sum_{k=2}^m \mathbf{e}_k \tag{5}$$

$$\mathbf{e}_{i}^{*} = \sum_{k=1}^{i-1} -(\mathbf{e}_{k}) + \sum_{k=i+1}^{m} \mathbf{e}_{k} (2 \le i \le m)$$
(6)

So

$$p_1 \mathbf{e}_1 + q_i \sum_{k=2}^m \mathbf{e}_k = a_1 \mathbf{u} + b_1 \mathbf{v}$$
(7)

$$q_{i}\sum_{k=1}^{i-1} -(\mathbf{e}_{k}) + p_{i}\mathbf{e}_{i} + q_{i}\sum_{k=i+1}^{m}\mathbf{e}_{k} = a_{i}\mathbf{u} + b_{i}\mathbf{v}(2 \le i \le m)$$
(8)

Define  $I = \{1 \leq i \leq m : q_i = 0 \text{ and so } \mathbf{e}_i \in \frac{1}{p_i}\Lambda_1\}$  and  $I' = \{1, 2, ...m\} \setminus I$ , and denote the indicator function of I' by  $\delta_{I'}$ . Then the above equation can be represented as follows:

$$q_{i}\sum_{k=1}^{i-1} -(\mathbf{e}_{k})\delta_{I'}(k) + \delta_{I'}(i)p_{i}\mathbf{e}_{i} + q_{i}\sum_{k=i+1}^{m}\delta_{I'}(k)\mathbf{e}_{k} = a'_{i}\mathbf{u} + b'_{i}\mathbf{v}$$
(9)

where  $a'_i, b'_i$  are rational numbers.

For convenience, those edges in  $\{\mathbf{e}_i\}_{i \in I'}$  can be re-enumerated in original order, and denote by  $\mathbf{e}'_i (1 \leq j \leq l')$ , where l' = |I'|. Then we have

$$\begin{bmatrix} p_1 & q_1 & \dots & q_1 \\ -q_2 & p_2 & \dots & q_2 \\ \dots & \dots & \dots & \dots \\ -q_{l'} & -q_{l'} & \dots & p_{l'} \end{bmatrix} \begin{bmatrix} \mathbf{e}'_1 \\ \mathbf{e}'_2 \\ \dots \\ \mathbf{e}'_{l'} \end{bmatrix} = B \begin{bmatrix} u \\ v \end{bmatrix}$$

where all entries of B are rational numbers.

By a series of linear matrix transformation, we get

$$\begin{bmatrix} p'_1 & 1 & \dots & 1\\ -1 & p'_2 & \dots & 1\\ \dots & \dots & \dots & \dots\\ -1 & -1 & \dots & p'_{l'} \end{bmatrix} \begin{bmatrix} \mathbf{e}'_1\\ \mathbf{e}'_2\\ \dots\\ \mathbf{e}'_{l'} \end{bmatrix} = B' \begin{bmatrix} u\\ v \end{bmatrix}$$

where  $p'_i(1 \le i \le l')$  are non-negative rational numbers and all entries of B' are rational numbers. Define

$$A(p'_1, ..., p'_{l'}) = \begin{bmatrix} p'_1 & 1 & .... & 1\\ -1 & p'_2 & .... & 1\\ ... & ... & ...\\ -1 & -1 & .... & p'_{l'} \end{bmatrix}$$
(10)

Next, we are going to prove that if  $p'_i \ge 0$  for each  $1 \le i \le l'$  and some  $p'_j > 0$ , then  $A(p'_1, ..., p'_{l'})$  is invertible. We are going to prove this assertion by induction on l'.

It is obvious that the assertion is true while l' = 1. Assume that the assertion holds for  $l' \leq n-1$ .

Next, we are going to prove this assertion holds when l' = n. The determinant of  $A(p'_1, ..., p'_{l'})$  can be seen as a function of variables  $p'_1, ..., p'_{l'}$ . It is easy to calculate that the partial derivative of  $A(p'_1, ..., p'_{l'})$  respect to  $p'_i(1 \le i \le l')$  is  $A(p'_1, ..., p'_{i-1}, p'_{i+1}, ..., p'_{l'})$ , by the assumption, which is greater than 0.

Clearly, A(0, 0, ...0) = 0 when l' is odd, otherwise A(0, 0, ...0) = 1. As a conclusion, we've proved the assertion.

When  $p'_i = 0$  for all  $1 \le i \le l'$ , then  $q_i \mathbf{e}_i^* \in \Lambda_1$  for  $i \in I'$  and  $\mathbf{e}_i \in \Lambda_1$  for each  $i \in I$ .

When  $p'_i \neq 0$  for some  $1 \leq i \leq l'$ , then  $A(p'_1, ..., p'_{l'})$  is invertible, so there exists an invertible matrix  $A(p'_1, ..., p'_{l'})^{-1}$  such that

$$\begin{bmatrix} \mathbf{e}_1' \\ \mathbf{e}_2' \\ \dots \\ \mathbf{e}_l' \end{bmatrix} = A(p_1', \dots, p_{l'}')^{-1} B' \begin{bmatrix} \mathbf{u} \\ \mathbf{v} \end{bmatrix}$$

Since all entries of  $A(p'_1, ..., p'_{l'})^{-1}$  and B' are rational numbers, there exists a positive integer  $\beta_1$  such that all entries of  $\beta_1 A(p'_1, ..., p'_{l'})^{-1}B'$  are integers, i.e.  $\mathbf{e}_i \in \frac{1}{\beta_1}\Lambda_1$  for each  $i \in I'$ .

As a conclusion, we've proved that there exist  $\beta_1$  such that for all  $1 \leq i \leq m$ , either  $\mathbf{e}_i$  or  $\mathbf{e}_i^*$  belongs to  $\frac{1}{\beta_1}\Lambda_1$ . Similarly, we can prove the case of any other lattice  $\Lambda_j(1 < j \leq l)$ , lemma 5 is proved.

By lemma 4 and lemma 5, in order to prove Theorem 1, it is suffice to show that for some  $\Lambda_j$ , there is a point of  $\frac{1}{2\beta_i}\Lambda_j$  in the relative interior of  $\mathbf{e}_i$  for each  $i(1 \le i \le m)$ .

**Lemma 6.** ii) With above notations, for some  $j(1 \le j \le l)$ , there exists a point of  $\frac{1}{2\beta_j}\Lambda_j$  in the relative interior of  $\mathbf{e}_i$  for each  $1 \le i \le m$ , where  $\beta_j$  is some positive integer.

**Proof.** To prove lemma 6, it is sufficient to discuss the following two cases:

**Case 1.** For some  $\Lambda_j$ , there exists some  $i(1 \leq i \leq m)$  such that  $p\mathbf{e}_i$  and  $q\mathbf{e}_i^* \in \Lambda_j$ , where  $p, q \in \mathbb{Q} \setminus \{0\}$ .

By lemma 5, it is easy to deduce that for all  $i(1 \le i \le m)$ ,  $q_i \mathbf{e}_i^* \in \Lambda_i$  for some positive  $q_i \in \mathbb{Q}$ , so there exists some positive integer  $\beta_j$  such that  $\mathbf{e}_i^* \in \frac{1}{\beta_j} \Lambda_j$  for all  $\mathbf{e}_i^*$ , the middle point of every edge  $\mathbf{e}_i$  of P belongs to  $\frac{1}{2\beta_i} \Lambda_j$ , lemma 6 holds.

**Case 2.** For each  $\Lambda_j (1 \le j \le l)$ ,  $p\mathbf{e}_i$  and  $q\mathbf{e}_i^*$  can not belong to  $\Lambda_j$  at the same time for each  $1 \le i \le m$ , where p, q are arbitrary non-zero rational numbers.

In this case, we are going to prove that lemma 6 holds for  $\Lambda_1$  (The argument for other lattice  $\Lambda_j$  is similar). If  $\mathbf{e}_i^*$  belongs to  $\frac{1}{\beta}\Lambda_1$  for some positive integer  $\beta$ , then the midpoint of  $\mathbf{e}_i$  belongs to the lattice  $\frac{1}{2\beta}\Lambda_1$ . Otherwise  $\mathbf{e}_i \in \frac{1}{\beta}\Lambda_1$  for some positive integer  $\beta$ , then it is suffice to show that there exist  $\mathbf{g} \in \frac{1}{\beta}\Lambda_1$  such that  $\mathbf{g} - \mathbf{e}_i^* = \lambda \mathbf{e}_i$  for some  $\lambda \in \mathbb{R}$ .

Assume that  $\mathbf{e}_1^* \notin \frac{1}{\beta} \Lambda_1$  for any positive integer  $\beta$ . Then we have  $\mathbf{e}_1 \in \frac{1}{\beta} \Lambda_1$  for some positive integer  $\beta$ , which means that  $\mu \mathbf{e}_1 \in \Lambda_1$  for some positive rational number  $\mu$ . Define  $\Omega = \{\Lambda_j + \mathbf{x}_j^k, 1 \leq j \leq l, 1 \leq k \leq l_j\}$ . For convenience, we may assume that  $\mu \mathbf{e}_1$  and  $\mathbf{u}$  are the basis vectors of lattice  $\Lambda_1$ . Denote the line containing  $\mathbf{e}_1 + \mathbf{x}_1^1 + n\mathbf{u}(n \in \mathbb{Z})$  by  $L(\mathbf{x}_1^1, n)$ .

Let  $\mathbf{p}_j^{1,n}$  be a normal point in the relative interior of  $\mathbf{e}_1 + \mathbf{x}_1^1 + j\mu\mathbf{e}_1 + n\mathbf{u}$  for each  $j \in \mathbb{Z}$ . Since  $n_1(\mathbf{p}_j^{1,n}) = n_2(\mathbf{p}_j^{1,n}) > 0$ , then we define

$$\Gamma_{1,n} = \{\mathbf{y}_j^{1,n} : \mathbf{y}_j^{1,n} \in N_2(\mathbf{p}_j^{1,n})\}_{j \in \mathbb{Z}}$$

Apparently, we can find an infinite subset  $\Gamma'_{1,n}$  of  $\Gamma_{1,n}$  which its elements are contained in the same translated lattice in  $\Omega$ , denote the corresponding lattice by  $\Lambda(\Gamma_{1,n})$ .

Define  $W_1 = \{n \in \mathbb{N} : p\mathbf{e}_1 \notin \Lambda(\Gamma_{1,n}) \text{ for any } p \in \mathbb{Q}\}$ . If  $|W_1| < \infty$ , then  $|\mathbb{N} \setminus W_1| = \infty$  and so we can find an infinite subset  $W'_1$  of  $\mathbb{N} \setminus W_1$  that the corresponding translated lattice  $\Lambda(\Gamma_{1,n})$  for every  $n \in W'_1$  is identical, denote this lattice by  $\Lambda(1)$ . For each  $n \in W'_1$ , let  $\mathbf{p}_j^{2,n}$  be a normal point in the relative interior of the edge  $\mathbf{e}_1 + \mathbf{y}_j^{1,n}$  for  $\mathbf{y}_j^{1,n} \in \Gamma'_{1,n}$ . Since  $n_1(\mathbf{p}_j^{2,n}) = n_2(\mathbf{p}_j^{2,n}) > 0$ , by analogy, define  $\Gamma_{2,n}$  as follows:

$$\Gamma_{2,n} = \{ \mathbf{y}_j^{2,n} : \mathbf{y}_j^{2,n} \in N_2(\mathbf{p}_j^{2,n}) \text{ for } j \text{ such that } \mathbf{y}_j^{1,n} \in \Gamma_{1,n}' \}$$
(11)

then we can find an infinite subset  $\Gamma'_{2,n}$  of  $\Gamma_{2,n}$  such that its elements are contained in the same translated lattice in  $\Omega$ , denote the corresponding lattice by  $\Lambda(\Gamma_{2,n})$ .



$$\Gamma'_{2,n} = \{ \mathbf{y}_j^{2,n} : \mathbf{y}_{j,n} \in \Lambda(\Gamma_{2,n}) \}$$
(12)

#### Figure 2

Similarly, we can define the set  $W_2 = \{n \in W'_1 : p\mathbf{e}_1 \notin \Lambda(\Gamma_{2,n}) \text{ for any } p \in \mathbb{Q}\}$ . If  $|W_2| < \infty$ , then we can define an infinite subset  $W'_2$  of  $W'_1 \setminus W_2$ . Next, we are going to use induction to define the notations appeared in the following argument.

Suppose that we have  $|W_k| < \infty (k \ge 2)$ , and  $|W'_{k-1}| = \infty$ . Then there is an infinite subset  $W'_k$  of  $W'_{k-1} \setminus W_k$  that the corresponding translated lattice  $\Lambda(\Gamma_{k,n})$  for every  $n \in W'_k$  is identical, denote this lattice by  $\Lambda(k)$ .

Let  $\mathbf{p}_{j}^{k+1,n}$  be a normal point in the relative interior of the edge  $\mathbf{e}_{1} + \mathbf{y}_{j}^{k,n}$  for  $\mathbf{y}_{j}^{k,n} \in \Gamma_{k,n}'$  and  $n \in W_{k}'$ .

Since 
$$n_1(\mathbf{p}_j^{k+1,n}) = n_2(\mathbf{p}_j^{k+1,n}) > 0$$
, define  $\Gamma_{k+1,n}$  as follows:  

$$\Gamma_{k+1,n} = \{\mathbf{y}_j^{k+1,n} : \mathbf{y}_j^{k+1,n} \in N_2(\mathbf{p}_j^{k+1,n}) \text{ for } j \text{ such that } \mathbf{y}_j^{k,n} \in \Gamma'_{k,n}\}$$
(13)

then we can find an infinite subset  $\Gamma'_{k+1,n}$  of  $\Gamma_{k+1,n}$  which its elements are contained in the same translated lattice in  $\Omega$ , denote the corresponding lattice denote by  $\Lambda(\Gamma_{k+1,n})$ .

$$\Gamma'_{k+1,n} = \{\mathbf{y}_j^{k+1,n} : \mathbf{y}_j^{k+1,n} \in \Lambda(\Gamma_{k+1,n}) \cap \Gamma'_{k+1,n}\}$$

and define  $W_{k+1} = \{ n \in W'_k : p\mathbf{e}_1 \notin \Lambda(k+1, n) \text{ for any } p \in \mathbb{Q} \}.$ 

Since  $|\Omega| < \infty$ , there are only two possible cases: (a)  $|W_{k^*}| = \infty$  for some positive integer  $k^*(k^* \ge 1)$ ; (b)  $\Lambda(k') = \Lambda(k'')$  for two different positive integers k', k''.

(a) If  $|W_{k^*}| = \infty$  for some positive integers  $k^*(k^* \ge 1)$ , then there exist two elements  $n_1, n_2$  of  $W_{k^*}$  such that  $\Lambda(k^*, n_1) = \Lambda(k^*, n_2)$ .

By the definition of  $\Lambda(k^*, n_1)$ ,  $\mu' * \mathbf{e}_1 \in \Lambda(k^*, n_1)$  for some  $\mu' \in \mathbb{R}$ , and by the definition of  $W_{k^*}$ ,  $\mu'$  is an irrational number. Again, by the definition of  $W_{k^*}$ ,  $q * \mathbf{e}_1^* \in \Lambda(k^*, n_1)$  for some positive rational number q.

By the definition of  $\Lambda(k^*, n_1)$ , there exist two elements  $\mathbf{y}_{j'}^{k^*, n_1} \in \Lambda(k^*, n_1)$  and  $\mathbf{y}_{j''}^{k^*, n_2} \in \Lambda(k^*, n_1)$  such that:

$$\mathbf{y}_{j'}^{k^*,n_1} = \mathbf{x}_1^1 + n_1 \mathbf{u} - k^* \mathbf{e}_1^* + \lambda_3 \mathbf{e}_1$$
(14)

$$\mathbf{y}_{j''}^{k^*,n_2} = \mathbf{x}_1^1 + n_2 \mathbf{u} - k^* \mathbf{e}_1^* + \lambda_4 \mathbf{e}_1$$
(15)

where  $\lambda_3, \lambda_4 \in \mathbb{R}$  and some  $j', j'' \in \mathbb{Z}$ .

Combined (14)-(15),

$$\mathbf{y}_{j'}^{k^*,n_1} - \mathbf{y}_{j''}^{k^*,n_2} = (n_1 - n_2)\mathbf{u} + (\lambda_3 - \lambda_4)\mathbf{e}_1$$
(16)

Since  $\mu' \mathbf{e}_1$  and  $q \mathbf{e}_1^*$  are linearly independent, then we have

$$\mathbf{y}_{j'}^{k^*,n_1} - \mathbf{y}_{j''}^{k^*,n_2} = a\mu'\mathbf{e}_1 + bq\mathbf{e}_1^*$$
(17)

where  $a, b \in \mathbb{Q}$  and  $bq \neq 0$  (since  $n_1 \neq n_2$ ).

Since  $\mathbf{u}$  and  $\mathbf{e}_1$  are linearly independent, then we have

$$\mathbf{e}_1^* = \lambda_5 \mathbf{u} + \lambda_6 \mathbf{e}_1 \tag{18}$$

When  $\lambda_5 \in \mathbb{Q}$ , then there is a point **g** in the lattice  $\frac{1}{\beta}\Lambda_1$  such that

$$\mathbf{g} - \mathbf{e}_1^* = \lambda_6 \mathbf{e}_1$$

which means that for some positive integer  $\beta$ , there is a lattice point of  $\frac{1}{\beta}\Lambda_1$  in the relative interior of  $\mathbf{e}_1$ .

When  $\lambda_5 \notin \mathbb{Q}$ , combined (16),(17) and (18), we have

$$\mathbf{y}_{j'}^{k^*,n_1} - \mathbf{y}_{j''}^{k^*,n_2} = a\mu'\mathbf{e}_1 + bq(\lambda_5\mathbf{u} + \lambda_6\mathbf{e}_1)$$
(19)

and

$$(n_1 - n_2)\mathbf{u} + (\lambda_3 - \lambda_4)\mathbf{e}_1 = a\mu'\mathbf{e}_1 + bq(\lambda_5\mathbf{u} + \lambda_6\mathbf{e}_1)$$
(20)

$$(n_1 - n_2 - bq\lambda_5)\mathbf{u} = (a\mu' + bq\lambda_6 + \lambda_4 - \lambda_3)\mathbf{e}_1$$
(21)

Since  $\lambda_5 \notin \mathbb{Q}$ , the left side of the equation (22) is not equal to zero. But **u** and **e**<sub>1</sub> is linearly independent over  $\mathbb{R}$ , then we get a contradiction.

(b)  $\Lambda(k') = \Lambda(k'')$ . Suppose that k' < k''. By the definition of W'(k') and W'(k''), we have that  $W'(k'') \subset W'(k')$ . Let  $n_1 \in W'(k'')$ , let  $\mathbf{y}_{j'}^{k',n_1} \in \Lambda(k'')$  and  $\mathbf{y}_{j''}^{k'',n_1} \in \Lambda(k'')$  for some positive integers j', j'',

$$\mathbf{y}_{j'}^{k',n_1} = \mathbf{x}_1^1 + n_1 \mathbf{u} - k' \mathbf{e}_1^* + \lambda_3 \mathbf{e}_1$$
(22)

$$\mathbf{y}_{j''}^{k'',n_1} = \mathbf{x}_1^1 + n_1 \mathbf{u} - k'' \mathbf{e}_1^* + \lambda_4 \mathbf{e}_1$$
(23)

for some real numbers  $\lambda_3, \lambda_4$ .

Then we have

$$\mathbf{y}_{j''}^{k'',n_1} - \mathbf{y}_{j'}^{k',n_1} = (k'' - k')\mathbf{e}_1^* + (\lambda_3 - \lambda_4)\mathbf{e}_1.$$
(24)

Let  $n_2 \in W'(k')$  and  $n_1 \neq n_2$ , and let  $\mathbf{y}_{j*}^{k',n_2} \in \Lambda(k')$ , then we have

$$\mathbf{y}_{j*}^{k',n_2} = \mathbf{x}_1^1 + n_2 \mathbf{u} - k' \mathbf{e}_1^* + \lambda_5 \mathbf{e}_1$$
(25)

$$\mathbf{y}_{j*}^{k',n_2} - \mathbf{y}_{j'}^{k',n_1} = (n_2 - n_1)\mathbf{u} + (\lambda_5 - \lambda_3)\mathbf{e}_1$$
(26)

Suppose that  $\frac{1}{h}(\mathbf{y}_{j*}^{k',n_2} - \mathbf{y}_{j'}^{k',n_1}) \in \Lambda(k')$  for some positive integer h, by the definition of  $\Lambda(k')$ ,  $\mu \mathbf{e}_1 \in \Lambda(k')$  for some real number  $\mu$ . Since  $\Lambda(k') = \Lambda(k'')$ ,  $\mathbf{y}_{j''}^{k'',n_1} - \mathbf{y}_{j'}^{k',n_1}$  can be represented as the linear combination of  $\frac{1}{h}(\mathbf{y}_{j*}^{k',n_2} - \mathbf{y}_{j'}^{k',n_1})$  and  $\mu \mathbf{e}_1$ :

$$\mathbf{y}_{j''}^{k'',n_1} - \mathbf{y}_{j'}^{k',n_1} = \frac{z_1}{h} (\mathbf{y}_{j*}^{k',n_2} - \mathbf{y}_{j'}^{k',n_1}) + z_2 \mu \mathbf{e}_1$$
(27)

where  $z_1, z_2 \in \mathbb{Q}$ .

Combined equations (24),(26) and (27), we have

$$(k'' - k')\mathbf{e}_1^* + (\lambda_3 - \lambda_4)\mathbf{e}_1 = z_1 * ((n_2 - n_1)\mathbf{u} + \frac{1}{h}(\lambda_5 - \lambda_3)\mathbf{e}_1) + z_2\mu\mathbf{e}_1$$
(28)

Simplify the equation (29), we have

$$\mathbf{e}_1^* = q_1 \mathbf{u} + \mu' \mathbf{e}_1$$

for some rational number  $q_1$  and some real number  $\mu'$ .

As a conclusion, we've proved that there is a positive integer  $\beta$  such that there is a lattice point **g** in the lattice  $\frac{1}{\beta}\Lambda_1$  such that

$$\mathbf{g} - \mathbf{e}_1^* = \lambda \mathbf{e}_1$$

for some real number  $\lambda$ .

Similarly, for each  $i(1 \le i \le m)$ , we can prove that there is a positive integer  $\beta_1$  such that there is a lattice point g in the lattice  $\frac{1}{\beta_1}\Lambda_1$  that

$$g - \mathbf{e}_i^* = \lambda_i \mathbf{e}_i$$

for some real number  $\lambda_i$ . As a conclusion of the above cases, we've proved lemma 6.

By Theorem 4, lemma 5 and 6, we've prove that there exist some j, for some positive integer  $\beta_j$ ,  $P + \frac{1}{\beta_j}\Lambda_j$  is a multiple lattice tiling, Theorem 1 is proved.

Acknowledgements. For helpful comments and suggestions, the author is grateful to Professor C.Zong. This work is supported by 973 Program 2013CB834201.

# References

- U.Bölle, On Multiple tiles in ℝ<sup>2</sup>, Intutive Geometry, Colloq. Math. Soc. J. Bolyai 63, North-Holland, Amsterdam, 1994.
- [2] P.Furtwängler, Über Gitter Konstanter Dichte, Monatsh. Math. Phys. 43(1936), 281-288.
- [3] N.Gravin, M.N.Kolountzakis, S.Robins and D.Shiryaev, Structual results for multiple tilings in 3D. Discrete Comput. Geom 32(2012),629-649.
- [4] N.Gravin, S.Robins and D.Shiryaev, Translational tilings by a polytope, with multiplicity. Combinatorica. 32(2012),629-649.
- [5] G.Hajós, Über einfache und mehrfache Bedeckung des n-dimensionalen Raumes mit einem Würfelgitter, Math. Z.47(1941), 427-467.
- [6] M.N.Kolountzakis, On the structure of multiple translational tilings by polygonal regions, *Discrete Comput. Geom.* 23(2000),537-553.
- [7] Bochen Liu, Periodic structure of translational mul-tiling in the plane. arXiv:1809.03440.
- [8] H.Minkowski, Allgemeine Lehrsätze über konvexen Polyeder, Nachr. K. Ges. Wiss. Göttingen, Math. Phys. KL. (1897), 198-219.
- [9] P.McMullen, Convex bodies which tiles space by translation, Mathematika 27(1980),113-121.
- [10] R.M.Robinson, Multiple tilings of n-dimensional space by unit cubes, Math.Z. 166(1979), 225-275.
- [11] B.A.Venkov, On a class of Euclidean polytopes, Vestnik Leningrad Univ, Ser.Mat.Fiz.Him.9(1954),11-31.
- [12] Q.Yang and C. Zong, Multiple translative tilings in euclidean spaces. arXiv:1711.02514.
- [13] Q.Yang and C. Zong, Characterization of the two-dimensional five-fold translative tiles. arXiv:1712.09732.
- [14] C.Zong. The cube: a window to convex and discrete geometry, volume 168 of *Cambridge Tracts* in *Mathematics*. Cambridge University Press, Cambridge, 2006.
- [15] C. Zong, Characterization of the two-dimensional five-fold lattice tiles. arXiv:1712.01122.

Qi Yang, School of Mathematical Science, Peking University, Beijing 100871. China *Email: Yangqi07@pku.edu.cn*