

ON KNÖRRER PERIODICITY FOR QUADRIC HYPERSURFACES IN SKEW PROJECTIVE SPACES

KENTA UEYAMA

ABSTRACT. We study the structure of the stable category $\underline{\mathbf{CM}}^{\mathbb{Z}}(S/(f))$ of graded maximal Cohen-Macaulay module over $S/(f)$ where S is a graded (± 1) -skew polynomial algebra in n variables of degree 1, and $f = x_1^2 + \cdots + x_n^2$. If S is commutative, then the structure of $\underline{\mathbf{CM}}^{\mathbb{Z}}(S/(f))$ is well-known by Knörrer's periodicity theorem. In this paper, we prove that if $n \leq 5$, then the structure of $\underline{\mathbf{CM}}^{\mathbb{Z}}(S/(f))$ is determined by the number of irreducible components of the point scheme of S which are isomorphic to \mathbb{P}^1 .

1. INTRODUCTION

Throughout this paper, we fix an algebraically closed field k of characteristic 0.

Knörrer's periodicity theorem ([5, Theorem 3.1]) plays an essential role in Cohen-Macaulay representation theory of Gorenstein rings. As a special case of Knörrer's periodicity theorem, the following result is well-known (see also [3]).

Theorem 1.1. *Let $S = k[x_1, \dots, x_n]$ be a graded polynomial algebra generated in degree 1, and $f = x_1^2 + x_2^2 + \cdots + x_n^2$. Let $\underline{\mathbf{CM}}^{\mathbb{Z}}(S/(f))$ denote the stable category of graded maximal Cohen-Macaulay module over $S/(f)$.*

- (1) *If n is odd, then $\underline{\mathbf{CM}}^{\mathbb{Z}}(S/(f)) \cong \underline{\mathbf{CM}}^{\mathbb{Z}}(k[x]/(x^2)) \cong \mathbf{D}^b(\text{mod } k)$.*
- (2) *If n is even, then $\underline{\mathbf{CM}}^{\mathbb{Z}}(S/(f)) \cong \underline{\mathbf{CM}}^{\mathbb{Z}}(k[x, y]/(x^2 + y^2)) \cong \mathbf{D}^b(\text{mod } k^2)$.*

The purpose of this paper is to study a “ (± 1) -skew” version of Theorem 1.1.

Definition 1.2. Let $n \in \mathbb{N}^+$.

- (1) We say that S is a graded skew polynomial algebra if

$$S = k\langle x_1, \dots, x_n \rangle / (x_i x_j - \alpha_{ij} x_j x_i)_{1 \leq i, j \leq n}$$

where $\alpha_{ii} = 1$ for every $1 \leq i \leq n$, $\alpha_{ij} \alpha_{ji} = 1$ for every $1 \leq i, j \leq n$, and $\deg x_i = 1$ for every $1 \leq i \leq n$.

- (2) We say that S is a graded (± 1) -skew polynomial algebra if

$$S = k\langle x_1, \dots, x_n \rangle / (x_i x_j - \varepsilon_{ij} x_j x_i)_{1 \leq i, j \leq n}$$

is a graded skew polynomial algebra such that ε_{ij} equals either 1 or -1 for every $1 \leq i, j \leq n, i \neq j$.

Clearly, a graded polynomial algebra $k[x_1, \dots, x_n]$ generated in degree 1 is an example of a graded (± 1) -skew polynomial algebra. Consider the element

$$f = x_1^2 + x_2^2 + \cdots + x_n^2$$

2010 *Mathematics Subject Classification.* 16G50, 16S38, 18E30, 14A22.

Key words and phrases. Knörrer periodicity, stable category, noncommutative quadric hypersurface, skew polynomial algebra, point scheme.

The author was supported by JSPS Grant-in-Aid for Early-Career Scientists 18K13381.

of a graded skew polynomial algebra $S = k\langle x_1, \dots, x_n \rangle / (x_i x_j - \alpha_{ij} x_j x_i)$. Then we notice that f is normal if and only if f is central if and only if S is a (± 1) -skew polynomial algebra.

Let $S = k\langle x_1, \dots, x_n \rangle / (x_i x_j - \varepsilon_{ij} x_j x_i)$ be a graded (± 1) -skew polynomial algebra so that $f = x_1^2 + x_2^2 + \dots + x_n^2 \in S$ is a homogeneous regular central element. Let A be the graded quotient algebra $S/(f)$. Since S is a noetherian AS-regular algebra of dimension n and A is a noetherian AS-Gorenstein algebra of dimension $n - 1$, A is regarded as a homogeneous coordinate ring of a quadric hypersurface in a (± 1) -skew projective space. The main focus of this paper is to determine the structure of $\underline{\mathbf{CM}}^{\mathbb{Z}}(A)$ from a geometric data associated to S called the point scheme of S . Based on our experiments, we propose the following conjecture.

Conjecture 1.3. Let $S = k\langle x_1, \dots, x_n \rangle / (x_i x_j - \varepsilon_{ij} x_j x_i)$ be a graded (± 1) -skew polynomial algebra, $f = x_1^2 + x_2^2 + \dots + x_n^2 \in S$, and $A = S/(f)$. Let ℓ be the number of irreducible components of the point scheme of S which are isomorphic to \mathbb{P}^1 .

(1) If n is odd, then

$$\binom{2m-1}{2} < \ell \leq \binom{2m+1}{2} \iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^{2^{2m}})$$

for $m \in \mathbb{N}$ where we consider $\binom{-1}{2} = -\infty$, $\binom{1}{2} = 0$.

(2) If n is even, then

$$\binom{2m}{2} < \ell \leq \binom{2m+2}{2} \iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^{2^{2m+1}})$$

for $m \in \mathbb{N}$ where we consider $\binom{0}{2} = -\infty$.

We prove the following result.

Theorem 1.4 (Theorem 3.10). *Let $S = k\langle x_1, \dots, x_n \rangle / (x_i x_j - \varepsilon_{ij} x_j x_i)$ be a graded (± 1) -skew polynomial algebra, $f = x_1^2 + x_2^2 + \dots + x_n^2 \in S$, and $A = S/(f)$. Assume that $n \leq 5$. Let ℓ be the number of irreducible components of the point scheme of S which are isomorphic to \mathbb{P}^1 .*

(1) *If n is odd, then $\ell \leq 10$ and*

$$\begin{aligned} \ell = 0 &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k), \\ 0 < \ell \leq 3 &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^4), \\ 3 < \ell \leq 10 &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^{16}). \end{aligned}$$

(2) *If n is even, then $\ell \leq 6$ and*

$$\begin{aligned} 0 \leq \ell \leq 1 &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^2), \\ 1 < \ell \leq 6 &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^8). \end{aligned}$$

This theorem asserts that Conjecture 1.3 is true if $n \leq 5$.

2. PRELIMINARIES

2.1. Notations. For an algebra A , we denote by $\text{Mod } A$ the category of right A -modules, and by $\text{mod } A$ the full subcategory consisting of finitely generated modules. The bounded derived category of $\text{mod } A$ is denoted by $\mathbf{D}^b(\text{mod } A)$.

For a connected graded algebra A , that is, $A = \bigoplus_{i \in \mathbb{N}} A_i$ with $A_0 = k$, we denote by $\text{GrMod } A$ the category of graded right A -modules with A -module homomorphisms of degree zero, and by $\text{grmod } A$ the full subcategory consisting of finitely generated graded modules.

Let A be a noetherian AS-Gorenstein algebra of dimension n (see [4, Section 1] for the definition). We define the local cohomology modules of $M \in \text{grmod } A$ by $\mathbf{H}_m^i(M) := \lim_{n \rightarrow \infty} \text{Ext}_A^i(A/A_{\geq n}, M)$.

It is well-known that $H_m^i(A) = 0$ for all $i \neq n$. We say that $M \in \text{grmod } A$ is graded maximal Cohen-Macaulay if $H_m^i(M) = 0$ for all $i \neq n$. We denote by $\text{CM}^{\mathbb{Z}}(A)$ the full subcategory of $\text{grmod } A$ consisting of graded maximal Cohen-Macaulay modules.

The stable category of graded maximal Cohen-Macaulay modules, denoted by $\underline{\text{CM}}^{\mathbb{Z}}(A)$, has the same objects as $\text{CM}^{\mathbb{Z}}(A)$ and the morphism set is given by

$$\text{Hom}_{\underline{\text{CM}}^{\mathbb{Z}}(A)}(M, N) = \text{Hom}_{\text{grmod } A}(M, N)/P(M, N)$$

for any $M, N \in \text{CM}^{\mathbb{Z}}(A)$, where $P(M, N)$ consists of degree zero A -module homomorphisms that factor through a projective module in $\text{grmod } A$. Since A is AS-Gorenstein, $\underline{\text{CM}}^{\mathbb{Z}}(A)$ is a triangulated category with respect to the translation functor $M[-1] = \Omega M$ (the syzygy of M) by [8, Theorem 3.1].

2.2. The algebra $C(A)$. The method we use is due to Smith and Van den Bergh [8]; it was originally developed by Buchweitz, Eisenbud, and Herzog [3].

Let S be an n -dimensional noetherian AS-regular algebra with the Hilbert series $H_S(t) = (1-t)^{-n}$. Then S is Koszul by [7, Theorem 5.11]. Let $f \in S$ be a homogeneous regular central element of degree 2, and let $A = S/(f)$. Then A is Koszul by [8, Lemma 5.1 (1)], and there exists a central regular element $w \in A_2^!$ such that $A^!/(w) \cong S^!$ by [8, Lemma 5.1 (2)]. We can define the algebra

$$C(A) := A^![w^{-1}]_0.$$

By [8, Lemma 5.1 (3)], we have $\dim_k C(A) = \dim_k(S^!)^{(2)} = 2^{n-1}$.

Theorem 2.1 ([8, Proposition 5.2]). *Let notation be as above. Then $\underline{\text{CM}}^{\mathbb{Z}}(A) \cong \text{D}^b(\text{mod } C(A))$.*

2.3. The point schemes of skew polynomial algebras. Let S be a quantum polynomial algebra of dimension n (see [6, Definition 2.1] for the definition).

Definition 2.2. A graded module $M \in \text{GrMod } S$ is called a point module if M is cyclic, generated in degree 0, and $H_M(t) = (1-t)^{-1}$.

If $M \in \text{GrMod } S$ is a point module, then M can be written as a quotient $S/(g_1S + g_2S + \cdots + g_{n-1}S)$ with linearly independent $g_1, \dots, g_{n-1} \in S_1$ by [6, Corollary 5.7, Theorem 3.8], so we can associate it with a unique point $p_M := \mathcal{V}(g_1, \dots, g_{n-1})$ in $\mathbb{P}(S_1^*) = \mathbb{P}^{n-1}$. Then the subset

$$E := \{p_M \in \mathbb{P}^{n-1} \mid M \in \text{GrMod } S \text{ is a point module}\}$$

has a k -scheme structure by [1], and it is called the point scheme of S . Point schemes have a pivotal role in noncommutative algebraic geometry.

Thanks to the following result, we can compute the point scheme of a graded skew polynomial algebra.

Theorem 2.3 ([9, Proposition 4.2], [2, Theorem 1 (1)]). *Let $S = k\langle x_1, \dots, x_n \rangle / (x_i x_j - \alpha_{ij} x_j x_i)$ be a graded skew polynomial algebra. Then the point scheme of S is given by*

$$E = \bigcap_{\substack{1 \leq i < j < k \leq n \\ \alpha_{ij} \alpha_{jk} \alpha_{ki} \neq 1}} \mathcal{V}(x_i x_j x_k) \subset \mathbb{P}^{n-1}.$$

For $1 \leq i_0, \dots, i_s \leq n$, we define the subspace

$$\mathbb{P}(i_1, \dots, i_s) := \bigcap_{\substack{1 \leq j \leq n \\ j \neq i_1, \dots, j \neq i_s}} \mathcal{V}(x_j) \subset \mathbb{P}^{n-1}.$$

It is easy to see that the point scheme of a graded skew polynomial algebra in 3 variables is isomorphic to \mathbb{P}^2 or $\mathbb{P}(2, 3) \cup \mathbb{P}(1, 3) \cup \mathbb{P}(1, 2)$. The following is the classification of the point schemes of graded skew polynomial algebras in 4 variables.

Proposition 2.4 ([9, Corollary 5.1], [2, Section 4.2]). *Let $S = k\langle x_1, x_2, x_3, x_4 \rangle / (x_i x_j - \alpha_{ij} x_j x_i)$ be a graded skew polynomial algebra in 4 variables. Then the point scheme of S is isomorphic one of the following:*

- \mathbb{P}^3 ;
- $\mathbb{P}(1, 2, 4) \cup \mathbb{P}(1, 2, 3) \cup \mathbb{P}(3, 4)$;
- $\mathbb{P}(2, 3, 4) \cup \mathbb{P}(1, 4) \cup \mathbb{P}(1, 3) \cup \mathbb{P}(1, 2)$;
- $\mathbb{P}(3, 4) \cup \mathbb{P}(2, 4) \cup \mathbb{P}(2, 3) \cup \mathbb{P}(1, 4) \cup \mathbb{P}(1, 3) \cup \mathbb{P}(1, 2)$.

3. RESULTS

Throughout this section,

- $S = k\langle x_1, \dots, x_n \rangle / (x_i x_j - \varepsilon_{ij} x_j x_i)$ is a graded (± 1) -skew polynomial algebra,
- E is the point scheme of S ,
- $f = x_1^2 + x_2^2 + \dots + x_n^2 \in S_2$ (a regular central element of S), and
- $A = S/(f)$.

Note that $\varepsilon_{ij} = \varepsilon_{ji}$ holds for every $1 \leq i, j \leq n$.

Lemma 3.1. (1) $A^!$ is isomorphic to $k\langle x_1, \dots, x_n \rangle / (\varepsilon_{ij} x_i x_j + x_j x_i, x_n^2 - x_i^2)_{1 \leq i, j \leq n, i \neq j}$.
 (2) $w = x_n^2 \in A_2^!$ is a central regular element such that $A^!/(w) \cong S^!$.
 (3) $C(A) := A^![w^{-1}]_0$ is isomorphic to

$$k\langle t_1, \dots, t_{n-1} \rangle / (t_i t_j + \varepsilon_{ni} \varepsilon_{ij} \varepsilon_{jn} t_j t_i, t_i^2 - 1)_{1 \leq i, j \leq n-1, i \neq j}.$$

Proof. (1) and (2) follow from direct calculation.

(3) Since S has a k -basis $\{x_1^{i_1} x_2^{i_2} \dots x_n^{i_n} \mid i_1, i_2, \dots, i_n \geq 0\}$, and

$$(x_n x_i w^{-1})(x_n x_j w^{-1}) = x_n x_i x_n x_j w^{-2} = -\varepsilon_{ni} x_n^2 x_i x_j w^{-2} = -\varepsilon_{ni} x_i x_j w^{-1}$$

in $C(A)$ for $1 \leq i \leq n-1, i \neq j$, it follows that $\{x_n x_1 w^{-1}, \dots, x_n x_{n-1} w^{-1}\}$ is a set of generators of $C(A)$. Put $t_i := x_n x_i w^{-1}$ for $1 \leq i \leq n-1$. Since

$$\begin{aligned} t_i t_j &= (x_n x_i w^{-1})(x_n x_j w^{-1}) = -\varepsilon_{ni} x_i x_j w^{-1} = \varepsilon_{ni} \varepsilon_{ji} x_j x_i w^{-1} \\ &= -\varepsilon_{ni} \varepsilon_{ij} \varepsilon_{jn} (-\varepsilon_{nj} x_j x_i w^{-1}) = -\varepsilon_{ni} \varepsilon_{ij} \varepsilon_{jn} (x_n x_j w^{-1})(x_n x_i w^{-1}) = -\varepsilon_{ni} \varepsilon_{ij} \varepsilon_{jn} t_j t_i, \end{aligned}$$

for $1 \leq i, j \leq n-1, i \neq j$, and

$$t_i^2 = (x_n x_i w^{-1})(x_n x_i w^{-1}) = -\varepsilon_{ni} x_i^2 w^{-1} = -\varepsilon_{ni} x_n^2 w^{-1} = -\varepsilon_{ni}$$

for $1 \leq i \leq n-1$, we have a surjection $k\langle t_1, \dots, t_{n-1} \rangle / (t_i t_j + \varepsilon_{ni} \varepsilon_{ij} \varepsilon_{jn} t_j t_i, t_i^2 + \varepsilon_{ni}) \rightarrow C(A)$. This is an isomorphism because the algebras have the same dimension. Since $\varepsilon_{ni} \neq 0$ for $1 \leq i \leq n-1$, the homomorphism defined by $t_i \rightarrow \sqrt{-\varepsilon_{ni}} t_i$ induces the isomorphism

$$k\langle t_1, \dots, t_{n-1} \rangle / (t_i t_j + \varepsilon_{ni} \varepsilon_{ij} \varepsilon_{jn} t_j t_i, t_i^2 + \varepsilon_{ni}) \xrightarrow{\sim} k\langle t_1, \dots, t_{n-1} \rangle / (t_i t_j + \varepsilon_{ni} \varepsilon_{ij} \varepsilon_{jn} t_j t_i, t_i^2 - 1).$$

□

Proposition 3.2. (1) If $E = \mathbb{P}^{n-1}$, then $C(A)$ is isomorphic to

$$C_+ := k\langle t_1, \dots, t_{n-1} \rangle / (t_i t_j + t_j t_i, t_i^2 - 1)_{1 \leq i, j \leq n-1, i \neq j}.$$

(2) $E = \bigcup_{1 \leq i < j \leq n} \mathbb{P}(i, j)$ if and only if $C(A)$ is isomorphic to

$$C_- := k\langle t_1, \dots, t_{n-1} \rangle / (t_i t_j - t_j t_i, t_i^2 - 1)_{1 \leq i, j \leq n-1, i \neq j}.$$

Proof. First note that

$$\varepsilon_{ij}\varepsilon_{jk}\varepsilon_{ki} = (\varepsilon_{ni}\varepsilon_{ij}\varepsilon_{jn})(\varepsilon_{nj}\varepsilon_{jk}\varepsilon_{kn})(\varepsilon_{nk}\varepsilon_{ki}\varepsilon_{in}) \quad (3.1)$$

for $1 \leq i < j < k \leq n$.

(1) By Theorem 2.3, (3.1), and Lemma 3.1 (3), it follows that

$$\begin{aligned} E = \mathbb{P}^{n-1} &\iff \varepsilon_{ij}\varepsilon_{jk}\varepsilon_{ki} = 1 \text{ for every } 1 \leq i < j < k \leq n \\ &\iff \varepsilon_{ni}\varepsilon_{ij}\varepsilon_{jn} = 1 \text{ for every } 1 \leq i < j \leq n \\ &\implies C(A) \cong C_+. \end{aligned}$$

(2) By Theorem 2.3, (3.1), and Lemma 3.1 (3), it follows that

$$\begin{aligned} E = \bigcup_{1 \leq i < j \leq n} \mathbb{P}(i, j) &\iff \varepsilon_{ij}\varepsilon_{jk}\varepsilon_{ki} \neq 1 \text{ for every } 1 \leq i < j < k \leq n \\ &\iff \varepsilon_{ij}\varepsilon_{jk}\varepsilon_{ki} = -1 \text{ for every } 1 \leq i < j < k \leq n \\ &\iff \varepsilon_{ni}\varepsilon_{ij}\varepsilon_{jn} = -1 \text{ for every } 1 \leq i < j \leq n \\ &\iff C(A) \cong C_-. \end{aligned}$$

Here the last \iff is by commutativity of $C(A)$. □

Theorem 3.3. (1) If $E = \mathbb{P}^{n-1}$ and n is odd, then $\underline{\mathbf{CM}}^{\mathbb{Z}}(S/(f)) \cong \mathbf{D}^b(\text{mod } k)$.

(2) If $E = \mathbb{P}^{n-1}$ and n is even, then $\underline{\mathbf{CM}}^{\mathbb{Z}}(S/(f)) \cong \mathbf{D}^b(\text{mod } k^2)$.

(3) $E = \bigcup_{1 \leq i < j \leq n} \mathbb{P}(i, j)$ if and only if $\underline{\mathbf{CM}}^{\mathbb{Z}}(S/(f)) \cong \mathbf{D}^b(\text{mod } k^{2^{n-1}})$.

Proof. Since C_+ is a Clifford algebra over k , it is known that

$$C_+ \cong \begin{cases} M_{2^{(n-1)/2}}(k) & \text{if } n \text{ is odd,} \\ M_{2^{(n-2)/2}}(k)^2 & \text{if } n \text{ is even,} \end{cases} \quad (3.2)$$

so

$$\text{mod } C_+ \cong \begin{cases} \text{mod } M_{2^{(n-1)/2}}(k) \cong \text{mod } k & \text{if } n \text{ is odd,} \\ \text{mod } M_{2^{(n-2)/2}}(k)^2 \cong \text{mod } k^2 & \text{if } n \text{ is even.} \end{cases}$$

Thus (1) and (2) follow from Theorem 2.1 and Proposition 3.2 (1).

We next show (3). If $E = \bigcup_{1 \leq i < j \leq n} \mathbb{P}(i, j)$, then $C(A) \cong C_-$ by Proposition 3.2 (2). Since C_- is isomorphic to the group algebra of $(\mathbb{Z}_2)^{n-1}$ over k , we have $C_- \cong k^{2^{n-1}}$, so it follows that $\underline{\mathbf{CM}}^{\mathbb{Z}}(S/(f)) \cong \mathbf{D}^b(\text{mod } k^{2^{n-1}})$ by Theorem 2.1. Conversely, if $\underline{\mathbf{CM}}^{\mathbb{Z}}(S/(f)) \cong \mathbf{D}^b(\text{mod } k^{2^{n-1}})$, then $\mathbf{D}^b(\text{mod } C(A)) \cong \mathbf{D}^b(\text{mod } k^{2^{n-1}})$ by Theorem 2.1. Since $\dim_k C(A) = 2^{n-1}$, it follows that $C(A) \cong k^{2^{n-1}} \cong C_-$. Hence $E = \bigcup_{1 \leq i < j \leq n} \mathbb{P}(i, j)$ by Proposition 3.2 (2). □

Note that Theorem 3.3 (1), (2) recover Theorem 1.1, and Theorem 3.3 (3) shows that a new phenomenon appears in the noncommutative case. We can now give an explicit classification of $\underline{\mathbf{CM}}^{\mathbb{Z}}(A)$ in the case $n \leq 3$ (the case $n = 1$ is clear; see Theorem 1.1 (1)).

Corollary 3.4. (1) If $n = 2$, then $E = \mathbb{P}^1$ and $\underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^2)$.

(2) If $n = 3$, then

$$\begin{aligned} E = \mathbb{P}^2 &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k), \\ E = \mathbb{P}(2, 3) \cup \mathbb{P}(1, 3) \cup \mathbb{P}(1, 2) &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^4). \end{aligned}$$

Proof. These follow from Theorem 2.3 and Theorem 3.3. □

As we will see later, the converse of Theorem 3.3 (1), (2) does not hold in general. So, in order to give a classification for the cases $n = 4$ and $n = 5$, we need a precise computation.

For a permutation $\sigma \in \mathfrak{S}_n$, we have an isomorphism

$$S = k\langle x_1, \dots, x_n \rangle / (x_i x_j - \varepsilon_{ij} x_j x_i) \xrightarrow[\varphi]{\sim} k\langle x_1, \dots, x_n \rangle / (x_{\sigma(i)} x_{\sigma(j)} - \varepsilon_{ij} x_{\sigma(j)} x_{\sigma(i)}) =: S_\sigma$$

between graded (± 1) -skew polynomial algebras, which we call a permutation isomorphism. Since φ preserves f , it induces an isomorphism

$$A = S/(f) \xrightarrow{\sim} S_\sigma/(f),$$

which we also call a permutation isomorphism.


Lemma 3.5. *If $n = 4$, then, via a permutation isomorphism, S is isomorphic to a graded (± 1) -skew polynomial algebra whose point scheme is one of the following:*

- (4a) \mathbb{P}^3 ;
- (4b) $\mathbb{P}(1, 2, 4) \cup \mathbb{P}(1, 2, 3) \cup \mathbb{P}(3, 4)$;
- (4c) $\mathbb{P}(3, 4) \cup \mathbb{P}(2, 4) \cup \mathbb{P}(2, 3) \cup \mathbb{P}(1, 4) \cup \mathbb{P}(1, 3) \cup \mathbb{P}(1, 2)$.

Proof. First, via a permutation isomorphism, S is isomorphic to one of the following:

- (4i) a graded (± 1) -skew polynomial algebra with $\varepsilon_{41}\varepsilon_{12}\varepsilon_{24} = \varepsilon_{41}\varepsilon_{13}\varepsilon_{34} = \varepsilon_{42}\varepsilon_{23}\varepsilon_{34} = 1$;
- (4ii) a graded (± 1) -skew polynomial algebra with $\varepsilon_{41}\varepsilon_{12}\varepsilon_{24} = \varepsilon_{41}\varepsilon_{13}\varepsilon_{34} = 1$, $\varepsilon_{42}\varepsilon_{23}\varepsilon_{34} = -1$;
- (4iii) a graded (± 1) -skew polynomial algebra with $\varepsilon_{41}\varepsilon_{12}\varepsilon_{24} = 1$, $\varepsilon_{41}\varepsilon_{13}\varepsilon_{34} = \varepsilon_{42}\varepsilon_{23}\varepsilon_{34} = -1$;
- (4iv) a graded (± 1) -skew polynomial algebra with $\varepsilon_{41}\varepsilon_{12}\varepsilon_{24} = \varepsilon_{41}\varepsilon_{13}\varepsilon_{34} = \varepsilon_{42}\varepsilon_{23}\varepsilon_{34} = -1$.

Note that the above follows from (3.1) and the classification of simple graphs of order 3:



(we define $\varepsilon_{4i}\varepsilon_{ij}\varepsilon_{j4} = -1$ if $\{i, j\}$ is an edge in the graph, and $\varepsilon_{4i}\varepsilon_{ij}\varepsilon_{j4} = 1$ otherwise).

The point scheme of an algebra in the case (4i) is \mathbb{P}^3 , so this is (4a).

The point scheme of an algebra in the case (4iii) is

$$\mathcal{V}(x_1 x_3 x_4) \cap \mathcal{V}(x_2 x_3 x_4) = \mathcal{V}(x_3) \cup \mathcal{V}(x_4) \cup \mathcal{V}(x_1, x_2),$$

so this is (4b). The point scheme of an algebra in the case (4ii) is $\mathcal{V}(x_1 x_2 x_3) \cap \mathcal{V}(x_2 x_3 x_4) = \mathcal{V}(x_2) \cup \mathcal{V}(x_3) \cup \mathcal{V}(x_1, x_4)$, so an algebra in the case (4ii) is isomorphic to an algebra in the case (4iii) via the permutation isomorphism induced by $\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 4 & 3 & 2 \end{pmatrix}$.

The point scheme of an algebra in the case (4iv) is $\bigcap_{1 \leq i < j < k \leq 4} \mathcal{V}(x_i x_j x_k) = \bigcup_{1 \leq i < j \leq 4} \mathcal{V}(x_i, x_j)$, so this is (4c). \square

Remark 3.6. It follows from Lemma 3.5 that not every point scheme in Proposition 2.4 appears as the point scheme of a graded (± 1) -skew polynomial algebra.

Lemma 3.7. *If $n = 5$, then, via a permutation isomorphism, S is isomorphic to a graded (± 1) -skew polynomial algebra whose point scheme is one of the following:*

- (5a) \mathbb{P}^4 ;
- (5b) $\mathbb{P}(1, 2, 3, 5) \cup \mathbb{P}(1, 2, 3, 4) \cup \mathbb{P}(4, 5)$;
- (5c) $\mathbb{P}(1, 2, 3, 4) \cup \mathbb{P}(3, 4, 5) \cup \mathbb{P}(1, 2, 5)$;
- (5d) $\mathbb{P}(3, 4, 5) \cup \mathbb{P}(1, 4, 5) \cup \mathbb{P}(1, 2, 5) \cup \mathbb{P}(1, 2, 3) \cup \mathbb{P}(2, 3, 4)$;
- (5e) $\mathbb{P}(1, 3, 5) \cup \mathbb{P}(1, 3, 4) \cup \mathbb{P}(1, 2, 5) \cup \mathbb{P}(1, 2, 4) \cup \mathbb{P}(4, 5) \cup \mathbb{P}(2, 3)$;
- (5f) $\mathbb{P}(1, 2, 5) \cup \mathbb{P}(1, 2, 4) \cup \mathbb{P}(1, 2, 3) \cup \mathbb{P}(4, 5) \cup \mathbb{P}(3, 5) \cup \mathbb{P}(3, 4)$;
- (5g) $\mathbb{P}(4, 5) \cup \mathbb{P}(3, 5) \cup \mathbb{P}(3, 4) \cup \mathbb{P}(2, 5) \cup \mathbb{P}(2, 4) \cup \mathbb{P}(2, 3) \cup \mathbb{P}(1, 5) \cup \mathbb{P}(1, 4) \cup \mathbb{P}(1, 3) \cup \mathbb{P}(1, 2)$.

Proof. First, via a permutation isomorphism, S is isomorphic to one of the following:

(5i) a graded (± 1) -skew polynomial algebra with

$$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = 1, \varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = 1, \varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = 1, \varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = 1, \varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = 1, \varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = 1;$$

(5ii) a graded (± 1) -skew polynomial algebra with

$$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = 1, \varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = 1, \varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = 1, \varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = 1, \varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = 1, \varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = -1;$$

(5iii) a graded (± 1) -skew polynomial algebra with

$$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = 1, \varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = 1, \varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = 1, \varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = 1, \varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = -1, \varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = -1;$$

(5iv) a graded (± 1) -skew polynomial algebra with

$$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = -1, \varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = 1, \varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = 1, \varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = 1, \varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = 1, \varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = -1;$$

(5v) a graded (± 1) -skew polynomial algebra with

$$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = 1, \varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = 1, \varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = -1, \varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = 1, \varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = -1, \varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = -1;$$

(5vi) a graded (± 1) -skew polynomial algebra with

$$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = 1, \varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = -1, \varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = 1, \varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = -1, \varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = -1, \varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = 1;$$

(5vii) a graded (± 1) -skew polynomial algebra with

$$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = 1, \varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = -1, \varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = -1, \varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = 1, \varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = 1, \varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = -1;$$

(5viii) a graded (± 1) -skew polynomial algebra with

$$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = 1, \varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = -1, \varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = -1, \varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = -1, \varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = -1, \varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = 1;$$

(5ix) a graded (± 1) -skew polynomial algebra with

$$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = 1, \varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = 1, \varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = -1, \varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = -1, \varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = -1, \varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = -1;$$


(5x) a graded (± 1) -skew polynomial algebra with

$$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = 1, \varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = -1, \varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = -1, \varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = -1, \varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = -1, \varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = -1;$$

(5xi) a graded (± 1) -skew polynomial algebra with

$$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = -1, \varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = -1, \varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = -1, \varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = -1, \varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = -1, \varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = -1;$$

Note that the above follows from (3.1) and the classification of simple graphs of order 4:



(we define $\varepsilon_{5i}\varepsilon_{ij}\varepsilon_{j5} = -1$ if $\{i, j\}$ is an edge in the graph, and $\varepsilon_{5i}\varepsilon_{ij}\varepsilon_{j5} = 1$ otherwise).

The point scheme of an algebra in the case (5i) is \mathbb{P}^4 , so this is (5a).

The point scheme of an algebra in the case (5v) is

$$\mathcal{V}(x_1x_4x_5) \cap \mathcal{V}(x_2x_4x_5) \cap \mathcal{V}(x_3x_4x_5) = \mathcal{V}(x_4) \cup \mathcal{V}(x_5) \cup \mathcal{V}(x_1, x_2, x_3),$$

so this is (5b). The point scheme of an algebra in the case (5ii) is $\mathcal{V}(x_1x_3x_4) \cap \mathcal{V}(x_2x_3x_4) \cap \mathcal{V}(x_3x_4x_5) = \mathcal{V}(x_3) \cup \mathcal{V}(x_4) \cup \mathcal{V}(x_1, x_2, x_5)$, so an algebra in the case (5ii) is isomorphic to an algebra in the case (5v) via the permutation isomorphism induced by $\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 5 & 4 & 3 \end{pmatrix}$.

The point scheme of an algebra in the case (5viii) is

$$\mathcal{V}(x_1x_3x_5) \cap \mathcal{V}(x_1x_4x_5) \cap \mathcal{V}(x_2x_3x_5) \cap \mathcal{V}(x_2x_4x_5) = \mathcal{V}(x_5) \cup \mathcal{V}(x_1, x_2) \cup \mathcal{V}(x_3, x_4),$$

so this is (5c). The point scheme of an algebra in the case (5iii) is $\mathcal{V}(x_1x_2x_4) \cap \mathcal{V}(x_1x_3x_4) \cap \mathcal{V}(x_2x_4x_5) \cap \mathcal{V}(x_3x_4x_5) = \mathcal{V}(x_4) \cup \mathcal{V}(x_1, x_5) \cup \mathcal{V}(x_2, x_3)$, so an algebra in the case (5iii) is isomorphic to an algebra in the case (5viii) via the permutation isomorphism induced by $\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 4 & 3 & 5 & 2 \end{pmatrix}$.

The point scheme of an algebra in the case (5vi) is

$$\begin{aligned} & \mathcal{V}(x_1x_2x_4) \cap \mathcal{V}(x_1x_3x_4) \cap \mathcal{V}(x_1x_3x_5) \cap \mathcal{V}(x_2x_3x_5) \cap \mathcal{V}(x_2x_4x_5) \\ & = \mathcal{V}(x_1, x_2) \cup \mathcal{V}(x_2, x_3) \cup \mathcal{V}(x_3, x_4) \cup \mathcal{V}(x_4, x_5) \cup \mathcal{V}(x_5, x_1), \end{aligned}$$

so this is (5d).

The point scheme of an algebra in the case (5ix) is

$$\begin{aligned} & \mathcal{V}(x_1x_2x_3) \cap \mathcal{V}(x_1x_4x_5) \cap \mathcal{V}(x_2x_3x_4) \cap \mathcal{V}(x_2x_3x_5) \cap \mathcal{V}(x_2x_4x_5) \cap \mathcal{V}(x_3x_4x_5) \\ & = \mathcal{V}(x_2, x_4) \cup \mathcal{V}(x_2, x_5) \cup \mathcal{V}(x_3, x_4) \cup \mathcal{V}(x_3, x_5) \cup \mathcal{V}(x_1, x_2, x_3) \cup \mathcal{V}(x_1, x_4, x_5), \end{aligned}$$

so this is (5e). The point scheme of an algebra in the case (5iv) is $\mathcal{V}(x_1x_2x_3) \cap \mathcal{V}(x_1x_2x_4) \cap \mathcal{V}(x_1x_2x_5) \cap \mathcal{V}(x_1x_3x_4) \cap \mathcal{V}(x_2x_3x_4) \cap \mathcal{V}(x_3x_4x_5) = \mathcal{V}(x_1, x_3) \cup \mathcal{V}(x_1, x_4) \cup \mathcal{V}(x_2, x_3) \cup \mathcal{V}(x_2, x_4) \cup \mathcal{V}(x_1, x_2, x_5) \cup \mathcal{V}(x_3, x_4, x_5)$, so an algebra in the case (5iv) is isomorphic to an algebra in the case (5ix) via the permutation isomorphism induced by $\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 2 & 5 & 4 & 1 \end{pmatrix}$.

The point scheme of an algebra in the case (5x) is

$$\begin{aligned} & \mathcal{V}(x_1x_3x_4) \cap \mathcal{V}(x_1x_3x_5) \cap \mathcal{V}(x_1x_4x_5) \cap \mathcal{V}(x_2x_3x_4) \cap \mathcal{V}(x_2x_3x_5) \cap \mathcal{V}(x_2x_4x_5) \cap \mathcal{V}(x_3x_4x_5) \\ & = \mathcal{V}(x_3, x_4) \cup \mathcal{V}(x_3, x_5) \cup \mathcal{V}(x_4, x_5) \cup \mathcal{V}(x_1, x_2, x_3) \cup \mathcal{V}(x_1, x_2, x_4) \cup \mathcal{V}(x_1, x_2, x_5), \end{aligned}$$

so this is (5f). The point scheme of an algebra in the case (5vii) is $\mathcal{V}(x_1x_2x_3) \cap \mathcal{V}(x_1x_2x_4) \cap \mathcal{V}(x_1x_3x_4) \cap \mathcal{V}(x_1x_3x_5) \cap \mathcal{V}(x_1x_4x_5) \cap \mathcal{V}(x_2x_3x_4) \cap \mathcal{V}(x_3x_4x_5) = \mathcal{V}(x_1, x_3) \cup \mathcal{V}(x_1, x_4) \cup \mathcal{V}(x_3, x_4) \cup \mathcal{V}(x_1, x_2, x_5) \cup \mathcal{V}(x_2, x_3, x_5) \cup \mathcal{V}(x_2, x_4, x_5)$, so an algebra in the case (5vii) is isomorphic to an algebra in the case (5x) via the permutation isomorphism induced by $\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 5 & 2 & 3 & 4 & 1 \end{pmatrix}$.

The point scheme of an algebra in the case (5xi) is $\bigcap_{1 \leq i < j < k \leq 5} \mathcal{V}(x_i x_j x_k) = \bigcup_{1 \leq i < j < k \leq 5} \mathcal{V}(x_i, x_j, x_k)$, so this is (5g). \square

To describe the algebras $C(A)$ appearing in Lemma 3.1, we show that the following algebras are isomorphic to algebras of the form $M_i(k)^j$.

- Lemma 3.8.** (1) $C_i := k\langle t_1, t_2, t_3 \rangle / (t_1t_2 + t_2t_1, t_1t_3 + t_3t_1, t_2t_3 - t_3t_2, t_1^2 - 1, t_2^2 - 1, t_3^2 - 1)$ is isomorphic to $M_2(k)^2$.
- (2) $C_{ii} := k\langle t_1, t_2, t_3 \rangle / (t_1t_2 + t_2t_1, t_1t_3 - t_3t_1, t_2t_3 - t_3t_2, t_1^2 - 1, t_2^2 - 1, t_3^2 - 1)$ is isomorphic to $M_2(k)^2$.
- (3) $C_{iii} := k\langle t_1, t_2, t_3, t_4 \rangle / (t_1t_2 + t_2t_1, t_1t_3 + t_3t_1, t_1t_4 - t_4t_1, t_2t_3 + t_3t_2, t_2t_4 - t_4t_2, t_3t_4 - t_4t_3, t_1^2 - 1, t_2^2 - 1, t_3^2 - 1, t_4^2 - 1)$ is isomorphic to $M_2(k)^4$.
- (4) $C_{iv} := k\langle t_1, t_2, t_3, t_4 \rangle / (t_1t_2 + t_2t_1, t_1t_3 - t_3t_1, t_1t_4 - t_4t_1, t_2t_3 - t_3t_2, t_2t_4 - t_4t_2, t_3t_4 + t_4t_3, t_1^2 - 1, t_2^2 - 1, t_3^2 - 1, t_4^2 - 1)$ is isomorphic to $M_4(k)$.
- (5) $C_v := k\langle t_1, t_2, t_3, t_4 \rangle / (t_1t_2 + t_2t_1, t_1t_3 - t_3t_1, t_1t_4 + t_4t_1, t_2t_3 - t_3t_2, t_2t_4 - t_4t_2, t_3t_4 + t_4t_3, t_1^2 - 1, t_2^2 - 1, t_3^2 - 1, t_4^2 - 1)$ is isomorphic to $M_4(k)$.
- (6) $C_{vi} := k\langle t_1, t_2, t_3, t_4 \rangle / (t_1t_2 + t_2t_1, t_1t_3 + t_3t_1, t_1t_4 - t_4t_1, t_2t_3 - t_3t_2, t_2t_4 - t_4t_2, t_3t_4 - t_4t_3, t_1^2 - 1, t_2^2 - 1, t_3^2 - 1, t_4^2 - 1)$ is isomorphic to $M_2(k)^4$.
- (7) $C_{vii} := k\langle t_1, t_2, t_3, t_4 \rangle / (t_1t_2 + t_2t_1, t_1t_3 - t_3t_1, t_1t_4 - t_4t_1, t_2t_3 - t_3t_2, t_2t_4 - t_4t_2, t_3t_4 - t_4t_3, t_1^2 - 1, t_2^2 - 1, t_3^2 - 1, t_4^2 - 1)$ is isomorphic to $M_2(k)^4$.

Proof. (1) Let

$$\begin{aligned} e_1 &= \frac{1}{4}(1 + t_2 + t_3 + t_2t_3), & e_2 &= \frac{1}{4}(1 - t_2 + t_3 - t_2t_3), \\ e_3 &= \frac{1}{4}(1 + t_2 - t_3 - t_2t_3), & e_4 &= \frac{1}{4}(1 - t_2 - t_3 + t_2t_3). \end{aligned}$$

Then they form a complete set of orthogonal idempotents of C_i . Since

$$\begin{aligned} e_1 t_1 &= \frac{1}{4}(1 + t_2 + t_3 + t_2 t_3) t_1 = \frac{1}{4} t_2 (1 - t_1 - t_3 + t_2 t_3) = t_1 e_4, \\ e_2 t_1 &= \frac{1}{4}(1 - t_2 + t_3 - t_2 t_3) t_1 = \frac{1}{4} t_2 (1 + t_1 - t_3 - t_2 t_3) = t_1 e_3, \\ e_3 t_1 &= \frac{1}{4}(1 + t_2 - t_3 - t_2 t_3) t_1 = \frac{1}{4} t_2 (1 - t_1 + t_3 - t_2 t_3) = t_1 e_2, \\ e_4 t_1 &= \frac{1}{4}(1 - t_2 - t_3 + t_2 t_3) t_1 = \frac{1}{4} t_2 (1 + t_1 + t_3 + t_2 t_3) = t_1 e_1, \end{aligned}$$

it follows that the map $M_2(k)^2 \rightarrow C_i$;

$$\left(\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}, \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} \right) \mapsto \begin{array}{cccc} a_{11}e_1 & +a_{12}e_1t_1e_4 & +b_{11}e_2 & +b_{12}e_2t_1e_3 \\ +a_{21}e_4t_1e_1 & +a_{22}e_4 & +b_{21}e_3t_1e_2 & +b_{22}e_3 \end{array}$$

is an isomorphism of algebras.

(2) Since t_3 commutes with t_1, t_2 in C_{ii} , we have

$$C_{ii} \cong k\langle t_1, t_2 \rangle / (t_1 t_2 + t_2 t_1, t_1^2 - 1, t_2^2 - 1) \otimes_k k[t_3] / (t_3^2 - 1) \cong M_2(k) \otimes_k k^2 \cong M_2(k)^2$$

by (3.2).

(3) Since t_4 commutes with t_1, t_2, t_3 in C_{iii} , we have

$$\begin{aligned} C_{iii} &\cong k\langle t_1, t_2, t_3 \rangle / (t_1 t_2 + t_2 t_1, t_1 t_3 + t_3 t_1, t_2 t_3 + t_3 t_2, t_1^2 - 1, t_2^2 - 1, t_3^2 - 1) \otimes_k k[t_4] / (t_4^2 - 1) \\ &\cong M_2(k)^2 \otimes_k k^2 \cong M_2(k)^4 \end{aligned}$$

by (3.2).

(4) Since t_3, t_4 commute with t_1, t_2 in C_{iv} , we have

$$\begin{aligned} C_{iv} &\cong k\langle t_1, t_2 \rangle / (t_1 t_2 + t_2 t_1, t_1^2 - 1, t_2^2 - 1) \otimes_k k\langle t_3, t_4 \rangle / (t_3 t_4 + t_4 t_3, t_3^2 - 1, t_4^2 - 1) \\ &\cong M_2(k) \otimes_k M_2(k) \cong M_4(k) \end{aligned}$$

by (3.2).

(5) Let

$$\begin{aligned} e_1 &= \frac{1}{4}(1 + t_1 + t_3 + t_1 t_3), & e_2 &= \frac{1}{4}(1 - t_1 + t_3 - t_1 t_3), \\ e_3 &= \frac{1}{4}(1 + t_1 - t_3 - t_1 t_3), & e_4 &= \frac{1}{4}(1 - t_1 - t_3 + t_1 t_3). \end{aligned}$$

Then they form a complete set of orthogonal idempotents of C_v . Similar to the proof of (1), we have

$$\begin{aligned} e_1 t_4 &= t_4 e_4, & e_1 t_2 &= t_2 e_2, & e_1 t_4 t_2 &= t_4 t_2 e_3, \\ e_2 t_4 &= t_4 e_3, & e_2 t_2 &= t_2 e_1, & e_2 t_4 t_2 &= t_4 t_2 e_4, \\ e_3 t_4 &= t_4 e_2, & e_3 t_2 &= t_2 e_4, & e_3 t_4 t_2 &= t_4 t_2 e_1, \\ e_4 t_4 &= t_4 e_1, & e_4 t_2 &= t_2 e_3, & e_4 t_4 t_2 &= t_4 t_2 e_2, \end{aligned}$$

so it follows that the map $M_4(k) \rightarrow C_v$;

$$\left(\begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix} \right) \mapsto \begin{array}{cccc} a_{11}e_1 & +a_{12}e_1t_4e_4 & +a_{13}e_1t_2e_2 & +a_{14}e_1t_4t_2e_3 \\ +a_{21}e_4t_4e_1 & +a_{22}e_4 & +a_{23}e_4t_4t_2e_2 & +a_{24}e_4t_2e_3 \\ +a_{31}e_2t_2e_1 & +a_{32}e_2t_4t_2e_4 & +a_{33}e_2 & +a_{34}e_2t_4e_3 \\ +a_{41}e_3t_4t_2e_1 & +a_{42}e_3t_2e_4 & +a_{43}e_3t_4e_2 & +a_{44}e_3 \end{array}$$

is an isomorphism of algebras.

(6) Since t_4 commutes with t_1, t_2, t_3 in C_{vi} , we have

$$\begin{aligned} C_{\text{vi}} &\cong k\langle t_1, t_2, t_3 \rangle / (t_1 t_2 + t_2 t_1, t_1 t_3 + t_3 t_1, t_2 t_3 - t_3 t_2, t_1^2 - 1, t_2^2 - 1, t_3^2 - 1) \otimes_k k[t_4] / (t_4^2 - 1) \\ &\cong M_2(k)^2 \otimes_k k^2 \cong M_2(k)^4 \end{aligned}$$

by (1).

(7) Since t_4 commutes with t_1, t_2, t_3 in C_{vii} , we have

$$\begin{aligned} C_{\text{vii}} &\cong k\langle t_1, t_2, t_3 \rangle / (t_1 t_2 + t_2 t_1, t_1 t_3 - t_3 t_1, t_2 t_3 - t_3 t_2, t_1^2 - 1, t_2^2 - 1, t_3^2 - 1) \otimes_k k[t_4] / (t_4^2 - 1) \\ &\cong M_2(k)^2 \otimes_k k^2 \cong M_2(k)^4 \end{aligned}$$

by (2). □

Theorem 3.9. (1) *If $n = 4$, then*

$$\begin{aligned} E \cong \mathbb{P}^3 \text{ or } \mathbb{P}(1, 2, 4) \cup \mathbb{P}(1, 2, 3) \cup \mathbb{P}(3, 4) &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^2), \\ E = \mathbb{P}(3, 4) \cup \mathbb{P}(2, 4) \cup \mathbb{P}(2, 3) \cup \mathbb{P}(1, 4) \cup \mathbb{P}(1, 3) \cup \mathbb{P}(1, 2) &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^8). \end{aligned}$$

(2) *If $n = 5$, then*

$$\begin{aligned} E \cong (5a), (5c), \text{ or } (5d) &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k), \\ E \cong (5b), (5e), \text{ or } (5f) &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^4), \\ E = (5g) &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^{16}), \end{aligned}$$

where

- (5a) \mathbb{P}^4
- (5b) $\mathbb{P}(1, 2, 3, 5) \cup \mathbb{P}(1, 2, 3, 4) \cup \mathbb{P}(4, 5)$
- (5c) $\mathbb{P}(1, 2, 3, 4) \cup \mathbb{P}(3, 4, 5) \cup \mathbb{P}(1, 2, 5)$
- (5d) $\mathbb{P}(3, 4, 5) \cup \mathbb{P}(1, 4, 5) \cup \mathbb{P}(1, 2, 5) \cup \mathbb{P}(1, 2, 3) \cup \mathbb{P}(2, 3, 4)$
- (5e) $\mathbb{P}(1, 3, 5) \cup \mathbb{P}(1, 3, 4) \cup \mathbb{P}(1, 2, 5) \cup \mathbb{P}(1, 2, 4) \cup \mathbb{P}(4, 5) \cup \mathbb{P}(2, 3)$
- (5f) $\mathbb{P}(1, 2, 5) \cup \mathbb{P}(1, 2, 4) \cup \mathbb{P}(1, 2, 3) \cup \mathbb{P}(4, 5) \cup \mathbb{P}(3, 5) \cup \mathbb{P}(3, 4)$
- (5g) $\mathbb{P}(4, 5) \cup \mathbb{P}(3, 5) \cup \mathbb{P}(3, 4) \cup \mathbb{P}(2, 5) \cup \mathbb{P}(2, 4) \cup \mathbb{P}(2, 3) \cup \mathbb{P}(1, 5) \cup \mathbb{P}(1, 4) \cup \mathbb{P}(1, 3) \cup \mathbb{P}(1, 2)$.

Proof. (1) By Lemma 3.5, there exists a graded (± 1) -skew polynomial algebra S' such that $A \cong S'/(f)$ and the point scheme E' of S' is $\mathbb{P}^3, \mathbb{P}(1, 2, 4) \cup \mathbb{P}(1, 2, 3) \cup \mathbb{P}(3, 4)$, or $\bigcup_{1 \leq i < j \leq 4} \mathbb{P}(i, j)$. (Note that $E \cong E'$.) By Theorem 3.3 (2), (3), we only consider the case $E' = \mathbb{P}(1, 2, 4) \cup \mathbb{P}(1, 2, 3) \cup \mathbb{P}(3, 4)$. In this case,

$$\varepsilon_{41}\varepsilon_{12}\varepsilon_{24} = 1, \varepsilon_{41}\varepsilon_{13}\varepsilon_{34} = -1, \varepsilon_{42}\varepsilon_{23}\varepsilon_{34} = -1$$

(see (4iii) in the proof of Lemma 3.5), so $C(S'/(f))$ is isomorphic to

$$k\langle t_1, t_2, t_3 \rangle / (t_1 t_2 + t_2 t_1, t_1 t_3 - t_3 t_1, t_2 t_3 - t_3 t_2, t_i^2 - 1) \cong M_2(k)^2$$

by Lemma 3.8 (2). Thus we have $\underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \underline{\mathbf{CM}}^{\mathbb{Z}}(S'/(f)) \cong \mathbf{D}^b(\text{mod } k^2)$ by Theorem 2.1.

(2) By Lemma 3.7, there exists a graded (± 1) -skew polynomial algebra S' such that $A \cong S'/(f)$ and the point scheme E' of S' is (5a), \dots , (5f), or (5g). By Theorem 3.3 (1), (3), we only consider the cases (5b) to (5f).

If E is (5b), then

$$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = 1, \varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = 1, \varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = -1, \varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = 1, \varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = -1, \varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = -1,$$

(see (5v) in the proof of Lemma 3.7), so $C(S'/(f))$ is isomorphic to

$$k\langle t_1, t_2, t_3, t_4 \rangle / (t_1 t_2 + t_2 t_1, t_1 t_3 + t_3 t_1, t_1 t_4 - t_4 t_1, t_2 t_3 + t_3 t_2, t_2 t_4 - t_4 t_2, t_3 t_4 - t_4 t_3, t_i^2 - 1) \cong M_2(k)^4$$

by Lemma 3.8 (3). Thus we have $\underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \underline{\mathbf{CM}}^{\mathbb{Z}}(S'/(f)) \cong \mathbf{D}^b(\text{mod } k^4)$ by Theorem 2.1.

If E is (5c), then

$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = 1$, $\varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = -1$, $\varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = -1$, $\varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = -1$, $\varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = -1$, $\varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = 1$,
(see (5viii) in the proof of Lemma 3.7), so $C(S'/(f))$ is isomorphic to

$$k\langle t_1, t_2, t_3, t_4 \rangle / (t_1t_2 + t_2t_1, t_1t_3 - t_3t_1, t_1t_4 - t_4t_1, t_2t_3 - t_3t_2, t_2t_4 - t_4t_2, t_3t_4 + t_4t_3, t_i^2 - 1) \cong M_4(k)$$

by Lemma 3.8 (4). Thus we have $\underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \underline{\mathbf{CM}}^{\mathbb{Z}}(S'/(f)) \cong \mathbf{D}^b(\text{mod } k)$ by Theorem 2.1.

If E is (5d), then

$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = 1$, $\varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = -1$, $\varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = 1$, $\varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = -1$, $\varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = -1$, $\varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = 1$,
(see (5vi) in the proof of Lemma 3.7), so $C(S'/(f))$ is isomorphic to

$$k\langle t_1, t_2, t_3, t_4 \rangle / (t_1t_2 + t_2t_1, t_1t_3 - t_3t_1, t_1t_4 + t_4t_1, t_2t_3 - t_3t_2, t_2t_4 - t_4t_2, t_3t_4 + t_4t_3, t_i^2 - 1) \cong M_4(k)$$

by Lemma 3.8 (5). Thus we have $\underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \underline{\mathbf{CM}}^{\mathbb{Z}}(S'/(f)) \cong \mathbf{D}^b(\text{mod } k)$ by Theorem 2.1.

If E is (5e), then

$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = 1$, $\varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = 1$, $\varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = -1$, $\varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = -1$, $\varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = -1$, $\varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = -1$,
(see (5ix) in the proof of Lemma 3.7), so $C(S'/(f))$ is isomorphic to

$$k\langle t_1, t_2, t_3, t_4 \rangle / (t_1t_2 + t_2t_1, t_1t_3 + t_3t_1, t_1t_4 - t_4t_1, t_2t_3 - t_3t_2, t_2t_4 - t_4t_2, t_3t_4 - t_4t_3, t_i^2 - 1) \cong M_2(k)^4$$

by Lemma 3.8 (6). Thus we have $\underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \underline{\mathbf{CM}}^{\mathbb{Z}}(S'/(f)) \cong \mathbf{D}^b(\text{mod } k^4)$ by Theorem 2.1.

If E is (5f), then

$\varepsilon_{51}\varepsilon_{12}\varepsilon_{25} = 1$, $\varepsilon_{51}\varepsilon_{13}\varepsilon_{35} = -1$, $\varepsilon_{51}\varepsilon_{14}\varepsilon_{45} = -1$, $\varepsilon_{52}\varepsilon_{23}\varepsilon_{35} = -1$, $\varepsilon_{52}\varepsilon_{24}\varepsilon_{45} = -1$, $\varepsilon_{53}\varepsilon_{34}\varepsilon_{45} = -1$,
(see (5x) in the proof of Lemma 3.7), so $C(S'/(f))$ is isomorphic to

$$k\langle t_1, t_2, t_3, t_4 \rangle / (t_1t_2 + t_2t_1, t_1t_3 - t_3t_1, t_1t_4 - t_4t_1, t_2t_3 - t_3t_2, t_2t_4 - t_4t_2, t_3t_4 - t_4t_3, t_i^2 - 1) \cong M_2(k)^4$$

by Lemma 3.8 (7). Thus we have $\underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \underline{\mathbf{CM}}^{\mathbb{Z}}(S'/(f)) \cong \mathbf{D}^b(\text{mod } k^4)$ by Theorem 2.1. \square

Let ℓ denote the number of irreducible components of E which are isomorphic to \mathbb{P}^1 , that is, the number of irreducible components of the form $\mathbb{P}(i, j)$. Corollary 3.4 and Theorem 3.9 imply the following result which states that Conjecture 1.3 is true for $n \leq 5$.

Theorem 3.10. *Assume that $n \leq 5$.*

(1) *If n is odd, then $\ell \leq 10$ and*

$$\begin{aligned} \ell = 0 &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k), \\ 0 < \ell \leq 3 &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^4), \\ 3 < \ell \leq 10 &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^{16}). \end{aligned}$$

(2) *If n is even, then $\ell \leq 6$ and*

$$\begin{aligned} 0 \leq \ell \leq 1 &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^2), \\ 1 < \ell \leq 6 &\iff \underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^8). \end{aligned}$$

At the end of paper, we collect some examples when $n = 6$ as further evidence for Conjecture 1.3.

Example 3.11. (1) Let $S = k\langle x_1, \dots, x_6 \rangle / (x_i x_j - \varepsilon_{ij} x_j x_i)$ with

$$\begin{aligned} \varepsilon_{12} = 1, \varepsilon_{13} = -1, \varepsilon_{14} = 1, \varepsilon_{15} = -1, \varepsilon_{16} = 1, \varepsilon_{23} = -1, \varepsilon_{24} = -1, \varepsilon_{25} = -1, \varepsilon_{26} = 1, \\ \varepsilon_{34} = 1, \varepsilon_{35} = -1, \varepsilon_{36} = 1, \varepsilon_{45} = -1, \varepsilon_{46} = 1, \varepsilon_{56} = 1. \end{aligned}$$

Then the point scheme of S is $\mathbb{P}(3, 4, 5) \cup \mathbb{P}(2, 3, 4) \cup \mathbb{P}(1, 4, 5) \cup \mathbb{P}(1, 2, 5) \cup \mathbb{P}(1, 2, 3) \cup \mathbb{P}(3, 4, 6) \cup \mathbb{P}(1, 4, 6) \cup \mathbb{P}(1, 2, 6) \cup \mathbb{P}(5, 6)$, so $\ell = 1$. On the other hand, one can check that $C(A) \cong M_4(k)^2$, so we have $\underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^2)$.

(2) Let $S = k\langle x_1, \dots, x_6 \rangle / (x_i x_j - \varepsilon_{ij} x_j x_i)$ with

$$\begin{aligned} \varepsilon_{12} = 1, \varepsilon_{13} = -1, \varepsilon_{14} = -1, \varepsilon_{15} = -1, \varepsilon_{16} = 1, \varepsilon_{23} = 1, \varepsilon_{24} = -1, \varepsilon_{25} = -1, \varepsilon_{26} = 1, \\ \varepsilon_{34} = -1, \varepsilon_{35} = -1, \varepsilon_{36} = 1, \varepsilon_{45} = 1, \varepsilon_{46} = 1, \varepsilon_{56} = 1. \end{aligned}$$

Then the point scheme of S is $\mathbb{P}(2, 3, 4, 5) \cup \mathbb{P}(1, 2, 4, 5) \cup \mathbb{P}(2, 3, 6) \cup \mathbb{P}(1, 2, 6) \cup \mathbb{P}(4, 5, 6) \cup \mathbb{P}(1, 3)$, so $\ell = 1$. On the other hand, one can check that $C(A) \cong M_4(k)^2$, so we have $\underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^2)$.

(3) Let $S = k\langle x_1, \dots, x_6 \rangle / (x_i x_j - \varepsilon_{ij} x_j x_i)$ with

$$\begin{aligned} \varepsilon_{12} = 1, \varepsilon_{13} = -1, \varepsilon_{14} = -1, \varepsilon_{15} = -1, \varepsilon_{16} = 1, \varepsilon_{23} = 1, \varepsilon_{24} = -1, \varepsilon_{25} = -1, \varepsilon_{26} = 1, \\ \varepsilon_{34} = -1, \varepsilon_{35} = -1, \varepsilon_{36} = 1, \varepsilon_{45} = -1, \varepsilon_{46} = 1, \varepsilon_{56} = 1. \end{aligned}$$

Then the point scheme of S is $\mathbb{P}(2, 3, 5) \cup \mathbb{P}(2, 3, 4) \cup \mathbb{P}(1, 2, 5) \cup \mathbb{P}(1, 2, 4) \cup \mathbb{P}(1, 2, 6) \cup \mathbb{P}(2, 3, 6) \cup \mathbb{P}(4, 5) \cup \mathbb{P}(1, 3) \cup \mathbb{P}(4, 6) \cup \mathbb{P}(5, 6)$, so $\ell = 4$. On the other hand, one can check that $C(A) \cong M_2(k)^8$, so we have $\underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^8)$.

(4) Let $S = k\langle x_1, \dots, x_6 \rangle / (x_i x_j - \varepsilon_{ij} x_j x_i)$ with

$$\begin{aligned} \varepsilon_{12} = 1, \varepsilon_{13} = -1, \varepsilon_{14} = -1, \varepsilon_{15} = -1, \varepsilon_{16} = 1, \varepsilon_{23} = -1, \varepsilon_{24} = -1, \varepsilon_{25} = -1, \varepsilon_{26} = 1, \\ \varepsilon_{34} = -1, \varepsilon_{35} = -1, \varepsilon_{36} = 1, \varepsilon_{45} = -1, \varepsilon_{46} = 1, \varepsilon_{56} = 1. \end{aligned}$$

Then the point scheme of S is $\mathbb{P}(1, 2, 5) \cup \mathbb{P}(1, 2, 4) \cup \mathbb{P}(1, 2, 3) \cup \mathbb{P}(1, 2, 6) \cup \mathbb{P}(4, 5) \cup \mathbb{P}(3, 5) \cup \mathbb{P}(3, 4) \cup \mathbb{P}(4, 6) \cup \mathbb{P}(3, 6) \cup \mathbb{P}(5, 6)$, so $\ell = 6$. On the other hand, one can check that $C(A) \cong M_2(k)^8$, so we have $\underline{\mathbf{CM}}^{\mathbb{Z}}(A) \cong \mathbf{D}^b(\text{mod } k^8)$.

ACKNOWLEDGMENT

The author thanks the referee for a careful reading of the manuscript and helpful comments.

REFERENCES

- [1] M. Artin, J. Tate, and M. Van den Bergh, Some algebras associated to automorphisms of elliptic curves, in *The Grothendieck Festschrift, vol. I*, Prog. Math., vol. 86, Birkhäuser, Boston, MA, 1990, pp. 33–85.
- [2] P. Belmans, K. De Laet, and L. Le Bruyn, The point variety of quantum polynomial rings, *J. Algebra* **463** (2016), 10–22.
- [3] R.-O. Buchweitz, D. Eisenbud, and J. Herzog, Cohen-Macaulay modules on quadrics, in *Singularities, representation of algebras, and vector bundles* (Lambrecht, 1985), Lecture Notes in Math., vol. 1273, Springer, Berlin, 1987, pp. 58–116.
- [4] P. Jørgensen, Local cohomology for non-commutative graded algebras, *Comm. Algebra* **25** (1997), no. 2, 575–591.
- [5] H. Knörrer, Cohen-Macaulay modules on hypersurface singularities I, *Invent. Math.* **88** (1987), 153–164.
- [6] I. Mori, Co-point modules over Koszul algebras, *J. London Math. Soc. (2)* **74** (2006), no. 3, 639–656.
- [7] S. P. Smith, Some finite-dimensional algebras related to elliptic curves, in *Representation theory of algebras and related topics* (Mexico City, 1994), CMS Conference Proceedings, vol. 19, Amer. Math. Soc., Providence, RI, 1996, pp. 315–348.
- [8] S. P. Smith and M. Van den Bergh, Noncommutative quadric surfaces, *J. Noncommut. Geom.* **7** (2013), no. 3, 817–856.
- [9] J. Vitoria, Equivalences for noncommutative projective spaces, preprint, [arXiv:1001.4400v3](https://arxiv.org/abs/1001.4400v3).

DEPARTMENT OF MATHEMATICS, FACULTY OF EDUCATION, HIROSAKI UNIVERSITY, 1 BUNKYOCHO, HIROSAKI, AOMORI 036-8560, JAPAN

E-mail address: k-ueyama@hirosaki-u.ac.jp