MODULATION IN TETRADIC HARMONY AND ITS ROLE IN JAZZ

OCTAVIO A. AGUSTÍN-AQUINO AND GUERINO MAZZOLA

ABSTRACT. After a quick exposition of Mazzola's quantum modulation model for the so-called triadic interpretation of the major scale within the equal temperament, we study the model for the tetradic interpretation of the same scale. It is known that tetrads are fundamental for jazz music, and some classical objects for this kind of music are recovered.

1. INTRODUCTION

Mazzola's quantum model of modulation has been successful to explain, for instance, the harmonic structure of the first movement of Beethoven's *Hammerklavier* sonata [11, Part VI, Chapter 28], and it also explains the importance of melodic, harmonic, and major diatonic scales as extremes of modulatory freedom [11, Part VI, Section 27.1.5]. Recently it has been used to guide the extension of traditional harmony to the 20-tone microtonal equal temperament ([5, 4]).

The notion of tonality in Western music is inextricably linked to the concept of *cadence*, which is a sequence of chords that has the intention of asserting tonality. Considering that chords are identified by their position within the scale with Roman numerals from the first (I) to the seventh (VII), prominent examples of cadences are the *authentic* V–I and the *plagal* IV–I cadences [8, Ch. 10] and Riemann's "grosse Cadenz" I–IV–I–V–I [13, p. 52].

Together with the concept of tonality, there is the one of *modulation*, that consists in the transit from a source tonality to a target tonality in a parsimonious way.

We now summarize Mazzola's theory of modulation ([11, Part VI, Chapter 27]) in order to explain how to extend it from the classical case of tonality with triads to tetrads or seventh chords, which are very important for jazz. First, we define an *scale* E as a subset of \mathbb{Z}_{12}

Date: 12 September 2018.

²⁰¹⁰ Mathematics Subject Classification. 00A65, 05E18.

Key words and phrases. harmony, modulation quanta, jazz, tetrads.

This work was partially supported by a grant from the Niels Hendrik Abel Board.

where k = |E| is its cardinality. A triadic interpretation of the scale E (or tonality of E), denoted by $E^{(3)}$, is a set of specific 3-subsets of E such that they cover E.

Each member of the triadic interpretation is called a *degree*, and takes its name according its position within the transversal in the natural order of E inherited from \mathbb{Z}_{12} , and it is usually denoted with a Roman numeral. This general definition is usually further restricted to take into account standard musical practice. More specifically, if $E = \{x_i\}_{i=0}^{k-1}$ is the indexing of E following its natural order, then the triadic interpretation

$$E^{(3)} = \{\{x_i, x_{i+2 \mod k}, x_{i+4 \mod k}\}\}_{i=0}^{k-1}$$

corresponds to stacked thirds on top of each member of the scale. This is the interpretation we will use from now on.

Example 1.1. The subset $C = \{0, 2, 4, 5, 7, 9, 11\} \subseteq \mathbb{Z}_{12}$ is the C major scale, and thus its triadic interpretation is

$$\{\{0, 4, 7\}, \{2, 5, 9\}, \{4, 7, 11\}, \{7, 11, 2\}, \{9, 0, 4\}, \{11, 2, 5\}\}.$$

For example, $I_C = \{0, 4, 7\}$, $II_C = \{2, 5, 9\}$, and so on.

A transposition in \mathbb{Z}_{12} is the map

$$T^a: \mathbb{Z}_{12} \to \mathbb{Z}_{12}: x \mapsto x + a,$$

and, in general, a *special affine symmetry* is a map

$$T^a.v: \mathbb{Z}_{12} \to \mathbb{Z}_{12}: x \mapsto vx + a$$

where $v \in \{1, -1\}$.

From this moment on, we will consider only the orbit of the C scale with respect to the group of transpositions extended pointwise to sets, and that we will call $Dia^{(3)}$. We have

$$Dia^{(3)} = \{T^a(C^{(3)}) : a \in \mathbb{Z}_{12}\}.$$

Remark 1. Observe that actually

$$Dia^{(3)} = \{g(C^{(3)}) : g \in T^{\mathbb{Z}_{12}}.\{\pm 1\}\}$$

because $C^{(3)}$ is invariant with respect to the inversion symmetry T^4 . -1.

2. Cadences

Definition 2.1. Given $Dia^{(3)}$, a set P of cadence parameters, and a map $\kappa : Dia^{(3)} \to P$, we say that κ is *cadential* in a tonality $E^{(3)} \in Dia^{(3)}$ if the fiber $\kappa^{-1}(\kappa(E^{(3)}))$ is the singleton $\{E^{(3)}\}$, i. e., κ is injective in $E^{(3)}$. The map κ is a *cadence* if it is cadential in every tonality $E^{(3)} \in Dia^{(3)}$. We will take P as covers of subsets of \mathbb{Z}_{12} by degrees¹ of $E^{(3)}$. Thus a cadence is completely determined by the indices of the degrees of its image, and these are called *cadential sets*. A cadential set is *minimal* if it does not have proper cadential subset.

For the case of $Dia^{(3)}$, in [10] the following minimal cadences are calculated

$$J_1 = \{II, III\}, J_2 = \{II, V\}, J_3 = \{III, IV\}, J_4 = \{IV, V\}, J_5 = \{VII\}.$$

Thus cadential sets are minimal amounts of information from which we can infer the tonality of a given musical fragment.

3. MODULATION

Arnold Schönberg, in his *Harmonielehre*, divides the process of modulation in three stages [14, p. 186]:

- (1) [...] introduction of such (neutral) chords as will permit the turn to the new tonality [...].
- (2) the actual modulatory part, that is, the modulatory chords [...];
- (3) the consolidation, with the cadence establishing the target tonality.

The chords mentioned in the second stage are the *pivots* (Wendepunkte), and a great deal of the modulation problem comes from finding them. Schönberg's definition is the point of departure for Mazzola's quantum modulation model, describing the connection between the source and the target tonalities through an affine symmetry. From this symmetry and the cadence that consolidates the new tonality, it defines a modulation *quantum*, analogous to the ones that carry the fundamental forces of nature and which are also are governed by symmetries.

Definition 3.1. A modulation between two tonalities $S^{(3)}$ and $T^{(3)}$ is a pair (g, μ) , where $g: S \to T$ is an special affine symmetry (called modulator) that induces an isomorphism between $S^{(3)}$ and $T^{(3)}$, and where μ is a minimal cadential set of $T^{(3)}$.

It can be proved that $g \in T^{\mathbb{Z}_{12}}$. $\{-1, 1\}$, i. e., it is an element of the special affine group of \mathbb{Z}_{12} .

Definition 3.2. A modulation quantum $Q \subseteq \mathbb{Z}_{12}$ of a modulation (g, μ) satisfies the following:

¹This is an special case of the concept of *global composition* [11, Section 13.2].



identification of two sides AB and BA of a rectangle.

- (1) the modulator g is a symmetry of Q, i. e., g(Q) = Q.
- (2) the set $T \cap Q$ is rigid (i. e., its symmetry group is trivial). The triads of the target tonality contained in this set are called the pivots.
- (3) The quantum Q is the smallest set with the aforementioned properties.

It is very remarkable that the model recuperates the pivots proposed by Schönberg in his treatise [11, p. 574].

4. The triadic nerve of the major scale

The nerve [1, pp. 128-129] of $E^{(3)}$ is the simplicial complex of $E^{(3)}$ where every degree is a point and a set of points define a simplex if their intersection is non-empty. Its geometrical realization (a topological space) turns out to be a Möbius strip (Figure 1), a fact discovered by Mazzola ([10]) and implied from Schönberg's concept of *Harmonisches Band* [14, p. 44]. Furthermore, it is the triangulation with the least number of vertices of the aforementioned space [2, p. 32]. This topological construction has consequences for the function theory proposed by Hugo Riemann, where he assigns to each cord its *parallel* degree (Parallelklang) and its *counterparallel* (Leittonwechselklang), globally [7, pp. 87-88]. The unorientability of the Möbius strip makes such a global assignment impossible [11, p. 324].

5. The tetradic case

In what follows, we extend Mazzola's model to seventh chords, that are tetrads of tones separated by thirds. More specifically,

$$E^{(4)} = \{\{x_i, x_{i+2 \mod 7}, x_{i+4 \mod 7}, x_{i+6 \mod 7}\}\}_{i=0}^6,$$

which is crucial for jazz music [12, p. 11,22]. Cadential sets are, in this case,

$$J_1 = \{I^7, II^7\}, J_2 = \{I^7, IV^7\}, J_3 = \{II^7, III^7\}, J_4 = \{III^7, IV^7\}, J_5 = \{V^7\}, J_6 = \{VII^7\}.$$

Remark 2. The set of chords $\{II^7, V^7\}$ is cadential, although not minimal. It is frequently used in jazz, like in *Giant Steps* by John Coltrane [12, p. 42].

With the following algorithm (which is described in more detail in [11, p. 573]) we can calculate all quanta.

Algorithm 1.

- **Input:** A modulation (g, μ) from the source tonality $E^{(3)}$ to the target tonality $F^{(3)}$.
- **Output:** The quantum Q of the modulation (g, μ) if the modulation is quantized; a message of failure otherwise.
 - 1: $G := \langle g \rangle, Q := \emptyset.$
 - 2: for all $h \in G$ do
 - 3: $Q := Q \cup g(\bigcup \mu).$
- 4: $M := Q \cap F$.
- 5: quantized := true.
- 6: for all $h \in T^{\mathbb{Z}_{12}}$. $\{-1,1\} \setminus \{1\}$ do
- 7: **if** h(M) = M **then**
- 8: quantized := false.
- 9: if quantized then
- 10: return Q.

11: else

12: **print** The modulation is not quantized.

The output for each quantized modulation is shown in Table 1 and leads us to the following insights.

- (1) All quantized modulations for the tetradic interpretations stem from the $\{V\}$ and $\{VII\}$ cadences.
- (2) In the theory as formulated in [11], it is required that $T \cap Q$ to be covered by triads, but this is not required in the original definition of [10]. In fact, the pivots marked with an asterisk in Table 1 are not enough to cover $T \cap Q$.
- (3) The modulation quanta of modulations

 $(T^{\{3,5\}}.11, \{V^7\})$ and $(T^{\{3,9\}}, \{V^7\})$

Tr.	Cadence	Quantum	Modulator	Pivots
1	$\{V^7\}$	$\{0, 2, 3, 5, 6, 8, 9, 11\}$	$T^{5}.11$	$\{III^7, V^7\}$
1	$\{VII^7\}$	$\{0, 2, 3, 5, 6, 7, 10, 11\}$	$T^{5}.11$	${VII^{7}}^{*}$
2	$\{V^7\}$	$\{1, 2, 4, 5, 7, 9, 11\}$	$T^{6}.11$	$\{II^7, V^7, VII^7\}$
2	$\{VII^7\}$	$\{1, 2, 4, 5, 7, 11\}$	$T^{6}.11$	$\{II^7, VII^7\}$
3	$\{V^7\}$	$\{1, 2, 4, 5, 7, 8, 10, 11\}$	T^3	$\{III^7, V^7\}$
3	$\{V^7\}$	$\{2, 5, 8, 9, 10, 11\}$	$T^{7}.11$	$\{V^7\}$
3	$\{VII^7\}$	$\{0, 2, 5, 7, 8, 11\}$	$T^{7}.11$	${VII^7}*$
4	$\{V^7\}$	$\{2, 3, 5, 6, 9, 11\}$	$T^{8}.11$	$\{V^7\}$
5	$\{VII^7\}$	$\{2, 4, 5, 7, 10, 11\}$	$T^{9}.11$	$\{II^7, VII^7\}$
6	$\{V^7\}$	$\{1, 2, 5, 7, 8, 11\}$	T^6	$\{V^7\}$
6	$\{V^7\}$	$\{1, 2, 5, 8, 9, 11\}$	$T^{10}.11$	$\{V^7\}$
6	$\{VII^7\}$	$\{2, 3, 5, 8, 9, 11\}$	T^6	$\{VII^7\}$
6	$\{VII^7\}$	$\{2, 3, 5, 7, 8, 11\}$	$T^{10}.11$	$\{VII^7\}$
7	$\{V^7\}$	$\{0, 2, 5, 6, 9, 11\}$	$T^{11}.11$	$\{III^7, V^7\}$
8	$\{VII^7\}$	$\{1, 2, 5, 7, 10, 11\}$	$T^{0}.11$	$\{VII^7\}$
9	$\{V^7\}$	$\{1, 2, 4, 5, 7, 8, 10, 11\}$	T^9	$\{III^7, V^7\}$
9	$\{V^7\}$	$\{2, 4, 5, 8, 9, 11\}$	$T^{1}.11$	$\{V^{7}\}*$
9	$\{VII^7\}$	$\{2, 5, 6, 7, 8, 11\}$	$T^{1}.11$	$\{VII^7\}$
10	$\{V^7\}$	$\{0, 2, 3, 5, 9, 11\}$	$T^{2}.11$	$\{III^7, V^7\}$
10	$\{\overline{VII^7}\}$	$\{0, \overline{2, 3, 5, 7, 9, 11}\}$	$T^{2}.11$	$\{III^7, V^7, VII^7\}$
11	$\{V^7\}$	$\{1, 2, 4, 5, 6, 9, 10, 11\}$	$T^{3}.11$	$\{V^{7}\}^{*}$
11	$\{VII^7\}$	$\{1, 2, 4, 5, 7, 8, 10, 11\}$	$T^{3}.11$	$\{II^7, VII^7\}$

TABLE 1. Summary of quantized modulations for the tetradic tonalities over the major scale.

are what in jazz are known as *diminished scales*, which are 8tone scales in which the notes ascend in alternating intervals of a whole step and a half step [12, p. 31].

- (4) Modulations with cadences {V⁷} and {VII⁷} to a distance of tritone correspond to the jazz artifice of tritone substitution [9, p. 147]. More specifically, the pivot of the target tonality is the tritone substitution of the same degree of the source tonality.
- (5) A particular case of a quick modulation towards a tonality a major second upwards known as *chaining* [12, p. 46] is recovered as a quantized modulation for cadence $\{V^7\}$ with pivots $\{II^7, V^7\}$.

 $\overline{7}$

6. The tetradic nerve of the major scale

If we calculate the nerve of E with the tetradic cover $E^{(4)}$, we see that all degrees intersect in at least one point, thus the 1-skeleton of such a nerve is the complete graph K_7 . This is a toroidal graph [3, p. 71], and its embedding in such a surface (Figure 2) is the simplicial triangulation with the smallest number of vertices [6, p. 277]. Moreover, there are seven 3-dimensional simplices or tetrahedra

$$S_{1} = \{I^{7}, II^{7}, IV^{7}, VI^{7}\}, S_{2} = \{II^{7}, III^{7}, V^{7}, VII^{7}\}, S_{3} = \{III^{7}, IV^{7}, VI^{7}, I^{7}\}, S_{4} = \{IV^{7}, V^{7}, VII^{7}, II^{7}\}, S_{5} = \{V^{7}, VI^{7}, I^{7}, III^{7}\}, S_{6} = \{VI^{7}, VII^{7}, II^{7}, IV^{7}\}, S_{7} = \{VII^{7}, I^{7}, III^{7}, V^{7}\}, S_{7} = \{VII^{7}, V^{7}, V^{7}\}, S_{7} = \{VII^{7}, V^{$$

which can be arranged in a cycle $S_5, S_7, S_2, S_4, S_6, S_1, S_3, S_5$ that forms a solid torus. Now, even although K_7 is embeddable in an orientable surface, it is not possible to formulate a Riemann global theory of chord functions: the obstacle is the fact that the 1-skeleton graph of the nerve contains the cycle $I^7, V^7, IV^7, VI^7, III^7, VII^7, II^7, I^7$, which makes impossible to define paralellism in a consistent way.

Acknowledgment

We thank Jesús David Gómez Téllez for detecting some mistakes in a preliminary version of this article and suggesting some improvements.

References

- [1] P. S. Aleksandrov. Combinatorial Topology, volume 1. Graylock Press, 1956.
- [2] F. H. Croom. Basic Concepts of Algebraic Topology. Springer, 1978.
- [3] A. Gibbons. Algorithmic Graph Theory. Cambridge University Press, 1985.
- [4] J. D. Gómez Téllez. La teoría de números y la teoría de grupos en el estudio de las escalas musicales microtonales. Tesis de licenciatura, Unam, 1999.
- [5] J. D. Gómez-Téllez, E. Lluis-Puebla, and M. Montiel. A symmetric quantum theory of modulation in Z₂₀. In O. A. Agustín-Aquino, E. Lluis-Puebla, and M. Montiel, editors, *Mathematics and Computation in Music*, pages 50–62. Springer, 2017.
- [6] I. Kantor, J. Matoušek, and R. Šámal. *Mathematics++*. American Mathematical Society, 2015.
- [7] D. Knopp. Chromatic Transformations in Nineteenth-Century Music, volume 17 of Cambridge Studies in Music Theory and Analysis. Cambridge University Press, 2006.
- [8] S. Kostka and D. Payne. Tonal Harmony. McGraw-Hill, 4th edition, 2000.
- [9] D. Latarski. Practical Theory for Guitar. Alfred Music Publishing, 1993.
- [10] G. Mazzola. Gruppen und Kategorien in der Musik. Heldermann, Berlin, 1985.
- [11] G. Mazzola. The Topos of Music. Birkhäuser, Basel, 2002.



FIGURE 2. Construction of the embedding of K_7 in the torus by identifications of the opposite sides of a rectangle.

- [12] R. Rawlins and N. Bahha. Jazzology: The Encyclopedia of Jazz Theory for All Musicians. Hal Leonard Corporation, 2005.
- [13] H. Riemann. Ueber das musikalische Hören. PhD thesis, Universtät Göttingen, 1874.
- [14] A. Schönberg. Harmonielehre. Universal-Edition, 3rd edition, 1922.

INSTITUTO DE FÍSICA Y MATEMÁTICAS, UNIVERSIDAD TECNOLÓGICA DE LA MIXTECA, HUAJUAPAN DE LEÓN, OAXACA, MÉXICO *E-mail address*: octavioalberto@mixteco.utm.mx

School of Music, University of Minnesota, Minneapolis, MN, USA *E-mail address*: mazzola@umn.edu

8