

SL(3, \mathbb{C})-character varieties of two-generator groups

Haimiao Chen *

Beijing Technology and Business University, Beijing, China

Abstract

We present an efficient method for computing the irreducible SL(3, \mathbb{C})-character varieties of two-generator groups.

Keywords: SL(3, \mathbb{C})-character variety; irreducible representation; two-generator group; double twist link; symmetric slice; asymmetric slice

MSC2020: 15A24, 57K31

1 Introduction

Given a finitely generated group Γ and a linear group $G \leq \mathrm{GL}(n, \mathbb{C})$, a G -representation of Γ is simply a homomorphism $\rho : \Gamma \rightarrow G$, and its *character* is the function $\chi_\rho : \Gamma \rightarrow \mathbb{C}$ sending x to $\mathrm{tr}(\rho(x))$. The G -representation variety of Γ is $\mathcal{R}_G(\Gamma) := \mathrm{hom}(\Gamma, G)$, on which G acts via conjugation, and the G -character variety $\mathcal{X}_G(\Gamma)$ is defined as the GIT quotient $\mathcal{R}_G(\Gamma)//G$. Let $\mathcal{R}_G^{\mathrm{irr}}(\Gamma)$ denote the open subset of $\mathcal{R}_G(\Gamma)$ consisting of irreducible representations, and let

$$\mathcal{X}_G^{\mathrm{irr}}(\Gamma) = \{\chi_\rho : \rho \in \mathcal{R}_G^{\mathrm{irr}}(\Gamma)\}.$$

It is known that $\mathcal{X}_G^{\mathrm{irr}}(\Gamma)$ can be identified with an open subset of $\mathcal{X}_G(\Gamma)$, and the map $\rho \mapsto \chi_\rho$ factors through to give an isomorphism $\mathcal{R}_G^{\mathrm{irr}}(\Gamma)/G \cong \mathcal{X}_G^{\mathrm{irr}}(\Gamma)$. Passing to character variety removes some redundancy; however, in some situations, it is necessary to work with representations. For more knowledge, see [18] and the references therein.

Since the seminal works [3, 19], character variety has been playing a significant role in low-dimensional topology, in that, much geometric and topological information on 3-manifolds M is carried by $\mathcal{X}_G(\pi_1(M))$. For instance, CR and complex hyperbolic structure involve $\mathrm{SU}(2, 1) \leq \mathrm{SL}(3, \mathbb{C})$ [4, 15, 16].

When $n \geq 3$, as pointed out by [8], a main concern is to collect a sufficient amount of examples which will give directions for further research. However, in general it seems to be difficult to do explicit computations. Even for $n = 3$, till now few results on representation/character varieties have been seen in the literature, except for [5–7, 9–11, 13, 14].

In this paper, we present a method to determine $\mathcal{X}_{\mathrm{SL}(3, \mathbb{C})}^{\mathrm{irr}}(\Gamma)$ for any two-generator group Γ , (the reducible part of $\mathcal{X}_{\mathrm{SL}(3, \mathbb{C})}(\Gamma)$ is, as usual, relatively easy to study). A representation $\Gamma \rightarrow \mathrm{SL}(3, \mathbb{C})$ is the same as a pair (\mathbf{a}, \mathbf{b}) satisfying

*Email: chenhm@math.pku.edu.cn

some equations $\mathbf{s}_j = \mathbf{0}$, each of which is equivalent to $\text{tr}(\mathbf{u}\mathbf{s}_j) = 0$ for all \mathbf{u} in the vector space \mathcal{M} of 3×3 matrices. We provide bases (in Theorem 3.6 and 3.11) in terms of \mathbf{a}, \mathbf{b} for \mathcal{M} , so as to reduce $\text{tr}(\mathbf{u}\mathbf{s}_j) = 0$ to equations of traces of words in \mathbf{a}, \mathbf{b} .

A key ingredient (see Theorem 3.8) is the observation that, $\text{tr}(\mathbf{a}\mathbf{b}\mathbf{a}^{-1}\mathbf{b}^{-1}) = \text{tr}(\mathbf{b}\mathbf{a}\mathbf{b}^{-1}\mathbf{a}^{-1})$ if and only if \mathbf{a}, \mathbf{b} can be simultaneously conjugated to be symmetric. A remarkable result of Lawton [12] describes $\mathcal{X}_{\text{SL}(3, \mathbb{C})}(F_2)$ as a hypersurface in \mathbb{C}^9 which is a double branched cover of \mathbb{C}^8 , and the branching locus exactly consists of the characters with $\text{tr}(\mathbf{a}\mathbf{b}\mathbf{a}^{-1}\mathbf{b}^{-1}) = \text{tr}(\mathbf{b}\mathbf{a}\mathbf{b}^{-1}\mathbf{a}^{-1})$. Our result gives a geometric interpretation for these characters. When $\text{tr}(\mathbf{a}\mathbf{b}\mathbf{a}^{-1}\mathbf{b}^{-1}) \neq \text{tr}(\mathbf{b}\mathbf{a}\mathbf{b}^{-1}\mathbf{a}^{-1})$, the situation turns out to be largely different, and the computation is much simpler.

As an application, the $\text{SL}(3, \mathbb{C})$ -character variety for each double twist link group is determined.

We expect that, results of this paper will promote the study of $\text{SL}(3, \mathbb{C})$ -character varieties for general groups in future.

Notation 1.1. For elements x, y of some group, let $\bar{x} = x^{-1}$, $x \lrcorner y = xy\bar{x}$, $[x, y] = xy\bar{x}\bar{y}$; be careful that $\bar{xy} = x^{-1}y^{-1}$, not $(xy)^{-1}$. (In this paper, notations for conjugate complex numbers are never used.) For a word r in x, y , let \overleftarrow{r} or r^{\leftarrow} denote the one obtained by writing r backwards. Let $\text{Cen}(x)$ denote the centralizer of x .

Let $\omega = e^{2\pi i/3}$, and let $\langle \omega \rangle = \{1, \omega, \omega^2\}$.

Let \mathcal{M} denote the set of 3×3 matrices with entries in \mathbb{C} , which is a 9-dimensional vector space over \mathbb{C} ; let \mathcal{M}^0 and \mathcal{M}^1 respectively denote the subspace consisting of symmetric and skew-symmetric matrices. Let \mathbf{e} denote the identity matrix. For $\mathbf{z} \in \mathcal{M}$, let \mathbf{z}^{tr} denote its transpose; let $\mathbf{z}^{\neg} = [z_{12}, z_{13}, z_{23}]^{\text{tr}}$.

Given elements $\mathbf{a}_1, \dots, \mathbf{a}_k$ of some vector space over \mathbb{C} , let $\mathbb{C}\langle \mathbf{a}_1, \dots, \mathbf{a}_k \rangle$ denote the subspace they span.

2 On 3×3 unimodular matrices

For each $\mathbf{x} \in \text{SL}(3, \mathbb{C})$, by Cayley-Hamilton Theorem,

$$\mathbf{x}^3 - \text{tr}(\mathbf{x}) \cdot \mathbf{x}^2 + \text{tr}(\bar{\mathbf{x}}) \cdot \mathbf{x} - \mathbf{e} = 0. \quad (1)$$

Consequently, for each $k \in \mathbb{Z}$,

$$\mathbf{x}^k = \alpha_k(\mathbf{x}) \cdot \mathbf{x} + \beta_k(\mathbf{x}) \cdot \mathbf{e} + \gamma_k(\mathbf{x}) \cdot \bar{\mathbf{x}}, \quad (2)$$

with $\alpha_k(\mathbf{x}), \beta_k(\mathbf{x}), \gamma_k(\mathbf{x})$ polynomials in $\text{tr}(\mathbf{x}), \text{tr}(\bar{\mathbf{x}})$. In particular,

$$\mathbf{x}^3 = (\text{tr}(\mathbf{x})^2 - \text{tr}(\bar{\mathbf{x}})) \cdot \mathbf{x} + (1 - \text{tr}(\mathbf{x})\text{tr}(\bar{\mathbf{x}})) \cdot \mathbf{e} + \text{tr}(\mathbf{x}) \cdot \bar{\mathbf{x}}. \quad (3)$$

Introduce the *standard forms*

$$\mathbf{d}_{\lambda_3}^{\lambda_1} = \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix}, \quad \text{where } \lambda_2 = (\lambda_1 \lambda_3)^{-1};$$

$$\mathbf{h}_\lambda = \begin{bmatrix} 2\lambda & 0 & i\lambda \\ 0 & \lambda^{-2} & 0 \\ i\lambda & 0 & 0 \end{bmatrix};$$

$$\mu \mathbf{k}, \quad \text{where } \mu \in \langle \omega \rangle, \quad \mathbf{k} = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & i \\ 0 & i & 1 \end{bmatrix}.$$

Clearly, each element of $\text{SL}(3, \mathbb{C})$ can be conjugated into standard form.
Put

$$\mathbf{g} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad \mathbf{f} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & i \\ 0 & i & 0 \end{bmatrix}, \quad \mathbf{f}^2 = \begin{bmatrix} 1 & 0 & i \\ 0 & 0 & 0 \\ i & 0 & -1 \end{bmatrix}.$$

We have $\mathbf{g}\mathbf{f} = \mathbf{f}\mathbf{g} = \mathbf{f}^3 = \mathbf{0}$, and

$$\mathbf{k} = \mathbf{e} + \mathbf{f}, \quad \mathbf{h}_\lambda = \lambda \mathbf{e} + (\lambda^{-2} - \lambda)\mathbf{g} + \lambda \mathbf{f}^2.$$

Definition 2.1. An element $\mathbf{x} \in \text{SL}(3, \mathbb{C})$ is called *ordinary* if it satisfies any one of the following conditions which are equivalent to each other:

- 1) \mathbf{x} is neither conjugate to $\mathbf{d}_\lambda^\lambda$ for any λ , nor to \mathbf{h}_κ with $\kappa \in \langle \omega \rangle$;
- 2) the degree of the minimal polynomial of \mathbf{x} is 3;
- 3) each eigenspace of \mathbf{x} is one-dimensional.

Call \mathbf{x} *special* if it is not ordinary.

Lemma 2.2. *If $\mathbf{x} \in \text{SL}(3, \mathbb{C})$ is ordinary, then each element of $\text{Cen}(\mathbf{x})$ is a linear combination of $\mathbf{e}, \mathbf{x}, \bar{\mathbf{x}}$, and has a square root.*

Proof. It suffices to consider standard forms.

- If $\mathbf{x} = \mathbf{d}_{\lambda_3}^{\lambda_1}$ with $\lambda_1 \neq \lambda_2 \neq \lambda_3 \neq \lambda_1$, then

$$\text{Cen}(\mathbf{x}) = \{\mathbf{d}_{\mu_3}^{\mu_1} : \mu_1 \mu_3 \neq 0\} \subset \mathbb{C}\langle \mathbf{e}, \mathbf{x}, \bar{\mathbf{x}} \rangle;$$

clearly each element has a square root.

- If $\mathbf{x} = \mathbf{h}_\lambda$ with $\lambda^3 \neq 1$, then $\bar{\mathbf{x}} = \lambda^{-1}\mathbf{e} + (\lambda^2 - \lambda^{-1})\mathbf{g} - \lambda^{-1}\mathbf{f}^2$, and

$$\text{Cen}(\mathbf{x}) = \{\mathbf{h}_{t,c} := t\mathbf{e} + (t^{-2} - t)\mathbf{g} + c\mathbf{f}^2 : t \neq 0\} \subset \mathbb{C}\langle \mathbf{e}, \mathbf{x}, \bar{\mathbf{x}} \rangle;$$

each element has a square root, as $(\mathbf{h}_{t,c})^2 = \mathbf{h}_{t^2, 2tc}$.

- If $\mathbf{x} = \mu \mathbf{k}$ with $\mu^3 = 1$, then

$$\text{Cen}(\mathbf{x}) = \{a\mathbf{e} + b\mathbf{f} + c\mathbf{f}^2 : a \in \langle \omega \rangle\} \subset \mathbb{C}\langle \mathbf{e}, \mathbf{x}, \bar{\mathbf{x}} \rangle;$$

each element has a square root: $(a\mathbf{e} + b\mathbf{f} + c\mathbf{f}^2)^2 = a^2\mathbf{e} + 2ab\mathbf{f} + (b^2 + 2ac)\mathbf{f}^2$.

□

Let $F_2 = \langle z_1, z_2 \mid - \rangle$ denote the rank two free group. Given $\mathbf{a}, \mathbf{b} \in \mathrm{SL}(3, \mathbb{C})$, let $\sigma_{\mathbf{a}, \mathbf{b}} : F_2 \rightarrow \mathrm{SL}(3, \mathbb{C})$ send z_1, z_2 to \mathbf{a}, \mathbf{b} , respectively, and let $\chi_{\mathbf{a}, \mathbf{b}}$ denote its character. Each representation is of this form. We usually do not distinguish (\mathbf{a}, \mathbf{b}) and $\sigma_{\mathbf{a}, \mathbf{b}}$. Adopt the notations $t_{\overline{1}^2 2^3} = \mathrm{tr}(\overline{\mathbf{a}}^2 \mathbf{b}^3)$, $t_{2\overline{1}^3 \overline{2}1^4} = \mathrm{tr}(\overline{\mathbf{b}} \overline{\mathbf{a}}^3 \overline{\mathbf{b}} \mathbf{a}^4)$, and so forth; in particular,

$$\begin{aligned} t_1 &= \mathrm{tr}(\mathbf{a}), & t_2 &= \mathrm{tr}(\mathbf{b}), & t_{\overline{1}} &= \mathrm{tr}(\overline{\mathbf{a}}), & t_{\overline{2}} &= \mathrm{tr}(\overline{\mathbf{b}}), \\ t_{12} &= \mathrm{tr}(\mathbf{ab}), & t_{\overline{1}\overline{2}} &= \mathrm{tr}(\overline{\mathbf{a}}\overline{\mathbf{b}}), & t_{1\overline{2}} &= \mathrm{tr}(\mathbf{a}\overline{\mathbf{b}}), & t_{\overline{1}2} &= \mathrm{tr}(\overline{\mathbf{a}}\mathbf{b}), \\ & & t_{12\overline{1}\overline{2}} &= \mathrm{tr}([\mathbf{a}, \mathbf{b}]), & t_{2\overline{1}\overline{2}\overline{1}} &= \mathrm{tr}([\mathbf{b}, \mathbf{a}]). \end{aligned}$$

According to [12] Theorem 8, the $\mathrm{SL}(3, \mathbb{C})$ -character variety of F_2 can be embedded in \mathbb{C}^9 as a hypersurface:

$$\begin{aligned} \mathcal{X}_{\mathrm{SL}(3, \mathbb{C})}(F_2) &\cong \{(s_1, s_{\overline{1}}, s_2, s_{\overline{2}}, s_3, s_{\overline{3}}, s_4, s_{\overline{4}}, s_5) : s_5^2 - Ps_5 + Q = 0\}, \\ \chi_{\mathbf{a}, \mathbf{b}} &\mapsto (t_1, t_{\overline{1}}, t_2, t_{\overline{2}}, t_{12}, t_{\overline{1}\overline{2}}, t_{1\overline{2}}, t_{\overline{1}2}, t_{12\overline{1}\overline{2}}), \end{aligned} \quad (4)$$

for certain $P, Q \in \mathbb{C}[s_1, s_{\overline{1}}, s_2, s_{\overline{2}}, s_3, s_{\overline{3}}, s_4, s_{\overline{4}}]$, with $\deg P = 4, \deg Q = 6$. Geometrically, (4) presents $\mathcal{X}_{\mathrm{SL}(3, \mathbb{C})}(F_2)$ as a branched double cover of \mathbb{C}^8 , with branching locus defined by $P^2 - 4Q = 0$.

The precise meaning is: for each word \mathbf{w} in \mathbf{a}, \mathbf{b} , $\mathrm{tr}(\mathbf{w})$ can be written as a polynomial in $t_1, t_{\overline{1}}, t_2, t_{\overline{2}}, t_{12}, t_{\overline{1}\overline{2}}, t_{1\overline{2}}, t_{\overline{1}2}$ (see Proposition 2.4). Actually,

$$\begin{aligned} t_{12\overline{1}\overline{2}} + t_{2\overline{1}\overline{2}\overline{1}} &= P(t_1, t_{\overline{1}}, t_2, t_{\overline{2}}, t_{12}, t_{\overline{1}\overline{2}}, t_{1\overline{2}}, t_{\overline{1}2}), \\ t_{12\overline{1}\overline{2}} \cdot t_{2\overline{1}\overline{2}\overline{1}} &= Q(t_1, t_{\overline{1}}, t_2, t_{\overline{2}}, t_{12}, t_{\overline{1}\overline{2}}, t_{1\overline{2}}, t_{\overline{1}2}). \end{aligned}$$

Notation 2.3. For a word \mathbf{w} in \mathbf{a}, \mathbf{b} , let $\theta(\mathbf{w}) = \mathbf{w} + \overleftarrow{\mathbf{w}}$, and $\delta(\mathbf{w}) = \mathbf{w} - \overleftarrow{\mathbf{w}}$.

By [15] Corollary 4.3 (to Proposition 4.2),

$$\begin{aligned} \theta(\mathbf{a} \lrcorner \mathbf{b}) &= t_1 \theta(\overline{\mathbf{a}}\mathbf{b}) + t_{\overline{1}} \theta(\mathbf{a}\mathbf{b}) - (1 + t_1 t_{\overline{1}}) \mathbf{b} + (t_{\overline{1}\overline{2}} - t_{\overline{1}} t_2) \mathbf{a} + (t_{12} - t_1 t_2) \overline{\mathbf{a}} \\ &\quad + (t_1 t_{\overline{1}} t_2 + t_2 - t_1 t_{\overline{1}\overline{2}} - t_{\overline{1}} t_{12}) \mathbf{e}, \end{aligned} \quad (5)$$

$$\mathbf{a} \mathbf{b} \mathbf{a} = -\theta(\mathbf{a}^2 \mathbf{b}) + t_1 \theta(\mathbf{a}\mathbf{b}) + t_2 \mathbf{a}^2 + (t_{12} - t_1 t_2) \mathbf{a} - t_{\overline{1}} \mathbf{b} + t_{\overline{1}\overline{2}} \mathbf{e}. \quad (6)$$

Another identity which is useful in practical computation is

$$\mathbf{a} \mathbf{b} \mathbf{a} = t_{12} \mathbf{a} - t_{\overline{1}\overline{2}} \overline{\mathbf{b}} + \overline{\mathbf{b}} \overline{\mathbf{a}} \overline{\mathbf{b}}. \quad (7)$$

It is obtained by applying (1) to $\mathbf{a}\mathbf{b}$ and then multiplying on the right by $\overline{\mathbf{b}} \overline{\mathbf{a}} \overline{\mathbf{b}}$.

Proposition 2.4. *Each word in \mathbf{a}, \mathbf{b} can be written as a linear combination of $\mathbf{e}, \mathbf{a}, \overline{\mathbf{a}}, \mathbf{b}, \overline{\mathbf{b}}, \mathbf{ab}, \mathbf{ba}, \mathbf{a}\overline{\mathbf{b}}, \overline{\mathbf{b}}\mathbf{a}, \overline{\mathbf{a}}\mathbf{b}, \mathbf{b}\overline{\mathbf{a}}, \overline{\mathbf{a}}\overline{\mathbf{b}}, \overline{\mathbf{b}}\overline{\mathbf{a}}, \mathbf{a} \lrcorner \mathbf{b}, \mathbf{a} \lrcorner \overline{\mathbf{b}}, \mathbf{b} \lrcorner \mathbf{a}, \mathbf{b} \lrcorner \overline{\mathbf{a}}, [\mathbf{a}, \mathbf{b}], [\mathbf{b}, \mathbf{a}]$, with coefficients in $\mathbb{C}[t_1, t_{\overline{1}}, t_2, t_{\overline{2}}, t_{12}, t_{\overline{1}\overline{2}}, t_{1\overline{2}}, t_{\overline{1}2}, t_{12\overline{1}\overline{2}}]$.*

Proof. By (2), it is sufficient to consider the words of the form $\mathbf{w} = \mathbf{x}_1^{\epsilon_1} \cdots \mathbf{x}_n^{\epsilon_n}$ with each $\epsilon_i \in \{\pm 1\}$ and $\{\mathbf{x}_j, \mathbf{x}_{j+1}\} = \{\mathbf{a}, \mathbf{b}\}$ for $1 \leq j < n$.

Whenever any of $\mathbf{a}, \overline{\mathbf{a}}, \mathbf{b}, \overline{\mathbf{b}}$ appears twice, we can apply (6) to write \mathbf{w} as a linear combination of shorter words. Hence we may just assume $n \leq 4$. Then the conclusion is deduced by applying (5) and the identity obtained by multiplying (5) on the right by $\overline{\mathbf{b}}$. □

Corollary 2.5. If \mathbf{a} is special, then each word in \mathbf{a}, \mathbf{b} can be written as a linear combination of $\mathbf{e}, \mathbf{a}, \mathbf{b}, \bar{\mathbf{b}}, \mathbf{ab}, \mathbf{ba}, \mathbf{a}\bar{\mathbf{b}}, \bar{\mathbf{b}}\mathbf{a}, \mathbf{b}\lrcorner\mathbf{a}$, with coefficients in $\mathbb{C}[t_1, t_2, t_{\bar{2}}, t_{12}, t_{1\bar{2}}]$.

Proof. By Proposition 2.4, it suffices to show this for

$$\bar{\mathbf{a}}\mathbf{b}, \mathbf{b}\bar{\mathbf{a}}, \bar{\mathbf{a}}\bar{\mathbf{b}}, \bar{\mathbf{b}}\bar{\mathbf{a}}, \mathbf{a}\lrcorner\mathbf{b}, \mathbf{a}\lrcorner\bar{\mathbf{b}}, [\mathbf{a}, \mathbf{b}], [\mathbf{b}, \mathbf{a}].$$

Since $\bar{\mathbf{a}} \in \mathbb{C}\langle \mathbf{e}, \mathbf{a} \rangle$, using (6) we can easily show that indeed each of these can be written as a linear combination of $\mathbf{e}, \mathbf{a}, \mathbf{b}, \bar{\mathbf{b}}, \mathbf{ab}, \mathbf{ba}, \mathbf{a}\bar{\mathbf{b}}, \bar{\mathbf{b}}\mathbf{a}, \mathbf{b}\lrcorner\mathbf{a}$, with coefficients in $\mathbb{C}[t_1, t_2, t_{\bar{2}}, t_{12}, t_{1\bar{2}}]$. \square

3 Irreducible representations of two-generator groups

Observe that for any $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{M}^1$,

$$\det[\mathbf{u}^\lrcorner, \mathbf{v}^\lrcorner, \mathbf{w}^\lrcorner] = \text{tr}(\mathbf{uvw}). \quad (8)$$

Lemma 3.1. *If $\mathbf{a} \in \mathcal{M}^0$, then*

$$\det[(\mathbf{b} - \mathbf{b}^{\text{tr}})^\lrcorner, (\mathbf{ab} - \mathbf{b}^{\text{tr}}\mathbf{a})^\lrcorner, (\bar{\mathbf{a}}\mathbf{b} - \mathbf{b}^{\text{tr}}\bar{\mathbf{a}})^\lrcorner] = t_{12\bar{1}\bar{2}} - t_{21\bar{2}\bar{1}}.$$

Proof. Expanding $\mathbf{q} := (\mathbf{b} - \mathbf{b}^{\text{tr}})(\mathbf{ab} - \mathbf{b}^{\text{tr}}\mathbf{a})(\bar{\mathbf{a}}\mathbf{b} - \mathbf{b}^{\text{tr}}\bar{\mathbf{a}})$ as

$$\begin{aligned} & (\mathbf{bab}\bar{\mathbf{a}}\mathbf{b} - (\mathbf{b}^{\text{tr}})^2\mathbf{ab}^{\text{tr}}\bar{\mathbf{a}}) + (\mathbf{bb}^{\text{tr}}\mathbf{ab}^{\text{tr}}\bar{\mathbf{a}} - \mathbf{babb}^{\text{tr}}\bar{\mathbf{a}}) \\ & + (\mathbf{b}^{\text{tr}}\mathbf{abb}^{\text{tr}}\bar{\mathbf{a}} - \mathbf{b}^{\text{tr}}\mathbf{ab}\bar{\mathbf{a}}\mathbf{b}) + ((\mathbf{b}^{\text{tr}})^2\mathbf{b} - \mathbf{bb}^{\text{tr}}\mathbf{b}), \end{aligned}$$

and using $\text{tr}(\mathbf{z}^{\text{tr}}) = \text{tr}(\mathbf{z})$, we obtain

$$\text{tr}(\mathbf{q}) = \text{tr}(\mathbf{bab}\bar{\mathbf{a}}\mathbf{b}) - \text{tr}(\bar{\mathbf{a}}\mathbf{bab}^2) = t_{12\bar{1}\bar{2}} - t_{21\bar{2}\bar{1}}.$$

Then the result follows from (8). \square

Definition 3.2. Call a pair (\mathbf{a}, \mathbf{b}) *irreducible* if $\sigma_{\mathbf{a}, \mathbf{b}} : F_2 \rightarrow \text{SL}(3, \mathbb{C})$ is irreducible, or equivalently, neither \mathbf{a}, \mathbf{b} nor $\mathbf{a}^{\text{tr}}, \mathbf{b}^{\text{tr}}$ share an eigenvector. Otherwise, call (\mathbf{a}, \mathbf{b}) *reducible*.

Lemma 3.3. *If (\mathbf{a}, \mathbf{b}) is irreducible, then*

- (i) *at least one of \mathbf{a}, \mathbf{b} is ordinary;*
- (ii) *any $\mathbf{u} \in \mathcal{M}$ commuting with both \mathbf{a} and \mathbf{b} is scalar, so that $\text{Cen}(\mathbf{a}) \cap \text{Cen}(\mathbf{b}) = \{\mu\mathbf{e} : \mu \in \langle \omega \rangle\}$.*

Proof. By Burnside's Theorem (see [1] p. 433), $\text{Im}(\sigma_{\mathbf{a}, \mathbf{b}})$ spans \mathcal{M} .

(i) Assume \mathbf{a}, \mathbf{b} are both special. Then from Corollary 2.5 and (6) we see $\text{Im}(\sigma_{\mathbf{a}, \mathbf{b}}) \subset \mathbb{C}\langle \mathbf{e}, \mathbf{a}, \mathbf{b}, \mathbf{ab}, \mathbf{ba} \rangle$, so $\text{Im}(\sigma_{\mathbf{a}, \mathbf{b}})$ could not span \mathcal{M} .

(ii) If $\mathbf{ua} = \mathbf{au}$ and $\mathbf{ub} = \mathbf{bu}$, then $\mathbf{uw} = \mathbf{wu}$ for any word \mathbf{w} in \mathbf{a}, \mathbf{b} ; by (i), \mathbf{u} is central in \mathcal{M} . \square

Proposition 3.4. *The following conditions are equivalent to each other: (i) $t_{12\bar{1}\bar{2}} = t_{21\bar{2}\bar{1}}$; (ii) 1 is an eigenvalue of $[\mathbf{a}, \mathbf{b}]$; (iii) $\det(\mathbf{ab} - \mathbf{ba}) = 0$.*

Proof. Notice that, for any $\mathbf{z} \in \mathrm{SL}(3, \mathbb{C})$, $\mathrm{tr}(\mathbf{z}) = \mathrm{tr}(\bar{\mathbf{z}})$ if and only if 1 is an eigenvalue of \mathbf{z} , which is equivalent to the existence of a vector $\mathbf{a} \neq 0$ such that $\mathbf{z}\mathbf{a} = \mathbf{a}$. For $\mathbf{z} = [\mathbf{a}, \mathbf{b}]$, this means $(\mathbf{a}\mathbf{b} - \mathbf{b}\mathbf{a})(\bar{\mathbf{a}}\bar{\mathbf{b}}\mathbf{a}) = 0$, and the existence of such \mathbf{a} is equivalent to $\det(\mathbf{a}\mathbf{b} - \mathbf{b}\mathbf{a}) = 0$. \square

Remark 3.5. Interestingly, as shown in [17] Theorem 4.1 and 4.2, when $\mathbf{a}, \mathbf{b} \in \mathrm{SU}(2, 1)$, the property that 1 is an eigenvalue of $[\mathbf{a}, \mathbf{b}]$ gives geometric information about the action of the subgroup $\langle \mathbf{a}, \mathbf{b} \rangle$.

3.1 The asymmetric part

Suppose $t_{12\bar{1}\bar{2}} \neq t_{21\bar{2}\bar{1}}$, then obviously (\mathbf{a}, \mathbf{b}) is irreducible. Fix (\mathbf{a}, \mathbf{b}) , and define a bilinear pairing on \mathcal{M} by

$$((\mathbf{u}, \mathbf{v})) = (t_{12\bar{1}\bar{2}} - t_{21\bar{2}\bar{1}})^{-1} \cdot \mathrm{tr}(\mathbf{u}\mathbf{v}).$$

Direct computations lead to the following table (with zeros in empty cases):

$((\mathbf{u}, \mathbf{v}))$	\mathbf{e}	\mathbf{a}	$\bar{\mathbf{a}}$	\mathbf{b}	$\bar{\mathbf{b}}$	$\theta(\mathbf{a}\mathbf{b})$	$\theta(\bar{\mathbf{a}}\bar{\mathbf{b}})$	$\theta(\bar{\mathbf{a}}\mathbf{b})$	$\theta(\mathbf{a}\bar{\mathbf{b}})$
$\delta([\mathbf{a}, \mathbf{b}])$	1	t_1			$t_{\bar{2}}$	t_{12}	$t_{1\bar{2}} + t_1 t_{\bar{2}}$		$t_{\bar{1}\bar{2}}$
$\delta(\mathbf{b}\bar{\mathbf{a}})$		1				t_2	$t_{\bar{2}}$		
$\delta(\mathbf{b}\bar{\mathbf{a}})$			-1					$-t_2$	$-t_{\bar{2}}$
$\delta(\mathbf{a}\bar{\mathbf{b}})$				-1		$-t_1$		$-t_{\bar{1}}$	
$\delta(\mathbf{a}\bar{\mathbf{b}})$					1		t_1		$t_{\bar{1}}$
$\delta(\bar{\mathbf{a}}\bar{\mathbf{b}})$						1			
$\delta(\bar{\mathbf{a}}\bar{\mathbf{b}})$							-1		
$\delta(\bar{\mathbf{a}}\mathbf{b})$								-1	
$\delta(\mathbf{a}\bar{\mathbf{b}})$									1

Theorem 3.6. If $t_{12\bar{1}\bar{2}} \neq t_{21\bar{2}\bar{1}}$, then $\mathfrak{B}_\delta, \mathfrak{B}_\theta$ are both bases of \mathcal{M} , where

$$\begin{aligned} \mathfrak{B}_\delta &= \{\delta(\mathbf{a}\mathbf{b}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\bar{\mathbf{a}}\mathbf{b}), \delta(\mathbf{a}\bar{\mathbf{b}}), \delta(\mathbf{a}\bar{\mathbf{b}}), \delta(\mathbf{a}\bar{\mathbf{b}}), \delta(\mathbf{b}\bar{\mathbf{a}}), \delta(\mathbf{b}\bar{\mathbf{a}}), \delta([\mathbf{a}, \mathbf{b}])\}, \\ \mathfrak{B}_\theta &= \{\mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}, \theta(\mathbf{a}\mathbf{b}), \theta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \theta(\bar{\mathbf{a}}\mathbf{b}), \theta(\mathbf{a}\bar{\mathbf{b}})\}. \end{aligned}$$

Proof. It is a special case of a general result: if $\langle \cdot, \cdot \rangle$ is a bilinear pairing on a vector space \mathcal{V} of dimension n , and $\mathfrak{B}_1 = \{\mathbf{u}_1, \dots, \mathbf{u}_n\}, \mathfrak{B}_2 = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ are subsets of \mathcal{V} such that the matrix $P = [\langle \mathbf{u}_i, \mathbf{v}_j \rangle]_{n \times n}$ is invertible, then $\mathfrak{B}_1, \mathfrak{B}_2$ are both bases of \mathcal{V} .

To show this, suppose $\sum_{i=1}^n a_i \mathbf{u}_i = 0$, with $a_i \in \mathbb{C}$. Then $\sum_{i=1}^n a_i \langle \mathbf{u}_i, \mathbf{v}_j \rangle = 0$ for each j . Hence $[a_1, \dots, a_n]P = 0$, implying $a_1 = \dots = a_n = 0$. Similarly, any linear relation $\sum_{j=1}^n b_j \mathbf{v}_j = 0$ holds only when $b_1 = \dots = b_n = 0$. \square

Remark 3.7. In \mathfrak{B}_δ , one may replace $\delta([\mathbf{a}, \mathbf{b}])$ by $\delta([\mathbf{b}, \mathbf{a}])$. Indeed,

$$\begin{aligned} \delta([\mathbf{a}, \mathbf{b}]) + \delta([\mathbf{b}, \mathbf{a}]) &= (t_{\bar{1}\bar{2}} - t_{\bar{1}} t_2) \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) + (t_1 t_{\bar{2}} - t_{1\bar{2}}) \delta(\bar{\mathbf{a}}\mathbf{b}) \\ &\quad + t_{\bar{2}} \delta(\mathbf{a}\bar{\mathbf{b}}) + t_2 \delta(\mathbf{a}\bar{\mathbf{b}}) + t_{\bar{1}} \delta(\mathbf{b}\bar{\mathbf{a}}) + t_1 \delta(\mathbf{b}\bar{\mathbf{a}}). \end{aligned}$$

This is obtained by writing

$$\delta([\mathbf{a}, \mathbf{b}]) + \delta([\mathbf{b}, \mathbf{a}]) = \theta(\mathbf{a}\bar{\mathbf{b}})\bar{\mathbf{b}} - \bar{\mathbf{b}}\theta(\mathbf{a}\bar{\mathbf{b}}) + \theta(\mathbf{b}\bar{\mathbf{a}})\bar{\mathbf{a}} - \bar{\mathbf{a}}\theta(\mathbf{b}\bar{\mathbf{a}})$$

and applying (5).

3.2 The symmetric slice

Theorem 3.8. *For any irreducible pair (\mathbf{a}, \mathbf{b}) , there exists $\mathbf{c} \in \mathrm{SL}(3, \mathbb{C})$ with $\mathbf{c} \lrcorner \mathbf{a}$, $\mathbf{c} \lrcorner \mathbf{b}$ both symmetric if and only if $t_{12\overline{12}} = t_{21\overline{21}}$.*

Proof. The “only if” part is trivial: if $\mathbf{c} \lrcorner \mathbf{a}$, $\mathbf{c} \lrcorner \mathbf{b} \in \mathcal{M}^0$, then

$$t_{12\overline{12}} = \mathrm{tr}([\mathbf{c} \lrcorner \mathbf{a}, \mathbf{c} \lrcorner \mathbf{b}]) = \mathrm{tr}([\mathbf{c} \lrcorner \mathbf{a}, \mathbf{c} \lrcorner \mathbf{b}]^{\mathrm{tr}}) = t_{21\overline{21}}.$$

For the “if” part, suppose $t_{12\overline{12}} = t_{21\overline{21}}$. By Lemma 3.3 and the symmetry between \mathbf{a}, \mathbf{b} , we can assume that \mathbf{a} is ordinary; also we can assume \mathbf{a} to be symmetric. By Lemma 3.1, there exists a nontrivial linear relation

$$\alpha(\mathbf{b} - \mathbf{b}^{\mathrm{tr}})^{\top} + \beta(\mathbf{a}\mathbf{b} - \mathbf{b}^{\mathrm{tr}}\mathbf{a})^{\top} + \gamma(\overline{\mathbf{a}}\mathbf{b} - \mathbf{b}^{\mathrm{tr}}\overline{\mathbf{a}})^{\top} = \mathbf{0};$$

equivalently, $\tilde{\mathbf{c}}\mathbf{b} = \mathbf{b}^{\mathrm{tr}}\tilde{\mathbf{c}}$, where $\tilde{\mathbf{c}} = \alpha\mathbf{e} + \beta\mathbf{a} + \gamma\overline{\mathbf{a}} \in \mathcal{M}$.

We show $\det(\tilde{\mathbf{c}}) \neq 0$. Assume on the contrary that $\det(\tilde{\mathbf{c}}) = 0$. Suppose $\det(t\mathbf{e} - \mathbf{a}) = (t - \lambda_1)(t - \lambda_2)(t - \lambda_3)$. Let $\mathcal{E}(\lambda_i)$ denote the eigenspace of \mathbf{a} with respect to λ_i , which is one-dimensional. Let $f(t) = \beta t^2 + \alpha t + \gamma$. Then $f(\lambda_i) = 0$ for at least one i , and $f(\mathbf{a}) = \mathbf{a}\tilde{\mathbf{c}}$.

- If the greatest common divisor of $f(t)$ and $\det(t\mathbf{e} - \mathbf{a})$ is $t - \lambda_i$ for some i , then taking a nonzero $\mathbf{a} \in \mathcal{E}(\lambda_i)$, we have

$$f(\mathbf{a})(\mathbf{b}\mathbf{a}) = \mathbf{a}\tilde{\mathbf{c}}\mathbf{b}\mathbf{a} = \mathbf{a}\mathbf{b}^{\mathrm{tr}}\tilde{\mathbf{c}}\mathbf{a} = \mathbf{a}\mathbf{b}^{\mathrm{tr}}\overline{\mathbf{a}}f(\mathbf{a})\mathbf{a} = 0.$$

Consequently, $(\mathbf{a} - \lambda_i\mathbf{e})(\mathbf{b}\mathbf{a}) = 0$, i.e. $\mathbf{b}\mathbf{a} \in \mathcal{E}(\lambda_i)$, showing that $\mathbf{b}\mathbf{a}$ is a multiple of \mathbf{a} , so \mathbf{a} is a common eigenvector of \mathbf{a} and \mathbf{b} . This contradicts the irreducibility of (\mathbf{a}, \mathbf{b}) .

- If $f(t) = \beta(t - \lambda_i)(t - \lambda_j)$ for $i \neq j$, (note that $\lambda_i = \lambda_j$ is allowed), then $\{\tilde{\mathbf{c}}\mathbf{c} : \mathbf{c} \in \mathbb{C}^3\} = \mathcal{E}(\lambda_k)$, where $\{k\} = \{1, 2, 3\} - \{i, j\}$. Taking \mathbf{b} with $\mathbf{a} := \tilde{\mathbf{c}}\mathbf{b} \neq 0$, we have

$$\mathbf{b}^{\mathrm{tr}}\mathbf{a} = \mathbf{b}^{\mathrm{tr}}(\tilde{\mathbf{c}}\mathbf{b}) = \tilde{\mathbf{c}}(\mathbf{b}\mathbf{b}) \in \mathcal{E}(\lambda_k) = \mathbb{C}\langle \mathbf{a} \rangle,$$

so \mathbf{a} is an eigenvector of both \mathbf{a} and \mathbf{b}^{tr} . This also contradicts the irreducibility of (\mathbf{a}, \mathbf{b}) .

Thus, $\det(\tilde{\mathbf{c}}) \neq 0$. Taking κ with $\kappa^2 = \det(\tilde{\mathbf{c}})$, we have $\kappa^{-1}\tilde{\mathbf{c}} \in \mathrm{Cen}(\mathbf{a})$. By Lemma 2.2, there exists $\mathbf{c} \in \mathrm{Cen}(\mathbf{a}) \subset \mathbb{C}\langle \mathbf{e}, \mathbf{a}, \overline{\mathbf{a}} \rangle \subset \mathcal{M}^0$ with $\mathbf{c}^2 = \kappa^{-1}\tilde{\mathbf{c}}$, which ensures $\mathbf{c}\mathbf{b}\overline{\mathbf{c}} = \overline{\mathbf{c}}\mathbf{b}^{\mathrm{tr}}\mathbf{c}$, i.e. $\mathbf{c} \lrcorner \mathbf{b} \in \mathcal{M}^0$. \square

Let $\mathcal{R}_{\mathrm{SL}(3, \mathbb{C})}^{\mathrm{irr}, \mathrm{sym}}(F_2)$ denote the set of irreducible pairs (\mathbf{a}, \mathbf{b}) with $\mathbf{a}, \mathbf{b} \in \mathcal{M}^0$. Let $\mathcal{X}_{\mathrm{SL}(3, \mathbb{C})}^{\mathrm{irr}, \mathrm{sym}}(F_2) = \{\chi_{\sigma} : \sigma \in \mathcal{R}_{\mathrm{SL}(3, \mathbb{C})}^{\mathrm{irr}, \mathrm{sym}}(F_2)\}$, which coincides with the subset of $\mathcal{X}_{\mathrm{SL}(3, \mathbb{C})}(F_2)$ consisting of irreducible characters with $t_{12\overline{12}} = t_{21\overline{21}}$.

Theorem 3.9. *The action of $\mathrm{SO}(3, \mathbb{C})$ on $\mathcal{R}_{\mathrm{SL}(3, \mathbb{C})}^{\mathrm{irr}, \mathrm{sym}}(F_2)$ via conjugation is free, and is transitive on each fiber of $\chi : \mathcal{R}_{\mathrm{SL}(3, \mathbb{C})}^{\mathrm{irr}, \mathrm{sym}}(F_2) \rightarrow \mathcal{X}_{\mathrm{SL}(3, \mathbb{C})}^{\mathrm{irr}, \mathrm{sym}}(F_2)$. Consequently,*

$$\mathcal{X}_{\mathrm{SL}(3, \mathbb{C})}^{\mathrm{irr}, \mathrm{sym}}(F_2) \cong \mathcal{R}_{\mathrm{SL}(3, \mathbb{C})}^{\mathrm{irr}, \mathrm{sym}}(F_2) / \mathrm{SO}(3, \mathbb{C}).$$

Proof. The first assertion follows from Lemma 3.3 (ii) and the fact that $\text{SO}(3, \mathbb{C}) \cap \langle \omega \rangle \mathbf{e} = \{\mathbf{e}\}$.

For the second one, suppose $(\mathbf{a}, \mathbf{b}), (\mathbf{a}', \mathbf{b}') \in \mathcal{R}_{\text{SL}(3, \mathbb{C})}^{\text{irr, sym}}(F_2)$ have the same character. Then there exists $\mathbf{c} \in \text{SL}(3, \mathbb{C})$ such that $\mathbf{c} \lrcorner \mathbf{a} = \mathbf{a}'$, $\mathbf{c} \lrcorner \mathbf{b} = \mathbf{b}'$. From $\mathbf{c} \lrcorner \mathbf{a} = \mathbf{a}' = (\mathbf{a}')^{\text{tr}} = (\mathbf{c} \lrcorner \mathbf{a})^{\text{tr}}$ we deduce $\mathbf{c}^{\text{tr}} \mathbf{c} \in \text{Cen}(\mathbf{a})$, and from $\mathbf{c} \lrcorner \mathbf{b} = (\mathbf{c} \lrcorner \mathbf{b})^{\text{tr}}$ we deduce $\mathbf{c}^{\text{tr}} \mathbf{c} \in \text{Cen}(\mathbf{b})$. Hence by Lemma 3.3 (ii), $\mathbf{c}^{\text{tr}} \mathbf{c} = \mu \mathbf{e}$ for some $\mu \in \langle \omega \rangle$. Then $\mu \mathbf{c} \in \text{SO}(3, \mathbb{C})$, and $(\mu \mathbf{c}) \lrcorner \mathbf{a} = \mathbf{a}'$, $(\mu \mathbf{c}) \lrcorner \mathbf{b} = \mathbf{b}'$. \square

We aim to find convenient bases for \mathcal{M}^0 , \mathcal{M}^1 , \mathcal{M} .

Lemma 3.10. *If \mathbf{a}, \mathbf{b} are symmetric and the pair (\mathbf{a}, \mathbf{b}) is irreducible, then*

$$\mathcal{M}^0 = \mathbb{C}\langle \mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}, \theta(\mathbf{ab}), \theta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \theta(\bar{\mathbf{a}}\mathbf{b}), \theta(\mathbf{a}\bar{\mathbf{b}}), \theta([\mathbf{a}, \mathbf{b}]), \theta([\bar{\mathbf{b}}, \mathbf{a}]) \rangle,$$

$$\mathcal{M}^1 = \mathbb{C}\langle \delta(\mathbf{ab}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\mathbf{a}\bar{\mathbf{b}}), \delta(\bar{\mathbf{a}}\mathbf{b}), \delta(\mathbf{a} \lrcorner \mathbf{b}), \delta(\mathbf{a} \lrcorner \bar{\mathbf{b}}), \delta(\mathbf{b} \lrcorner \mathbf{a}), \delta(\mathbf{b} \lrcorner \bar{\mathbf{a}}), \delta([\mathbf{a}, \mathbf{b}]) \rangle.$$

Proof. Note the basic fact: if \mathcal{M} is spanned by $\mathbf{x}_1, \dots, \mathbf{x}_n$, then \mathcal{M}^0 is spanned by $\mathbf{x}_1 + \mathbf{x}_1^{\text{tr}}, \dots, \mathbf{x}_n + \mathbf{x}_n^{\text{tr}}$, and \mathcal{M}^1 is spanned by $\mathbf{x}_1 - \mathbf{x}_1^{\text{tr}}, \dots, \mathbf{x}_n - \mathbf{x}_n^{\text{tr}}$.

The first equation follows from Proposition 2.4 together with (5), and the second one follows from Proposition 2.4 together with Remark 3.7. \square

Theorem 3.11. *Suppose \mathbf{a}, \mathbf{b} are symmetric and the pair (\mathbf{a}, \mathbf{b}) is irreducible.*

(i) *If \mathbf{a} is special (so that \mathbf{b} is ordinary), then $\{\mathbf{e}, \mathbf{a}, \mathbf{b}, \bar{\mathbf{b}}, \theta(\mathbf{ab}), \theta(\bar{\mathbf{a}}\bar{\mathbf{b}})\}$ is a basis of \mathcal{M}^0 , and $\{\delta(\mathbf{ab}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\mathbf{b} \lrcorner \mathbf{a})\}$ is a basis of \mathcal{M}^1 .*

(ii) *If \mathbf{a}, \mathbf{b} are both ordinary, then at least one of the following occurs:*

- 1) *$\{\mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}, \theta(\mathbf{ab})\}$ is a basis of \mathcal{M}^0 , and $\{\delta(\mathbf{ab}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\mathbf{a}\bar{\mathbf{b}})\}$ is a basis of \mathcal{M}^1 , so $\mathfrak{B}_1^{\text{sym}} = \{\mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}, \mathbf{ab}, \mathbf{ba}, \bar{\mathbf{a}}\bar{\mathbf{b}}, \bar{\mathbf{a}}\mathbf{b}\}$ is a basis of \mathcal{M} ;*
- 2) *$\{\mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}, \theta(\bar{\mathbf{a}}\bar{\mathbf{b}})\}$ is a basis of \mathcal{M}^0 , and $\{\delta(\mathbf{ab}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\bar{\mathbf{a}}\mathbf{b})\}$ is a basis of \mathcal{M}^1 , so $\mathfrak{B}_2^{\text{sym}} = \{\mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}, \mathbf{ab}, \mathbf{b}\bar{\mathbf{a}}, \bar{\mathbf{a}}\bar{\mathbf{b}}, \bar{\mathbf{a}}\mathbf{b}\}$ is a basis of \mathcal{M} .*

Therefore, in any case,

$$\mathcal{M}^0 = \mathbb{C}\langle \mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}, \theta(\mathbf{ab}), \theta(\bar{\mathbf{a}}\bar{\mathbf{b}}) \rangle,$$

$$\mathcal{M}^1 = \mathbb{C}\langle \delta(\mathbf{ab}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\mathbf{a}\bar{\mathbf{b}}), \delta(\bar{\mathbf{a}}\mathbf{b}) \rangle,$$

$$\mathcal{M} = \mathbb{C}\langle \mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}, \mathbf{ab}, \mathbf{ba}, \bar{\mathbf{a}}\bar{\mathbf{b}}, \bar{\mathbf{a}}\mathbf{b}, \bar{\mathbf{a}}\bar{\mathbf{b}} \rangle.$$

Proof. (i) This is an immediate consequence of Corollary 2.5 and the basic fact stated in the proof of Lemma 3.10.

(ii) The result is established by proving a series of assertions.

1. $\mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}$ are linearly independent.

Suppose

$$\alpha_0 \mathbf{e} + \alpha_1 \mathbf{a} + \alpha_2 \bar{\mathbf{a}} + \beta_1 \mathbf{b} + \beta_2 \bar{\mathbf{b}} = \mathbf{0}.$$

Then $\beta_1 \mathbf{b} + \beta_2 \bar{\mathbf{b}}$ commutes with \mathbf{a} and \mathbf{b} , so by Lemma 3.3 (ii), $\beta_1 \mathbf{b} + \beta_2 \bar{\mathbf{b}} = \gamma \mathbf{e}$ for some γ . Since \mathbf{b} is ordinary, we have $\beta_1 = \beta_2 = \gamma = 0$, so that $\alpha_0 \mathbf{e} + \alpha_1 \mathbf{a} + \alpha_2 \bar{\mathbf{a}} = \mathbf{0}$. This forces $\alpha_0 = \alpha_1 = \alpha_2 = 0$, as \mathbf{a} is ordinary.

2. Let $\mathcal{V}^0 = \mathbb{C}\langle \mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}} \rangle$. Then either $\theta(\mathbf{ab}) \notin \mathcal{V}^0$ or $\theta(\bar{\mathbf{a}}\bar{\mathbf{b}}) \notin \mathcal{V}^0$.

Assume on the contrary that $\theta(\mathbf{ab}), \theta(\bar{\mathbf{a}}\bar{\mathbf{b}}) \in \mathcal{V}^0$. Then

$$\theta(\mathbf{a}^2\mathbf{b}) = t_1\theta(\mathbf{ab}) - 2t_1\mathbf{b} + \theta(\bar{\mathbf{a}}\bar{\mathbf{b}}) \in \mathcal{V}^0;$$

by (5), $\theta(\mathbf{a}\lrcorner\mathbf{b}) \in \mathcal{V}^0$; by (6), $\mathbf{aba} \in \mathcal{V}^0$. Suppose

$$\theta(\mathbf{ab}) = \alpha_0\mathbf{e} + \alpha_1\mathbf{a} + \alpha_2\bar{\mathbf{a}} + \beta_1\mathbf{b} + \beta_2\bar{\mathbf{b}}. \quad (9)$$

Multiplying this by \mathbf{b} on the left and right, we respectively obtain

$$\mathbf{bab} + \mathbf{b}^2\mathbf{a} = \alpha_0\mathbf{b} + \alpha_1\mathbf{ba} + \alpha_2\mathbf{b}\bar{\mathbf{a}} + \beta_1\mathbf{b}^2 + \beta_2\mathbf{e}, \quad (10)$$

$$\mathbf{ab}^2 + \mathbf{bab} = \alpha_0\mathbf{b} + \alpha_1\mathbf{ab} + \alpha_2\bar{\mathbf{a}}\mathbf{b} + \beta_1\mathbf{b}^2 + \beta_2\mathbf{e}. \quad (11)$$

Their sum yields $\theta(\mathbf{ab}^2) + 2\mathbf{bab} \in \mathcal{V}^0$. By (6), $\theta(\mathbf{ab}^2) \in \mathcal{V}^0$. Furthermore, since $\theta(\mathbf{ab}^2) = t_2\theta(\mathbf{ab}) - 2t_2\mathbf{a} + \theta(\bar{\mathbf{a}}\bar{\mathbf{b}})$, we have $\theta(\bar{\mathbf{a}}\bar{\mathbf{b}}) \in \mathcal{V}^0$.

A similar argument leads to $\theta(\bar{\mathbf{a}}\bar{\mathbf{b}}^2), \theta(\bar{\mathbf{a}}\bar{\mathbf{b}}) \in \mathcal{V}^0$, and moreover,

$$\theta(\bar{\mathbf{a}}^2\bar{\mathbf{b}}) = t_1\theta(\bar{\mathbf{a}}\bar{\mathbf{b}}) - 2t_1\bar{\mathbf{b}} + \theta(\bar{\mathbf{a}}\bar{\mathbf{b}}) \in \mathcal{V}^0.$$

Also by (5), $\theta(\mathbf{a}\lrcorner\bar{\mathbf{b}}), \theta(\mathbf{b}\lrcorner\mathbf{a}), \theta(\mathbf{b}\lrcorner\bar{\mathbf{a}}) \in \mathcal{V}^0$.

Multiplying (9) by $\bar{\mathbf{a}}\bar{\mathbf{b}}$ on the right and by $\bar{\mathbf{b}}\bar{\mathbf{a}}$ on the left, and then adding the resulted equations, we obtain

$$\theta([\mathbf{a}, \mathbf{b}]) + 2\mathbf{e} = \alpha_0\theta(\bar{\mathbf{a}}\bar{\mathbf{b}}) + 2\alpha_1\bar{\mathbf{b}} + \alpha_2\theta(\bar{\mathbf{a}}^2\bar{\mathbf{b}}) + \beta_1\theta(\mathbf{b}\lrcorner\bar{\mathbf{a}}) + 2\beta_2\bar{\mathbf{b}}\bar{\mathbf{a}},$$

implying $\theta([\mathbf{a}, \mathbf{b}]) \in \mathcal{V}^0$. Similarly, $\theta([\mathbf{b}, \mathbf{a}]) \in \mathcal{V}^0$.

By Lemma 3.10, $\mathcal{M}^0 = \mathcal{V}^0$ which is absurd, so the assumption $\theta(\mathbf{ab}), \theta(\bar{\mathbf{a}}\bar{\mathbf{b}}) \in \mathcal{V}^0$ is false.

Therefore, $\mathcal{M}^0 = \mathbb{C}\langle \mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}, \theta(\mathbf{ab}) \rangle$ or $\mathcal{M}^0 = \mathbb{C}\langle \mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}, \theta(\bar{\mathbf{a}}\bar{\mathbf{b}}) \rangle$.

3. Let $\mathcal{V}^1 = \mathbb{C}\langle \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) \rangle$. Then either $\delta(\mathbf{ab}) \notin \mathcal{V}^1$ or $\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) \notin \mathcal{V}^1$.

Assume on the contrary that $\delta(\mathbf{ab}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) \in \mathcal{V}^1$. Suppose

$$\delta(\mathbf{ab}) = \eta_1\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) + \eta_2\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}). \quad (12)$$

Note that $\eta_1 \neq 0$: otherwise, $\mathbf{a}(\mathbf{b} - \eta_2\bar{\mathbf{b}}) = (\mathbf{b} - \eta_2\bar{\mathbf{b}})\mathbf{a}$, so that $\mathbf{b} - \eta_2\bar{\mathbf{b}} \in \text{Cen}(\mathbf{a}) \cap \text{Cen}(\mathbf{b})$, which would contradict the assumption that \mathbf{b} is ordinary. Similarly, $\eta_2 \neq 0$.

Multiplying (12) by \mathbf{a} on the left and right, and then taking sum, we obtain $\delta(\mathbf{a}^2\mathbf{b}) = -\eta_1\delta(\mathbf{a}\lrcorner\mathbf{b}) + \eta_2\delta(\mathbf{a}^2\bar{\mathbf{b}})$; since $\delta(\mathbf{a}^2\mathbf{b}^{\pm 1}) = t_1\delta(\mathbf{ab}^{\pm 1}) + \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}^{\pm 1})$, we have $\delta(\mathbf{a}\lrcorner\mathbf{b}) \in \mathcal{V}^1$.

Multiplying (12) by $\bar{\mathbf{a}}$ on the left and right, and then taking sum, we obtain $\delta(\mathbf{a}\lrcorner\mathbf{b}) = \eta_1\delta(\bar{\mathbf{a}}^2\bar{\mathbf{b}}) + \eta_2\delta(\mathbf{a}\lrcorner\bar{\mathbf{b}})$, implying $\delta(\mathbf{a}\lrcorner\bar{\mathbf{b}}) \in \mathcal{V}^1$.

Similarly, $\delta(\mathbf{b}\lrcorner\mathbf{a}), \delta(\mathbf{b}\lrcorner\bar{\mathbf{a}}) \in \mathcal{V}^1$.

Multiplying (12) by $\bar{\mathbf{a}}\bar{\mathbf{b}}$ on the right and $\bar{\mathbf{b}}\bar{\mathbf{a}}$ on the left, and then taking sum, we obtain $\delta([\mathbf{a}, \mathbf{b}]) = \eta_1\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}\bar{\mathbf{a}}\bar{\mathbf{b}}) + \eta_2\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}\bar{\mathbf{a}}\bar{\mathbf{b}})$. Since

$$\begin{aligned}\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}\bar{\mathbf{a}}\bar{\mathbf{b}}) &\stackrel{(7)}{=} \delta((t_{12}\bar{\mathbf{a}} - t_{12}\bar{\mathbf{b}} + \bar{\mathbf{b}}\bar{\mathbf{a}}\bar{\mathbf{b}})\bar{\mathbf{b}}) = t_{12}\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) + \delta(\bar{\mathbf{b}}\bar{\mathbf{a}}\bar{\mathbf{b}}^2) \\ &= t_{12}\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) + \delta(\bar{\mathbf{b}}\bar{\mathbf{a}}(t_2\bar{\mathbf{b}} - t_2\mathbf{e} + \mathbf{b})) \\ &= t_{12}\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) + t_2\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) - \delta(\mathbf{b}\lrcorner\bar{\mathbf{a}}) \in \mathcal{V}^1, \\ \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}\bar{\mathbf{a}}\bar{\mathbf{b}}) &\stackrel{(7)}{=} \delta(\mathbf{a}(t_{12}\bar{\mathbf{b}} - t_{12}\mathbf{a} + \bar{\mathbf{a}}\bar{\mathbf{b}}\bar{\mathbf{a}})) = t_{12}\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) + \delta(\mathbf{a}^2\bar{\mathbf{b}}\bar{\mathbf{a}}) \\ &= t_{12}\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) + \delta((t_1\mathbf{a} - t_1\mathbf{e} + \bar{\mathbf{a}})\bar{\mathbf{b}}\bar{\mathbf{a}}) \\ &= t_{12}\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) + t_1\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) - \delta(\mathbf{a}\lrcorner\bar{\mathbf{b}}) \in \mathcal{V}^1,\end{aligned}$$

we have $\delta([\mathbf{a}, \mathbf{b}]) \in \mathcal{V}^1$.

By Lemma 3.10, $\mathcal{M}^1 = \mathcal{V}^1$ which is absurd.

Therefore, $\mathcal{M}^1 = \mathbb{C}\langle\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}})\rangle$ or $\mathcal{M}^1 = \mathbb{C}\langle\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}})\rangle$.

4. Finally, if $\mathcal{M}^1 = \mathbb{C}\langle\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}})\rangle$, then $\mathcal{M}^0 = \mathbb{C}\langle\mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}, \theta(\bar{\mathbf{a}}\bar{\mathbf{b}})\rangle$: otherwise, supposing (9), the difference between (10) and (11) would imply

$$\alpha_1\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) + \alpha_2\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) = \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}^2) = t_1\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) + \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}),$$

so that $\dim \mathcal{M}^1 \leq 2$ which is absurd.

Similarly, if $\mathcal{M}^1 = \mathbb{C}\langle\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}), \delta(\bar{\mathbf{a}}\bar{\mathbf{b}})\rangle$, then $\mathcal{M}^0 = \mathbb{C}\langle\mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}, \theta(\bar{\mathbf{a}}\bar{\mathbf{b}})\rangle$.

The proof is complete. \square

3.3 Computing $\mathcal{X}_{\text{SL}(3, \mathbb{C})}^{\text{irr}}(\Gamma)$ for general Γ

For any two-generator group $\Gamma = \langle z_1, z_2 \mid r_1, \dots, r_k \rangle$, we can decompose

$$\begin{aligned}\mathcal{R}_{\text{SL}(3, \mathbb{C})}^{\text{irr}}(\Gamma) &= \mathcal{R}_{\text{SL}(3, \mathbb{C})}^{\text{asym}}(\Gamma) \cup \mathcal{R}_{\text{SL}(3, \mathbb{C})}^{\text{irr, sym}}(\Gamma), \\ \mathcal{X}_{\text{SL}(3, \mathbb{C})}^{\text{irr}}(\Gamma) &= \mathcal{X}_{\text{SL}(3, \mathbb{C})}^{\text{asym}}(\Gamma) \cup \mathcal{X}_{\text{SL}(3, \mathbb{C})}^{\text{irr, sym}}(\Gamma);\end{aligned}$$

the meaning of each term is self-evident.

For each j , the condition $\sigma_{\mathbf{a}, \mathbf{b}}(r_j) = \mathbf{e}$ can be transformed into an equation of the form $\mathbf{s}_j = \mathbf{0}$, where \mathbf{s}_j is a noncommutative polynomial in $\mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}$.

The asymmetric part $\mathcal{X}_{\text{SL}(3, \mathbb{C})}^{\text{asym}}(\Gamma)$ is relatively easy to determine: for $j = 1, \dots, k$, $\mathbf{s}_j = \mathbf{0}$ is equivalent to $((\mathbf{u}, \mathbf{s}_j)) = 0$ for each $\mathbf{u} \in \mathfrak{B}_\theta$ or $((\mathbf{u}, \mathbf{s}_j)) = 0$ for each $\mathbf{u} \in \mathfrak{B}_\delta$; a system of polynomial equations in traces can be written down.

To find the symmetric part $\mathcal{X}_{\text{SL}(3, \mathbb{C})}^{\text{irr, sym}}(\Gamma)$, we may assume \mathbf{a}, \mathbf{b} to be symmetric. When \mathbf{a} is special, by Corollary 2.5, each \mathbf{s}_j can be written as a linear combination of $\mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}, \mathbf{a}\bar{\mathbf{b}}, \bar{\mathbf{a}}\mathbf{b}, \mathbf{b}\lrcorner\bar{\mathbf{a}}$ which form a basis of \mathcal{M} , so $\mathbf{s}_j = \mathbf{0}$ is equivalent to the simultaneous vanishing of the coefficients. When \mathbf{b} is special, the situation is similar. When \mathbf{a}, \mathbf{b} are both ordinary, requiring $\text{tr}(\mathbf{u}\mathbf{s}_j) = 0$ for each

$$\mathbf{u} \in \{\mathbf{e}, \mathbf{a}, \bar{\mathbf{a}}, \mathbf{b}, \bar{\mathbf{b}}, \mathbf{a}\bar{\mathbf{b}}, \bar{\mathbf{a}}\mathbf{b}, \mathbf{b}\lrcorner\bar{\mathbf{a}}, \bar{\mathbf{b}}\lrcorner\bar{\mathbf{a}}\},$$

the result is a system of polynomial equations in traces.

When the relation r_j admits a certain symmetry (this occurs for rational link groups; see [2] Exercise 12.1), the computation may be dramatically simplified by investigating $\text{tr}(\mathbf{u}\mathbf{s}_j) = 0$ for \mathbf{u} in the bases given by Theorem 3.6 and 3.11.

Explicit representations can be found without much difficulty, once the corresponding characters are determined.

Example 3.12. Let $\Gamma = \langle x, y \mid xy\bar{x}yx = yx\bar{y}xy \rangle$, the group of the figure eight knot. Given $\mathbf{x}, \mathbf{y} \in \mathrm{SL}(3, \mathbb{C})$, there exists a representation $\rho : \Gamma \rightarrow \mathrm{SL}(3, \mathbb{C})$ with $\rho(x) = \mathbf{x}, \rho(y) = \mathbf{y}$ if and only if

$$\mathbf{c} := \mathbf{xy}\bar{\mathbf{x}}\mathbf{yx} - \mathbf{yx}\bar{\mathbf{y}}\mathbf{xy} = \mathbf{0}.$$

Suppose ρ is irreducible. Then \mathbf{x}, \mathbf{y} must be both ordinary, as they are conjugate.

In the case $t_{\bar{1}2} = t_{1\bar{2}} = s$ (the reason for considering this will soon be clear),

$$\begin{aligned} \mathbf{xy}\bar{\mathbf{x}}\mathbf{yx} &\stackrel{(7)}{=} \mathbf{x}(t_{\bar{1}2}\mathbf{y} - t_{1\bar{2}}\mathbf{x} + \mathbf{x}\bar{\mathbf{y}}\mathbf{x}) = t_{\bar{1}2}\mathbf{xyx} - t_{1\bar{2}}\mathbf{x}^3 + \mathbf{x}^2\bar{\mathbf{y}}\mathbf{x}^2 \\ &= -\theta(\mathbf{x}\bar{\mathbf{y}}) - t_{\bar{1}2}\theta(\bar{\mathbf{x}}\mathbf{y}) + \bar{t}t_{\bar{1}2}\mathbf{y} - \bar{t}^2\bar{\mathbf{y}} + (\bar{t} - \bar{t}^2 + \bar{t}t_{\bar{1}2} + t_{1\bar{2}}t_{\bar{1}2} + tt_{\bar{1}2})\mathbf{x} \\ &\quad + (t_{\bar{1}2} + tt_{\bar{1}2} - 2\bar{t}^2)\bar{\mathbf{x}} + (1 - \bar{t})t_{\bar{1}2}\mathbf{e} + (\star), \end{aligned}$$

where (\star) stands for the sum of terms that are symmetric between x and y , and we have omit some intermediate steps that can be easily filled in. So

$$\mathbf{c} = (1 - s)(\theta(\bar{\mathbf{x}}\mathbf{y}) - \theta(\mathbf{x}\bar{\mathbf{y}})) + (\bar{t} - \bar{t}^2 + st_{1\bar{2}} + tt_{\bar{1}2})(\mathbf{x} - \mathbf{y}) + (t_{\bar{1}2} + st - \bar{t}^2)(\bar{\mathbf{x}} - \bar{\mathbf{y}}).$$

(i) If $t_{1\bar{2}\bar{1}2} \neq t_{2\bar{1}2\bar{1}}$, then it follows from

$$\begin{aligned} 0 &= \mathrm{tr}(\mathbf{c}\delta(\bar{\mathbf{x}}\mathbf{y})) = (t_{1\bar{2}\bar{1}2} - t_{2\bar{1}2\bar{1}})(1 - t_{\bar{1}2}), \\ 0 &= \mathrm{tr}(\mathbf{c}\delta(\mathbf{x}\bar{\mathbf{y}})) = (t_{1\bar{2}\bar{1}2} - t_{2\bar{1}2\bar{1}})(t_{\bar{1}2} - 1) \end{aligned}$$

that $t_{\bar{1}2} = t_{1\bar{2}} = 1$. Hence by (3), $(\bar{\mathbf{x}}\mathbf{y})^4 = \mathbf{e}$, and

$$\mathbf{0} = \mathbf{c} = (\bar{t} - \bar{t}^2 + t_{1\bar{2}} + tt_{\bar{1}2})(\mathbf{x} - \mathbf{y}) + (t_{\bar{1}2} + t - \bar{t}^2)(\bar{\mathbf{x}} - \bar{\mathbf{y}}).$$

Similarly as in the first step of proving Theorem 3.11 (ii),

$$\bar{t} - \bar{t}^2 + t_{1\bar{2}} + tt_{\bar{1}2} = t_{\bar{1}2} + t - \bar{t}^2 = 0,$$

i.e. $t_{\bar{1}2} = \bar{t}^2 - t$, $t_{1\bar{2}} = t^2 - \bar{t}$. This implies $t_{1\bar{2}\bar{1}2} = t_{\bar{1}2\bar{1}2} = t_{\bar{1}2^2} = t_{1\bar{2}^2} = 0$, which is equivalent to $(\mathbf{x}^2\bar{\mathbf{y}})^3 = (\mathbf{y}^2\bar{\mathbf{x}})^3 = \mathbf{e}$. Thus, ρ factors through

$$\Gamma \rightarrow D(3, 3, 4) := \langle a, b \mid a^3 = b^3 = (ab)^4 = 1 \rangle, \quad x \mapsto bab, \quad y \mapsto \bar{a}\bar{b}\bar{a},$$

which is surjective, as witnessed by $\bar{y}x\bar{y} \mapsto a$ and $x\bar{y}x \mapsto b$.

(ii) If $t_{\bar{1}2} \neq t_{1\bar{2}}$, then with $x = \bar{y}zy\bar{z}y$, the presentation of Γ becomes $\langle z, y \mid zy\bar{z}yz = yz\bar{y}zy \rangle$. We have $z = x \lrcorner y$. Hence $\mathrm{tr}([\mathbf{y}, \mathbf{z}]) = t_{\bar{1}2}$ and $\mathrm{tr}([\mathbf{z}, \mathbf{y}]) = t_{1\bar{2}}$. Now that the relation has the same form as the previous one and $\mathrm{tr}([\mathbf{y}, \mathbf{z}]) \neq \mathrm{tr}([\mathbf{z}, \mathbf{y}])$, an argument same as above leads to

$$(\mathbf{z}^2\bar{\mathbf{y}})^3 = (\mathbf{y}^2\bar{\mathbf{z}})^3 = (\bar{\mathbf{z}}\mathbf{y})^4 = \mathbf{e}.$$

This time ρ factors through the epimorphism $\Gamma \twoheadrightarrow D(3, 3, 4)$, $x \mapsto \bar{b}a$, $y \mapsto \bar{a}\bar{b}\bar{a}$.

(iii) Now suppose $t_{12\overline{12}} = t_{21\overline{21}}$ and $t_{\overline{12}} = t_{1\overline{2}} = s$. Then $\mathbf{c} = \mathbf{0}$ is equivalent to $\text{tr}(\mathbf{c}\mathbf{u}) = 0$ for $\mathbf{u} \in \{\mathbf{e}, \mathbf{x}, \overline{\mathbf{x}}, \mathbf{y}, \overline{\mathbf{y}}, \theta(\mathbf{x}\mathbf{y}), \theta(\overline{\mathbf{x}}\mathbf{y})\}$. It is easy to verify

$$\text{tr}(\mathbf{c}) = \text{tr}(\mathbf{c}\theta(\mathbf{x}\mathbf{y})) = 0, \quad \text{tr}(\mathbf{c}\mathbf{x}) = \text{tr}(\mathbf{c}\mathbf{y}), \quad \text{tr}(\mathbf{c}\overline{\mathbf{x}}) = \text{tr}(\mathbf{c}\overline{\mathbf{y}}).$$

Thus, $\mathcal{X}_{\text{SL}(3, \mathbb{C})}^{\text{irr, sym}}(\Gamma)$ is given by

$$\text{tr}(\mathbf{c}\mathbf{x}) = \text{tr}(\mathbf{c}\overline{\mathbf{x}}) = \text{tr}(\mathbf{c}\theta(\overline{\mathbf{x}}\mathbf{y})) = 0,$$

which can be written as a system of polynomial equations in $t, \bar{t}, t_{12}, t_{\overline{12}}, s$.

These can be compared with the components V_0, V_1, V_2 of $\mathcal{X}_{\text{SL}(3, \mathbb{C})}^{\text{irr}}(\Gamma)$, given in [11]; S, T there respectively correspond to x, y here. It is easy to see that V_1, V_2 of [11] are reproduced exactly by (i),(ii). We believe that V_0 can be identified with the result of (iii), but some more efforts are needed.

4 Application to the double twist link groups




Figure 1: Left: the double twist link $J(2m+1, 2n+1)$. Right: when $m = n = 1$, it is the Whitehead link.

Let $L = J(2m+1, 2n+1)$ be the double twist link as shown in Figure 1. Let $\Gamma = \pi_1(E_L)$, where E_L is the complement of a tubular neighborhood of L in S^3 . Adopting Wirtinger presentation, we have $x_1 = z^m \lrcorner x$ and $y_1 = z^{m+1} \lrcorner y$, where $z = xy$; writing y_1 in another way leads to the unique relation $(x_1 \overline{y})^{n+1} \lrcorner y = z^{m+1} \lrcorner y$. Substituting $x = z \overline{y}$ results in

$$\Gamma = \langle y, z \mid \overline{y} z^{m+1} w^{n+1} = z^{m+1} w^{n+1} \overline{y} \rangle, \quad \text{with } w = \overline{z}^{m+1} y z^m y.$$

Note that $z^{m+1} w^{n+1} = (z^{m+1} w^{n+1})^\leftarrow$, so the relation is the same as

$$\overline{y} z^{m+1} w^{n+1} = (\overline{y} z^{m+1} w^{n+1})^\leftarrow.$$

Given $\mathbf{y}, \mathbf{z} \in \text{SL}(3, \mathbb{C})$, there exists a representation $\rho : \Gamma \rightarrow \text{SL}(3, \mathbb{C})$ with $\rho(y) = \mathbf{y}, \rho(z) = \mathbf{z}$ if and only if

$$\mathbf{c} := \delta(\overline{\mathbf{y}} \mathbf{z}^{m+1} \mathbf{w}^{n+1}) = \mathbf{0}, \quad \text{with } \mathbf{w} = \overline{\mathbf{z}}^{m+1} \mathbf{y} \mathbf{z}^m \mathbf{y}. \quad (13)$$

Abbreviate $\alpha_k(\mathbf{z}), \beta_k(\mathbf{z}), \gamma_k(\mathbf{z})$ to $\alpha_k, \beta_k, \gamma_k$, respectively. Let $\tilde{\alpha} = \alpha_{n+1}(\mathbf{w}), \tilde{\beta} = \beta_{n+1}(\mathbf{w}), \tilde{\gamma} = \gamma_{n+1}(\mathbf{w})$. Then

$$\begin{aligned} \mathbf{c} &= \delta(\overline{\mathbf{y}} \mathbf{z}^{m+1} (\tilde{\alpha} \mathbf{w} + \tilde{\beta} \mathbf{e} + \tilde{\gamma} \overline{\mathbf{w}})) \\ &= \tilde{\alpha} \delta(\mathbf{z}^m \mathbf{y}) + \tilde{\beta} \delta(\overline{\mathbf{y}} \mathbf{z}^{m+1}) + \tilde{\gamma} \delta(\overline{\mathbf{y}} \mathbf{z}^{m+1} \overline{\mathbf{y}} \mathbf{z}^m \overline{\mathbf{y}} \mathbf{z}^{m+1}). \end{aligned} \quad (14)$$

To shorten the writing, let $p_k = \text{tr}(\mathbf{y}\mathbf{z}^k)$, $q_k = \text{tr}(\bar{\mathbf{y}}\bar{\mathbf{z}}^k)$. Then

$$\begin{aligned}\bar{\mathbf{y}}\mathbf{z}^{m+1}\bar{\mathbf{y}}\mathbf{z}^m\bar{\mathbf{y}}\mathbf{z}^{m+1} &= \bar{\mathbf{y}}\mathbf{z}^{m+1}(q_{-m}\bar{\mathbf{y}} - p_m\mathbf{z}^m + \mathbf{z}^m\mathbf{y}\mathbf{z}^m)\mathbf{z}^{m+1} \\ &= q_{-m}(\bar{\mathbf{y}}\mathbf{z}^{m+1})^2 - p_m\bar{\mathbf{y}}\mathbf{z}^{3m+2} + \bar{\mathbf{y}}\mathbf{z}^{2m+1}\mathbf{y}\mathbf{z}^{2m+1} \\ &= q_{-m}(q_{m+1}\bar{\mathbf{y}}\mathbf{z}^{m+1} - p_{-m-1}\mathbf{e} + \bar{\mathbf{z}}^{m+1}\mathbf{y}) - p_m\bar{\mathbf{y}}\mathbf{z}^{3m+2} \\ &\quad + \bar{\mathbf{y}}(p_{2m+1}\mathbf{z}^{2m+1} - q_{-2m-1}\bar{\mathbf{y}} + \bar{\mathbf{y}}\mathbf{z}^{2m+1}\bar{\mathbf{y}}).\end{aligned}$$

Hence

$$\begin{aligned}&\delta(\bar{\mathbf{y}}\mathbf{z}^{m+1}\bar{\mathbf{y}}\mathbf{z}^m\bar{\mathbf{y}}\mathbf{z}^{m+1}) \\ &= q_{-m}q_{m+1}\delta(\bar{\mathbf{y}}\mathbf{z}^{m+1}) + q_{-m}\delta(\bar{\mathbf{z}}^{m+1}\mathbf{y}) - p_m\delta(\bar{\mathbf{y}}\mathbf{z}^{3m+2}) + p_{2m+1}\delta(\bar{\mathbf{y}}\mathbf{z}^{2m+1}) \\ &\quad + p_0\delta(\bar{\mathbf{y}}\mathbf{z}^{2m+1}) + \delta(\mathbf{y}\bar{\mathbf{z}}^{2m+1}\bar{\mathbf{y}}) \\ &= q_{-m}q_{m+1}(\alpha_{m+1}\delta(\bar{\mathbf{y}}\mathbf{z}) + \gamma_{m+1}\delta(\bar{\mathbf{y}}\bar{\mathbf{z}})) + q_{-m}(\alpha_{-m-1}\delta(\mathbf{z}\mathbf{y}) + \gamma_{-m-1}\delta(\bar{\mathbf{z}}\mathbf{y})) \\ &\quad - p_m(\alpha_{3m+2}\delta(\bar{\mathbf{y}}\mathbf{z}) + \gamma_{3m+2}\delta(\bar{\mathbf{y}}\bar{\mathbf{z}})) + p_{2m+1}(\alpha_{2m+1}\delta(\bar{\mathbf{y}}\mathbf{z}) + \gamma_{2m+1}\delta(\bar{\mathbf{y}}\bar{\mathbf{z}})) \\ &\quad + p_0(\alpha_{-2m-1}\delta(\bar{\mathbf{y}}\mathbf{z}) + \gamma_{-2m-1}\delta(\bar{\mathbf{y}}\bar{\mathbf{z}})) + \alpha_{-2m-1}\delta(\mathbf{y}\lrcorner\mathbf{z}) + \gamma_{-2m-1}\delta(\mathbf{y}\lrcorner\bar{\mathbf{z}}) \\ &= -q_{-m}\alpha_{-m-1}\delta(\mathbf{y}\mathbf{z}) + \varsigma_1\delta(\bar{\mathbf{y}}\mathbf{z}) - q_{-m}\gamma_{-m-1}\delta(\mathbf{y}\bar{\mathbf{z}}) + \varsigma_2\delta(\bar{\mathbf{y}}\bar{\mathbf{z}}) \\ &\quad + \alpha_{-2m-1}\delta(\mathbf{y}\lrcorner\mathbf{z}) + \gamma_{-2m-1}\delta(\mathbf{y}\lrcorner\bar{\mathbf{z}}),\end{aligned}$$

with

$$\varsigma_1 = q_{-m}q_{m+1}\alpha_{m+1} - p_m\alpha_{3m+2} + p_{2m+1}\alpha_{2m+1} + p_0\alpha_{-2m-1}, \quad (15)$$

$$\varsigma_2 = q_{-m}q_{m+1}\gamma_{m+1} - p_m\gamma_{3m+2} + p_{2m+1}\gamma_{2m+1} + p_0\gamma_{-2m-1}. \quad (16)$$

By (14),

$$\mathbf{c} = \eta_1\delta(\mathbf{y}\mathbf{z}) + \eta_2\delta(\bar{\mathbf{y}}\mathbf{z}) + \eta_3\delta(\mathbf{y}\bar{\mathbf{z}}) + \eta_4\delta(\bar{\mathbf{y}}\bar{\mathbf{z}}) + \eta_5\delta(\mathbf{y}\lrcorner\mathbf{z}) + \eta_6\delta(\mathbf{y}\lrcorner\bar{\mathbf{z}}), \quad (17)$$

with

$$\eta_1 = -\tilde{\gamma}q_{-m}\alpha_{-m-1} - \tilde{\alpha}\alpha_m, \quad \eta_2 = \tilde{\gamma}\varsigma_1 + \tilde{\beta}\alpha_{m+1}, \quad (18)$$

$$\eta_3 = -\tilde{\gamma}q_{-m}\gamma_{-m-1} - \tilde{\alpha}\gamma_m, \quad \eta_4 = \tilde{\gamma}\varsigma_2 + \tilde{\beta}\gamma_{m+1}, \quad (19)$$

$$\eta_5 = \tilde{\gamma}\alpha_{-2m-1}, \quad \eta_6 = \tilde{\gamma}\gamma_{-2m-1}. \quad (20)$$

4.1 The asymmetric slice: $\text{tr}(\mathbf{y}\mathbf{z}\bar{\mathbf{y}}\bar{\mathbf{z}}) \neq \text{tr}(\mathbf{z}\mathbf{y}\bar{\mathbf{z}}\bar{\mathbf{y}})$

The table given in Section 3.1 will be repeatedly applied.

By Theorem 3.6, $\mathbf{c} = \mathbf{0}$ if and only if

$$((\mathbf{u}, \mathbf{c})) = 0, \quad \text{for } \mathbf{u} \in \{\mathbf{e}, \mathbf{y}, \bar{\mathbf{y}}, \mathbf{z}, \bar{\mathbf{z}}, \theta(\mathbf{y}\mathbf{z}), \theta(\mathbf{y}\bar{\mathbf{z}}), \theta(\bar{\mathbf{y}}\mathbf{z}), \theta(\bar{\mathbf{y}}\bar{\mathbf{z}})\}.$$

From (17) it is clear that $((\mathbf{u}, \mathbf{c})) = 0$ for $\mathbf{u} \in \{\mathbf{e}, \mathbf{y}, \bar{\mathbf{y}}\}$. As necessary conditions, $0 = ((\mathbf{z}, \mathbf{c})) = -\tilde{\gamma}\gamma_{-2m-1}$, and $0 = ((\bar{\mathbf{z}}, \mathbf{c})) = \tilde{\gamma}\alpha_{-2m-1}$.

- If $\tilde{\gamma} \neq 0$, then $\alpha_{-2m-1} = \gamma_{-2m-1} = 0$, implying $\mathbf{z}^{2m+1} = \mu\mathbf{e}$ with $\mu \in \langle \omega \rangle$, so $\mathbf{w} = \mu^{-1}(\mathbf{z}^m\mathbf{y})^2$, and (13) becomes

$$\mathbf{0} = \mathbf{c}' := \mu^n \mathbf{c} = \delta((\mathbf{z}^m\mathbf{y})^{2n+1}).$$

Let $\alpha' = \alpha_{2n+1}(\mathbf{z}^m \mathbf{y})$, $\gamma' = \gamma_{2n+1}(\mathbf{z}^m \mathbf{y})$. Then

$$\begin{aligned} \mathbf{c}' &= \alpha' \delta(\mathbf{z}^m \mathbf{y}) + \gamma' \delta(\overline{\mathbf{y}} \mathbf{z}^m) \\ &= \alpha' (\alpha_m \delta(\mathbf{z} \mathbf{y}) + \gamma_m \delta(\overline{\mathbf{z}} \mathbf{y})) + \gamma' (\alpha_{-m} \delta(\overline{\mathbf{y}} \mathbf{z}) + \gamma_{-m} \delta(\overline{\mathbf{y}} \mathbf{z})). \end{aligned}$$

We easily obtain

$$\begin{aligned} 0 &= ((\theta(\mathbf{y} \mathbf{z}), \mathbf{c}')) = \gamma_{-m} \gamma', & 0 &= ((\theta(\mathbf{y} \overline{\mathbf{z}}), \mathbf{c}')) = -\alpha_{-m} \gamma', \\ 0 &= ((\theta(\overline{\mathbf{y}} \mathbf{z}), \mathbf{c}')) = \gamma_m \alpha', & 0 &= ((\theta(\overline{\mathbf{y}} \overline{\mathbf{z}}), \mathbf{c}')) = -\alpha_m \alpha'. \end{aligned}$$

If $\alpha_m = \gamma_m = 0$, then $\mathbf{z}^m = \kappa \mathbf{e}$, $\kappa \in \langle \omega \rangle$, but this together with $\mathbf{z}^{2m+1} = \mu \mathbf{e}$ would imply $\mathbf{z} = \kappa \mu \mathbf{e}$. Similarly, $\alpha_{-m} = \gamma_{-m} = 0$ is neither possible. Hence $\alpha' = \gamma' = 0$, implying $(\mathbf{z}^m \mathbf{y})^{2n+1} = \nu \mathbf{e}$, $\nu \in \langle \omega \rangle$.

- If $\tilde{\gamma} = 0$, then

$$\begin{aligned} \mathbf{c} &= \tilde{\alpha} \delta(\mathbf{z}^m \mathbf{y}) + \tilde{\beta} \delta(\overline{\mathbf{y}} \mathbf{z}^{m+1}) \\ &= \tilde{\alpha} (\alpha_m \delta(\mathbf{z} \mathbf{y}) + \gamma_m \delta(\overline{\mathbf{z}} \mathbf{y})) + \tilde{\beta} (\alpha_{m+1} \delta(\overline{\mathbf{y}} \mathbf{z}) + \gamma_{m+1} \delta(\overline{\mathbf{y}} \mathbf{z})). \end{aligned}$$

As is easy to verify, $((\mathbf{z}, \mathbf{c})) = ((\overline{\mathbf{z}}, \mathbf{c})) = 0$, and

$$\begin{aligned} 0 &= ((\theta(\mathbf{y} \mathbf{z}), \mathbf{c})) = \gamma_{m+1} \tilde{\beta}, & 0 &= ((\theta(\mathbf{y} \overline{\mathbf{z}}), \mathbf{c})) = -\alpha_{m+1} \tilde{\beta}, \\ 0 &= ((\theta(\overline{\mathbf{y}} \mathbf{z}), \mathbf{c})) = \gamma_m \tilde{\alpha}, & 0 &= ((\theta(\overline{\mathbf{y}} \overline{\mathbf{z}}), \mathbf{c})) = -\alpha_m \tilde{\alpha}. \end{aligned}$$

There are two possibilities.

- If $\alpha_m = \gamma_m = 0$, then $\mathbf{z}^m = \mu \mathbf{e}$, $\mu \in \langle \omega \rangle$. Consequently, $\alpha_{m+1} \neq 0$ or $\gamma_{m+1} \neq 0$, either implying $\tilde{\beta} = 0$. Then $\mathbf{w}^{n+1} = \tilde{\alpha} \mathbf{w}$, which can be rewritten as $\mathbf{w}^n = \nu \mathbf{e}$, $\nu \in \langle \omega \rangle$; equivalently, $(\overline{\mathbf{z}} \mathbf{y}^2)^n = \nu \mathbf{e}$.
- Otherwise, $\tilde{\alpha} = 0$, so that $\mathbf{w}^{n+1} = \nu \mathbf{e}$, $\nu \in \langle \omega \rangle$, and $\tilde{\beta} = \nu$. Hence $\alpha_{m+1} = \gamma_{m+1} = 0$, implying $\mathbf{z}^{m+1} = \mu \mathbf{e}$, $\mu \in \langle \omega \rangle$. Then $\mathbf{w}^{n+1} = \nu \mathbf{e}$ becomes $(\overline{\mathbf{z}} \mathbf{y}^2)^{n+1} = \nu \mathbf{e}$.

Lemma 4.1. *Let k be a nonzero integer. If \mathbf{z} is ordinary and $\mathbf{z}^k = \mu \mathbf{e}$ with $\mu \in \langle \omega \rangle$, then $|k| \geq 3$, \mathbf{z} is conjugate to $\mathbf{d}_{\lambda \zeta}^\lambda$ for some λ, ζ such that $\lambda^k = \mu$, $\zeta^k = 1$, and $\zeta \notin \{1, \lambda^{-3}, \pm \sqrt{\lambda^{-3}}\}$. In particular, $\mu = 1$ when $k = \pm 3$.*

Proof. Now that $\mathbf{z}^k = \mu \mathbf{e}$ and \mathbf{z} is ordinary, $|k|$ must be at least 3. It is impossible for \mathbf{z} to be conjugate to \mathbf{h}_λ or $\nu \mathbf{k}$, as

$$\begin{aligned} (\mathbf{h}_\lambda)^k &= (\mathbf{d}_\lambda^\lambda + \lambda \mathbf{f}^2)^k = (\mathbf{d}_\lambda^\lambda)^k + k\lambda (\mathbf{d}_\lambda^\lambda)^{k-1} \mathbf{f}^2 \notin \mathbb{C} \langle \mathbf{e} \rangle, \\ \mathbf{k}^k &= (\mathbf{e} + \mathbf{f} + \mathbf{f}^2)^k = \mathbf{e} + k\mathbf{f} + \frac{k(k+1)}{2} \mathbf{f}^2 \notin \mathbb{C} \langle \mathbf{e} \rangle. \end{aligned}$$

Hence \mathbf{z} is conjugate to $\mathbf{d}_{\lambda \zeta}^\lambda$ for some λ, ζ such that $\lambda^k = \mu$, $\zeta^k = 1$. Since \mathbf{z} is ordinary, $\lambda, \lambda \zeta, (\lambda^2 \zeta)^{-1}$ should be distinct; equivalently, $\zeta \notin \{1, \lambda^{-3}, \pm \sqrt{\lambda^{-3}}\}$. This constraint forces $\mu = 1$ when $k = \pm 3$. \square

Put $\Lambda_k = \emptyset$ if $|k| \leq 2$, and put

$$\Lambda_k = \{(\lambda, \zeta) : \lambda^{3k} = \zeta^k = 1, \zeta \neq 1, \lambda^{-3}, \pm \sqrt{\lambda^{-3}}\}$$

if $|k| \geq 3$. Recall (4):

$$\mathcal{X}_{\text{SL}(3, \mathbb{C})}(F_2) \cong \{(s_1, s_{\overline{1}}, s_2, s_{\overline{2}}, s_3, s_{\overline{3}}, s_4, s_{\overline{4}}, s_5) : s_5^2 - P s_5 + Q = 0\}.$$

Theorem 4.2. *The asymmetric slice $\mathcal{X}_{\text{SL}(3,\mathbb{C})}^{\text{asym}}(\Gamma)$ is the disjoint union of*

$$\begin{aligned}\mathcal{X}_{\lambda_1, \zeta_1; \lambda_2, \zeta_2}^1, & \quad (\lambda_1, \zeta_1) \in \Lambda_{2m+1}, \quad (\lambda_2, \zeta_2) \in \Lambda_{2n+1}, \\ \mathcal{X}_{\lambda_1, \zeta_1; \lambda_2, \zeta_2}^2, & \quad (\lambda_1, \zeta_1) \in \Lambda_m, \quad (\lambda_2, \zeta_2) \in \Lambda_n, \\ \mathcal{X}_{\lambda_1, \zeta_1; \lambda_2, \zeta_2}^3, & \quad (\lambda_1, \zeta_1) \in \Lambda_{m+1}, \quad (\lambda_2, \zeta_2) \in \Lambda_{n+1},\end{aligned}$$

where $\mathcal{X}_{\lambda_1, \zeta_1; \lambda_2, \zeta_2}^j$ is isomorphic to the subspace of $\mathcal{X}_{\text{SL}(3,\mathbb{C})}(F_2)$ determined by

$$s_i = \lambda_i + \lambda_i \zeta_i + (\lambda_i^2 \zeta_i)^{-1}, \quad s_{\bar{i}} = \lambda_i^{-1} + (\lambda_i \zeta_i)^{-1} + \lambda_i^2 \zeta_i, \quad i = 1, 2, \quad P^2 \neq 4Q.$$

Proof. According to the above discussions, there are three possibilities for \mathbf{y}, \mathbf{z} to define a representation: 1) $\mathbf{z}^{2m+1} = \mu \mathbf{e}$, $(\mathbf{z}^m \mathbf{y})^{2n+1} = \nu \mathbf{e}$; 2) $\mathbf{z}^m = \mu \mathbf{e}$, $(\bar{\mathbf{z}} \mathbf{y}^2)^n = \nu \mathbf{e}$; 3) $\mathbf{z}^{m+1} = \mu \mathbf{e}$, $(\bar{\mathbf{z}} \mathbf{y}^2)^{n+1} = \nu \mathbf{e}$. In each case, $\mu, \nu \in \langle \omega \rangle$.

In case 1), let $\mathbf{a} = \mathbf{z}$, $\mathbf{b} = \mathbf{z}^m \mathbf{y}$. Then $\text{tr}(\mathbf{y} \mathbf{z} \bar{\mathbf{y}} \bar{\mathbf{z}}) \neq \text{tr}(\mathbf{z} \bar{\mathbf{y}} \bar{\mathbf{z}} \mathbf{y})$ is equivalent to $\text{tr}(\mathbf{a} \bar{\mathbf{b}} \bar{\mathbf{a}} \mathbf{b}) \neq \text{tr}(\mathbf{b} \bar{\mathbf{a}} \bar{\mathbf{b}} \mathbf{a})$. By Lemma 4.1, \mathbf{a} is conjugate to $\mathbf{d}_{\lambda_1 \zeta_1}^{\lambda_1}$ for some $(\lambda_1, \zeta_1) \in \Lambda_{2m+1}$, and \mathbf{b} is conjugate to $\mathbf{d}_{\lambda_2 \zeta_2}^{\lambda_2}$ for some $(\lambda_2, \zeta_2) \in \Lambda_{2n+1}$. Hence

$$\begin{aligned}\text{tr}(\mathbf{a}) &= \lambda_1 + \lambda_1 \zeta_1 + (\lambda_1^2 \zeta_1)^{-1}, & \text{tr}(\bar{\mathbf{a}}) &= \lambda_1^{-1} + (\lambda_1 \zeta_1)^{-1} + \lambda_1^2 \zeta_1, \\ \text{tr}(\mathbf{b}) &= \lambda_2 + \lambda_2 \zeta_2 + (\lambda_2^2 \zeta_2)^{-1}, & \text{tr}(\bar{\mathbf{b}}) &= \lambda_2^{-1} + (\lambda_2 \zeta_2)^{-1} + \lambda_2^2 \zeta_2.\end{aligned}$$

Cases 2) and 3) are dealt with similarly. \square

4.2 The symmetric slice: $\text{tr}(\mathbf{y} \mathbf{z} \bar{\mathbf{y}} \bar{\mathbf{z}}) = \text{tr}(\mathbf{z} \bar{\mathbf{y}} \bar{\mathbf{z}} \mathbf{y})$

Suppose \mathbf{y}, \mathbf{z} are symmetric and (\mathbf{y}, \mathbf{z}) is irreducible. Let $t_1 = \text{tr}(\mathbf{y})$, $t_2 = \text{tr}(\mathbf{z})$, and so forth. Recall (15)–(20).

Theorem 4.3. *The symmetric slice $\mathcal{X}_{\text{SL}(3,\mathbb{C})}^{\text{sym,irr}}(\Gamma) = \mathcal{X}_{\text{sp,od}} \cup \mathcal{X}_{\text{od,sp}} \cup \mathcal{X}_{\text{od,od}}$, where $\mathcal{X}_{\text{sp,od}}$ is determined by*

$$\begin{aligned}\lambda &\in \mathbb{C}^*, \quad t_1 = 2\lambda + \lambda^{-2}, \quad t_{\bar{1}} = 2\lambda^{-1} + \lambda^2, \\ t_{\bar{1}2} &= (\lambda^2 + \lambda^{-1})t_2 - \lambda t_{12}, \quad t_{\bar{1}\bar{2}} = (\lambda^2 + \lambda^{-1})t_{\bar{2}} - \lambda t_{\bar{1}\bar{2}}, \\ \eta_1 - \lambda \eta_2 + (\lambda^2 + \lambda^{-1})\eta_5 &= \eta_3 - \lambda \eta_4 + (\lambda^2 + \lambda^{-1})\eta_6 = 0,\end{aligned}\tag{21}$$

$\mathcal{X}_{\text{od,sp}}$ is determined by

$$\begin{aligned}\lambda &\in \mathbb{C}^*, \quad t_2 = 2\lambda + \lambda^{-2}, \quad t_{\bar{2}} = 2\lambda^{-1} + \lambda^2, \\ t_{\bar{1}2} &= (\lambda^2 + \lambda^{-1})t_1 - \lambda t_{12}, \quad t_{\bar{1}\bar{2}} = (\lambda^2 + \lambda^{-1})t_{\bar{1}} - \lambda t_{\bar{1}\bar{2}}, \\ \eta_1 - \lambda \eta_3 &= \eta_2 - \lambda \eta_4 = \eta_5 - \lambda \eta_6 = 0,\end{aligned}\tag{22}$$

and $\mathcal{X}_{\text{od,od}}$ is determined by $[\eta_1, \dots, \eta_6]A = 0$, with

$$A = \begin{bmatrix} t_{1212} - t_1^2 t_2^2 & t_{12\bar{1}2} - t_2^2 & t_{121\bar{2}} - t_1^2 & t_{12\bar{1}\bar{2}} - 3 \\ t_{12\bar{1}2} - t_2^2 & t_{\bar{1}2\bar{1}2} - t_{\bar{1}}^2 t_2^2 & t_{12\bar{1}\bar{2}} - 3 & t_{\bar{1}2\bar{1}\bar{2}} - t_{\bar{1}}^2 \\ t_{121\bar{2}} - t_1^2 & t_{12\bar{1}\bar{2}} - 3 & t_{\bar{1}21\bar{2}} - t_1 t_2^2 & t_{12\bar{1}\bar{2}} - t_{\bar{2}}^2 \\ t_{12\bar{1}\bar{2}} - 3 & t_{\bar{1}21\bar{2}} - t_{\bar{1}}^2 & t_{12\bar{1}\bar{2}} - t_{\bar{2}}^2 & t_{\bar{1}21\bar{2}} - t_{\bar{1}}^2 t_2^2 \\ t_{12^2} - t_1^2 t_{\bar{1}2} & t_{\bar{1}2\bar{1}^2} - t_{\bar{1}}^2 t_2^2 & t_1 - t_1^2 t_{\bar{1}2} & t_{12\bar{1}^2} - t_{\bar{1}} \\ t_1 - t_1^2 t_{\bar{1}2} & t_{\bar{1}2\bar{1}^2} - t_{\bar{1}} & t_{\bar{1}2^2} - t_1^2 t_{\bar{1}2} & t_{\bar{1}2\bar{1}^2} - t_{\bar{1}}^2 \end{bmatrix}.\tag{23}$$

Proof. If \mathbf{y} is special, with multiple eigenvalue λ , so that $\bar{\mathbf{y}} = (\lambda^2 + \lambda^{-1})\mathbf{e} - \lambda\mathbf{y}$, then $\delta(\bar{\mathbf{y}}\mathbf{z}^{\pm 1}) = -\lambda\delta(\mathbf{y}\mathbf{z}^{\pm 1})$, and $\delta(\mathbf{y}\lrcorner\mathbf{z}^{\pm 1}) = (\lambda^2 + \lambda^{-1})\delta(\mathbf{y}\mathbf{z}^{\pm 1})$, implying

$$\mathbf{c} = (\eta_1 - \lambda\eta_2 + (\lambda^2 + \lambda^{-1})\eta_5)\delta(\mathbf{y}\mathbf{z}) + (\eta_3 - \lambda\eta_4 + (\lambda^2 + \lambda^{-1})\eta_6)\delta(\mathbf{y}\bar{\mathbf{z}}).$$

By Theorem 3.11 (i), $\mathbf{c} = \mathbf{0}$ is equivalent to (21).

If \mathbf{z} is special, with multiple eigenvalue λ , so that $\bar{\mathbf{z}} = (\lambda^2 + \lambda^{-1})\mathbf{e} - \lambda\mathbf{z}$, then $\delta(\mathbf{y}^{\pm 1}\bar{\mathbf{z}}) = -\lambda\delta(\mathbf{y}^{\pm 1}\mathbf{z})$, and $\delta(\mathbf{y}\lrcorner\bar{\mathbf{z}}) = -\lambda\delta(\mathbf{y}\lrcorner\mathbf{z})$, implying

$$\mathbf{c} = (\eta_1 - \lambda\eta_3)\delta(\mathbf{y}\mathbf{z}) + (\lambda^2 - \lambda\eta_4)\delta(\bar{\mathbf{y}}\mathbf{z}) + (\eta_5 - \lambda\eta_6)\delta(\mathbf{y}\lrcorner\mathbf{z}).$$

Hence $\mathbf{c} = \mathbf{0}$ is equivalent to (22).

Now suppose \mathbf{y} , \mathbf{z} are both ordinary. Let $\mathbf{u}_1 = \mathbf{v}_1 = \mathbf{y}\mathbf{z}$, $\mathbf{u}_2 = \mathbf{v}_2 = \bar{\mathbf{y}}\mathbf{z}$, $\mathbf{u}_3 = \mathbf{v}_3 = \mathbf{y}\bar{\mathbf{z}}$, $\mathbf{u}_4 = \mathbf{v}_4 = \bar{\mathbf{y}}\bar{\mathbf{z}}$, $\mathbf{u}_5 = \mathbf{y}\lrcorner\mathbf{z}$, $\mathbf{u}_6 = \mathbf{y}\lrcorner\bar{\mathbf{z}}$. Direct computation shows $[\text{tr}(\delta(\mathbf{u}_i)\delta(\mathbf{v}_j))/2]_{6 \times 4} = A$ as given by (23). By Theorem 3.11 (ii), $\mathbf{c} = \mathbf{0}$ is equivalent to $[\eta_1, \dots, \eta_6]A = 0$. \square

4.3 The Whitehead link

When $m = n = 1$, L is the Whitehead link. Now

$$\Gamma = \langle y, z \mid yzy\bar{z}^2yz = zy\bar{z}^2yzy \rangle.$$

With $a = z, b = zy$, one has an alternative presentation $\langle a, b \mid \bar{a}b^2\bar{a}^3b = b\bar{a}^3b^2\bar{a} \rangle$, which is the same as given by (3) in [9]. Let $\mathbf{a} = \mathbf{z}, \mathbf{b} = \mathbf{zy}$, and let $t_1 = \text{tr}(\mathbf{a}), t_2 = \text{tr}(\mathbf{b})$ and so forth.

For the asymmetric slice, by Lemma 4.1, only case 1) in the proof of Theorem 4.2 occurs: $\mathbf{a}^3 = \mathbf{b}^3 = \mathbf{e}$, which is equivalent to $t_1 = t_{\bar{1}} = t_2 = t_{\bar{2}} = 0$. This recovers the result of [9]. With (4) recalled, $\mathcal{X}_{\text{SL}(3, \mathbb{C})}^{\text{asym}}(\Gamma)$ is identified with the subspace of $\mathcal{X}_{\text{SL}(3, \mathbb{C})}(F_2)$ given by

$$\{(0, 0, 0, 0, s_3, s_{\bar{3}}, s_4, s_{\bar{4}}, s_5) : s_5^2 - Ps_5 + Q = 0, P^2 \neq 4Q\}.$$

For the symmetric slice, let $\check{\mathbf{c}} = \delta(\bar{\mathbf{a}}\mathbf{b}^2\bar{\mathbf{a}}^3\mathbf{b})$. Since

$$\bar{\mathbf{a}}\mathbf{b}^2\bar{\mathbf{a}}^3\mathbf{b} = (t_{\bar{1}}^2 - t_1)\bar{\mathbf{a}}\mathbf{b}^2\bar{\mathbf{a}}\mathbf{b} + (1 - t_1t_{\bar{1}})\bar{\mathbf{a}}\mathbf{b}^3 + t_{\bar{1}}\bar{\mathbf{a}}\mathbf{b}^2\mathbf{a}\mathbf{b},$$

we have

$$\begin{aligned} \check{\mathbf{c}} &= (t_{\bar{1}}^2 - t_1)(t_{\bar{1}2}\delta(\bar{\mathbf{a}}\mathbf{b}) + \delta(\bar{\mathbf{b}}^2\bar{\mathbf{a}}\mathbf{b})) + (1 - t_1t_{\bar{1}})((t_{\bar{2}}^2 - t_2)\delta(\bar{\mathbf{a}}\mathbf{b}) + t_2\delta(\bar{\mathbf{a}}\bar{\mathbf{b}})) \\ &\quad + t_{\bar{1}}(t_2\delta(\bar{\mathbf{a}}\mathbf{b}\mathbf{a}\mathbf{b}) + \delta(\bar{\mathbf{a}}\mathbf{b}\mathbf{a}\mathbf{b})) \\ &= ((t_{\bar{1}}^2 - t_1)t_{\bar{1}2} + (1 - t_1t_{\bar{1}})(t_{\bar{2}}^2 - t_2))\delta(\bar{\mathbf{a}}\mathbf{b}) + (1 - t_1t_{\bar{1}})t_2\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) + t_{\bar{1}}\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}\mathbf{a}\mathbf{b}) \\ &\quad + (t_{\bar{1}}^2 - t_1)(-t_2\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) + \delta(\bar{\mathbf{b}}\mathbf{a}\bar{\mathbf{b}})) + t_{\bar{1}}t_2(t_{12}\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) - t_1\delta(\bar{\mathbf{b}}\bar{\mathbf{a}}) + \delta(\bar{\mathbf{a}}\bar{\mathbf{b}}\bar{\mathbf{a}})) \\ &= \eta\delta(\bar{\mathbf{a}}\mathbf{b}) + t_2(t_1 - t_{\bar{1}}^2)\delta(\bar{\mathbf{a}}\mathbf{b}) + t_2\delta(\bar{\mathbf{a}}\bar{\mathbf{b}}) + t_{\bar{1}}t_2\delta(\mathbf{a}\lrcorner\bar{\mathbf{b}}) + (t_{\bar{1}}^2 - t_1)\delta(\mathbf{b}\lrcorner\mathbf{a}) - t_{\bar{1}}\delta([\mathbf{b}, \mathbf{a}]), \end{aligned}$$

with

$$\eta = (t_{\bar{1}}^2 - t_1)(t_2t_{\bar{1}2} - t_{\bar{1}}t_{\bar{2}} + t_{\bar{1}\bar{2}}) + (1 - t_1t_{\bar{1}})(t_{\bar{2}}^2 - t_2) + t_{\bar{1}}t_2t_{12}.$$

In the last step, we have expanded $t_{\bar{1}2}$ as $t_2t_{\bar{1}2} - t_{\bar{1}}t_{\bar{2}} + t_{\bar{1}\bar{2}}$. Similarly as in Theorem 4.3, we can decompose $\mathcal{X}_{\text{SL}(3, \mathbb{C})}^{\text{sym, irr}}(\Gamma)$ into three parts, corresponding respectively to three cases: \mathbf{a} is special; \mathbf{b} is special; \mathbf{a}, \mathbf{b} are both ordinary.

References

- [1] W. Burnside, *On the condition of reducibility of any group of linear substitutions*. Proc. Lond. Math. Soc. (1905), no. 3, 430–434.
- [2] G. Burder, H. Zieschang, *Knots*. de Gruyter Studies in Mathematics, vol. 5, Walter de Gruyter, Berlin, New York, 2003.
- [3] M. Culler, P.B. Shalen, *Varieties of group representations and splittings of 3-manifolds*. Ann. Math. (2), 117 (1983), no. 1, 109–146.
- [4] M. Deraux, *On spherical CR uniformization of 3-manifolds*. Experimental Math. 24 (2015), no. 3, 355–370.
- [5] M. Deraux, E. Falbel, *Complex hyperbolic geometry of the figure eight knot*. Geometry & Topology, 19 (2015), no. 1, 237–293.
- [6] E. Falbel, A. Guilloux, P.-V. Koseleff, F. Rouillier, M. Thistlethwaite, *Character varieties for $SL(3; \mathbb{C})$: the figure eight knot*. Exp. Math 25 (2016), no. 2, 219–235.
- [7] E. Falbel, P.V. Koseleff, F. Rouillier, *Representations of fundamental groups of 3-manifolds into $PGL(3, \mathbb{C})$: exact computations in low complexity*. Geome. Dedicata, 177 (2015), no. 1, 229–255.
- [8] E. Falbel, Q.-X. Wang, *Duality and invariants of representations of fundamental groups of 3-manifolds into $PGL(3, \mathbb{C})$* . J. Lond. Math. Soc. 95 (2017), no. 1, 1–22.
- [9] A. Guilloux, P. Will, *On $SL(3, \mathbb{C})$ -representations of the Whitehead link group*. Geom. Dedicata (2018). <https://doi.org/10.1007/s10711-018-0404-8>
- [10] M. Heusener, *Some recent results about the $SL_n(\mathbb{C})$ -representation spaces of knot groups*. Séminaire de théorie spectrale et géométrie 32 (2014–2015), 137–161.
- [11] M. Heusener, V. Muñoz, J. Porti, *The $SL(3, \mathbb{C})$ -character variety of the figure eight knot*. Illinois J. Math. 60 (2016), 55–98.
- [12] S. Lawton, *Generators, relations and symmetries in pairs of 3×3 unimodular matrices*. J. Algebra 313 (2007), no. 2, 782–801.
- [13] V. Muñoz, J. Porti, *Geometry of the $SL(3, \mathbb{C})$ -character variety of torus knots*. Algebr. Geom. Topol. 16 (2016), 397–426.
- [14] V. Muñoz, J. Sánchez, *$SL(3, \mathbb{C})$ -character variety of torus knots*. Marco Castrillón López, (2016): 515–526.
- [15] J.R. Parker, *Traces in complex hyperbolic geometry*. In: Geometry, topology and dynamics of character varieties, Lecture notes series (National University of Singapore, Institute for Mathematical Sciences), vol. 23, World Scientific Publishing Company, 2012.
- [16] J.R. Parker, P. Will, *A complex hyperbolic Riley slice*. Geom. Topol. 21 (2017), no. 6, 3391–3451.

- [17] J. Paupert, P. Will, *Real reflections, commutators, and cross-ratios in complex hyperbolic space*. Groups Geom. Dyn. 11 (2017), no. 1, 311–352.
- [18] A.S. Sikora, *Character varieties*. Trans. Amer. Math. Soc. 364 (2012), no. 10, 5173–5208.
- [19] W.P. Thurston, *Three-dimensional manifolds, Kleinian groups and hyperbolic geometry*. Bull. Amer. Math. Soc. (N.S.), 6 (1982), no. 3, 357–381.