Generalized quasi-statistical structures

Adara M. Blaga and Antonella Nannicini

Abstract

Given a non-degenerate (0, 2)-tensor field h on a smooth manifold M, we consider a natural generalized complex and a generalized product structure on the generalized tangent bundle $TM \oplus T^*M$ of M and we show that they are ∇ -integrable, for ∇ an affine connection on M, if and only if (M, h, ∇) is a quasi-statistical manifold. We introduce the notion of generalized quasi-statistical structure and we prove that any quasi-statistical structure on M induces generalized quasi-statistical structures on $TM \oplus T^*M$. In this context, dual connections are considered and some of their properties are established. The results are described in terms of Patterson-Walker and Sasaki metrics on T^*M , horizontal lift and Sasaki metrics on TM and, when the connection ∇ is flat, we define prolongation of quasi-statistical structures on manifolds to their cotangent and tangent bundles via generalized geometry. Moreover, Norden and Para-Norden structures are defined on T^*M and TM.

1 Introduction

Statistical manifolds were introduced in [1], [7]. They are manifolds of probability distributions, used in Information Geometry and related to Codazzi tensors and Affine Geometry. Let h be a pseudo-Riemannian metric and let ∇ be a torsion-free affine connection on a smooth manifold M. Then (M, h, ∇) is called a *statistical manifold* if

²⁰¹⁰ Mathematics Subject Classification. 53C15, 53B05, 53D05.

Key words and phrases. Quasi-statistical structures, generalized geometry, Patterson-Walker metric, Sasaki metric, Norden structures.

 $(\nabla_X h)(Y,Z) = (\nabla_Y h)(X,Z)$, for all $X, Y, Z \in C^{\infty}(TM)$. The definition can be extended to (0,2)-tensor fields and affine connections ∇ with torsion, T^{∇} . In this case, (h,∇) is called a *quasi-statistical structure* on M if $d^{\nabla}h = 0$, where $(d^{\nabla}h)(X,Y,Z) :=$ $(\nabla_X h)(Y,Z) - (\nabla_Y h)(X,Z) + h(T^{\nabla}(X,Y),Z)$, for all $X, Y, Z \in C^{\infty}(TM)$, and the triple (M,h,∇) is called a *quasi-statistical manifold*.

In this paper, given a non-degenerate (0, 2)-tensor field h and an affine connection ∇ on a smooth manifold M, we consider a natural generalized complex and a generalized product structure on the generalized tangent bundle $TM \oplus T^*M$ of M and we show that they are ∇ -integrable if and only if (M, h, ∇) is a quasi-statistical manifold. We introduce the notion of generalized quasi-statistical structure and we prove that any quasi-statistical structure on M, defined by a symmetric or skew-symmetric tensor, induces two natural generalized quasi-statistical structures on $TM \oplus T^*M$. We compute the dual connections and study some of their properties. The results are described in terms of Patterson-Walker and Sasaki metrics on T^*M , horizontal lift and Sasaki metrics on TM. In the case, the connection ∇ is flat we can define prolongation of quasi-statistical structures on manifolds to their cotangent and tangent bundles via generalized geometry. Moreover, in the last section, we construct Norden and Para-Norden structures on T^*M and TM.

2 Quasi-statistical structures and generalized structures induced

Let M be a smooth manifold and h a non-degenerate (0, 2)-tensor field on M. On the generalized tangent bundle $TM \oplus T^*M$ of M, we shall consider the generalized complex structure

(1)
$$\hat{J}_c := \begin{pmatrix} 0 & -h^{-1} \\ h & 0 \end{pmatrix}$$

and the generalized product structure

(2)
$$\hat{J}_p := \begin{pmatrix} 0 & h^{-1} \\ h & 0 \end{pmatrix},$$

where we denoted by h the musical isomorphism, $\flat_h : TM \to T^*M$, $\flat_h(X) := i_X h$, and by h^{-1} its inverse, $\sharp_h : T^*M \to TM$. Let

(3)
$$\langle X + \alpha, Y + \beta \rangle := -\frac{1}{2}(\alpha(Y) + \beta(X))$$

be the natural indefinite metric on $TM\oplus T^*M$ and

(4)
$$(X + \alpha, Y + \beta) := -\frac{1}{2}(\alpha(Y) - \beta(X))$$

be the natural symplectic structure on $TM \oplus T^*M$.

REMARK 2.1. i) If h is symmetric, then:

$$< \hat{J}_c \sigma, \hat{J}_c \tau >= - < \sigma, \tau > \quad and \quad (\hat{J}_c \sigma, \hat{J}_c \tau) = (\sigma, \tau),$$

$$< \hat{J}_p \sigma, \hat{J}_p \tau >= < \sigma, \tau > \quad and \quad (\hat{J}_p \sigma, \hat{J}_p \tau) = -(\sigma, \tau),$$

or, equivalently:

$$< \hat{J}_c \sigma, \tau > = < \sigma, \hat{J}_c \tau > and (\hat{J}_c \sigma, \tau) = -(\sigma, \hat{J}_c \tau),$$
$$< \hat{J}_p \sigma, \tau > = < \sigma, \hat{J}_p \tau > and (\hat{J}_p \sigma, \tau) = -(\sigma, \hat{J}_p \tau),$$

for any $\sigma, \tau \in C^{\infty}(TM \oplus T^*M)$.

i) If h is skew-symmetric, then:

$$\langle \hat{J}_{c}\sigma, \hat{J}_{c}\tau \rangle = \langle \sigma, \tau \rangle \quad and \quad (\hat{J}_{c}\sigma, \hat{J}_{c}\tau) = -(\sigma, \tau),$$
$$\langle \hat{J}_{p}\sigma, \hat{J}_{p}\tau \rangle = -\langle \sigma, \tau \rangle \quad and \quad (\hat{J}_{p}\sigma, \hat{J}_{p}\tau) = (\sigma, \tau),$$

or, equivalently:

$$<\hat{J}_{c}\sigma,\tau>=-<\sigma,\hat{J}_{c}\tau> \quad and \quad (\hat{J}_{c}\sigma,\tau)=(\sigma,\hat{J}_{c}\tau),$$
$$<\hat{J}_{p}\sigma,\tau>=-<\sigma,\hat{J}_{p}\tau> \quad and \quad (\hat{J}_{p}\sigma,\tau)=(\sigma,\hat{J}_{p}\tau),$$

for any $\sigma, \tau \in C^{\infty}(TM \oplus T^*M)$.

On $TM \oplus T^*M$ we consider the bilinear form:

(5)
$$\check{h}(X+\alpha,Y+\beta) := h(X,Y) + h(h^{-1}(\alpha),h^{-1}(\beta)),$$

for any $X, Y \in C^{\infty}(TM)$ and $\alpha, \beta \in C^{\infty}(T^*M)$.

A direct computation gives the following:

LEMMA 2.2. The structures \hat{J}_c and \hat{J}_p satisfy respectively:

$$\begin{split} &\check{h}(\hat{J}_c\sigma,\tau)=2(\sigma,\tau),\\ &\check{h}(\sigma,\hat{J}_p\tau)=2<\sigma,\tau>, \end{split}$$

for any $\sigma, \tau \in C^{\infty}(TM \oplus T^*M)$.

For ∇ an affine connection on M, we consider the bracket $[\cdot, \cdot]_{\nabla}$ on $C^{\infty}(TM \oplus T^*M)$ [6]:

$$[X + \alpha, Y + \beta]_{\nabla} := [X, Y] + \nabla_X \beta - \nabla_Y \alpha,$$

for all $X, Y \in C^{\infty}(TM)$ and $\alpha, \beta \in C^{\infty}(T^*M)$.

A generalized complex or product structure \hat{J} is called ∇ -integrable if its Nijenhuis tensor field $N_{\hat{J}}^{\nabla}$ with respect to ∇ :

$$N_{\hat{J}}^{\nabla}(\sigma,\tau) := [\hat{J}\sigma,\hat{J}\tau]_{\nabla} - \hat{J}[\hat{J}\sigma,\tau]_{\nabla} - \hat{J}[\sigma,\hat{J}\tau]_{\nabla} + \hat{J}^2[\sigma,\tau]_{\nabla}$$

vanishes for all $\sigma, \tau \in C^{\infty}(TM \oplus T^*M)$.

Let M be a smooth manifold with a non-degenerate (0, 2)-tensor field h and an affine connection ∇ .

DEFINITION 2.3. [4] We call (h, ∇) a quasi-statistical structure (respectively, (M, h, ∇) a quasi-statistical manifold) if $d^{\nabla}h = 0$, where

$$(d^{\nabla}h)(X,Y,Z) := (\nabla_X h)(Y,Z) - (\nabla_Y h)(X,Z) + h(T^{\nabla}(X,Y),Z),$$

for any $X, Y, Z \in C^{\infty}(TM)$ and $T^{\nabla}(X, Y) := \nabla_X Y - \nabla_Y X - [X, Y].$

We can state:

PROPOSITION 2.4. The structures \hat{J}_c and \hat{J}_p are integrable if and only if (M, h, ∇) is a quasi-statistical manifold.

PROOF. In this proof we will shortly denote \hat{J}_{\mp} for $\hat{J}_c =: \hat{J}_-$ and $\hat{J}_p =: \hat{J}_+$. Let us compute:

$$N_{\hat{J}_{\mp}}^{\nabla}(X,Y) = [\hat{J}_{\mp}X, \hat{J}_{\mp}Y]_{\nabla} - \hat{J}_{\mp}[\hat{J}_{\mp}X,Y]_{\nabla} - \hat{J}_{\mp}[X, \hat{J}_{\mp}Y]_{\nabla} + \hat{J}_{\mp}^{2}[X,Y]_{\nabla} =$$
$$= \pm h^{-1}((\nabla_{X}h)Y - (\nabla_{Y}h)X + h(\nabla_{X}Y - \nabla_{Y}X - [X,Y])) =$$

$$= \pm h^{-1}((d^{\nabla}h)(X,Y))$$

$$N_{\hat{J}_{\mp}}^{\nabla}(X,h(Y)) = [\hat{J}_{\mp}X,\hat{J}_{\mp}h(Y)]_{\nabla} - \hat{J}_{\mp}[\hat{J}_{\mp}X,h(Y)]_{\nabla} - \hat{J}_{\mp}[X,\hat{J}_{\mp}h(Y)]_{\nabla} + \hat{J}_{\mp}^{2}[X,h(Y)]_{\nabla} =$$

$$= \mp((\nabla_{X}h)Y - (\nabla_{Y}h)X + h(\nabla_{X}Y - \nabla_{Y}X - [X,Y])) =$$

$$= \mp(d^{\nabla}h)(X,Y)$$

$$N_{\hat{J}_{\mp}}^{\nabla}(h(X),h(Y)) = [\hat{J}_{\mp}h(X),\hat{J}_{\mp}h(Y)]_{\nabla} - \hat{J}_{\mp}[\hat{J}_{\mp}h(X),h(Y)]_{\nabla} - \hat{J}_{\mp}[h(X),\hat{J}_{\mp}h(Y)]_{\nabla} +$$

$$+ \hat{J}_{\mp}^{2}[h(X),h(Y)]_{\nabla} =$$

$$= -h^{-1}((\nabla_{X}h)Y - (\nabla_{Y}h)X + h(\nabla_{X}Y - \nabla_{Y}X - [X,Y])) =$$

$$= -h^{-1}((d^{\nabla}h)(X,Y)),$$

for any $X, Y \in C^{\infty}(TM)$. Therefore the proof is complete.

3 Generalized quasi-statistical structures

DEFINITION 3.1. We call $D: C^{\infty}(TM \oplus T^*M) \times C^{\infty}(TM \oplus T^*M) \to C^{\infty}(TM \oplus T^*M)$ an *affine connection* on $TM \oplus T^*M$ if it is \mathbb{R} -bilinear and for any $f \in C^{\infty}(M)$ and $\sigma, \tau \in C^{\infty}(TM \oplus T^*M)$, we have:

- 1. $D_{f\sigma}\tau = f D_{\sigma}\tau$,
- 2. $D_{\sigma}(f\tau) = \sigma(f)\tau + fD_{\sigma}\tau,$

where $(X + \alpha)(f) := X(f)$, for $X + \alpha \in C^{\infty}(TM \oplus T^*M)$.

Let \hat{h} be a non-degenerate (0, 2)-tensor field and D an affine connection on the generalized tangent bundle $TM \oplus T^*M$ of the smooth manifold M.

DEFINITION 3.2. We call (\hat{h}, D) a generalized quasi-statistical structure if $d^D \hat{h} = 0$, where

$$(d^D\hat{h})(\sigma,\tau,\nu) := (D_{\sigma}\hat{h})(\tau,\nu) - (D_{\tau}\hat{h})(\sigma,\nu) + \hat{h}(T^D(\sigma,\tau),\nu),$$

for any $\sigma, \tau, \nu \in C^{\infty}(TM \oplus T^*M)$ and $T^D(\sigma, \tau) := D_{\sigma}\tau - D_{\tau}\sigma - [\sigma, \tau]_{\nabla}$, with ∇ a given connection on M.

3.1 Generalized quasi-statistical structures induced by quasistatistical structures

Let h be a non-degenerate (0, 2)-tensor field and let ∇ be an affine connection on M. We define the affine connection $\hat{\nabla}$ on $TM \oplus T^*M$ by:

(6)
$$\hat{\nabla}_{X+\alpha}Y + \beta := \nabla_X Y + h(\nabla_X h^{-1}(\beta)),$$

for any $X, Y \in C^{\infty}(TM)$ and $\alpha, \beta \in C^{\infty}(T^*M)$.

THEOREM 3.3. $(TM \oplus T^*M, \hat{h}, \hat{\nabla})$ is a generalized quasi-statistical manifold if and only if (M, h, ∇) is a quasi-statistical manifold, where \hat{h} is precisely $\langle \cdot, \cdot \rangle$ or (\cdot, \cdot) given by (3) and (4) respectively, according as h is symmetric or skew-symmetric, and $\hat{\nabla}$ is given by (6).

PROOF. First notice that the torsion of $\hat{\nabla}$ equals to

$$T^{\nabla}(X+\alpha,Y+\beta) := \hat{\nabla}_{X+\alpha}Y+\beta - \hat{\nabla}_{Y+\beta}X+\alpha - [X+\alpha,Y+\beta]_{\nabla} =$$
$$= T^{\nabla}(X,Y) + h(\nabla_X h^{-1}(\beta) - \nabla_Y h^{-1}(\alpha)) - \nabla_X \beta + \nabla_Y \alpha.$$

We have:

$$\begin{split} (d^{\hat{\nabla}}\hat{h})(X+\alpha,Y+\beta,Z+\gamma) &:= (\hat{\nabla}_{X+\alpha}\hat{h})(Y+\beta,Z+\gamma) - (\hat{\nabla}_{Y+\beta}\hat{h})(X+\alpha,Z+\gamma) + \\ &+ \hat{h}(T^{\hat{\nabla}}(X+\alpha,Y+\beta),Z+\gamma) := \\ &:= X(\hat{h}(Y+\beta,Z+\gamma)) - \hat{h}(\hat{\nabla}_{X+\alpha}Y+\beta,Z+\gamma) - \hat{h}(Y+\beta,\hat{\nabla}_{X+\alpha},Z+\gamma) - \\ &- Y(\hat{h}(X+\alpha,Z+\gamma)) + \hat{h}(\hat{\nabla}_{Y+\beta}X+\alpha,Z+\gamma) + \hat{h}(X+\alpha,\hat{\nabla}_{Y+\beta},Z+\gamma) + \\ &+ \hat{h}(T^{\hat{\nabla}}(X+\alpha,Y+\beta),Z+\gamma) := \\ &:= -\frac{1}{2}[X(\beta(Z)\pm\gamma(Y)) - h(\nabla_Xh^{-1}(\beta),Z) \mp \gamma(\nabla_XY) - \beta(\nabla_XZ) \mp h(\nabla_Xh^{-1}(\gamma),Y)] + \\ &+ \frac{1}{2}[Y(\alpha(Z)\pm\gamma(X)) - h(\nabla_Yh^{-1}(\alpha),Z) \mp \gamma(\nabla_YX) - \alpha(\nabla_YZ) \mp h(\nabla_Yh^{-1}(\gamma),X)] - \\ &- \frac{1}{2}[h(\nabla_Xh^{-1}(\beta),Z) - h(\nabla_Yh^{-1}(\alpha),Z) - (\nabla_X\beta)Z + (\nabla_Y\alpha)Z] \mp \frac{1}{2}\gamma(T^{\nabla}(X,Y)) := \\ &:= -\frac{1}{2}[\pm X(\gamma(Y)) \mp \gamma(\nabla_XY) \mp h(\nabla_Xh^{-1}(\gamma),Y) \mp \\ &\mp Y(\gamma(X)) \pm \gamma(\nabla_YX) \pm h(\nabla_Yh^{-1}(\gamma),X) \pm \gamma(T^{\nabla}(X,Y)) := \\ &:= -\frac{1}{2}(d^{\nabla}h)(X,Y,h^{-1}(\gamma)). \end{split}$$

Therefore the proof is complete.

The couple $(\hat{h}, \hat{\nabla})$ with \hat{h} given by (3) or (4) respectively (according as h is symmetric or skew-symmetric) and $\hat{\nabla}$ given by (6) will be called the *generalized quasi-statistical* structure induced by (h, ∇) .

A direct computation gives the expression of the generalized dual quasi-statistical connection of $\hat{\nabla}$, precisely:

PROPOSITION 3.4. Let (M, h, ∇) be a quasi-statistical manifold and let $(\hat{h}, \hat{\nabla})$ be the generalized quasi-statistical structure induced on $TM \oplus T^*M$. Then the generalized dual quasi-statistical connection, $\hat{\nabla}^*$, defined by:

$$\hat{h}(Y+\beta,\hat{\nabla}^*_{X+\alpha}Z+\gamma) = X(\hat{h}(Y+\beta,Z+\gamma)) - \hat{h}(\hat{\nabla}_{X+\alpha}Y+\beta,Z+\gamma),$$

for all $X, Y, Z \in C^{\infty}(TM)$ and $\alpha, \beta, \gamma \in C^{\infty}(T^*M)$, is given by:

$$\hat{\nabla}_{X+\alpha}^* Z + \gamma = h^{-1}(\nabla_X h(Z)) + \nabla_X \gamma.$$

PROOF. From the definition of the generalized dual quasi-statistical connection and using the definition of $\hat{\nabla}$, we get:

$$\hat{h}(Y+\beta,\hat{\nabla}^*_{X+\alpha}Z+\gamma) = X(\beta(Z)) \pm X(\gamma(Y)) - h(\nabla_X h^{-1}(\beta), Z) \mp \gamma(\nabla_X Y),$$

for any $X, Y, Z \in C^{\infty}(TM)$ and $\alpha, \beta, \gamma \in C^{\infty}(T^*M)$.

Let us denote $\hat{\nabla}^*_{X+\alpha}Z + \gamma =: V + \eta$. Then we have:

$$\beta(V) \pm \eta(Y) = X(\beta(Z)) \pm X(\gamma(Y)) - h(\nabla_X h^{-1}(\beta), Z) \mp \gamma(\nabla_X Y),$$

for any $X, Y, Z \in C^{\infty}(TM)$ and $\beta, \gamma \in C^{\infty}(T^*M)$.

Taking $\beta := 0$, we obtain:

$$\eta(Y) = X(\gamma(Y)) - \gamma(\nabla_X Y) := (\nabla_X \gamma)Y$$

and taking Y := 0, we obtain:

$$\beta(V) = X(\beta(Z)) - h(\nabla_X h^{-1}(\beta), Z)$$

which is equivalent to:

$$h(V, h^{-1}(\beta)) = X(h(Z, h^{-1}(\beta))) - h(Z, \nabla_X h^{-1}(\beta)) := (\nabla_X h)(Z, h^{-1}(\beta)) + h(\nabla_X Z, h^{-1}(\beta))$$

and to:

$$h(V) = (\nabla_X h)(Z, \cdot) + h(\nabla_X Z) = \nabla_X h(Z)$$

and to:

$$V = h^{-1}(\nabla_X h(Z)).$$

Therefore the proof is complete.

PROPOSITION 3.5. Let (M, h, ∇) be a quasi-statistical manifold. Then $\hat{\nabla}^*$ is torsion-free.

PROOF. For all $X + \alpha, Y + \beta \in C^{\infty}(TM \oplus T^*M)$, we have:

$$T^{\nabla^*}(X + \alpha, Y + \beta) = \hat{\nabla}^*_{X + \alpha} Y + \beta - \hat{\nabla}^*_{Y + \beta} X + \alpha - [X + \alpha, Y + \beta]_{\nabla} =$$

= $h^{-1}(\nabla_X h(Y)) - h^{-1}(\nabla_Y h(X)) - [X, Y] =$
= $h^{-1}((\nabla_X h)Y - (\nabla_Y h)X + h(T^{\nabla}(X, Y))) =$
= $h^{-1}(d^{\nabla}(X, Y)) = 0.$

Let h be a non-degenerate, symmetric or skew-symmetric (0, 2)-tensor field on M and let ∇ be an affine connection on M. We have the following:

THEOREM 3.6. $(TM \oplus T^*M, \check{h}, \hat{\nabla})$ is a generalized quasi-statistical manifold if and only if (M, h, ∇) is a quasi-statistical manifold, where \check{h} is given by (5) and $\hat{\nabla}$ is given by (6).

PROOF. We have:

$$\begin{split} (d^{\hat{\nabla}}\check{h})(X+\alpha,Y+\beta,Z+\gamma) &:= (\hat{\nabla}_{X+\alpha}\check{h})(Y+\beta,Z+\gamma) - (\hat{\nabla}_{Y+\beta}\check{h})(X+\alpha,Z+\gamma) + \\ &+\check{h}(T^{\hat{\nabla}}(X+\alpha,Y+\beta),Z+\gamma) := \\ &:= X(\check{h}(Y+\beta,Z+\gamma)) - \check{h}(\hat{\nabla}_{X+\alpha}Y+\beta,Z+\gamma) - \check{h}(Y+\beta,\hat{\nabla}_{X+\alpha},Z+\gamma) - \\ &-Y(\check{h}(X+\alpha,Z+\gamma)) + \check{h}(\hat{\nabla}_{Y+\beta}X+\alpha,Z+\gamma) + \check{h}(X+\alpha,\hat{\nabla}_{Y+\beta},Z+\gamma) + \\ &+\check{h}(T^{\hat{\nabla}}(X+\alpha,Y+\beta),Z+\gamma) := \\ &:= X(h(Y,Z)) + X(\beta(h^{-1}(\gamma))) - \check{h}(\nabla_XY+h(\nabla_Xh^{-1}(\beta)),Z+\gamma) - \\ &-\check{h}(Y+\beta,\nabla_XZ+h(\nabla_Xh^{-1}(\gamma))) - \\ &-Y(h(X,Z)) - Y(\alpha(h^{-1}(\gamma))) + \check{h}(\nabla_YX+h(\nabla_Yh^{-1}(\alpha)),Z+\gamma) + \end{split}$$

$$+\check{h}(X+\alpha,\nabla_Y Z+h(\nabla_Y h^{-1}(\gamma)))+h(T^{\nabla}(X,Y),Z)\pm\gamma((\nabla_X h^{-1})\beta-(\nabla_Y h^{-1})\alpha):=$$
$$:=(d^{\nabla}h)(X,Y,Z)+(\nabla_X\beta)h^{-1}(\gamma)-(\nabla_Y\alpha)h^{-1}(\gamma)\pm\gamma(h^{-1}(\nabla_Y\alpha)-h^{-1}(\nabla_X\beta)):=$$
$$:=(d^{\nabla}h)(X,Y,Z),$$

where the sign + is for h symmetric, - is for h skew-symmetric. Therefore the proof is complete.

PROPOSITION 3.7. Let (M, h, ∇) be a quasi-statistical manifold and let $(\check{h}, \hat{\nabla})$ be the generalized quasi-statistical structure induced on $TM \oplus T^*M$. Then the generalized dual quasi-statistical connection, $\hat{\nabla}^*_{\check{h}}$, defined by:

$$\check{h}(Y+\beta,(\hat{\nabla}^*_{\check{h}})_{X+\alpha}Z+\gamma) = X(\check{h}(Y+\beta,Z+\gamma)) - \check{h}(\hat{\nabla}_{X+\alpha}Y+\beta,Z+\gamma),$$

for all $X, Y, Z \in C^{\infty}(TM)$ and $\alpha, \beta, \gamma \in C^{\infty}(T^*M)$, is given by:

$$(\hat{\nabla}^*_{\check{h}})_{X+\alpha}Z + \gamma = h^{-1}(\nabla_X h(Z)) + \nabla_X \gamma.$$

Therefore:

$$\hat{\nabla}^*_{\check{h}} = \hat{\nabla}^*$$

PROOF. We get:

$$\check{h}(Y+\beta,(\hat{\nabla}_{\check{h}}^*)_{X+\alpha}Z+\gamma) = X(h(Y,Z)) + X(\beta(h^{-1}(\gamma))) - h(\nabla_X Y,Z) \mp \gamma(\nabla_X h^{-1}(\beta)) =$$
$$= X(h(Y,Z)) - h(\nabla_X Y,Z) \mp \gamma(\nabla_X h^{-1}(\beta)) \pm X(\gamma(h^{-1}(\beta))),$$

for any $X, Y, Z \in C^{\infty}(TM)$ and $\alpha, \beta, \gamma \in C^{\infty}(T^*M)$.

Let us denote $(\hat{\nabla}^*_{\check{h}})_{X+\alpha}Z + \gamma =: V + \eta$. Then we have:

$$h(Y,V) \pm \eta(h^{-1}(\beta)) = X(h(Y,Z)) - h(\nabla_X Y,Z) \pm (\nabla_X \gamma)h^{-1}(\beta),$$

for any $X, Y, Z \in C^{\infty}(TM)$ and $\beta \in C^{\infty}(T^*M)$.

Taking Y := 0, we obtain:

$$\eta(h^{-1}(\beta)) = \nabla_X h^{-1}(\beta)$$

and taking $\beta := 0$, we obtain:

$$h(Y,V) = (\nabla_X h)(Y,Z) + h(Y,\nabla_X Z)$$

which is equivalent to:

$$h(V) = (\nabla_X h)(Z, \cdot) + h(\nabla_X Z) = \nabla_X h(Z)$$

and to:

$$V = h^{-1}(\nabla_X h(Z)).$$

Therefore the proof is complete.

Given an affine connection D on $TM \oplus T^*M$, we define the curvature operator of D, $R^D : C^{\infty}(TM \oplus T^*M) \times C^{\infty}(TM \oplus T^*M) \times C^{\infty}(TM \oplus T^*M) \to C^{\infty}(TM \oplus T^*M)$, on $\sigma, \tau, \nu \in C^{\infty}(TM \oplus T^*M)$, as in the following:

$$R^{D}(\sigma,\tau)\nu = (D_{\sigma}D_{\tau} - D_{\tau}D_{\sigma} - D_{[\sigma,\tau]_{\nabla}})\nu,$$

where ∇ is a given connection on M.

PROPOSITION 3.8. Let (M, h, ∇) be a quasi-statistical manifold and let $(\hat{h}, \hat{\nabla})$ be the generalized quasi-statistical structure induced on $TM \oplus T^*M$. Then the curvature operators of $\hat{\nabla}$ and $\hat{\nabla}^*$ are given respectively by:

$$R^{\hat{\nabla}}(X+\alpha,Y+\beta)Z+\gamma = R^{\nabla}(X,Y)Z + h(R^{\nabla}(X,Y)h^{-1}(\gamma))$$
$$R^{\hat{\nabla}^*}(X+\alpha,Y+\beta)Z+\gamma = h^{-1}(R^{\nabla}(X,Y)h(Z)) + R^{\nabla}(X,Y)\gamma,$$

where $X, Y, Z \in C^{\infty}(TM)$, $\alpha, \beta, \gamma \in C^{\infty}(T^*M)$ and R^{∇} is the curvature operator of ∇ . In particular, $\hat{\nabla}$ and its dual $\hat{\nabla}^*$ are flat if and only if ∇ is flat.

PROOF. Let us compute:

$$\begin{split} \hat{\nabla}_{X+\alpha} \hat{\nabla}_{Y+\beta} Z + \gamma - \hat{\nabla}_{Y+\beta} \hat{\nabla}_{X+\alpha} Z + \gamma - \hat{\nabla}_{[X+\alpha,Y+\beta]_{\nabla}} Z + \gamma := \\ &:= \hat{\nabla}_{X+\alpha} (\nabla_Y Z + h(\nabla_Y h^{-1}(\gamma))) - \hat{\nabla}_{Y+\beta} (\nabla_X Z + h(\nabla_X h^{-1}(\gamma))) - \\ &- \nabla_{[X,Y]} Z - h(\nabla_{[X,Y]} h^{-1}(\gamma)) := \\ &:= \nabla_X \nabla_Y Z + h(\nabla_X \nabla_Y h^{-1}(\gamma)) - \nabla_Y \nabla_X Z - h(\nabla_Y \nabla_X h^{-1}(\gamma)) - \\ &- \nabla_{[X,Y]} Z - h(\nabla_{[X,Y]} h^{-1}(\gamma)) := \\ &:= R^{\nabla} (X,Y) Z + h(R^{\nabla} (X,Y) h^{-1}(\gamma)) \end{split}$$

and:

$$\hat{\nabla}_{X+\alpha}^* \hat{\nabla}_{Y+\beta}^* Z + \gamma - \hat{\nabla}_{Y+\beta}^* \hat{\nabla}_{X+\alpha}^* Z + \gamma - \hat{\nabla}_{[X+\alpha,Y+\beta]_{\nabla}}^* Z + \gamma :=$$

$$\begin{split} &:= \hat{\nabla}_{X+\alpha}^* (h^{-1}(\nabla_Y h(Z)) + \nabla_Y \gamma) - \hat{\nabla}_{Y+\beta}^* (h^{-1}(\nabla_X h(Z)) + \nabla_X \gamma) - \\ &-h^{-1}(\nabla_{[X,Y]} h(Z)) - \nabla_{[X,Y]} \gamma := \\ &:= h^{-1}(\nabla_X \nabla_Y h(Z)) + \nabla_X \nabla_Y \gamma - h^{-1}(\nabla_Y \nabla_X h(Z)) - \nabla_Y \nabla_X \gamma - \\ &-h^{-1}(\nabla_{[X,Y]} h(Z)) - \nabla_{[X,Y]} \gamma := \\ &:= h^{-1}(R^{\nabla}(X,Y) h(Z)) + R^{\nabla}(X,Y) \gamma. \end{split}$$

Therefore the proof is complete.

PROPOSITION 3.9. The structures \hat{J}_c and \hat{J}_p are $\hat{\nabla}$ -parallel and $\hat{\nabla}^*$ -parallel.

PROOF. In this proof we will shortly denote \hat{J}_{\mp} for $\hat{J}_c =: \hat{J}_-$ and $\hat{J}_p =: \hat{J}_+$. Let us compute:

$$(\hat{\nabla}_{X+\alpha}\hat{J}_{\mp})Y + \beta := \hat{\nabla}_{X+\alpha}(\mp h^{-1}(\beta) + h(Y)) - \hat{J}_{\mp}(\hat{\nabla}_{X+\alpha}Y + \beta) :=$$

$$:= \mp \nabla_X h^{-1}(\beta) + h(\nabla_X h^{-1}(h(Y))) - \hat{J}_{\mp}(\nabla_X Y + h(\nabla_X h^{-1}(\beta))) :=$$

$$:= \mp \nabla_X h^{-1}(\beta) + h(\nabla_X Y) \pm h^{-1}(h(\nabla_X h^{-1}(\beta))) - h(\nabla_X Y) = 0;$$

moreover:

$$\begin{aligned} (\hat{\nabla}_{X+\alpha}^* \hat{J}_{\mp})Y + \beta &:= \hat{\nabla}_{X+\alpha}^* (\mp h^{-1}(\beta) + h(Y)) - \hat{J}_{\mp}(\hat{\nabla}_{X+\alpha}^* Y + \beta) := \\ &:= \mp h^{-1}(\nabla_X \beta) + \nabla_X h(Y) - \hat{J}_{\mp}(h^{-1}(\nabla_X h(Y)) + \nabla_X \beta) := \\ &:= \mp h^{-1}(\nabla_X \beta) + \nabla_X h(Y) - \nabla_X h(Y) \pm h^{-1}(\nabla_X \beta) = 0, \end{aligned}$$

for any $X, Y \in C^{\infty}(TM)$ and $\alpha, \beta \in C^{\infty}(T^*M)$. Therefore the proof is complete.

3.2 Generalized quasi-statistical structures induced by torsionfree connections

Another affine connection on the generalized tangent bundle $TM \oplus T^*M$ is naturally defined by an affine connection ∇ on M by:

(7)
$$\check{\nabla}_{X+\alpha}Y + \beta := \nabla_X Y + \nabla_X \beta,$$

for any $X, Y \in C^{\infty}(TM)$ and $\alpha, \beta \in C^{\infty}(T^*M)$.

REMARK 3.10. One can check that if h is a non-degenerate (0, 2)-tensor field on Mwhich is ∇ -parallel, then the connections $\hat{\nabla}$ and $\check{\nabla}$ coincide (since we have $\hat{\nabla}_{X+\alpha}Y + \beta - \check{\nabla}_{X+\alpha}Y + \beta = -(\nabla_X h)(h^{-1}(\beta), \cdot)$, for any $X, Y \in C^{\infty}(TM)$ and $\alpha, \beta \in C^{\infty}(T^*M)$). In particular, $\hat{\nabla}^* = \hat{\nabla} = \check{\nabla}$.

We have the following:

PROPOSITION 3.11. $(TM \oplus T^*M, \hat{h}, \check{\nabla})$ is a generalized quasi-statistical manifold if and only if ∇ is torsion-free, where \hat{h} is precisely $\langle \cdot, \cdot \rangle$ or (\cdot, \cdot) given by (3) and (4) respectively and $\check{\nabla}$ is given by (7).

PROOF. First notice that the torsion of $\check{\nabla}$ equals to

$$T^{\nabla}(X+\alpha,Y+\beta) := \check{\nabla}_{X+\alpha}Y + \beta - \check{\nabla}_{Y+\beta}X + \alpha - [X+\alpha,Y+\beta]_{\nabla} = T^{\nabla}(X,Y).$$

We have:

$$\begin{split} (d^{\bar{\nabla}}\hat{h})(X+\alpha,Y+\beta,Z+\gamma) &:= (\check{\nabla}_{X+\alpha}\hat{h})(Y+\beta,Z+\gamma) - (\check{\nabla}_{Y+\beta}\hat{h})(X+\alpha,Z+\gamma) + \\ &\quad + \hat{h}(T^{\bar{\nabla}}(X+\alpha,Y+\beta),Z+\gamma) := \\ &:= X(\hat{h}(Y+\beta,Z+\gamma)) - \hat{h}(\check{\nabla}_{X+\alpha}Y+\beta,Z+\gamma) - \hat{h}(Y+\beta,\check{\nabla}_{X+\alpha},Z+\gamma) - \\ &- Y(\hat{h}(X+\alpha,Z+\gamma)) + \hat{h}(\check{\nabla}_{Y+\beta}X+\alpha,Z+\gamma) + \hat{h}(X+\alpha,\check{\nabla}_{Y+\beta},Z+\gamma) + \\ &\quad + \hat{h}(T^{\bar{\nabla}}(X+\alpha,Y+\beta),Z+\gamma) := \\ &:= -\frac{1}{2}[X(\beta(Z)\pm\gamma(Y)) - (\nabla_X\beta)Z\mp\gamma(\nabla_XY) - \beta(\nabla_XZ)\mp(\nabla_X\gamma)Y] + \\ &\quad + \frac{1}{2}[Y(\alpha(Z)\pm\gamma(X)) - (\nabla_Y\alpha)Z\mp\gamma(\nabla_YX) - \alpha(\nabla_YZ)\mp(\nabla_Y\gamma)X]\mp\frac{1}{2}\gamma(T^{\nabla}(X,Y)) := \\ &:= -\frac{1}{2}[X(\beta(Z))\pm X(\gamma(Y)) - X(\beta(Z)) + \beta(\nabla_XZ)\mp\gamma(\nabla_XY) - \beta(\nabla_XZ)\mp X(\gamma(Y))\pm\gamma(\nabla_XY)] + \\ &\quad + \frac{1}{2}[Y(\alpha(Z))\pm Y(\gamma(X)) - Y(\alpha(Z)) + \alpha(\nabla_YZ)\mp\gamma(\nabla_YX) - \alpha(\nabla_YZ)\mp Y(\gamma(X))\pm\gamma(\nabla_YX)] \mp \\ &\quad \mp\frac{1}{2}\gamma(T^{\nabla}(X,Y)) = \mp\frac{1}{2}\gamma(T^{\nabla}(X,Y)). \end{split}$$

Therefore the proof is complete.

PROPOSITION 3.12. Let ∇ be a torsion-free affine connection on M and let $(\langle \cdot, \cdot \rangle, \check{\nabla})$ and $((\cdot, \cdot), \check{\nabla})$ be the canonical generalized quasi-statistical structures defined in Proposition 3.11. Then $\check{\nabla}$ and its generalized dual quasi-statistical connection, $\check{\nabla}^*$, coincide.

PROOF. Let us denote $\check{\nabla}^*_{X+\alpha}Z + \gamma =: V + \eta$. From the definition of the generalized dual quasi-statistical connection and using the definition of $\check{\nabla}$, we get:

$$\beta(V) \pm \eta(Y) = X(\beta(Z)) \pm X(\gamma(Y)) - (\nabla_X \beta)Z \mp \gamma(\nabla_X Y),$$

for any $X, Y, Z \in C^{\infty}(TM)$ and $\beta, \gamma \in C^{\infty}(T^*M)$.

Taking $\beta := 0$, we obtain:

$$\pm \eta(Y) = \pm X(\gamma(Y)) \mp \gamma(\nabla_X Y) := \pm (\nabla_X \gamma) Y$$

and taking Y := 0, we obtain:

$$\beta(V) = X(\beta(Z)) - (\nabla_X \beta)Z := \beta(\nabla_X Z).$$

Therefore the proof is complete.

PROPOSITION 3.13. If ∇ is a torsion-free affine connection and h is a ∇ -parallel (0,2)-tensor field on M, then $(\check{h},\check{\nabla})$ is a generalized quasi-statistical structure, where \check{h} is given by (5) and $\check{\nabla}$ is given by (7).

PROOF. We have:

$$\begin{split} (d^{\tilde{\nabla}}\check{h})(X+\alpha,Y+\beta,Z+\gamma) &:= (\check{\nabla}_{X+\alpha}\check{h})(Y+\beta,Z+\gamma) - (\check{\nabla}_{Y+\beta}\check{h})(X+\alpha,Z+\gamma) + \\ &+\check{h}(T^{\tilde{\nabla}}(X+\alpha,Y+\beta),Z+\gamma) := \\ &:= X(\check{h}(Y+\beta,Z+\gamma)) - \check{h}(\check{\nabla}_{X+\alpha}Y+\beta,Z+\gamma) - \check{h}(Y+\beta,\check{\nabla}_{X+\alpha},Z+\gamma) - \\ &-Y(\check{h}(X+\alpha,Z+\gamma)) + \check{h}(\check{\nabla}_{Y+\beta}X+\alpha,Z+\gamma) + \check{h}(X+\alpha,\check{\nabla}_{Y+\beta},Z+\gamma) + \\ &+\check{h}(T^{\tilde{\nabla}}(X+\alpha,Y+\beta),Z+\gamma) := \\ &:= X(h(Y,Z) + h(h^{-1}(\beta),h^{-1}(\gamma))) - \\ &-h(\nabla_XY,Z) - h(h^{-1}(\nabla_X\beta),h^{-1}(\gamma)) - h(Y,\nabla_XZ) - h(h^{-1}(\beta),h^{-1}(\nabla_X\gamma)) - \\ &-Y(h(X,Z) + h(h^{-1}(\alpha),h^{-1}(\gamma))) + \\ &+h(\nabla_YX,Z) + h(h^{-1}(\nabla_Y\alpha),h^{-1}(\gamma)) + h(X,\nabla_YZ) + h(h^{-1}(\alpha),h^{-1}(\nabla_Y\gamma)) + h(T^{\nabla}(X,Y),Z) := \\ &:= (\nabla_Xh)(Y,Z) - (\nabla_Yh)(X,Z) + h(T^{\nabla}(X,Y),Z) + \\ &+X(\beta(h^{-1}(\gamma))) - (\nabla_X\beta)h^{-1}(\gamma) - \beta(h^{-1}(\nabla_X\gamma)) - \\ &-Y(\alpha(h^{-1}(\gamma))) + (\nabla_Y\alpha)h^{-1}(\gamma) + \alpha(h^{-1}(\nabla_Y\gamma)) = \\ \end{split}$$

$$= (\nabla_X h)(Y, Z) - (\nabla_Y h)(X, Z) + h(T^{\nabla}(X, Y), Z) + \\ + \beta(\nabla_X h^{-1}(\gamma)) - \beta(h^{-1}(\nabla_X \gamma)) - \alpha(\nabla_Y h^{-1}(\gamma)) + \alpha(h^{-1}(\nabla_Y \gamma)).$$

Also, for any $V \in C^{\infty}(TM)$, we have:

$$h(h^{-1}(\nabla_X \gamma) - \nabla_X h^{-1}(\gamma), V) = h(h^{-1}(\nabla_X \gamma), V) - h(\nabla_X h^{-1}(\gamma), V) =$$

= $(\nabla_X \gamma)V - h(\nabla_X h^{-1}(\gamma), V) := X(\gamma(V)) - \gamma(\nabla_X V) - h(\nabla_X h^{-1}(\gamma), V) =$
= $X(h(h^{-1}(\gamma), V)) - h(h^{-1}(\gamma), \nabla_X V) - h(\nabla_X h^{-1}(\gamma), V) := (\nabla_X h)(h^{-1}(\gamma), V) = 0,$

hence $h^{-1}(\nabla_X \gamma) - \nabla_X h^{-1}(\gamma) = 0$, for any $X \in C^{\infty}(TM)$ and $\gamma \in C^{\infty}(T^*M)$. Therefore, $d^{\check{\nabla}}\check{h} = 0$ and the proof is complete.

PROPOSITION 3.14. Let (M, h, ∇) be a quasi-statistical manifold with ∇ a torsion-free affine connection, $h \ a \ \nabla$ -parallel (0, 2)-tensor field on M and let $(\check{h}, \check{\nabla})$ be the generalized quasi-statistical structure on $TM \oplus T^*M$, with \check{h} given by (5) and $\check{\nabla}$ given by (7). Then $\check{\nabla}$ and its generalized dual quasi-statistical connection, $\check{\nabla}^*_{\check{h}}$, coincide.

PROOF. We get:

$$\check{h}(Y + \beta, (\check{\nabla}_{\check{h}}^{*})_{X+\alpha}Z + \gamma) = X(h(Y, Z)) + X(h(h^{-1}(\beta), h^{-1}(\gamma))) - -h(\nabla_X Y, Z) - h(h^{-1}(\nabla_X \beta), h^{-1}(\gamma)) = = h(Y, \nabla_X Z) + \beta(\nabla_X h^{-1}(\gamma)),$$

for any $X, Y, Z \in C^{\infty}(TM)$ and $\alpha, \beta, \gamma \in C^{\infty}(T^*M)$. Let us denote $(\check{\nabla}^*_{\check{h}})_{X+\alpha}Z + \gamma =: V + \eta$. Then we have:

$$h(Y,V) + h(h^{-1}(\beta), h^{-1}(\eta)) = h(Y, \nabla_X Z) + \beta(\nabla_X h^{-1}(\gamma)),$$

for any $X, Y, Z \in C^{\infty}(TM)$ and $\beta, \gamma \in C^{\infty}(T^*M)$.

Taking Y := 0, we obtain:

$$\beta(h^{-1}(\eta)) = \beta(\nabla_X h^{-1}(\gamma))$$

which is equivalent to:

$$\eta = h(\nabla_X h^{-1}(\gamma))$$

and taking $\beta := 0$, we obtain:

$$h(Y,V) = h(Y,\nabla_X Z)$$

which is equivalent to:

$$V = \nabla_X Z.$$

From Remark 3.10 we get $\check{\nabla}^*_{\check{h}} = \hat{\nabla} = \check{\nabla}$. Therefore the proof is complete.

 \Box

4 The pull-back tensors on $TM \oplus T^*M$ of horizontal lifts, Sasaki and Patterson-Walker metrics

4.1 Patterson-Walker and Sasaki metrics on T^*M

Let M be a smooth manifold and let ∇ be an affine connection on M.

Let $\pi : T^*M \to M$ be the canonical projection and $\pi_* : T(T^*M) \to TM$ be the tangent map of π . If $a \in T^*M$ and $A \in T_a(T^*M)$, then $\pi_*(A) \in T_{\pi(a)}M$ and we denote by χ_a the standard identification between $T^*_{\pi(a)}M$ and its tangent space $T_a(T^*_{\pi(a)}M)$.

Let $\Phi^{\nabla}: TM \oplus T^*M \to T(T^*M)$ be the bundle morphism defined by [5]:

(8)
$$\Phi^{\nabla}(X+\alpha) := X_a^H + \chi_a(\alpha)$$

where $a \in T^*M$ and X_a^H is the horizontal lift of $X \in T_{\pi(a)}M$.

Let $\{x^1, ..., x^n\}$ be local coordinates on M, let $\{\tilde{x}^1, ..., \tilde{x}^n, y_1, ..., y_n\}$ be respectively the corresponding local coordinates on T^*M and let $\{X_1, ..., X_n, \frac{\partial}{\partial y_1}, ..., \frac{\partial}{\partial y_n}\}$ be a local frame on $T(T^*M)$, where $X_i = \frac{\partial}{\partial \tilde{x}^i}$. The horizontal lift of $\frac{\partial}{\partial x^i}$ is defined by:

$$\left(\frac{\partial}{\partial x^i}\right)^H := X_i + y_k \Gamma^k_{il} \frac{\partial}{\partial y_l}$$

and we will denote $X_i^H =: (\frac{\partial}{\partial x^i})^H$. Moreover, the vertical lift of $\frac{\partial}{\partial x^i}$ is defined by:

$$(\frac{\partial}{\partial x^i})^V := \frac{\partial}{\partial y_i}$$

where i, k, l run from 1 to n and Γ_{il}^k are the Christoffel's symbols of ∇ .

Let $\Phi^{\nabla} : TM \oplus T^*M \to T(T^*M)$ be the bundle morphism defined before (which is an isomorphism on the fibres). In local coordinates, we have the following expressions:

$$\Phi^{\nabla}\left(\frac{\partial}{\partial x^{i}}\right) = X_{i}^{H}$$
$$\Phi^{\nabla}\left(dx^{j}\right) = \frac{\partial}{\partial y_{j}}.$$

In [9], starting from a torsion-free affine connection on M, the Patterson-Walker metric, \tilde{h} , on T^*M is defined as in the following:

$$\tilde{h}(X^H, Y^H) = 0$$

$$\tilde{h}(X^V, Y^V) = 0$$

 $\tilde{h}(Y^V, X^H) = \tilde{h}(X^H, Y^V) = ((\Phi^{\nabla})^{-1}(Y^V))(X)$

where $X, Y \in C^{\infty}(T^*M), X^H, Y^H$ are the horizontal lifts and X^V, Y^V are the vertical lifts of X, Y respectively.

The definition can also be given if ∇ has torsion and we define \tilde{h}_{\pm} on T^*M as in the following:

$$\tilde{h}_{\pm}(X^{H}, Y^{H}) = 0$$
$$\tilde{h}_{\pm}(X^{V}, Y^{V}) = 0$$
$$\tilde{h}_{\pm}(Y^{V}, X^{H}) = ((\Phi^{\nabla})^{-1}(Y^{V}))(X)$$
$$\tilde{h}_{\pm}(X^{H}, Y^{V}) = \pm ((\Phi^{\nabla})^{-1}(Y^{V}))(X),$$

where $X, Y \in C^{\infty}(T^*M)$, X^H, Y^H are the horizontal lifts and X^V, Y^V are the vertical lifts of X, Y respectively.

We denote by \tilde{h}_{\pm} the pull-back tensors of \tilde{h}_{\pm} on $TM \oplus T^*M$:

$$\tilde{\tilde{h}}_{\pm}(\sigma,\tau) := (\Phi^{\nabla})^*(\tilde{h}_{\pm})(\sigma,\tau) := \tilde{h}_{\pm}(\Phi^{\nabla}(\sigma),\Phi^{\nabla}(\tau)),$$

for any $\sigma, \tau \in C^{\infty}(TM \oplus T^*M)$. Remark that $\tilde{\tilde{h}}_{\pm}$ are related to the indefinite metric or to the symplectic structure of $TM \oplus T^*M$ as follows.

PROPOSITION 4.1.

$$\begin{split} \tilde{\tilde{h}}_+ &= -2 < \cdot, \cdot > \\ \tilde{\tilde{h}}_- &= -2(\cdot, \cdot). \end{split}$$

PROOF. Let $\sigma = X + \alpha$, $\tau = Y + \beta$, $X, Y \in C^{\infty}(TM)$, $\alpha, \beta \in C^{\infty}(T^*M)$. Then:

$$\tilde{\tilde{h}}_{\pm}(\sigma,\tau) = \tilde{h}_{\pm}(X^H + \Phi^{\nabla}(\alpha), Y^H + \Phi^{\nabla}(\beta)) =$$
$$= \tilde{h}_{\pm}(\Phi^{\nabla}(\alpha), Y^H) + \tilde{h}_{\pm}(X^H, \Phi^{\nabla}(\beta)) =$$
$$= \alpha(Y) \pm \beta(X).$$

Then we get the statement.

16

Let h be a non-degenerate (0, 2)-tensor field on M. The Sasaki (0, 2)-tensor field h^{S^*} on T^*M , with respect to ∇ , is naturally defined by:

$$h^{S^{*}}(X^{H}, Y^{H}) = h(X, Y)$$
$$h^{S^{*}}(\alpha^{V}, \beta^{V}) = h(h^{-1}(\alpha), h^{-1}(\beta))$$
$$h^{S^{*}}(\alpha^{V}, Y^{H}) = 0,$$

where $X, Y \in C^{\infty}(TM)$, $\alpha, \beta \in C^{\infty}(T^*M)$, X^H, Y^H are the horizontal lifts of X, Y and α^V, β^V are the vertical lifts of α, β respectively.

We denote by \tilde{h}^{S^*} the pull-back tensor of h^{S^*} on $TM \oplus T^*M$:

$$\tilde{h}^{S^*}(\sigma,\tau) := (\Phi^{\nabla})^*(h^{S^*})(\sigma,\tau) := h^{S^*}(\Phi^{\nabla}(\sigma), \Phi^{\nabla}(\tau)),$$

for any $\sigma, \tau \in C^{\infty}(TM \oplus T^*M)$.

PROPOSITION 4.2. If h is a non-degenerate (0, 2)-tensor field on M, then:

$$\tilde{h}^{S^*}(X + \alpha, Y + \beta) = \hat{h}(X + \alpha, Y + \beta) = h(X, Y) + h(h^{-1}(\alpha), h^{-1}(\beta)),$$

for any $X, Y \in C^{\infty}(TM)$ and $\alpha, \beta \in C^{\infty}(T^*M)$.

PROOF. Let $\sigma = X + \alpha$, $\tau = Y + \beta$, $X, Y \in C^{\infty}(TM)$, $\alpha, \beta \in C^{\infty}(T^*M)$. Then:

$$\begin{split} \tilde{h}^{S^*}(\sigma,\tau) &= h^{S^*}(X^H + \Phi^{\nabla}(\alpha), Y^H + \Phi^{\nabla}(\beta)) = \\ &= h^{S^*}(X^H, Y^H) + h^{S^*}(\Phi^{\nabla}(\alpha), \Phi^{\nabla}(\beta)) = \\ &= h(X,Y) + h^{S^*}(\Phi^{\nabla}(\alpha), \Phi^{\nabla}(\beta)). \end{split}$$

In local coordinates, let $\alpha = \alpha_k dx^k$, $\beta = \beta_l dx^l$ and we get:

$$h^{S^*}(\Phi^{\nabla}(\alpha), \Phi^{\nabla}(\beta)) = h^{S^*}(\alpha_k \frac{\partial}{\partial y_k}, \beta_l \frac{\partial}{\partial y_l}) =$$
$$= \alpha_k \beta_l h_{kl} = h(h^{-1}(\alpha), h^{-1}(\beta)).$$

Then we get the statement.

4.2 Horizontal lift and Sasaki metrics on TM

Let M be a smooth manifold, let h be a non-degenerate (0, 2)-tensor field on M, and let ∇ be an affine connection on M. The horizontal lift h^H of h on TM with respect to ∇ is defined by:

$$h^{H}(X^{H}, Y^{H}) = 0$$
$$h^{H}(X^{V}, Y^{V}) = 0$$
$$h^{H}(X^{H}, Y^{V}) = h(X, Y)$$

where $X, Y \in C^{\infty}(TM), X^{H}, Y^{H}$ are the horizontal lifts and X^{V}, Y^{V} are the vertical lifts of X, Y respectively.

Let $\pi : TM \to M$ be the canonical projection and $\pi_* : T(TM) \to TM$ be the tangent map of π . If $a \in TM$ and $A \in T_a(TM)$, then $\pi_*(A) \in T_{\pi(a)}M$ and we denote by χ_a the standard identification between $T_{\pi(a)}M$ and its tangent space $T_a(T_{\pi(a)}M)$.

Let $\Psi^{\nabla} : TM \oplus T^*M \to T(TM)$ be the bundle morphism defined by:

(9)
$$\Psi^{\nabla}(X+\alpha) := X_a^H + \chi_a(h^{-1}(\alpha)),$$

where $a \in TM$ and X_a^H is the horizontal lift of $X \in T_{\pi(a)}M$.

Let $\{x^1, ..., x^n\}$ be local coordinates on M, let $\{\tilde{x}^1, ..., \tilde{x}^n, y^1, ..., y^n\}$ be respectively the corresponding local coordinates on TM and let $\{X_1, ..., X_n, \frac{\partial}{\partial y^1}, ..., \frac{\partial}{\partial y^n}\}$ be a local frame on T(TM), where $X_i = \frac{\partial}{\partial \tilde{x}^i}$. The horizontal lift of $\frac{\partial}{\partial x^i}$ is defined by:

$$(\frac{\partial}{\partial x^i})^H := X_i - y^k \Gamma^l_{ik} \frac{\partial}{\partial y^l}$$

and we will denote $X_i^H =: (\frac{\partial}{\partial x^i})^H$. Moreover, the vertical lift of $\frac{\partial}{\partial x^i}$ is defined by:

$$(\frac{\partial}{\partial x^i})^V := \frac{\partial}{\partial y^i},$$

where i, k, l run from 1 to n and Γ_{il}^k are the Christoffel's symbols of ∇ .

Let $\Psi^{\nabla} : TM \oplus T^*M \to T(TM)$ be the bundle morphism defined before (which is an isomorphism on the fibres). In local coordinates, we have the following expressions:

$$\Psi^{\nabla}\left(\frac{\partial}{\partial x^i}\right) = X_i^H$$

Generalized quasi-statistical structures

$$\Psi^{\nabla}\left(dx^{j}\right) = h^{jk}\frac{\partial}{\partial y^{k}}.$$

We denote by \overline{h} the pull-back tensor of h^H on $TM \oplus T^*M$:

$$\bar{h}(\sigma,\tau):=(\Psi^{\nabla})^*(h^H)(\sigma,\tau):=h^H(\Psi^{\nabla}(\sigma),\Psi^{\nabla}(\tau)),$$

for any $\sigma, \tau \in C^{\infty}(TM \oplus T^*M)$. Remark that \bar{h} is related to the indefinite metric or to the symplectic structure of $TM \oplus T^*M$ as follows.

PROPOSITION 4.3. If h is a symmetric tensor, then:

$$\bar{h} = -2 < \cdot, \cdot > .$$

If h is a skew-symmetric tensor, then:

$$\bar{h} = -2(\cdot, \cdot).$$

PROOF. Let $\sigma = X + \alpha$, $\tau = Y + \beta$, $X, Y \in C^{\infty}(TM)$, $\alpha, \beta \in C^{\infty}(T^*M)$. Then:

$$\bar{h}(\sigma,\tau) = h^H (X^H + \Psi^{\nabla}(\alpha), Y^H + \Psi^{\nabla}(\beta)) =$$
$$= h^H (\Psi^{\nabla}(\alpha), Y^H) + h^H (X^H, \Psi^{\nabla}(\beta)).$$

In local coordinates, let $X = X^i \frac{\partial}{\partial x^i}$, $Y = Y^j \frac{\partial}{\partial x^j}$, $\alpha = \alpha_k dx^k$, $\beta = \beta_l dx^l$ and we get:

$$\bar{h}(\sigma,\tau) = h^{H}(\alpha_{k}h^{kr}\frac{\partial}{\partial y^{r}}, Y^{j}X^{H}_{j}) + h^{H}(X^{i}X^{H}_{i}, \beta_{l}h^{ls}\frac{\partial}{\partial y^{s}}) =$$
$$= \alpha_{k}Y^{j}h^{kr}h_{rj} + X^{i}\beta_{l}h^{ls}h_{is} = \alpha_{k}Y^{j}\delta^{k}_{j} \pm X^{i}\beta_{l}\delta^{l}_{i} =$$
$$= \alpha(Y) \pm \beta(X),$$

where we denoted by δ the Kronecker's symbol and the sign + is for h symmetric, - is for h skew-symmetric. Then we get the statement.

The Sasaki (0,2)-tensor field h^S on TM, with respect to ∇ , is naturally defined by:

$$h^{S}(X^{H}, Y^{H}) = h(X, Y)$$
$$h^{S}(X^{V}, Y^{V}) = h(X, Y)$$
$$h^{S}(X^{H}, Y^{V}) = 0,$$

where $X, Y \in C^{\infty}(TM), X^{H}, Y^{H}$ are the horizontal lifts and X^{V}, Y^{V} are the vertical lifts of X, Y respectively.

We denote by \bar{h}^S the pull-back tensor of h^S on $TM \oplus T^*M$:

$$\bar{h}^S(\sigma,\tau):=(\Psi^\nabla)^*(h^S)(\sigma,\tau):=h^S(\Psi^\nabla(\sigma),\Psi^\nabla(\tau)),$$

for any $\sigma, \tau \in C^{\infty}(TM \oplus T^*M)$.

PROPOSITION 4.4. If h is a non-degenerate (0, 2)-tensor field on M, then:

$$\bar{h}^S(X+\alpha,Y+\beta) = \check{h}(X+\alpha,Y+\beta) = h(X,Y) + h(h^{-1}(\alpha),h^{-1}(\beta)),$$

for any $X, Y \in C^{\infty}(TM)$ and $\alpha, \beta \in C^{\infty}(T^*M)$.

PROOF. Let
$$\sigma = X + \alpha, \tau = Y + \beta, X, Y \in C^{\infty}(TM), \alpha, \beta \in C^{\infty}(T^*M)$$
. Then:
 $\bar{h}^S(\sigma, \tau) = h^S(X^H + \Psi^{\nabla}(\alpha), Y^H + \Psi^{\nabla}(\beta)) =$
 $= h^S(X^H, Y^H) + h^S(\Psi^{\nabla}(\alpha), \Psi^{\nabla}(\beta)) =$
 $= h(X, Y) + h^S(\Psi^{\nabla}(\alpha), \Psi^{\nabla}(\beta)).$

In local coordinates, let $\alpha = \alpha_k dx^k$, $\beta = \beta_l dx^l$ and we get:

$$h^{S}(\Psi^{\nabla}(\alpha),\Psi^{\nabla}(\beta)) = h^{S}(\alpha_{k}h^{kr}\frac{\partial}{\partial y^{r}},\beta_{l}h^{ls}\frac{\partial}{\partial y^{s}}) =$$
$$= \alpha_{k}\beta_{l}h^{kr}h^{ls}h_{rs} = h(h^{-1}(\alpha),h^{-1}(\beta)).$$

Then we get the statement.

4.3 Quasi-statistical structures on cotangent bundles

Given an affine connection on M, the splitting in horizontal and vertical subbundles identifies $T(T^*M)$ with the pull-back bundle $\pi^*(TM \oplus T^*M)$, where $\pi : T^*M \to M$ is the canonical projection map. In particular, given a connection on $TM \oplus T^*M$, we can define the pull-back connection on $\pi^*(TM \oplus T^*M)$.

A direct computation gives the following:

PROPOSITION 4.5. The pull-back connection $\tilde{\nabla}$ of $\hat{\nabla}$ on T^*M is defined, in local coordinates, by:

$$\begin{split} \tilde{\nabla}_{(\frac{\partial}{\partial x^{i}})^{H}} (\frac{\partial}{\partial x^{j}})^{H} &= \Gamma_{ij}^{k} (\frac{\partial}{\partial x^{k}})^{H} \\ \tilde{\nabla}_{(\frac{\partial}{\partial x^{i}})^{H}} \frac{\partial}{\partial y_{j}} &= (\frac{\partial h^{jk}}{\partial x^{i}} + h^{jl} \Gamma_{il}^{k}) h_{rk} \frac{\partial}{\partial y_{r}} \\ \tilde{\nabla}_{\frac{\partial}{\partial y_{j}}} (\frac{\partial}{\partial x^{i}})^{H} &= 0 \\ \tilde{\nabla}_{\frac{\partial}{\partial y_{i}}} \frac{\partial}{\partial y_{j}} &= 0. \end{split}$$

In local coordinates, the torsion $T^{\tilde{\nabla}}$ of $\tilde{\nabla}$ is:

$$T^{\tilde{\nabla}}((\frac{\partial}{\partial x^{i}})^{H},(\frac{\partial}{\partial x^{j}})^{H}) = (\Gamma_{ij}^{k} - \Gamma_{ji}^{k})(\frac{\partial}{\partial x^{k}})^{H} - y_{l}R_{ijk}^{l}\frac{\partial}{\partial y_{k}}$$
$$T^{\tilde{\nabla}}(\frac{\partial}{\partial y_{i}},(\frac{\partial}{\partial x^{j}})^{H}) = -((\frac{\partial h^{ik}}{\partial x^{j}} + h^{il}\Gamma_{jl}^{k})h_{rk} + \Gamma_{jk}^{i})\frac{\partial}{\partial y_{r}}$$
$$T^{\tilde{\nabla}}(\frac{\partial}{\partial y_{i}},\frac{\partial}{\partial y_{j}}) = 0$$

and the curvature $R^{\tilde{\nabla}}$ of $\tilde{\nabla}$, which is the pull-back of $R^{\hat{\nabla}}$, is:

$$\begin{split} R^{\tilde{\nabla}}(\frac{\partial}{\partial y_i}, \frac{\partial}{\partial y_j}) &= 0\\ R^{\tilde{\nabla}}((\frac{\partial}{\partial x^i})^H, \frac{\partial}{\partial y_j}) &= 0\\ R^{\tilde{\nabla}}((\frac{\partial}{\partial x^i})^H, (\frac{\partial}{\partial x^j})^H)\frac{\partial}{\partial y_k} &= h^{kr}R^l_{ijr}h_{ls}\frac{\partial}{\partial y_s}\\ R^{\tilde{\nabla}}((\frac{\partial}{\partial x^i})^H, (\frac{\partial}{\partial x^j})^H)(\frac{\partial}{\partial x^k})^H &= (R^{\nabla}(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j})\frac{\partial}{\partial x^k})^H. \end{split}$$

Therefore we get:

PROPOSITION 4.6. ∇ is flat if and only if $\tilde{\nabla}$ is flat.

THEOREM 4.7. Let (M, h, ∇) be a quasi-statistical manifold such that ∇ is flat. Then $(T^*M, h^{S^*}, \tilde{\nabla})$ is a flat quasi-statistical manifold.

PROOF. Let us compute $d^{\tilde{\nabla}}h^{S^*}$. From the definition of h^{S^*} and $\tilde{\nabla}$ we get immediately:

$$(d^{\tilde{\nabla}}h^{S^*})(\frac{\partial}{\partial y_i},\frac{\partial}{\partial y_j}) = 0$$

$$(d^{\tilde{\nabla}}h^{S^*})((\frac{\partial}{\partial x^i})^H,\frac{\partial}{\partial y_j}) = 0$$

$$(d^{\tilde{\nabla}}h^{S^*})((\frac{\partial}{\partial x^i})^H,(\frac{\partial}{\partial x^j})^H)(\frac{\partial}{\partial y_k}) = -y_l R^l_{ijr}h^{kr}$$

$$(d^{\tilde{\nabla}}h^{S^*})((\frac{\partial}{\partial x^i})^H,(\frac{\partial}{\partial x^j})^H)(\frac{\partial}{\partial x^k})^H = (d^{\nabla}h)(\frac{\partial}{\partial x^i},\frac{\partial}{\partial x^j})(\frac{\partial}{\partial x^k}).$$

Then we get the statement.

(

Moreover, considering the Patterson-Walker metric, \tilde{h}_{\pm} , we get the following:

THEOREM 4.8. Let (M, h, ∇) be a quasi-statistical manifold such that ∇ is flat. Then $(T^*M, \tilde{h}_{\pm}, \tilde{\nabla})$ is a quasi-statistical manifold.

PROOF. Let us compute $d^{\tilde{\nabla}}\tilde{h}_{\pm}$. From the definition of \tilde{h}_{\pm} and $\tilde{\nabla}$ we get immediately:

$$\begin{split} (d^{\tilde{\nabla}}\tilde{h}_{\pm})(\frac{\partial}{\partial y_{i}},\frac{\partial}{\partial y_{j}}) &= 0\\ (d^{\tilde{\nabla}}\tilde{h}_{\pm})((\frac{\partial}{\partial x^{i}})^{H},\frac{\partial}{\partial y_{j}}) &= 0\\ (d^{\tilde{\nabla}}\tilde{h}_{\pm})((\frac{\partial}{\partial x^{i}})^{H},(\frac{\partial}{\partial x^{j}})^{H})(\frac{\partial}{\partial x^{k}})^{H} &= -y_{l}R_{ijk}^{l}\\ (d^{\tilde{\nabla}}\tilde{h}_{\pm})((\frac{\partial}{\partial x^{i}})^{H},(\frac{\partial}{\partial x^{j}})^{H})(\frac{\partial}{\partial y_{k}}) &= \pm h^{kl}(d^{\nabla}h)(\frac{\partial}{\partial x^{i}},\frac{\partial}{\partial x^{j}})(\frac{\partial}{\partial x^{l}}), \end{split}$$

where the sign + is for h symmetric, - is for h skew-symmetric. Then we get the statement.

DEFINITION 4.9. A quasi-statistical manifold (M, h, ∇) such that ∇ is flat is called a *Hessian manifold*.

Therefore we get:

COROLLARY 4.10. If (M, h, ∇) is a Hessian manifold, then $(T^*M, h^{S^*}, \tilde{\nabla})$ and $(T^*M, \tilde{h}_{\pm}, \tilde{\nabla})$ are Hessian manifolds.

4.4 Quasi-statistical structures on tangent bundles

Given a non-degenerate (0, 2)-tensor field h on M, we have an isomorphism between $T(T^*M)$ and T(TM). The connection $\tilde{\nabla}$ on TM corresponding to $\tilde{\nabla}$ on T^*M , is the following:

$$\begin{split} \tilde{\tilde{\nabla}}_{(\frac{\partial}{\partial x^{i}})^{H}} (\frac{\partial}{\partial x^{j}})^{H} &= \Gamma_{ij}^{k} (\frac{\partial}{\partial x^{k}})^{H} \\ \tilde{\tilde{\nabla}}_{(\frac{\partial}{\partial x^{i}})^{H}} \frac{\partial}{\partial y^{j}} &= \Gamma_{ij}^{k} \frac{\partial}{\partial y^{k}} \\ \tilde{\tilde{\nabla}}_{\frac{\partial}{\partial y^{j}}} (\frac{\partial}{\partial x^{i}})^{H} &= 0 \\ \tilde{\tilde{\nabla}}_{\frac{\partial}{\partial y^{i}}} \frac{\partial}{\partial y^{j}} &= 0. \end{split}$$

In local coordinates, the torsion $T^{\tilde{\tilde{\nabla}}}$ of $\tilde{\tilde{\nabla}}$ is:

$$T^{\tilde{\nabla}}((\frac{\partial}{\partial x^{i}})^{H}, (\frac{\partial}{\partial x^{j}})^{H}) = (\Gamma_{ij}^{k} - \Gamma_{ji}^{k})(\frac{\partial}{\partial x^{k}})^{H} - y^{l}R_{ijl}^{k}\frac{\partial}{\partial y^{k}}$$
$$T^{\tilde{\nabla}}(\frac{\partial}{\partial y^{i}}, (\frac{\partial}{\partial x^{j}})^{H}) = 0$$
$$T^{\tilde{\nabla}}(\frac{\partial}{\partial y^{i}}, \frac{\partial}{\partial y^{j}}) = 0$$

and the curvature $R^{\tilde{\tilde{\nabla}}}$ of $\tilde{\tilde{\nabla}}$ is:

$$\begin{split} R^{\tilde{\nabla}}(\frac{\partial}{\partial y^{i}},\frac{\partial}{\partial y^{j}}) &= 0\\ R^{\tilde{\nabla}}((\frac{\partial}{\partial x^{i}})^{H},\frac{\partial}{\partial y^{j}}) &= 0\\ R^{\tilde{\nabla}}((\frac{\partial}{\partial x^{i}})^{H},(\frac{\partial}{\partial x^{j}})^{H})\frac{\partial}{\partial y^{k}} &= R^{l}_{ijk}\frac{\partial}{\partial y^{l}}\\ R^{\tilde{\nabla}}((\frac{\partial}{\partial x^{i}})^{H},(\frac{\partial}{\partial x^{j}})^{H})(\frac{\partial}{\partial x^{k}})^{H} &= R^{\nabla}(\frac{\partial}{\partial x^{i}},\frac{\partial}{\partial x^{j}})\frac{\partial}{\partial x^{k}}. \end{split}$$

Therefore we get:

PROPOSITION 4.11. ∇ is flat if and only if $\tilde{\tilde{\nabla}}$ is flat.

THEOREM 4.12. Let (M, h, ∇) be a quasi-statistical manifold such that ∇ is flat. Then $(TM, h^S, \tilde{\nabla})$ is a flat quasi-statistical manifold.

PROOF. Let us compute $d^{\tilde{\nabla}}h^S$. From the definition of h^S and $\tilde{\tilde{\nabla}}$ we get immediately:

$$(d^{\tilde{\nabla}}h^{S})(\frac{\partial}{\partial y^{i}},\frac{\partial}{\partial y^{j}}) = 0$$

$$(d^{\tilde{\nabla}}h^{S})((\frac{\partial}{\partial x^{i}})^{H},\frac{\partial}{\partial y^{j}}) = 0$$

$$(d^{\tilde{\nabla}}h^{S})((\frac{\partial}{\partial x^{i}})^{H},(\frac{\partial}{\partial x^{j}})^{H})(\frac{\partial}{\partial y^{k}}) = -y^{l}R^{r}_{ijl}h_{rk}$$

$$(d^{\tilde{\nabla}}h^{S})((\frac{\partial}{\partial x^{i}})^{H},(\frac{\partial}{\partial x^{j}})^{H})(\frac{\partial}{\partial x^{k}})^{H} = (d^{\nabla}h)(\frac{\partial}{\partial x^{i}},\frac{\partial}{\partial x^{j}})(\frac{\partial}{\partial x^{k}}).$$
e statement.

Then we get the statement.

Moreover, considering the horizontal lift metric, h^H , we get the following:

THEOREM 4.13. Let (M, h, ∇) be a quasi-statistical manifold such that ∇ is flat. Then $(TM, h^H, \tilde{\nabla})$ is a quasi-statistical manifold if and only if $\nabla h = 0$.

PROOF. Let us compute $d^{\tilde{\nabla}}h^H$. From the definition of h^H and $\tilde{\nabla}$ we get immediately:

$$\begin{split} (d^{\tilde{\nabla}}h^{H})(\frac{\partial}{\partial y^{i}},\frac{\partial}{\partial y^{j}}) &= 0\\ (d^{\tilde{\nabla}}h^{H})((\frac{\partial}{\partial x^{i}})^{H},\frac{\partial}{\partial y^{j}}) &= \pm (\nabla_{\underline{\partial}}h)\frac{\partial}{\partial x^{j}}\\ (d^{\tilde{\nabla}}h^{H})((\frac{\partial}{\partial x^{i}})^{H},(\frac{\partial}{\partial x^{j}})^{H})(\frac{\partial}{\partial x^{k}})^{H} &= -y^{l}R^{s}_{ijl}h_{sk}\\ (d^{\tilde{\nabla}}h^{H})((\frac{\partial}{\partial x^{i}})^{H},(\frac{\partial}{\partial x^{j}})^{H})(\frac{\partial}{\partial y^{k}}) &= (d^{\nabla}h)(\frac{\partial}{\partial x^{i}},\frac{\partial}{\partial x^{j}})(\frac{\partial}{\partial x^{k}}), \end{split}$$

where the sign + is for h symmetric, - is for h skew-symmetric. Then we get the statement.

Therefore we get:

COROLLARY 4.14. If (M, h, ∇) is a Hessian manifold, then $(TM, h^S, \tilde{\nabla})$ is a Hessian manifold. Moreover, if $\nabla h = 0$, then $(TM, h^H, \tilde{\nabla})$ is a Hessian manifold.

5 Norden and Para-Norden structures on cotangent and tangent bundles

Norden manifolds, also called almost complex manifolds with B-metric, were introduced in [8]. They have applications in mathematics and in theoretical physics.

DEFINITION 5.1. A Norden manifold, (M, J, h), is an almost complex manifold (M, J) with a pseudo-Riemannian metric, h (called Norden metric), such that J is h-symmetric.

Moreover, if J is integrable, then (M, J, h) is called *complex Norden manifold*.

DEFINITION 5.2. An almost Para-complex Norden manifold (or simply, almost Para-Norden manifold), (M, J, h), is a real even dimensional smooth manifold M with a pseudo-Riemannian metric, h, and a (1, 1)-tensor field, J, such that $J^2 = I$, the two eigenbundles T^+M , T^-M , associated to the two eigenvalues +1, -1, of J respectively have the same rank and J is h-symmetric.

Moreover, if J is integrable, then (M, J, h) is called *Para-Norden manifold*.

5.1 Norden and Para-Norden structures on cotangent bundles

Let (M, h) be a pseudo-Riemannian manifold and let \hat{J}_c , \hat{J}_p be the generalized complex structure and the generalized product structure defined by h in (1) and (2) respectively. Again we will denote \hat{J}_{\mp} for $\hat{J}_c =: \hat{J}_-$ and $\hat{J}_p =: \hat{J}_+$.

Let ∇ be an affine connection on M and let $\Phi^{\nabla} : TM \oplus T^*M \to T(T^*M)$ be the bundle morphism defined by (8). We define:

$$\tilde{J}^{\nabla}_{\mp} =: \Phi^{\nabla} \circ \hat{J}_{\mp} \circ (\Phi^{\nabla})^{-1}.$$

We have immediately that $(\tilde{J}^{\nabla}_{\pm})^2 = \pm I$.

PROPOSITION 5.3. Let \tilde{h} be the Patterson-Walker metric on T^*M . Then $(T^*M, \tilde{J}_{-}^{\nabla}, \tilde{h})$ is a Norden manifold and $(T^*M, \tilde{J}_{+}^{\nabla}, \tilde{h})$ is an almost Para-Norden manifold. Moreover, if (M, h, ∇) is a flat quasi-statistical manifold, then $(T^*M, \tilde{J}_{-}^{\nabla}, \tilde{h})$ is a complex Norden manifold and $(T^*M, \tilde{J}_{+}^{\nabla}, \tilde{h})$ is a Para-Norden manifold.

PROOF. In local coordinates, we get the following:

$$\tilde{J}^{\nabla}_{\mp}(X^H_i) =: h_{ik} \frac{\partial}{\partial y_k}$$

Generalized quasi-statistical structures

$$\tilde{J}^{\nabla}_{\mp}(\frac{\partial}{\partial y_j}) =: \mp h^{jk} X^H_k.$$

In particular, we have:

$$\begin{split} \tilde{h}(\tilde{J}^{\nabla}_{\mp}(X^{H}_{i}), X^{H}_{j}) &= h_{ij} \\ \tilde{h}(\tilde{J}^{\nabla}_{\mp}(\frac{\partial}{\partial y_{i}}), \frac{\partial}{\partial y_{j}}) &= \mp h^{ij} \\ \tilde{h}(\tilde{J}^{\nabla}_{\mp}(X^{H}_{i}), \frac{\partial}{\partial y_{j}}) &= 0 \\ \tilde{h}(\tilde{J}^{\nabla}_{\mp}(\frac{\partial}{\partial y_{i}}), X^{H}_{j}) &= 0, \end{split}$$

therefore, from the symmetry of h, we get the first statement.

Moreover, if we compute the Nijenhuis tensor field of \tilde{J}^{∇}_{\mp} , we have:

$$N_{\tilde{J}^{\nabla}_{\mp}}(X_{i}^{H}, X_{j}^{H}) = \pm (h^{kl}(d^{\nabla}h)(\frac{\partial}{\partial x^{i}}, \frac{\partial}{\partial x^{j}})(\frac{\partial}{\partial x^{k}}) - y_{k}R_{ijl}^{k})\frac{\partial}{\partial y_{l}}$$

$$N_{\tilde{J}^{\nabla}_{\mp}}(X_{i}^{H}, \frac{\partial}{\partial y_{j}}) = h^{jl}(h^{sr}y_{k}R_{ils}^{k}X_{r}^{H} \mp (d^{\nabla}h)(\frac{\partial}{\partial x^{i}}, \frac{\partial}{\partial x^{l}})(\frac{\partial}{\partial x^{r}})\frac{\partial}{\partial y_{r}})$$

$$N_{\tilde{J}^{\nabla}_{\mp}}(\frac{\partial}{\partial y_{i}}, \frac{\partial}{\partial y_{j}}) = h^{ik}h^{jl}(h^{ps}(d^{\nabla}h)(\frac{\partial}{\partial x^{l}}, \frac{\partial}{\partial x^{k}})(\frac{\partial}{\partial x^{p}})X_{s}^{H} + y_{s}R_{klr}^{s}\frac{\partial}{\partial y^{r}}).$$
proof is complete.

Then the proof is complete.

REMARK 5.4. If h is a non-degenerate skew-symmetric (0, 2)-tensor field on M, then the same construction gives rise to a Hermitian, respectively Para-Hermitian, structure on T^*M .

Norden and Para-Norden structures on tangent bundles 5.2

Let (M,h) be a pseudo-Riemannian manifold and let \hat{J}_c , \hat{J}_p be the generalized complex structure and the generalized product structure defined by h in (1) and (2) respectively. Again we will denote \hat{J}_{\mp} for $\hat{J}_c =: \hat{J}_-$ and $\hat{J}_p =: \hat{J}_+$.

Let ∇ be an affine connection on M and let $\Psi^{\nabla}: TM \oplus T^*M \to T(TM)$ be the bundle morphism defined by (9). We define:

$$\bar{J}^{\nabla}_{\mp} =: \Psi^{\nabla} \circ \hat{J}_{\mp} \circ (\Psi^{\nabla})^{-1}.$$

Let $X \in C^{\infty}(TM)$ and let X^{H} , X^{V} be respectively the horizontal and vertical lift of X. We have immediately that

$$\bar{J}^{\nabla}_{\mp}(X^H) = X^V$$
$$\bar{J}^{\nabla}_{\mp}(X^V) = \mp X^H.$$

A direct computation gives the following:

PROPOSITION 5.5. Let h^H be the horizontal lift metric of h on TM. Then $(TM, \overline{J}^{\nabla}_{-}, h^H)$ is a Norden manifold and $(TM, \overline{J}^{\nabla}_{+}, h^H)$ is an almost Para-Norden manifold.

REMARK 5.6. The almost complex structure $\overline{J}_{-}^{\nabla}$ is the canonical almost complex structure of TM defined in [3]. In particular, it is integrable if and only if ∇ is flat and torsion-free.

References

- S. Amari, Differential-Geometrical Methods in Statistics, Lecture Notes in Statistics, 28 Springer (1985).
- [2] A. M. Blaga, A. Nannicini, Generalized metallic structures, arXiv:1807.08308, 2018.
- [3] P. Dombrowski, On the geometry of tangent bundles, J. reine angew. Math. 210 (1962), 73-88.
- [4] H. Matsuzoe, Quasi-statistical manifolds and geometry of affine distributions, Pure and Applied Differential Geometry 2012: In Memory of Franki Dillen, Berichte aus der Mathematik, ed. Joeri Van der Veken, Ignace Van de Woestyne, Leopold Verstraelen, Luc Vrancken, Shaker Verlag, 2013.
- [5] A. Nannicini, Almost complex structures on cotangent bundles and generalized geometry, J. Geom. Phys. 60 (2010), 1781-1791.
- [6] A. Nannicini, Calibrated complex structures on the generalized tangent bundle of a Riemannian manifold, J. Geom. Phys. 56 (2006), 903-916.
- [7] M. Nogushi, Geometry of statistical manifolds, Differ. Geom. Appl. 2 (1992), 197-222.
- [8] A. P. Norden, On a class of four-dimensional A-spaces, Russian Math. (Izv VUZ) 17 (4) (1960), 145-157.

[9] E. M. Patterson, A. G. Walker, *Riemann extension*, Quart. J. Math. Oxford 3 (1952), 19-28.

Adara M. Blaga Department of Mathematics West University of Timişoara Bld. V. Pârvan nr. 4, 300223, Timişoara, România adarablaga@yahoo.com

Antonella Nannicini Department of Mathematics and Informatics "U. Dini" University of Florence Viale Morgagni, 67/a, 50134, Firenze, Italy antonella.nannicini@unifi.it