

# Extensions of modules over a class of Lie conformal algebras $\mathscr{W}(b)$

Kaijing Ling, Lamei Yuan \*

Department of Mathematics, Harbin Institute of Technology, Harbin 150001, China

E-mails: kjling\_edu@126.com, lmyuan@hit.edu.cn

**Abstract:** Let  $\mathscr{W}(b)$  be a class of free Lie conformal algebras of rank 2 with  $\mathbb{C}[\partial]$ -basis  $\{L, H\}$  and relations

$$[L_\lambda L] = (\partial + 2\lambda)L, \quad [L_\lambda H] = (\partial + (1 - b)\lambda)H, \quad [H_\lambda H] = 0,$$

where  $b$  is a nonzero complex number. In this paper, we classify extensions between two finite irreducible conformal modules over the Lie conformal algebras  $\mathscr{W}(b)$ .

**Key words:** Lie conformal algebra, conformal modules, extensions

**MR(2000) Subject Classification:** 17B65, 17B68

## 1 Introduction

Conformal modules of Lie conformal algebras were introduced in [4]. Cheng and Kac studied finite conformal modules over the Virasoro, the Neveu-Schwarz and the current conformal algebras in [1]. In general, conformal modules are not completely reducible. Thus it is necessary to study the extension problem. In [2, 3], Cheng, Kac and Wakimoto classified extensions between finite conformal modules over the Virasoro, the current, the Neveu-Schwarz and the semi-direct sum of the Virasoro and the current conformal algebras. In [5], Ngau Lam solved the same problem for the supercurrent conformal algebras by using the techniques developed in [2].

In this paper, we aim to study extensions between conformal modules over a class of Lie conformal algebras  $\mathscr{W}(b)$ , which was introduced in [7] as the Lie conformal algebras associated with the Lie algebras  $W(a, b)$  with  $a, b \in \mathbb{C}$ . By definition, the Lie conformal algebra  $\mathscr{W}(b) = \mathbb{C}[\partial]L \oplus \mathbb{C}[\partial]H$  with the  $\lambda$ -bracket

$$[L_\lambda L] = (\partial + 2\lambda)L, \quad [L_\lambda H] = (\partial + (1 - b)\lambda)H, \quad [H_\lambda H] = 0, \quad (1.1)$$

where  $b \in \mathbb{C}$ . Finite irreducible conformal modules over  $\mathscr{W}(b)$  were classified in [8]. It turns out that any finite irreducible conformal  $\mathscr{W}(b)$ -module is free of rank one and of the form  $M(\alpha, \beta, \Delta) = \mathbb{C}[\partial]v_\Delta$  with  $(\Delta, \beta) \neq (0, 0)$  if  $b = 0$  or  $M(\alpha, \Delta) = \mathbb{C}[\partial]v_\Delta$  with  $\Delta \neq 0$  if  $b \neq 0$  (see Proposition 2.5).

For each  $b$ , the Lie conformal algebra  $\mathscr{W}(b)$  is a semidirect product of the Virasoro conformal algebra  $\text{Vir} = \mathbb{C}[\partial]L$  with  $[L_\lambda L] = (\partial + 2\lambda)L$  and the conformal Vir-module  $\mathbb{C}[\partial]H$  on which the action of  $L$  is given by  $L_\lambda H = (\partial + (1 - b)\lambda)H$ . Two special cases of  $\mathscr{W}(b)$  should be pointed out. One is  $\mathscr{W}(0)$ , which is called Heisenberg-Virasoro conformal algebra in [7]. Its cohomology was studied in [9]. The other one is  $\mathscr{W}(-1)$ , which is exactly the  $W(2, 2)$  Lie conformal algebra. Its conformal derivations and central extensions were studied in [10].

In this paper, we will consider extensions of conformal  $\mathscr{W}(b)$ -modules with  $b \neq 0$ , since this problem for  $\mathscr{W}(0)$ -modules have been solved in [6]. The rest paper is organized as follows. In Section 2, we recall

---

\*Corresponding author: lmyuan@hit.edu.cn (Lamei Yuan)

the notions of conformal modules and their extensions, and some known results. In Section 3, we consider extensions of conformal  $\mathscr{W}(b)$ -modules of the following three types:

$$0 \longrightarrow \mathbb{C}c_\gamma \longrightarrow E \longrightarrow M(\alpha, \Delta) \longrightarrow 0, \quad (1.2)$$

$$0 \longrightarrow M(\alpha, \Delta) \longrightarrow E \longrightarrow \mathbb{C}c_\gamma \longrightarrow 0, \quad (1.3)$$

$$0 \longrightarrow M(\bar{\alpha}, \bar{\Delta}) \longrightarrow E \longrightarrow M(\alpha, \Delta) \longrightarrow 0. \quad (1.4)$$

where  $M(\alpha, \Delta)$  and  $M(\bar{\alpha}, \bar{\Delta})$  are irreducible  $\mathscr{W}(b)$ -modules, and  $\mathbb{C}c_\gamma$  is the 1-dimensional  $\mathscr{W}(b)$ -module with  $L_\lambda c_\gamma = H_\lambda c_\gamma = 0$  and  $\partial c_\gamma = \gamma c_\gamma$ . During the whole process, we use some results of the Virasoro conformal algebra from [2]. The main results of our paper are summarized in Theorems 3.2, 3.4 and 3.7.

## 2 Preliminaries

In this section, we recall some definitions, notations, and related results for later use. For more details, the reader is referred to [1, 2, 8].

**Definition 2.1.** A conformal module  $M$  over a Lie conformal algebra  $R$  is a  $\mathbb{C}[\partial]$ -module endowed with a  $\mathbb{C}$ -linear map  $R \rightarrow \text{End}_{\mathbb{C}} M \otimes_{\mathbb{C}} \mathbb{C}[\lambda]$ ,  $a \mapsto a_\lambda^M$ , satisfying the following conditions for all  $a, b \in R$ :

$$\begin{aligned} [a_\lambda^M, b_\mu^M] &= a_\lambda^M b_\mu^M - b_\mu^M a_\lambda^M = [a_\lambda b]_{\lambda+\mu}^M, \\ (\partial a)_\lambda^M &= [\partial, a_\lambda^M] = -\lambda a_\lambda^M. \end{aligned}$$

A module  $M$  over a conformal algebra  $R$  is called *finite*, if  $M$  is finitely generated over  $\mathbb{C}[\partial]$ . A module  $M$  over a conformal algebra  $R$  is called *irreducible* if there is no nontrivial invariant subspace.

The vector space  $\mathbb{C}$  is viewed as a trivial module with trivial actions of  $\partial$  and  $R$ . For a fixed nonzero complex constant  $\alpha$ , there is a natural  $\mathbb{C}[\partial]$ -module structure on  $\mathbb{C}c_\alpha$ , such that  $\mathbb{C}c_\alpha = \mathbb{C}$  and  $\partial c_\alpha = \alpha c_\alpha$ . Then  $\mathbb{C}c_\alpha$  becomes an  $R$ -module with trivial action of  $R$ .

Let  $V$  and  $W$  be two modules over a Lie conformal algebra (or a Lie algebra)  $R$ . An *extension* of  $W$  by  $V$  is an exact sequence of  $R$ -modules of the form

$$0 \longrightarrow V \xrightarrow{i} E \xrightarrow{p} W \longrightarrow 0. \quad (2.1)$$

Two extensions  $0 \longrightarrow V \xrightarrow{i} E \xrightarrow{p} W \longrightarrow 0$  and  $0 \longrightarrow V \xrightarrow{i'} E' \xrightarrow{p'} W \longrightarrow 0$  are said to be *equivalent* if there exists a commutative diagram of the form

$$\begin{array}{ccccccccc} 0 & \longrightarrow & V & \xrightarrow{i} & E & \xrightarrow{p} & W & \longrightarrow & 0 \\ & & 1_V \downarrow & & \psi \downarrow & & 1_W \downarrow & & \\ 0 & \longrightarrow & V & \xrightarrow{i'} & E' & \xrightarrow{p'} & W & \longrightarrow & 0, \end{array}$$

where  $1_V : V \rightarrow V$  and  $1_W : W \rightarrow W$  are the respective identity maps and  $\psi : E \rightarrow E'$  is a homomorphism of modules.

The direct sum of modules  $V \oplus W$  obviously gives rise to an extension. Extensions equivalent to it are called *trivial extensions*. In general, an extension can be thought of as the direct sum of vector spaces  $E = V \oplus W$ , where  $V$  is a submodule of  $E$ , while for  $w$  in  $W$  we have:

$$a \cdot w = aw + \phi_a(w), \quad a \in R,$$

where  $\phi_a : W \rightarrow V$  is a linear map satisfying the cocycle condition:

$$\phi_{[a,b]}(w) = \phi_a(bw) + a\phi_b(w) - \phi_b(aw) - b\phi_a(w), \quad b \in R.$$

The set of these cocycles form a vector space over  $\mathbb{C}$ . Cocycles equivalent to the trivial extension are called *coboundaries*. They form a subspace and the quotient space by it is denoted by  $\text{Ext}(W, V)$ .

For the Virasoro conformal algebra  $\text{Vir}$ , it was shown in [1] that all free nontrivial  $\text{Vir}$ -modules of rank one over  $\mathbb{C}[\partial]$  are the following ones ( $\Delta, \alpha \in \mathbb{C}$ ):

$$M(\alpha, \Delta) = \mathbb{C}[\partial]v_\Delta, \quad L_\lambda v = (\partial + \alpha + \Delta\lambda)v_\Delta. \quad (2.2)$$

The module  $M(\alpha, \Delta)$  is irreducible if and only if  $\Delta \neq 0$ . The module  $M(\alpha, 0)$  contains a unique nontrivial submodule  $(\partial + \alpha)M(\alpha, 0)$  isomorphic to  $M(\alpha, 1)$ . Moreover, the modules  $M(\alpha, \Delta)$  with  $\Delta \neq 0$  exhaust all finite irreducible non-1-dimensional  $\text{Vir}$ -modules.

In [2], extensions over the Virasoro conformal modules of the following types have been classified:

$$0 \longrightarrow \mathbb{C}c_\gamma \longrightarrow E \longrightarrow M(\alpha, \Delta) \longrightarrow 0, \quad (2.3)$$

$$0 \longrightarrow M(\alpha, \Delta) \longrightarrow E \longrightarrow \mathbb{C}c_\gamma \longrightarrow 0, \quad (2.4)$$

$$0 \longrightarrow M(\bar{\alpha}, \bar{\Delta}) \longrightarrow E \longrightarrow M(\alpha, \Delta) \longrightarrow 0. \quad (2.5)$$

The following are the corresponding results.

**Theorem 2.2.** (cf. Ref. [2]) *Nontrivial extensions of the form (2.3) exist if and only if  $\alpha + \gamma = 0$  and  $\Delta = 1$  or  $2$ . In these cases, they are given (up to equivalence) by*

$$L_\lambda v_\Delta = (\partial + \alpha + \Delta\lambda)v_\Delta + f(\lambda)c_\gamma,$$

where

$$(i) \quad f(\lambda) = c_2\lambda^2, \text{ for } \Delta = 1 \text{ and } c_2 \neq 0.$$

$$(ii) \quad f(\lambda) = c_3\lambda^3, \text{ for } \Delta = 2 \text{ and } c_3 \neq 0.$$

Furthermore, all trivial cocycles are given by scalar multiples of the polynomial  $f(\lambda) = \alpha + \gamma + \Delta\lambda$ .

**Theorem 2.3.** (cf. Ref. [2]) *There are nontrivial extensions of Virasoro conformal modules of the form (2.4) if and only if  $\alpha + \gamma = 0$  and  $\Delta = 1$ . These extensions are given (up to equivalence) by*

$$L_\lambda c_\gamma = f(\partial, \lambda)v_\Delta, \quad \partial c_\gamma = \gamma c_\gamma + h(\partial)v_\Delta,$$

where  $f(\partial, \lambda) = h(\partial) = a_0$ , and  $a_0 \neq 0$ .

**Theorem 2.4.** (cf. Ref. [2]) Nontrivial extensions of the form (2.5) of Virasoro conformal modules exist only if  $\alpha = \bar{\alpha}$  and  $\Delta - \bar{\Delta} = 0, 1, 2, 3, 4, 5, 6$ . In these cases, they are given (up to equivalence) by

$$L_\lambda v_\Delta = (\partial + \alpha + \Delta\lambda)v_\Delta + f(\partial, \lambda)v_{\bar{\Delta}}.$$

And the following is a complete list of values of  $\Delta$  and  $\bar{\Delta}$  along with the corresponding polynomials  $f(\partial, \lambda)$ , whose nonzero scalar multiples give rise to nontrivial extensions (by replacing  $\partial$  by  $\partial + \alpha$ ):

- (i)  $\Delta = \bar{\Delta}, f_1(\partial, \lambda) = c_0 + c_1\lambda, (c_0, c_1) \neq (0, 0)$ .
- (ii)  $\Delta = 1, \bar{\Delta} = 0, f_2(\partial, \lambda) = c_0\partial + c_1\partial\lambda + c_2\lambda^2$ , where  $(c_0, c_1, c_2) \neq (0, 0, 0)$ .
- (iii)  $\Delta - \bar{\Delta} = 2, f_3(\partial, \lambda) = \lambda^2(2\partial + \lambda)$ .
- (iv)  $\Delta - \bar{\Delta} = 3, f_4(\partial, \lambda) = \partial\lambda^2(\partial + \lambda)$ .
- (v)  $\Delta - \bar{\Delta} = 4, f_5(\partial, \lambda) = \lambda^2(4\partial^3 + 6\partial^2\lambda - \partial\lambda^2 + \bar{\Delta}\lambda^3)$ .
- (vi)  $\Delta = 5, \bar{\Delta} = 0, f_6(\partial, \lambda) = 5\partial^4\lambda^2 + 10\partial^2\lambda^4 - \partial\lambda^5$ .
- (vi')  $\Delta = 1, \bar{\Delta} = -4, f_{6'}(\partial, \lambda) = \partial^4\lambda^2 - 10\partial^2\lambda^4 - 17\partial\lambda^5 - 8\lambda^6$ .
- (vii)  $\Delta = \frac{7}{2} \pm \frac{\sqrt{19}}{2}, \Delta - \bar{\Delta} = 6, f_7(\partial, \lambda) = \partial^4\lambda^3 - (2\bar{\Delta} + 3)\partial^3\lambda^4 - 3\bar{\Delta}\partial^2\lambda^5 - (3\bar{\Delta} + 1)\partial\lambda^6 - (\bar{\Delta} + \frac{9}{28})\lambda^7$ .

The following result, due to [8], gives all finite free irreducible conformal modules over the Lie conformal algebras  $\mathscr{W}(b)$ .

**Proposition 2.5.** (cf. Ref. [8]) Any finite free irreducible conformal module over the Lie conformal algebra  $\mathscr{W}(b)$  is of the form

$$M = \mathbb{C}[\partial]v_\Delta, L_\lambda v_\Delta = (\partial + \alpha + \Delta\lambda)v_\Delta, H_\lambda v_\Delta = \delta_{b,0}v_\Delta,$$

with  $\Delta, \alpha, \beta \in \mathbb{C}$ , and  $(\Delta, \beta) \neq (0, 0)$ .

In this paper, we denote the  $\mathscr{W}(b)$ -module  $M$  from Proposition 2.5 by  $M(\alpha, \beta, \Delta)$  if  $b = 0$ , and  $M(\alpha, \Delta)$  if  $b \neq 0$ , respectively.

### 3 Extensions of conformal $\mathscr{W}(b)$ -modules

In this section, we always assume that  $b \neq 0$ . In this case, it follows from Proposition 2.5 that any finite free irreducible conformal module over the Lie conformal algebra  $\mathscr{W}(b)$  is of the form

$$M(\alpha, \Delta) = \mathbb{C}[\partial]v_\Delta, L_\lambda v_\Delta = (\partial + \alpha + \Delta\lambda)v_\Delta, H_\lambda v_\Delta = 0,$$

where  $\Delta, \alpha \in \mathbb{C}$ , and  $\Delta \neq 0$ .

Let  $M$  be a  $\mathbb{C}[\partial]$ -module. By Definition 2.1, a  $\mathscr{W}(b)$ -module structure on  $M$  is given by  $L_\lambda^M, H_\mu^M \in \text{End}_{\mathbb{C}}(M)[\lambda]$  such that

$$[L_\lambda^M, L_\mu^M] = (\lambda - \mu)L_{\lambda+\mu}^M, \quad (3.1)$$

$$[L_\lambda^M, H_\mu^M] = -(b\lambda + \mu)H_{\lambda+\mu}^M, \quad (3.2)$$

$$[H_\lambda^M, L_\mu^M] = (\lambda + b\mu)H_{\lambda+\mu}^M, \quad (3.3)$$

$$[H_\lambda^M, H_\mu^M] = 0, \quad (3.4)$$

$$[\partial, L_\lambda^M] = -\lambda L_\lambda^M, \quad (3.5)$$

$$[\partial, H_\lambda^M] = -\lambda H_\lambda^M. \quad (3.6)$$

First, we consider extensions of  $\mathscr{W}(b)$ -modules of the form

$$0 \longrightarrow \mathbb{C}c_\gamma \longrightarrow E \longrightarrow M(\alpha, \Delta) \longrightarrow 0. \quad (3.7)$$

As a module over  $\mathbb{C}[\partial]$ ,  $E$  in (3.7) is isomorphic to  $\mathbb{C}c_\gamma \oplus M(\alpha, \Delta)$ , where  $\mathbb{C}c_\gamma$  is a  $\mathscr{W}(b)$ -submodule, and  $M(\alpha, \Delta) = \mathbb{C}[\partial]v_\Delta$  such that the following identities hold in  $E$ :

$$L_\lambda v_\Delta = (\partial + \alpha + \Delta\lambda)v_\Delta + f(\lambda)c_\gamma, \quad H_\lambda v_\Delta = g(\lambda)c_\gamma, \quad (3.8)$$

where  $f(\lambda) = \sum_{n \geq 0} f_n \lambda^n$ ,  $g(\lambda) = \sum_{n \geq 0} g_n \lambda^n$ , with  $f_n, g_n \in \mathbb{C}$ .

**Lemma 3.1.** *All trivial extensions of the form (3.7) are given by (3.8), where  $f(\lambda)$  is a scalar multiple of  $\alpha + \gamma + \Delta\lambda$ , and  $g(\lambda) = 0$ .*

*Proof.* Suppose that (3.7) represents a trivial cocycle. This means that the exact sequence (3.7) is split and hence there exists  $v'_\Delta = \varphi(\partial)v_\Delta + ac_\gamma \in E$ , where  $a \in \mathbb{C}$ , such that

$$L_\lambda v'_\Delta = (\partial + \alpha + \Delta\lambda)v'_\Delta = (\partial + \alpha + \Delta\lambda)\varphi(\partial)v_\Delta + a(\gamma + \alpha + \Delta\lambda)c_\gamma,$$

and  $H_\lambda v'_\Delta = 0$ . On the other hand, it follows from (3.8) that

$$L_\lambda v'_\Delta = \varphi(\partial + \lambda)(\partial + \alpha + \Delta\lambda)v_\Delta + \varphi(\partial + \lambda)f(\lambda)c_\gamma,$$

$$H_\lambda v'_\Delta = \varphi(\partial + \lambda)g(\lambda)c_\gamma.$$

Comparing both expressions for  $L_\lambda v'_\Delta$  and  $H_\lambda v'_\Delta$  respectively, we see that  $\varphi(\partial)$  is a constant. Therefore  $f(\lambda)$  is a scalar multiple of  $\alpha + \gamma + \Delta\lambda$ , and  $g(\lambda) = 0$ .  $\square$

**Theorem 3.2.** *There are nontrivial extensions of  $\mathscr{W}(b)$ -modules of the form (3.7) if and only if  $\alpha + \gamma = 0$ . Moreover, these nontrivial extensions are given (up to equivalence) by (3.8), where, if  $g(\lambda) = 0$ , then  $\Delta = 1, 2$  and  $f(\lambda)$  is from the nonzero polynomials of Theorem 2.2 for all  $0 \neq b \in \mathbb{C}$ , or else  $g(\lambda) = a$  for some  $0 \neq a \in \mathbb{C}$ ,  $\Delta = b$  and*

$$f(\lambda) = \begin{cases} c_2 \lambda^2, & \text{for } b = 1, \\ c_3 \lambda^3, & \text{for } b = 2, \\ 0, & \text{otherwise,} \end{cases} \quad (3.9)$$

with  $c_2, c_3 \in \mathbb{C}$ .

*Proof.* Applying both sides of (3.1) and (3.2) to  $v_\Delta$ , we obtain

$$(\alpha + \gamma + \lambda + \Delta\mu)f(\lambda) - (\alpha + \gamma + \mu + \Delta\lambda)f(\mu) = (\lambda - \mu)f(\lambda + \mu), \quad (3.10)$$

$$(\alpha + \gamma + \mu + \Delta\lambda)g(\mu) = (b\lambda + \mu)g(\lambda + \mu). \quad (3.11)$$

Setting  $\lambda = 0$  in (3.11) gives

$$(\alpha + \gamma)g(\mu) = 0. \quad (3.12)$$

**Case 1.**  $\alpha + \gamma \neq 0$ .

By (3.12),  $g(\mu) = 0$ . Setting  $\mu = 0$  in (3.10), we see that  $f(\lambda)$  is a scalar multiple of  $\alpha + \gamma + \Delta\lambda$ . Thus, the extension is trivial by Lemma 3.1.

**Case 2.**  $\alpha + \gamma = 0$ .

Putting  $\mu = 0$  in (3.11), we obtain  $g(\lambda) = \frac{\Delta}{b}g(0)$ , which gives  $g(\lambda)$  is a constant. If  $g(\lambda) = a \neq 0$ , then  $\Delta = b$ . By Theorem 2.2,  $f(\lambda) = c_2\lambda^2$  if  $b = 1$ ,  $f(\lambda) = c_3\lambda^3$  if  $b = 2$ , and  $f(\lambda) = 0$  if  $b \neq 1$  or  $2$ . In these cases, the corresponding extensions are nontrivial. If  $g(\lambda) = 0$ , then it reduces to the Virasoro case. By Theorem 2.2, we obtain the result.  $\square$

Next we consider extensions of  $\mathscr{W}(b)$ -modules of the form

$$0 \longrightarrow M(\alpha, \Delta) \longrightarrow E \longrightarrow \mathbb{C}c_\gamma \longrightarrow 0. \quad (3.13)$$

As a vector space,  $E$  in (3.13) is isomorphic to  $M(\alpha, \Delta) \oplus \mathbb{C}c_\gamma$ . Here  $M(\alpha, \Delta) = \mathbb{C}[\partial]v_\Delta$  is a  $\mathscr{W}(b)$ -submodule and we have

$$L_\lambda c_\gamma = f(\partial, \lambda)v_\Delta, \quad H_\lambda c_\gamma = g(\partial, \lambda)v_\Delta, \quad \partial c_\gamma = \gamma c_\gamma + h(\partial)v_\Delta, \quad (3.14)$$

where  $f(\partial, \lambda) = \sum_{n \geq 0} f_n(\partial)\lambda^n$ ,  $g(\partial, \lambda) = \sum_{n \geq 0} g_n(\partial)\lambda^n$ ,  $f_n(\partial), g_n(\partial), h(\partial) \in \mathbb{C}[\partial]$ .

**Lemma 3.3.** *All trivial extensions of the form (3.13) are given by (3.14) with  $f(\partial, \lambda) = (\partial + \alpha + \Delta\lambda)\phi(\partial + \lambda)$ ,  $g(\partial, \lambda) = 0$  and  $h(\partial) = (\partial - \gamma)\phi(\partial)$ , where  $\phi$  is a polynomial.*

*Proof.* Suppose that (3.13) represents a trivial cocycle. This means that the exact sequence (3.13) is split and hence there exists  $c'_\gamma = ac_\gamma + \phi(\partial)v_\Delta \in E$ , where  $a \in \mathbb{C}$ , such that  $L_\lambda c'_\gamma = H_\lambda c'_\gamma = 0$  and  $\partial c'_\gamma = \gamma c'_\gamma$ . On the other hand, a short computation shows that

$$\begin{aligned} L_\lambda c'_\gamma &= (\partial + \alpha + \Delta\lambda)\phi(\partial + \lambda)v_\Delta + af(\partial, \lambda)v_\Delta, \\ H_\lambda c'_\gamma &= ag(\partial, \lambda)v_\Delta, \\ \partial c'_\gamma &= a\gamma c_\gamma + (ah(\partial) + \partial\phi(\partial))v_\Delta. \end{aligned}$$

Comparing both expressions for  $L_\lambda c'_\gamma$ ,  $H_\lambda c'_\gamma$  and  $\partial c'_\gamma$  respectively, we obtain the result.  $\square$

**Theorem 3.4.** *There are nontrivial extensions of  $\mathscr{W}(b)$ -modules of the form (3.13) if and only if  $\alpha + \gamma = 0$  and  $\Delta = 1$ . In this case,  $\dim_{\mathbb{C}}(\mathbb{C}c_{-\alpha}, \text{Ext}(M(\alpha, 1))) = 1$ , and the unique (up to a scalar) nontrivial extension is given by*

$$L_\lambda c_\gamma = av_\Delta, \quad H_\lambda c_\gamma = 0, \quad \partial c_\gamma = \gamma c_\gamma + av_\Delta,$$

where  $a$  is a nonzero complex number.

*Proof.* Applying both sides of (3.1), (3.5) and (3.6) to  $c_\gamma$  gives the following functional equations:

$$(\partial + \alpha + \Delta\lambda)f(\partial + \lambda, \mu) - (\partial + \alpha + \Delta\mu)f(\alpha + \mu, \lambda) = (\lambda - \mu)f(\partial, \lambda + \mu), \quad (3.15)$$

$$(\partial + \lambda - \gamma)f(\partial, \lambda) = (\partial + \alpha + \Delta\lambda)h(\partial + \lambda), \quad (3.16)$$

$$(\partial + \lambda - \gamma)g(\partial, \lambda) = 0. \quad (3.17)$$

Obviously,  $g(\partial, \lambda) = 0$  by (3.17). Replacing  $\partial$  by  $\bar{\partial} = \partial + \alpha$  and letting  $\bar{f}(\bar{\partial}, \lambda) = f(\bar{\partial} - \alpha, \lambda)$ , and  $\bar{h}(\bar{\partial}) = h(\bar{\partial} - \alpha)$ , we can rewrite (3.15) and (3.16) in a homogeneous form

$$(\bar{\partial} + \Delta\lambda)\bar{f}(\bar{\partial} + \lambda, \mu) - (\bar{\partial} + \Delta\mu)\bar{f}(\bar{\partial} + \mu, \lambda) = (\lambda - \mu)\bar{f}(\bar{\partial}, \lambda + \mu), \quad (3.18)$$

$$(\bar{\partial} - \alpha + \lambda - \gamma)\bar{f}(\bar{\partial}, \lambda) = (\bar{\partial} + \Delta\lambda)\bar{h}(\bar{\partial} + \lambda). \quad (3.19)$$

Taking  $\mu = 0$  in (3.18), we can obtain that, if degree of  $\bar{f}$  is positive,  $\bar{f}(\bar{\partial}, \lambda)$  is a scalar multiple of  $(\bar{\partial} + \Delta\lambda)\bar{f}(\bar{\partial} + \lambda)$ , where  $\bar{f}(\bar{\partial} + \lambda)$  is a polynomial in  $\bar{\partial} + \lambda$ , or else  $\bar{f}(\bar{\partial}, \lambda) = a \in \mathbb{C}$ .

Assume that  $\bar{f}(\bar{\partial}, \lambda)$  is a scalar multiple of  $(\bar{\partial} + \Delta\lambda)\bar{f}(\bar{\partial} + \lambda)$ . Letting  $\lambda = 0$  in (3.19), we have  $\bar{h}(\bar{\partial})$  is a scalar multiple of  $(\bar{\partial} - \alpha - \gamma)\bar{f}(\bar{\partial})$ . However, they correspond to the trivial extension by Lemma 3.3.

It is left to consider the case  $\bar{f}(\bar{\partial}, \lambda) = a$ . Substituting this into (3.19) gives  $\bar{h}(\bar{\partial}) = \bar{f}(\bar{\partial}, \lambda) = a$ , and in particular,  $\alpha + \gamma = 0$ ,  $\Delta = 1$  if  $a \neq 0$ . The proof is finished.  $\square$

Finally, we study extensions of conformal  $\mathscr{W}(b)$ -modules of the form

$$0 \longrightarrow M(\bar{\alpha}, \bar{\Delta}) \longrightarrow E \longrightarrow M(\alpha, \Delta) \longrightarrow 0, \quad (3.20)$$

where  $\Delta, \bar{\Delta} \neq 0$ . As a  $\mathbb{C}[\partial]$ -module,  $E \cong \mathbb{C}[\partial]v_{\bar{\Delta}} \oplus \mathbb{C}[\partial]v_{\Delta}$ . The following identities hold in  $E$

$$L_\lambda v_{\Delta} = (\partial + \alpha + \Delta\lambda)v_{\Delta} + f(\partial, \lambda)v_{\bar{\Delta}}, \quad H_\lambda v_{\Delta} = g(\partial, \lambda)v_{\bar{\Delta}}, \quad (3.21)$$

where  $f(\partial, \lambda) = \sum_{n \geq 0} f_n(\partial)\lambda^n$ ,  $g(\partial, \lambda) = \sum_{n \geq 0} g_n(\partial)\lambda^n$ ,  $f_n(\partial), g_n(\partial) \in \mathbb{C}[\partial]$  and  $f_n(\partial) = g_n(\partial) = 0$  for  $n \gg 0$ .

**Lemma 3.5.** *All trivial extensions of the form (3.20) are given by (3.21), where  $f(\partial, \lambda)$  is a scalar multiple of  $(\partial + \alpha + \Delta\lambda)\phi(\partial) - (\partial + \bar{\alpha} + \bar{\Delta}\lambda)\phi(\partial + \lambda)$  and  $g(\partial, \lambda) = 0$ , where  $\phi$  is a polynomial.*

*Proof.* Suppose that (3.20) represents a trivial cocycle. This means that the exact sequence (3.20) is split and hence there exists  $v'_\Delta = \psi(\partial)v_{\Delta} + \phi(\partial)v_{\bar{\Delta}} \in E$ , such that

$$L_\lambda v'_\Delta = (\partial + \alpha + \Delta\lambda)v'_\Delta = (\partial + \alpha + \Delta\lambda)(\psi(\partial)v_{\Delta} + \phi(\partial)v_{\bar{\Delta}}),$$

and  $H_\lambda v'_\Delta = 0$ . On the other hand, a short computation shows that

$$\begin{aligned} L_\lambda v'_\Delta &= L_\lambda(\psi(\partial)v_{\Delta} + \phi(\partial)v_{\bar{\Delta}}) \\ &= \psi(\partial + \lambda)L_\lambda v_{\Delta} + \phi(\partial + \lambda)L_\lambda v_{\bar{\Delta}} \\ &= (\partial + \alpha + \Delta\lambda)\psi(\partial + \lambda)v_{\Delta} + (\psi(\partial + \lambda)f(\partial, \lambda) + \phi(\partial + \lambda)(\partial + \bar{\alpha} + \bar{\Delta}\lambda))v_{\bar{\Delta}} \end{aligned}$$

Comparing both expressions for  $L_\lambda v'_\Delta$  gives

$$\begin{aligned} (\partial + \alpha + \Delta\lambda)\psi(\partial) &= (\partial + \alpha + \Delta\lambda)\psi(\partial + \lambda), \\ (\partial + \alpha + \Delta\lambda)\phi(\partial) &= f(\partial, \lambda)\psi(\partial + \lambda) + (\partial + \bar{\alpha} + \bar{\Delta}\lambda)\phi(\partial + \lambda), \end{aligned}$$

which imply that  $\psi(\partial) = a$  with  $a$  being a nonzero complex number and hence  $f(\partial, \lambda)$  is a scalar multiple of  $(\partial + \alpha + \Delta\lambda)\phi(\partial) - (\partial + \bar{\alpha} + \bar{\Delta}\lambda)\phi(\partial + \lambda)$ . Similarly, we have

$$H_\lambda v'_\Delta = ag(\partial, \lambda)v_{\bar{\Delta}} = 0,$$

leading to  $g(\partial, \lambda) = 0$ .  $\square$

Applying both sides of (3.1) and (3.2) to  $v_\Delta$  gives the following functional equations:

$$\begin{aligned} (\lambda - \mu)f(\partial, \lambda + \mu) &= (\partial + \lambda + \Delta\mu + \alpha)f(\partial, \lambda) + (\partial + \bar{\Delta}\lambda + \bar{\alpha})f(\partial + \lambda, \mu) \\ &\quad - (\partial + \mu + \Delta\lambda + \alpha)f(\partial, \mu) - (\partial + \bar{\Delta}\mu + \bar{\alpha})f(\partial + \mu, \lambda), \end{aligned} \quad (3.22)$$

$$-(b\lambda + \mu)g(\partial, \lambda + \mu) = (\partial + \bar{\Delta}\lambda + \bar{\alpha})g(\partial + \lambda, \mu) - (\partial + \mu + \Delta\lambda + \alpha)g(\partial, \mu). \quad (3.23)$$

Putting  $\lambda = 0$  in (3.22) and (3.23) gives that

$$(\alpha - \bar{\alpha})f(\partial, \mu) = (\partial + \Delta\mu + \alpha)f(\partial, 0) - (\partial + \bar{\Delta}\mu + \bar{\alpha})f(\partial + \mu, 0), \quad (3.24)$$

$$(\alpha - \bar{\alpha})g(\partial, \mu) = 0. \quad (3.25)$$

**Case 1.**  $\alpha \neq \bar{\alpha}$ .

By (3.24) and (3.25), we have  $f(\partial, \mu) = \frac{1}{\alpha - \bar{\alpha}}((\partial + \Delta\mu + \alpha)f(\partial, 0) - (\partial + \bar{\Delta}\mu + \bar{\alpha})f(\partial + \mu, 0))$  and  $g(\partial, \mu) = 0$ . This corresponds to the trivial extension by Lemma 3.5.

**Case 2.**  $\alpha = \bar{\alpha}$ .

For convenience, we put  $\bar{\partial} = \partial + \alpha$  and let  $\bar{f}(\bar{\partial}, \lambda) = f(\bar{\partial} - \alpha, \lambda)$ ,  $\bar{g}(\bar{\partial}, \lambda) = g(\bar{\partial} - \alpha, \lambda)$ . In what follows we will continue to write  $\partial$  for  $\bar{\partial}$ ,  $f$  for  $\bar{f}$  and  $g$  for  $\bar{g}$ . Now we can rewrite (3.22) and (3.23) as follows:

$$\begin{aligned} (\lambda - \mu)f(\partial, \lambda + \mu) &= (\partial + \lambda + \Delta\mu)f(\partial, \lambda) + (\partial + \bar{\Delta}\lambda)f(\partial + \lambda, \mu) \\ &\quad - (\partial + \mu + \Delta\lambda)f(\partial, \mu) - (\partial + \bar{\Delta}\mu)f(\partial + \mu, \lambda), \end{aligned} \quad (3.26)$$

$$-(b\lambda + \mu)g(\partial, \lambda + \mu) = (\partial + \bar{\Delta}\lambda)g(\partial + \lambda, \mu) - (\partial + \mu + \Delta\lambda)g(\partial, \mu). \quad (3.27)$$

Setting  $\mu = 0$  in (3.27) gives

$$b\lambda g(\partial, \lambda) = (\partial + \Delta\lambda)g(\partial, 0) - (\partial + \bar{\Delta}\lambda)g(\partial + \lambda, 0). \quad (3.28)$$

By the nature of (3.27), we may assume that a solution is a homogeneous polynomial in  $\partial$  and  $\lambda$  of degree  $m$ . Hence we will assume from now on that  $g(\partial, \lambda) = \sum_{i=0}^m a_i \partial^{m-i} \lambda^i$ ,  $a_i \in \mathbb{C}$ . Substituting this into (3.28) gives

$$\sum_{i=0}^m b a_i \partial^{m-i} \lambda^{i+1} = (\partial + \Delta\lambda) a_0 \partial^m - (\partial + \bar{\Delta}\lambda) a_0 (\partial + \lambda)^m. \quad (3.29)$$

We see that  $a_i = 0$  for  $i = 0, 1, \dots, m$  if  $a_0 = 0$ . Then  $g(\partial, \lambda) = 0$ , and it reduces to the Virasoro case.

Now we assume that  $a_0 \neq 0$ . Comparing the coefficients of  $\partial^m \lambda$  in (3.29) gives

$$\Delta - \bar{\Delta} = m + b. \quad (3.30)$$

Equating the coefficients of  $\partial^{m-i} \lambda^{i+1}$  in both sides of (3.29), we have

$$b a_i = -a_0 \binom{m}{i+1} - a_0 \bar{\Delta} \binom{m}{i}, \quad 1 \leq i \leq m. \quad (3.31)$$

Plugging  $g(\partial, \lambda) = \sum_{i=0}^m a_i \partial^{m-i} \lambda^i$  into (3.27) gives

$$-(b\lambda + \mu) \sum_{i=0}^m a_i \partial^{m-i} (\lambda + \mu)^i = (\partial + \bar{\Delta}\lambda) \sum_{i=0}^m a_i (\partial + \lambda)^{m-i} \mu^i - (\partial + \mu + \Delta\lambda) \sum_{i=0}^m a_i \partial^{m-i} \mu^i \quad (3.32)$$

Comparing the coefficients of  $\lambda^m \mu$  in (3.32) with  $m \geq 2$  gives  $(-bm - 1)a_m = \bar{\Delta}a_1$  and hence by (3.31),

$$(-bm - 1)\bar{\Delta} = \binom{m}{2} + m\bar{\Delta}. \quad (3.33)$$

Since  $\bar{\Delta} \neq 0$ ,

$$-bm - 1 = \binom{m}{2} + m\bar{\Delta}. \quad (3.34)$$

Similarly, collecting the coefficients of  $\partial \lambda^{m-1} \mu$  in (3.32) with  $m \geq 3$  and using (3.31) again, we have

$$(-b(m-1) - 1)(1 + m\bar{\Delta}) = (1 + (m-1)\bar{\Delta}) \left( \binom{m}{2} + m\bar{\Delta} \right). \quad (3.35)$$

Combining (3.34) with (3.35) gives  $\bar{\Delta} = b$ . Plugging this into (3.34), we obtain

$$\binom{m}{2} + 2bm + 1 = 0. \quad (3.36)$$

Assume that  $m \geq 4$ . In this case the coefficients of  $\partial^{m-2} \lambda^2 \mu$  in (3.32) gives [using  $\bar{\Delta} = b$  and (3.31) again]

$$(-2b - 1) \left( \binom{m}{3} + \binom{m}{2} b \right) = \left( \binom{m-1}{2} + (m-1)b \right) \left( \binom{m}{2} + bm \right). \quad (3.37)$$

Combining (3.36) with (3.37), we have  $m = -2, -1, 2$  or  $3$ , a contradiction. Therefore,  $m$  is at most  $3$ , i.e.,  $m = 0, 1, 2, 3$ .

For  $m = 0$ , we have  $\Delta - \bar{\Delta} = b$  and  $g(\partial, \lambda) = a_0$  is a solution to (3.27).

For  $m = 1$ , we have  $\Delta - \bar{\Delta} = 1 + b$ . And it follows from (3.31) that  $g(\partial, \lambda) = a_0(\partial - \frac{1}{b}\bar{\Delta}\lambda)$ , which is easily checked to be a solution to (3.27).

Now assume that  $m = 2$ . By (3.30) and (3.34), we have  $\bar{\Delta} = -1 - b$  and  $\Delta = 1$ . By (3.31), we may assume that  $g(\partial, \lambda) = \partial^2 - (\frac{1}{b} + \frac{2}{b}\bar{\Delta})\partial\lambda - \frac{1}{b}\bar{\Delta}\lambda^2$ . Plugging this back into (3.27), we obtain

$$\begin{aligned} & -(b\lambda + \mu) \left( \partial^2 - \left( \frac{1}{b} + \frac{2}{b}\bar{\Delta} \right) \partial(\lambda + \mu) - \frac{1}{b}\bar{\Delta}(\lambda + \mu)^2 \right) \\ &= (\partial + \bar{\Delta}\lambda) \left( (\partial + \lambda)^2 - \left( \frac{1}{b} + \frac{2}{b}\bar{\Delta} \right) (\partial + \lambda)\mu - \frac{1}{b}\bar{\Delta}\mu^2 \right) \\ & \quad - (\partial + \mu + \Delta\lambda) \left( \partial^2 - \left( \frac{1}{b} + \frac{2}{b}\bar{\Delta} \right) \partial\mu - \frac{1}{b}\bar{\Delta}\mu^2 \right), \end{aligned}$$

which holds provided that  $\bar{\Delta} = -1 - b$  and  $\Delta = 1$ .

Finally we consider the case  $m = 3$ . By (3.36) and (3.30), we have  $\bar{\Delta} = b = -\frac{2}{3}$  and  $\Delta = \frac{5}{3}$ . By (3.31), we may assume that  $g(\partial, \lambda) = \partial^3 + \frac{3}{2}\partial^2\lambda - \frac{3}{2}\partial\lambda^2 - \lambda^3$ . Plugging this back into (3.27), we obtain that it is a solution.

The above discussions prove the following:

**Lemma 3.6.** *Let  $g(\partial, \lambda)$  be a nonzero homogeneous polynomial of degree  $m$  satisfying (3.27). Then  $\Delta - \bar{\Delta} = m + b$  and  $m \leq 3$ . Furthermore, we have*

- (1) For  $b = -\frac{2}{3}$ , all the solutions (up to a scalar) to (3.27) are given by

- (i)  $m = 0, \Delta - \bar{\Delta} = -\frac{2}{3},$  and  $g(\partial, \lambda) = 1;$
- (ii)  $m = 1, \Delta - \bar{\Delta} = \frac{1}{3},$  and  $g(\partial, \lambda) = \partial + \frac{3}{2}\bar{\Delta}\lambda;$
- (iii)  $m = 2, \Delta = 1, \bar{\Delta} = -\frac{1}{3},$  and  $g(\partial, \lambda) = \partial^2 + \frac{1}{2}\partial\lambda - \frac{1}{2}\lambda^2;$
- (iv)  $m = 3, \Delta = \frac{5}{3}, \bar{\Delta} = -\frac{2}{3},$  and  $g(\partial, \lambda) = \partial^3 + \frac{3}{2}\partial^2\lambda - \frac{3}{2}\partial\lambda^2 - \lambda^3,$

(2) For  $b \neq -\frac{2}{3},$  all the solutions (up to a scalar) to (3.27) are given by

- (i)  $m = 0, \Delta - \bar{\Delta} = b,$  and  $g(\partial, \lambda) = 1;$
- (ii)  $m = 1, \Delta - \bar{\Delta} = 1 + b,$  and  $g(\partial, \lambda) = \partial - \frac{1}{b}\bar{\Delta}\lambda;$
- (iii)  $m = 2, \Delta = 1, \bar{\Delta} = -1 - b$  and  $g(\partial, \lambda) = \partial^2 - (\frac{1}{b} + \frac{2}{b}\bar{\Delta})\partial\lambda - \frac{1}{b}\bar{\Delta}\lambda^2.$

By Lemma 3.6 and Theorem 2.4, we obtain the following.

**Theorem 3.7.** *Nontrivial extensions of  $\mathscr{W}(b)$ -modules of the form (3.20) exist only if  $\alpha = \bar{\alpha}$ . For each  $\alpha \in \mathbb{C},$  these extensions are given, up to equivalence, by*

$$L_\lambda v_\Delta = (\partial + \alpha + \Delta\lambda)v_\Delta + f(\partial, \lambda)v_{\bar{\Delta}}, \quad H_\lambda v_\Delta = g(\partial, \lambda)v_{\bar{\Delta}},$$

where  $g(\partial, \lambda) = 0$  and  $f(\partial, \lambda)$  is from the nonzero polynomials of Theorem 2.4 with  $\Delta, \bar{\Delta} \neq 0,$  or the values of  $\Delta$  and  $\bar{\Delta}$  along with the pairs of polynomials  $g(\partial, \lambda)$  and  $f(\partial, \lambda),$  whose nonzero scalar multiples give rise to nontrivial extensions, are listed as follows (by replacing  $\partial$  by  $\partial + \alpha$ ):

- (1) When  $b = -1,$  we have  $\Delta - \bar{\Delta} = -1$  or  $0.$  And
  - (i) In the case  $\Delta - \bar{\Delta} = -1,$   $f(\partial, \lambda) = 0$  and  $g(\partial, \lambda) = a_0,$  where  $a_0 \neq 0.$
  - (ii) In the case  $\Delta - \bar{\Delta} = 0,$   $f(\partial, \lambda) = c_0 + c_1\lambda,$  and  $g(\partial, \lambda) = \partial + \bar{\Delta}\lambda,$  where  $c_0, c_1 \in \mathbb{C}.$
- (2) When  $b = 1,$  we have  $\Delta - \bar{\Delta} = 1, 2$  or  $\Delta = 1, \bar{\Delta} = -2.$  And
  - (i) In the case  $\Delta - \bar{\Delta} = 1,$   $f(\partial, \lambda) = 0$  and  $g(\partial, \lambda) = a_0,$  where  $a_0 \neq 0.$
  - (ii) In the case  $\Delta - \bar{\Delta} = 2,$   $f(\partial, \lambda) = c_0\lambda^2(2\partial + \lambda)$  and  $g(\partial, \lambda) = \partial - \bar{\Delta}\lambda,$  where  $c_0 \in \mathbb{C}.$
  - (iii) In the case  $\Delta = 1, \bar{\Delta} = -2,$   $f(\partial, \lambda) = c_0\partial\lambda^2(\partial + \lambda)$  and  $g(\partial, \lambda) = \partial^2 + 3\partial\lambda + 2\lambda^2,$  where  $c_0 \in \mathbb{C}.$
- (3) When  $b = 2,$  we have  $\Delta - \bar{\Delta} = 2, 3$  or  $\Delta = 1, \bar{\Delta} = -3.$  And
  - (i) In the case  $\Delta - \bar{\Delta} = 2,$   $f(\partial, \lambda) = c_0\lambda^2(2\partial + \lambda)$  and  $g(\partial, \lambda) = a_0,$  where  $c_0 \in \mathbb{C}, a_0 \neq 0.$
  - (ii) In the case  $\Delta - \bar{\Delta} = 3,$   $f(\partial, \lambda) = c_0\partial\lambda^2(\partial + \lambda)$  and  $g(\partial, \lambda) = \partial - \frac{1}{2}\bar{\Delta}\lambda,$  where  $c_0 \in \mathbb{C}.$
  - (iii) In the case  $\Delta = 1, \bar{\Delta} = -3,$   $f(\partial, \lambda) = c_0\lambda^2(4\partial^3 + 6\partial^2\lambda - \partial\lambda^2 + 3\lambda^3)$  and  $g(\partial, \lambda) = \partial^2 + \frac{5}{2}\partial\lambda + \frac{3}{2}\lambda^2,$  where  $c_0 \in \mathbb{C}.$
- (4) When  $b = 3,$  we have  $\Delta - \bar{\Delta} = 3, 4$  or  $\Delta = 1, \bar{\Delta} = -4.$  And
  - (i) In the case  $\Delta - \bar{\Delta} = 3,$   $f(\partial, \lambda) = c_0\partial\lambda^2(\partial + \lambda)$  and  $g(\partial, \lambda) = a_0,$  where  $c_0 \in \mathbb{C}, a_0 \neq 0.$
  - (ii) In the case  $\Delta - \bar{\Delta} = 4,$   $f(\partial, \lambda) = c_0\lambda^2(4\partial^3 + 6\partial^2\lambda - \partial\lambda^2 + \bar{\Delta}\lambda^3)$  and  $g(\partial, \lambda) = \partial - \frac{1}{3}\bar{\Delta}\lambda,$  where  $c_0 \in \mathbb{C}.$

(iii) In the case  $\Delta = 1, \bar{\Delta} = -4$ ,  $f(\partial, \lambda) = c_0(\partial^4\lambda^2 - 10\partial^2\lambda^4 - 17\partial\lambda^5 - 8\lambda^6)$  and  $g(\partial, \lambda) = \partial^2 + \frac{7}{3}\partial\lambda + \frac{4}{3}\lambda^2$ , where  $c_0 \in \mathbb{C}$ .

(5) When  $b = 4$ , we have  $\Delta - \bar{\Delta} = 4, 5$  or  $\Delta = 1, \bar{\Delta} = -5$ . And

(i) In the case  $\Delta - \bar{\Delta} = 4$ ,  $f(\partial, \lambda) = c_0\lambda^2(4\partial^3 + 6\partial^2\lambda - \partial\lambda^2 + \bar{\Delta}\lambda^3)$  and  $g(\partial, \lambda) = a_0$ , where  $c_0 \in \mathbb{C}, a_0 \neq 0$ .

(ii) In the case  $\Delta - \bar{\Delta} = 5$ ,  $f(\partial, \lambda) = 0$  and  $g(\partial, \lambda) = \partial - \frac{1}{4}\bar{\Delta}\lambda$ .

(ii') In the case  $\Delta = 1, \bar{\Delta} = -4$ ,  $f(\partial, \lambda) = c_0(\partial^4\lambda^2 - 10\partial^2\lambda^4 - 17\partial\lambda^5 - 8\lambda^6)$  and  $g(\partial, \lambda) = \partial + \lambda$ , where  $c_0 \in \mathbb{C}$ .

(iii) In the case  $\Delta = 1, \bar{\Delta} = -5$ ,  $f(\partial, \lambda) = 0$  and  $g(\partial, \lambda) = \partial^2 + \frac{9}{4}\partial\lambda + \frac{5}{4}\lambda^2$ .

(6) When  $b = 5$ , we have  $\Delta - \bar{\Delta} = 5, 6$  or  $\Delta = 1, \bar{\Delta} = -6$ . And

(i) In the case  $\Delta - \bar{\Delta} = 5$ ,  $f(\partial, \lambda) = 0$  and  $g(\partial, \lambda) = a_0$ , where  $a_0 \neq 0$ .

(i') In the case  $\Delta = 1, \bar{\Delta} = -4$ ,  $f(\partial, \lambda) = c_0(\partial^4\lambda^2 - 10\partial^2\lambda^4 - 17\partial\lambda^5 - 8\lambda^6)$  and  $g(\partial, \lambda) = a_0$ , where  $c_0 \in \mathbb{C}, a_0 \neq 0$ .

(ii) In the case  $\Delta - \bar{\Delta} = 6$ ,  $f(\partial, \lambda) = 0$  and  $g(\partial, \lambda) = \partial - \frac{1}{5}\bar{\Delta}\lambda$ .

(ii') In the case  $\Delta = \frac{7}{2} \pm \frac{\sqrt{19}}{2}, \Delta - \bar{\Delta} = 6$ ,  $f(\partial, \lambda) = c_0(\partial^4\lambda^3 - (2\bar{\Delta} + 3)\partial^3\lambda^4 - 3\bar{\Delta}\partial^2\lambda^5 - (3\bar{\Delta} + 1)\partial\lambda^6 - (\bar{\Delta} + \frac{9}{28})\lambda^7)$  and  $g(\partial, \lambda) = \partial - \frac{1}{5}\bar{\Delta}\lambda$ , where  $c_0 \in \mathbb{C}$ .

(iii) In the case  $\Delta = 1, \bar{\Delta} = -6$ ,  $f(\partial, \lambda) = 0$  and  $g(\partial, \lambda) = \partial^2 + \frac{11}{5}\partial\lambda + \frac{6}{5}\lambda^2$ .

(7) When  $b = 6$ , we have  $\Delta - \bar{\Delta} = 6, 7$  or  $\Delta = 1, \bar{\Delta} = -7$ . And

(i) In the case  $\Delta - \bar{\Delta} = 6$ ,  $f(\partial, \lambda) = 0$  and  $g(\partial, \lambda) = a_0$ , where  $a_0 \neq 0$ .

(i') In the case  $\Delta = \frac{7}{2} \pm \frac{\sqrt{19}}{2}, \bar{\Delta} = -\frac{5}{2} \pm \frac{\sqrt{19}}{2}$ ,  $f(\partial, \lambda) = c_0(\partial^4\lambda^3 - (2\bar{\Delta} + 3)\partial^3\lambda^4 - 3\bar{\Delta}\partial^2\lambda^5 - (3\bar{\Delta} + 1)\partial\lambda^6 - (\bar{\Delta} + \frac{9}{28})\lambda^7)$  and  $g(\partial, \lambda) = a_0$ , where  $c_0 \in \mathbb{C}, a_0 \neq 0$ .

(ii) In the case  $\Delta - \bar{\Delta} = 7$ ,  $f(\partial, \lambda) = 0$  and  $g(\partial, \lambda) = \partial - \frac{1}{6}\bar{\Delta}\lambda$ .

(iii) In the case  $\Delta = 1, \bar{\Delta} = -7$ ,  $f(\partial, \lambda) = 0$  and  $g(\partial, \lambda) = \partial^2 + \frac{13}{6}\partial\lambda + \frac{7}{6}\lambda^2$ .

(8) When  $b = -\frac{2}{3}$ , we have  $f(\partial, \lambda) = 0$  and the values  $\Delta$  and  $\bar{\Delta}$  along with  $g(\partial, \lambda)$  are from Lemma 3.6(1).

(9) When  $b \neq -1, 1, 2, 3, 4, 5, 6$  or  $-\frac{2}{3}$ ,  $f(\partial, \lambda) = 0$  and the values  $\Delta$  and  $\bar{\Delta}$  along with  $g(\partial, \lambda)$  are from Lemma 3.6(2).

**Acknowledgements.** This work was supported by National Natural Science Foundation of China (11301109) and China Scholarship Council.

## References

- [1] Cheng S.-J., Kac V., Conformal modules, *Asian J. Math.*, **1**(1) (1997) 181–193.
- [2] Cheng S.-J., Kac V., Wakimoto, M., Extensions of conformal modules, *In Topological Field Theory, Primitive Forms and Related Topics*, *Progr. Math.* Vol. 160., Kashiwara, M., et al. Eds. Boston: Birkhäuser, 1998; 79-129.

- [3] Cheng S.-J., Kac V., Wakimoto, M., Extensions of Neveu-Schwarz conformal modules, *J. Math. Phys.*, **41**(4) (2000) 2271–2294.
- [4] D’Andrea A., Kac V., Structure theory of finite conformal algebras, *Sel. Math., New Ser.*, **4** (1998) 377–418.
- [5] Lam Ngau, Extensions of modules over supercurrent conformal algebras, *Comm. Algebra*, **29**(7) (2001) 3061–3068.
- [6] Ling K., Yuan L., Extensions of modules over Heisenberg-Virasoro conformal algebra, *Internat. J. Math.*, **28**(5) (2017) 1750036, 13pp.
- [7] Xu Y., Yue X.,  $W(a,b)$  Lie conformal algebra and its conformal module of rank one, *Algebra Colloq.*, **22**(3) (2015) 405–412.
- [8] Wu H., Yuan L., Classification of finite irreducible conformal modules over some Lie conformal algebras related to the Virasoro conformal algebra, *J. Math. Phys.*, **58** (2017) 041701, 10pp.
- [9] Yuan L., Wu H., Cohomology of Heisenberg-Virasoro conformal algebra, *J. Lie Theory*, **26** (2016) 1187–1197.
- [10] Yuan L., Wu H., Structures of  $W(2,2)$  Lie conformal algebra, *Open Math.*, **14** (2016) 629–640.