

KSBA COMPACTIFICATION OF THE MODULI SPACE OF K3 SURFACES WITH PURELY NON-SYMPLECTIC AUTOMORPHISM OF ORDER FOUR

HAN-BOM MOON AND LUCA SCHAFFLER

ABSTRACT. We describe a compactification by KSBA stable pairs of the five-dimensional moduli space of K3 surfaces with purely non-symplectic automorphism of order four and $U(2) \oplus D_4^{\oplus 2}$ lattice polarization. These K3 surfaces can be realized as the minimal resolution of the double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ branched along a specific $(4, 4)$ curve. We show that, up to a finite group action, this stable pair compactification is isomorphic to Kirwan's partial desingularization of the GIT quotient $(\mathbb{P}^1)^8 // \mathrm{SL}_2$ with the symmetric linearization.

1. INTRODUCTION

Recent advances in algebraic geometry including the minimal model program and the boundedness for stable pairs, enables us to investigate compactifications of moduli spaces of higher dimensional algebraic varieties, in particular the Kollár, Shepherd-Barron, and Alexeev (KSBA) moduli space of stable pairs [KSB88, Ale96, Kol18]. However, the geometry of the moduli spaces of higher dimensional varieties is extremely complicated. For instance, these moduli spaces are rarely irreducible, and they may even have arbitrary singularity types [Mnë85, Vak06].

Nonetheless, sometimes we may understand in detail the geometry of moduli spaces of special algebraic varieties of interest. These explicit moduli spaces are beneficial because in many cases the generalities are out of reach, and also they reveal interesting geometric behaviors (see [AP09, Sch16, GMGZ18, AB19, AET19, DH21]). In this paper, we study one of such explicit examples: the moduli space of K3 surfaces with a purely non-symplectic automorphism of order four (see Definition 2.2). The geometry of K3 surfaces with purely non-symplectic automorphism of order four was studied in [AS15], and for a survey on the subject in general we refer to [Zha07].

Here we state our main result. For the precise definitions and terminology, see §2. Consider the moduli space of K3 surfaces \tilde{X} with a purely non-symplectic automorphism of order four together with a $U(2) \oplus D_4^{\oplus 2}$ lattice polarization. By Kondō's work [Kon07], there is a five dimensional irreducible moduli space \mathbf{M} of such K3 surfaces. These K3 surfaces can be obtained by taking the minimal resolution of the double cover X of $\mathbb{P}^1 \times \mathbb{P}^1$ branched along an appropriate divisor B of class $(4, 4)$.

We adopt the KSBA theory to compactify \mathbf{M} . Let $\overline{\mathbf{K}}$ be the normalization of the closure in the KSBA moduli space of stable pairs of the locus parametrizing $(X, \epsilon R)$ where X is the double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ branched along B , R is the ramification divisor, and $0 < \epsilon \ll 1$.

Mathematics Subject Classification (2010): 14J10, 14J28, 14D06.

Theorem 1.1. *The KSBA compactification $\overline{\mathbf{K}}$ is isomorphic to \mathbf{P}/H where $H \cong (S_4 \times S_4) \rtimes S_2$ and \mathbf{P} is the partial desingularization of the GIT quotient $(\mathbb{P}^1)^8 // \mathrm{SL}_2$ with the symmetric linearization.*

Note that \mathbf{P} has another moduli theoretic interpretation: \mathbf{P} is isomorphic to the Hassett's moduli space of weighted pointed curves $\overline{\mathbf{M}}_{0,(\frac{1}{4}+\epsilon)^8}$ [KM11, Theorem 1.1].

1.1. K3 surfaces from eight points on \mathbb{P}^1 and GIT. Here we elaborate more on the construction of these K3 surfaces. Fix eight distinct points $([\lambda_1 : 1], \dots, [\lambda_8 : 1])$ on \mathbb{P}^1 . Let $[x_0 : x_1], [y_0 : y_1]$ be the homogeneous coordinates of $\mathbb{P}^1 \times \mathbb{P}^1$, and define B to be the following curve of the class $(4, 4)$:

$$y_0 y_1 \left(y_0^2 \prod_{i=1}^4 (x_0 - \lambda_i x_1) + y_1^2 \prod_{i=5}^8 (x_0 - \lambda_i x_1) \right) = 0.$$

Let X be the double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ branched along B . In [Kon07], it was shown that 1) the minimal resolution \tilde{X} of X is a K3 surface with a purely non-symplectic automorphism of order four, so $\mathbf{U} \subseteq (\mathbb{P}^1)^8$ parametrizing eight distinct points also parametrizes K3 surfaces, 2) the construction is SL_2 -invariant and S_8 -invariant (see [Kon07, §2.1] and §2.3 of this paper), so $\mathbf{M} = \mathbf{U}/\mathrm{SL}_2/S_8$ can be regarded as a parameter space of such K3 surfaces. The involution $([x_0 : x_1], [y_0 : y_1]) \mapsto ([x_0 : x_1], [y_0 : -y_1])$ is lifted to a purely non-symplectic automorphism σ of order four on \tilde{X} .

Let $H^2(\tilde{X}, \mathbb{Z})^+$ be the invariant subspace of $H^2(\tilde{X}, \mathbb{Z})$ with respect to the $(\sigma^2)^*$ -action. Then for any $\mathbf{p} \in \mathbf{M}$, the associated $H^2(\tilde{X}, \mathbb{Z})^+$ is a primitive sublattice of $\mathrm{NS}(\tilde{X})$ isometric to $U(2) \oplus D_4^{\oplus 2}$. In summary, the GIT quotient $(\mathbb{P}^1)^8 // \mathrm{SL}_2/S_8$ with the symmetric linearization can be thought of as a compactification of the moduli space of K3 surfaces in analysis.

1.2. KSBA compactification. To adopt the KSBA theory in this context, one has to choose an ample divisor A on \tilde{X} and make a pair (\tilde{X}, A) . However, we make two minor modifications to the moduli problem we consider. First of all, instead of taking an ample divisor A , we choose a big and nef divisor on \tilde{X} , which is the pull-back of the ramification divisor R on X . This makes the description of the parameter space more accessible by using the theory of abelian covers [AP12]. Secondly, instead of taking the entire linear system of R , we just take R to make a five-dimensional moduli space of pairs.

Technically, the resulting moduli space $\overline{\mathbf{K}}$ is a compactification of a finite cover of \mathbf{M} because the same K3 surface with a purely non-symplectic automorphism of order four may be regarded as a minimal resolution of the double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ in several ways. However, compared to other ways of compactifying \mathbf{M} , the compactification $\overline{\mathbf{K}}$ has the merit to have an explicit moduli theoretic meaning.

1.3. Relation to Hodge theoretic compactifications. Hodge theory is another standard tool one could use to compactify a given moduli space of varieties. Here we leave some related works. In [DM86], Deligne and Mostow proved that the symmetric GIT quotient $(\mathbb{P}^1)^8 // \mathrm{SL}_2$ is isomorphic to the Satake–Baily–Borel compactification $\overline{\mathcal{B}}/\Gamma^*$ of the quotient

of a five-dimensional complex ball \mathcal{B} by an arithmetic group Γ . This is proved using periods of a family of curves arising as the $\mathbb{Z}/4\mathbb{Z}$ -covers of \mathbb{P}^1 branched along eight points.

There are two interesting problems. One may wonder how GIT compactifications and Hodge theoretic compactifications are related. In the literature, one celebrated example of interaction between Hodge theory and GIT is the case of K3 surfaces with a degree two polarization. More precisely, in [Loo86] it is shown that a small partial resolution of the Satake–Baily–Borel compactification for such K3 surfaces is isomorphic to a partial Kirwan desingularization of the GIT quotient for sextic plane curves. The moduli space of K3 surfaces which are double covers of $\mathbb{P}^1 \times \mathbb{P}^1$ branched along a curve of class $(4, 4)$ were recently investigated in [LO18] from the point of view of GIT and Hodge theory. In our context of moduli space of K3 surfaces with purely non-symplectic automorphism of order four, the GIT compactification and the Satake–Baily–Borel compactification give the same answer after quotienting by S_8 [Kon07].

Usually, GIT and Hodge theoretic compactifications do not have a strong modular interpretation. Thus we have the second interesting problem: finding a modular compactification of the given moduli space. From the perspective of moduli theory, the KSBA compactification is arguably the best known theoretical approach. Motivated by this, in [GKS21] it is shown that the KSBA compactification $\overline{\mathcal{K}}$ is isomorphic to the quotient by $(S_4 \times S_4) \rtimes S_2$ of the unique toroidal compactification of the above Deligne–Mostow ball quotient \mathcal{B}/Γ .

1.4. Structure of the paper. The paper is organized as follows. In §2 we review Kondō’s construction of the 5-dimensional family of K3 surfaces with purely non-symplectic automorphism of order four and $U(2) \oplus D_4^{\oplus 2}$ lattice polarization. In §3 we recall the notion of stable pair, their moduli functor, and the theory of abelian covers. §4 contains a brief summary of Kirwan’s partial desingularization [Kir85], which is applied to the case of $(\mathbb{P}^1)^8//\mathrm{SL}_2$. In §5 we study the KSBA limits of specific one-parameter degenerations of stable pairs $(\mathbb{P}^1 \times \mathbb{P}^1, \frac{1+\epsilon}{2}B)$. These calculations are then used in §6 to finally prove Theorem 1.1.

We work over \mathbb{C} .

ACKNOWLEDGEMENTS

We would like to express our gratitude to Valery Alexeev, Maksym Fedorchuk, Paul Hacking, Giovanni Inchiostro, Jenia Tevelev, and Alan Thompson for helpful conversations and suggestions. We also thank the anonymous referee for the valuable comments. Most part of this work was done while the first author was a member of the Institute for Advanced Study. The first author was partially supported by the Minerva Research Foundation.

2. K3 SURFACES FROM EIGHT POINTS ON \mathbb{P}^1

2.1. Kondō’s construction.

Definition 2.1. A normal surface X is called an *ADE K3 surface* if its minimal resolution is a smooth K3 surface, or equivalently,

- (1) X has only ADE singularities (thus it is Gorenstein);
- (2) $\omega_X \cong \mathcal{O}_X$;
- (3) $H^1(X, \mathcal{O}_X) = 0$.

In [Kon07], a K3 surface is constructed from the data of eight distinct points on \mathbb{P}^1 as follows. Up to the natural SL_2 -action on \mathbb{P}^1 , we may assume that the eight points are in the form $[\lambda_1 : 1], \dots, [\lambda_8 : 1]$. Let C be the curve in $\mathbb{P}^1 \times \mathbb{P}^1$ given by

$$(1) \quad y_0^2 \prod_{i=1}^4 (x_0 - \lambda_i x_1) + y_1^2 \prod_{i=5}^8 (x_0 - \lambda_i x_1) = 0,$$

where $([x_0 : x_1], [y_0 : y_1])$ are coordinates in $\mathbb{P}^1 \times \mathbb{P}^1$. If all λ_i 's are distinct, C is a smooth curve. Let L_i be the line $y_i = 0, i = 1, 2$. The double cover $\pi : X \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ branched along $C + L_0 + L_1$, which has bidegree $(4, 4)$, has eight A_1 singularities which lie above $C \cap L_0$ and $C \cap L_1$. The minimal resolution $\rho : \tilde{X} \rightarrow X$ of this double cover is a K3 surface. Thus X is an ADE K3 surface with a polarization $F := \pi^* \mathcal{O}(1, 1)$ of degree 4. Also \tilde{X} carries a natural big and nef polarization $\rho^* F$.

Definition 2.2. An automorphism σ of a K3 surface is *non-symplectic* if the induced automorphism on the global sections of the canonical sheaf is not the identity. In addition, we say that σ is *purely non-symplectic* if all its non-trivial powers are non-symplectic.

As Kondō described in [Kon07, §2], a K3 surface \tilde{X} as above admits a purely non-symplectic automorphism σ of order four given by the lift of the involution

$$([x_0 : x_1], [y_0 : y_1]) \mapsto ([x_0 : x_1], [y_0 : -y_1]).$$

The lattice $H^2(\tilde{X}, \mathbb{Z})^+ := \{x \in H^2(\tilde{X}, \mathbb{Z}) \mid (\sigma^2)^*(x) = x\}$, which embeds primitively into $NS(\tilde{X})$, is isometric to $U(2) \oplus D_4^{\oplus 2}$, and $H^2(\tilde{X}, \mathbb{Z})^- := \{x \in H^2(\tilde{X}, \mathbb{Z}) \mid (\sigma^2)^*(x) = -x\}$ is isometric to $U \oplus U(2) \oplus D_4^{\oplus 2}$ [Kon07, Lemma 2.2].

Definition 2.3. Let M be the coarse moduli space of K3 surfaces \tilde{X} with a purely non-symplectic automorphism of order four such that $H^2(\tilde{X}, \mathbb{Z})^+$ is isometric to $U(2) \oplus D_4^{\oplus 2}$.

Once $M := H^2(\tilde{X}, \mathbb{Z})^+$ is identified with $U(2) \oplus D_4^{\oplus 2}$, the lattice $N := H^2(\tilde{X}, \mathbb{Z})^- \cong U \oplus U(2) \oplus D_4^{\oplus 2}$ is given because $N = M^\perp$. On $N \otimes \mathbb{C}$, the linear map σ^* has minimal polynomial $x^2 + 1$, which implies that σ^* is diagonalizable. Moreover, the only possible eigenvalues of σ^* are $\pm\sqrt{-1}$. These both occur with the same multiplicity because σ^* is a real operator. Hence, if V denotes the eigenspace for $\sqrt{-1}$, then $\dim V = \frac{1}{2} \dim N \otimes \mathbb{C} = 6$. Thus from [DK07, §11] (see also [AS15, §1]), M is the quotient of the ball $\{[z] \in \mathbb{P}(V) \mid z \cdot \bar{z} > 0\}$ by an appropriate arithmetic group. In particular, it is a 5-dimensional irreducible analytic variety.

Remark 2.4. Note that the family of K3 surfaces parametrized by M in Definition 2.3 is the same as the family of K3 surfaces with a purely non-symplectic automorphism of order four and a $U(2) \oplus D_4^{\oplus 2}$ lattice polarization. This is true because the very general member \tilde{X} of the latter family has $H^2(\tilde{X}, \mathbb{Z})^+ = NS(\tilde{X}) \cong U(2) \oplus D_4^{\oplus 2}$.

The above construction of ADE K3 surfaces can be relativized. Let $([a_1 : b_1], \dots, [a_8 : b_8])$ be coordinates in $(\mathbb{P}^1)^8$. Consider the hypersurface $\mathcal{C} \subseteq (\mathbb{P}^1)^8 \times \mathbb{P}^1 \times \mathbb{P}^1$ given by

$$y_0^2 \prod_{i=1}^4 (b_i x_0 - a_i x_1) + y_1^2 \prod_{i=5}^8 (b_i x_0 - a_i x_1) = 0,$$

which has multidegree $(4, \dots, 4, 2, 2)$. It can be understood as a family of curves over $(\mathbb{P}^1)^8$. Let $\mathbf{U} \subseteq (\mathbb{P}^1)^8$ be the open subset consisting of 8-tuples of distinct points. Let $\mathcal{X} \rightarrow \mathbf{U} \times \mathbb{P}^1 \times \mathbb{P}^1$ be the double cover branched along $(\mathcal{C} + \mathcal{L}_0 + \mathcal{L}_1)|_{\mathbf{U}}$, where $\mathcal{L}_i := V(y_i) \subseteq \mathbf{U} \times \mathbb{P}^1 \times \mathbb{P}^1$ for $i = 0, 1$. Since an SL_2 -orbit in \mathbf{U} parametrizes isomorphic ADE K3 surfaces, \mathbf{U}/SL_2 is a five dimensional parameter space of ADE K3 surfaces with a purely non-symplectic automorphism of order four.

Note that there is a natural S_8 action on \mathbf{U}/SL_2 which permutes the eight points.

Definition 2.5. Let $H \cong (S_4 \times S_4) \rtimes S_2$ be the subgroup of permutations of S_8 which is generated by the permutations of the first four points, the permutations of the last four points, and the involution which exchanges the set of first four points and the set of last four points.

Any ADE K3 surfaces parametrized by an H -orbit are isomorphic to each other because $C + L_0 + L_1$ is $(S_4 \times S_4)$ -invariant, and the S_2 -action induces an isomorphism of associated surfaces derived by the involution $[y_0 : y_1] \rightarrow [y_1 : y_0]$.

Furthermore, it was shown by Kondō that two ADE K3 surfaces as above are isomorphic if and only if the associated points on \mathbf{U}/SL_2 are in the same S_8 -orbit [Kon07, §3.7]. We will come back to this S_8 -invariance in §2.3, where we discuss it when some of the eight points on \mathbb{P}^1 collide. In particular, $\mathbf{U}/\mathrm{SL}_2/S_8$ is a dense open subset of \mathbf{M} .

2.2. Degenerate point configurations and GIT. Consider the diagonal SL_2 -action on $(\mathbb{P}^1)^8$ together with the natural symmetric linearization $\mathcal{O}(1, \dots, 1)$. A point in $(\mathbb{P}^1)^8$ is stable (resp. semi-stable) if and only if at most three (resp. four) points coincide. We denote the semi-stable locus (resp. stable locus) by $((\mathbb{P}^1)^8)^{ss}$ (resp. $((\mathbb{P}^1)^8)^s$) [MFK94, §3].

Lemma 2.6. *The construction in §2.1 yields an ADE K3 surface for any $\mathbf{p} \in ((\mathbb{P}^1)^8)^s$.*

Proof. Let $\mathbf{p} = (p_1, \dots, p_8)$ be a stable point configuration with at least one collision. Up to H -symmetry, it is clear that Table 1 describes all the possibilities. By a local computation, we see that the double cover X of $\mathbb{P}^1 \times \mathbb{P}^1$ branched along $C + L_0 + L_1$ has only ADE singularities. The two conditions $\omega_X \cong \mathcal{O}_X$ and $H^1(X, \mathcal{O}_X) = 0$ are easy to check using the fact that X is the double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ branched along a divisor of class $(4, 4)$. \square

If \mathbf{p} is strictly semi-stable, then the associated double cover has worse singularities and it is not an ADE K3 surface. Up to the H -action, there are three cases:

- (1) $p_1 = p_2 = p_5 = p_6$;
- (2) $p_1 = p_2 = p_3 = p_5$;
- (3) $p_1 = p_2 = p_3 = p_4$.

TABLE 1. Degenerate point configurations and singularities on the double cover.

collision	analytic local equation	singularity of double cover
$p_1 = p_2 = [0 : 1]$	$y_1(x_0^2 + y_1^2) = 0$	D_4
$p_1 = p_5 = [0 : 1]$		A_1 singularities
$p_1 = p_2 = p_5 = [0 : 1]$	$x_0 y_1(x_0 + y_1^2) = 0$	D_6
$p_1 = p_2 = p_3 = [0 : 1]$	$y_1(x_0^3 + y_1^2) = 0$	E_7

For cases (2) and (3), we may compute the minimal resolution of the double cover $X \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ branched along $C + L_0 + L_1$ using the *canonical resolution* method [BHPV04, III.7]. The exceptional locus in case (2) is a genus one curve of self-intersection -2 , and in case (3) is a genus one curve of self intersection -1 . These singularities are known as \tilde{E}_7 and \tilde{E}_8 respectively (see [Ish18, §7.6]). In case (1), the ramification divisor is not even reduced.

For later purpose, the strictly semi-stable points with maximal dimensional stabilizer group are important. There are three types of semi-stable points with positive dimensional stabilizer group, which is isomorphic to \mathbb{C}^* .

Definition 2.7. Let $\mathbf{p} = (p_1, \dots, p_8)$ be a strictly semi-stable point configuration. We say that \mathbf{p} is of type a if, up to H -action, it is of the form $p_1 = p_2 = p_5 = p_6$ and $p_3 = p_4 = p_7 = p_8$. Similarly, we say that \mathbf{p} is of type b if it is of the form $p_1 = p_2 = p_3 = p_5$ and $p_4 = p_6 = p_7 = p_8$. Finally, we say that \mathbf{p} is of type c if it is of the form $p_1 = p_2 = p_3 = p_4$ and $p_5 = p_6 = p_7 = p_8$.

The corresponding curve $C \subseteq \mathbb{P}^1 \times \mathbb{P}^1$ is given by:

- (type a) $(x_0 - \lambda_1 x_1)^2(x_0 - \lambda_3 x_1)^2(y_0^2 + y_1^2) = 0$;
- (type b) $(x_0 - \lambda_1 x_1)(x_0 - \lambda_4 x_1)(y_0^2(x_0 - \lambda_1 x_1)^2 + y_1^2(x_0 - \lambda_4 x_1)^2) = 0$;
- (type c) $y_0^2(x_0 - \lambda_1 x_1)^4 + y_1^2(x_0 - \lambda_5 x_1)^4 = 0$.

Each connected component of such strictly semi-stable points is isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1 \setminus \Delta$, where Δ is the diagonal. The locus of type a point configurations has 18 connected components, that of type b configurations has 16 connected components, and that of type c configurations has a unique component.

Remark 2.8. The associated double covers of $\mathbb{P}^1 \times \mathbb{P}^1$ branched along $C + L_0 + L_1$ with C of type a, type b, and type c appear in Shah's list [Sha81, Theorem 4.8, B, Type II, (i)–(iii)].

2.3. S_8 -invariance of Kondō's K3 surfaces. Let X be an ADE K3 surface associated to 8 distinct points on \mathbb{P}^1 . In [Kon07] it is observed that the isomorphism class of X is independent from the ordering of the eight points. Here we give a proof which is valid for all stable configurations.

Proposition 2.9. *Let X be an ADE K3 surface which is the double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ branched along the curve*

$$B : y_0 y_1 \left(y_0^2 \prod_{i=1}^4 (x_0 - \lambda_i x_1) + y_1^2 \prod_{i=5}^8 (x_0 - \lambda_i x_1) \right) = 0.$$

Let $\tau \in S_8$ and define X_τ to be the ADE K3 surface obtained as the double cover of $\mathbb{P}^1 \times \mathbb{P}^1$ branched along the curve

$$B_\tau : y_0 y_1 \left(y_0^2 \prod_{i=1}^4 (x_0 - \lambda_{\tau(i)} x_1) + y_1^2 \prod_{i=5}^8 (x_0 - \lambda_{\tau(i)} x_1) \right) = 0.$$

Then $X \cong X_\tau$.

Proof. It is enough to prove the case where τ is a transposition (ij) . Because of the symmetry of B , we may assume that $(ij) = (15)$. Since smooth K3 surfaces have trivial canonical class, it is sufficient to show that X and $X_{(15)}$ are birational.

Over the affine patch $x_1 y_1 \neq 0$, X is defined in $\mathbb{A}_{(x,y,z)}^3$ by

$$z^2 = y \left(y^2 \prod_{i=1}^4 (x - \lambda_i) + \prod_{i=5}^8 (x - \lambda_i) \right),$$

where we set $y = y_0/y_1$ and $x = x_0/x_1$. Consider the birational transformation

$$\begin{aligned} \mathbb{A}_{(\xi,\eta,\zeta)}^3 &\dashrightarrow \mathbb{A}_{(x,y,z)}^3 \\ (\xi, \eta, \zeta) &\mapsto \left(\xi, \frac{\xi - \lambda_5}{\xi - \lambda_1} \eta, \frac{\xi - \lambda_5}{\xi - \lambda_1} \zeta \right). \end{aligned}$$

Under this birational transformation, the pull-back of X satisfies

$$\begin{aligned} \frac{(\xi - \lambda_5)^2}{(\xi - \lambda_1)^2} \zeta^2 &= \frac{\xi - \lambda_5}{\xi - \lambda_1} \eta \left(\frac{(\xi - \lambda_5)^2}{(\xi - \lambda_1)^2} \eta^2 \prod_{i=1}^4 (\xi - \lambda_i) + \prod_{i=5}^8 (\xi - \lambda_i) \right) \\ \implies \zeta^2 &= \eta \left(\frac{\xi - \lambda_5}{\xi - \lambda_1} \eta^2 \prod_{i=1}^4 (\xi - \lambda_i) + \frac{\xi - \lambda_1}{\xi - \lambda_5} \prod_{i=5}^8 (\xi - \lambda_i) \right), \end{aligned}$$

which is the equation for $X_{(15)}$ over the affine patch $x_1 y_1 \neq 0$. Thus they are birational. \square

3. STABLE PAIRS AND KSBA COMPACTIFICATION

In this section we recall the definition of stable pair, their moduli spaces, and the theory of abelian covers. Our main references are [Ale15, AP12, Kol13, Kol18].

3.1. Definition of stable pair.

Definition 3.1. Let X be a variety and let D be a \mathbb{Q} -divisor on X with coefficients in $(0, 1]$. A pair (X, D) is *semi-log canonical* if:

- (1) X is demi-normal (that is, X is S_2 and its codimension 1 points are either regular or ordinary nodes);
- (2) If $\nu : X^\nu \rightarrow X$ is the normalization with conductors $E \subseteq X$ and $E^\nu \subseteq X^\nu$, then the support of E does not contain any irreducible component of D ;
- (3) $K_X + D$ is \mathbb{Q} -Cartier;

- (4) The pair $(X^\nu, E^\nu + \nu_*^{-1}D)$ is log canonical. More precisely, for each connected component Z of X^ν the pair $(Z, (E^\nu + \nu_*^{-1}D)|_Z)$ is log canonical, where $\nu_*^{-1}D$ denotes the strict transform of D . For the definition of log canonical we refer to [Koll13, Definition 2.8].

Definition 3.2. A pair (X, D) is *stable* if the following conditions are satisfied:

- (1) (X, D) is a semi-log canonical pair;
- (2) $K_X + D$ is ample.

Let ϵ be a sufficiently small positive rational number.

Lemma 3.3. Let $([\lambda_1 : 1], \dots, [\lambda_8 : 1]) \in (\mathbb{P}^1)^8$ be a stable point and let $C \subseteq \mathbb{P}^1 \times \mathbb{P}^1$ as in Equation (1). Let $B := C + L_0 + L_1$. Then

- (1) $(\mathbb{P}^1 \times \mathbb{P}^1, \frac{1+\epsilon}{2}B)$ is a stable pair;
- (2) $(K_{\mathbb{P}^1 \times \mathbb{P}^1} + \frac{1+\epsilon}{2}B)^2 = 8\epsilon^2$.

In short, $(\mathbb{P}^1)^8$ parametrizes stable pairs $(\mathbb{P}^1 \times \mathbb{P}^1, \frac{1+\epsilon}{2}B)$.

Proof. If the eight points are distinct, then the divisor B is simple normal crossing, hence $(\mathbb{P}^1 \times \mathbb{P}^1, \frac{1+\epsilon}{2}B)$ is semi-log canonical. If some of the eight points coincide, then the semi-log canonicity of the pair follows by inspecting all cases in Table 1. The ampleness of $K_{\mathbb{P}^1 \times \mathbb{P}^1} + \frac{1+\epsilon}{2}B$ and the equality in (2) follow from $K_{\mathbb{P}^1 \times \mathbb{P}^1} + \frac{1+\epsilon}{2}B \sim 2\epsilon(1, 1)$. \square

3.2. Stable pairs and finite abelian covers. For the reader's convenience, we recall the following well known facts about stable pairs and finite abelian covers. For a reference, see [AP12].

Definition 3.4. A morphism of pairs $f : (X, B_X) \rightarrow (P, B_P)$ is a morphism $f : X \rightarrow P$ mapping $\text{Supp}(B_X)$ to $\text{Supp}(B_P)$. If G is a finite abelian group, then a morphism $\pi : (X, B_X) \rightarrow (P, B_P)$ is called a G -cover if:

- $\pi : X \rightarrow P$ is the quotient morphism for a generically faithful action of G ;
- π is branched along $\text{Supp}(B_P)$ and ramified at $\text{Supp}(B_X)$;
- $K_X + B_X = \pi^*(K_P + B_P)$.

Lemma 3.5 ([AP12, Lemma 2.3]). Let $\pi : (X, B_X) \rightarrow (P, B_P)$ be a G -cover for a finite abelian group G . Then (X, B_X) is stable if and only if (P, B_P) is stable.

Remark 3.6. More precisely, [AP12, Lemma 2.3] guarantees that (X, B_X) is semi-log canonical if and only if (P, B_P) is semi-log canonical. We have that $K_X + B_X$ is ample if and only if $\pi^*(K_P + B_P)$ is because π is a finite covering (see [Laz04, Proposition 1.2.13 and Corollary 1.2.28]).

Example 3.7. Let $P := \mathbb{P}^1 \times \mathbb{P}^1$ and let $B := C + L_0 + L_1$ as in § 2.1. Let X be the double cover of P branched along B and let R be the ramification divisor. Then it is straightforward to check that

$$\pi : (X, \epsilon R) \rightarrow \left(P, \frac{1+\epsilon}{2}B \right)$$

is a $(\mathbb{Z}/2\mathbb{Z})$ -cover. In particular, $(K_X + \epsilon R)^2 = \pi^* \left(K_P + \frac{1+\epsilon}{2}B \right)^2 = 16\epsilon^2$.

3.3. The moduli functor.

Definition 3.8. We fix constants $d, N \in \mathbb{Z}_{>0}$, $C \in \mathbb{Q}_{>0}$, and $\underline{b} = (b_1, \dots, b_n)$ with $b_i \in (0, 1] \cap \mathbb{Q}$ and $Nb_i \in \mathbb{Z}$ for all $i = 1, \dots, n$. The Viehweg's moduli stack $\overline{\mathcal{V}} := \overline{\mathcal{V}}_{d,N,C,\underline{b}}$ is defined as follows. For any reduced \mathbb{C} -scheme S , $\overline{\mathcal{V}}_{d,N,C,\underline{b}}(S)$ is the set of proper flat families $\mathcal{X} \rightarrow S$ together with a divisor $\mathcal{B} = \sum_i b_i \mathcal{B}_i$ satisfying:

- (1) For all $i = 1, \dots, n$, \mathcal{B}_i is a codimension one closed subscheme such that $\mathcal{B}_i \rightarrow S$ is flat at the generic points of $\mathcal{X}_s \cap \text{Supp}(\mathcal{B}_i)$ for every $s \in S$;
- (2) Every geometric fiber (X, B) is a stable pair of dimension d with $(K_X + B)^d = C$;
- (3) There exists an invertible sheaf \mathcal{L} on \mathcal{X} such that for every geometric fiber (X, B) one has $\mathcal{L}|_X \cong \mathcal{O}_X(N(K_X + B))$.

In what follows, we refer to such families $(\mathcal{X}, \mathcal{B}) \rightarrow S$ as *families of stable pairs*.

Remark 3.9. Over characteristic zero, an algebraic stack is Deligne–Mumford if and only if all parametrized objects have finite automorphism groups [Ols16, Remark 8.3.4]. Since every stable pair parametrized by $\overline{\mathcal{V}}$ has a finite automorphism group [Fuj14, Theorem 1.20], $\overline{\mathcal{V}}$ is a Deligne–Mumford stack.

Let \underline{b} be very general [Ale15, §1.5.3]. For a suitably chosen positive integer N depending on d, C , and \underline{b} (which does not need to be specified, see [Ale96, §3.13]), the stack $\overline{\mathcal{V}}$ above is coarsely represented by a projective scheme by [Ale15, Theorem 1.6.1].

We now describe our special case of interest.

Definition 3.10. Let $\overline{\mathcal{V}}$ be the Viehweg's moduli stack for $d = 2$, $C = 16\epsilon^2$ for a small general $\epsilon > 0$, and $\underline{b} = (b_1) = (\epsilon)$. The pairs $(X, \epsilon R)$ where X is an ADE K3 surface in §2.1 and $R = \frac{1}{2}\pi^*B$ is the ramification divisor, are parametrized by $\overline{\mathcal{V}}$. Now consider the family of stable pairs

$$\left(\mathcal{Y} := \mathbf{U} \times \mathbb{P}^1 \times \mathbb{P}^1, \frac{1+\epsilon}{2}(\mathcal{C} + \mathcal{L}_0 + \mathcal{L}_1)|_{\mathbf{U}} \right) \rightarrow \mathbf{U}.$$

(For the definitions of $\mathbf{U}, \mathcal{C}, \mathcal{L}_0, \mathcal{L}_1$ we refer to §2.1.) Let \mathcal{X} be the double cover of \mathcal{Y} branched along $(\mathcal{C} + \mathcal{L}_0 + \mathcal{L}_1)|_{\mathbf{U}}$ and let \mathcal{R} be the ramification divisor. Then we obtain a family of stable pairs $(\mathcal{X}, \epsilon \mathcal{R}) \rightarrow \mathbf{U}$ where the fibers X are ADE K3 surfaces with a purely non-symplectic automorphism of order four and R is the ramification divisor. Thus we obtain a morphism $\mathbf{U} \rightarrow \overline{\mathcal{V}}$ and denote by $\overline{\mathcal{K}}'$ the closure of its image in $\overline{\mathcal{V}}$. Let $\overline{\mathbf{K}}'$ be the coarse moduli space corresponding to $\overline{\mathcal{K}}'$, and denote by $\overline{\mathbf{K}}$ its normalization. Observe that $\overline{\mathbf{K}}$ is compactifying $\mathbf{U}/\text{SL}_2/H$, and we call it the *KSBA compactification* of the moduli space of ADE K3 pairs with purely non-symplectic automorphism of order four and $U(2) \oplus D_4^{\oplus 2}$ lattice polarization. As we already pointed out in the introduction, $\overline{\mathbf{K}}$ is compactifying a finite cover of \mathbf{M} in Definition 2.3, and more precisely a (S_8/H) -cover.

Our ultimate goal is to study the geometry of the compactified moduli space $\overline{\mathbf{K}}$. To do so, we consider the following other projective moduli space.

Definition 3.11. Let $\overline{\mathcal{T}}$ be the Viehweg's moduli stack for $d = 2$, $C = 8\epsilon^2$, $\underline{b} = (b_1) = (\frac{1+\epsilon}{2})$. Consider again the family of stable pairs $(\mathcal{Y}, \frac{1+\epsilon}{2}(\mathcal{C} + \mathcal{L}_0 + \mathcal{L}_1)|_{\mathbf{U}}) \rightarrow \mathbf{U}$. There is an induced

morphism $U \rightarrow \overline{\mathcal{T}}$ and denote by $\overline{\mathcal{J}'}$ the closure of its image in $\overline{\mathcal{T}}$. Let $\overline{\mathcal{J}'}$ be the coarse moduli space corresponding to $\overline{\mathcal{J}'}$, and denote by $\overline{\mathcal{J}}$ its normalization. We have that $\overline{\mathcal{J}}$ is compactifying $U/\mathrm{SL}_2/H$, and hence it is birational to $\overline{\mathcal{K}}$.

In Proposition 3.13 we show that $\overline{\mathcal{K}} \cong \overline{\mathcal{J}}$. (More precisely, we will prove that there is a bijective morphism $\overline{\mathcal{K}'} \rightarrow \overline{\mathcal{J}'}$. This does not imply that $\overline{\mathcal{K}'} \cong \overline{\mathcal{J}'}$, but we can conclude that their normalizations are isomorphic.) Thus it is enough to investigate the geometry of $\overline{\mathcal{J}}$.

Remark 3.12.

- (1) Note that, however, the associated stacks $\overline{\mathcal{K}'}$ and $\overline{\mathcal{J}'}$ are not isomorphic because of their stacky structure. Recall that the former parametrizes stable pairs $(X, \epsilon R)$ where X is an ADE K3 surface with a purely non-symplectic automorphism of order four, and the latter stable pairs $(\mathbb{P}^1 \times \mathbb{P}^1, \frac{1+\epsilon}{2}B)$. Therefore, a general object parametrized by $\overline{\mathcal{K}'}$ has an extra $\mathbb{Z}/2\mathbb{Z}$ -action which comes from its double covering structure.
- (2) The proof of Proposition 3.13 tells us that there is a bijective morphism $\overline{\mathcal{K}'} \rightarrow \overline{\mathcal{J}'}$ between the coarse moduli spaces.

Recall that $U \subseteq (\mathbb{P}^1)^8$ is the locus of eight distinct points. In Definitions 3.10 and 3.11 we introduced two compactifications of $U/\mathrm{SL}_2/H$, which we denoted by $\overline{\mathcal{K}}$ and $\overline{\mathcal{J}}$. The remaining part of this section is devoted to proving the following.

Proposition 3.13. *The compactifications $\overline{\mathcal{K}}$ and $\overline{\mathcal{J}}$ are isomorphic.*

To prove the claim above we need a lemma.

Lemma 3.14. *Let S be a scheme and let $(\mathcal{X}, \mathcal{B}_{\mathcal{X}}) \rightarrow S$ be a family of stable pairs. Let G be a finite abelian group and assume there exists a dense open subset $U \subseteq S$ such that $(\mathcal{X}, \mathcal{B}_{\mathcal{X}})|_U$ has a fiberwise generically faithful action of G which ramifies at $\mathrm{Supp}(\mathcal{B}_{\mathcal{X}}|_U)$. Then the G -action extends to the whole $(\mathcal{X}, \mathcal{B}_{\mathcal{X}})$ giving a fiberwise generically faithful action which ramifies at $\mathrm{Supp}(\mathcal{B}_{\mathcal{X}})$.*

Proof. Let $g \in G$ be arbitrary. We show that the corresponding action $\alpha_g : \mathcal{X}|_U \rightarrow \mathcal{X}|_U$ extends to \mathcal{X} . Consider a resolution of indeterminacies

$$\begin{array}{ccc} & \mathcal{X}' & \\ \swarrow & & \searrow \alpha'_g \\ \mathcal{X} & \text{-----} & \mathcal{X} \\ & \alpha_g & \end{array}$$

Let $\mathcal{B}'_{\mathcal{X}}$ be the strict transform of $\mathcal{B}_{\mathcal{X}}$ under $\mathcal{X}' \rightarrow \mathcal{X}$. Then α'_g induces a morphism α_g^{lc} from the log canonical model of $(\mathcal{X}', \mathcal{B}'_{\mathcal{X}})$ to \mathcal{X} [Kol13, Definition 1.19]. Since the log canonical model of $(\mathcal{X}', \mathcal{B}'_{\mathcal{X}})$ is $(\mathcal{X}, \mathcal{B}_{\mathcal{X}})$, it follows that α_g^{lc} is the desired extension of α_g . \square

Proof of Proposition 3.13. Since $\overline{\mathcal{K}'}$ is a Deligne–Mumford stack (see Remark 3.9), there exists a scheme A and a surjective étale morphism $\alpha : A \rightarrow \overline{\mathcal{K}'}$. Therefore, there exists a

family $\mathcal{A} \rightarrow A$ (we omit the datum of the divisor for simplicity of notation), and every object parametrized by $\overline{\mathcal{K}}'$ appears as a fiber of $\mathcal{A} \rightarrow A$ because α is surjective. In particular, there exists a dense open subset $U \subseteq A$ such that $\mathcal{A}|_U \rightarrow U$ admits a fiberwise $(\mathbb{Z}/2\mathbb{Z})$ -action. So we can apply Lemma 3.14 to extend the $(\mathbb{Z}/2\mathbb{Z})$ -action to the whole family $\mathcal{A} \rightarrow A$.

Now, let S be a normal scheme and let $\mathcal{X} \rightarrow S$ be a family for the stack $\overline{\mathcal{K}}'$. Consider the algebraic space $S' := S \times_{\overline{\mathcal{K}}'} A$, which we may assume is a scheme by replacing it by its atlas, if necessary. We have that S' comes with a family $\mathcal{X}' \rightarrow S'$ obtained by pulling-back $\mathcal{A} \rightarrow A$ along the morphism $S' \rightarrow A$. Observe that $\mathcal{X}' \rightarrow S'$ equals the pull-back of $\mathcal{X} \rightarrow S$ along the morphism $S' \rightarrow S$ because the two compositions $S' \rightarrow A \rightarrow \overline{\mathcal{K}}'$ and $S' \rightarrow S \rightarrow \overline{\mathcal{K}}'$ are equal. We can summarize these considerations in the following commutative diagrams of cartesian squares:

$$\begin{array}{ccccc}
 & & \mathcal{X}' & & \\
 & \swarrow & \downarrow & \searrow & \\
 \mathcal{X} & & S' & & A \\
 \downarrow & \swarrow & \downarrow & \searrow & \downarrow \\
 S & & A & & A \\
 & \searrow & \swarrow & & \\
 & & \overline{\mathcal{K}}' & &
 \end{array}$$

Since $\mathcal{X}' \rightarrow S'$ has a fiberwise $(\mathbb{Z}/2\mathbb{Z})$ -action, also $\mathcal{X} \rightarrow S$ does. The quotient of $\mathcal{X} \rightarrow S$ by this $(\mathbb{Z}/2\mathbb{Z})$ -action gives an object for the stack $\overline{\mathcal{J}}'$. This guarantees the existence of a morphism of stacks $\overline{\mathcal{K}}' \rightarrow \overline{\mathcal{J}}'$ which induces a morphism $f : \overline{\mathcal{K}}' \rightarrow \overline{\mathcal{J}}'$ of the underlying coarse moduli spaces.

The morphism f is surjective because it is the identity on $\mathbf{U}/\mathrm{SL}_2/H$. On the other hand, assume $f(p) = q$ and let $(X, \epsilon R)$ (resp. $(\mathbb{P}^1 \times \mathbb{P}^1, \frac{1+\epsilon}{2}B)$) be the stable pair parametrized by p (resp. q). Then $(X, \epsilon R)$ is the double cover of $(\mathbb{P}^1 \times \mathbb{P}^1, \frac{1+\epsilon}{2}B)$, and since such double cover is unique, we have that f is injective as well. Thus f is bijective.

Now consider the associated morphism $\overline{\mathcal{K}} \rightarrow \overline{\mathcal{J}}$ between normalizations. Since it is a quasi-finite surjective birational morphism between normal varieties, by Zariski's Main Theorem it is an isomorphism. \square

4. PARTIAL DESINGULARIZATION OF GIT QUOTIENTS

4.1. Review on partial desingularization. In this section, we recall the partial desingularization of GIT quotient developed by Kirwan in [Kir85].

Let G be a reductive group and let X be a smooth projective variety equipped with an L -linearized G -action. Suppose that the stable locus $X^s(L)$ is nonempty. If X has strictly semi-stable points, so that $X^s(L) \subsetneq X^{ss}(L)$, then the GIT quotient $X//_L G$ may have non-finite quotient singularities. Such a singularity can be partially G -equivariantly resolved

by using Kirwan's partial desingularization. The outcome of the partial desingularization process is a new algebraic variety X' equipped with an L' -linearized G -action such that:

- (1) There is a dominant projective birational map $X'//_{L'}G \rightarrow X//_L G$;
- (2) $X'//_{L'}G$ has finite quotient singularities only.

Assume that $X^{ss}(L) \neq X^s(L)$. Let Y be the closed G -invariant subvariety of $X^{ss}(L)$ with a maximal dimensional stabilizer group. Let \tilde{X} be the blow-up of $X^{ss}(L)$ along Y and let $\pi : \tilde{X} \rightarrow X^{ss}(L)$ be the blow-up morphism. Let E be the exceptional divisor. \tilde{X} is a smooth quasi-projective variety since Y is smooth. For a small $\epsilon > 0$, $L_\epsilon := \pi^*L \otimes \mathcal{O}(-\epsilon E)$ is an ample line bundle on \tilde{X} and the G -action is naturally extended to \tilde{X} and to L_ϵ .

The stable and semi-stable loci of X and \tilde{X} are related as follows:

$$\pi^{-1}X^s(L) \subseteq \tilde{X}^s(L_\epsilon) \subseteq \tilde{X}^{ss}(L_\epsilon) \subseteq \pi^{-1}X^{ss}(L).$$

Thus there is a natural G -equivariant morphism $\tilde{X}^{ss}(L_\epsilon) \rightarrow X^{ss}(L)$ which induces a morphism $\bar{\pi} : \tilde{X}//_{L_\epsilon}G \rightarrow X//_L G$ between quotients. On \tilde{X} , a point $x \in \tilde{X}$ is unstable if the orbit of $\pi(x)$ is not closed in $X^{ss}(L)$ [Kir85, Lemma 6.6]. Furthermore, the maximal dimension of the stabilizer group strictly decreases. After replacing X by \tilde{X}^{ss} and L by L_ϵ , we can continue this process and it terminates. The result of this procedure is X' and L' .

4.2. Partial resolution of the parameter space of eight points on the projective line. Let $\mathbf{X}_0 := (\mathbb{P}^1)^8$ equipped with a diagonal SL_2 -action. Let $L = \mathcal{O}(1, \dots, 1)$ be the symmetric linearization. We explicitly describe the partial desingularization of $\mathbf{X}_0//_L \mathrm{SL}_2$. For the detail, see [Kir85, §9].

Recall that for a point configuration $\mathbf{p} \in \mathbf{X}_0$, $\mathbf{p} \in \mathbf{X}_0^{ss}$ if and only if at most four points collide and $\mathbf{p} \in \mathbf{X}_0^s$ if and only if at most three points collide.

If $\binom{[8]}{4}$ denotes the set of 4-element subsets of $[8] := \{1, \dots, 8\}$, for any $I \in \binom{[8]}{4}$, let $\Delta_I := \{\mathbf{p} = (p_i) \in \mathbf{X}_0^{ss} \mid p_i = p_j \text{ for all } i, j \in I\}$. It is a five dimensional smooth subvariety of \mathbf{X}_0^{ss} . The set of strictly semi-stable points is

$$\mathbf{X}_0^{ss} \setminus \mathbf{X}_0^s = \bigcup_{I \in \binom{[8]}{4}} \Delta_I.$$

For $I \in \binom{[8]}{4}$, let $\Delta_{I,I^c} := \Delta_I \cap \Delta_{I^c}$. Note that $\Delta_{I,I^c} \cong (\mathbb{P}^1)^2 \setminus \mathbb{P}^1$, where the last \mathbb{P}^1 is the diagonal. For any $J \neq I, I^c$, we have that $\Delta_{I,I^c} \cap \Delta_{J,J^c} = \emptyset$. So if we let $\Delta_{4,4} = \bigcup_{I \in \binom{[8]}{4}} \Delta_{I,I^c}$, then $\Delta_{4,4}$ is a disjoint union of 35 copies of two dimensional smooth closed subvarieties of \mathbf{X}_0^{ss} . $\Delta_{4,4}$ is precisely the set of semi-stable points with positive dimensional stabilizer, which is isomorphic to \mathbb{C}^* .

Let \mathbf{X}_1 be the blow-up of \mathbf{X}_0^{ss} along $\Delta_{4,4}$. Let E_{I,I^c} be the exceptional divisor over Δ_{I,I^c} . For each Δ_I , let $\tilde{\Delta}_I$ be the proper transform of Δ_I . Then \mathbf{X}_1 is a smooth variety equipped with a linearized SL_2 -action. Let $\rho : \mathbf{X}_1 \rightarrow \mathbf{X}_0^{ss}$ be the blow-up morphism. By the recipe of the partial desingularization, it is straightforward to check the following:

- (1) If $\rho(\mathbf{p}) \in \mathbf{X}_0^s$, then \mathbf{p} is stable;
- (2) If $\mathbf{p} \in E_{I,I^c} \setminus (\tilde{\Delta}_I \cup \tilde{\Delta}_{I^c})$, then \mathbf{p} is stable;

(3) If $\mathbf{p} \in \tilde{\Delta}_I$, then \mathbf{p} is unstable.

In particular, $\mathbf{X}_1^{ss} = \mathbf{X}_1^s$. One may check that for every $\mathbf{p} \in \mathbf{X}_1^s$, the stabilizer group is isomorphic to $\{\pm 1\}$. Therefore, the GIT quotient $\mathbf{X}_1^s/\mathrm{SL}_2$ is a smooth variety.

Definition 4.1. Let \mathbf{P} be the partial desingularization of $(\mathbb{P}^1)^8/\mathrm{SL}_2$ with symmetric linearization, that is, $\mathbf{P} := \mathbf{X}_1^s/\mathrm{SL}_2$.

Note that both \mathbf{X}_0 and \mathbf{X}_1 have \mathbf{U} as an open subset. So there is an open embedding $\mathbf{U}/\mathrm{SL}_2 \hookrightarrow \mathbf{P} = \mathbf{X}_1^s/\mathrm{SL}_2$.

Remark 4.2.

- (1) By [KM11, Theorem 1.1], \mathbf{P} is isomorphic to $\overline{\mathbf{M}}_{0,(\frac{1}{4}+\epsilon)^8}$, the moduli space of stable rational curves with eight marked points of weight $\frac{1}{4} + \epsilon$ (see [Has03]).
- (2) An irreducible component \overline{E}_{I,I^c} of the exceptional divisor of $\tilde{\pi} : \mathbf{P} \rightarrow (\mathbb{P}^1)^8/\mathrm{SL}_2$ is isomorphic to $\mathbb{P}^2 \times \mathbb{P}^2$. Indeed, because SL_2 acts on Δ_{I,I^c} transitively with stabilizer group \mathbb{C}^* , the exceptional set is isomorphic to $\pi^{-1}(\Delta_{I,I^c})/\mathrm{SL}_2 \cong \mathbb{P}^5/\mathbb{C}^*$. Here $\mathbb{P}^5 \cong \mathbb{P}(\mathcal{N}_{\Delta_{I,I^c}/\mathbf{X}_0^{ss}}|_x)$ for some $x \in \Delta_{I,I^c}$. One can check that the weight decomposition of $\mathcal{N}_{\Delta_{I,I^c}/\mathbf{X}_0^{ss}}|_x$ is $(2)^3 \oplus (-2)^3$. Thus the GIT quotient $\mathbb{P}^5/\mathbb{C}^*$ is isomorphic to $\mathbb{P}^2 \times \mathbb{P}^2$.
Alternatively, we may adopt the moduli theoretic meaning (again, [KM11, Theorem 1.1]) to obtain the same result: $\overline{E}_{I,I^c} \cong \overline{\mathbf{M}}_{0,(1,(\frac{1}{4}+\epsilon)^4)} \times \overline{\mathbf{M}}_{0,(1,(\frac{1}{4}+\epsilon)^4)} \cong \mathbb{P}^2 \times \mathbb{P}^2$.

5. EXPLICIT CALCULATIONS OF STABLE REPLACEMENTS

The first step toward the proof of Theorem 1.1 is the construction of an extension $(\tilde{\mathcal{Y}}, \frac{1+\epsilon}{2}\tilde{\mathcal{B}}) \rightarrow \mathbf{X}_1^s$ of the family of pairs $(\mathcal{Y}, \frac{1+\epsilon}{2}\mathcal{B}) \rightarrow \mathbf{U}$, where recall $\mathcal{Y} = \mathbf{U} \times (\mathbb{P}^1 \times \mathbb{P}^1)$ and we set $\mathcal{B} := (\mathcal{C} + \mathcal{L}_0 + \mathcal{L}_1)|_{\mathbf{U}}$ (see Definition 3.10). This would induce a functorial morphism $\mathbf{X}_1^s \rightarrow \overline{\mathbf{K}}$. The above extension is obtained by modifying $(\mathcal{Y}_1, \frac{1+\epsilon}{2}\mathcal{B}_1) \rightarrow \mathbf{X}_1^s$, where $\mathcal{Y}_1 := \mathbf{X}_1^s \times (\mathbb{P}^1 \times \mathbb{P}^1)$ and \mathcal{B}_1 is obtained by pulling-back $\mathcal{C} + \mathcal{L}_0 + \mathcal{L}_1$ under the appropriate morphism. We postpone the global analysis of this modification to §6. In this section, we describe how the modification of the family goes with concrete examples of one-parameter degenerations.

Throughout the whole section we adopt the following notation. For a stable pair (X, B) , if $\nu : \mathrm{II}_i X_i \rightarrow X$ is the normalization map, then we denote by D_{X_i} the conductor divisor on X_i . We use the letter P to denote $\mathbb{P}^1 \times \mathbb{P}^1$, and Δ to denote the germ of a curve whose uniformizing parameter is t . On $P \times \Delta$, we let $\mathcal{L}_0, \mathcal{L}_1$ denote the divisors $V(y_0), V(y_1)$ respectively. For simplicity of notation, we denote the fibers $(\mathcal{L}_0)_0, (\mathcal{L}_1)_0$ over $t = 0$ by L_0, L_1 respectively. \mathcal{B} denotes the divisor $\mathcal{C} + \mathcal{L}_0 + \mathcal{L}_1$, where \mathcal{C} is appropriately defined in each one of the examples that follow, and it depends on the choice of eight general complex numbers $\vec{\lambda} := (\lambda_1, \dots, \lambda_8) \in \mathbb{C}^8 \subseteq (\mathbb{P}^1)^8$.

5.1. Semi-stable points of type a.

Example 5.1. Consider the one-parameter family of divisors \mathcal{C} on $P \times \Delta$ given by

$$\begin{aligned} & y_0^2(x_0 - t\lambda_1 x_1)(x_0 - t\lambda_2 x_1)(x_0 - \lambda_3 x_1)(x_0 - \lambda_4 x_1) \\ & + y_1^2(x_0 - t\lambda_5 x_1)(x_0 - t\lambda_6 x_1)(x_0 - \lambda_7 x_1)(x_0 - \lambda_8 x_1) = 0, \end{aligned}$$

which is associated to $\vec{\lambda}_t = (t\lambda_1, t\lambda_2, \lambda_3, \lambda_4, t\lambda_5, t\lambda_6, \lambda_7, \lambda_8) \in (\mathbb{P}^1)^8$. We have a family of pairs $(P \times \Delta, \frac{1+\epsilon}{2}\mathcal{B}) \rightarrow \Delta$. When $t = 0$, $\vec{\lambda}_0$ is strictly semi-stable. For a general $t \neq 0$, the fiber $(P, \frac{1+\epsilon}{2}\mathcal{B}_t)$ is a stable pair, thus its double cover branched along \mathcal{B}_t is also a stable ADE K3 pair. On the other hand, note that $(P, \frac{1+\epsilon}{2}\mathcal{B}_0)$ is not a stable pair because $\mathcal{B}_0 = 2L + C' + L_0 + L_1$ has a double line where L is the line $V(x_0)$. Here C' is the $(2, 2)$ divisor given by

$$y_0^2(x_0 - \lambda_3x_1)(x_0 - \lambda_4x_1) + y_1^2(x_0 - \lambda_7x_1)(x_0 - \lambda_8x_1) = 0.$$

The intersection $L \cap C'$ consists of the two distinct solutions of $y_0^2\lambda_3\lambda_4 + y_1^2\lambda_7\lambda_8 = 0$ (recall we assumed that the λ_i are general).

We blow-up the double locus $x_0 = t = 0$. Let E be the exceptional divisor, which is isomorphic to P . Let C'' denote the restriction to E of the strict transform of \mathcal{C} . On the affine patch $x_1y_0 \neq 0$, the equation of C'' is given by the smallest degree terms with respect to x_0 and t in the equation for \mathcal{C} . Note that y_1 is regarded as a constant during this computation. Thus, after homogenizing back, C'' is the $(2, 2)$ curve on E given by

$$y_0^2(x_0 - t\lambda_1)(x_0 - t\lambda_2)\lambda_3\lambda_4 + y_1^2(x_0 - t\lambda_5)(x_0 - t\lambda_6)\lambda_7\lambda_8 = 0,$$

where $([x_0 : t], [y_0 : y_1])$ are the coordinates of E . The restriction of the strict transform of $\mathcal{L}_0 + \mathcal{L}_1$ to E also consists of the two lines $y_0 = 0$ and $y_1 = 0$. The resulting limit is described in Figure 1. The central fiber is semi-log canonical, and one may check that

$$K_P + D_P + \frac{1+\epsilon}{2}(L_0 + L_1 + C') \sim (-2, -2) + (1, 0) + \frac{1+\epsilon}{2}(2, 4) = \epsilon(1, 2),$$

which is ample. By symmetry, this is enough to show that the limit is stable.

To conclude this example, observe that $\vec{\lambda}_0$ is discarded in the partial desingularization process. However, the study we carried out is preliminary to the next Example 5.2.




FIGURE 1. Stable limit pair in Example 5.1.

Example 5.2. Consider the one-parameter family of divisors \mathcal{C} on $P \times \Delta$ given by

$$\begin{aligned} & y_0^2(x_0 - t\lambda_1x_1)(x_0 - t\lambda_2x_1)(tx_0 - \lambda_3x_1)(tx_0 - \lambda_4x_1) \\ & + y_1^2(x_0 - t\lambda_5x_1)(x_0 - t\lambda_6x_1)(tx_0 - \lambda_7x_1)(tx_0 - \lambda_8x_1) = 0, \end{aligned}$$

whose special fiber \mathcal{C}_0 is

$$(\lambda_3\lambda_4y_0^2 + \lambda_7\lambda_8y_1^2)x_0^2x_1^2 = 0.$$

In this case, $\vec{\lambda}_0$ is a strictly semi-stable point with a closed orbit. Thus the normal direction $\vec{\lambda}_t$ corresponds to a stable point on the exceptional divisor of the partial desingularization. When $t = 0$, \mathcal{B}_0 is the union of four distinct horizontal lines $y_0 = 0$, $y_1 = 0$, $y_0 = \pm\sqrt{-\lambda_7\lambda_8/(\lambda_3\lambda_4)}y_1$, and two non-reduced vertical lines $x_0 = 0$, $x_1 = 0$ with multiplicity two.

As we did in Example 5.1, let E_0 (resp. E_1) be the exceptional divisor of the blow-up along $x_0 = t = 0$ (resp. $x_1 = t = 0$). Observe that these two exceptional divisors are isomorphic to P . Let $\tilde{\mathcal{C}}$ be the strict transform of \mathcal{C} . On E_0 , which has coordinates $([x_0 : t], [y_0 : y_1])$, the restriction of $\tilde{\mathcal{C}}$ has equation

$$C_0 : y_0^2(x_0 - t\lambda_1)(x_0 - t\lambda_2)\lambda_3\lambda_4 + y_1^2(x_0 - t\lambda_5)(x_0 - t\lambda_6)\lambda_7\lambda_8 = 0.$$

Similarly, on E_1 , which has coordinates $([t : x_1], [y_0 : y_1])$, the restriction of $\tilde{\mathcal{C}}$ has equation

$$C_1 : y_0^2(t - \lambda_3x_1)(t - \lambda_4x_1) + y_1^2(t - \lambda_7x_1)(t - \lambda_8x_1) = 0.$$

The limit pair is pictured in Figure 2. On the central component,

$$K_P + D_P + \frac{1+\epsilon}{2}B_P = (-2, -2) + (2, 0) + \frac{1+\epsilon}{2}(0, 4) = \epsilon(0, 2),$$

which is not ample. Therefore, we need to contract P in the central fiber horizontally, obtaining a surface with two components E_0 and E_1 . The resulting stable pair is the same to that in Figure 1.




FIGURE 2. Semi-log canonical limit pair in Example 5.2. The stable limit is obtained after contracting horizontally the middle component.

Remark 5.3. Let us describe the double cover $\pi : X = X_0 \cup X_1 \rightarrow E_0 \cup E_1$ of the degeneration in Example 5.2. Consider the double cover $\pi_0 : X_0 \rightarrow E_0$, whose branch divisor is of class $(2, 4)$ and has four isolated singularities of type A_1 . We have that

$$K_{X_0} = \pi_0^*(K_{E_0} + (1, 2)) = \pi_0^*(-1, 0),$$

which implies that $-K_{X_0}$ is nef and $K_{X_0}^2 = 0$. From $\pi_{0*}\mathcal{O}_{X_0} \cong \mathcal{O}_{E_0} \oplus \mathcal{O}_{E_0}(-1, -2)$ we can argue that $H^1(\mathcal{O}_{X_0}) = 0$. Moreover, by the projection formula, we have that

$$\pi_{0*}\mathcal{O}_{X_0}(2K_{X_0}) = \pi_{0*}\pi_0^*\mathcal{O}_{E_0}(-2, 0) \cong \mathcal{O}_{E_0}(-2, 0) \oplus \mathcal{O}_{E_0}(-3, -2),$$

so $H^0(\mathcal{O}_{X_0}(2K_{X_0})) = 0$. Therefore, by Castelnuovo's rationality criterion, X_0 is rational. Note that X_0 has an elliptic fibration induced by the projection on E_0 given by $([x_0 : t], [y_0 :$

$y_1]) \mapsto [x_0 : t]$. The same calculations work for X_1 . Observe that the gluing locus $X_0 \cap X_1$ is a genus one curve.

In Example 5.2 we started with eight general points. Note that the same construction is valid for all the degenerate point configurations of interest, as we clarify in the following lemma.

Lemma 5.4. *Consider a semi-stable choice of $\lambda_1, \dots, \lambda_8 \in \mathbb{C}^8 \subseteq (\mathbb{P}^1)^8$ such that $\lambda_3\lambda_4\lambda_7\lambda_8 \neq 0$, at least one of $\lambda_1, \lambda_2, \lambda_5, \lambda_6$ is different from the others, and at least one of $\lambda_3, \lambda_4, \lambda_7, \lambda_8$ is different from the others. Then the irreducible components $(E_i, \frac{1+\epsilon}{2}(C_i + L_0 + L_1) + D_{E_i})$ of the limit pair are stable, $i = 0, 1$.*

Remark 5.5. The reason why we exclude the cases $\lambda_3\lambda_4\lambda_7\lambda_8 = 0$ and $\lambda_1 = \lambda_2 = \lambda_5 = \lambda_6$ is because, in the former case, $\vec{\lambda}_0$ is an unstable point configuration. In the latter case, $\vec{\lambda}_t$ is a curve along the strictly semi-stable locus in $(\mathbb{P}^1)^8$, and this case can be discarded in the partial desingularization process, as $\vec{\lambda}_t$ with $t \neq 0$ is unstable on the blow-up of $(\mathbb{P}^1)^8$. For this same reason, we also exclude $\lambda_3 = \lambda_4 = \lambda_7 = \lambda_8$.

Proof of Lemma 5.4. Let us prove the claim for the pair corresponding to $i = 0$, since the other one is analogous. We only have to prove that $(E_0, \frac{1+\epsilon}{2}(C_0 + L_0 + L_1) + D_{E_0})$ is log canonical. Up to symmetries, the significant cases are $\lambda_1, \lambda_2, \lambda_5, \lambda_6$ distinct (where C_0 is smooth), $\lambda_1 = \lambda_2$, $\lambda_1 = \lambda_5$, and $\lambda_1 = \lambda_2 = \lambda_5$. In each case we conclude that $C_0 + L_0 + L_1$ produces planar singularities that appear on Table 1 only, and on D_{E_0} they do not have any double point (we omit these simple computations for brevity). So the pair $(E_0, \frac{1+\epsilon}{2}(C_0 + L_0 + L_1) + D_E)$ is log canonical. \square

5.2. Semi-stable points of type b.

Example 5.6. Consider the family of pairs $(P \times \Delta, \frac{1+\epsilon}{2}\mathcal{B}) \rightarrow \Delta$ where \mathcal{C} on $P \times \Delta$ is given by

$$y_0^2(x_0 - t\lambda_1x_1)(x_0 - t\lambda_2x_1)(x_0 - t\lambda_3x_1)(x_0 - \lambda_4x_1) \\ + y_1^2(x_0 - t\lambda_5x_1)(x_0 - \lambda_6x_1)(x_0 - \lambda_7x_1)(x_0 - \lambda_8x_1) = 0.$$

For $t \neq 0$ the pair is stable. For $t = 0$, the only non-log canonical singularity is at $x_0 = y_1 = 0$, and it is locally analytically four distinct concurrent lines with weight $\frac{1+\epsilon}{2}$. So we restrict to the affine patch $x_1y_0 \neq 0$. Take the blow-up of $P \times \Delta$ at $t = x_0 = y_1 = 0$ and let $E \cong \mathbb{P}^2$ be the exceptional divisor of the blow-up. Denote by C_E the restriction to E of the strict transform of \mathcal{C} . On E , the divisor C_E has equation

$$(x_0 - t\lambda_1)(x_0 - t\lambda_2)(x_0 - t\lambda_3)\lambda_4 + y_1^2(x_0 - t\lambda_5)\lambda_6\lambda_7\lambda_8 = 0,$$

which is a smooth cubic curve if $\vec{\lambda}$ is general, which we assumed. C does not have an irreducible component $V(y_1)$. If h denotes the class of a line on E , then

$$K_E + D_E + \frac{1+\epsilon}{2}B_E = \left(-3 + 1 + \frac{1+\epsilon}{2}4\right)h = 2\epsilon h,$$

which is ample. See the left hand side of Figure 3 for the limit pair.

Let $\pi : P' \rightarrow P$ be the blow-up of the central fiber, where the exceptional divisor is the conductor $D_{P'}$ and $B_{P'}$ is the strict transform of B_0 . We have $K_{P'} = \pi^*(-2, -2) + D_{P'}$ and $B_{P'} = \pi^*(4, 4) - 4D_{P'}$. We can see that the divisor $K_{P'} + D_{P'} + \frac{1+\epsilon}{2}B_{P'}$ is not ample on P' . Consider the proper transform of the line $V(y_1)$, whose class is $\pi^*(0, 1) - D_{P'}$. It is simple to check that the intersection number $(\pi^*(0, 1) - D_{P'}) \cdot (K_{P'} + D_{P'} + \frac{1+\epsilon}{2}B_{P'})$ equals zero. The same computation yields the intersection with the proper transform of $V(x_0)$ is also zero. After contracting these two lines, we obtain a stable limit with two irreducible components isomorphic to \mathbb{P}^2 , as in the right hand side of Figure 3.




FIGURE 3. Stable limit pair in Example 5.6.

Example 5.7. Consider the family of pairs $(P \times \Delta, \frac{1+\epsilon}{2}\mathcal{B}) \rightarrow \Delta$, where \mathcal{C} is given by

$$y_0^2(x_0 - t\lambda_1x_1)(x_0 - t\lambda_2x_1)(x_0 - t\lambda_3x_1)(tx_0 - \lambda_4x_1) \\ + y_1^2(x_0 - t\lambda_5x_1)(tx_0 - \lambda_6x_1)(tx_0 - \lambda_7x_1)(tx_0 - \lambda_8x_1) = 0.$$

Observe that the limit 8-point configuration is a strictly semi-stable point of type b. On the central fiber, there are two non-log canonical singularities at $x_0 = y_1 = 0$ and at $x_1 = y_0 = 0$. By taking two blow-ups along those two points, which introduces the exceptional divisors E_0 and E_1 respectively, we have a configuration of three surfaces as in Figure 4. On E_0 , the restriction of the strict transform of \mathcal{C} is given by

$$C_0 : (x_0 - t\lambda_1)(x_0 - t\lambda_2)(x_0 - t\lambda_3)\lambda_4 + y_1^2(x_0 - t\lambda_5)\lambda_6\lambda_7\lambda_8 = 0.$$

On E_1 , the restriction of the strict transform of \mathcal{C} is given by

$$C_1 : y_0^2(t - \lambda_4x_1) + (t - \lambda_6x_1)(t - \lambda_7x_1)(t - \lambda_8x_1) = 0.$$

The central component P'' is a ruled surface and

$$K_{P''} + D_{P''} + \frac{1+\epsilon}{2}B_{P''} = (\pi^*(-2, -2) + D_{E_0} + D_{E_1}) \\ + D_{E_0} + D_{E_1} + \frac{1+\epsilon}{2}(\pi^*(4, 4) - 4D_{E_0} - 4D_{E_1}) \\ = \epsilon(\pi^*(2, 2) - 2D_{E_0} - 2D_{E_1})$$

is not ample. P'' is contracted diagonally, thus the resulting stable limit is the same as the picture on the right hand side in Figure 3.




FIGURE 4. Semi-log canonical limit pair in Example 5.7. The stable limit is obtained after contracting diagonally the middle component.

Remark 5.8. For $i = 0, 1$, the double cover $\pi_i : X_i \rightarrow E_i$ of an irreducible component $E_i \cong \mathbb{P}^2$ is branched along a reducible quartic curve B_i with three nodal singularities. Since $K_{X_i} = \pi_i^* K_{E_i} + \frac{1}{2} \pi_i^* B_i = -\pi_i^* h$ where h is the class of a line, we have that K_{X_i} is anti-ample. Since X_i has only three A_1 singularities, the minimal resolution of X_i is a weak del Pezzo surface of degree 2. The gluing locus $X_0 \cap X_1$ is a genus one curve.

The same construction is valid for all the degenerate point configurations of interest in view of the partial desingularization. The reasoning behind the restrictions on the λ_i in the next lemma is analogous to what we already explained in Remark 5.5.

Lemma 5.9. Consider a semi-stable choice of $\lambda_1, \dots, \lambda_8 \in \mathbb{C}^8 \subseteq (\mathbb{P}^1)^8$ such that $\lambda_4 \lambda_6 \lambda_7 \lambda_8 \neq 0$, at least one of $\lambda_1, \lambda_2, \lambda_3, \lambda_5$ is different from the others, and at least one of $\lambda_4, \lambda_6, \lambda_7, \lambda_8$ is different from the others. Then the irreducible components $(E_0, \frac{1+\epsilon}{2}(C_0 + L_1) + D_{E_0}), (E_1, \frac{1+\epsilon}{2}(C_1 + L_0) + D_{E_1})$ of the limit pair are stable.

Proof. By symmetry, it is enough to check the log canonicity of $(E_0, \frac{1+\epsilon}{2}(C_0 + L_1) + D_{E_0})$. By checking degenerate point configurations, under the assumption that one of $\lambda_1, \lambda_2, \lambda_3, \lambda_5$ is distinct from the others, one may conclude that $(C_0 + L_1)$ has only singularities that appear on Table 1, and along $D_E = V(t)$ it has no singularities at all. So the pair $(E_0, \frac{1+\epsilon}{2}(C_0 + L_1) + D_{E_0})$ is log canonical. \square

5.3. Semi-stable points of type c.

Example 5.10. Consider the one-parameter family of divisors \mathcal{C} on $\mathcal{P} := P \times \Delta$ given by

$$y_0^2 \prod_{i=1}^4 (x_0 - t\lambda_i x_1) + y_1^2 \prod_{i=5}^8 (x_0 - \lambda_i x_1) = 0.$$

The divisor \mathcal{B}_0 has five multiple points, but only $([0 : 1], [1 : 0])$ is not a simple double point. In the affine patch $x_1 y_0 \neq 0$, locally at $(0, 0)$, the singularity of \mathcal{B}_0 is isomorphic to $y_1(x_0^4 + y_1^2) = 0$, which is not log canonical for the weight $(1 + \epsilon)/2$. To find the stable replacement, let us restrict to the affine patch $x_1 y_0 \neq 0$, so that the equation for \mathcal{B} becomes

$$y_1 \left(\prod_{i=1}^4 (x_0 - t\lambda_i) + y_1^2 \prod_{i=5}^8 (x_0 - \lambda_i) \right) = 0.$$

We perform the following birational modifications (see Figure 5).

- (1) Let $\mathcal{P}' \rightarrow \mathcal{P}$ be the weighted blow-up at $(t, x_0, y_1) = (0, 0, 0)$ with weights $(1, 1, 2)$, which is the blow-up of the ideal (t^2, tx_0, x_0^2, y_1) . The exceptional divisor E_1 is isomorphic to the weighted projective plane $\mathbb{P}(1, 1, 2)$, which is isomorphic to \mathbb{F}_2^0 , a cone over a smooth conic. This weighted blow-up introduces an A_1 singularity on each irreducible component of \mathcal{P}'_0 . Let \mathcal{B}' be the strict transform of the divisor \mathcal{B} . Observe that the equation for \mathcal{B}'_{E_1} , which is the smallest degree part with respect to x_0, y_1, t of the equation for \mathcal{B} , is given by

$$y_1 \left(\prod_{i=1}^4 (x_0 - t\lambda_i) + y_1^2 \prod_{i=5}^8 \lambda_i \right) = 0.$$

It is the union of a 2-section and a section of \mathbb{F}_2^0 .

- (2) The strict transform of the line $L_1 = V(y_1)$ on P intersects $K_{\mathcal{P}'_0} + \frac{1+\epsilon}{2}\mathcal{B}'_0$ negatively. We flip this curve by first blowing it up, introducing an exceptional divisor E_2 isomorphic to \mathbb{F}_1 .
- (3) Blow-up the strict transform of L_1 on E_2 introducing the exceptional divisor E_3 isomorphic to \mathbb{F}_0 .
- (4) Contract E_3 along the ruling intersecting E_2 in the exceptional divisor. The strict transform of E_2 is now isomorphic to \mathbb{P}^2 .
- (5) Contract the strict transform of E_2 to a point. This introduces an A_1 singularity on the strict transform of P and on the strict transform of E_1 .




FIGURE 5. Stable replacement in Example 5.10.

The central fiber is log canonical (it was already log canonical at step (1) – the modification that followed did not make the singularities of the pair worse), so we only need to check that the numerical condition is satisfied. Denote by Y_1, Y_2 the two irreducible components of the central fiber, where Y_1 is the strict transform of P . Let B_{Y_i} be the restriction of the strict transform of \mathcal{B} to Y_i . We want to show that $K_{Y_i} + D_{Y_i} + \frac{1+\epsilon}{2}B_{Y_i}$ is ample for $i = 1, 2$.

- (1) On Y_1 , let M_0, M_1 be the two vertical boundary lines, where $M_0^2 = -\frac{1}{2}$ and $M_1^2 = \frac{1}{2}$. Note that L_0 is the horizontal boundary line disjoint from Y_2 . It is a straightforward exercise to compute the following intersection numbers (observe that L_0 is contained in the support of B_{Y_1}):

\cdot	M_0	M_1	L_0	D_{Y_1}	B_{Y_1}
M_0	$-1/2$	0	1	$1/2$	1
M_1		$1/2$	1	$1/2$	3
L_0			0	0	4
D_{Y_1}				0	2

Moreover, $K_{Y_1} = -M_0 - M_1 - L_0 - D_{Y_1}$. One may check that $K_{Y_1} + D_{Y_1} + \frac{1+\epsilon}{2}B_{Y_1}$ intersects positively with M_0, M_1, L_0 , and D_{Y_1} . Since Y_1 is a toric variety, this implies $K_{Y_1} + D_{Y_1} + \frac{1+\epsilon}{2}B_{Y_1}$ is ample.

- (2) On Y_2 , let N_0, N_1 be the two vertical boundary lines, where $N_0^2 = -\frac{1}{2}$ and $N_1^2 = \frac{1}{2}$. Denote by T the top boundary line. We have the following intersection numbers:

\cdot	N_0	N_1	T	D_{Y_2}	B_{Y_2}
N_0	$-1/2$	0	1	$1/2$	1
N_1		$1/2$	1	$1/2$	3
T			0	0	4
D_{Y_2}				0	2

Then $K_{Y_2} = -N_0 - N_1 - T - D_{Y_2}$ and $K_{Y_2} + D_{Y_2} + \frac{1+\epsilon}{2}B_{Y_2}$ is ample because it intersects positively with N_0, N_1, T, D_{Y_2} .

Note that two intersection matrices for Y_1 and Y_2 are same. Indeed, they are isomorphic toric surfaces. But the non-toric divisors B_{Y_1} and B_{Y_2} are in general different.

Example 5.11. Finally, consider the one-parameter family of divisors \mathcal{C} on $\mathcal{P} := P \times \Delta$ given by

$$y_0^2 \prod_{i=1}^4 (x_0 - t\lambda_i x_1) + y_1^2 \prod_{i=5}^8 (tx_0 - \lambda_i x_1) = 0.$$

We have that $(\mathcal{P}_0, \frac{1+\epsilon}{2}\mathcal{B}_0)$ is not log canonical at $([0 : 1], [1 : 0]), ([1 : 0], [0 : 1])$ where the divisor is locally isomorphic to $y(x^4 + y^2)$. To obtain the stable replacement we may repeat the procedure in Example 5.10 for both singularities. Let F_0 be the exceptional divisor of the weighted blow-up at $([0 : 1], [1 : 0])$. Then the restriction to F_0 of the strict transform of \mathcal{C} has equation

$$C_0 : \prod_{i=1}^4 (x_0 - t\lambda_i) + y_1^2 \prod_{i=5}^8 \lambda_i = 0.$$

Similarly, if F_1 be the exceptional divisor of the weighted blow-up at $([1 : 0], [0 : 1])$, then the restriction to F_1 of the strict transform of \mathcal{C} has equation

$$C_1 : y_0^2 + \prod_{i=5}^8 (t - \lambda_i x_1) = 0.$$

Continuing with the flip described in Example 5.10 (which we perform once for each of the two singularities), the semi-stable replacement has three irreducible components: let Y_1 be the strict transform of P and let Y_2, Y_3 be the strict transforms of the exceptional divisors of the two exceptional divisors F_0, F_1 respectively (see left picture in Figure 6). Observe that Y_1 has four A_1 singularities and Y_2 and Y_3 have two A_1 singularities each. Let B_{Y_i} be the restriction to Y_i of the strict transform of \mathcal{B} .




FIGURE 6. On the left, the semi-log canonical limit pair in Example 5.11, and on the right, the corresponding stable limit obtained after contracting the middle component.

On Y_1 , let M_i be the strict transform of the line $x_i = 0$, $i = 0, 1$. The conductor divisor D_{Y_1} consists of the two irreducible components $D_{12} := Y_1 \cap Y_2$ and $D_{13} := Y_1 \cap Y_3$. We have the following intersection numbers:

\cdot	M_0	M_1	D_{12}	D_{13}	B_{Y_1}
M_0	0	0	1/2	1/2	0
M_1		0	1/2	1/2	0
D_{12}			0	0	2
D_{13}				0	2

One can compute that $K_{Y_1} = -2M_0 - 2D_{13}$. Therefore, $K_{Y_1} + D_{Y_1} + \frac{1+\epsilon}{2}B_{Y_1}$ intersects M_0 and M_1 trivially and Y_1 can be contracted vertically to obtain a stable pair. The resulting pair is the same as the right picture in Figure 6.

Remark 5.12. We compute the double cover of the degeneration in Figure 6 after contracting the middle component Y_1 . Let $\pi_2 : X_2 \rightarrow Y_2$ be the double cover (analogous considerations hold for Y_3). Using that $B_{Y_2} = 4N_0 + 3T$ and $K_{Y_2} = -(N_0 + N_1 + T + D_{Y_2}) = -2N_0 - 2T$ (notice that Y_2 has Picard number 2), and following the same strategy as in Remark 5.3,

by Castelnuovo's rationality criterion one can conclude that X_2 is a rational surface with $K_{X_2}^2 = 0$, which also comes with an elliptic fibration. The limit stable surface is the union of two rational surfaces glued along a genus one curve.

As in type a and type b, we may generalize this computation to all the degenerate point configurations of interest.

Lemma 5.13. *Consider a semi-stable choice of $\lambda_1, \dots, \lambda_8 \in \mathbb{C}^8 \subseteq (\mathbb{P}^1)^8$ such that $\lambda_5\lambda_6\lambda_7\lambda_8 \neq 0$, at least one of $\lambda_1, \lambda_2, \lambda_3, \lambda_4$ is different from the others, and at least one of $\lambda_5, \lambda_6, \lambda_7, \lambda_8$ is different from the others. Then the irreducible components $(Y_2, \frac{1+\epsilon}{2}B_{Y_2} + D_{Y_2}), (Y_3, \frac{1+\epsilon}{2}B_{Y_3} + D_{Y_3})$ of the limit pair are stable.*

Proof. To check that the pairs are stable, it is sufficient to check the singularities of the pairs corresponding to the exceptional divisors of the first two weighted blow-ups, because the flip, which was described in Example 5.10, does not produce any new singularities on the curve. With reference to the notation introduced in Example 5.11, we have to show that $(F_0, \frac{1+\epsilon}{2}(C_0 + L_1) + D_{F_0})$ and $(F_1, \frac{1+\epsilon}{2}(C_1 + L_0) + D_{F_1})$ are log canonical. We only show this for the former pair since the proof for the latter is analogous.

Note that from the equation of C_0 in Example 5.11, it is clear that C_0 does not intersect two coordinate points $V(t, x_0)$ and $V(t, y_1)$. In particular, $C_0 + L_1$ is supported on the smooth locus of F_0 . If all λ_i 's for $i \leq 4$ are distinct, then C_0 is nonsingular and intersects with L_1 and D_{F_0} transversally. By checking the allowed collisions of the λ_i with $i \leq 4$, one can conclude that $(C_0 + L_1)$ has only singularities on Table 1, and it intersects with D_{F_0} at three points transversally. Thus the pair is log canonical. \square

Remark 5.14. Let $\mathcal{X}^0 \rightarrow \Delta \setminus \{0\}$ be a proper flat family of smooth K3 surfaces with a purely non-symplectic automorphism of order four and a $U(2) \oplus D_4^{\oplus 2}$ lattice polarization as in § 2.1. Let \mathcal{X} be a semistable completion of \mathcal{X}^0 over Δ with \mathcal{X} smooth and $K_{\mathcal{X}} \sim 0$ (see [Kul77, PP81]). If the central fiber \mathcal{X}_0 is not smooth, then \mathcal{X}_0 has Kulikov type II according to [Kul77, Theorem II]. This follows from the study of degenerations we carried out in § 5, together with [Sch16, Theorem 7.4].

Remark 5.15. Motivated by the study of the KSBA compactification of the moduli space of K3 surfaces of degree two, in [AT21], Alexeev and Thompson introduced classes of combinatorially defined surfaces called *ADE* surfaces and $\tilde{A}\tilde{D}\tilde{E}$ surfaces [AT21, § 3]. These are log canonical non-klt del Pezzo surfaces with reduced boundary. An *ADE* (resp. $\tilde{A}\tilde{D}\tilde{E}$) cover is a double cover of an *ADE* (resp. $\tilde{A}\tilde{D}\tilde{E}$) surface.

The three K3 degenerations of type a, b, and c in this section are reducible surfaces whose irreducible components are of type II according to [AT21, Lemma 2.5]. With reference to the notation introduced in [AT21, § 3A], before taking the \mathbb{Z}_2 -cover, we have that for a degeneration of type a each irreducible component is a \tilde{D}_8 surface. For a type b degeneration each irreducible component is an \tilde{E}_7 surface. Finally, for a type c degeneration, the first weighted blow-up in Figure 5 produces a component which is an \tilde{E}_8^- surface. The next four blow-ups/downs correspond to the 'priming' operation in [AT21, Definition 3.13]. The resulting degeneration has two irreducible components which are \tilde{E}_8^+ surfaces.

A stable ADE (resp. $\tilde{A}\tilde{D}\tilde{E}$) cover is obtained by gluing ADE (resp. $\tilde{A}\tilde{D}\tilde{E}$) covers along boundaries. Alexeev and Thompson constructed moduli spaces of ADE covers [AT21, Theorem C]. Moduli spaces of $\tilde{A}\tilde{D}\tilde{E}$ covers, which are related to our situation, are not completely studied yet.

6. PROOF OF THE MAIN THEOREM

We are ready to prove Theorem 1.1. Let $(\mathcal{Y}, \frac{1+\epsilon}{2}\mathcal{B}) \rightarrow \mathbf{U}$ as in the beginning of §5. The first step is the following statement:

Proposition 6.1. *There is a flat family of stable pairs $(\tilde{\mathcal{Y}}, \frac{1+\epsilon}{2}\tilde{\mathcal{B}}) \rightarrow \mathbf{X}_1^s$ which is an extension of $(\mathcal{Y}, \frac{1+\epsilon}{2}\mathcal{B}) \rightarrow \mathbf{U}$.*

Proof. We subdivide the proof in different parts.

Setup and notation: Let $([a_1 : b_1], \dots, [a_8 : b_8])$ be the homogeneous coordinates of $\mathbf{X}_0 := ((\mathbb{P}^1)^8)$. Let $\mathcal{Y}_0 := \mathbf{X}_0 \times (\mathbb{P}^1 \times \mathbb{P}^1)$ and let $\pi_0 : \mathcal{Y}_0 \rightarrow \mathbf{X}_0$ be the natural projection. For $\mathbb{P}^1 \times \mathbb{P}^1$, let $([x_0 : x_1], [y_0 : y_1])$ be the homogeneous coordinates.

As usual, let \mathcal{C} be the divisor on \mathcal{Y}_0 defined by

$$y_0^2 \prod_{i=1}^4 (b_i x_0 - a_i x_1) + y_1^2 \prod_{i=5}^8 (b_i x_0 - a_i x_1) = 0.$$

Let $\mathcal{L}_0 := V(y_0)$, $\mathcal{L}_1 := V(y_1)$, $\mathcal{B}_0 := \mathcal{C} + \mathcal{L}_0 + \mathcal{L}_1$. Then $(\mathcal{Y}_0, \frac{1+\epsilon}{2}\mathcal{B}_0)$ is a family of pairs over \mathbf{X}_0 . Note that these pairs are stable over \mathbf{X}_0^s .

Recall that for $I \in \binom{[8]}{4}$, $\Delta_{I,I^c} := \Delta_I \cap \Delta_{I^c}$ is a connected component of the locus of strictly semi-stable points with closed orbits. Consult §4.2 for the detail. We say that Δ_{I,I^c} or the exceptional divisor E_{I,I^c} is of type a if Δ_{I,I^c} parametrizes type a point configurations. In the same way we define type b and type c components.

Let $\mathbf{X}_1 \rightarrow \mathbf{X}_0^{ss}$ be the blow-up along $\cup \Delta_{I,I^c}$ and let ρ_1 be the composition $\mathbf{X}_1^s \hookrightarrow \mathbf{X}_1 \rightarrow \mathbf{X}_0^{ss}$. Let $(\mathcal{Y}_1, \frac{1+\epsilon}{2}\mathcal{B}_1)$ be the pulled-back family over \mathbf{X}_1^s , that is, $\mathcal{Y}_1 := \mathbf{X}_1^s \times (\mathbb{P}^1 \times \mathbb{P}^1)$ and $\mathcal{B}_1 := \mathcal{C}_1 + \mathcal{L}_{0,1} + \mathcal{L}_{1,1} := (\rho_1 \times \text{id})^*(\mathcal{C} + \mathcal{L}_0 + \mathcal{L}_1)|_{\mathbf{X}_0^{ss} \times (\mathbb{P}^1 \times \mathbb{P}^1)}$. Let $\pi_1 : \mathcal{Y}_1 \rightarrow \mathbf{X}_1^s$ be the natural projection.

Main idea: The claimed extension $(\tilde{\mathcal{Y}}, \frac{1+\epsilon}{2}\tilde{\mathcal{B}}) \rightarrow \mathbf{X}_1^s$ is obtained from $(\mathcal{Y}_1, \frac{1+\epsilon}{2}\mathcal{B}_1) \rightarrow \mathbf{X}_1^s$ by first appropriately blowing-up \mathcal{Y}_1 , and then applying the relative minimal model program. It is clear that the fibers of $(\mathcal{Y}_1, \frac{1+\epsilon}{2}\mathcal{B}_1) \rightarrow \mathbf{X}_1^s$ that are not stable lie above the divisors E_{I,I^c}^s . In what follows, we blow-up appropriate sub-loci of $\pi_1^{-1}(E_{I,I^c}^s)$ mimicking in a relative setting the blow-ups described in §5.

Type a modification: For a type a divisor E_{I,I^c} , let $\mathcal{Z}_I^a := \pi_1^{-1}(E_{I,I^c}^s) \cap V(b_i x_0 - a_i x_1 \mid i \in I)$. Note that we intentionally abuse our notation: it is understood that the vanishing locus $V(b_i x_0 - a_i x_1 \mid i \in I) \subseteq \mathcal{Y}_0$ is pulled-back to \mathcal{Y}_1 under the appropriate morphism. This is a smooth codimension two subvariety of \mathcal{Y}_1 , which is a \mathbb{P}^1 -bundle over E_{I,I^c}^s . Note that $\mathcal{Z}_{I^c}^a$ is also in $\pi_1^{-1}(E_{I,I^c}^s)$ but it is disjoint from \mathcal{Z}_I^a . Let $\mathcal{Y}_2 \rightarrow \mathcal{Y}_1$ be the blow-up along $\cup(\mathcal{Z}_I^a \sqcup \mathcal{Z}_{I^c}^a)$, where the union is over all type a divisors E_{I,I^c} . Fiberwisely, this is exactly the blow-up of the two double lines as in Example 5.2. Let \mathcal{C}_2 (resp. $\mathcal{L}_{i,2}$) be the

proper transform of \mathcal{C}_1 (resp. $\mathcal{L}_{i,1}$) and $\mathcal{B}_2 := \mathcal{C}_2 + \mathcal{L}_{0,2} + \mathcal{L}_{1,2}$. Let π_2 be the composition $\mathcal{Y}_2 \rightarrow \mathcal{Y}_1 \rightarrow \mathbf{X}_1^s$. Then $\pi_2 : (\mathcal{Y}_2, \frac{1+\epsilon}{2}\mathcal{B}_2) \rightarrow \mathbf{X}_1^s$ is a flat family of pairs. Over a type a divisor E_{I,I^c}^s , each fiber of $(\mathcal{Y}_2, \frac{1+\epsilon}{2}\mathcal{B}_2)$ is semi-log canonical and it is isomorphic to the pair in Figure 2.

Type b modification: Over a type b divisor E_{I,I^c} , we may assume that $|I \cap \{1, \dots, 4\}| = 3$ (hence $|I^c \cap \{5, 6, 7, 8\}| = 3$). Let $\mathcal{Z}_I^b := \pi_2^{-1}(E_{I,I^c}^s) \cap \mathcal{L}_1 \cap V(b_i x_0 - a_i x_1 \mid i \in I)$ and let $\mathcal{Z}_{I^c}^b := \pi_2^{-1}(E_{I,I^c}^s) \cap \mathcal{L}_0 \cap V(b_i x_0 - a_i x_1 \mid i \in I^c)$. Then \mathcal{Z}_I^b and $\mathcal{Z}_{I^c}^b$ are both disjoint sections of E_{I,I^c}^s . Let $\mathcal{Y}_3 \rightarrow \mathcal{Y}_2$ be the blow-up along $\cup(\mathcal{Z}_I^b \sqcup \mathcal{Z}_{I^c}^b)$ for all type b divisors E_{I,I^c}^s . Let \mathcal{C}_3 (resp. $\mathcal{L}_{i,3}$) be the proper transform of \mathcal{C}_2 (resp. $\mathcal{L}_{i,2}$) and let $\mathcal{B}_3 := \mathcal{C}_3 + \mathcal{L}_{0,3} + \mathcal{L}_{1,3}$. Let π_3 be the composition $\mathcal{Y}_3 \rightarrow \mathcal{Y}_2 \rightarrow \mathbf{X}_1^s$. Then $\pi_3 : (\mathcal{Y}_3, \frac{1+\epsilon}{2}\mathcal{B}_3) \rightarrow \mathbf{X}_1^s$ is a family of pairs and over a type b divisor, each fiber of $(\mathcal{Y}_3, \frac{1+\epsilon}{2}\mathcal{B}_3)$ is semi-log canonical and it is isomorphic to the pair in Figure 4.

Type c modification: Finally, over the type c divisor E_{I,I^c} , we may assume that $I = \{1, 2, 3, 4\}$. Let $\mathcal{P} := \pi_3^{-1}(E_{I,I^c}^s) \cong E_{I,I^c}^s \times \mathbb{P}^1 \times \mathbb{P}^1$. Let $\mathcal{Z}_I^c := \pi_3^{-1}(E_{I,I^c}^s) \cap \mathcal{L}_1 \cap V(b_i x_0 - a_i x_1 \mid i \in I)$ and $\mathcal{Z}_{I^c}^c := \pi_3^{-1}(E_{I,I^c}^s) \cap \mathcal{L}_0 \cap V(b_i x_0 - a_i x_1 \mid i \in I^c)$. Then \mathcal{Z}_I^c and $\mathcal{Z}_{I^c}^c$ are disjoint sections of E_{I,I^c}^s . Let $\mathcal{Y}'_4 \rightarrow \mathcal{Y}_3$ be the weighted blow-up along $\cup(\mathcal{Z}_I^c \sqcup \mathcal{Z}_{I^c}^c)$ where the normal subbundles $\mathcal{N}_{\mathcal{Z}_I^c/\mathcal{L}_{1,3}}$ and $\mathcal{N}_{\mathcal{Z}_{I^c}^c/\mathcal{L}_{0,3}}$ have weight two. Let \mathcal{C}'_4 (resp. $\mathcal{L}'_{i,4}$) be the proper transform of \mathcal{C}_3 (resp. $\mathcal{L}_{i,3}$). Set $\mathcal{B}'_4 := \mathcal{C}'_4 + \mathcal{L}'_{0,4} + \mathcal{L}'_{1,4}$. Let \mathcal{P}' be the proper transform of \mathcal{P} .

Let $\mathcal{W}_i := \mathcal{L}'_{i,4} \cap \mathcal{P}'$, which is a smooth codimension two subvariety of \mathcal{Y}'_4 . Note that \mathcal{W}_0 and \mathcal{W}_1 are disjoint. Let $\mathcal{Y}''_4 \rightarrow \mathcal{Y}'_4$ be the blow-up along $\mathcal{W}_0 \sqcup \mathcal{W}_1$. The two exceptional divisors are denoted by E_0 and E_1 . Let $\mathcal{C}''_4, \mathcal{L}''_{i,4}$ be the proper transforms of $\mathcal{C}'_4, \mathcal{L}'_{i,4}$ respectively. Set $\mathcal{B}''_4 := \mathcal{C}''_4 + \mathcal{L}''_{0,4} + \mathcal{L}''_{1,4}$. Note that $\mathcal{C}''_4 \cong \mathcal{C}'_4$ because \mathcal{C}'_4 and \mathcal{W}_i are disjoint.

Let $\mathcal{V}_i := \mathcal{L}''_{i,4} \cap E_i$, which is a smooth codimension two subvariety of \mathcal{Y}''_4 . \mathcal{V}_0 is disjoint from \mathcal{V}_1 . Let $\mathcal{Y}_4 \rightarrow \mathcal{Y}''_4$ be the blow-up along $\mathcal{V}_0 \sqcup \mathcal{V}_1$. Let $\mathcal{C}_4, \mathcal{L}_{i,4}$ be the proper transforms of $\mathcal{C}''_4, \mathcal{L}''_{i,4}$ respectively and let $\mathcal{B}_4 := \mathcal{C}_4 + \mathcal{L}_{0,4} + \mathcal{L}_{1,4}$. The family of pairs $\pi : (\mathcal{Y}_4, \frac{1+\epsilon}{2}\mathcal{B}_4) \rightarrow \mathbf{X}_1^s$ is semi-log canonical over the type c divisor. Over E_{I,I^c}^s , the fiber of \mathcal{Y}_4 has seven irreducible components. There is a ‘central’ component, and two ‘tails’ consisting of three irreducible components whose configurations are the same to the top three components of the fourth step in Figure 5.

Contraction: Consider the resulting family of pairs $\pi : (\mathcal{Y}_4, \frac{1+\epsilon}{2}\mathcal{B}_4) \rightarrow \mathbf{X}_1^s$. For each $x \in \mathbf{X}_1^s$, the fiber $(\mathcal{Y}_{4x}, \frac{1+\epsilon}{2}\mathcal{B}_{4x})$ is either irreducible stable pair or one of semi-stable pairs described in Figures 2, 4, and 6. In any case, each fiber is a semi-log canonical pair and $K_{\mathcal{Y}_{4x}} + \frac{1+\epsilon}{2}\mathcal{B}_{4x}$ is a nef and big divisor. Thus, by [Fuj14, Theorem 1.10], $H^1(\mathcal{Y}_{4x}, \mathcal{O}(m(K_{\mathcal{Y}_{4x}} + \frac{1+\epsilon}{2}\mathcal{B}_{4x}))) = 0$ for $m \gg 0$. Moreover, by [Fuj14, Theorem 1.16], $K_{\mathcal{Y}_{4x}} + \frac{1+\epsilon}{2}\mathcal{B}_{4x}$ is semi-ample. Then by the standard cohomology and base change [Har77, Theorem 12.11], $\pi_* \mathcal{O}(m(K_{\mathcal{Y}_4/\mathbf{X}_1^s} + \frac{1+\epsilon}{2}\mathcal{B}_4))$ is locally free and we obtain a new family of varieties

$$\tilde{\mathcal{Y}} := \text{Proj} \bigoplus_{m \geq 0} \pi_* \mathcal{O} \left(m \left(K_{\mathcal{Y}_4/\mathbf{X}_1^s} + \frac{1+\epsilon}{2}\mathcal{B}_4 \right) \right) \rightarrow \mathbf{X}_1^s,$$

and there is a contraction \mathbf{X}_1^s -morphism $\mathcal{Y}_4 \rightarrow \tilde{\mathcal{Y}}$. By taking the push-forward of \mathcal{B}_4 , we obtain $(\tilde{\mathcal{Y}}, \frac{1+\epsilon}{2}\tilde{\mathcal{B}}) \rightarrow \mathbf{X}_1^s$. The resulting morphism $\tilde{\mathcal{Y}} \rightarrow \mathbf{X}_1^s$ is flat by [HKT09, Lemma 10.12].

Then by [Kol18, Theorem 4.3], $(\tilde{\mathcal{Y}}, \frac{1+\epsilon}{2}\tilde{\mathcal{B}}) \rightarrow \mathbf{X}_1^s$ is a well-defined family of pairs. Finally, Lemmas 5.4, 5.9, and 5.13 tell us that each fiber of $(\tilde{\mathcal{Y}}, \frac{1+\epsilon}{2}\tilde{\mathcal{B}}) \rightarrow \mathbf{X}_1^s$ is stable. \square

Proof of Theorem 1.1. In Proposition 3.13 we showed that $\bar{\mathbf{K}} \cong \bar{\mathbf{J}}$, so we focus on the latter. By Proposition 6.1 we have a family of stable pairs over \mathbf{X}_1^s for the functor $\bar{\mathcal{J}}'$ in Definition 3.11, hence there is a functorial morphism $f : \mathbf{X}_1^s \rightarrow \bar{\mathbf{J}}$. Since \mathbf{U} is an open dense subset of \mathbf{X}_1^s , the image of f is precisely $\bar{\mathbf{J}}$. Clearly the morphism f is SL_2 -invariant, so there is a quotient morphism $\bar{f} : \mathbf{P} = \mathbf{X}_1^s/\mathrm{SL}_2 \rightarrow \bar{\mathbf{J}}$. This map is also H -invariant, thus we can obtain yet another quotient map $\tilde{f} : \mathbf{P}/H \rightarrow \bar{\mathbf{J}}$ which we show is an isomorphism.

Since \tilde{f} is a birational morphism between normal varieties, it is enough to show it is finite, or equivalently that \bar{f} is finite. Given an exceptional divisor E_{I,I^c} of the blow-up $\mathbf{X}_1 \rightarrow \mathbf{X}_0^{ss}$, denote by \bar{E}_{I,I^c} the quotient $E_{I,I^c}^s/\mathrm{SL}_2$. It is sufficient to show that \bar{E}_{I,I^c} , which is isomorphic to $\mathbb{P}^2 \times \mathbb{P}^2$ by Remark 4.2, is not contracted by \bar{f} . This is explained in Lemma 6.2 below. \square

Lemma 6.2. *With the notation introduced in the proof of Theorem 1.1, the divisors $\bar{E}_{I,I^c} \subseteq \mathbf{P}$ are not contracted by \bar{f} .*

Proof. If $\bar{E}_{I,I^c} \cong \mathbb{P}^2 \times \mathbb{P}^2$ is contracted, then at least one of the two copies of \mathbb{P}^2 , say the first component, is contracted to a point. We show that we can find two points $(\bar{p}_1, \bar{q}), (\bar{p}_2, \bar{q}) \in \bar{E}_{I,I^c}$ with $\bar{p}_1 \neq \bar{p}_2$ parametrizing non-isomorphic stable pairs, obtaining a contradiction. There are three cases to consider corresponding to the type of \bar{E}_{I,I^c} , which we define to be equal to the type of E_{I,I^c} .

Assume \bar{E}_{I,I^c} is of type a. Up to relabeling, we may assume $I = \{1, 2, 5, 6\}, I^c = \{3, 4, 7, 8\}$. The stable pair parametrized by a point in E_{I,I^c}^s has two irreducible components isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$ (see Example 5.2). Consider the irreducible component with divisor in the form $C_0 + L_0 + L_1$, where C_0 is given by

$$y_0^2(x_0 - t\lambda_1)(x_0 - t\lambda_2)\lambda_3\lambda_4 + y_1^2(x_0 - t\lambda_5)(x_0 - t\lambda_6)\lambda_7\lambda_8 = 0.$$

Recall from Remark 5.5 that $\lambda_3\lambda_4\lambda_7\lambda_8 \neq 0$ and at least one of $\lambda_1, \lambda_2, \lambda_5, \lambda_6$ is different from the others. So pick any point $(p_1, q) \in E_{I,I^c}^s$ such that the corresponding $\lambda_1, \lambda_2, \lambda_5, \lambda_6$ are distinct. Consider the projection from $\mathbb{P}^1 \times \mathbb{P}^1$ on the $[x_0 : t]$ coordinate. The images of the four points $C_0 \cap L_0, C_0 \cap L_1$ are $[\lambda_1 : 1], [\lambda_2 : 1], [\lambda_5 : 1], [\lambda_6 : 1]$, which are distinct points on \mathbb{P}^1 . Denote by β their cross-ratio. Choose $[\mu_1 : 1], [\mu_2 : 1], [\mu_5 : 1], [\mu_6 : 1]$ distinct points on \mathbb{P}^1 such that the corresponding cross-ratio is different from β . Then the stable pair obtained by replacing $\lambda_1, \lambda_2, \lambda_5, \lambda_6$ with $\mu_1, \mu_2, \mu_5, \mu_6$ respectively and keeping $\lambda_3, \lambda_4, \lambda_7, \lambda_8$ unchanged is parametrized by a point $(p_2, q) \in E_{I,I^c}^s$ with $p_1 \neq p_2$. The images $(\bar{p}_1, \bar{q}), (\bar{p}_2, \bar{q})$ in \bar{E}_{I,I^c} are also distinct because the cross-ratio is SL_2 -invariant, showing what we needed.

The cases of type b and type c are handled similarly, but with the following differences. For type b, given a stable pair parametrized by E_{I,I^c}^s , one can consider the cross-ratio of the four points on L_1 given by $[\lambda_1 : 1], [\lambda_2 : 1], [\lambda_3 : 1], [1 : 0]$, where the last point is the intersection of L_1 with the conductor divisor. For type c, look at $[\lambda_1 : 1], [\lambda_2 : 1], [\lambda_3 : 1], [\lambda_4 : 1]$ on the curve T in Figure 5. \square

REFERENCES

- [Ale96] Valery Alexeev. Moduli spaces $M_{g,n}(W)$ for surfaces. *Higher-Dimensional Complex Varieties (Trento, 1994)*, de Gruyter, Berlin, 1996, pp. 1–22. [1](#), [9](#)
- [Ale15] Valery Alexeev. *Moduli of weighted hyperplane arrangements*. Edited by Gilberto Bini, Martí Lahoz, Emanuele Macrì and Paolo Stellari. Advanced Courses in Mathematics. CRM Barcelona. Birkhäuser/Springer, Basel, 2015. [7](#), [9](#)
- [AP09] Valery Alexeev and Rita Pardini. Explicit compactifications of moduli spaces of Campedelli and Burniat surfaces. (2009) arXiv:0901.4431. [1](#)
- [AP12] Valery Alexeev and Rita Pardini. Non-normal abelian covers. *Compos. Math.* 148 (2012), no. 4, 1051–1084. [2](#), [7](#), [8](#)
- [AT21] Valery Alexeev and Alan Thompson. ADE surfaces and their moduli. *J. Algebraic Geom.* 30 (2021), no. 2, 331–405. [22](#), [23](#)
- [AET19] Valery Alexeev, Philip Engel, and Alan Thompson. Stable pair compactification of moduli of K3 surfaces of degree 2. (2019) arXiv:1903.09742. [1](#)
- [AS15] Michela Artebani and Alessandra Sarti. Symmetries of order four on K3 surfaces. *J. Math. Soc. Japan* 67 (2015), no. 2, 503–533. [1](#), [4](#)
- [AB19] Kenneth Ascher and Dori Bejleri. Compact moduli of elliptic K3 surfaces. (2019) arXiv:1902.10686. [1](#)
- [BHPV04] Wolf Barth, Klaus Hulek, Chris Peters, and Antonius Van de Ven. *Compact complex surfaces*. Second edition. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. Springer-Verlag, Berlin, 2004. xii+436 pp. [6](#)
- [DM86] Pierre Deligne and George D. Mostow. Monodromy of hypergeometric functions and nonlattice integral monodromy. *Inst. Hautes Études Sci. Publ. Math.* No. 63 (1986), 5–89. [2](#)
- [DH21] Anand Deopurkar and Changho Han. Stable log surfaces, admissible covers, and canonical curves of genus 4. *Trans. Amer. Math. Soc.* 374 (2021), 589–641. [1](#)
- [DK07] Igor Dolgachev and Shigeyuki Kondō. Moduli of K3 surfaces and complex ball quotients. *Arithmetic and geometry around hypergeometric functions*, 43–100, Progr. Math., 260, Birkhäuser, Basel, 2007. [4](#)
- [Fuj14] Osamu Fujino. Fundamental theorems for semi log canonical pairs. *Algebr. Geom.* 1 (2014), no. 2, 194–228. [9](#), [24](#)
- [GKS21] Patricio Gallardo, Matt Kerr, and Luca Schaffler. Geometric interpretation of toroidal compactifications of moduli of points in the line and cubic surfaces. To appear in *Advances in Mathematics*, 381 (2021), <https://doi.org/10.1016/j.aim.2021.107632>. [3](#)
- [GMGZ18] Patricio Gallardo, Jesus Martinez-Garcia, and Zheng Zhang. Compactifications of the moduli space of plane quartics and two lines. *Eur. J. Math.* 4, No. 3, 1000–1034 (2018). [1](#)
- [HKT09] Paul Hacking, Sean Keel, and Jenia Tevelev. Stable pair, tropical, and log canonical compactifications of moduli spaces of del Pezzo surfaces. *Invent. Math.* 178 (2009), no. 1, 173–227. [24](#)
- [Har77] Robin Hartshorne. *Algebraic geometry*. Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977. xvi+496 pp. [24](#)
- [Has03] Brendan Hassett. Moduli spaces of weighted pointed stable curves. *Adv. Math.* 173 (2003), 316–352. [13](#)
- [Ish18] Shihoko Ishii. *Introduction to singularities*. Second edition. Springer, Tokyo, 2018. x+236 pp. [6](#)
- [KM11] Young-Hoon Kiem and Han-Bom Moon. Moduli spaces of weighted pointed stable rational curves via GIT. *Osaka J. Math.* 48 (2011), no. 4, 1115–1140. [2](#), [13](#)
- [Kir85] Frances Kirwan. Partial desingularisations of quotients of nonsingular varieties and their Betti numbers. *Ann. of Math. (2)* 122 (1985), no. 1, 41–85. [3](#), [11](#), [12](#)
- [Kol13] János Kollár. *Singularities of the minimal model program*. With a collaboration of Sándor Kovács. Cambridge Tracts in Mathematics, 200. Cambridge University Press, Cambridge, 2013. [7](#), [8](#), [10](#)
- [Kol18] János Kollár. *Families of varieties of general type*. Book in preparation, 2018. [1](#), [7](#), [25](#)
- [KSB88] János Kollár and Nick Shepherd-Barron. Threefolds and deformations of surface singularities. *Invent. Math.* 91 (1988), no. 2, 299–338. [1](#)

- [Kon07] Shigeyuki Kondō. The moduli space of 8 points on \mathbb{P}^1 and automorphic forms. *Algebraic geometry, 89–106, Contemp. Math.*, 422, Amer. Math. Soc., Providence, RI, 2007. [1](#), [2](#), [3](#), [4](#), [5](#), [6](#)
- [Kul77] Viktor S. Kulikov. Degenerations of K3 surfaces and Enriques surfaces. *Math. USSR Izv.* 11 (1977), 957–989. [22](#)
- [LO18] Radu Laza and Kieran O’Grady. GIT versus Baily–Borel compactification for K3’s which are double covers of $\mathbb{P}^1 \times \mathbb{P}^1$. (2018) arXiv:1801.04845. [3](#)
- [Laz04] Robert Lazarsfeld. *Positivity in algebraic geometry. I. Classical setting: line bundles and linear series.* Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], 48, Springer-Verlag, Berlin, 2004. [8](#)
- [Loo86] Eduard Looijenga. New compactifications of locally symmetric varieties. *Proceedings of the 1984 Vancouver conference in algebraic geometry, 341–364, CMS Conf. Proc.*, 6, Amer. Math. Soc., Providence, RI, 1986. [3](#)
- [Mnë85] Nikolai Mnëv. Varieties of combinatorial types of projective configurations and convex polyhedra. *Dokl. Akad. Nauk SSSR* 283 (1985), 1312–1314. [1](#)
- [MFK94] David Mumford, John Fogarty, and Francis Kirwan. *Geometric invariant theory.* Third edition. Ergebnisse der Mathematik und ihrer Grenzgebiete (2), 34. Springer-Verlag, Berlin, 1994. xiv+292 pp. [5](#)
- [Ols16] Martin Olsson. *Algebraic spaces and stacks.* American Mathematical Society Colloquium Publications, 62. American Mathematical Society, Providence, RI, 2016. xi+298 pp. [9](#)
- [PP81] Ulf Persson and Henry Pinkham. Degeneration of surfaces with trivial canonical bundle. *Ann. of Math. (2)* 113 (1981), no. 1, 45–66. [22](#)
- [Sch16] Luca Schaffler. The KSBA compactification of the moduli space of $D_{1,6}$ -polarized Enriques surfaces. (2016) arXiv:1608.02564. [1](#), [22](#)
- [Sha81] Jayant Shah. Degenerations of K3 surfaces of degree 4. *Trans. Amer. Math. Soc.* 263 (1981), 271–308. [6](#)
- [Vak06] Ravi Vakil. Murphy’s law in algebraic geometry: badly-behaved deformation spaces. *Invent. Math.* 164 (2006), no. 3, 569–590. [1](#)
- [Zha07] De-Qi Zhang. Automorphisms of K3 surfaces. *Proceedings of the International Conference on Complex Geometry and Related Fields, 379–392, AMS/IP Stud. Adv. Math.*, 39, Amer. Math. Soc., Providence, RI, 2007. [1](#)

DEPARTMENT OF MATHEMATICS, FORDHAM UNIVERSITY, NEW YORK, NY 10023

Email address: hmoon8@fordham.edu

DEPARTMENT OF MATHEMATICS, KTH ROYAL INSTITUTE OF TECHNOLOGY, SE-100 44 STOCKHOLM, SWEDEN

Email address: lucsch@math.kth.se