

# A class of non-weight modules over the Schrödinger-Virasoro algebras

Yumei Wang and Honglian Zhang\*

ABSTRACT. We construct and classify the free  $U(\mathbb{C}L_0 \oplus \mathbb{C}M_0 \oplus \mathbb{C}Y_0)$ -modules of rank 1 over the Schrödinger-Virasoro algebra  $\mathfrak{sv}(s)$  for  $s = 0$ . Moreover, we show that the class of free  $U(\mathbb{C}L_0 \oplus \mathbb{C}M_0)$ -modules of rank 1 over the Schrödinger-Virasoro algebra  $\mathfrak{sv}(s)$  for  $s = \frac{1}{2}$  is nonexistent.

## 1. Introduction

In the 1960s, physicist Virasoro gave an important infinite dimensional Lie algebra—the Virasoro algebra with a basis  $\{L_n, C | n \in \mathbb{Z}\}$  and the following relations

$$[L_n, L_m] = (m - n)L_{m+n} + \delta_{m+n} \frac{n^3 - n}{12} C, \quad m, n \in \mathbb{Z},$$

$$[L_n, C] = 0, \quad n \in \mathbb{Z},$$

which is a universal central extension of the Witt algebra. The Virasoro algebra is closely related to string theory [17] and conformal field theory [15]. Additionally, the representation theory of the Virasoro algebra is widely applied in many branches of mathematics and physics, such as vertex algebras [7, 19] and quantum physics [10]. Furthermore, the generalizations of the Virasoro algebra are extensively studied, such as the Schrödinger-Virasoro algebras [4, 16, 20], the Virasoro-like algebras [9, 14], the affine Virasoro algebras [8] and so on.

The Schrödinger-Virasoro algebra is one of the natural generalizations of the Virasoro algebra, which was introduced by M. Henkel in [11] during his study of the free Schrödinger equations in non-equilibrium statistical physics. The Schrödinger-Virasoro algebra  $\mathfrak{sv}(s)$  for  $s = 0$  or  $\frac{1}{2}$  (c.f. [11], [13]) is an infinite-dimensional Lie algebra over  $\mathbb{C}$  with a basis  $\{L_n, M_n, Y_p | n \in \mathbb{Z}, p \in \mathbb{Z} + s\}$  and satisfying the following relations

$$[L_n, L_m] = (m - n)L_{m+n},$$

$$[L_n, Y_p] = (p - \frac{n}{2})Y_{p+n},$$

$$[L_n, M_m] = mM_{m+n},$$

$$[Y_p, Y_q] = (q - p)M_{p+q},$$

---

2010 *Mathematics Subject Classification*. Primary 17B10, 17B65, 17B68.

*Key words and phrases*. the Schrödinger-Virasoro algebra, non-weight module, free module.

\* H. Zhang, Corresponding Author.

$$[Y_p, M_m] = [M_n, M_m] = 0,$$

where  $m, n \in \mathbb{Z}$  and  $p, q \in \mathbb{Z} + s$ . The Schrödinger-Virasoro algebras play an important role in many fields of mathematics and physics. Furthermore, there were a number of works on the Schrödinger-Virasoro algebras and their representations theory (see [4, 16, 20–22] ect.). Actually, the weight modules of the Schrödinger-Virasoro algebras have been studied extensively. For instance, Li and Su studied the weight modules with finite-dimensional weight spaces over the Schrödinger-Virasoro algebras in [16].

On the other hand, it is well known that the theory of non-weight modules has been extensively studied in the past few years. In [1], Batra and Mazorchuk gave a general set-up for studying Whittaker modules, from which Nilsson considered to construct families of simple  $\mathfrak{sl}_{n+1}$ -modules. In [18], Nilsson determined and classified the free  $U(\mathfrak{h})$ -modules of rank 1 over  $\mathfrak{sl}_{n+1}$ , in which  $\mathfrak{h}$  is the standard Cartan subalgebra of  $\mathfrak{sl}_{n+1}$ . Furthermore, the idea of Nilsson in [18] has been generalized and applied into many infinite dimensional algebras. The non-weight modules which are free of rank 1 over the Kac-Moody algebras and the classical Lie superalgebras were studied in [5] and [6] respectively. Moreover, such modules for the Virasoro algebra and its related algebras were investigated in [2] and [12]. In addition, Chen and Guo constructed the free modules of rank 1 and determined the simplicity over the Heisenberg-Virasoro algebra and the  $W(2, 2)$  algebra in [3].

Our goal of the present paper is to focus on the non-weight modules over the Schrödinger-Virasoro algebra  $\mathfrak{sv}(s)$  for  $s = 0$  or  $\frac{1}{2}$ . We construct a class of free  $U(\mathbb{C}L_0 \oplus \mathbb{C}M_0 \oplus \mathbb{C}Y_0)$ -modules of rank 1 over the algebra  $\mathfrak{sv}(0)$ , denoted by  $\Phi(\lambda, \alpha)$  for  $\lambda \in \mathbb{C}^*$  and  $\alpha \in \mathbb{C}$ . Moreover, we classify all such modules and obtain the main result of the present paper: Let  $\mathfrak{sv}(0)$  be the Schrödinger-Virasoro algebra  $\mathfrak{sv}(s)$  for  $s = 0$ , if  $M$  is a free  $U(\mathbb{C}L_0 \oplus \mathbb{C}M_0 \oplus \mathbb{C}Y_0)$ -module of rank 1 over the algebra  $\mathfrak{sv}(0)$ , then there exists some  $\lambda \in \mathbb{C}^*$  and  $\alpha \in \mathbb{C}$  such that  $M \cong \Phi(\lambda, \alpha)$ . In addition, we prove that the class of free  $U(\mathbb{C}L_0 \oplus \mathbb{C}M_0)$ -modules of rank 1 over the algebra  $\mathfrak{sv}(\frac{1}{2})$  is nonexistent.

The paper is organized as follows. In Section 2, we demonstrate the definition and properties of the algebra  $\mathfrak{sv}(0)$ . Besides, we construct a class of free  $U(\mathbb{C}L_0 \oplus \mathbb{C}M_0 \oplus \mathbb{C}Y_0)$ -modules of rank 1 over the algebra  $\mathfrak{sv}(0)$ . Section 3 is aimed to classify the free  $U(\mathbb{C}L_0 \oplus \mathbb{C}M_0 \oplus \mathbb{C}Y_0)$ -modules of rank 1 over the algebra  $\mathfrak{sv}(0)$ . In Section 4, we mainly prove the nonexistence of free  $U(\mathbb{C}L_0 \oplus \mathbb{C}M_0)$ -modules of rank 1 over the algebra  $\mathfrak{sv}(\frac{1}{2})$ .

## 2. Preliminaries

In this section, we list some basic notations and useful results for our purpose. Throughout the paper,  $\mathbb{N}$ ,  $\mathbb{C}$ ,  $\mathbb{C}^*$ ,  $\mathbb{Z}$  and  $\mathbb{Z}^*$  stand for the set of all natural numbers, complex numbers, nonzero complex numbers, integers and nonzero integers, respectively. Note that we focus on the Schrödinger-Virasoro algebra  $\mathfrak{sv}(s)$  for  $s = 0$  in Section 2 and Section 3.

The Schrödinger-Virasoro algebra  $\mathfrak{sv}(0)$  has a  $\mathbb{C}$ -basis  $\{L_n, Y_n, M_n | n \in \mathbb{Z}\}$  with the following relations

$$(2.1) \quad [L_n, L_m] = (m - n)L_{m+n},$$

$$(2.2) \quad [L_n, Y_m] = (m - \frac{n}{2})Y_{m+n},$$

$$(2.3) \quad [L_n, M_m] = mM_{m+n},$$

$$(2.4) \quad [Y_n, Y_m] = (m - n)M_{m+n},$$

$$(2.5) \quad [Y_n, M_m] = [M_n, M_m] = 0,$$

where  $m \in \mathbb{Z}, n \in \mathbb{Z}$ .

From the above relations (2.1)-(2.5), we obtain some important formulas that will be needed later.

**PROPOSITION 2.1.** *For  $i \in \mathbb{Z}, n \in \mathbb{Z}$  and  $i \geq 1$ , the following formulas hold:*

- (1)  $M_n L_0^i = (L_0 - n)^i M_n.$
- (2)  $M_n M_0^i = M_0^i M_n.$
- (3)  $M_n Y_0^i = Y_0^i M_n.$
- (4)  $Y_n L_0^i = (L_0 - n)^i Y_n.$
- (5)  $Y_n M_0^i = M_0^i Y_n.$
- (6)  $Y_n Y_0^i = Y_0^i Y_n - n i Y_0^{i-1} M_n.$
- (7)  $L_n L_0^i = (L_0 - n)^i L_n.$
- (8)  $L_n M_0^i = M_0^i L_n.$
- (9)  $L_n Y_0^i = Y_0^i L_n - \frac{n i}{2} Y_0^{i-1} Y_n + \frac{n^2 i(i-1)}{4} Y_0^{i-2} M_n.$

**PROOF.** Here we only check formula (6) and formula (9), others can be easily shown by induction on  $i$ . For formula (6), it is obvious for  $i = 1$  according to  $Y_n Y_0 = Y_0 Y_n + [Y_n, Y_0] = Y_0 Y_n - n M_n$ . Now assume that formula (6) holds for  $i = k - 1$ , that is,

$$Y_n Y_0^{k-1} = Y_0^{k-1} Y_n - (k-1)n Y_0^{k-2} M_n.$$

For  $i = k$ , by the inductive assumption we have

$$\begin{aligned} Y_n Y_0^k &= Y_n Y_0^{k-1} Y_0 \\ &= Y_0^{k-1} Y_n Y_0 - (k-1)n Y_0^{k-2} M_n Y_0 \\ &= Y_0^k Y_n - n Y_0^{k-1} M_n - (k-1)n Y_0^{k-1} M_n \\ &= Y_0^k Y_n - k n Y_0^{k-1} M_n, \end{aligned}$$

which implies that formula (6) holds.

For formula (9), we first have  $L_n Y_0 = Y_0 L_n + [L_n, Y_0] = Y_0 L_n - \frac{n}{2} Y_n$ , formula (9) holds for  $i = 1$ . Next suppose that formula (9) holds for  $i = k - 1$ , one has

$$L_n Y_0^{k-1} = Y_0^{k-1} L_n - \frac{(k-1)n}{2} Y_0^{k-2} Y_n + \frac{(k-1)(k-2)n^2}{4} Y_0^{k-3} M_n.$$

For  $i = k$ , we immediately obtain that

$$\begin{aligned}
L_n Y_0^k &= L_n Y_0^{k-1} Y_0 \\
&= Y_0^{k-1} L_n Y_0 - \frac{(k-1)n}{2} Y_0^{k-2} Y_n Y_0 + \frac{(k-1)(k-2)n^2}{4} Y_0^{k-3} M_n Y_0 \\
&= Y_0^k L_n - \frac{n}{2} Y_0^{k-1} Y_n - \frac{(k-1)n}{2} Y_0^{k-1} Y_n + \frac{(k-1)n^2}{2} Y_0^{k-2} M_n \\
&\quad + \frac{(k-1)(k-2)n^2}{4} Y_0^{k-2} M_n \\
&= Y_0^k L_n - \frac{nk}{2} Y_0^{k-1} Y_n + \frac{k(k-1)n^2}{4} Y_0^{k-2} M_n.
\end{aligned}$$

Therefore formula (9) holds.  $\square$

**REMARK 2.2.** Note that these formulas in Proposition 2.1 also hold in the case of  $i = 0$ .

**DEFINITION 2.3.** For  $\lambda \in \mathbb{C}^*$  and  $\alpha \in \mathbb{C}$ , define the action of a basis of the algebra  $\mathfrak{sv}(0)$  on  $\Phi(\lambda, \alpha) := \mathbb{C}[s, t, v]$ , in which  $\mathbb{C}[s, t, v]$  is the polynomial algebra in three indeterminates  $s, t$  and  $v$ , as follows:

$$\begin{aligned}
(2.6) \quad L_m \cdot f(s, t, v) &= \lambda^m \left( (s + m\alpha) f(s - m, t, v) - \frac{m}{2} v \frac{\partial f(s - m, t, v)}{\partial v} \right. \\
&\quad \left. + \frac{m^2}{4} t \frac{\partial^2 f(s - m, t, v)}{\partial v^2} \right),
\end{aligned}$$

$$(2.7) \quad M_m \cdot f(s, t, v) = \lambda^m t f(s - m, t, v),$$

$$(2.8) \quad Y_m \cdot f(s, t, v) = \lambda^m \left( v f(s - m, t, v) - m t \frac{\partial f(s - m, t, v)}{\partial v} \right),$$

where  $m \in \mathbb{Z}$  and  $f(s, t, v) \in \mathbb{C}[s, t, v]$ .

**PROPOSITION 2.4.** For  $\lambda \in \mathbb{C}^*$  and  $\alpha \in \mathbb{C}$ ,  $\Phi(\lambda, \alpha)$  is a class of free modules of rank 1 over the algebra  $\mathfrak{sv}(0)$  under the action defined by (2.6)-(2.8).

**PROOF.** For  $m \in \mathbb{Z}$  and  $n \in \mathbb{Z}$ , according to (2.6), one has the following

$$\begin{aligned}
&L_n \cdot L_m \cdot f(s, t, v) \\
&= L_n \cdot \left[ \lambda^m \left( (s + m\alpha) f(s - m, t, v) - \frac{m}{2} v \frac{\partial f(s - m, t, v)}{\partial v} + \frac{m^2}{4} t \frac{\partial^2 f(s - m, t, v)}{\partial v^2} \right) \right] \\
&= \lambda^{m+n} \left\{ (s + n\alpha)(s - n + m\alpha) f(s - m - n, t, v) + \left[ -\frac{n}{2} v (s - n + m\alpha) \right. \right. \\
&\quad \left. \left. - \frac{m}{2} (s + n\alpha) v + \frac{mn}{4} v \right] \frac{\partial f(s - m - n, t, v)}{\partial v} + \left[ \frac{n^2}{4} t (s - n + m\alpha) \right. \right. \\
&\quad \left. \left. + \frac{mn}{4} v^2 - \frac{mn^2}{4} t + \frac{m^2}{4} (s + n\alpha) t \right] \frac{\partial^2 f(s - m - n, t, v)}{\partial v^2} \right. \\
&\quad \left. + \left[ -\frac{mn^2}{8} vt - \frac{m^2 n}{8} vt \right] \frac{\partial^3 f(s - m - n, t, v)}{\partial v^3} \right. \\
&\quad \left. + \frac{m^2 n^2}{16} t^2 \frac{\partial^4 f(s - m - n, t, v)}{\partial v^4} \right\},
\end{aligned}$$

which implies that

$$\begin{aligned}
& L_n \cdot L_m \cdot f(s, t, v) - L_m \cdot L_n \cdot f(s, t, v) \\
&= \lambda^{m+n} \left\{ (m-n) \left[ (s + (m+n)\alpha) f(s-m-n, t, v) - \frac{(m+n)}{2} v \frac{\partial f(s-m-n, t, v)}{\partial v} \right. \right. \\
&\quad \left. \left. + \frac{(m+n)^2}{4} t \frac{\partial^2 f(s-m-n, t, v)}{\partial v^2} \right] \right\} \\
&= (m-n) L_{m+n} \cdot f(s, t, v) \\
&= [L_n, L_m] \cdot f(s, t, v).
\end{aligned}$$

By (2.6) and (2.8), one easily gets

$$\begin{aligned}
& L_n \cdot Y_m \cdot f(s, t, v) \\
&= L_n \cdot \left[ \lambda^m \left( v f(s-m, t, v) - m t \frac{\partial f(s-m, t, v)}{\partial v} \right) \right] \\
&= \lambda^{m+n} \left\{ \left( (s+n\alpha)v - \frac{n}{2}v \right) f(s-m-n, t, v) + \left( -\frac{n}{2}v^2 + \frac{n^2}{2}t - m(s+n\alpha)t \right) \right. \\
&\quad \frac{\partial f(s-m-n, t, v)}{\partial v} + \left( \frac{n^2}{4}vt + \frac{mn}{2}vt \right) \frac{\partial^2 f(s-m-n, t, v)}{\partial v^2} \\
(2.9) \quad & \left. - \frac{mn^2}{4}t^2 \frac{\partial^3 f(s-m-n, t, v)}{\partial v^3} \right\}.
\end{aligned}$$

Similarly, it immediately holds that

$$\begin{aligned}
& Y_m \cdot L_n \cdot f(s, t, v) \\
&= Y_m \cdot \left[ \lambda^n \left( (s+n\alpha) f(s-n, t, v) - \frac{n}{2} v \frac{\partial f(s-n, t, v)}{\partial v} + \frac{n^2}{4} t \frac{\partial^2 f(s-n, t, v)}{\partial v^2} \right) \right] \\
&= \lambda^{m+n} \left\{ v(s-m+n\alpha) f(s-m-n, t, v) + \left( -mt(s-m+n\alpha) - \frac{n}{2}v^2 + \frac{n}{2}mt \right) \right. \\
&\quad \frac{\partial f(s-m-n, t, v)}{\partial v} + \left( \frac{n}{2}mtv + \frac{n^2}{4}vt \right) \frac{\partial^2 f(s-m-n, t, v)}{\partial v^2} \\
(2.10) \quad & \left. - \frac{n^2}{4}mt^2 \frac{\partial^3 f(s-m-n, t, v)}{\partial v^3} \right\}.
\end{aligned}$$

Subtracting (2.10) from (2.9) gives rise to

$$\begin{aligned}
& L_n \cdot Y_m \cdot f(s, t, v) - Y_m \cdot L_n \cdot f(s, t, v) \\
&= \lambda^{m+n} \left\{ \left( m - \frac{n}{2} \right) \left[ v f(s-m-n, t, v) - (m+n)t \frac{\partial f(s-m-n, t, v)}{\partial v} \right] \right\} \\
&= \left( m - \frac{n}{2} \right) Y_{m+n} \cdot f(s, t, v) \\
&= [L_n, Y_m] \cdot f(s, t, v).
\end{aligned}$$

Using (2.6) and (2.7), one obtains

$$\begin{aligned}
& L_n \cdot M_m \cdot f(s, t, v) \\
&= L_n \cdot \left[ \lambda^m t f(s-m, t, v) \right] \\
&= \lambda^{m+n} \left\{ (s+n\alpha) t f(s-m-n, t, v) - \frac{n}{2} v t \frac{\partial f(s-m-n, t, v)}{\partial v} \right. \\
(2.11) \quad & \left. + \frac{n^2}{4} t^2 \frac{\partial^2 f(s-m-n, t, v)}{\partial v^2} \right\}.
\end{aligned}$$

Similarly, it is easy to see that

$$\begin{aligned}
& M_m \cdot L_n \cdot f(s, t, v) \\
&= M_m \cdot \left[ \lambda^n \left( (s + n\alpha)f(s - n, t, v) - \frac{n}{2}v \frac{\partial f(s - n, t, v)}{\partial v} \right. \right. \\
&\quad \left. \left. + \frac{n^2}{4}t \frac{\partial^2 f(s - n, t, v)}{\partial v^2} \right) \right] \\
&= \lambda^{m+n} \left\{ t(s - m + n\alpha)f(s - m - n, t, v) - \frac{n}{2}tv \frac{\partial f(s - m - n, t, v)}{\partial v} \right. \\
(2.12) \quad &\quad \left. + \frac{n^2}{4}t^2 \frac{\partial^2 f(s - m - n, t, v)}{\partial v^2} \right\}.
\end{aligned}$$

Considering (2.11) and (2.12), we have

$$\begin{aligned}
& L_n \cdot M_m \cdot f(s, t, v) - M_m \cdot L_n \cdot f(s, t, v) \\
&= \lambda^{m+n} m t f(s - m - n, t, v) \\
&= m M_{m+n} \cdot f(s, t, v) \\
&= [L_n, M_m] \cdot f(s, t, v).
\end{aligned}$$

It follows from (2.8)

$$\begin{aligned}
& Y_n \cdot Y_m \cdot f(s, t, v) \\
&= Y_n \cdot \left[ \lambda^m \left( v f(s - m, t, v) - m t \frac{\partial f(s - m, t, v)}{\partial v} \right) \right] \\
&= \lambda^{m+n} \left\{ (v^2 - m t) f(s - m - n, t, v) + m n t \frac{\partial^2 f(s - m, t, v)}{\partial v^2} \right\},
\end{aligned}$$

which implies that

$$\begin{aligned}
& Y_n \cdot Y_m \cdot f(s, t, v) - Y_m \cdot Y_n \cdot f(s, t, v) \\
&= \lambda^{m+n} (v^2 - m t - v^2 + n t) f(s - m - n, t, v) \\
&= (n - m) M_{m+n} \cdot f(s, t, v) \\
&= [Y_n, Y_m] \cdot f(s, t, v).
\end{aligned}$$

Finally, due to (2.8) and (2.7), it is straightforward to verify that

$$\begin{aligned}
& Y_n \cdot M_m \cdot f(s, t, v) - M_m \cdot Y_n \cdot f(s, t, v) \\
&= Y_n \cdot \left[ \lambda^m t f(s - m, t, v) \right] - M_m \cdot \left[ \lambda^n \left( v f(s - n, t, v) - n t \frac{\partial f(s - n, t, v)}{\partial v} \right) \right] \\
&= 0 = [Y_n, M_m] \cdot f(s, t, v)
\end{aligned}$$

and

$$\begin{aligned}
& M_n \cdot M_m \cdot f(s, t, v) - M_m \cdot M_n \cdot f(s, t, v) \\
&= M_n \cdot (\lambda^m t f(s - m, t, v)) - M_m \cdot (\lambda^n t f(s - n, t, v)) \\
&= 0 = [M_n, M_m] \cdot f(s, t, v).
\end{aligned}$$

This completes the proof.  $\square$

**REMARK 2.5.** For  $\lambda \in \mathbb{C}^*$  and  $\alpha \in \mathbb{C}$ ,  $\Phi(\lambda, \alpha)$  is reducible over the Schrödinger-Virasoro algebra  $\mathfrak{sv}(s)$  for  $s = 0$  as a consequence of Definition 2.3. In fact, it is effortless to get that  $t^i \mathbb{C}[s, t, v]$  is a submodule of  $\Phi(\lambda, \alpha)$  for  $i \in \mathbb{Z}$ .

### 3. Modules over $\mathfrak{sv}(0)$

In this section, we classify the free  $U(\mathbb{C}L_0 \oplus \mathbb{C}M_0 \oplus \mathbb{C}Y_0)$ -modules of rank 1 over the algebra  $\mathfrak{sv}(0)$ . Indeed, we get the following main result.

**THEOREM 3.1.** *Let  $\mathfrak{sv}(0)$  be the Schrödinger-Virasoro algebra  $\mathfrak{sv}(s)$  for  $s = 0$ , if there exists a free  $U(\mathbb{C}L_0 \oplus \mathbb{C}M_0 \oplus \mathbb{C}Y_0)$ -module of rank 1 over the algebra  $\mathfrak{sv}(0)$ , denoted by  $M$ , then*

$$M \cong \Phi(\lambda, \alpha)$$

for some  $\lambda \in \mathbb{C}^*$  and  $\alpha \in \mathbb{C}$ .

In order to prove Theorem 3.1, we first show several important lemmas.

**LEMMA 3.2.** *Let  $M$  be a free  $U(\mathbb{C}L_0 \oplus \mathbb{C}M_0 \oplus \mathbb{C}Y_0)$ -module of rank 1 over the algebra  $\mathfrak{sv}(0)$ . For  $m \in \mathbb{Z}$ , assume that*

$$L_m \cdot 1 = g_m(L_0, M_0, Y_0),$$

$$M_m \cdot 1 = a_m(L_0, M_0, Y_0),$$

$$Y_m \cdot 1 = p_m(L_0, M_0, Y_0),$$

in which  $g_m(L_0, M_0, Y_0), a_m(L_0, M_0, Y_0), p_m(L_0, M_0, Y_0) \in M$ , then  $g_m(L_0, M_0, Y_0), a_m(L_0, M_0, Y_0)$  and  $p_m(L_0, M_0, Y_0)$  completely determine the action of  $L_m, M_m$  and  $Y_m$  on  $M$ .

**PROOF.** Now take any  $u(L_0, M_0, Y_0) = \sum_{i,j,k \geq 0} a_{i,j,k} L_0^i M_0^j Y_0^k \in M$  and according to formulas (7)-(9) in Proposition 2.1, one has

$$\begin{aligned} L_m \cdot u(L_0, M_0, Y_0) &= L_m \cdot \sum_{i,j,k \geq 0} a_{i,j,k} L_0^i M_0^j Y_0^k \\ &= \sum_{i,j,k \geq 0} a_{i,j,k} (L_0 - m)^i M_0^j \left[ Y_0^k g_m(L_0, M_0, Y_0) - \frac{mk}{2} Y_0^{k-1} p_m(L_0, M_0, Y_0) \right. \\ &\quad \left. + \frac{k(k-1)m^2}{4} Y_0^{k-2} a_m(L_0, M_0, Y_0) \right] \\ &= u(L_0 - m, M_0, Y_0) g_m(L_0, M_0, Y_0) - \frac{m}{2} \frac{\partial u(L_0 - m, M_0, Y_0)}{\partial Y_0} p_m(L_0, M_0, Y_0) \\ &\quad + \frac{m^2}{4} \frac{\partial^2 u(L_0 - m, M_0, Y_0)}{\partial Y_0^2} a_m(L_0, M_0, Y_0). \end{aligned}$$

By formulas (4)-(6) in Proposition 2.1, it is easy to deduce that

$$\begin{aligned} Y_m \cdot u(L_0, M_0, Y_0) &= Y_m \cdot \sum_{i,j,k \geq 0} a_{i,j,k} L_0^i M_0^j Y_0^k \\ &= \sum_{i,j,k \geq 0} a_{i,j,k} (L_0 - m)^i M_0^j \left[ Y_0^k p_m(L_0, M_0, Y_0) - mk Y_0^{k-1} a_m(L_0, M_0, Y_0) \right] \\ &= u(L_0 - m, M_0, Y_0) p_m(L_0, M_0, Y_0) - m \frac{\partial u(L_0 - m, M_0, Y_0)}{\partial Y_0} a_m(L_0, M_0, Y_0). \end{aligned}$$

Using formulas (1)-(3) in Proposition 2.1, one gets

$$\begin{aligned}
M_m \cdot u(L_0, M_0, Y_0) &= M_m \cdot \sum_{i,j,k \geq 0} a_{i,j,k} L_0^i M_0^j Y_0^k \\
&= \sum_{i,j,k \geq 0} a_{i,j,k} (L_0 - m)^i M_0^j Y_0^k a_m(L_0, M_0, Y_0) \\
(3.1) \quad &= u(L_0 - m, M_0, Y_0) a_m(L_0, M_0, Y_0).
\end{aligned}$$

As a consequence, Lemma 3.2 holds.  $\square$

LEMMA 3.3.  $a_m(L_0, M_0, Y_0) \neq 0$  for all  $m \in \mathbb{Z}$ .

**PROOF.** Suppose that  $a_m(L_0, M_0, Y_0) = 0$  for some  $m \in \mathbb{Z}^*$ , then  $M_m \cdot M = 0$  owing to (3.1). Now according to relation (2.3), one has

$$mM_0 \cdot M = [L_{-m}, M_m] \cdot M = 0.$$

It suggests that  $M_0 \cdot M = 0$ , hence  $a_0(L_0, M_0, Y_0) = 0$ , which is a contradiction because of  $a_0(L_0, M_0, Y_0) = M_0$ . Therefore, Lemma 3.3 holds.  $\square$

LEMMA 3.4. For all  $m \in \mathbb{Z}$ , we have  $a_m(L_0, M_0, Y_0) \in \mathbb{C}[Y_0, M_0]$ .

**PROOF.** For  $m \in \mathbb{Z}$ , now assume that  $a_m(L_0, M_0, Y_0) = \sum_{i=0}^{k_m} b_{m,i} L_0^i$ , in which  $b_{m,i} \in \mathbb{C}[Y_0, M_0]$ ,  $k_m \in \mathbb{N}$  and  $b_{m,k_m} \neq 0$ . Due to relation (2.5) and Proposition 2.1, one obtains the following

$$\begin{aligned}
0 &= [M_n, M_m] \cdot 1 \\
&= \sum_{i=0}^{k_m} b_{m,i} (L_0 - n)^i \sum_{j=0}^{k_n} b_{n,j} L_0^j - \sum_{j=0}^{k_n} b_{n,j} (L_0 - m)^j \sum_{i=0}^{k_m} b_{m,i} L_0^i \\
&\equiv b_{m,k_m} b_{n,k_n} (mk_n - nk_m) L_0^{k_m+k_n-1} \pmod{\bigoplus_{i=0}^{k_m+k_n-2} \mathbb{C}[Y_0, M_0] L_0^i},
\end{aligned}$$

yielding that  $mk_n - nk_m = 0$ , thus  $k_m = mk_1$ . If  $k_1 > 0$ , then it is nonsense for  $k_m$  when  $m$  is negative, which forces  $k_1 = 0$ , hence  $k_m = 0$  for  $m \in \mathbb{Z}$ , therefore  $a_m(L_0, M_0, Y_0) \in \mathbb{C}[Y_0, M_0]$ .  $\square$

LEMMA 3.5. For  $m \in \mathbb{Z}$ , one has  $p_m(L_0, M_0, Y_0) \in \mathbb{C}[Y_0, M_0]$  and  $a_m(L_0, M_0, Y_0) \in \mathbb{C}[M_0]$ .

**PROOF.** For all  $m \in \mathbb{Z}$ , according to Lemma 3.4 assume that  $a_m(L_0, M_0, Y_0) = \sum_{i=0}^{h_m} l_{m,i} Y_0^i$ , where  $l_{m,i} \in \mathbb{C}[M_0]$ ,  $h_m \in \mathbb{N}$ ,  $l_{m,h_m} \neq 0$ . Moreover, let  $p_m(L_0, M_0, Y_0) = \sum_{i=0}^{t_m} d_{m,i} L_0^i$ , in which  $d_{m,i} \in \mathbb{C}[Y_0, M_0]$ ,  $t_m \in \mathbb{N}$  and  $d_{m,t_m} \neq 0$ . Applying relation (2.5) and Proposition 2.1, one has

$$\begin{aligned}
0 &= [Y_n, M_m] \cdot 1 \\
&= \sum_{i=0}^{h_m} l_{m,i} \left( Y_0^i \sum_{j=0}^{t_n} d_{n,j} L_0^j - ni Y_0^{i-1} \sum_{j=0}^{h_n} l_{n,j} Y_0^j \right)
\end{aligned}$$



$$\begin{aligned}
& - \sum_{j=0}^{t_n} d_{n,j} (L_0 - m)^j \sum_{i=0}^{h_m} l_{m,i} Y_0^i \\
& = \sum_{i=0}^{h_m} \sum_{j=0}^{t_n} l_{m,i} d_{n,j} L_0^j Y_0^i - \sum_{i=1}^{h_m} \sum_{j=0}^{h_n} l_{m,i} l_{n,j} n^i Y_0^{i-1} Y_0^j \\
(3.2) \quad & - \sum_{i=0}^{h_m} \sum_{j=0}^{t_n} l_{m,i} d_{n,j} (L_0 - m)^j Y_0^i.
\end{aligned}$$

Observing all terms of  $L_0$  in (3.2), one knows that  $t_n \leq 1$ , that is  $t_n = 0$  or  $1$ .

We claim that  $t_n = 0$  for all  $n \in \mathbb{Z}$ . In fact, suppose  $t_k = 1$  for  $k \in \mathbb{Z}^*$ , then (3.2) becomes

$$0 = m \sum_{i=0}^{h_m} l_{m,i} d_{k,1} Y_0^i - k \sum_{i=1}^{h_m} \sum_{j=0}^{h_k} l_{m,i} l_{k,j} i Y_0^{i-1} Y_0^j,$$

taking  $m = k$  and  $m = -k$  respectively, one has

$$\begin{aligned}
0 & = k \sum_{i=0}^{h_k} l_{k,i} d_{k,1} Y_0^i - k \sum_{i=1}^{h_k} \sum_{j=0}^{h_k} l_{k,i} l_{k,j} i Y_0^{i-1} Y_0^j \\
& = k \sum_{i=0}^{h_k} l_{k,i} d_{k,1} Y_0^i - k \sum_{j=1}^{h_k} \sum_{i=0}^{h_k} l_{k,j} l_{k,i} j Y_0^{j-1} Y_0^i \\
& = k \sum_{i=0}^{h_k} l_{k,i} Y_0^i \left( d_{k,1} - \sum_{j=1}^{h_k} l_{k,j} j Y_0^{j-1} \right)
\end{aligned}$$

and

$$\begin{aligned}
0 & = -k \sum_{i=0}^{h_{-k}} l_{-k,i} d_{k,1} Y_0^i - k \sum_{i=1}^{h_{-k}} \sum_{j=0}^{h_k} l_{-k,i} l_{k,j} i Y_0^{i-1} Y_0^j \\
& = -k \sum_{i=0}^{h_{-k}} l_{-k,i} \left( d_{k,1} Y_0 + \sum_{j=0}^{h_k} l_{k,j} j Y_0^j \right) Y_0^{i-1},
\end{aligned}$$

thus  $d_{k,1} = \sum_{j=1}^{h_k} l_{k,j} j Y_0^{j-1}$  and  $d_{k,1} Y_0 + \sum_{j=0}^{h_k} l_{k,j} h_{-k} Y_0^j = 0$  which suggests  $l_{k,0} h_{-k} = 0$ ,

thereby  $d_{k,1} = \sum_{j=1}^{h_k} l_{k,j} j Y_0^{j-1} = - \sum_{j=1}^{h_k} l_{k,j} h_{-k} Y_0^{j-1} \neq 0$ , which is impossible. Therefore, we get  $t_n = 0$  for  $n \in \mathbb{Z}$ .

In addition, for  $n \neq 0$  and  $t_n = 0$ , it is easy to find that (3.2) becomes

$$0 = -n \sum_{i=0}^{h_m} \sum_{j=0}^{h_n} l_{m,i} l_{n,j} i Y_0^{i-1} Y_0^j,$$

whence  $h_m = 0$  for all  $m \in \mathbb{Z}$ , then  $a_m(L_0, M_0, Y_0) \in \mathbb{C}[M_0]$ .  $\square$

LEMMA 3.6.  $\deg_{Y_0} p_m(L_0, M_0, Y_0) = 1$  for all  $m \in \mathbb{Z}$ .

**PROOF.** For  $m \in \mathbb{Z}$  assume that  $p_m(L_0, M_0, Y_0) = \sum_{i=0}^{f_m} q_{m,i} Y_0^i$ ,  $a_m(L_0, M_0, Y_0) = \sum_{i=0}^{e_m} s_{m,i} M_0^i$ , in which  $q_{m,i} \in \mathbb{C}[M_0]$ ,  $s_{m,i} \in \mathbb{C}$  and  $q_{m,f_m} \neq 0$ ,  $s_{m,e_m} \neq 0$ ,  $f_m \in \mathbb{N}$ ,

$e_m \in \mathbb{N}$ . Due to relation (2.4), one easily finds

$$(3.3) \quad [Y_n, Y_m] \cdot 1 = (m - n)M_{m+n} \cdot 1.$$

By straightforward calculations, the left hand side of (3.3) is

$$\begin{aligned} & Y_n Y_m \cdot 1 - Y_m Y_n \cdot 1 \\ &= \sum_{i=0}^{f_m} q_{m,i} (Y_0^i Y_n \cdot 1 - n i Y_0^{i-1} M_n \cdot 1) - \sum_{j=0}^{f_n} q_{n,j} (Y_0^j Y_m \cdot 1 - m j Y_0^{j-1} M_m \cdot 1) \\ &= \sum_{i=0}^{f_m} q_{m,i} \left( Y_0^i \sum_{j=0}^{f_n} q_{n,j} Y_0^j - n i Y_0^{i-1} \sum_{k=0}^{e_n} s_{n,k} M_0^k \right) \\ &\quad - \sum_{j=0}^{f_n} q_{n,j} \left( Y_0^j \sum_{i=0}^{f_m} q_{m,i} Y_0^i - m j Y_0^{j-1} \sum_{k=0}^{e_m} s_{m,k} M_0^k \right) \\ &= \sum_{j=0}^{f_n} \sum_{k=0}^{e_m} s_{m,k} q_{n,j} m j Y_0^{j-1} M_0^k - \sum_{i=0}^{f_m} \sum_{k=0}^{e_n} s_{n,k} q_{m,i} n i Y_0^{i-1} M_0^k. \end{aligned}$$

While the right hand side of (3.3) does not contain the terms of  $Y_0$  and is not zero from Lemma 3.5 and Lemma 3.3, thus we have  $f_n = f_m \neq 0$ , thereby one directly obtains  $f_m = f_0 = 1$ .  $\square$

**LEMMA 3.7.** *For all  $m \in \mathbb{Z}$ , one has  $\deg_{L_0} g_m(L_0, M_0, Y_0) = 1$  and  $g_m(L_0, M_0, Y_0) \in \mathbb{C}[L_0, M_0]$ .*

**PROOF.** Let  $g_n(L_0, M_0, Y_0) = \sum_{i=0}^{r_n} c_{n,i} L_0^i$ , in which  $c_{n,i} \in \mathbb{C}[M_0, Y_0]$  and  $c_{n,r_n} \neq 0$ ,  $r_n \in \mathbb{N}$ . Thanks to Lemma 3.5, for  $m \in \mathbb{Z}$  assume that  $a_m(L_0, M_0, Y_0) = \sum_{i=0}^{e_m} s_{m,i} M_0^i$ , where  $s_{m,i} \in \mathbb{C}$  and  $s_{m,e_m} \neq 0$ ,  $e_m \in \mathbb{N}$ . Owing to relation (2.3), it is easy to see that

$$(3.4) \quad [L_n, M_m] \cdot 1 = m M_{m+n} \cdot 1.$$

Then the left hand side of (3.4) directly becomes

$$\begin{aligned} & L_n M_m \cdot 1 - M_m L_n \cdot 1 \\ &= \sum_{i=0}^{e_m} s_{m,i} M_0^i \sum_{j=0}^{r_n} c_{n,j} L_0^j - \sum_{j=0}^{r_n} c_{n,j} (L_0 - m)^j \sum_{i=0}^{e_m} s_{m,i} M_0^i \\ (3.5) \quad &= - \sum_{i=0}^{e_m} \sum_{j=1}^{r_n} s_{m,i} c_{n,j} M_0^i \left( -j m L_0^{j-1} + C_j^2 (-m)^2 L_0^{j-2} + \cdots + C_j^j (-m)^j \right). \end{aligned}$$

Note that the right hand side of (3.4) does not contain the terms of  $L_0$ , thereby one deduces that  $r_n \leq 1$ . If  $r_k = 0$  for some  $k \in \mathbb{Z}^*$ , then the left hand side of (3.4) vanishes, however the right hand side of (3.4) is not equal to zero if  $m \neq 0$  as a result of Lemma 3.3, hence  $r_n = 1$  for  $n \in \mathbb{Z}$ .

In addition, it is clearly that the right hand side of (3.4) does not contain the terms of  $Y_0$  according to Lemma 3.5, hence from (3.5) one knows  $c_{n,1} \in \mathbb{C}[M_0]$ .

Next suppose that  $g_n(L_0, M_0, Y_0) = c_{n,1}L_0 + \sum_{i=0}^{\lambda_n} \alpha_{n,i}Y_0^i$ , in which  $c_{n,1} \neq 0$  and  $\alpha_{n,i} \in \mathbb{C}[M_0]$ ,  $\alpha_{n,\lambda_n} \neq 0$ ,  $\lambda_n \in \mathbb{N}$ . On the one hand,

$$\begin{aligned}
& L_n L_m \cdot 1 - L_m L_n \cdot 1 \\
= & L_n(c_{m,1}L_0 + \sum_{i=0}^{\lambda_m} \alpha_{m,i}Y_0^i) - L_m(c_{n,1}L_0 + \sum_{i=0}^{\lambda_n} \alpha_{n,i}Y_0^i) \\
= & -\sum_{i=0}^{\lambda_n} \alpha_{n,i}(Y_0^i L_m \cdot 1 - \frac{mi}{2}Y_0^{i-1}Y_m \cdot 1 + \frac{m^2i(i-1)}{4}Y_0^{i-2}M_m \cdot 1) \\
& + \sum_{i=0}^{\lambda_m} \alpha_{m,i}(Y_0^i L_n \cdot 1 - \frac{ni}{2}Y_0^{i-1}Y_n \cdot 1 + \frac{n^2i(i-1)}{4}Y_0^{i-2}M_n \cdot 1) \\
& + c_{m,1}(L_0 - n)L_n \cdot 1 - c_{n,1}(L_0 - m)L_m \cdot 1 \\
= & mc_{n,1} \sum_{i=0}^{\lambda_m} \alpha_{m,i}Y_0^i - nc_{m,1} \sum_{i=0}^{\lambda_n} \alpha_{n,i}Y_0^i - \sum_{i=0}^{\lambda_m} \alpha_{m,i} \frac{ni}{2}Y_0^{i-1}Y_n \cdot 1 \\
& + \sum_{i=0}^{\lambda_n} \alpha_{n,i} \frac{mi}{2}Y_0^{i-1}Y_m \cdot 1 + \sum_{i=0}^{\lambda_m} \alpha_{m,i} \frac{n^2i(i-1)}{4}Y_0^{i-2}M_n \cdot 1 \\
& - \sum_{i=0}^{\lambda_n} \alpha_{n,i} \frac{m^2i(i-1)}{4}Y_0^{i-2}M_m \cdot 1 + (m-n)c_{m,1}c_{n,1}L_0,
\end{aligned}$$

on the other hand,

$$\begin{aligned}
& (m-n)L_{m+n} \cdot 1 \\
= & (m-n)(c_{m+n,1}L_0 + \sum_{i=0}^{\lambda_{m+n}} \alpha_{m+n,i}Y_0^i).
\end{aligned}$$

According to relation (2.1), it is obvious that  $\lambda_m = \lambda_n = \lambda_{m+n} = \lambda_0 = 0$ , that is,  $g_m(L_0, M_0, Y_0) \in \mathbb{C}[L_0, M_0]$ .  $\square$

From what has been discussed above, now we turn to the proof of Theorem 3.1.

### The Proof of Theorem 3.1.

Lemma 3.2 suggests that  $g_m(L_0, M_0, Y_0)$ ,  $a_m(L_0, M_0, Y_0)$  and  $p_m(L_0, M_0, Y_0)$  completely determine the action of  $L_m$ ,  $M_m$  and  $Y_m$  on  $M$ . Thus the next task is just to determine  $g_m(L_0, M_0, Y_0)$ ,  $a_m(L_0, M_0, Y_0)$ , and  $p_m(L_0, M_0, Y_0)$  for all  $m \in \mathbb{Z}$ .

According to Lemma 3.5, Lemma 3.7 and Lemma 3.6, here we need to rewrite the preceding assumptions. For  $m \in \mathbb{Z}$ , suppose that

$$(3.6) \quad g_m(L_0, M_0, Y_0) = a_m + b_m L_0,$$

$$(3.7) \quad p_m(L_0, M_0, Y_0) = c_m + d_m Y_0,$$

in which  $a_m, b_m, c_m, d_m \in \mathbb{C}[M_0]$  and  $b_m \neq 0, d_m \neq 0$ . In particular,  $a_0 = c_0 = 0$  and  $b_0 = d_0 = 1$ . Thanks to relation (2.1), one gets directly

$$(3.8) \quad [L_n, L_m] \cdot 1 = (m-n)L_{m+n} \cdot 1.$$

Substituting (3.6) into (3.8), one immediately finds that the left hand side of (3.8) becomes

$$L_n L_m \cdot 1 - L_m L_n \cdot 1$$

$$\begin{aligned}
&= L_n(a_m + b_m L_0) - L_m(a_n + b_n L_0) \\
(3.9) \quad &= ma_m b_n - na_n b_m + (m - n)b_n b_m L_0
\end{aligned}$$

and the right hand side of (3.8) becomes

$$\begin{aligned}
(3.10) \quad &(m - n)L_{m+n} \cdot 1 \\
&= (m - n)(a_{m+n} + b_{m+n} L_0).
\end{aligned}$$

Comparing the coefficients of  $L_0$  of (3.9) and (3.10), one has  $b_n b_m = b_{m+n}$ , yielding that  $b_m = \lambda^m (\lambda := b_1 \in \mathbb{C}^*)$  for all  $m \in \mathbb{Z}$ . Moreover, observing the constant term in (3.9) and (3.10), one obtains that  $ma_m b_n - na_n b_m = (m - n)a_{m+n}$ . Choosing  $m = 1$  in the formula gives  $\lambda^n a_1 - n\lambda a_n = (1 - n)a_{n+1}$ , from which we have  $a_m = m\lambda^m \alpha$  by induction on  $n$ , where  $\alpha = \frac{a_1}{\lambda}$  and  $a_1 \in \mathbb{C}$ . Therefore,  $g_m(L_0, M_0, Y_0) = \lambda^m(L_0 + m\alpha)$ .

Similarly, according to relation (2.2), it is effortless to see that

$$(3.11) \quad [L_n, Y_m] \cdot 1 = (m - \frac{n}{2})Y_{m+n} \cdot 1.$$

Thanks to (3.6) and (3.7) it is easy to deduce that the left hand side of (3.11) is

$$\begin{aligned}
&L_n Y_m \cdot 1 - Y_m L_n \cdot 1 \\
&= L_n(c_m + d_m Y_0) - Y_m(a_n + b_n L_0) \\
(3.12) \quad &= mc_m b_n - \frac{n}{2}d_m c_n + mb_n d_m Y_0 - \frac{n}{2}d_m d_n Y_0
\end{aligned}$$

and applying (3.7) one finds that the right hand side of (3.11) becomes

$$\begin{aligned}
(3.13) \quad &(m - \frac{n}{2})Y_{m+n} \cdot 1 \\
&= (m - \frac{n}{2})(c_{m+n} + d_{m+n} Y_0).
\end{aligned}$$

Hence  $mc_m b_n - \frac{n}{2}d_m c_n = (m - \frac{n}{2})c_{m+n}$  and  $mb_n d_m - \frac{n}{2}d_m d_n = (m - \frac{n}{2})d_{m+n}$ . Taking  $m = 1$  in the latter formula one gets  $\lambda^n d_1 - \frac{n}{2}d_1 d_n = (1 - \frac{n}{2})d_{n+1}$ , which implies that  $d_m = \lambda^m$  for  $m \in \mathbb{Z}$ . Besides, choosing  $m = \pm 1$  in the former formula gives

$$(3.14) \quad \lambda^n c_1 - \frac{n}{2}\lambda c_n = (1 - \frac{n}{2})c_{n+1}$$

and

$$(3.15) \quad -c_{-1}\lambda^n - \frac{n}{2}\lambda^{-1}c_n = (-1 - \frac{n}{2})c_{n-1}.$$

Setting  $n = \pm 1$  in (3.14), one deduces that  $c_2 = \lambda c_1$  and  $c_{-1} = -2\lambda^{-2}c_1$ . In addition, taking  $n = 2$  in (3.15), one easily has

$$-c_{-1}\lambda^2 - \lambda^{-1}c_2 = -2c_1.$$

It yields that  $c_2 = 4\lambda c_1$ , whence  $c_1 = 0$ . Thus from (3.14) or (3.15), it is easy to find that  $c_m = 0$  for  $m \in \mathbb{Z}$ . Therefore  $p_m(L_0, M_0, Y_0) = \lambda^m Y_0$ .

Now it remains to consider how to determine  $a_m(L_0, M_0, Y_0)$ . For  $m \neq 0$ , continue to use the assumption  $a_m(L_0, M_0, Y_0) = \sum_{i=0}^{e_m} s_{m,i} M_0^i$ , in which  $s_{m,i} \in \mathbb{C}$  and  $s_{m,e_m} \neq 0$ . Due to relation (2.3), it is clearly to see that

$$(3.16) \quad [L_n, M_m] \cdot 1 = mM_{m+n} \cdot 1.$$

Now calculating the left hand side of (3.16), one always has

$$\begin{aligned} & L_n M_m \cdot 1 - M_m L_n \cdot 1 \\ &= \sum_{i=0}^{e_m} s_{m,i} \lambda^n (L_0 + n\alpha) M_0^i - \sum_{i=0}^{e_m} s_{m,i} \lambda^n (L_0 - m + n\alpha) M_0^i \\ &= m \sum_{i=0}^{e_m} s_{m,i} \lambda^n M_0^i, \end{aligned}$$

thus one immediately finds that

$$(3.17) \quad m \sum_{i=0}^{e_m} s_{m,i} \lambda^n M_0^i = m \sum_{i=0}^{e_{m+n}} s_{m+n,i} M_0^i.$$

Whence one knows  $e_m = e_{m+n} = e_0 = 1$  for all  $m \in \mathbb{Z}$ . Therefore (3.17) becomes

$$m \lambda^n (s_{m,0} + s_{m,1} M_0) = m (s_{m+n,0} + s_{m+n,1} M_0).$$

Hence one always has  $s_{m+n,0} = \lambda^n s_{m,0}$  and  $s_{m+n,1} = \lambda^n s_{m,1}$ , from which it is direct to obtain that  $s_{m,0} = \lambda^m s_{0,0} = 0$  and  $s_{m,1} = \lambda^m s_{0,1} = \lambda^m$  for all  $m \in \mathbb{Z}$ .

By putting everything together, it can be seen that

$$\begin{aligned} g_m(L_0, M_0, Y_0) &= \lambda^m (L_0 + m\alpha), \\ p_m(L_0, M_0, Y_0) &= \lambda^m Y_0, \\ a_m(L_0, M_0, Y_0) &= \lambda^m M_0, \end{aligned}$$

for  $\lambda \in \mathbb{C}^*$ ,  $\alpha \in \mathbb{C}$  and all  $m \in \mathbb{Z}$ . Thus according to Lemma 3.2, Theorem 3.1 holds.  $\square$

#### 4. Modules over $\mathfrak{sv}(\frac{1}{2})$

In this section, we focus on the Schrödinger-Virasoro algebra  $\mathfrak{sv}(s)$  for  $s = \frac{1}{2}$  and show that the class of free  $U(\mathbb{C}L_0 \oplus \mathbb{C}M_0)$ -modules of rank 1 over the algebra  $\mathfrak{sv}(\frac{1}{2})$  is nonexistent.

The Schrödinger-Virasoro algebra  $\mathfrak{sv}(\frac{1}{2})$  is an infinite-dimensional Lie algebra over  $\mathbb{C}$  with a basis

$$\{L_n, M_n, Y_p \mid n \in \mathbb{Z}, p \in \mathbb{Z} + \frac{1}{2}\}$$

and has the following relations

$$(4.1) \quad [L_n, L_m] = (m - n)L_{m+n},$$

$$(4.2) \quad [L_n, Y_p] = (p - \frac{n}{2})Y_{p+n},$$

$$(4.3) \quad [L_n, M_m] = mM_{m+n},$$

$$(4.4) \quad [Y_p, Y_q] = (q - p)M_{p+q},$$

$$(4.5) \quad [Y_p, M_m] = [M_n, M_m] = 0,$$

where  $m, n \in \mathbb{Z}, p, q \in \mathbb{Z} + \frac{1}{2}$ .

From the above relations (4.1)-(4.5), there are some useful formulas that are similar to Proposition 2.1.

**PROPOSITION 4.1.** For  $i \in \mathbb{Z}, n \in \mathbb{Z}, p \in \mathbb{Z} + \frac{1}{2}$  and  $i \geq 0$ , the following formulas hold:

- (1)  $M_n L_0^i = (L_0 - n)^i M_n.$
- (2)  $M_n M_0^i = M_0^i M_n.$
- (3)  $Y_p L_0^i = (L_0 - p)^i Y_p.$
- (4)  $Y_p M_0^i = M_0^i Y_p.$
- (5)  $L_n L_0^i = (L_0 - n)^i L_n.$
- (6)  $L_n M_0^i = M_0^i L_n.$

It is straightforward to check these formulas by induction on  $i$  for all  $i \geq 0$ .

**THEOREM 4.2.** Let  $\mathfrak{sv}(\frac{1}{2})$  be the Schrödinger-Virasoro algebra  $\mathfrak{sv}(s)$  for  $s = \frac{1}{2}$ , denote by  $\mathcal{M}$  the set of all free  $U(\mathbb{C}L_0 \oplus \mathbb{C}M_0)$ -modules of rank 1 over  $\mathfrak{sv}(\frac{1}{2})$ , then

$$\mathcal{M} = \emptyset.$$

**PROOF.** Assume on the contrary that  $\mathcal{M} \neq \emptyset$  and let  $N \in \mathcal{M}$ , so  $N = U(\mathbb{C}L_0 \oplus \mathbb{C}M_0)$ . Take any  $w(L_0, M_0) = \sum_{i,j \geq 0} a_{i,j} L_0^i M_0^j \in N$  and suppose that

$$L_m \cdot 1 = g_m(L_0, M_0),$$

$$M_m \cdot 1 = a_m(L_0, M_0),$$

$$Y_p \cdot 1 = h_p(L_0, M_0),$$

where  $m \in \mathbb{Z}, p \in \mathbb{Z} + \frac{1}{2}$  and  $g_m(L_0, M_0), a_m(L_0, M_0), h_p(L_0, M_0) \in N$ .

Next we will prove that  $g_m(L_0, M_0), a_m(L_0, M_0)$  and  $h_p(L_0, M_0)$  completely determine the action of  $L_m, M_m$  and  $Y_p$  on  $N$ .

It follows from formulas (5)-(6) in Proposition 4.1

$$\begin{aligned} L_m \cdot w(L_0, M_0) &= L_m \cdot \left( \sum_{i,j \geq 0} a_{i,j} L_0^i M_0^j \right) \\ &= \sum_{i,j,k \geq 0} a_{i,j,k} (L_0 - m)^i M_0^j g_m(L_0, M_0) \\ (4.6) \quad &= w(L_0 - m, M_0) g_m(L_0, M_0). \end{aligned}$$

According to formulas (3)-(4) in Proposition 4.1, one always has

$$\begin{aligned} Y_p \cdot w(L_0, M_0) &= Y_p \cdot \left( \sum_{i,j \geq 0} a_{i,j} L_0^i M_0^j \right) \\ &= \sum_{i,j \geq 0} a_{i,j} (L_0 - p)^i M_0^j h_p(L_0, M_0) \\ (4.7) \quad &= w(L_0 - p, M_0) h_p(L_0, M_0). \end{aligned}$$

By formulas (1)-(2) in Proposition 4.1, it is easy to obtain that

$$M_m \cdot w(L_0, M_0) = M_m \cdot \left( \sum_{i,j \geq 0} a_{i,j} L_0^i M_0^j \right)$$

$$\begin{aligned}
&= \sum_{i,j \geq 0} a_{i,j}(L_0 - m)^i M_0^j a_m(L_0, M_0) \\
(4.8) \quad &= w(L_0 - m, M_0) a_m(L_0, M_0).
\end{aligned}$$

The above three formulas (4.6)-(4.8) suggest that one only needs to consider how to determine  $g_m(L_0, M_0)$ ,  $a_m(L_0, M_0)$ , and  $h_p(L_0, M_0)$  for  $m \in \mathbb{Z}$  and  $p \in \mathbb{Z} + \frac{1}{2}$  if one wants to know the action of  $L_m, M_m$  and  $Y_p$  on  $M$ . Whence we will discuss how to determine them in the next step.

For  $m \in \mathbb{Z}$ , assume that  $a_m(L_0, M_0) = \sum_{i=0}^{k'_m} b'_{m,i} L_0^i$ , in which  $b'_{m,i} \in \mathbb{C}[M_0]$ ,  $k'_m \in \mathbb{N}$  and  $b'_{m,k'_m} \neq 0$ . According to relation (4.5) and Proposition 4.1, one obtains  $k'_m = 0$  referring to the proof of Lemma 3.4, therefore  $a_m(L_0, M_0) \in \mathbb{C}[M_0]$ .

In addition, we claim that  $a_m(L_0, M_0) \neq 0$  for all  $m \in \mathbb{Z}$ . In fact, if there exists  $m \in \mathbb{Z}^*$  such that  $a_m(L_0, M_0) = 0$ , then  $M_m \cdot M = 0$  due to formula (4.8). According to relation (4.3), one immediately deduces that

$$mM_0 \cdot M = [L_{-m}, M_m] \cdot M = 0,$$

which implies that  $a_0(L_0, M_0) = 0$ . This is a contradiction.

For  $m \in \mathbb{Z}^*$  and  $p \in \mathbb{Z} + \frac{1}{2}$ , suppose that  $a_m(L_0, M_0) = \sum_{i=0}^{e'_m} s'_{m,i} M_0^i$  and  $h_p(L_0, M_0) = \sum_{i=0}^{t'_p} d'_{p,i} L_0^i$ , in which  $s'_{m,i} \in \mathbb{C}$ ,  $e'_m \in \mathbb{N}$ ,  $s'_{m,e'_m} \neq 0$  and  $d'_{p,i} \in \mathbb{C}[M_0]$ ,  $t'_p \in \mathbb{N}$ ,  $d'_{p,t'_p} \neq 0$ . Combining relation (4.5) with Proposition 4.1, one has

$$\begin{aligned}
0 &= [Y_p, M_m] \cdot 1 \\
&= \sum_{i=0}^{e'_m} s'_{m,i} M_0^i \sum_{j=0}^{t'_p} d'_{p,j} L_0^j - \sum_{j=0}^{t'_p} d'_{p,j} (L_0 - m)^j \sum_{i=0}^{e'_m} s'_{m,i} M_0^i \\
&= - \sum_{i=0}^{e'_m} \sum_{j=0}^{t'_p} s'_{m,i} d'_{p,j} M_0^i \left( -jm L_0^{j-1} + C_j^2 (-m)^2 L_0^{j-2} + \dots + C_j^j (-m)^j \right),
\end{aligned}$$

hence  $t'_p = 0$ , that is  $h_p(L_0, M_0) \in \mathbb{C}[M_0]$ . Thus from (4) in Proposition 4.1 and relation (4.4), we always have

$$[Y_p, Y_q] \cdot 1 = 0 \neq (q - p)M_{p+q}$$

for  $p \neq q$ , which is impossible.

From what has been discussed above, we draw a conclusion that the class of free  $U(\mathbb{C}L_0 \oplus \mathbb{C}M_0)$ -modules of rank 1 over the algebra  $\mathfrak{sv}(\frac{1}{2})$  is not existent, which completes the proof of Theorem 4.2.  $\square$

#### ACKNOWLEDGMENT

This work was supported by the NSFC (Grant No. 11871325 and 11726016).

## References

- [1] Batra P., Mazorchuk V., Block and modules for Whittaker pairs, *J. Pure Appl. Algebra* 215 (2011), 1552-1568.
- [2] Chen Q., Cai Y., Module over algebras related to the Virasoro algebra, *Internat. J. Math.* 26 (2015), 99-155.
- [3] Chen H., Guo X., Non-weight modules over the Heisenberg-Virasoro algebra and the  $W$  algebra  $W(2, 2)$ , *J. Algebra Appl.* 16 (2017), 1750097.
- [4] Chen H., Hong Y., Su Y., A family of new simple modules over the Schrödinger-Virasoro algebra, *J. Pure Appl. Algebra* 222 (2018), 900-913.
- [5] Cai Y., Tan H., Zhao K., Module structure on  $U(\mathfrak{h})$  for Kac-Moody algebras, arXiv:1606.01891v2.
- [6] Cai Y., Zhao K., Module structure on  $\mathcal{U}(H)$  for basic Lie superalgebras, *Toyama Math. J.* 37 (2015), 55-72.
- [7] Frenkel I., Zhu M., Vertex algebras associated to modified regular representations of the Virasoro algebra, *Adv. Math.* 229 (2012), 3468-3507.
- [8] Gao Y., Hu N., Liu D., Representations of the affine-Virasoro algebra of type  $A_1$ , *J. Geom. Phys.* 106 (2016), 102-107.
- [9] Guo X., Liu X., Wei Z., Irreducible weight modules for the Virasoro-like algebra and its  $q$ -analog, *Comm. Algebra* 45 (2017), 666-676.
- [10] Goddard P., Olive D., Kac-Moody and Virasoro algebras in relation to quantum physics, *Internat. J. Modern Phys. A* 1 (1986), 303-414.
- [11] Henkel M., Schrödinger invariance and strongly anisotropic critical systems, *J. Statist. Phys.* 75 (1994), 1023-1061.
- [12] Han J., Chen Q., Su Y., Modules over algebra  $Vir(a, b)$ , *Linear Algebra Appl.* 515 (2017), 11-23.
- [13] Henkel M., Unterberger J., Schrödinger invariance and space-time symmetries, *Nucl. Phys. B* 660 (2003), 407-435.
- [14] Jiang J., Lin W., Partial classification of cuspidal simple modules for Virasoro-like algebra, *J. Algebra* 464 (2016), 266-278.
- [15] Krohn M., Flohr M., Ghost systems revisited: modified Virasoro generators and logarithmic conformal field theories, *J. High Energy Phys.* 7 (2003), 431-449.
- [16] Li J., Su Y., Representations of the Schrödinger-Virasoro algebras, *J. Math. Phys.* 49 (2008), 053512.
- [17] Mertes N., Schnabl, M., String field representation of the Virasoro algebra, *J. High Energy Phys.* 12 (2016), 1-14.
- [18] Nilsson J., Simple  $\mathfrak{sl}_{n+1}$ -module structures on  $U(\mathfrak{h})$ , *J. Algebra* 424 (2015), 294-329.
- [19] Radobolja G., Application of vertex algebras to the structure theory of certain representations over the Virasoro algebra, *Algebr. Represent. Theory* 17 (2014), 1013-1034.
- [20] Wang D., Yu X., Biderivations and linear commuting maps on the Schrödinger-Virasoro Lie algebra, *Comm. Algebra* 41 (2013), 2166-2173.
- [21] Zhang X., Tan S., Unitary representations for the Schrödinger-Virasoro Lie algebra, *J. Algebra Appl.* 12 (2013), 1250132.
- [22] Zhang X., Tan S., Whittaker modules for the Schrödinger-Virasoro algebra, arXiv:0812.3245.

DEPARTMENT OF MATHEMATICS, SHANGHAI UNIVERSITY, SHANGHAI 200444, CHINA  
*E-mail address:* yumei.wang@shu.edu.cn

DEPARTMENT OF MATHEMATICS, SHANGHAI UNIVERSITY, SHANGHAI 200444, CHINA  
*E-mail address:* hlzhangmath@shu.edu.cn