

Sharp conditions for the existence of an even $[a, b]$ -factor in a graph

Eun-Kyung Cho* Jong Yoon Hyun† Suil O ‡
 Jeong Rye Park§

Abstract

Let a and b be positive integers. An even $[a, b]$ -factor of a graph G is a spanning subgraph H such that for every vertex $v \in V(G)$, $d_H(v)$ is even and $a \leq d_H(v) \leq b$. Matsuda conjectured that if G is an n -vertex 2-edge-connected graph such that $n \geq 2a + b + \frac{a^2-3a}{b} - 2$, $\delta(G) \geq a$, and $\sigma_2(G) \geq \frac{2an}{a+b}$, then G has an even $[a, b]$ -factor. In this paper, we provide counterexamples, which are highly connected. Furthermore, we give sharp sufficient conditions for a graph to have an even $[a, b]$ -factor. For even an , we conjecture a lower bound for $\lambda_1(G)$ in an n -vertex graph to have an $[a, b]$ -factor, where $\lambda_1(G)$ is the largest eigenvalue of G .

Keywords. Even $[a, b]$ -factor; edge-connectivity; vertex-connectivity; spectral radius.

AMS subject classifications. 05C70, 05C40, 05C50.

1 Introduction

Throughout all sections, a graph G is finite, simple, and undirected. We denote by $V(G)$ the set of vertices of G and by $E(G)$ the set of edges of G . For $S \subseteq V(G)$, we denote by $G - S$ the subgraph of G obtained from G by deleting the vertices in S together with the edges incident to vertices in S . For $S, T \subseteq V(G)$, we denote by $[S, T]$ the set of edges joining S and T . The *degree* of a vertex v in G , written $d_G(v)$ (or $d(v)$ if G is clear from the context), is the number of edges in $E(G)$ incident to the vertex v . The *minimum* and *maximum degree* of a graph G are denoted by $\delta(G)$ and $\Delta(G)$, respectively. For a subgraph H of G , the degree of a vertex v in H , written $d_H(v)$, is the number of edges in

*Department of Mathematics, Pusan National University, Busan, 46241, Republic of Korea; ekcho@pusan.ac.kr.

†Korea Institute for Advanced Study, Seoul, 02455, Republic of Korea; hyun33@kias.re.kr. Research supported by NRF-2017R1D1A1B05030707.

‡Applied Mathematics and Statistics, The State University of New York, Korea, Incheon, 21985; suil.o@sunkorea.ac.kr (corresponding author). Research supported by NRF-2017R1D1A1B03031758.

§Finance.Fishery.Manufacture Industrial Mathematics Center on Big Data, Pusan National University, Busan, 46241, Republic of Korea; parkjr@pusan.ac.kr. Research supported by NRF-2018R1D1A1B07048197.

$E(H)$ incident to the vertex v . An *even* $[a, b]$ -factor of a graph is a spanning subgraph H such that $d_H(v)$ is even and $a \leq d_H(v) \leq b$ for all $v \in V(G)$. If $a = b$, then we call it an a -factor. A graph G is k -edge-connected if for $S \subseteq E(G)$ with $|S| < k$, $G - S$ is connected. The *edge-connectivity* of G , denoted $\kappa'(G)$, is the maximum k such that G is k -edge-connected. A graph G is k -vertex-connected if $|V(G)| \geq k + 1$ and for $S \subseteq V(G)$ with $|S| < k$, $G - S$ is connected. The *vertex-connectivity* of G , denoted $\kappa(G)$, is the maximum k such that G is k -vertex-connected.

Kouider and Vestaargard [4], [5] had explored sufficient conditions for a graph to have an even $[a, b]$ -factor. In 2005, Matsuda [7] gave a sharp sufficient condition for a graph to have an even $[2, b]$ -factor and proposed a conjecture for the existence of an even $[a, b]$ -factor in a graph as follows:

Conjecture 1.1. *Let $2 \leq a \leq b$ be even integers. If G is a graph with n vertices such that (i) $\kappa'(G) \geq 2$, (ii) $n \geq 2a + b + \frac{a^2 - 3a}{b} - 2$, (iii) $\delta(G) \geq a$, and (iv) $\sigma_2(G) \geq \frac{2an}{a+b}$, where $\sigma_2(G) = \min_{uv \in E(G)} (d(u) + d(v))$, then G contains an even $[a, b]$ -factor.*

However, Conjecture 1.1 is not true even when $a = 2$. Remark 3 in [7] says that if $n = b + 2$, then Conjecture 1.1 does not hold. Theorem 8 in [7] says that if we replace $n \geq b + 2$ by $n \geq b + 3$, then G contains an even $[2, b]$ -factor. A result of Iida and Nishimura [2] implies that Conjecture 1.1 is true when $a = b$.

For $a \geq 4$, all other conditions in the conjecture are sharp, except $\kappa'(G) \geq 2$. In Section 2, we provide counterexamples, which are $(a-1)$ -edge-connected. Furthermore, there are also $(a-1)$ -vertex-connected graphs satisfying all conditions in Conjecture 1.1, which do not contain an even $[a, b]$ -factor. Thus to guarantee the existence of an even $[a, b]$ -factor in a graph, we need high vertex-(or edge-)connectivity. By reinforcing the condition $\sigma_2(G) \geq \frac{2an}{a+b}$ to $\delta(G) \geq \frac{an}{a+b}$, we give sufficient conditions for a graph to have an even $[a, b]$ -factor in Theorem 1.2.

Theorem 1.2 (Main Theorem). *Let $4 \leq a \leq b$ be even integers. If G is a graph with n vertices such that (i) $\kappa(G) \geq a$, (ii) $n \geq 2a + b + \frac{a^2 - 3a}{b} - 2$, and (iii) $\delta(G) \geq \frac{an}{a+b}$, then G contains an even $[a, b]$ -factor.*

Katerinis [3], and Egawa and Enomoto [1] independently showed that Theorem 1.2 is true when $a = b$. In this paper, we prove for all $4 \leq a \leq b$ including the case $a = b$. In the papers [3, 1], to have an $[a, a]$ -factor (or a -factor), one of the sufficient conditions is just “connected”. However, if there is an enough gap between a and b , then to have an even $[a, b]$ -factor, a graph must be highly connected (See Section 2).

Note that Condition (ii) and (iii) in Theorem 1.2 imply that $\delta(G) \geq a + 1$. If $\delta(G) \leq a$, then Condition (iii) says $\frac{an}{a+b} \leq a$. Thus we have $n \leq a + b$, which contradicts Condition (ii).

The examples in Section 2 show that the conditions in Theorem 1.2 are sharp. In Section 3, we prove Theorem 1.2 by using Corollary 1.4 of Lovasz’s (g, f) -factor Theory.

Theorem 1.3 (Lovasz’s (g, f) -factor Theory [6]). *Let G be a graph and let g, f be two integer valued functions defined on $V(G)$ such that $0 \leq g(x) \leq f(x) \leq d_G(x)$ for all $x \in V(G)$. Then G has a (g, f) -factor if and only if*

$$\sum_{v \in T} (d(v) - g(v)) + \sum_{u \in S} f(u) - |[S, T]| - q(S, T) \geq 0$$

for all disjoint subsets S and T of $V(G)$, where $q(S, T)$ is the number of components Q of $G - (S \cup T)$ such that $g(v) = f(v)$ for all $v \in V(Q)$ and

$$|[Q, T]| + \sum_{v \in V(Q)} f(v) \equiv 1 \pmod{2}.$$

Corollary 1.4. *Let a and b be even integers with $2 \leq a \leq b$. A graph G has an even $[a, b]$ -factor if*

$$q(S, T) - b|S| + a|T| - \sum_{v \in T} d_{G-S}(v) \leq 0$$

for all disjoint choices $S, T \subseteq V(G)$, where $q(S, T)$ is the number of components Q of $G - (S \cup T)$ such that $|[Q, T]|$ is odd.

By applying Theorem 1.3 when $g(x) = a$ and $f(x) = b$, we have Corollary 1.4.

We point out that Tutte [9] proved that the Lovasz's (g, f) -factor Theory [6] can be demonstrated by using Tutte's f -factor Theory [8].

The Parity Lemma is also used in the proof of our main result.

Lemma 1.5 (Parity Lemma). *Let a and b be positive integers with the same parity. Then $q(S, T) - b|S| + a|T| - \sum_{v \in T} d_{G-S}(v)$ has the same parity as a and b for any disjoint sets $S, T \subseteq V(G)$.*

2 Sharp Examples

In this section, by providing Example 2.1 and Example 2.3, we show why high edge-(or vertex-)connectivity in Theorem 1.2 requires. Note that Matsuda [7] showed in the last section that Condition (ii) and (iii) in Theorem 1.2 are sharp.

Example 2.1 shows that if a graph satisfying Condition (ii), (iii), and (iv) in Conjecture 1.1 is not a -edge-connected, then we cannot guarantee the existence of even $[a, b]$ -factor in the graph. Thus the graph in Example 2.1 is a counterexample to Conjecture 1.1, which has edge-connectivity equal to $a - 1$.

Example 2.1. Let a and b be even integers such that $12 \leq 3a \leq b$, and let t be an integer such that $t \geq \frac{(a+b)^2 - 3a - 4b}{2b} (= \frac{2a+b-4}{2} + \frac{a^2-3a}{2b} > a)$. For $i \in \{1, 2\}$, let H_i be a copy of the complete graph on t vertices, and let $V(H_i) = \{x_{i1}, \dots, x_{it}\}$. Let H_3 be a copy of the complete graph on 2 vertices and let $V(H_3) = \{y, z\}$. Suppose that H is the graph obtained from H_1, H_2 , and H_3 by adding edges between y and $x_{11}, \dots, x_{1(\frac{a}{2}-1)}, x_{2\frac{a}{2}}, \dots, x_{2(a-1)}$, and between z and $x_{21}, \dots, x_{2(\frac{a}{2}-1)}, x_{1\frac{a}{2}}, \dots, x_{1(a-1)}$ (see Figure 1).

Proposition 2.2. *The graph in Example 2.1 has edge-connectivity equal to $a - 1$ and satisfies all conditions in Conjecture 1.1. Furthermore, it does not contain an even $[a, b]$ -factor.*

Proof. Since there are $\frac{a}{2} - 1$ edges between y and H_1 and $\frac{a}{2}$ edges between y and H_2 , and H_1 and H_2 are both complete graphs, there are exactly $a - 1$ edge-disjoint $y - z$ paths including the yz edge. Also, since there are exactly $a - 1$ edges between H_i to H_3 , we have $\kappa'(H) = a - 1$.




Figure 1: The graph H in Example 2.1

The order of H is

$$|V(H)| = 2t + 2 \geq \frac{(a+b)^2 - 3a - 4b}{b} + 2 = 2a + b + \frac{a^2 - 3a}{b} - 2.$$

Since every vertex in $V(H_1) \cup V(H_2)$ has degree at least a and $d_H(y) = d_H(z) = a$, we have $\delta(H) = a$.

Since $\sigma_2(H) = a + (t - 1)$ and $t = \frac{|V(H)| - 2}{2}$, we have

$$\sigma_2(H) = a + \frac{|V(H)|}{2} - 2 \geq \frac{|V(H)|}{2} = \frac{2a|V(H)|}{4a} \geq \frac{2a|V(H)|}{a+b}.$$

Thus H satisfies all conditions in Conjecture 1.1.

Now, we prove that H does not contain an even $[a, b]$ -factor. Assume to the contrary that H has an even $[a, b]$ -factor F . Since $d_H(y) = a$, all edges incident to y must be in F . Since $H_1 \cap F$ is also a graph, $\sum_{v \in V(H_1 \cap F)} d_{H_1 \cap F}(v)$ must be even by the degree-sum formula. Note that the $a - 1$ edges incident to both H_1 and H_2 in F are not in $H_1 \cap F$. Thus we have

$$\sum_{v \in V(H_1 \cap F)} d_{H_1 \cap F}(v) = \sum_{v \in V(H_1 \cap F)} d_F(v) - (a - 1).$$

However, the degree sum is odd since $a - 1$ is odd and every vertex in F has even degree. Thus we have the desired result. \square

Example 2.3 shows that if a graph satisfying Condition (ii), (iii), and (iv) in Conjecture 1.1 is not a -vertex-connected, then we cannot guarantee the existence of even $[a, b]$ -factor in the graph. Thus the graph in Example 2.3 is also a counterexample to Conjecture 1.1, which is $(a - 1)$ -vertex-connected. Note that in Example 2.1, we require $b \geq 3a$ while in Example 2.3, we require $b \geq \frac{a^2 - 3a + a\sqrt{(a-3)(a+1)}}{2}$.

Example 2.3. Let a and b be even integers at least 4 with $b \geq \frac{a^2 - 3a + a\sqrt{(a-3)(a+1)}}{2}$. Let L_0 be the trivial graph on $a - 1$ vertices, and let $V(L_0) = \{y_1, \dots, y_{(a-1)}\}$.

For $1 \leq i \leq a$, let L_i be a copy of the complete graph on $a + 2$ vertices and let $V(L_i) = \{x_{i1}, \dots, x_{i(a+2)}\}$. Let t be a positive integer such that $(a + 2) - a^2 - a + b + \frac{a^2 - 3a}{b} - 1 \leq t \leq -a^2 - 2a + b + \frac{b}{a} + 2$. Let L_{a+1} be a copy of the complete graph on t vertices and let $V(L_{a+1}) = \{x_{(a+1)1}, \dots, x_{(a+1)t}\}$. Suppose that L is the graph obtained from L_0, \dots, L_{a+1} by adding edges between y_j and x_{ij} for all $i \in \{1, \dots, a + 1\}$ and for all $j \in \{1, \dots, a - 1\}$ (see Figure 2).

Proposition 2.4. *The graph in Example 2.3 has vertex-connectivity equal to $a - 1$ and satisfies all conditions in Conjecture 1.1. Furthermore, it does not contain an even $[a, b]$ -factor.*

Proof. For each $i \in [a + 1]$, there are $a - 1$ vertex-disjoint paths between any vertex in L_i and L_0 by using the vertex $x_{i1}, x_{i2}, \dots, x_{i(a-1)}$. Also, for $i \neq j$, there are $a + 1$ vertex-disjoint paths between y_i and y_j by using the path $y_i x_{ki} x_{kj} y_j$ for $k \in [a + 1]$. Thus we have $\kappa(L) = (a - 1)$.

Since $t \geq -a^2 - a + b + \frac{a^2 - 3a}{b} - 1$, the order of L is

$$|V(L)| = a - 1 + (a + 2)a + t \geq 2a + b + \frac{a^2 - 3a}{b} - 2.$$

Since for $i \in \{0, 1, \dots, a + 1\}$, every vertex in $V(L_i)$ has degree at least $a + 1$ and every vertex in L_0 has degree $a + 1$, we have $\delta(L) = a + 1$.

Since $\sigma_2(L) = 2(a + 1)$ and $t \leq -a^2 - 2a + b + \frac{b}{a} + 2$, we have

$$\frac{2a|V(L)|}{a + b} = \frac{2a(a^2 + 3a - 1 + t)}{a + b} \leq \frac{2a(a + b + \frac{b}{a} + 1)}{a + b} = 2(a + 1) = \sigma_2(L)$$

Thus F satisfies all conditions in Conjecture 1.1.

Now, we prove that L does not contain an even $[a, b]$ -factor. Assume to the contrary that L has an even $[a, b]$ -factor F . Then we have $d_F(v) = a$ for every vertex in L_0 since a is even. Since $\sum_{v \in V(L_i \cap F)} d_{L_i \cap F}(v)$ must be even by the degree-sum formula, there are at most $a - 2$ edges coming out from $V(L_i)$ in F . Thus we have

$$a(a - 1) \leq (a - 2)(a + 1),$$

which is a contradiction. \square

Proposition 2.4 shows that Condition (i) in Theorem 1.2 is tight.

3 Proof of Theorem 1.2

When $a = 2$, Conjecture 1.1 is true if we replace Condition (ii) by $n \geq b + 3$. So is Theorem 1.2 for $a = 2$. There is a counterexample to Conjecture 1.1 if $n = b + 2$ (see Remark 3 [7]). From now, we assume that $a \geq 4$. The examples in Section 2 and the last section in [7] say that we require the conditions in Theorem 1.2 for a graph to have an even $[a, b]$ -factor.

In this section, we prove Theorem 1.2. Note that for $a = b$, Theorem 1.2 is true by Katerinis [3], Egawa and Enomoto [1], and Iida and Nishimura [2]. In this paper, we prove for all $4 \leq a \leq b$ including the case $a = b$. To prove Case 3 and Case 4-1 in the proof of Theorem 1.2, we use Proposition 3.1.




Figure 2: The graph L in Example 2.3

Proposition 3.1. *Let a, b, n , and p be integers such that $4 \leq a \leq b$ and $p > 0$, and let $f(x) = n + (a - 1 - \frac{an}{a+b})x + (x - 1 - b)\frac{ax-p}{b}$.*

(i) If $n \geq 2a + b + \frac{a^2-3a}{b} - 2$, then $f(b+1) < 0$ and $f(a+b-3) < 0$.

(ii) If $n \geq 2a + b + \frac{a^2-3a}{b} + 1$, then $f(a+b-1) < 0$ and $f(a+b-2) < 0$.

Proof. (i) Assume that $n \geq 2a + b + \frac{a^2-3a}{b} - 2$. Then we have

$$\begin{aligned} f(b+1) &= n + (a - 1 - \frac{an}{a+b})(b+1) = (1-a)(\frac{bn}{a+b} - b - 1) \\ &\leq (1-a) \left[\frac{b}{a+b} (2a + b + \frac{a^2-3a}{b} - 2) - b - 1 \right] \\ &= (1-a) \left[\frac{b(a-3) + a(a-4)}{a+b} \right] < 0 \end{aligned}$$

and

$$\begin{aligned} f(a+b-3) &= n + (a - 1 - \frac{an}{a+b})(a+b-3) + [(a+b-3) - 1 - b] \frac{a(a+b-3) - p}{b} \\ &= (4a + b - a^2 - ab) \left(\frac{n}{a+b} - \frac{a+b-3}{b} \right) - \frac{(a-4)p}{b} \\ &\leq (4a + b - a^2 - ab) \left(\frac{2a + b + \frac{a^2-3a}{b} - 2}{a+b} - \frac{a+b-3}{b} \right) - \frac{(a-4)p}{b} \\ &= \frac{a(4-a) + b(1-a)}{a+b} - \frac{(a-4)p}{b} < 0. \end{aligned}$$

(ii) Assume that $n \geq 2a + b + \frac{a^2-3a}{b} + 1$. Then we have

$$\begin{aligned}
f(a+b-1) &= n + (a-1 - \frac{an}{a+b})(a+b-1) + [(a+b-1) - 1 - b] \frac{a(a+b-1) - p}{b} \\
&= (2a+b-a^2-ab)(\frac{n}{a+b} - \frac{a+b-1}{b}) - \frac{(a-2)p}{b} \\
&\leq (2a+b-a^2-ab)(\frac{2a+b + \frac{a^2-3a}{b} + 1}{a+b} - \frac{a+b-1}{b}) - \frac{(a-2)p}{b} \\
&= \frac{(a(2-a) + b(1-a))(-2a+2b)}{b(a+b)} - \frac{(a-2)p}{b} < 0.
\end{aligned}$$

Since $f(x)$ is a quadratic function which has a positive leading coefficient and $f(b+1) < 0$ by (i), we have $f(x) < 0$ for all $x \in [b+1, a+b-1]$ so that $f(a+b-2) < 0$. \square

Now, we are ready to prove Theorem 1.2.

Proof of Theorem 1.2. Assume to the contrary that G has no even $[a, b]$ -factor. Then there exist disjoint subsets S and T in $V(G)$ such that

$$0 < q(S, T) - b|S| + a|T| - \sum_{v \in T} d_{G-S}(v)$$

by Corollary 1.4. Let $p = -b|S| + a|T|$ so that

$$0 < q(S, T) + p - \sum_{v \in T} d_{G-S}(v).$$

Note that $p > 0$ since $q(S, T) - \sum_{v \in T} d_{G-S}(v) \leq 0$.

We consider four cases depending on $|T|$.

$$|T| = \frac{b}{a}|S| \quad \begin{array}{c} \text{Case 2} \\ | \\ b \end{array} \quad \begin{array}{c} \text{Case 3} \\ | \\ a+b-3 \end{array} \quad \begin{array}{c} \text{Case 4} \\ | \\ a+b-1 \end{array} \quad \begin{array}{c} \text{Case 1} \\ | \\ n \end{array}$$

In Case 4, we consider two subcases Case 4-1 and Case 4-2 depending on n . To prove Case 1, Case 3, and Case 4-1, we use the same argument as in [4]. For Case 2 and Case 4-2, we prove using a new technique.

Case 1: $|T| \geq a+b$. Since $n \geq |S| + |T| + q(S, T)$, we have

$$\begin{aligned}
|S| = \frac{a|T| - p}{b} &\leq \frac{a(n - |S| - q(S, T)) - p}{b} \iff |S| \leq \frac{a(n - q(S, T)) - p}{a+b} \\
\implies |[S, T]| &\leq |S||T| \leq \frac{a(n - q(S, T)) - p}{a+b} |T|. \tag{1}
\end{aligned}$$

With Inequality (1), we have

$$\begin{aligned}
0 < q(S, T) - b|S| + a|T| - \sum_{v \in T} d_{G-S}(v) &= q(S, T) + p - \sum_{v \in T} d_G(v) + |[S, T]| \\
&\leq q(S, T) + p - \delta(G)|T| + \frac{a(n - q(S, T)) - p}{a+b} |T| \\
&\leq q(S, T) + p - \frac{an}{a+b} |T| + \frac{a(n - q(S, T)) - p}{a+b} |T| \\
&= q(S, T) + p - \frac{aq(S, T) + p}{a+b} |T| \leq (1-a)q(S, T) < 0,
\end{aligned}$$

which is a contradiction.

Case 2: $|T| \leq b$. Since $\delta(G) \geq a + 1$, we have

$$\begin{aligned} 0 &< q(S, T) - b|S| + a|T| - \sum_{v \in T} d_{G-S}(v) = q(S, T) - b|S| + a|T| - \sum_{v \in T} d_G(v) + |[S, T]| \\ &\leq q(S, T) - b|S| + a|T| - \delta(G)|T| + |S||T| \leq q(S, T) - b|S| + [a - (a + 1) + |S|]|T| \\ &\leq q(S, T) - b|S| + b|S| - |T| = q(S, T) - |T|, \end{aligned}$$

which implies $q(S, T) > |T| \geq 0$. Let l be the minimum of $|[Q, T]|$ over all components Q of $G - (S \cup T)$ such that $|[Q, T]|$ is odd. Then we have $l \geq 1$. Also, we have $\sum_{v \in T} d_{G-S}(v) \geq lq(S, T) \iff \frac{1}{l} \sum_{v \in T} d_{G-S}(v) \geq q(S, T)$. Thus we have

$$\begin{aligned} 0 &< q(S, T) - b|S| + a|T| - \sum_{v \in T} d_{G-S}(v) \leq \frac{1-l}{l} \sum_{v \in T} d_{G-S}(v) - b|S| + a|T| \\ &= \frac{1-l}{l} \sum_{v \in T} d_G(v) - \frac{1-l}{l} |[S, T]| - b|S| + a|T| \\ &\leq \frac{1-l}{l} \delta(G)|T| - \frac{1-l}{l} |S||T| - b|S| + a|T| \\ &\leq \frac{(1-l)(a+1)}{l} |T| + |S| \left(\frac{l-1}{l} |T| - b \right) + a|T| \\ &\leq \frac{a+1-l}{l} |T| - |S| \frac{b}{l}. \end{aligned} \tag{2}$$

If $a+1-l \leq 0$ in Inequality (2), then it is a contradiction. Thus $a+1-l > 0$, and since $|T| \leq b$, we have

$$0 < \frac{a+1-l}{l} |T| - |S| \frac{b}{l} \leq \frac{a+1-l-|S|}{l} b.$$

This gives $a+1-l-|S| > 0$ so that $|S| \leq a-l$.

Claim 1: $|S| = a-l$. Assume to the contrary that $|S| < a-l$. Then each component Q in $G-S-T$ with $|[Q, T]| = l$ can have at most $a-1$ neighbors in $S \cup T$. Since G is a -vertex-connected, there is only one such a component and every vertex of T must have a neighbor in Q . Thus $|T| \leq l$. For $v \in V(T)$, we have

$$d(v) = |[{\{v\}, S}]| + (d_{G[T]}(v) + |[{\{v\}, Q}]|) \leq (a-l-1) + l = a-1 < a+1 \leq \delta(G),$$

which is a contradiction. Thus $|S| = a-l$, which implies $S \neq \emptyset$ since a is even and l is odd. Since $|S| \geq 1$ and $|T| \leq b$, we have

$$\begin{aligned} 0 &< q(S, T) - b|S| + a|T| - \sum_{v \in T} d_{G-S}(v) \\ &= q(S, T) - b|S| + a|T| - \sum_{v \in T} d_G(v) + |[S, T]| \\ &\leq q(S, T) - b|S| + a|T| - \delta(G)|T| + |S||T| \\ &\leq q(S, T) - b|S| + [a - (a + 1) + |S|]|T| \\ &\leq q(S, T) - b|S| + (|S| - 1)b = q(S, T) - b. \end{aligned} \tag{3}$$

By Inequality (3), we have $q(S, T) \geq b + 1$. Let $q(S, T) = b + \alpha$ for some $\alpha \geq 1$.

Let q_l be the number of components Q of $G - (S \cup T)$ such that $||[Q, T]|| = l$. Since $|S| = a - l$, $|T| \leq b$, and $q(S, T) = b + \alpha$, we have

$$\begin{aligned}
0 &< q(S, T) - b|S| + a|T| - \sum_{v \in T} d_{G-S}(v) \\
&\leq q(S, T) - b(a - l) + a|T| - [lq_l + (l + 2)(q(S, T) - q_l)] \\
&\leq (-l - 1)q(S, T) + bl + 2q_l = (-l - 1)(b + \alpha) + bl + 2q_l \\
&= -b - (l + 1)\alpha + 2q_l.
\end{aligned} \tag{4}$$

By Inequality (4), we have $q_l > \frac{b + (1+l)\alpha}{2}$. Note that b and $1 + l$ are even integers so that $\frac{b + (1+l)\alpha}{2}$ is an integer. Thus $q_l \geq \frac{b + (1+l)\alpha}{2} + 1$.

Let m be the minimum of $|Q|$ over all components Q in $G - (S \cup T)$ such that $||[Q, T]|| = l$. There exists a vertex v in $V(Q)$ such that $||[\{v\}, T]|| \leq \frac{l}{m}$ by the pigeonhole principle. Then we have

$$\begin{aligned}
\delta(G) &\leq d(v) = d_Q(v) + ||[\{v\}, T]|| + ||[\{v\}, S]|| \leq (m - 1) + \left(\frac{l}{m}\right) + (a - l) \\
\implies m &\leq \frac{\delta(G) + l + 1 - a - \sqrt{(\delta(G) + l + 1 - a)^2 - 4l}}{2} \\
\text{or } m &\geq \frac{\delta(G) + l + 1 - a + \sqrt{(\delta(G) + l + 1 - a)^2 - 4l}}{2}.
\end{aligned}$$

Note that we have

$$\frac{\delta(G) + l + 1 - a - \sqrt{(\delta(G) + l + 1 - a)^2 - 4l}}{2} < 1 \iff \delta(G) \geq a + 1. \tag{5}$$

Since $m \geq 1$, we have $m \geq \frac{\delta(G) + l + 1 - a + \sqrt{(\delta(G) + l + 1 - a)^2 - 4l}}{2}$ by Inequality (5). Note that we have

$$\begin{aligned}
\frac{\delta(G) + l + 1 - a + \sqrt{(\delta(G) + l + 1 - a)^2 - 4l}}{2} &\geq \delta(G) - |S| \\
\iff (\delta(G) + l + 1 - a)^2 - 4l &\geq (\delta(G) - 2|S| - 1 + a - l)^2 \\
\iff (\delta(G) - |S|)(1 - a + l + |S|) &\geq l \\
\iff \delta(G) - a + l \geq l &\iff \delta(G) \geq a.
\end{aligned} \tag{6}$$

By Inequality (6), we have $m \geq \delta(G) - |S|$, implying $n \geq |S| + |T| + q_l(\delta(G) - |S|)$. Since $q_l \geq \frac{b + (1+l)\alpha}{2} + 1$, $\delta(G) \geq \frac{an}{a+b}$, $2b - a(b + (1+l)\alpha) = (2-a)b - (1+l)\alpha < 0$, and $a|T| - b|S| > 0$, we have

$$\begin{aligned}
\frac{a+b}{a}\delta(G) &\geq n \geq |S| + |T| + \left[\frac{b + (1+l)\alpha}{2} + 1\right](\delta(G) - |S|) \\
\iff \delta(G) &\leq \frac{2a|T| - a[b + (1+l)\alpha]|S|}{2b - a[b + (1+l)\alpha]} = \frac{2(a|T| - b|S|)}{2b - a[b + (1+l)\alpha]} + |S| < |S|,
\end{aligned}$$

which is a contradiction.

Case 3: $b+1 \leq |T| \leq a+b-3$. Since $q(S, T) \leq n - |S| - |T|$ and $\delta(G) \geq \frac{an}{a+b}$, we have

$$\begin{aligned}
0 &< q(S, T) - b|S| + a|T| - \sum_{v \in T} d_{G-S}(v) \\
&\leq (n - |S| - |T|) - b|S| + a|T| - \frac{an}{a+b}|T| + |S||T| \\
&= n + (a - 1 - \frac{an}{a+b})|T| + (|T| - 1 - b)|S| \\
&= n + (a - 1 - \frac{an}{a+b})|T| + (|T| - 1 - b)\frac{a|T| - p}{b}. \tag{7}
\end{aligned}$$

Let $f(|T|) = n + (a - 1 - \frac{an}{a+b})|T| + (|T| - 1 - b)\frac{a|T| - p}{b}$. Since f is a quadratic function which has a positive leading coefficient, the maximum value of f occurs when $|T| = b+1$ or $|T| = a+b-3$. By Proposition 3.1, both $f(b+1)$ and $f(a+b-3)$ are negative, which contradicts Inequality (7).

Case 4: $|T| = a+b-2$ or $a+b-1$.

Case 4-1: $n \geq 2a+b + \frac{a^2-3a}{b} + 1$. By using the same argument with Case 3 and Proposition 3.1, we have the desired result.

Case 4-2: $2a+b + \frac{a^2-3a}{b} - 2 \leq n < 2a+b + \frac{a^2-3a}{b} + 1$. Let $|T| = a+b-k$ and $2a+b + \frac{a^2-3a}{b} - j \leq n < 2a+b + \frac{a^2-3a}{b} - j + 1$ where $k \in \{1, 2\}$ and $j \in \{0, 1, 2\}$. Let $n = 2a+b + \frac{a^2-3a}{b} - j + \epsilon$, where $0 \leq \epsilon < 1$.

Claim 2: If $\delta(G) \geq j - k + i + n - |T| = i + a + \frac{a^2-3a}{b} + \epsilon$, then $a(3-k) - \epsilon b \geq (a-k)(j-k) + i(a+b-k) + (a-k-1)q(S, T) + 2$, where i is an integer. By Lemma 1.5, we have

$$\begin{aligned}
2 &\leq q(S, T) - b|S| + a|T| - \sum_{v \in T} d_G(v) + |[S, T]| \\
&\leq q(S, T) - b|S| + a|T| - \delta(G)|T| + |S||T| \\
&\leq q(S, T) - b|S| + a|T| - (i + a + \frac{a^2-3a}{b} + \epsilon)|T| + |S||T| \\
&\leq q(S, T) + (a-k)(n - |T| - q(S, T)) - (i + \frac{a^2-3a}{b} + \epsilon)(a+b-k) \\
&= (k+1-a)q(S, T) + (a-k)(a-i-j+k) - b(i + \frac{a^2-3a}{b} + \epsilon) \\
&= (k+1-a)q(S, T) + (a-k)(k-j) - i(a+b-k) + a(3-k) - \epsilon b.
\end{aligned}$$

Thus we have the desired result.

Since $\delta(G) \geq \frac{an}{a+b}$, we have

$$\begin{aligned}
\delta(G) - n + |T| &\geq \frac{an}{a+b} - n + |T| = -\frac{bn}{a+b} + (a+b-k) \\
&= \frac{-b(2a+b + \frac{a^2-3a}{b} - j + \epsilon) + (a+b-k)(a+b)}{a+b} \\
&= \frac{(3-k)a + (j-k)b - \epsilon b}{a+b} \tag{8} \\
&> j - k - 1,
\end{aligned}$$

which is true for $j \in \{0, 1, 2\}$ and $\epsilon \in [0, 1)$. Since $\delta(G) - n + |T|$ is an integer, we obtain $\delta(G) - n + |T| \geq j - k$, which satisfies the condition on $\delta(G)$ when $i = 0$ in Claim 2. Thus we have $a(3 - k) - \epsilon b \geq (a - k)(j - k) + (a - k - 1)q(S, T) + 2$. By Inequality (8), we have

$$\begin{aligned} \delta(G) - n + |T| &\geq \frac{(3 - k)a + (j - k)b - \epsilon b}{a + b} \\ &\geq \frac{(j - k)b + (a - k)(j - k) + (a - k - 1)q(S, T) + 2}{a + b} \\ &= j - k + \frac{-k(j - k) + (a - k - 1)q(S, T) + 2}{a + b} > j - k, \end{aligned}$$

which is true for $k \in \{1, 2\}$ and $j \in \{0, 1, 2\}$ and $a \geq 4$. Since $\delta(G) - n + |T|$ is an integer, we obtain $\delta(G) - n + |T| \geq j - k + 1$, which satisfies the condition on $\delta(G)$ when $i = 1$ in Claim 2. Thus we have

$$a(3 - k) - \epsilon b \geq (a - k)(j - k) + (a + b - k) + (a - k - 1)q(S, T) + 2. \quad (9)$$

When $k = j = 1$, Inequality (9) becomes $a - (\epsilon + 1)b \geq (a - 2)q(S, T) + 1$ which is a contradiction since $a - (\epsilon + 1)b \leq 0$ and $(a - 2)q(S, T) + 1 > 0$. Similarly, we have a contradiction when $(k, j) \in \{(1, 2), (2, 1), (2, 2)\}$ by using Inequality (9). The remaining case is when $j = 0$. By Inequality (8) and (9), we improve $\delta(G)$ as follows:

$$\begin{aligned} \delta(G) - n + |T| &\geq \frac{(3 - k)a + (j - k)b - \epsilon b}{a + b} \\ &\geq \frac{(j - k)b + (a - k)(j - k) + (a + b - k) + (a - k - 1)q(S, T) + 2}{a + b} \\ &= j - k + 1 + \frac{-k(j - k + 1) + (a - k - 1)q(S, T) + 2}{a + b} > j - k + 1, \end{aligned}$$

which is true for $(k, j) \in \{(1, 0), (2, 0)\}$ and $a \geq 4$. Since $\delta(G) - n + |T|$ is an integer, we obtain $\delta(G) - n + |T| \geq j - k + 2$, which satisfies the condition on $\delta(G)$ when $i = 2$ in Claim 2. Thus we have

$$a(3 - k) - \epsilon b \geq (a - k)(j - k) + 2(a + b - k) + (a - k - 1)q(S, T) + 2. \quad (10)$$

When $k = 1$ and $j = 0$, Inequality (10) becomes $a - (\epsilon + 2)b \geq (a - 2)q(S, T) + 2$ which is a contradiction since $a - (\epsilon + 2)b < 0$ and $(a - 2)q(S, T) + 2 > 0$. Similarly, we get a contradiction when $k = 2$ and $j = 0$, which completes the proof. \square

4 Concluding Remarks

In this section, we provide some questions and conjecture.

Question 4.1. *If we replace “ $\kappa(G)$ ” in Theorem 1.2 by “ $\kappa'(G)$ ”, then do we have the same conclusion?*

Question 4.2. *If we replace “ $\delta(G) \geq \frac{an}{a+b}$ ” in Theorem 1.2 by “ $\sigma_2(G) \geq \frac{2an}{a+b}$ ”, then do we have the same conclusion?*

We might be also interested in some sufficient conditions for a certain eigenvalue in a certain graph G to have an even $[a, b]$ -factor. If G has an even $[a, b]$ -factor, then we have $\lambda_1(G) \geq a$ since $\lambda_1(G) \geq \delta(G)$, where $\lambda_1(G)$ is the largest eigenvalue of G . Is there a sufficient condition for $\lambda_1(G)$ in a graph G to have an even $[a, b]$ -factor? If we restrict our attention to a complete bipartite graph, which looks the simplest case, then it is easy to get a sufficient condition for the largest eigenvalue.

Observation 4.3. *Let G be the complete bipartite graph $K_{x, n-x}$ such that $n \geq 2x > 0$. Then G has an $[a, b]$ -factor if and only if*

$$\lambda_1(G) \geq \begin{cases} \sqrt{a(n-a)} & \text{if } n < a+b \\ \frac{\sqrt{ab}}{a+b}n & \text{if } n \geq a+b. \end{cases}$$

Proof. G has an $[a, b]$ -factor F if and only if

$$x \geq a \text{ and } (n-x-b)x \leq (n-x)(x-a) (\Leftrightarrow x \geq \frac{an}{a+b})$$

since $\delta(F) \geq a$ and $\Delta(F) \leq b$.

Thus we have the desired result with $\lambda_1(G) = \sqrt{x(n-x)}$. \square

Among n -vertex graphs G without $[a, b]$ -factor, we guess that the n -vertex graph $H_{n,a}$ obtained from one vertex and a copy of K_{n-1} by adding $a-1$ edges between them has the largest eigenvalue. Note that there are $n-a$ vertices with degree $n-2$, $a-1$ vertices with degree $n-1$, and 1 vertex with degree $a-1$ in the graph H_a . Thus H_a cannot have an $[a, b]$ -factor.

Conjecture 4.4. *Let an be an even integer at least 2, where $n \geq a+1$, and let $\rho(n, a)$ be the largest eigenvalue of $H_{n,a}$. If G is an n -vertex graph with $\lambda_1(G) > \rho(n, a)$, then G has an $[a, b]$ -factor.*

We mention that $\lambda_1(H_{n,a})$ equals the largest root of $x^3 - (n-3)x^2 - (a+n-3)x - a^2 + (a-1)n + 1 = 0$ without giving a reason in detail.

References

- [1] Y. Egawa and H. Enomoto, Sufficient conditions for the existence of k -factors, *Recent Studies in Graph Theory*, V. R. Kulli, Ed., Vishwa International Publication, India (1989) 96–105.
- [2] T. Iida and T. Nishimura, An ore-type condition for the existence of k -factors in graphs, *Graphs Combin.* **7** Issue 4, (1991), 353–361.
- [3] P. Katerinis, Minimum degree of a graph and the existence of k -factors. *Proc. Indian Acad. Sci (Math. Sci.)* **94** (1985), 123–127.
- [4] M. Kouider, P.D. Vestergaard, Even $[a, b]$ -factors in graphs, *Discuss. Math. Graph Theory* **24** (2004), 431–441.
- [5] M. Kouider, P.D. Vestergaard, Sufficient condition for the existence of an even $[a, b]$ -factor in graph, *Graphs Combin.* **29** Issue 4 (2013) 1051–1057.

- [6] L. Lovasz, Subgraphs with prescribed valencies, *J. of Combinatorial Theory B* (1970) 391–416.
- [7] H. Matsuda, Ore-type conditions for the existence of even $[2, b]$ -factors in graphs, *Disc. Math.* **304** (2005) 51–61.
- [8] W. T. Tutte, The factors of graphs, *Canad. J. Math.* **4** (1952) 314–328.
- [9] W. T. Tutte, Graph factors, *Combinatorica* **1** (1981) 79–97.