

**JACOB'S LADDERS AND GRAFTING OF THE COMPLETE
HYBRID FORMULAS INTO ζ -SYNERGETIC
META-FUNCTIONAL EQUATION FOR RIEMANN'S
ZETA-FUNCTION**

JAN MOSER

ABSTRACT. In this paper we prove that there is a grafting of complete hybrid formula on meta-functional equation. This synergetic formula gives the result of a ζ -chemical reaction between different sets of values of Riemann's zeta-function on the critical strip.

Dedicated to recalling on A.S. Eddington's - manipulations with set of world-constants.

1. INTRODUCTION

In this paper we obtain the first set of meta-functional equations. It is an example of entirely new kind of formulas in the theory of Riemann's zeta-function. Namely: there are infinite sets

$$(1.1) \quad \begin{aligned} & \{\alpha_1^{1,1}\}, \{\alpha_1^{2,1}\}, \{\alpha_1^{3,1}\} \in (L, +\infty), L > 0, \\ & \{w_1\}, \{w_2\}, \{w_3\} \in S_{\sigma_1, \sigma_2} = (\sigma_1, \sigma_2) \times (0, +\infty), \\ & \frac{1}{2} < \sigma_1 < \sigma_2 < 1, \\ & w_1, w_2, w_3 \in \mathbb{C} \end{aligned}$$

of values such that the following meta-functional equation

$$(1.2) \quad \begin{aligned} & \left| \zeta \left(\frac{1}{2} + i\alpha_1^{2,1} \right) \right|^2 |\zeta(w_2)| - \left| \zeta \left(\frac{1}{2} + i\alpha_1^{3,1} \right) \right|^2 |\zeta(w_1)| \sim \\ & \sim \left| \zeta \left(\frac{1}{2} + i\alpha_1^{3,1} \right) \right|^2 |\zeta(w_3)|, L \rightarrow \infty \end{aligned}$$

holds true.

Remark 1. The formula (1.2) is:

- (a) the simplest element of meta-functional equations obtained in this paper,
- (b) synergetic one since it is a result of cooperative interactions (see our interpretation in [8]) between the following sets of values

$$(1.3) \quad \left\{ \left| \zeta \left(\frac{1}{2} + it \right) \right|^2 \right\}, t \in (L, +\infty), |\zeta(w)|, w \in S_{\sigma_1, \sigma_2}$$

Key words and phrases. Riemann zeta-function.

where, of course,

$$(1.4) \quad \left\{ \frac{1}{2} + it : t > L \right\} \cap S_{\sigma_1, \sigma_2} = \emptyset.$$

Remark 2. Let us notice explicitly that in the case (1.3) we have synergetic (cooperative) interactions between the following parts of the function

$$\zeta(s), \quad s \in \mathbb{C} \setminus \{1\}, \quad s = \sigma + it,$$

namely, between the part

$$\left\{ \left| \zeta \left(\frac{1}{2} + it \right) \right|^2 \right\}, \quad t > L$$

on the critical line $\sigma = \frac{1}{2}$, and the part (see (1.1))

$$\{ |\zeta(\sigma + it)| \}, \quad \sigma, \quad t \in S_{\sigma_1, \sigma_2}$$

on the strip (see also (1.4)).

1.1. Next, we give the following

Remark 3. The proof of meta-functional equation (comp. (1.2)) is based:

- (a) on new notions and methods in the theory of Riemann's zeta-function we have introduced in our series of 46 papers concerning Jacob's ladders (these can be found in arXiv [math.CA] starting with the paper [1]),
- (b) on the classical H. Bohr's theorem in analysis (1914) about the roots of the equation

$$\zeta(s) = a, \quad a \neq 0, \quad a \in \mathbb{C}, \quad \frac{1}{2} < \sigma_1 < \sigma < \sigma_2 < 1,$$

- (c) and on our presently introduced notion of *gifting* of complete hybrid formulas that serves for synthesis of the conceptions (a) and (b).

From our notions and methods (comp. (a)) we use here especially: Jacob's ladder (see [1]), ζ -disconnected set (see [3]), algorithm for generating the ζ -factorization formulas (see [4]) crossbreeding, secondary crossbreeding, exact and asymptotic complete hybrid formula (see [6], [8]). Short surveys of these notions can be found in [5] and [8].

Remark 4. We notice explicitly that the notion of meta-functional equation introduced here constitutes also the generic complement to the Riemann's functional equation on the critical strip (comp. also [7]).

2. ZETA-FACTORIZATION FORMULAS

Here we obtain, by making use of our algorithm for generating ζ -factorization formulas (see [5], (3.1) – (3.11), comp. [4]) the following.

2.1. First, we have the following lemmas (comp. [6], (2.1) – (2.9)):

Since

$$\frac{1}{U} \int_{\pi L}^{\pi L + U} \sin^2 t dt = \frac{1}{2} \left(1 - \frac{\sin 2U}{2U} \right)$$

then we have the following

Lemma 1. For the function

$$(2.1) \quad f_1(t) = \sin^2 t \in \tilde{C}_0[\pi L, \pi L + U], \quad U \in (0, \pi/2), \quad L \in \mathbb{N}$$

there are vector-valued functions

$$(2.2) \quad (\alpha_0^{1,k_1}, \alpha_1^{1,k_1}, \dots, \alpha_{k_1}^{1,k_1}, \beta_1^{k_1}, \dots, \beta_{k_1}^{k_1}), \\ 1 \leq k_1 \leq k_0, \quad k_1, k_0 \in \mathbb{N}$$

(here we fix arbitrary k_0) such that the following exact ζ -factorization formula

$$(2.3) \quad \prod_{r=1}^{k_1} \frac{\tilde{Z}^2(\alpha_r^{1,k_1})}{\tilde{Z}^2(\beta_r^{k_1})} = \frac{1}{2} \left(1 - \frac{\sin 2U}{2U} \right) \frac{1}{\sin^2(\alpha_0^{1,k_1})}, \quad \forall L \geq L_0 > 0$$

(where L_0 is sufficiently big) holds true, where

$$(2.4) \quad \alpha_r^{1,k_1} = \alpha_r(U, \pi L, k_1; f_1), \quad r = 0, 1, \dots, k_1, \\ \beta_r^{k_1} = \beta_r(U, \pi L, k_1), \quad r = 1, \dots, k_1, \\ \alpha_0^{1,k_1} \in (\pi L, \pi L + U), \\ \alpha_r^{1,k_1}, \beta_r^{k_1} \in (\overbrace{\pi L}^r, \overbrace{\pi L + U}^r), \quad r = 1, \dots, k_1,$$

and the segment

$$(\overbrace{\pi L}^r, \overbrace{\pi L + U}^r)$$

is the r -th reverse iteration (by means of the Jacob's ladder, see [3]) of the basic segment

$$[\pi L, \pi L + U] = [\overbrace{\pi L}^0, \overbrace{\pi L + U}^0].$$

Next,

$$(2.5) \quad \tilde{Z}^2(t) = \frac{|\zeta(\frac{1}{2} + it)|^2}{\omega(t)}, \quad \omega(t) = \left\{ 1 + \mathcal{O}\left(\frac{\ln \ln t}{\ln t}\right) \right\} \ln t,$$

see [2], (6.1), (6.7), (7.7), (7.8) and (9.1).

Since

$$\frac{1}{U} \int_{\pi L}^{\pi L + U} \cos^2 t dt = \frac{1}{2} \left(1 + \frac{\sin 2U}{2U} \right)$$

then we have the following

Lemma 2. For the function

$$(2.6) \quad f_2(t) = \cos^2 t \in \tilde{C}_0[\pi L, \pi L + U], \quad U \in (0, \pi/2), \quad L \in \mathbb{N}$$

there are vector-valued functions

$$(2.7) \quad (\alpha_0^{2,k_2}, \alpha_1^{2,k_2}, \dots, \alpha_{k_2}^{2,k_2}, \beta_1^{k_2}, \dots, \beta_{k_2}^{k_2}), \\ 1 \leq k_2 \leq k_0, \quad k_2 \in \mathbb{N}$$

such that the following exact ζ -factorization formula

$$(2.8) \quad \prod_{r=1}^{k_2} \frac{\tilde{Z}^2(\alpha_r^{2,k_2})}{\tilde{Z}^2(\beta_r^{k_2})} = \frac{1}{2} \left(1 + \frac{\sin 2U}{2U} \right) \frac{1}{\cos^2(\alpha_0^{2,k_2})}, \quad \forall L \geq L_0$$

holds true, where

$$(2.9) \quad \begin{aligned} \alpha_r^{2,k_2} &= \alpha_r(U, \pi L, k_2; f_2), \quad r = 0, 1, \dots, k_2, \\ \beta_r^{k_2} &= \beta_r(U, \pi L, k_2), \quad r = 1, \dots, k_2, \\ \alpha_0^{2,k_2} &\in (\pi L, \pi L + U), \\ \alpha_r^{2,k_2}, \beta_r^{k_2} &\in (\overbrace{\pi L}^r, \overbrace{\pi L + U}^r), \quad r = 1, \dots, k_2. \end{aligned}$$

2.2. Secondly, since

$$\frac{1}{U} \int_{\pi L}^{\pi L + U} \cos 2t dt = \frac{\sin 2U}{2U}$$

then we have the following

Lemma 3. For the function

$$(2.10) \quad f_3(t) = \cos 2t \in \tilde{C}_0[\pi L, \pi L + U], \quad U \in (0, \pi/4), \quad L \in \mathbb{N}$$

there are vector-valued functions

$$(2.11) \quad \begin{aligned} &(\alpha_0^{3,k_3}, \alpha_1^{3,k_3}, \dots, \alpha_{k_3}^{3,k_3}, \beta_1^{k_3}, \dots, \beta_{k_3}^{k_3}), \\ &1 \leq k_3 \leq k_0, \quad k_3 \in \mathbb{N} \end{aligned}$$

such that the following exact ζ -factorization formula

$$(2.12) \quad \prod_{r=1}^{k_3} \frac{\tilde{Z}^2(\alpha_r^{3,k_3})}{\tilde{Z}^2(\beta_r^{k_3})} = \frac{\sin 2U}{2U} \frac{1}{\cos(2\alpha_0^{3,k_3})}, \quad \forall L \geq L_0$$

holds true, where

$$(2.13) \quad \begin{aligned} \alpha_r^{3,k_3} &= \alpha_r(U, \pi L, k_3; f_3), \quad r = 0, 1, \dots, k_3, \\ \beta_r^{k_3} &= \beta_r(U, \pi L, k_3), \quad r = 1, \dots, k_3, \\ \alpha_0^{2,k_3} &\in (\pi L, \pi L + U), \\ \alpha_r^{3,k_3}, \beta_r^{k_3} &\in (\overbrace{\pi L}^r, \overbrace{\pi L + U}^r), \quad r = 1, \dots, k_3. \end{aligned}$$

Remark 5. Let us remind that

$$\begin{aligned} f(t) \in \tilde{C}_0[\pi L, \pi L + U] &\Leftrightarrow \\ f(t) \in C[\pi L, \pi L + U] \wedge f(t) \geq 0, \neq 0 \wedge U &= o\left(\frac{L}{\ln L}\right), \quad L \rightarrow \infty. \end{aligned}$$

3. THE SET OF COMPLETE HYBRID FORMULAS

3.1. Crossbreeding in the set

$$(3.1) \quad \{(2.3), (2.8)\}, \quad U \in (0, \pi/4)$$

of two ζ -factorization formulas (one stage is sufficient) gives the following (comp. [5])

Exact Complete Hybrid Formula 1.

$$(3.2) \quad \left\{ \prod_{r=1}^{k_1} \frac{\tilde{Z}^2(\alpha_r^{1,k_1})}{\tilde{Z}^2(\beta_r^{k_1})} \right\} \sin^2(\alpha_0^{1,k_1}) + \left\{ \prod_{r=1}^{k_2} \frac{\tilde{Z}^2(\alpha_r^{2,k_2})}{\tilde{Z}^2(\beta_r^{k_2})} \right\} \cos^2(\alpha_0^{2,k_2}) = 1, \\ \forall L \geq L_0, \quad 1 \leq k_1, k_2 \leq k_0, \quad U \in (0, \pi/4).$$

If we use (2.7) and a little of algebra (comp. [8], subsection 8.2) in (3.2) then we obtain the following

Asymptotic Complete Hybrid Formula 1.

$$(3.3) \quad \left\{ \prod_{r=1}^{k_1} \frac{|\zeta(\frac{1}{2} + i\alpha_r^{1,k_1})|^2}{|\zeta(\frac{1}{2} + i\beta_r^{k_1})|^2} \right\} \sin^2(\alpha_0^{1,k_1}) + \left\{ \prod_{r=1}^{k_2} \frac{|\zeta(\frac{1}{2} + i\alpha_r^{2,k_2})|^2}{|\zeta(\frac{1}{2} + i\beta_r^{k_2})|^2} \right\} \cos^2(\alpha_0^{2,k_2}) \\ \sim 1, L \rightarrow \infty.$$

3.2. Next, we make the following crossbreeding in the set

$$(3.4) \quad \{(2.3), (2.8), (2.12)\}, U \in (0, \pi/4)$$

of three ζ -factorization formulas. Namely:

(a) first (2.8) \ominus (2.3) \Rightarrow

$$(3.5) \quad \left\{ \prod_{r=1}^{k_2} \frac{\tilde{Z}^2(\alpha_r^{2,k_2})}{\tilde{Z}^2(\beta_r^{k_2})} \right\} \cos^2(\alpha_0^{2,k_2}) - \left\{ \prod_{r=1}^{k_1} \frac{\tilde{Z}^2(\alpha_r^{1,k_1})}{\tilde{Z}^2(\beta_r^{k_1})} \right\} \sin^2(\alpha_0^{1,k_1}) = \\ = \frac{\sin 2U}{2U},$$

(b) second, the formula (3.5) together with the formula (2.12) imply the following

Exact Complete Hybrid Formula 2.

$$(3.6) \quad \left\{ \prod_{r=1}^{k_2} \frac{\tilde{Z}^2(\alpha_r^{2,k_2})}{\tilde{Z}^2(\beta_r^{k_2})} \right\} \cos^2(\alpha_0^{2,k_2}) - \left\{ \prod_{r=1}^{k_1} \frac{\tilde{Z}^2(\alpha_r^{1,k_1})}{\tilde{Z}^2(\beta_r^{k_1})} \right\} \sin^2(\alpha_0^{1,k_1}) = \\ = \left\{ \prod_{r=1}^{k_3} \frac{\tilde{Z}^2(\alpha_r^{3,k_3})}{\tilde{Z}^2(\beta_r^{k_3})} \right\} \cos(2\alpha_0^{3,k_3}), \\ \forall L \geq L_0, 1 \leq k_1, k_2, k_3 \leq k_0, U \in (0, \pi/4).$$

If we rewrite (3.6) into the form

$$\{\dots\} \cos^2(\alpha_0^{2,k_2}) = \{\dots\} \sin^2(\alpha_0^{1,k_1}) + \{\dots\} \cos(2\alpha_0^{3,k_3})$$

then we obtain, similarly to (3.3) the following

Asymptotic Complete Hybrid Formula 2.

$$(3.7) \quad \left\{ \prod_{r=1}^{k_2} \frac{|\zeta(\frac{1}{2} + i\alpha_r^{2,k_2})|^2}{|\zeta(\frac{1}{2} + i\beta_r^{k_2})|^2} \right\} \cos^2(\alpha_0^{2,k_2}) - \left\{ \prod_{r=1}^{k_1} \frac{|\zeta(\frac{1}{2} + i\alpha_r^{1,k_1})|^2}{|\zeta(\frac{1}{2} + i\beta_r^{k_1})|^2} \right\} \sin^2(\alpha_0^{1,k_1}) \\ \sim \left\{ \prod_{r=1}^{k_3} \frac{|\zeta(\frac{1}{2} + i\alpha_r^{3,k_3})|^2}{|\zeta(\frac{1}{2} + i\beta_r^{k_3})|^2} \right\} \cos(2\alpha_0^{3,k_3}), L \rightarrow \infty.$$

3.3. In connection with the crossbreeding we give the following

Remark 6. Our description of the operation of crossbreeding (see [6], comp. [8], subsection 3.3) contains the following expression:

... that is: after the finite number of eliminations of external functions

$$E_m(U, T), \quad m = 1, \dots, M, \dots$$

However, this is not exact. Exact phrase should be as follows:

... that is: after the finite number of eliminations of the variables U, T from the set of external functions ...

We call these U, T as external variables.

4. THEOREMS ABOUT GRAFTING OF COMPLETE HYBRID FORMULAS INTO META-FUNCTIONAL EQUATIONS

We use the following consequence of the classical H. Bohr's result (see [9], pp. 76, 99).

Lemma 4. There is the infinite set of roots of equation

$$(4.1) \quad \zeta(s) = a, \quad a \neq 0, \quad a \in \mathbb{C}, \quad s = \sigma + it$$

in the strip

$$(4.2) \quad (\sigma_1, \sigma_2) \times (0, +\infty), \quad \frac{1}{2} < \sigma_1 < \sigma_2 < 1.$$

4.1. The formula (3.6) contains the following functions

$$(4.3) \quad \sin^2(\alpha_0^{1,k_1}), \quad \cos^2(\alpha_0^{2,k_2}), \quad \cos(2\alpha_0^{3,k_3})$$

where (see (2.4),(2.9), (2.13))

$$(4.4) \quad \begin{aligned} \alpha_0^{1,k_1} &= \alpha_0(U, \pi L, k_1; f_1) \in (\pi L, \pi L + U), \quad U \in (0, \pi/4), \\ \alpha_0^{2,k_2} &= \alpha_0(U, \pi L, k_2; f_2) \in (\pi L, \pi L + U), \quad U \in (0, \pi/4), \\ \alpha_0^{3,k_3} &= \alpha_0(U, \pi L, k_3; f_3) \in (\pi L, \pi L + U), \quad U \in (0, \pi/4). \end{aligned}$$

Now, (4.3) and (4.4) imply

$$(4.5) \quad \sin^2(\alpha_0^{1,k_1}) \in (0, 1/2), \quad \cos^2(\alpha_0^{2,k_2}) \in (1/2, 1), \quad \cos(2\alpha_0^{3,k_3}) \in (0, 1),$$

i.e. it is true that

$$(4.6) \quad \sin^2(\alpha_0^{1,k_1}), \cos^2(\alpha_0^{2,k_2}), \cos(2\alpha_0^{3,k_3}) \in (0, 1)$$

for every

$$U \in (0, \pi/4), L \geq L_0, (k_1, k_2, k_3) : 1 \leq k_1, k_2, k_3 \leq k_0.$$

4.2. Next, we choose (see (4.1))

$$\sigma_0^1, \sigma_0^2, \sigma_0^3 : \frac{1}{2} < \sigma_1 < \sigma_0^1 < \sigma_0^2 < \sigma_0^3 < \sigma_2 < 1$$

such that

$$(4.7) \quad \begin{aligned} \sigma_1 < \sigma_0^1 - \delta, \quad \sigma_0^3 + \delta < \sigma_2, \\ \sigma_0^1 + \delta < \sigma_0^2 - \delta, \quad \sigma_0^2 + \delta < \sigma_0^3 - \delta \end{aligned}$$

for some sufficiently small $\delta > 0$. Now, in the strip

$$(4.8) \quad S_{\sigma_0^1, \delta}^1 = (\sigma_0^1 - \delta, \sigma_0^1 + \delta) \times (0, +\infty)$$

there is (see Bohr's Lemma 4) for every fixed and admissible parameters

$$U, \pi L, k_1, f_1, \sigma_0^1 - \delta, \sigma_0^1 + \delta$$

infinite set

$$(4.9) \quad \begin{aligned} W_1 &= W_1\{\sin^2[\alpha_0(U, \pi L, k_1, f_1)], \sigma_0^1 - \delta, \sigma_0^1 + \delta\} = \\ &= W_1(U, \pi L, k_1, f_1, \sigma_0^1 - \delta, \sigma_0^1 + \delta) \end{aligned}$$

of elements

$$(4.10) \quad w_1 = w_1(U, \pi L, k_1, f_1, \sigma_0^1 - \delta, \sigma_0^1 + \delta) \in W_1 \subset S_{\sigma_0^1, \delta}^1$$

for which (see (4.1), (4.6))

$$(4.11) \quad \zeta(w_1) = \sin^2(\alpha_0^{1, k_1}) > 0.$$

Since (4.11) \Rightarrow

$$\operatorname{Re}\{\zeta(w_1)\} = \sin^2(\alpha_0^{1, k_1}), \quad \operatorname{Im}\{\zeta(w_1)\} = \sin^2(\alpha_0^{1, k_1}) = 0,$$

then we have the first infinite set of grafts (=equalities)

$$(4.12) \quad \sin^2(\alpha_0^{1, k_1}) = |\zeta(w_1)|, \quad w_1 \in W_1$$

generated by the function

$$\sin^2(\alpha_0^{1, k_1})$$

for every set of fixed and admissible parameters

$$U, \pi L, k_1, f_1, \sigma_0^1 - \delta, \sigma_0^1 + \delta.$$

4.3. Next, similarly to (4.7) – (4.12) we obtain the following objects

$$(4.13) \quad S_{\sigma_0^l, \delta}^l, W_l \subset S_{\sigma_0^l, \delta}^l, \quad l = 2, 3$$

and two other infinite set of grafts

$$(4.14) \quad \begin{aligned} \cos^2(\alpha_0^{2, k_2}) &= |\zeta(w_2)|, \quad w_2 \in W_2, \\ \cos(2\alpha_0^{3, k_3}) &= |\zeta(w_3)|, \quad w_3 \in W_3, \end{aligned}$$

where

$$\begin{aligned} w_2 &= w_2(U, \pi L, k_2, f_2, \sigma_0^2 - \delta, \sigma_0^2 + \delta), \\ w_3 &= w_3(U, \pi L, k_3, f_3, \sigma_0^3 - \delta, \sigma_0^3 + \delta) \end{aligned}$$

for every set of fixed and admissible corresponding parameters.

Remark 7. Of course,

$$\begin{aligned} S_{\sigma_0^l, \delta}^l \cap S_{\sigma_0^k, \delta}^k &= \emptyset, \quad k \neq l; \quad k, l = 1, 2, 3 \Rightarrow \\ W_l \cap W_k &= \emptyset, \quad l \neq k. \end{aligned}$$

4.4. Next, we choose an arbitrary finite set

$$\{U_n\} : 0 < U_1 < U_2 < \dots < U_{n_0} < \frac{\pi}{4}, \quad n_0 \in \mathbb{N}.$$

Every fixed U_n generates the following three infinite sets

$$(4.15) \quad W_1(U_n), \quad W_2(U_n), \quad W_3(U_n), \quad n = 1, \dots, n_0.$$

Now, for every fixed n we choose a single element

$$(4.16) \quad w_1^n, w_2^n, w_3^n, \quad n = 1, \dots, n_0$$

from every corresponding set in (4.15). Consequently, we have (comp. (4.13), (4.14)) the following grafts (=equalities)

$$(4.17) \quad \begin{aligned} \sin^2(\alpha_0^{1, k_1, n}) &= |\zeta(w_1^n)|, \\ \cos^2(\alpha_0^{2, k_2, n}) &= |\zeta(w_2^n)|, \\ \cos(2\alpha_0^{3, k_3, n}) &= |\zeta(w_3^n)|, \\ n &= 1, \dots, n_0 \end{aligned}$$

where (comp. (4.4))

$$\alpha_0^{1, k_1, n} = \alpha_0(U_n, \pi L, k_1; f_1), \dots$$

4.5. Finally, we make a grafting (we use the equalities (4.17) as substitutions) on the exact complete hybrid formula (3.6) (=our tree). As the result of these operations we obtain the following

Theorem 1. For every finite set

$$\{U_n\} : 0 < U_1 < U_2 < \dots < U_{n_0} < \frac{\pi}{4}$$

there are the following elements

$$w_1^n \in S_{\sigma_0^1, \delta}^1, \quad w_2^n \in S_{\sigma_0^2, \delta}^2, \quad w_3^n \in S_{\sigma_0^3, \delta}^3, \quad n = 1, \dots, n_0$$

where

$$(4.18) \quad \begin{aligned} w_1^n &= w_1(U_n, \pi L, k_1, f_1, \sigma_0^1 - \delta, \sigma_0^1 + \delta), \\ w_2^n &= w_2(U_n, \pi L, k_2, f_2, \sigma_0^2 - \delta, \sigma_0^2 + \delta), \\ w_3^n &= w_3(U_n, \pi L, k_3, f_3, \sigma_0^3 - \delta, \sigma_0^3 + \delta), \end{aligned}$$

such that the following exact meta-functional equation holds true:

$$(4.19) \quad \begin{aligned} &\left\{ \prod_{r=1}^{k_2} \frac{\tilde{Z}^2(\alpha_r^{2, k_2, n})}{\tilde{Z}^2(\beta_r^{k_2, n})} \right\} |\zeta(w_2^n)| - \left\{ \prod_{r=1}^{k_1} \frac{\tilde{Z}^2(\alpha_r^{1, k_1, n})}{\tilde{Z}^2(\beta_r^{k_1, n})} \right\} |\zeta(w_1^n)| = \\ &= \left\{ \prod_{r=1}^{k_3} \frac{\tilde{Z}^2(\alpha_r^{3, k_3, n})}{\tilde{Z}^2(\beta_r^{k_3, n})} \right\} |\zeta(w_3^n)|, \\ &\forall L \geq L_0, \quad 1 \leq k_1, k_2, k_3 \leq k_0, \end{aligned}$$

where

$$(4.20) \quad \begin{aligned} \alpha_r^{1,k_1,n} &= \alpha_r(U_n, \pi L, k_1; f_1), \\ \beta_r^{k_1,n} &= \beta_r(U_n, \pi L, k_1), \\ &\dots, \\ n &= 1, \dots, n_0. \end{aligned}$$

4.6. Next, we obtain by the method (4.1) – (4.17) other new-type formula.

Theorem 2. Under the same assumptions as in our Theorem 1 we have the following exact meta-functional equation: (3.2) \Rightarrow

$$(4.21) \quad \left\{ \prod_{r=1}^{k_1} \frac{\tilde{Z}^2(\alpha_r^{1,k_1,n})}{\tilde{Z}^2(\beta_r^{k_1,n})} \right\} |\zeta(w_1^n)| + \left\{ \prod_{r=1}^{k_2} \frac{\tilde{Z}^2(\alpha_r^{2,k_2,n})}{\tilde{Z}^2(\beta_r^{k_2,n})} \right\} |\zeta(w_1^n)| = 1$$

$\forall L \geq L_0, 1 \leq k_1, k_2 \leq k_0.$

5. ASYMPTOTIC META-FUNCTIONAL EQUATIONS

5.1. First, the method (4.1) – (4.17) applied on the asymptotic formula (3.3) gives the following

Corollary 1. The following asymptotic meta-functional equation

$$(5.1) \quad \left\{ \prod_{r=1}^{k_1} \frac{|\zeta(\frac{1}{2} + i\alpha_r^{1,k_1,n})|^2}{|\zeta(\frac{1}{2} + i\beta_r^{k_1,n})|^2} \right\} |\zeta(w_1^n)| + \left\{ \prod_{r=1}^{k_2} \frac{|\zeta(\frac{1}{2} + i\alpha_r^{2,k_2,n})|^2}{|\zeta(\frac{1}{2} + i\beta_r^{k_2,n})|^2} \right\} |\zeta(w_2^n)| \sim$$

$\sim 1, L \rightarrow \infty$

holds true.

5.2. Second, the method (4.1) – (4.17) applied on the asymptotic formula (3.7) gives the following

Corollary 2.

$$(5.2) \quad \left\{ \prod_{r=1}^{k_2} \frac{|\zeta(\frac{1}{2} + i\alpha_r^{2,k_2,n})|^2}{|\zeta(\frac{1}{2} + i\beta_r^{k_2,n})|^2} \right\} |\zeta(w_2^n)| - \left\{ \prod_{r=1}^{k_1} \frac{|\zeta(\frac{1}{2} + i\alpha_r^{1,k_1,n})|^2}{|\zeta(\frac{1}{2} + i\beta_r^{k_1,n})|^2} \right\} |\zeta(w_1^n)| \sim$$

$$\sim \left\{ \prod_{r=1}^{k_3} \frac{|\zeta(\frac{1}{2} + i\alpha_r^{3,k_3,n})|^2}{|\zeta(\frac{1}{2} + i\beta_r^{k_3,n})|^2} \right\} |\zeta(w_3^n)|,$$

$L \rightarrow \infty.$

5.3. Now we obtain (see (5.1), (5.2)) the following

Corollary 3.

$$(5.3) \quad \left\{ \prod_{r=1}^{k_2} \frac{|\zeta(\frac{1}{2} + i\alpha_r^{2,k_2,n})|^2}{|\zeta(\frac{1}{2} + i\beta_r^{k_2,n})|^2} \right\} |\zeta(w_2^n)| \sim$$

$$\sim \frac{1}{2} + \frac{1}{2} \left\{ \prod_{r=1}^{k_3} \frac{|\zeta(\frac{1}{2} + i\alpha_r^{3,k_3,n})|^2}{|\zeta(\frac{1}{2} + i\beta_r^{k_3,n})|^2} \right\} |\zeta(w_3^n)|, L \rightarrow \infty,$$

$$\begin{aligned}
(5.4) \quad & \left\{ \prod_{r=1}^{k_1} \frac{|\zeta(\frac{1}{2} + i\alpha_r^{1,k_1,n})|^2}{|\zeta(\frac{1}{2} + i\beta_r^{k_1,n})|^2} \right\} |\zeta(w_1^n)| \sim \\
& \sim \frac{1}{2} - \frac{1}{2} \left\{ \prod_{r=1}^{k_3} \frac{|\zeta(\frac{1}{2} + i\alpha_r^{3,k_3,n})|^2}{|\zeta(\frac{1}{2} + i\beta_r^{k_3,n})|^2} \right\} |\zeta(w_3^n)|, \quad L \rightarrow \infty.
\end{aligned}$$

Remark 8. The set of asymptotic meta-functional equations (5.1) – (5.4) represents the higher level of ζ -encoding (in comparison with this one that represents the corresponding set of asymptotic complete hybrid formulas) of the set of elementary trigonometric formulas

$$\begin{aligned}
\sin^2 x + \cos^2 x &= 1, \quad \cos^2 x - \sin^2 x = \cos 2x, \\
\cos^2 x &= \frac{1}{2} + \frac{1}{2} \cos 2x, \quad \sin^2 x = \frac{1}{2} - \frac{1}{2} \cos 2x.
\end{aligned}$$

5.4. Putting in (5.2)

$$(5.5) \quad k_1 = k_2 = k_3 = k; \quad 1 \leq k \leq k_0,$$

we obtain the following

Corollary 4.

$$\begin{aligned}
(5.6) \quad & \left\{ \prod_{r=1}^k \left| \zeta\left(\frac{1}{2} + i\alpha_r^{2,k,n}\right) \right|^2 \right\} |\zeta(w_2^n)| - \left\{ \prod_{r=1}^k \left| \zeta\left(\frac{1}{2} + i\alpha_r^{1,k,n}\right) \right|^2 \right\} |\zeta(w_1^n)| \sim \\
& \sim \left\{ \prod_{r=1}^k \left| \zeta\left(\frac{1}{2} + i\alpha_r^{3,k,n}\right) \right|^2 \right\} |\zeta(w_3^n)|, \quad L \rightarrow \infty.
\end{aligned}$$

Remark 9. We shall call the formula (5.6) as the secondary asymptotic meta-functional equation since it is the result of immediate metamorphosis of the formula (5.2) in the case (5.5).

Remark 10. The case $k = 1$ in (5.5) gives the formula whose simple variant is the formula (1.2) in the Introduction.

6. SYNERGETIC FORMULAS ONLY FOR RIEMANN'S ZETA-FUNCTION: INTERACTIONS BETWEEN SOME PARTS OF THIS FUNCTION

6.1. Let us remind that every element from the complete hybrid formulas set (3.2), (3.3), (3.5) – (3.7) is synergetic one - see our interpretation of this phenomenon in our paper [8]. Really, every from these formulas is the result of an interaction controlled by the Jacob's ladder $\varphi_1(t)$ between following sets

$$\left\{ \left| \zeta\left(\frac{1}{2} + it\right) \right|^2 \right\}, \quad \{\sin^2 t\}, \quad \{\cos^2 t\}, \quad \{\cos 2t\}, \quad t \geq L_0.$$

6.2. In this paper, we have introduced some interactions of Riemann's zeta-function with itself on the critical strip. Namely, the interactions between following four parts

$$(6.1) \quad \left\{ \left| \zeta\left(\frac{1}{2} + it\right) \right|^2 \right\}, \quad t \geq L_0, \quad |\zeta(s)|, \quad s \in S_{\sigma_0^l, \delta}^l, \quad l = 1, 2, 3$$

of the Riemann's zeta-function

$$\zeta(s), \quad s \in \mathbb{C} \setminus \{1\}.$$

Here, of course,

$$S_{\sigma_0, \delta}^l \subset (\sigma_1, \sigma_2) \times (0, +\infty), \quad S_{\sigma_0, \delta}^l \cap S_{\sigma_0, \delta}^k = \emptyset, \quad k \neq l, \quad k, l = 1, 2, 3.$$

Let us call the set of these interactions as the ζ -chemical reaction of the *substances* (6.1) - this is some analogue to what we have based on the classical Belousov-Zhabotinski chemical reactions.

Remark 11. The result of these ζ -chemical reactions (= ζ -chemical compounds) are our synergetic meta-functional equations (4.19), (4.21) as well as the formulae (5.1) - (5.4), (5.6).

Remark 12. Since the Planck length reads

$$L_P = 8.1 \times 10^{-35} \text{ cm}$$

and the Planck time reads

$$T_P = 2.7 \times 10^{-43} \text{ s}$$

then, from a point of view of the sciences, we may use (for example) the following complementary condition on the set

$$\{U_n\} : U_{n+1} - U_n > 10^{-43}, \quad n = 1, \dots, n_0 - 1, U_1, \frac{\pi}{4} - U_{n_0} > 10^{-43}.$$

This condition corresponds with the point of view of Jakov Zeldovich for using maths on study of real-world phenomena.

I would like to thank Michal Demetrian for his moral support of my study of Jacob's ladders.

REFERENCES

- [1] J. Moser, 'Jacob's ladders and almost exact asymptotic representation of the Hardy-Littlewood integral', Math. Notes 88, (2010), 414-422, arXiv: 0901.3937.
- [2] J. Moser, 'Jacob's ladders, structure of the Hardy-Littlewood integral and some new class of nonlinear integral equations', Proc. Steklov Inst. 276 (2011), 208-221, arXiv: 1103.0359.
- [3] J. Moser, 'Jacob's ladders, reverse iterations and new infinite set of L_2 -orthogonal systems generated by the Riemann zeta-function, arXiv: 1402.2098.
- [4] J. Moser, 'Jacob's ladders, factorization and metamorphoses as an appendix to the Riemann functional equation for $\zeta(s)$ on the critical line', Proc. Steklov Inst. 296 (2017), pp. 92-102, arXiv: 1506.00442v1.
- [5] J. Moser, 'Jacob's ladders, interactions between ζ -oscillating systems and ζ -analogue of an elementary trigonometric identity', arXiv: 1609.09293v1, Proc. Steklov Inst. 299, 189-204, 2017.
- [6] J. Moser, 'Jacob's ladders, crossbreeding in the set of ζ -factorization formulas and the selection of families of ζ -kindred real continuous functions', arXiv: 1710.04428 .
- [7] J. Moser, 'Jacob ladders and new families of ζ -kindred real continuous functions', arXiv: 1801.09425v1.
- [8] J. Moser, 'Jacob ladders, crossbreeding, secondary crossbreeding and synergetic phenomena generated by the Riemann's zeta-function and some elementary functions on disconnected sets of the critical line', arXiv: 1806.07095v1.
- [9] E.C. Titchmarsh, 'The zeta-function of Riemann', Cambridge, 1930.

DEPARTMENT OF MATHEMATICAL ANALYSIS AND NUMERICAL MATHEMATICS, COMENIUS UNIVERSITY, MLYNSKA DOLINA M105, 842 48 BRATISLAVA, SLOVAKIA

E-mail address: jan.mozer@fmph.uniba.sk