NON-REMOVABILITY OF SIERPIŃSKI CARPETS

DIMITRIOS NTALAMPEKOS

ABSTRACT. We prove that all Sierpiński carpets in the plane are non-removable for (quasi)conformal maps. More precisely, we show that for any two Sierpiński carpets $S, S' \subset \widehat{\mathbb{C}}$ there exists a homeomorphism $f \colon \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ that is conformal in $\widehat{\mathbb{C}} \setminus S$ and it maps S onto S'. As a corollary, we obtain a partial answer to a question of Bishop [Bis94], whether any planar compact set with empty interior and positive measure can be mapped to a set of measure zero with an exceptional homeomorphism of the plane, conformal off that set.

1. Introduction

In recent work [Nta19, Theorem 1.8] the current author proved that the Sierpiński gasket, also called the Sierpiński triangle, is non-removable for conformal maps. In the same paper the author provided some evidence that *all* homeomorphic copies of the Sierpiński gasket should be non-removable for conformal maps; see [Nta19, Theorem 1.7]. This gave birth to the conception that "some sets should be non-removable for topological reasons". Sierpiński carpets are topologically "larger" than gaskets and provide a perfect candidate to test this heuristic. In this work we prove that this is actually the case:

Theorem 1.1. All Sierpiński carpets $S \subset \widehat{\mathbb{C}}$ are non-removable for conformal maps.

This result resolves a conjecture of the author [Nta19, Conjecture 1]. Here, we pose another more general conjecture:

Conjecture 1.1. Every planar compact set containing a homeomorphic copy of $C \times [0,1]$ is non-removable for conformal maps, where C is the middle-thirds Cantor set.

We provide some necessary definitions before stating our other results.

Definition 1.2. Let $K \subset \widehat{\mathbb{C}}$ be a compact set. We say that K is *conformally removable* or *removable for conformal maps* if any homeomorphism $f : \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ that is conformal in $\widehat{\mathbb{C}} \setminus K$ is actually conformal in $\widehat{\mathbb{C}}$ and thus, it is Möbius.

An equivalent notion is that of quasiconformal removability. We direct the reader to [You15], [Nta19], and the references therein for more background.

The standard Sierpiński carpet is constructed by subdividing the unit square $[0,1]^2$ into nine squares of sidelength 1/3 and removing the middle square, and

Date: September 5, 2019.

²⁰¹⁰ Mathematics Subject Classification. Primary 30C20, 30C35; Secondary 30C62.

Key words and phrases. Sierpiński carpets, Removability, Conformal maps, Quasiconformal maps.

The author was partially supported by NSF Grant DMS-1506099.

then proceeding inductively in each of the remaining eight squares. It can be easily proved that the standard Sierpiński carpet is non-removable for conformal maps; see e.g. the discussion in [Nta19, Section 1.1].

In general, a Sierpiński carpet $S \subset \widehat{\mathbb{C}}$ is a set homeomorphic to the standard carpet. It is a fundamental result of Whyburn [Why58] that a set S is a Sierpiński carpet if and only if S has empty interior and $S = \widehat{\mathbb{C}} \setminus \bigcup_{i \in \mathbb{N}} Q_i$, where $\{Q_i\}_{i \in \mathbb{N}}$ is a family of Jordan regions with disjoint closures and (spherical) diameters converging to 0. The regions Q_i , $i \in \mathbb{N}$, are called the peripheral disks and the boundaries ∂Q_i , $i \in \mathbb{N}$, are called the peripheral circles of S. Our main result is the following.

Theorem 1.3. Let $S, S' \subset \widehat{\mathbb{C}}$ be Sierpiński carpets. Then there exists a homeomorphism $f \colon \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ with f(S) = S' such that f is conformal on $\widehat{\mathbb{C}} \setminus S$. Moreover, the image of any finite collection of peripheral circles can be prescribed: if C_1, \ldots, C_N and C'_1, \ldots, C'_N are peripheral circles of S and S', respectively, then f can be chosen such that $f(C_i) = C'_i$ for all $i = 1, \ldots, N$.

One can easily construct carpets S' with positive Lebesgue measure. The theorem, combined with this fact, proves immediately Theorem 1.1, since a map f that sends a carpet S of measure zero or positive measure to a carpet S' of positive measure or measure zero, respectively, cannot be Möbius. Our proof of Theorem 1.3 is topological and utilizes the ideas of Whyburn [Why58].

Using Theorem 1.3 we obtain a partial answer to a question raised by Bishop [Bis94, Question 3]. He asked whether any compact set $K \subset \mathbb{C}$ with empty interior and positive area can be mapped to a set of measure zero with a homeomorphism $f: \mathbb{C} \to \mathbb{C}$ that is conformal on $\mathbb{C} \setminus K$. A partial answer to that question was given by Kaufman and Wu [KW96, Theorem 3], where the authors proved that there exists a *subset* of K with positive, strictly smaller measure such that the answer to the question is positive for that subset. We prove that one can "enlarge" K to a set L by attaching to it a small set of measure zero and Hausdorff dimension arbitrarily close to 1, so that the answer is positive for L:

Proposition 1.4. Let $K \subset \mathbb{C}$ be a compact set with empty interior. Then for each $\varepsilon > 0$ there exists a compact set $L \supset K$ such that the Hausdorff dimension of $L \setminus K$ is at most $1 + \varepsilon$ and the following holds: there exists a homeomorphism $f : \mathbb{C} \to \mathbb{C}$ that is conformal on $\mathbb{C} \setminus L$ such that f(L) has Hausdorff dimension at most $1 + \varepsilon$ and in particular Lebesque measure zero.

In fact L can be taken to lie in the ε -neighborhood of K if we allow f to be quasiconformal, but we will not go into details for the sake of brevity.

Acknowledgments. I would like to thank my PhD advisor Mario Bonk at UCLA for many useful conversations on the problem of non-removability of carpets and for his notes on the proof of Whyburn, which were very enlightening. I also thank Stanislav Smirnov for reigniting my interest on the problem by asking me the question whether carpets are removable and the anonymous referee for reading the manuscript carefully and providing useful comments.

2. Proof of Theorem 1.3

In the entire section we only use the spherical metric of $\widehat{\mathbb{C}}$. We start with the following definition adapted from [Why58].

Definition 2.1. Let $\varepsilon > 0$ and Y be a closed subset of $\widehat{\mathbb{C}}$. A partition $\{Y_{\alpha}\}$ of Y is called an ε -subdivision if the collection $\{Y_{\alpha}\}$ consists of finitely many closed Jordan regions, also called 2-cells, with disjoint interiors and diameter less than ε .

A partition $\{S_{\alpha}\}$ of a Sierpiński carpet $S \subset \widehat{\mathbb{C}}$ into Sierpiński carpets is an ε -subdivision (rel. Q_1, \ldots, Q_N) if there exist peripheral disks Q_1, \ldots, Q_N of S and an ε -subdivision $\{Y_{\alpha}\}$ of the closed domain $Y := \widehat{\mathbb{C}} \setminus \bigcup_{i=1}^N Q_i$, for which

- (1) the boundaries of the 2-cells Y_{α} are contained in $S \setminus \bigcup_{i=N+1}^{\infty} \partial Q_i$, where Q_i , $i \geq N+1$, are the remaining peripheral disks of S, and
- (2) $\{S_{\alpha}\}=\{S\cap Y_{\alpha}\}$, i.e., the Sierpiński carpets in the subdivision $\{S_{\alpha}\}$ arise as the intersections of the 2-cells in the subdivision $\{Y_{\alpha}\}$ with the original Sierpiński carpet S.

We remark that each peripheral disk of S is either one of the disks Q_1, \ldots, Q_N , or it is a peripheral disk of one of the carpets S_{α} in the ε -subdivision of S.

Lemma 2.2. Let $S, S' \subset \widehat{\mathbb{C}}$ be Sierpiński carpets. Fix peripheral disks Q_1, \ldots, Q_N and Q'_1, \ldots, Q'_N of S and S', respectively. Then for each $\varepsilon > 0$ there exists M > N and peripheral disks Q_{N+1}, \ldots, Q_M and Q'_{N+1}, \ldots, Q'_M of S and S', respectively, such that there exists an ε -subdivision of S rel. Q_1, \ldots, Q_M and an ε -subdivision of S' rel. Q'_1, \ldots, Q'_M . Moreover, for each family of orientation-preserving homeomorphisms $g_i \colon \overline{Q_i} \to \overline{Q'_i}$, $i \in \{1, \ldots, M\}$, the ε -subdivisions can be taken in such a way that they correspond to each other under a homeomorphism $g \colon \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ that extends $g_i, i \in \{1, \ldots, M\}$.

Here the correspondence of the subdivisions under g is be understood in the sense that each 2-cell Y_{α} of the subdivision of S is mapped by g onto a 2-cell Y'_{α} of the subdivision of S'.

A weaker version of this lemma is used by Whyburn in his topological characterization of the carpet; see [Why58, Lemma 1]. For the sake of completeness and to make this work self-contained, we include an outline of the proof of Lemma 2.2 later in Section 4.

Proof of Theorem 1.3. Let $\varepsilon_k=1/k$. We fix peripheral circles C_1,\ldots,C_N and C'_1,\ldots,C'_N of S and S', respectively, as in the statement of the theorem, and we consider conformal maps $g_i,\ i\in\{1,\ldots,N\}$, from the peripheral disk Q_i of S bounded by C_i onto the peripheral disk Q'_i of S' bounded by C'_i . These conformal maps extend to the boundaries by Carathéodory's theorem and provide orientation-preserving homeomorphisms $g_i\colon \overline{Q_i}\to \overline{Q'_i}$. By Lemma 2.2, we can find peripheral disks Q_{N+1},\ldots,Q_M and Q'_{N+1},\ldots,Q'_M of S and S', respectively, with conformal maps $g_i\colon \overline{Q_i}\to \overline{Q'_i},\ i\in\{N+1,\ldots,M\}$, and we can find ε_1 -subdivisions of S and S' that correspond to each other under a global homeomorphic extension $f_1\colon \widehat{\mathbb{C}}\to \widehat{\mathbb{C}}$ of the maps $g_i\colon \overline{Q_i}\to \overline{Q'_i}$. Observe that $\mathrm{dist}(f_1(S),S')\leq \varepsilon_1$ and $\mathrm{dist}(S,f_1^{-1}(S'))\leq \varepsilon_1$.

Next, we fix one of the Sierpiński carpets S_{α} in the ε_1 -subdivision of S that corresponds to a carpet S'_{α} in the ε_1 -subdivision of S'; that is, the corresponding 2-cells are mapped to each other under f_1 (however, we do not necessarily have $f_1(S_{\alpha}) = S'_{\alpha}$). Note that the peripheral circles of S_{α} and S'_{α} necessarily have diameters bounded above by ε_1 .

There is a distinguished peripheral circle of S_{α} (S'_{α}) that separates it from the other carpets in the subdivision of S (S'). We call R_1 (R'_1) the peripheral disk of S_{α} (S'_{α}) bounded by that peripheral circle. Consider the orientation-preserving

homeomorphism $h_1 := f_1|_{\overline{R_1}} : \overline{R_1} \to \overline{R'_1}$. Now we apply Lemma 2.2 on S_{α} and S'_{α} , to obtain peripheral disks R_2, \ldots, R_K and R'_2, \ldots, R'_K of S_{α} and S'_{α} , respectively, together with conformal maps $h_i : \overline{R_i} \to \overline{R'_i}$, $i \in \{2, \ldots, K\}$, and we can find ε_2 -subdivisions of S_{α} and S'_{α} (rel. R_i and R'_i , $i = 1, \ldots, K$, respectively) that correspond to each other under a homeomorphic extension $f_{\alpha,2} : \widehat{\mathbb{C}} \setminus R_1 \to \widehat{\mathbb{C}} \setminus R'_1$ of the maps h_i , $i \in \{2, \ldots, K\}$.

We repeat this procedure for each of the carpets S_{α} in the ε_1 -subdivision of S. If we collect the ε_2 -subdivisions of the carpets of the form $S_{\alpha} \subset S$ and $S'_{\alpha} \subset S'$, we obtain ε_2 -subdivisions of the carpets S and S'. Patching together the resulting maps of the form $f_{\alpha,2}$ yields a homeomorphism $f_2 \colon \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ under which the ε_1 - and ε_2 -subdivisions of S, S' are in correspondence. The map f_2 agrees with f_1 on $\bigcup_{i=1}^M \overline{Q_i}$ and maps conformally the peripheral disks of S having diameter larger than ε_2 onto some peripheral disks of S'. Note that by construction f_2^{-1} maps conformally the peripheral disks of S' having diameter larger than ε_2 onto some peripheral disks of S. Moreover, the L^{∞} -distance of f_1 and f_2 is bounded by ε_1 ; the same statement holds for the inverses of the maps. Also, note that $\operatorname{dist}(f_2(S), S') \leq \varepsilon_2$ and $\operatorname{dist}(S, f_2^{-1}(S')) \leq \varepsilon_2$.

We repeat the procedure of the last two paragraphs for each carpet in the ε_2 -subdivision of S, S'. Inductively, for each $k \in \mathbb{N}$ we obtain a homeomorphism f_k of $\widehat{\mathbb{C}}$ with the following properties:

- (1) The L^{∞} -distance of f_k , f_m and the L^{∞} -distance of f_k^{-1} , f_m^{-1} are bounded by ε_k for $m \geq k$.
- (2) $\operatorname{dist}(f_k(S), S') \leq \varepsilon_k$ and $\operatorname{dist}(S, f_k^{-1}(S')) \leq \varepsilon_k$ for all $k \in \mathbb{N}$.
- (3) For each $k \in \mathbb{N}$ the map f_k maps conformally peripheral disks Q_{i_1}, \ldots, Q_{i_k} of S onto corresponding peripheral disks $Q'_{i_1}, \ldots, Q'_{i_k}$ of S'. In fact, for each $l \in \mathbb{N}$ the sequence $\{f_k\}_{k \in \mathbb{N}}$ is eventually constant on Q_{i_l} .
- (4) The sequences of peripheral disks $Q_{i_k}, Q'_{i_k}, k \in \mathbb{N}$, exhaust the sequences of peripheral disks of S, S', respectively.

Since $\varepsilon_k \to 0$, (1) implies that f_k converges uniformly in $\widehat{\mathbb{C}}$ to a homeomorphism f of $\widehat{\mathbb{C}}$. By (2), we have f(S) = S'. Finally, (3) and (4) imply that each peripheral disk of S is mapped conformally onto a peripheral disk of S'. In particular, f is conformal on $\widehat{\mathbb{C}} \setminus S$.

3. Proof of Proposition 1.4

An easily verifiable fact that we will use here is that for each $\varepsilon > 0$ there exists a self-similar square carpet $S \subset \mathbb{C}$ with Hausdorff dimension bounded above by $1 + \varepsilon$.

We fix $\varepsilon > 0$. Suppose that $K \subset \mathbb{C}$ is a compact set with empty interior and let B be a large ball containing K. We consider a dyadic square decomposition of the open set $B \setminus K$. In each dyadic square $Q \subset B \setminus K$ in this decomposition we consider a square carpet S_Q with Hausdorff dimension bounded by $1 + \varepsilon$, scaled so that it "fits" in the square S; that is to say that the boundary of the unbounded peripheral disk of S_Q is precisely the boundary of Q. Then we take L to be the closure of the union of all carpets S_Q .

Note that $L \supset K$, since $\operatorname{int}(K) = \emptyset$, and that L is a Sierpiński carpet, as one can see by Whyburn's criterion (see the Introduction): $\operatorname{int}(L) = \emptyset$ (since $\operatorname{int}(K) = \emptyset$) and the complement of L in $\widehat{\mathbb{C}}$ consists of countably many Jordan regions with disjoint closures and diameters shrinking to 0. Moreover, $L \setminus K$ has Hausdorff

dimension at most $1+\varepsilon$, since it is the union of the circle ∂B with countably many carpets S_Q , each having Hausdorff dimension at most $1+\varepsilon$.

Now, we apply Theorem 1.3, which provides us with a homeomorphism $f\colon \widehat{\mathbb{C}} \to \widehat{\mathbb{C}}$ that maps the carpet L onto a fixed square carpet with Hausdorff dimension at most $1+\varepsilon$. In particular, f(L) has Lebesgue measure zero. By post-composing with a Möbius transformation of $\widehat{\mathbb{C}}$, which does not affect the Hausdorff dimension, we may assume that $f(\infty) = \infty$ and thus, we have a homeomorphism $f\colon \mathbb{C} \to \mathbb{C}$ that is conformal in $\mathbb{C} \setminus L$, as desired.

4. Outline of proof of Lemma 2.2

The proof relies on the following version of Moore's theorem [Moo25]:

Theorem 4.1. [Dav86, Corollary 6A, p. 56] Let $\{R_i\}_{i\in\mathbb{N}}$ be a sequence of Jordan regions in the sphere $\widehat{\mathbb{C}}$ with mutually disjoint closures and diameters converging to 0, and consider an open set $\Omega \supset \bigcup_{i\in\mathbb{N}} \overline{R_i}$. Then there exists a continuous, surjective map $F: \widehat{\mathbb{C}} \to S^2$ that is the identity outside Ω and it induces the decomposition of $\widehat{\mathbb{C}}$ into the sets $\{\overline{R_i}\}_{i\in\mathbb{N}}$ and points. In other words, there are countably many distinct points $p_i \in S^2$, $i \in \mathbb{N}$, such that $F^{-1}(p_i) = \overline{R_i}$ for $i \in \mathbb{N}$ and F is injective on $\widehat{\mathbb{C}} \setminus \bigcup_{i\in\mathbb{N}} \overline{R_i}$ with $F(\widehat{\mathbb{C}} \setminus \bigcup_{i\in\mathbb{N}} \overline{R_i}) = S^2 \setminus \{p_i : i \in \mathbb{N}\}$.

Here S^2 is identical to $\widehat{\mathbb{C}}$, but it will be more convenient to have different notation for the target and view S^2 mostly from a topological point of view.

First, we consider the peripheral disks Q_1, \ldots, Q_N of S and Q'_1, \ldots, Q'_N of S' given in the statement of Lemma 2.2 and fix $\varepsilon > 0$. We append to these collections some peripheral disks Q_{N+1}, \ldots, Q_M of S and Q'_{N+1}, \ldots, Q'_M of S' so that the remaining peripheral disks of S and S' have diameters less than ε . Now, we apply Theorem 4.1 to the region $\Omega = \widehat{\mathbb{C}} \setminus \bigcup_{i=1}^M \overline{Q_i}$ and to the remaining peripheral disks $\{R_i\}_{i\in\mathbb{N}}$ of S contained in Ω , all of which have diameter less than ε . This yields a "collapsing" map $F \colon \widehat{\mathbb{C}} \to S^2$ that collapses each R_i to a point p_i , but is injective otherwise. We apply analogously Theorem 4.1 to the carpet S' and obtain a map F' that collapses peripheral disks R'_i to points p'_i .

Now, we consider given orientation-preserving homeomorphisms $g_i \colon \overline{Q_i} \to \overline{Q'_i}$, $i \in \{1, ..., M\}$, as in the statement of Lemma 2.2. These homeomorphisms "project" down to homeomorphisms

$$\widetilde{g}_i := F' \circ g_i \circ \left(F |_{\overline{Q_i}} \right)^{-1} : F(\overline{Q_i}) \to F'(\overline{Q_i'}), \quad 1 \le i \le M.$$

These are orientation-preserving homeomorphisms between finitely many disjoint Jordan regions in S^2 . This implies that there exists a homeomorphic extension $\tilde{g} \colon S^2 \to S^2$ of \tilde{g}_i , $i \in \{1, \dots, M\}$.

Consider the countable set $A' = \{\widetilde{g}(p_i) : i \in \mathbb{N}\} \cup \{p_i' : i \in \mathbb{N}\} \subset S^2$. This is the union of the set of points p_i' arising from collapsing the peripheral disks of S' (under F') that are different from Q_1', \ldots, Q_M' , together with the images $\widetilde{g}(p_i)$ of the corresponding points p_i arising from collapsing the peripheral disks of S (under F). We fix a $\delta > 0$ (to be specified) and a δ -subdivision $\{Y_\alpha'\}$ of the closed domain $Y' = S^2 \setminus \bigcup_{i=1}^M F'(Q_i')$ so that the boundaries $\partial Y_\alpha'$ avoid the countable set A'. We fix a Jordan region Y_α' and consider the preimage $(F')^{-1}(\partial Y_\alpha')$, which is a

We fix a Jordan region Y'_{α} and consider the preimage $(F')^{-1}(\partial Y'_{\alpha})$, which is a Jordan curve since F' is injective on it. Let $X'_{\alpha} \subset \widehat{\mathbb{C}}$ be the Jordan region bounded by that Jordan curve with the property that $(F')^{-1}(p'_i) \subset X'_{\alpha}$ for some $p'_i \in Y'_{\alpha}$;

in fact, this will hold for all $p_i' \in Y_\alpha'$, as one can see using homotopy arguments. Then the Jordan regions $\{X_\alpha'\}$ give a subdivision of the closed domain $\widehat{\mathbb{C}} \setminus \bigcup_{i=1}^M \overline{Q_i'}$ in the sense of Definition 2.1 and the boundaries of the 2-cells X_α' are contained in $S' \setminus \bigcup_{i=M+1}^\infty \partial Q_i'$ (here Q_i' , $i \geq M+1$, are the remaining peripheral disks of S'). Finally, the sets $S_\alpha' := S' \cap Y_\alpha'$ are Sierpiński carpets by Whyburn's criterion (stated before Theorem 1.3), since $S_\alpha' \subset S'$ has empty interior and the complement of S_α' consists of Jordan regions with disjoint closures, whose diameters converge to 0; what is important here is that $\partial X_\alpha' = (F')^{-1}(\partial Y_\alpha')$ does not intersect any peripheral circle $\partial Q_i'$, $i \geq M+1$. Therefore, the collection $\{S_\alpha'\}$ is a subdivision of the carpet S', as in Definition 2.1. One can ensure that this subdivision is an ε -subdivision by choosing a sufficiently small $\delta > 0$ in the previous paragraph and using the following elementary lemma:

Lemma 4.2. Suppose that X and Y are compact metric spaces and $h: X \to Y$ is a continuous, surjective map with the property that the preimage of each point has diameter at most a given number $\varepsilon > 0$. Then there exists a $\delta > 0$ such that for each set $B \subset Y$ with $\operatorname{diam}(B) < \delta$ we have $\operatorname{diam}(h^{-1}(B)) < \varepsilon$.

Now, we consider the preimages Y_{α} under \widetilde{g} of the Jordan regions Y'_{α} lying in the partition of $Y' = S^2 \setminus \bigcup_{i=1}^M F'(Q'_i)$. The sets Y_{α} provide a partition of $Y = S^2 \setminus \bigcup_{i=1}^M F(Q_i)$ into Jordan regions and the boundaries ∂Y_{α} of the Jordan regions in the partition avoid the countable set $A = \{p_i : i \in \mathbb{N}\}$. One now takes preimages under F of the Jordan curves ∂Y_{α} ; these provide a subdivision of S as in the previous paragraph. We denote by $X_{\alpha} \subset \widehat{\mathbb{C}}$ the Jordan region bounded by $F^{-1}(\partial Y_{\alpha})$ that contains a carpet S_{α} in the subdivision of S. Again, if δ is chosen to be sufficiently small, then by continuity the regions Y_{α} will be small enough, so that the preceding lemma can guarantee that the sets X_{α} yield an ε -subdivision of S.

The map \widetilde{g} (composed appropriately with F and $(F')^{-1}$) provides a homeomorphism g from the union of the Jordan curves of the form ∂X_{α} onto the union of the Jordan curves of the form $\partial X'_{\alpha}$. This homeomorphism extends the maps $g_i \colon \partial Q_i \to \partial Q'_i$, $i \in \{1, \ldots, M\}$. The union of the Jordan curves ∂X_{α} is the 1-skeleton of a 2-cell decomposition of $\widehat{\mathbb{C}}$, in which the 2-cells are the sets X_{α} , together with the sets $\overline{Q_i}$, $i \in \{1, \ldots, M\}$. The analogous statement holds for X'_{α} and $\overline{Q'_i}$, $i \in \{1, \ldots, M\}$. Now, one can extend the homeomorphism g to a homeomorphism of $\widehat{\mathbb{C}}$; see e.g. [BM17, Chapter 5.2]. By construction, the ε -subdivisions of S and S' are in correspondence under g, in the sense of the comments following the statement of Lemma 2.2.

References

- [Bis94] C. J. Bishop, Some homeomorphisms of the sphere conformal off a curve, Ann. Acad. Sci. Fenn. Ser. A I Math. 19 (1994), no. 2, 323–338.
- [BM17] M. Bonk and D. Meyer, Expanding Thurston maps, Mathematical Surveys and Monographs, vol. 225, American Mathematical Society, Providence, RI, 2017.
- [Dav86] R. J. Daverman, Decompositions of manifolds, Pure and Applied Mathematics, vol. 124, Academic Press, Inc., Orlando, FL, 1986.
- [KW96] R. Kaufman and J.-M. Wu, On removable sets for quasiconformal mappings, Ark. Mat. 34 (1996), no. 1, 141–158.
- [Moo25] R. L. Moore, Concerning upper semi-continuous collections of continua, Trans. Amer. Math. Soc. 27 (1925), no. 4, 416–428.

- [Nta19] D. Ntalampekos, Non-removability of the Sierpiński gasket, Invent. Math. 216 (2019), no. 2, 519–595.
- [Why58] G. T. Whyburn, Topological characterization of the Sierpiński curve, Fund. Math. 45 (1958), 320–324.
- [You15] M. Younsi, On removable sets for holomorphic functions, EMS Surv. Math. Sci. 2 (2015), no. 2, 219–254.

Institute for Mathematical Sciences, Stony Brook University, Stony Brook, NY 11794, USA.

 $E\text{-}mail\ address: \verb|dimitrios.ntalampekos@stonybrook.edu||$