

## A REMARK ON THE EXTENSION OF $L^2$ HOLOMORPHIC FUNCTIONS

QI'AN GUAN

ABSTRACT. In this note, we answer a question on the extension of  $L^2$  holomorphic functions posed by Ohsawa.

### 1. AN ANSWER TO A QUESTION POSED BY OHSAWA

In [7], Ohsawa gave a survey on a recent "remarkable" progress (c.f. [1, 2, 3, 4, 5, 6]) around the famous Ohsawa-Takegoshi  $L^2$  extension theorem [8]. After that, Ohsawa recalled the following consequence of the main result in [4], and presented a shorter proof based on a general criterion for the extendibility in [7].

**Theorem 1.1.** ([4], see also Theorem 0.1 in [7]) *Let  $D \subset \mathbb{C}^n$  be a pseudoconvex domain, and let  $\varphi$  be a plurisubharmonic function on  $D$  and  $H = \{z_n = 0\}$ . Then for any holomorphic function  $f$  on  $H$  satisfying*

$$\int_H |f|^2 e^{-\varphi} dV_H < \infty,$$

*there exists a holomorphic function  $F$  on  $D$  satisfying  $F = f$  on  $H$  and*

$$\int_D |F|^2 e^{-\varphi - (1+\varepsilon) \log(1+|z_n|^2)} dV_D \leq \frac{\pi}{\varepsilon} \int_H |f|^2 e^{-\varphi} dV_H.$$

In [7], considering general plurisubharmonic function  $\psi(z_n)$  instead of  $(1 + \varepsilon) \log(1 + |z_n|^2)$  in Theorem 1.3, Ohsawa posed the following question on the extension of  $L^2$  holomorphic functions.

**Question 1.2.** *Given a subharmonic function  $\psi$  on  $\mathbb{C}$  such that  $\int_{\mathbb{C}} e^{-\psi} < +\infty$ , for any subharmonic function  $\varphi$  on  $\mathbb{C}$ , can one find a holomorphic function  $f$  on  $\mathbb{C}$  satisfying  $f(0) = 1$ , and*

$$\int_{\mathbb{C}} |f|^2 e^{-\varphi - \psi} \leq e^{-\varphi(0)} \int_{\mathbb{C}} e^{-\psi}?$$

When  $\psi$  does not depend on  $\arg z$ , Sha Yao gave a positive answer to Question 1.2 in her Ph.D thesis by using the main result in [4].

In the present article, we give the following (negative) answer to Question 1.2.

**Theorem 1.3.** *There exist subharmonic functions  $\psi$  and  $\varphi$  on  $\mathbb{C}$  satisfying*

- (1)  $\int_{\mathbb{C}} e^{-\psi} < +\infty$ ;
- (2)  $\varphi(0) \in (-\infty, +\infty)$ ;
- (3) *for any holomorphic function  $f$  on  $\mathbb{C}$  satisfying  $f(0) = 1$ ,  $\int_{\mathbb{C}} |f|^2 e^{-\varphi - \psi} = +\infty$  holds.*

2010 *Mathematics Subject Classification.* 32D15, 32E10, 32L10, 32U05, 32W05.

*Key words and phrases.* plurisubharmonic functions, holomorphic functions,  $L^2$  extension.

The authors were partially supported by NSFC.

## 2. PROOF OF THEOREM 1.3

Let  $\psi = 2 \max\{c_1 \log |z - 1|, c_2 \log |z - 1|\}$  and  $\varphi = (1 - c_1) \log |z - 1|$ , where  $c_1 \in (\frac{1}{2}, 1)$  and  $c_2 \in (1, \frac{3}{2})$ .

We prove Theorem 1.3 by contradiction: if not, then there exists holomorphic function  $f$  on  $\mathbb{C}$  satisfying  $f(0) = 1$ , and

$$\int_{\mathbb{C}} |f|^2 e^{-\varphi - \psi} < +\infty. \quad (2.1)$$

Note that  $(\psi + \varphi)|_{\{|z| < 1\}} = 2 \log |z - 1|$ , then inequality (2.1) implies that  $f(1) = 0$ .

Note that  $\psi + \varphi - 2(1 - c_1 + c_2) \log |z|$  is bounded near  $\infty$ , then inequality (2.1) implies that  $f$  is a polynomial. Furthermore, it follows from  $1 - c_1 + c_2 < 2$  and inequality (2.1) that the degree of  $f$  must be 0, which contradicts  $f(1) = 0$ . This proves the present theorem.

*Acknowledgements.* The author would like to thank Professor Takeo Ohsawa for giving us series talks in Peking University and sharing his recent work.

## REFERENCES

- [1] B. Berndtsson and L. Lempert, A proof of the Ohsawa-Takegoshi theorem with sharp estimates. *J. Math. Soc. Japan* 68 (2016), no. 4, 1461–1472.
- [2] Z. Blocki, Suita conjecture and the Ohsawa-Takegoshi extension theorem, *Invent. Math.* 193 (2013) 149–158.
- [3] Q.A. Guan, X.Y. Zhou, Optimal constant problem in the  $L^2$  extension theorem. *C. R. Math. Acad. Sci. Paris* 350 (2012), no. 15–16, 753–756.
- [4] Q.A. Guan, X.Y. Zhou, A solution of an  $L^2$  extension problem with an optimal estimate and applications, *Ann. of Math. (2)* 181 (2015), no. 3, 1139–1208.
- [5] Q.A. Guan, X.Y. Zhou, Optimal constant in an  $L^2$  extension problem and a proof of a conjecture of Ohsawa, *Sci. China Math.* 58(1) (2015) 35–59.
- [6] Q.-A. Guan, X.-Y. Zhou and L. Zhu, On the Ohsawa-Takegoshi  $L^2$  extension theorem and the twisted Bochner-Kodaira identity, *C. R. Math. Acad. Sci. Paris* 349(13C14) (2011) 797–800.
- [7] T. Ohsawa, On the extension of  $L^2$  holomorphic functions VIII—a remark on a theorem of Guan and Zhou. *Internat. J. Math.* 28 (2017), no. 9, 1740005, 12 pp.
- [8] T. Ohsawa, K. Takegoshi, *On the extension of  $L^2$  holomorphic functions*, *Math. Z.* 195 (1987), 197–204.

QI'AN GUAN: SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING 100871, CHINA.

*E-mail address:* guanqian@math.pku.edu.cn