The Structure of the Realizations of the Causal Information Rate-Distortion Function for Markovian Sources:

Realizations with Densities

Milan S. Derpich

Department of Electronic Engineering

Universidad Técnica Federico Santa María.

Abstract

The main purpose of this note is to show that in a realization (x_1^n, y_1^n) of the causal information rate-distortion function (IRDF) for a κ -th order Markovian source x_1^n , under a single letter sum distortion constraint, the smallest integer ℓ for which $y_k \leftrightarrow y_1^{k-1}, x_{k-\ell+1}^k \leftrightarrow x_1^{k-\ell}$ holds is $\ell = \kappa$. This result is derived under the assumption that the sequences (x_1^n, y_1^n) have a joint probability density function.

I. INTRODUCTION

Consider the causal information rate-distortion function (IRDF) for a random source $\mathbf{x}_1^n = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$, defined as

$$R_{c,n}^{it}(D) \triangleq \frac{1}{n} \inf I(\mathbf{x}_1^n; \mathbf{y}_1^n), \tag{1}$$

where the minimization is over all conditional PDFs $f_{y_1^n \mid x_1^n}$ satisfying the distortion constraint

$$\frac{1}{n} E \left[\sum_{i=1}^{n} \rho(\mathbf{x}_i, \mathbf{y}_i) \right] \le D \tag{2}$$

and the causality Markov chains

$$y_1^i \longleftrightarrow x_1^i \longleftrightarrow y_{i+1}^n, \quad i = 1, \dots, n.$$
 (3)

If the infimum is achieved by some conditional distribution, the associated pair of sequences $\mathbf{x}_1^n, \mathbf{y}_1^n$ is called a realization of $R_{c,n}^{it}(D)$. Here we assume that such distribution exists and that the corresponding

realization has a joint PDF. This assumption is satisfied if, for example, \mathbf{x}_1^n is Gaussian and $\rho(x,y) = (x-y)^2$.

The first purpose of this note is to show that in a realization of the causal IRDF for a κ -th order Markovian source x_1^n , under the average distortion constraint (2), and supposing that in such realization the sequences have a joint PDF, it holds that

$$f_{\mathbf{y}_{k}\mid\mathbf{x}_{1}^{n},\mathbf{y}_{1}^{k-1}}(y_{k}|x_{1}^{n},y_{1}^{k-1}) = \frac{e^{-s\rho(x_{k},y_{k})}\,\breve{F}_{k}(x_{k-\kappa+1}^{k},y_{1}^{k})}{\int e^{-s\rho(x_{k},y_{k})}\,\breve{F}_{k}(x_{k-\kappa+1}^{k},y_{1}^{k})dy_{k}}$$
(4a)

where $f_{\mathbf{x}_1^n}$ is the PDF of \mathbf{x}_1^n and

$$\breve{F}_{k}(x_{k-\kappa+1}^{k}, y_{1}^{k}) = e^{\int \ln\left(\int e^{-s\rho(x_{k+1}, y_{k+1})} \breve{F}_{k+1}(x_{k-\kappa+2}^{k+1}, y_{1}^{k+1}) dy_{k+1}\right) f_{x_{k+1}^{n} | x_{k-\kappa+1}^{k}}(x_{k+1}^{n} | x_{k-\kappa+1}^{k}) dx_{k+1}^{n}}$$
(4b)

The expressions given in (4) are a special case of the ones given by [1, equations (16),(17),(18)] for abstract spaces, where their derivation is not included. The value of our first result resides in that

- We provide a proof for the validity of (4) (absent in [1]).
- In this proof, we pose the causal IRDF optimization problem with $f_{\mathbf{y}_1^n \mid \mathbf{x}_1^n}$ as the decision variable (instead of the collection $\{f_{\mathbf{y}_i \mid \mathbf{x}_1^i, \mathbf{y}_1^{i-1}}\}_{i=1}^n$ as would be the case in [1] for probability measures having an associated PDF). Accordingly, we impose an explicit causality constraint on $f_{\mathbf{y}_1^n \mid \mathbf{x}_1^n}$, instead of enforcing causality structurally by restricting $f_{\mathbf{y}_1^n \mid \mathbf{x}_1^n}$ to be the product of $\{f_{\mathbf{y}_i \mid \mathbf{x}_1^i, \mathbf{y}_1^{i-1}}\}_{i=1}^n$, as done in [1], [2].

The second (and main) goal of this document is to note that from (4a) it is clear that

$$y_k \longleftrightarrow y_1^{k-1}, x_{k-\kappa+1}^k \longleftrightarrow x_1^{k-\kappa}$$
 (5)

holds, and that

$$y_k \longleftrightarrow y_1^{k-1}, x_k \longleftrightarrow x_1^{k-1}$$
 (6)

does not hold, except for $\kappa = 1$. Crucially, (6) does not become true by supposing that the joint PDF of x_1^k, y_1^k is stationary, thus contradicting [2, Remark IV.5] and what is stated in the discussion paragraph at the end of [1, Section V].

II. PROOF

The causal IRDF under the above conditions is yielded by the solution to the following optimization problem:

minimize:
$$I(\mathbf{x}_1^n; \mathbf{y}_1^n)$$
 (7a)

subject to:
$$\left(\int f_{y_1^n \mid x_1^n}(y_1^n \mid x_1^n) dy_1^n - 1 \right) f_{x_1^n}(x_1^n) = 0, \quad \forall x_1^n$$
 (7b)

$$\iint f_{\mathbf{y}_1^n \mid \mathbf{x}_1^n}(y_1^n | x_1^n) f_{\mathbf{x}_1^n}(x_1^n) \sum_{k=1}^n \rho(x_k, y_k) dy_1^n dx_1^n \le D$$
 (7c)

$$(f_{\mathbf{v}_1^k \mid \mathbf{x}_1^k}(y_1^k \mid x_1^k) - f_{\mathbf{v}_1^k \mid \mathbf{x}_1^n}(y_1^k \mid x_1^n)) f_{\mathbf{x}_1^n}(x_1^n) = 0, \quad \forall y_1^k, x_1^n, \ k = 1, \dots, n.$$
 (7d)

where the minimization is over the conditional PDF $f_{y_1^n \mid x_1^n}$. Notice that (7d) is an explicit causality constraint equivalent to (3).

Let $f_{\mathbf{y}_1^n \mid \mathbf{x}_1^n}' : \mathbb{R}^{n \times n} \to [0,1]$ be any conditional PDF, and define

$$g_{y_1^n \mid x_1^n} \triangleq (f'_{y_1^n \mid x_1^n} - f_{y_1^n \mid x_1^n}) \tag{8}$$

$$g_{\mathbf{y}_{1}^{n}}(y_{1}^{n}) \triangleq \int g_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}(y_{1}^{n} \mid x_{1}^{n}) f_{\mathbf{x}_{1}^{n}}(x_{1}^{n}) dx_{1}^{n} \tag{9}$$

$$f_{y_1^n \mid x_1^n}^{\epsilon} \triangleq f_{y_1^n \mid x_1^n} + \epsilon g_{y_1^n \mid x_1^n} \tag{10}$$

$$f_{\mathbf{y}_{1}^{n}}^{\epsilon}(y_{1}^{n}) \triangleq \int f_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}^{\epsilon}(y_{1}^{n} \mid x_{1}^{n}) f_{\mathbf{x}_{1}^{n}}(x_{1}^{n}) dx_{1}^{n} \tag{11}$$

where $\epsilon \in [0, 1]$.

Before writing the Lagrangian and taking its Gateaux differential, let us obtain the Gateaux differential of $I(\mathbf{x}_1^n; \mathbf{y}_1^n)$ in the direction $g_{\mathbf{y}_1^n \mid \mathbf{x}_1^n}$, given by

$$\frac{dI(\mathbf{x}_{1}^{n}; \mathbf{y}_{1}^{n})}{d\epsilon}\Big|_{\epsilon=0} = \frac{d}{d\epsilon} \left[\iint f_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}^{\epsilon}(y_{1}^{n} \mid x_{1}^{n}) f_{\mathbf{x}_{1}^{n}}(x_{1}^{n}) \ln \left(\frac{f_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}^{\epsilon}(y_{1}^{n} \mid x_{1}^{n})}{f_{\mathbf{y}_{1}^{n}}^{\epsilon}(y_{1}^{n})} \right) dy_{1}^{n} dx_{1}^{n} \right]\Big|_{\epsilon=0}$$
(12)

$$= \iint g_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}(y_{1}^{n} \mid x_{1}^{n}) f_{\mathbf{x}_{1}^{n}}(x_{1}^{n}) \ln \left(\frac{f_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}(y_{1}^{n} \mid x_{1}^{n})}{f_{\mathbf{y}_{1}^{n}}(y_{1}^{n})} \right) dy_{1}^{n} dx_{1}^{n} + R$$
(13)

where

$$R \triangleq \iint f_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}(y_{1}^{n} \mid x_{1}^{n}) f_{\mathbf{x}_{1}^{n}}(x_{1}^{n}) \left(\frac{g_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}(y_{1}^{n} \mid x_{1}^{n})}{f_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}(y_{1}^{n} \mid x_{1}^{n})} - \frac{g_{\mathbf{y}_{1}^{n}}(y_{1}^{n})}{f_{\mathbf{y}_{1}^{n}}(y_{1}^{n})} \right) dy_{1}^{n} dx_{1}^{n}$$

$$(14)$$

$$= \iint g_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}(y_{1}^{n} \mid x_{1}^{n}) f_{\mathbf{x}_{1}^{n}}(x_{1}^{n}) dy_{1}^{n} dx_{1}^{n} - \iint \frac{f_{\mathbf{y}_{1}^{n}, \mathbf{x}_{1}^{n}}(y_{1}^{n}, x_{1}^{n}) g_{\mathbf{y}_{1}^{n}}(y_{1}^{n})}{f_{\mathbf{y}_{1}^{n}}(y_{1}^{n})} dy_{1}^{n} dx_{1}^{n}$$

$$(15)$$

$$= \int g_{\mathbf{y}_{1}^{n}}(y_{1}^{n})dy_{1}^{n} - \int \frac{g_{\mathbf{y}_{1}^{n}}(y_{1}^{n})}{f_{\mathbf{y}_{1}^{n}}(y_{1}^{n})} \left(\int f_{\mathbf{y}_{1}^{n},\mathbf{x}_{1}^{n}}(y_{1}^{n},x_{1}^{n})dx_{1}^{n} \right) dy_{1}^{n}$$

$$(16)$$

$$=0 (17)$$

On the other hand, for each $i=1,\ldots,n$, the causality constraint (7d) appears in the Lagrangian as

$$\iint \lambda_i(x_1^n, y_1^i) \left[f_{y_1^i \mid x_1^i}(y_1^i \mid x_1^i) - f_{y_1^i \mid x_1^n}(y_1^i \mid x_1^n) \right] f_{x_1^n}(x_1^n) dy_1^i dx_1^n \tag{18}$$

$$= \iint \lambda_i(x_1^n, y_1^i) \left(\int \left[f_{\mathbf{y}_1^n \mid \mathbf{x}_1^i}(y_1^n | x_1^i) - f_{\mathbf{y}_1^n \mid \mathbf{x}_1^n}(y_1^n | x_1^n) \right] dy_{i+1}^n \right) f_{\mathbf{x}_1^n}(x_1^n) dy_1^i dx_1^n$$
(19)

$$= \int \left(\int \lambda_i(x_1^n, y_1^i) f_{\mathbf{y}_1^n \mid \mathbf{x}_1^i}(y_1^n \mid x_1^i) f_{\mathbf{x}_1^n}(x_1^n) dx_1^n - \int \lambda_i(x_1^n, y_1^i) f_{\mathbf{y}_1^n \mid \mathbf{x}_1^n}(y_1^n \mid x_1^n) f_{\mathbf{x}_1^n}(x_1^n) dx_1^n \right) dy_1^n \quad (20)$$

It will be convenient to manipulate this expression so as to give it a structure similar to the other terms in the Lagrangian. For this purpose, notice that

$$\int \lambda_i(x_1^n, y_1^i) f_{y_1^n \mid x_1^i}(y_1^n \mid x_1^i) f_{x_1^n}(x_1^n) dx_1^n \tag{21}$$

$$= \int \lambda_i(x_1^n, y_1^i) f_{\mathbf{y}_1^n, \mathbf{x}_1^i}(y_1^n, x_1^i) f_{\mathbf{x}_{i+1}^n \mid \mathbf{x}_1^i}(x_{i+1}^n \mid x_1^i) dx_1^n$$
(22)

$$= \int f_{\mathbf{y}_{1}^{n}, \mathbf{x}_{1}^{i}}(y_{1}^{n}, x_{1}^{i}) \left(\int \lambda_{i}(x_{1}^{n}, y_{1}^{i}) f_{\mathbf{x}_{i+1}^{n} \mid \mathbf{x}_{1}^{i}}(x_{i+1}^{n} \mid x_{1}^{i}) dx_{i+1}^{n} \right) dx_{1}^{i}$$
(23)

$$= \int f_{\mathbf{y}_1^n, \mathbf{x}_1^i}(y_1^n, x_1^i) \bar{\lambda}(x_1^i, y_1^i) dx_1^i$$
 (24)

$$= \int \left(\int f_{\mathbf{y}_{1}^{n}, \mathbf{x}_{1}^{n}}(y_{1}^{n}, x_{1}^{n}) dx_{i+1}^{n} \right) \bar{\lambda}(x_{1}^{i}, y_{1}^{i}) dx_{1}^{i}$$
(25)

$$= \int f_{\mathbf{y}_1^n \mid \mathbf{x}_1^n}(y_1^n \mid x_1^n) f_{\mathbf{x}_1^n}(x_1^n) \bar{\lambda}(x_1^i, y_1^i) dx_1^n$$
(26)

where

$$\bar{\lambda}_i(x_1^i, y_1^i) \triangleq \int \lambda_i(x_1^n, y_1^i) f_{\mathbf{x}_{i+1}^n \mid \mathbf{x}_1^i}(x_{i+1}^n \mid x_1^i) dx_{i+1}^n, \quad i = 1, \dots, n.$$
 (27)

Substituting this into (20) we obtain

$$\int \lambda_i(x_1^n, y_1^i) (f_{y_1^i \mid x_1^i}(y_1^i \mid x_1^i) - f_{y_1^i \mid x_1^n}(y_1^i \mid x_1^n)) f_{x_1^n}(x_1^n) dy_1^i dx_1^n$$
(28)

$$= \int \left(\bar{\lambda}_i(x_1^i, y_1^i) - \lambda_i(x_1^n, y_1^i)\right) f_{y_1^n \mid x_1^n}(y_1^n \mid x_1^n) f_{x_1^n}(x_1^n) dy_1^n dx_1^n \tag{29}$$

We can now write the Lagrangian associated with optimization problem (7) as

$$\mathcal{L}(f_{\mathbf{y}_{1}^{n}\mid\mathbf{x}_{1}^{n}}) \triangleq I(\mathbf{x}_{1}^{n};\mathbf{y}_{1}^{n}) + \int \eta(x_{1}^{n}) \left(\int f_{\mathbf{y}_{1}^{n}\mid\mathbf{x}_{1}^{n}}(y_{1}^{n}|x_{1}^{n})dy_{1}^{n} - 1 \right) f_{\mathbf{x}_{1}^{n}}(x_{1}^{n})dx_{1}^{n}$$
(30)

$$+ s \left(\int f_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}(y_{1}^{n} \mid x_{1}^{n}) f_{\mathbf{x}_{1}^{n}}(x_{1}^{n}) \left(\sum_{i=1}^{n} \rho(x_{i}, y_{i}) \right) dx_{1}^{n} dy_{1}^{n} - D \right)$$
(31)

$$+\sum_{i=1}^{n} \int \left(\bar{\lambda}_{i}(x_{1}^{i}, y_{1}^{i}) - \lambda_{i}(x_{1}^{n}, y_{1}^{i})\right) f_{y_{1}^{n} \mid x_{1}^{n}}(y_{1}^{n} \mid x_{1}^{n}) f_{x_{1}^{n}}(x_{1}^{n}) dy_{1}^{n} dx_{1}^{n}$$
(32)

From the theory of Lagrangian optimization on vector spaces [3], $f_{y_1^n \mid x_1^n}$ is a solution to Optimization Problem (7) only if

$$0 = \frac{d}{d\epsilon} \mathcal{L}(f_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}^{\epsilon}) \Big|_{\epsilon=0}$$

$$= \sum_{y_{1}^{n}, x_{1}^{n}} \left[\ln \left(\frac{f_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}(y_{1}^{n} \mid x_{1}^{n})}{f_{\mathbf{y}_{1}^{n}}(y_{1}^{n})} \right) + \eta(x_{1}^{n}) + \sum_{i=1}^{n} \left(s\rho(x_{i}, y_{i}) + \bar{\lambda}_{i}(x_{1}^{i}, y_{1}^{i}) - \lambda_{i}(x_{1}^{n}, y_{1}^{i}) \right) \right]$$

$$\times g_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}(y_{1}^{n} \mid x_{1}^{n}) f_{\mathbf{x}_{1}^{n}}(x_{1}^{n})$$

$$(33)$$

$$\times g_{\mathbf{y}_{1}^{n} \mid \mathbf{x}_{1}^{n}}(y_{1}^{n} \mid x_{1}^{n}) f_{\mathbf{x}_{1}^{n}}(x_{1}^{n})$$

for every function $g_{y_1^n \mid x_1^n}$ as defined in (8), i.e., for every conditional PDF $f'_{y_1^n \mid x_1^n}$. This holds if and only if for every x_1^n, y_1^n :

$$\ln\left(\frac{f_{\mathbf{y}_{1}^{n}|\mathbf{x}_{1}^{n}}(y_{1}^{n}|x_{1}^{n})}{f_{\mathbf{y}_{1}^{n}}(y_{1}^{n})}\right) = -\eta(x_{1}^{n}) - \sum_{i=1}^{n} \left(s\rho(x_{i}, y_{i}) + \bar{\lambda}_{i}(x_{1}^{i}, y_{1}^{i}) - \lambda_{i}(x_{1}^{n}, y_{1}^{i})\right)$$
(35)

$$\iff f_{y_1^n \mid x_1^n}(y_1^n \mid x_1^n) = e^{-\eta(x_1^n) - \sum_{i=1}^n \left(s\rho(x_i, y_i) + \bar{\lambda}_i(x_1^i, y_1^i) - \lambda_i(x_1^n, y_1^i) \right)} f_{y_1^n}(y_1^n)$$
(36)

The Lagrange multiplier function $\eta(x_1^n)$ must enforce the constraint (7b). Hence,

$$f_{\mathbf{y}_{1}^{n}\mid\mathbf{x}_{1}^{n}}(y_{1}^{n}|x_{1}^{n}) = \frac{e^{-\sum_{i=1}^{n} \left(s\rho(x_{i},y_{i}) + \bar{\lambda}_{i}(x_{1}^{i},y_{1}^{i}) - \lambda_{i}(x_{1}^{n},y_{1}^{i})\right)} f_{\mathbf{y}_{1}^{n}}(y_{1}^{n})}{K_{1}(x_{1}^{n})},$$
(37)

where

$$K_1(x_1^n) \triangleq \int e^{-\sum_{i=1}^n \left(s\rho(x_i, y_i) + \bar{\lambda}_i(x_1^i, y_1^i) - \lambda_i(x_1^n, y_1^i)\right)} f_{y_1^n}(y_1^n) dy_1^n$$
(38)

Marginalizing over y_{k+1}^n we obtain

$$f_{\mathbf{y}_{1}^{k}\mid\mathbf{x}_{1}^{n}}(y_{1}^{k}|x_{1}^{n}) = \frac{e^{-\sum_{i=1}^{k}(s\rho(x_{i},y_{i})+\bar{\lambda}_{i}(x_{1}^{i},y_{1}^{i})-\lambda_{i}(x_{1}^{n},y_{1}^{i}))} \int e^{-\sum_{i=k+1}^{n}\left(s\rho(x_{i},y_{i})+\bar{\lambda}_{i}(x_{1}^{i},y_{1}^{i})-\lambda_{i}(x_{1}^{n},y_{1}^{i})\right)} f_{\mathbf{y}_{1}^{n}}(y_{1}^{n})dy_{k+1}^{n}}{K_{1}(x_{1}^{n})}$$
(39)

Using Bayes' rule we can write

$$f_{\mathbf{y}_{k}\mid\mathbf{x}_{1}^{n},y_{1}^{k-1}}(y_{k}|x_{1}^{n},y_{1}^{k-1}) = \frac{f_{\mathbf{y}_{1}^{k}\mid\mathbf{x}_{1}^{n}}(y_{1}^{k}|x_{1}^{n})}{f_{\mathbf{y}_{1}^{k-1}\mid\mathbf{x}_{1}^{n}}(y_{1}^{k-1}|x_{1}^{n})}$$
(40)

$$= \frac{e^{-s\rho(x_k, y_k)} F_k(x_1^n, y_1^k)}{\int e^{-s\rho(x_k, y_k)} F_k(x_1^n, y_1^k) dy_k}$$
(41)

where

$$F_k(x_1^n, y_1^k) \triangleq e^{-(\bar{\lambda}_k(x_1^k, y_1^k) - \lambda_k(x_1^n, y_1^k))} \int e^{-\sum_{i=k+1}^n \left(s\rho(x_i, y_i) + \bar{\lambda}_i(x_1^i, y_1^i) - \lambda_i(x_1^n, y_1^i)\right)} f_{y_1^n}(y_1^n) dy_{k+1}^n$$
(42)

These functions can be written recursively as

$$F_n(y_1^n) = f_{y_1^n}(y_1^n) (43a)$$

$$F_k(x_1^n, y_1^k) = e^{-(\bar{\lambda}_k(x_1^k, y_1^k) - \lambda_k(x_1^n, y_1^k))} \int e^{-s\rho(x_{k+1}, y_{k+1})} F_{k+1}(x_1^n, y_1^{k+1}) dy_{k+1}$$
(43b)

In order attain causality in (41), the functions $F_k(x_1^n,y_1^k)$ must depend only on x_1^k and y_1^k . Since for each k, the function F_{k+1} does not depend on terms $(\bar{\lambda}_i(x_1^i,y_1^i)-\lambda_i(x_1^n,y_1^i))$ with $i\leq k$, the causality constraint is met if and only if we choose $(\bar{\lambda}_i(x_1^k,y_1^k)-\lambda_i(x_1^n,y_1^k))$ in (43b) such that, for each $k=1,\ldots,n$

$$F_k(x_1^n, y_1^k) = e^{-(\bar{\lambda}_i(x_1^k, y_1^k) - \lambda_i(x_1^n, y_1^k))} \int e^{-s\rho(x_{k+1}, y_{k+1})} F_{k+1}(x_1^n, y_1^{k+1}) dy_{k+1} = \breve{F}_k(x_1^k, y_1^k)$$
(44)

for some function \check{F}_k .

For k=n, the causality constraint is satisfied automatically since $F_n(x_1^n,y_1^n)=\breve{F}_n(y_1^k)\triangleq f_{y_1^n}(y_1^n)$ (see (43a)). Suppose now that (44) (i.e., causality) is satisfied for k+1, for some k>n. In such case, one can replace $F_{k+1}(x_1^n,y_1^{k+1})$ in (44) by $\breve{F}_{k+1}(x_1^{k+1},y_1^{k+1})$ and, defining

$$K_{k+1}(x_1^{k+1}, y_1^k) \triangleq \int e^{-s\rho(x_{k+1}, y_{k+1})} \breve{F}_{k+1}(x_1^{k+1}, y_1^{k+1}) dy_{k+1},$$

write (44) as

$$\bar{\lambda}_k(x_1^k, y_1^k) - \lambda_k(x_1^n, y_1^k) = \ln K_{k+1}(x_1^n, y_1^k) - \ln \check{F}_k(x_1^k, y_1^k)$$
(45)

Multiplying both sides by $f_{\mathbf{x}_{k+1}^n \mid \mathbf{x}_1^k}(x_{k+1}^n | x_1^k)$ and integrating over x_{k+1}^n we obtain

$$0 = \int \left(\bar{\lambda}_k(x_1^k, y_1^k) - \lambda_k(x_1^n, y_1^k)\right) f_{\mathbf{x}_{k+1}^n \mid \mathbf{x}_1^k}(x_{k+1}^n \mid x_1^k) dx_{k+1}^n \tag{46}$$

$$= \int \left(\ln K_{k+1}(x_1^n, y_1^k) - \ln \check{F}_k(x_1^k, y_1^k) \right) f_{\mathbf{x}_{k+1}^n \mid \mathbf{x}_1^k}(x_{k+1}^n \mid x_1^k) dx_{k+1}^n \tag{47}$$

$$\iff \ln \breve{F}_k(x_1^k, y_1^k) = \int \ln K_{k+1}(x_1^n, y_1^k) f_{\mathbf{x}_{k+1}^n \mid \mathbf{x}_1^k}(x_{k+1}^n \mid x_1^k) dx_{k+1}^n \tag{48}$$

This yields that the recursion (43) takes the form

$$\check{F}_n(x_1^n, y_1^n) = f_{y_1^n}(y_1^n)$$
(49)

$$\breve{F}_{k}(x_{1}^{k}, y_{1}^{k}) = e^{\int \ln\left(\sum_{y_{k+1}} e^{-s\rho(x_{k+1}, y_{k+1})} \breve{F}_{k+1}(x_{1}^{k+1}, y_{1}^{k+1})\right) f_{x_{k+1}^{n} \mid x_{1}^{k}}(x_{k+1}^{n} \mid x_{1}^{k}) dx_{k+1}^{n}}$$
(50)

If \mathbf{x}_1^n is κ -th order Markovian, then $f_{\mathbf{x}_k^n \mid \mathbf{x}_1^{k-1}}(x_k^n \mid x_1^{k-1}) = f_{\mathbf{x}_k^n \mid \mathbf{x}_{k-\kappa}^{k-1}}(x_k^n \mid x_{k-\kappa}^{k-1})$, for all $k = 1, \ldots, n$, in which case (50) becomes (4b). Substituting the latter into (44) and then in (41) yields (4a).

Finally, from (4a), it follows that in a realization of the causal IRDF it must hold that

$$y_k \longleftrightarrow y_1^{k-1}, x_{k-\kappa+1}^k \longleftrightarrow x_1^{k-\kappa}$$
 (51)

and that

$$y_k \longleftrightarrow y_1^{k-1}, x_k \longleftrightarrow x_1^{k-1}$$
 (52)

does not hold, except for k = n. This completes the proof.

¹ This reflects the fact that there is no need to enforce the causality constraint for k = n, since there are no source samples for time k > n.

REFERENCES

- [1] P. Stavrou, C. D. Charalambous, and C. K. Kourtellaris, "Optimal nonstationary reproduction distribution for nonanticipative RDF on abstract alphabets," *CoRR*, vol. abs/1301.6522, 2013. [Online]. Available: http://arxiv.org/abs/1301.6522
- [2] C. Charalambous, P. Stavrou, and N. Ahmed, "Nonanticipative rate distortion function and relations to filtering theory," *IEEE Transactions on Automatic Control*, vol. 59, no. 4, pp. 937–952, Apr. 2014.
- [3] D. Luenberger, Optimization by Vector Space Methods. London: John Wiley and Sons, Inc., 1969.