

ON GENERALIZED ERDŐS-GINZBURG-ZIV CONSTANTS OF C_n^r

DONGCHUN HAN AND HANBIN ZHANG

ABSTRACT. Let G be an additive finite abelian group with exponent $\exp(G) = m$. For any positive integer k , the k -th generalized Erdős-Ginzburg-Ziv constant $\mathfrak{s}_{km}(G)$ is defined as the smallest positive integer t such that every sequence S in G of length at least t has a zero-sum subsequence of length km . It is easy to see that $\mathfrak{s}_{kn}(C_n^r) \geq (k+r)n - r$ where $n, r \in \mathbb{N}$. Kubertin conjectured that the equality holds for any $k \geq r$. In this paper, we mainly prove the following results:

- (1) For every positive integer $k \geq 6$, we have

$$\mathfrak{s}_{kn}(C_n^3) = (k+3)n + O\left(\frac{n}{\ln n}\right).$$

- (2) For every positive integer $k \geq 18$, we have

$$\mathfrak{s}_{kn}(C_n^4) = (k+4)n + O\left(\frac{n}{\ln n}\right).$$

- (3) For $n \in \mathbb{N}$, assume that the largest prime power divisor of n is p^a for some $a \in \mathbb{N}$. For any fixed $r \geq 5$, if $p^t \geq r$ for some $t \in \mathbb{N}$, then for any $k \in \mathbb{N}$ we have

$$\mathfrak{s}_{kp^t n}(C_n^r) \leq (kp^t + r)n + c_r \frac{n}{\ln n},$$

where c_r is a constant depends on r .

Note that the main terms in our results are consistent with the conjectural values proposed by Kubertin.

1. INTRODUCTION

Let G be an additive finite abelian group with exponent $\exp(G) = m$. Let $S = g_1 \cdot \dots \cdot g_k$ be a sequence over G (repetition is allowed), where $g_i \in G$ for $1 \leq i \leq k$, k is called the length of the sequence S . We call S a zero-sum sequence if $\sum_{i=1}^k g_i = 0$. The classical direct zero-sum problem studies conditions (mainly refer to lengths) which ensure that given sequences have non-empty zero-sum subsequences with prescribed properties (also mainly refer to lengths). For example, the Davenport constant, denoted by $D(G)$, is the smallest positive integer t such that every sequence S over G of length at least t has a nonempty zero-sum subsequence. It is easy to prove that $D(C_n) = n$, where C_n is the cyclic group of order n . For any positive integer k , the k -th generalized Erdős-Ginzburg-Ziv constant $\mathfrak{s}_{km}(G)$ is defined as the smallest positive integer t such that every sequence S over G of length at least t has a zero-sum subsequence of length km . In particular, for $k = 1$, $\mathfrak{s}_m(G)$ is called the Erdős-Ginzburg-Ziv constant, which is a classical invariant in combinatorial number theory. In 1961, Erdős, Ginzburg and Ziv [6] proved that $\mathfrak{s}_n(C_n) = 2n - 1$ which is usually regarded as a starting point of zero-sum theory (see [2] for other different proofs of this result). We refer to [10] for a survey of zero-sum problems. In this paper, we will focus on $\mathfrak{s}_{km}(G)$.

Let $G = C_n^r = \langle e_1 \rangle \oplus \dots \oplus \langle e_r \rangle$. Assume that T consists of $n - 1$ copies of e_i for $1 \leq i \leq r$. Let S consist of $kn - 1$ copies of 0 and T , then it is easy to show

that S is a sequence over C_n^r of length $(k+r)n - r - 1$ and S contains no zero-sum subsequences of length kn . Consequently we have

$$(1.1) \quad \mathfrak{s}_{kn}(C_n^r) \geq (k+r)n - r.$$

For general finite abelian group G with $\exp(G) = m$, similar construction can be used to show that $\mathfrak{s}_{km}(G) \geq km + D(G) - 1$ holds for $k \geq 1$. In 1996, Gao [9] proved that $\mathfrak{s}_{km}(G) = km + D(G) - 1$, provided that $km \geq |G|$. In [13], Gao and Thangadurai proved that if $km < D(G)$, then $\mathfrak{s}_{km}(G) > km + D(G) - 1$. Define $l(G)$ as the smallest integer t such that $\mathfrak{s}_{km}(G) = km + D(G) - 1$ holds for every $k \geq t$. From the above we know that

$$\frac{D(G)}{m} \leq l(G) \leq \frac{|G|}{m}.$$

Recently, Gao, Han, Peng and Sun conjectured ([11], Conjecture 4.7) that

$$l(G) = \lceil \frac{D(G)}{m} \rceil.$$

Clearly we have $l(C_n) = 1$ by the Erdős-Ginzburg-Ziv theorem. For finite abelian groups G of rank two, $l(G) = 2$ (see [11]). Let p be a prime and q a power of p , the above conjecture was verified for C_q^r where $1 \leq r \leq 4$ (also more generally for abelian p -group G with $D(G) \leq 4m$) except for some cases when p is rather small, see [13, 19, 22]. For the studies of $l(G)$ for the general cases, we refer to [11, 20, 22].

Recall (1.1) that $\mathfrak{s}_{kn}(C_n^r) \geq (k+r)n - r$, in [22], Kubertin conjectured that the equality actually holds for any $k \geq r$.

Conjecture 1.1. For any positive integers k, n with $k \geq r$, we have

$$\mathfrak{s}_{kn}(C_n^r) = (k+r)n - r.$$

According to the results in [13, 19, 22], Conjecture 1.1 has been verified for $r \leq 4$ except for some cases when p is rather small ($p \leq 3$). Recently, Sidorenko [23, 24] verified Conjecture 1.1 for C_2^r . He [23] also applied his results to prove new bounds for the codegree Turán density of complete r -graphs. Moreover, he [24] established connections between $\mathfrak{s}_{2k}(C_2^r)$ and linear binary codes. Actually, he showed that the problem of determining $\mathfrak{s}_{2k}(C_2^r)$ is essentially equivalent to finding the lowest redundancy of a linear binary code of given length which does not contain words of Hamming weight $2k$.

Towards Conjecture 1.1, Kubertin [22] proved that

$$\mathfrak{s}_{kq}(C_q^r) \leq (k + \frac{3}{8}r^2 + \frac{3}{2}r - \frac{3}{8})q - r,$$

where $p > \min\{2k, 2r\}$ is a prime and q is a power of p . By extending the method of Kubertin, He [20] improved the above upper bound and obtained that

$$\mathfrak{s}_{kq}(C_q^r) \leq (k + 5r - 2)q - 3r$$

when $2p \geq 7r - 3$ and $k \geq r$. He also proved that $\mathfrak{s}_{kn}(C_n^r) \leq 6kn$ for n with large prime factors and k sufficiently large. More precisely, he showed that for $r, l > 0$, $n = p_1^{\alpha_1} \cdots p_l^{\alpha_l}$ with distinct prime factors $p_1, \dots, p_l \geq \frac{7}{2}r - 3$ and $k = a_1 \cdots a_l$ a product of positive integers $a_1, \dots, a_l \geq r$, $\mathfrak{s}_{kn}(C_n^r) \leq 6kn$. We also refer to [3, 14] for some recent results on the lower bound of $\mathfrak{s}_{kn}(C_n^r)$ when k is much smaller than the rank r , note that in this case $\mathfrak{s}_{kn}(C_n^r) > (k+r)n - r$ (see [13]).

For $n \in \mathbb{N}$, let

$$M(n) = \max\{p^k \mid \text{with } p^k \mid n \text{ where } p \text{ is a prime and } k \in \mathbb{N}\},$$

i.e., the largest prime power divisor of n . For convenience, let $M(1) = 1$. For any $n, r \in \mathbb{N}$, we define

$$p(n, r) = \min\{p^t \mid M(n) = p^a \text{ and } p^t \geq r\}.$$

In this paper, we focus on the Conjecture 1.1 and prove the following results.

Theorem 1.2. *Let $k \in \mathbb{N}$. We have*

(1) *For every $k \geq 6$,*

$$s_{kn}(C_n^3) = (k+3)n + O\left(\frac{n}{\ln n}\right);$$

(2) *For every $k \geq 18$,*

$$s_{kn}(C_n^4) = (k+4)n + O\left(\frac{n}{\ln n}\right);$$

(3) *For every $k \in \mathbb{N}$ and fixed $r \geq 5$,*

$$s_{kp(n,r)n}(C_n^r) = (kp(n,r) + r)n + O_r\left(\frac{n}{\ln n}\right),$$

where O_r depends on r .

Note that the main terms in Theorem 1.2 are consistent with the conjectural values in Conjecture 1.1. Moreover, the error term can be improved in some cases. By some further studies of $M(n)$, roughly speaking, for any real number $A \geq 1$, we can improve the order of the error term from $\frac{n}{\ln n}$ to $\frac{n}{(\ln n)^A}$ for almost every $n \geq 1$. Furthermore, when the number of distinct prime divisors of n is a given integer m , we can even improve the order of the error term to $n^{1-\frac{1}{m}}$.

The following sections are organized as follows. In Section 2, we shall introduce some notations and preliminary results. In Section 3, we will prove our main results. In Section 4, we will provide further studies on $M(n)$ and then apply these results to improve our main results.

2. PRELIMINARIES

This section will provide more rigorous definitions and notations. We also introduce some preliminary results that will be used repeatedly below.

Let \mathbb{N} denote the set of positive integers, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ and \mathbb{R} the field of real numbers. Let f and g be real valued functions, both defined on \mathbb{N} , such that $g(x)$ is strictly positive for all large enough values of x . Then we denote $f(x) = O(g(x))$ if and only if there exists a positive real number M and a positive integer x_0 such that

$$|f(x)| \leq M|g(x)| \quad \text{for all } x \geq x_0.$$

We also use the notation O_r (resp. $O_{A,\epsilon}$) which means that the above M depends on r (resp. A and ϵ), where $r \in \mathbb{N}_0$, $A, \epsilon \in \mathbb{R}$. Similarly, we denote $f(x) = o(g(x))$ if and only if for every positive constant ϵ , there exists a positive integer x_0 such that

$$|f(x)| \leq \epsilon g(x) \quad \text{for all } x \geq x_0.$$

Let G be an additive finite abelian group. By the fundamental theorem of finite abelian groups we have

$$G \cong C_{n_1} \oplus \cdots \oplus C_{n_r}$$

where $r = r(G) \in \mathbb{N}_0$ is the rank of G , $n_1 | \cdots | n_r \in \mathbb{N}$ are positive integers. Moreover, n_1, \dots, n_r are uniquely determined by G , and $n_r = \exp(G)$ is called the *exponent* of G .

We define a *sequence* over G to be an element of the free abelian monoid $(\mathcal{F}(G), \cdot)$, see Chapter 5 of [17] for detailed explanation. Our notations of sequences follow the notations in the paper [15]. In particular, in order to avoid confusion between exponentiation of the group operation in G and exponentiation of the sequence operation \cdot in $\mathcal{F}(G)$, we define:

$$g^{[k]} = \underbrace{g \cdots g}_k \in \mathcal{F}(G) \quad \text{and} \quad T^{[k]} = \underbrace{T \cdots T}_k \in \mathcal{F}(G),$$

for $g \in G$, $T \in \mathcal{F}(G)$ and $k \in \mathbb{N}_0$.

We write a sequence S in the form

$$S = \prod_{g \in G} g^{\nu_g(S)} \quad \text{with } \nu_g(S) \in \mathbb{N}_0 \text{ for all } g \in G.$$

We call

- $\nu_g(S)$ the *multiplicity* of g in S ,
- $|S| = l = \sum_{g \in G} \nu_g(S) \in \mathbb{N}_0$ the *length* of S ,
- $T = \prod_{g \in G} g^{\nu_g(T)}$ a *subsequence* of S if $\nu_g(T) \leq \nu_g(S)$ for all $g \in G$, and denote by $T|S$,
- $\sigma(S) = \sum_{i=1}^l g_i = \sum_{g \in G} \nu_g(S)g \in G$ the *sum* of S ,
- S a *zero-sum sequence* if $\sigma(S) = 0$,
- S a *zero-sum free sequence* if $\sigma(T) \neq 0$ for every $T|S$,
- S a *short zero-sum sequence* if it is a zero-sum sequence of length $|S| \in [1, \exp(G)]$.

Using these concepts, we can define

- $D(G)$ as the smallest integer $l \in \mathbb{N}$ such that every sequence S over G of length $|S| \geq l$ has a non-empty zero-sum subsequence. We call $D(G)$ the *Davenport constant* of G .
- $\mathfrak{s}_{k \exp(G)}(G)$ as the smallest integer $l \in \mathbb{N}$ such that every sequence S over G of length $|S| \geq l$ has a non-empty zero-sum subsequence T of length $|T| = k \exp(G)$, where $k \in \mathbb{N}$. We call $\mathfrak{s}(G) := \mathfrak{s}_{\exp(G)}(G)$ the *Erdős-Ginzburg-Ziv constant* and $\mathfrak{s}_{k \exp(G)}(G)$ the k -th *generalized Erdős-Ginzburg-Ziv constant*.

Lemma 2.1. ([17], Theorem 5.5.9) *Let G be a finite abelian p -group and $G = C_{p^{n_1}} \oplus \cdots \oplus C_{p^{n_r}}$, then*

$$D(G) = \sum_{i=1}^r (p^{n_i} - 1) + 1.$$

Lemma 2.2. *Let G be a finite abelian group with $\exp(G) = m$, then*

$$\mathfrak{s}_{km}(G) \geq km + D(G) - 1$$

holds for every $k \geq 1$.

Proof. By the definition of $D(G)$, there exists a zero-sum free sequence T of length $|T| = D(G) - 1$. Let $S = T \cdot 0^{[km-1]}$. It is easy to know that S is a sequence over G

of length $|S| = km + D(G) - 2$ and S contains no zero-sum subsequence of length km . This completes the proof. \square

Lemma 2.3. ([8], Theorem 3.2) *Let G be a finite abelian p -group and $\exp(G) = p^{nr}$. If $p^{m+nr} \geq D(G)$ for some $m \in \mathbb{N}$, then*

$$s_{kp^m p^{nr}}(G) = k \cdot p^{m+nr} + D(G) - 1,$$

holds for any $k \in \mathbb{N}$.

The following classical result of Alon and Dubiner is crucial in our proof.

Lemma 2.4 ([1], Theorem 1.1). *There exists an absolute constant $c > 0$ such that*

$$s(C_n^r) \leq (cr \log_2 r)^r n.$$

Although the precise values of $s(G)$ for general C_n^r are not known, some cases (when n is a power of a small prime) have been determined. We list some of these results which are very useful in our proof.

Lemma 2.5. *Let $n \in \mathbb{N}$.*

- (1) $s(C_{2^n}^3) = 8 \cdot 2^n - 7$;
- (2) $s(C_{3^n}^3) = 9 \cdot 3^n - 8$;
- (3) $s(C_{2^n}^4) = 16 \cdot 2^n - 15$;
- (4) $s(C_{3^n}^4) = 20 \cdot 3^n - 19$.

Proof. (1) See [4], Corollary 4.4. (2) See [12], Theorem 1.7. (3) See [4], Corollary 4.4. (4) See [4], Theorem 1.3, 1.4 and Section 5. \square

In the rest of this section, we provide some results about $M(n)$ which are useful in this paper. Recall that, for any $n \in \mathbb{N}$, let

$$M(n) = \max\{p^k \mid \text{with } p^k \mid n \text{ where } p \text{ is a prime and } k \in \mathbb{N}\}$$

be the largest prime power divisor of n . For convenience, let $M(1) = 1$. For example, we have $M(20) = M(2^2 5) = 5$, $M(40) = M(2^3 5) = 2^3$ and $M(200) = M(2^3 5^2) = 5^2$. Unlike the widely studied largest prime divisor function

$$P(n) = \max\{p \mid \text{with } p \mid n \text{ and } p \text{ is a prime}\},$$

as far as we know, $M(n)$ has not received much attention. As $M(p) = p$ where p is a prime, certainly we have $\limsup_{n \rightarrow \infty} \frac{M(n)}{n} = 1$. It is known and easy to prove that

$$(2.1) \quad \liminf_{n \rightarrow \infty} \frac{M(n)}{\ln n} = 1,$$

consequently

$$(2.2) \quad \lim_{n \rightarrow \infty} M(n) = \infty.$$

Recently, Girard [18] used (2.2) to show that $D(C_n^r) = rn + o(n)$, which is an important result in zero-sum theory and also can be regarded as an example of application of $M(n)$. In this paper, we will continue to employ the estimates of $M(n)$ to the zero-sum problems. Although the proof of (2.1) is simple and elementary, it is hard to find this result in literatures or standard textbooks. So we decide to provide a proof here for the convenience of the reader.

Let

$$\pi(x) = \#\{p \mid p \leq x\}$$

be the prime-counting function and

$$\vartheta(x) = \sum_{p \leq x} \ln p$$

the Chebyshev ϑ function. The result in the following lemma is very classical and can be easily found in [25].

Lemma 2.6. *For any $x \geq 2$, we have*

$$\pi(x) \leq 2 \frac{x}{\ln x}.$$

Proof. This result is an easy consequence of Theorem 3, Page 11 in [25]. \square

Lemma 2.7. *For any $n \in \mathbb{N}$, we have*

$$M(n) \geq \frac{1}{2} \ln n.$$

Proof. When $n = p^m$ is a prime power, the result is obvious. If n is not a prime power, we may assume that

$$n = q_1^{r_1} \cdots q_k^{r_k} p^m,$$

where $q_1 < \cdots < q_k$ and p are distinct prime numbers, $r_1, \dots, r_k, m \in \mathbb{N}$ with $M(n) = p^m$. By the definition of $M(n)$, clearly we have $n \leq p^m p^{km}$. Moreover we have $k < \pi(p^m)$. For otherwise if $k \geq \pi(p^m)$, then we have

$$k = \pi(p_k) \geq \pi(p^m) \geq \pi(p)$$

and consequently $p_k \geq p$. As $q_1 < \cdots < q_k$ and p are distinct prime numbers, we have $q_k > p_k$. Therefore,

$$\pi(q_k) > \pi(p_k) = k \geq \pi(p^m)$$

and consequently $q_k > p^m$. By the definition of $M(n)$, we have $M(n) \geq q_k^{r_k} > p^m$, but this contradicts $M(n) = p^m$. Therefore $n \leq p^{m\pi(p^m)}$, and by Lemma 2.6 we have

$$\frac{M(n)}{\ln n} = \frac{p^m}{\ln n} \geq \frac{p^m}{\ln p^{m\pi(p^m)}} = \frac{p^m}{\pi(p^m) \ln p^m} \geq \frac{1}{2}.$$

This completes the proof. \square

For sufficiently large number n , $P(n)$ may be rather small, for example $P(2^m) = 2$ for any $m \in \mathbb{N}$. However, Lemma 2.7 means that $M(n)$ cannot be too small for sufficiently large n . We shall use Lemma 2.7 to prove our main results in the next section. In the following, we will prove (2.1) which shows that actually $\ln n$ is the minimal order of $M(n)$.

Let p_k denote the k -th prime and $n_k = p_1 \cdots p_k \in \mathbb{N}$. Clearly, we have $M(n_k) = p_k$ and $\ln n_k = \vartheta(p_k)$. By the Prime Number Theorem, for any $\epsilon > 0$ there exists $k_0(\epsilon) > 0$ such that for all $k > k_0(\epsilon)$ we have

$$\frac{M(n_k)}{\ln n_k} = \frac{p_k}{\vartheta(p_k)} \leq (1 + \epsilon).$$

Therefore we have $\liminf_{n \rightarrow \infty} \frac{M(n)}{\ln n} \leq 1$.

Similarly, from Lemma 2.7, together with the Prime Number Theorem, for any $\epsilon > 0$ there exists $n_0(\epsilon) > 0$ such that for all $n > n_0(\epsilon)$ we have

$$\frac{M(n)}{\ln n} \geq \frac{p^m}{\pi(p^m) \ln p^m} \geq (1 - \epsilon).$$

Therefore we have $\liminf_{n \rightarrow \infty} \frac{M(n)}{\ln n} \geq 1$ and

$$\liminf_{n \rightarrow \infty} \frac{M(n)}{\ln n} = 1.$$

This completes the proof of (2.1).

3. PROOF OF THE MAIN RESULTS

In this section, we shall prove our main results, Theorem 1.2. Firstly, we have to verify Conjecture 1.1 for some small primes which are the remaining cases in [13, 19, 22].

Lemma 3.1. *For any $n \in \mathbb{N}$, we have*

- (1) $\mathfrak{s}_{k2^n}(C_{2^n}^3) = (k+3)2^n - 3$, holds for $k \geq 4$;
- (2) $\mathfrak{s}_{k3^n}(C_{3^n}^3) = (k+3)3^n - 3$, holds for $k \geq 6$;
- (3) $\mathfrak{s}_{k2^n}(C_{2^n}^4) = (k+4)2^n - 4$, holds for $k \geq 12$;
- (4) $\mathfrak{s}_{k3^n}(C_{3^n}^4) = (k+4)3^n - 4$, holds for $k \geq 18$.

Proof. By Lemma 2.2, it suffices to prove $\mathfrak{s}_{k \exp(G)} \leq k \exp(G) + D(G) - 1$.

(1) We prove by induction on k . Since $2^{2+n} = 4 \cdot 2^n \geq D(C_{2^n}^3) = 3 \cdot 2^n - 2$, by Lemma 2.3 we have $\mathfrak{s}_{4 \cdot 2^n}(C_{2^n}^3) = 7 \cdot 2^n - 3$. This proves the case $k = 4$. Suppose that $k \geq 5$ and the result holds for all positive integers n with $4 \leq n \leq k$. Now we need to prove $\mathfrak{s}_{(k+1)2^n}(C_{2^n}^3) = (k+1+3)2^n - 3$.

Let S be any sequence over $C_{2^n}^3$ of length

$$|S| = (k+1+3)2^n - 3.$$

By Lemma 2.5.(1) and the fact that $|S| \geq 8 \cdot 2^n - 7$, we have S contains a zero-sum subsequence T of length $|T| = 2^n$. Since

$$|S \cdot T^{-1}| = (k+3)2^n - 3 = \mathfrak{s}_{k2^n}(C_{2^n}^3),$$

we have $S \cdot T^{-1}$ contains a zero-sum subsequence U of length $|U| = k2^n$. Consequently, $T \cdot U$ is a zero-sum subsequence of S of length $|T \cdot U| = (k+1)2^n$. This completes the proof.

(2) We prove by induction on k . Since $3^{1+n} = 3 \cdot 3^n \geq D(C_{3^n}^3) = 3 \cdot 3^n - 2$, by Lemma 2.3 we have $\mathfrak{s}_{6 \cdot 3^n}(C_{3^n}^3) = 9 \cdot 3^n - 3$. This proves the case $k = 6$. Suppose that $k \geq 7$ and the result holds for all positive integers n with $6 \leq n \leq k$. Now we need to prove $\mathfrak{s}_{(k+1)3^n}(C_{3^n}^3) = (k+1+3)3^n - 3$.

Let S be any sequence over $C_{3^n}^3$ of length

$$|S| = (k+1+3)3^n - 3.$$

By Lemma 2.5.(2) and the fact that $|S| \geq 9 \cdot 3^n - 8$, we have S contains a zero-sum subsequence T of length $|T| = 3^n$. Since

$$|S \cdot T^{-1}| = (k+3)3^n - 3 = \mathfrak{s}_{k3^n}(C_{3^n}^3),$$

we have $S \cdot T^{-1}$ contains a zero-sum subsequence U of length $|U| = k3^n$. Consequently, $T \cdot U$ is a zero-sum subsequence of S of length $|T \cdot U| = (k+1)3^n$. This completes the proof.

(3) We prove by induction on k . Since $2^{2+n} = 4 \cdot 2^n \geq D(C_{2^n}^4) = 4 \cdot 2^n - 3$, by Lemma 2.3 we have $\mathfrak{s}_{12 \cdot 2^n}(C_{2^n}^4) = 16 \cdot 2^n - 4$. This proves the case $k = 12$. Suppose that $k \geq 13$ and the result holds for all positive integers n with $12 \leq n \leq k$. Now we need to prove $\mathfrak{s}_{(k+1)2^n}(C_{2^n}^4) = (k+1+4)2^n - 4$.

Let S be any sequence over $C_{2^n}^4$ of length

$$|S| = (k + 1 + 4)2^n - 4.$$

By Lemma 2.5.(3) and the fact that $|S| \geq 16 \cdot 2^n - 15$, we have S contains a zero-sum subsequence T of length $|T| = 2^n$. Since

$$|S \cdot T^{-1}| = (k + 4)2^n - 4 = \mathfrak{s}_{k2^n}(C_{2^n}^4),$$

we have $S \cdot T^{-1}$ contains a zero-sum subsequence U of length $|U| = k2^n$. Consequently, $T \cdot U$ is a zero-sum subsequence of S of length $|T \cdot U| = (k + 1)2^n$. This completes the proof.

(4) We prove by induction on k . Since $3^{2+n} = 9 \cdot 3^n \geq D(C_{3^n}^4) = 4 \cdot 3^n - 4$, by Lemma 2.3 we have $\mathfrak{s}_{18 \cdot 3^n}(C_{3^n}^4) = 22 \cdot 3^n - 4$. This proves the case $k = 18$. Suppose that $k \geq 19$ and the result holds for all positive integers n with $18 \leq n \leq k$. Now we need to prove $\mathfrak{s}_{(k+1)3^n}(C_{3^n}^4) = (k + 1 + 4)3^n - 4$.

Let S be any sequence over $C_{3^n}^4$ of length

$$|S| = (k + 1 + 4)3^n - 4.$$

By Lemma 2.5.(4) and the fact that $|S| \geq 20 \cdot 3^n - 19$, we have S contains a zero-sum subsequence T of length $|T| = 3^n$. Since

$$|S \cdot T^{-1}| = (k + 4)3^n - 4 = \mathfrak{s}_{k3^n}(C_{3^n}^4),$$

we have $S \cdot T^{-1}$ contains a zero-sum subsequence U of length $|U| = k3^n$. Consequently, $T \cdot U$ is a zero-sum subsequence of S of length $|T \cdot U| = (k + 1)3^n$. This completes the proof. \square

Corollary 3.2. Let $n, m \in \mathbb{N}$ and p be any prime. we have

- (1) $\mathfrak{s}_{kp^m}(C_{p^m}^3) = (k + 3)p^m - 3$ holds for $k \geq 6$;
- (2) $\mathfrak{s}_{kp^m}(C_{p^m}^4) = (k + 4)p^m - 4$ holds for $k \geq 18$;

Proof. By Lemma 3.1, the results hold for the cases $p = 2, 3$. For $p \geq 5$, see Theorem 1.(3) in [22] and Theorem 1.2.(3) in [19]. \square

The following crucial lemma is based on a standard argument in zero-sum theory (we refer to [17], Proposition 5.7.11).

Lemma 3.3. Let $n, m, p, k, r \in \mathbb{N}$ and p a prime. Assume that $\mathfrak{s}_{kp^m}(C_{p^m}^r) = (k + r)p^m - r$. Then we have

$$\mathfrak{s}_{knp^m}(C_{np^m}^r) \leq (k + r)np^m + a_r n,$$

where a_r is a constant depends on r .

Proof. Let S be a sequence of length $|S| = ((k + r)p^m - r)n + \mathfrak{s}(C_n^r)$ over C_n^r . Consider the following map:

$$\varphi : C_{np^m}^r \rightarrow C_n^r.$$

Then $\varphi(S)$ is a sequence over C_n^r of length $((k + r)p^m - r)n + \mathfrak{s}(C_n^r)$. By the definition of $\mathfrak{s}(C_n^r)$, we have $\varphi(S)$ contains at least $(k + r)p^m - r$ zero-sum subsequences $S_1, \dots, S_{(k+r)p^m-r}$ over C_n^r with $|S_i| = n$ for $1 \leq i \leq (k + r)p^m - r$. This means that

$$\sigma(S_1), \dots, \sigma(S_{(k+r)p^m-r}) \in \ker(\varphi) = C_{p^m}^r.$$

By the assumption that $\mathfrak{s}_{kp^m}(C_{p^m}^r) = (k + r)p^m - r$, there exist

$$\{i_1, \dots, i_{kp^m}\} \subset \{1, \dots, (k + r)p^m - r\}$$

such that $\sigma(S_{i_1}) + \dots + \sigma(S_{i_{k_p^m}}) = 0$ and this implies that $S_{i_1} \cdot \dots \cdot S_{i_{k_p^m}}$ is a zero-sum subsequence of S over $C_{np^m}^r$ of length knp^m . Therefore

$$s_{knp^m}(C_{np^m}^r) \leq ((k+r)p^m - r)n + s(C_n^r).$$

Moreover, by Lemma 2.4, there exists an absolute constant c such that

$$((k+r)p^m - r)n + s(C_n^r) \leq ((k+r)p^m - r)n + (cr \log_2 r)^r n.$$

Let $a_r = (cr \log_2 r)^r - r$, then we have the desired result. \square

For any fixed $r \in \mathbb{N}$, we denote $a_r = (cr \log_2 r)^r - r$, where c is the absolute constant mentioned in Lemma 2.4. The following corollary is an easy consequence of the above lemma.

Corollary 3.4. Let $n, k, r \in \mathbb{N}$. Assume that $M(n) = p^m$ and

$$s_{kp^m}(C_{p^m}^r) = (k+r)p^m - r.$$

Then we have

$$s_{kn}(C_n^r) \leq (k+r)n + a_r \frac{n}{M(n)}.$$

By Corollary 3.4, in order to prove the main results, it suffices to combine the results about $M(n)$ in Section 2.

Proof of the Theorem 1.2. (1) By Corollary 3.2.(1) and 3.4, for $k \geq 6$, we have

$$s_{kn}(C_n^3) \leq (k+3)n + a_3 \frac{n}{M(n)}.$$

By lemma 2.7, for $k \geq 6$, actually we have

$$s_{kn}(C_n^3) \leq (k+3)n + 2a_3 \frac{n}{\ln n}$$

and we get the desired result.

(2) By Corollary 3.2.(2) and 3.4, for $k \geq 18$, we have

$$s_{kn}(C_n^4) \leq (k+4)n + a_4 \frac{n}{M(n)}.$$

By lemma 2.7, for $k \geq 18$, actually we have

$$s_{kn}(C_n^4) \leq (k+4)n + 2a_4 \frac{n}{\ln n}$$

and we get the desired result.

(3) By Lemma 2.3 and Corollary 3.4, for any $k \in \mathbb{N}$, we have

$$s_{kp(n,r)n}(C_n^r) \leq (kp(n,r) + r)n + a_r \frac{n}{M(n)}.$$

By lemma 2.7, for any $k \in \mathbb{N}$, actually we have

$$s_{kp(n,r)n}(C_n^r) \leq (kp(n,r) + r)n + 2a_r \frac{n}{\ln n}$$

and we get the desired result. \square

4. FURTHER STUDIES ABOUT $M(n)$ AND SOME IMPROVEMENTS

In this section, we will provide some further estimates for $M(n)$ in some special cases. With these further estimates, we can improve our main results. All these results can be seen as some applications of $M(n)$.

Note that, by (2.1), it is impossible to improve the order of the lower bound in Lemma 2.7 of $M(n)$ for every n any more. However, we can get some better estimates in some special cases.

We denote

$$E(x, y) = \{n \leq x \mid M(n) \leq y\}$$

and $\overline{E}(x, y) = \{n \leq x \mid n \notin E(x, y)\}$. Let $A > 1$ be any real number, in the following we shall consider

$$E(x, (\ln x)^A) = \{n \leq x \mid M(n) \leq (\ln x)^A\},$$

where $A \geq 1$. Actually, we have the following lemma.

Lemma 4.1. *For any positive integer $A \geq 1$ and $\epsilon > 0$, we have*

$$|E(x, (\ln x)^A)| = O_{A, \epsilon}(x^{1 - \frac{1}{A} + \epsilon}).$$

Proof. Clearly we have $|E(x, (\ln x)^A)| = \sum_{\substack{n \leq x \\ M(n) \leq (\ln x)^A}} 1$, then for any $\delta > 0$, we have

$$\sum_{\substack{n \leq x \\ M(n) \leq (\ln x)^A}} 1 \leq \sum_{\substack{n \leq x \\ M(n) \leq (\ln x)^A}} \left(\frac{x}{n}\right)^\delta.$$

Similar to the Euler product of the Riemann zeta function, by the fundamental theorem of arithmetic, we have

$$\begin{aligned} \sum_{\substack{n \leq x \\ M(n) \leq (\ln x)^A}} \left(\frac{x}{n}\right)^\delta &\leq x^\delta \prod_{p \leq (\ln x)^A} \left(1 - \frac{1}{p^\delta}\right)^{-1} \\ &= x^\delta \prod_{p \leq (\ln x)^A} \left(1 + \frac{1}{p^\delta - 1}\right). \end{aligned}$$

If we take $c_\delta = \frac{2^\delta}{2^\delta - 1}$, then $\frac{1}{p^\delta - 1} \leq \frac{c_\delta}{p^\delta}$ and we have

$$x^\delta \prod_{p \leq (\ln x)^A} \left(1 + \frac{1}{p^\delta - 1}\right) \leq x^\delta \prod_{p \leq (\ln x)^A} \left(1 + \frac{c_\delta}{p^\delta}\right).$$

As $1 + x \leq e^x$ for any $x \geq 0$, we have

$$\begin{aligned} x^\delta \prod_{p \leq (\ln x)^A} \left(1 + \frac{c_\delta}{p^\delta}\right) &\leq x^\delta \prod_{p \leq (\ln x)^A} \exp\left(\frac{c_\delta}{p^\delta}\right) \\ &= x^\delta \exp\left(\sum_{p \leq (\ln x)^A} \frac{c_\delta}{p^\delta}\right). \end{aligned}$$

To estimate the last sum, we employ the relation between the sum and the integral,

$$\begin{aligned} x^\delta \exp\left(\sum_{p \leq (\ln x)^A} \frac{c_\delta}{p^\delta}\right) &\leq x^\delta \exp\left(\sum_{2 \leq n \leq (\ln x)^A} \frac{c_\delta}{n^\delta}\right) \\ &\leq x^\delta \exp\left(c_\delta \int_1^{(\ln x)^A} \frac{1}{t^\delta} dt\right). \end{aligned}$$

Therefore

$$\begin{aligned} x^\delta \exp(c_\delta \int_1^{(\ln x)^A} \frac{1}{t^\delta} dt) &= x^\delta \exp\left(\frac{c_\delta}{1-\delta}((\ln x)^{A(1-\delta)} - 1)\right) \\ &= \exp\left(\frac{c_\delta}{\delta-1}\right) x^\delta \exp\left(\frac{c_\delta}{1-\delta}((\ln x)^{A(1-\delta)})\right). \end{aligned}$$

Now, we take $\delta = 1 - \frac{1}{A} + \frac{\epsilon}{2}$. Since

$$A(1-\delta) = 1 - \frac{A\epsilon}{2} < 1$$

and

$$\exp\left(\frac{c_\delta}{1-\delta}((\ln x)^{A(1-\delta)})\right) = O_{A,\epsilon}(x^{\frac{\epsilon}{2}}),$$

we have

$$\exp\left(\frac{c_\delta}{\delta-1}\right) x^\delta \exp\left(\frac{c_\delta}{1-\delta}((\ln x)^{A(1-\delta)})\right) = O_{A,\epsilon}(x^{1-\frac{1}{A}+\epsilon}).$$

This completes the proof. \square

Recall that for any fixed $r \in \mathbb{N}$, we denote $a_r = (cr \log_2 r)^r - r$, where c is the absolute constant mentioned in Lemma 2.4. Let

$$\mathbb{S}_k^r(x, A) = \begin{cases} \{n \leq x \mid \mathfrak{s}_{kn}(C_n^r) \leq (k+r)n + a_r \frac{n}{(\ln n)^A}\}, & \text{if } r = 3 \text{ or } 4, \\ \{n \leq x \mid \mathfrak{s}_{kp(n,r)n}(C_n^r) \leq (kp(n,r) + r)n + a_r \frac{n}{(\ln n)^A}\}, & \text{if } r \geq 5 \end{cases}$$

and

$$\overline{\mathbb{S}_k^r(x, A)} = \{n \leq x \mid n \notin \mathbb{S}_k^r(x, A)\}.$$

Theorem 4.2. *For any $A \geq 1$ and $\epsilon > 0$, we have*

- (1) $|\overline{\mathbb{S}_k^3(x, A)}| = O_{A,\epsilon}(x^{1-\frac{1}{A}+\epsilon})$, holds for $k \geq 6$;
- (2) $|\overline{\mathbb{S}_k^4(x, A)}| = O_{A,\epsilon}(x^{1-\frac{1}{A}+\epsilon})$, holds for $k \geq 18$;
- (3) $|\overline{\mathbb{S}_k^r(x, A)}| = O_{A,\epsilon}(x^{1-\frac{1}{A}+\epsilon})$, holds for $r \geq 5$ and any $k \in \mathbb{N}$.

In particular,

- (1) For $k \geq 6$, we have

$$\lim_{x \rightarrow \infty} \frac{|\overline{\mathbb{S}_k^3(x, A)}|}{x} = 0;$$

- (2) For $k \geq 18$, we have

$$\lim_{x \rightarrow \infty} \frac{|\overline{\mathbb{S}_k^4(x, A)}|}{x} = 0;$$

- (3) For $r \geq 5$ and any $k \in \mathbb{N}$, we have

$$\lim_{x \rightarrow \infty} \frac{|\overline{\mathbb{S}_k^r(x, A)}|}{x} = 0;$$

Proof. By the definition of $\overline{\mathbb{E}(x, (\ln x)^A)}$ and Corollary 3.4, it is easy to see that

$$\overline{\mathbb{E}(x, (\ln x)^A)} \subset \overline{\mathbb{S}_k^3(x, A)}.$$

Therefore, we have

$$\overline{\mathbb{S}_k^3(x, A)} \subset \overline{\mathbb{E}(x, (\ln x)^A)}.$$

The desired result follows from Lemma 4.1. In particular, let $\epsilon = \frac{1}{2A}$ in the above result, we have

$$\lim_{x \rightarrow \infty} \frac{|\overline{\mathbb{S}_k^3(x, A)}|}{x} \leq \lim_{x \rightarrow \infty} \frac{|\mathbb{E}(x, (\ln x)^A)|}{x} = \lim_{x \rightarrow \infty} \frac{O(x^{1-\frac{1}{2A}})}{x} = 0.$$

This completes the proof of (1). The proofs of (2) and (3) are similar. \square

Remark 4.3. According to Theorem 4.2, for any $A \geq 1$, roughly speaking, for almost every $n \geq 1$ we have

(1) For $k \geq 6$,

$$s_{kn}(C_n^3) \leq (k+3)n + a_3 \frac{n}{(\ln n)^A};$$

(2) For $k \geq 18$,

$$s_{kn}(C_n^4) \leq (k+4)n + a_4 \frac{n}{(\ln n)^A};$$

(3) For any $k \in \mathbb{N}$ and fixed $r \geq 5$,

$$s_{kp(n,r)n}(C_n^r) \leq (kp(n,r) + r)n + a_r \frac{n}{(\ln n)^A}.$$

Let $\omega(n)$ denote the number of distinct prime divisors of n . In the following, we can improve the error terms for some $n \in \mathbb{N}$ when $\omega(n)$ is a given integer m .

Lemma 4.4. *For any $n \in \mathbb{N}$, we have*

$$M(n) \geq n^{\frac{1}{\omega(n)}}.$$

Proof. For any $n \in \mathbb{N}$, we assume that $n = q_1^{r_1} \cdots q_m^{r_m}$, where $q_1^{r_1} < \cdots < q_m^{r_m}$ and q_1, \dots, q_m are distinct prime numbers, $r_1, \dots, r_m \in \mathbb{N}$. Then by the definition of $M(n)$, we have $M(n) = q_m^{r_m}$.

Clearly we have $\omega(n) = m$ and

$$n = q_1^{r_1} \cdots q_m^{r_m} \leq q_m^{mr_m} = M(n)^{\omega(n)},$$

consequently $M(n) \geq n^{\frac{1}{\omega(n)}}$. \square

Theorem 4.5. *Let $n, r, m \in \mathbb{N}$ with $\omega(n) = m$, we have*

(1) For $k \geq 6$,

$$s_{kn}(C_n^3) \leq (k+3)n + a_3 n^{1-\frac{1}{m}};$$

(2) For $k \geq 18$,

$$s_{kn}(C_n^4) \leq (k+4)n + a_4 n^{1-\frac{1}{m}};$$

(3) For any $k \in \mathbb{N}$ and fixed $r \geq 5$,

$$s_{kp(n,r)n}(C_n^r) \leq (kp(n,r) + r)n + a_r n^{1-\frac{1}{m}}.$$

Proof. (1) By Corollary 3.2 and 3.4, Lemma 4.4 and $\omega(n) = m$, for $k \geq 6$ we have

$$\begin{aligned} s_{kn}(C_n^3) &\leq (k+3)n + a_3 \frac{n}{M(n)} = (k+3)n + a_3 \frac{n}{n^{\frac{1}{\omega(n)}}} \\ &= (k+3)n + a_3 \frac{n}{n^{\frac{1}{m}}} = (k+3)n + a_3 n^{1-\frac{1}{m}}. \end{aligned}$$

This completes the proof. The proofs of (2) and (3) are similar. \square

Remark 4.6. Compared with the previous error terms $\frac{n}{\ln n}$ and $\frac{n}{(\ln n)^A}$, the error term $n^{1-\frac{1}{m}}$ is a large improvement and it is valid for every $n \in \mathbb{N}$ with $\omega(n) = m$.

Acknowledgments. D.C. Han was supported by the National Science Foundation of China Grant No.11601448 and the Fundamental Research Funds for the Central Universities Grant No.2682016CX121. H.B. Zhang was supported by the National Science Foundation of China Grant No.11671218 and China Postdoctoral Science Foundation Grant No. 2017M620936. The authors would like to thank Prof. Weidong Gao for many useful comments and corrections.

REFERENCES

1. N. Alon and M. Dubiner, *A lattice point problem and additive number theory*, *Combinatorica* **15**(1995) 301-309.
2. N. Alon and M. Dubiner, *Zero-sum sets of prescribed size*, in: *Combinatorics, Paul Erdős is Eighty*, Bolyai Society, Mathematical studies, Keszthely, Hungary, 1993, 33-50.
3. J. Bitz, C. Griffith and X. He, *Exponential Lower Bounds on the Generalized Erdős-Ginzburg-Ziv Constant*, arXiv:1712.00861.
4. Y. Edel, C. Elsholtz, A. Geroldinger, S. Kubertin and L. Rackham, *Zero-sum problems in finite abelian groups and affine caps*, *Q. J. Math.* **58** (2007) 159-186.
5. C. Elsholtz, *Lower bounds for multidimensional zero sums*, *Combinatorica* **24**(2004) 351-358.
6. P. Erdős, A. Ginzburg and A. Ziv, *Theorem in the additive number theory*, *Bull. Res. Council Israel* **10**(1961) 41-43.
7. Y. Fan, W. Gao and Q. Zhong, *On the Erdős-Ginzburg-Ziv constant of finite abelian groups of high rank*, *J. Number Theory* **131**(2011) 1864-1874.
8. W. Gao, *On zero-sum subsequences of restricted size. II*. *Discrete Math.* **271**(2003) 51-59.
9. W. Gao, *A combinatorial problem on finite abelian groups*, *J. Number Theory*, **58**(1996) 100-103.
10. W. Gao and A. Geroldinger, *Zero-sum problems in finite abelian groups: A survey*, *Expo. Math.* **24**(2006) 337-369.
11. W. Gao, D. Han, J. Peng and F. Sun, *On zero-sum subsequences of length $k \exp(G)$* , *J. Combin. Theory Ser. A* **125**(2014), 240-253.
12. W. Gao, Q. Hou, W. Schmid, and R. Thangadurai, *On short zero-sum subsequences II*, *Integers: Electronic Journal of Combinatorial Number Theory*, **7**(2007), paper A21, 22 pp.
13. W. Gao and R. Thangadurai, *On zero-sum sequences of prescribed length*, *Aequationes Math.* **72**(2006) 201-212.
14. J. Geneson, *Improved lower bound on generalized Erdős-Ginzburg-Ziv constants*, arXiv:1712.02069.
15. A. Geroldinger and D.J. Grynkiewicz, *The large Davenport constant I: groups with a cyclic, index 2 subgroup*, *J. Pure. Appl. Algebra* **217** (2013), 863-885.
16. A. Geroldinger, D. Grynkiewicz and W. Schmid, *Zero-sum problems with congruence conditions*, *Acta Math. Hungarica.* **131**(2011) 323-345.
17. A. Geroldinger and F. Halter-Koch, *Non-unique factorizations. Algebraic, Combinatorial and Analytic Theory*, *Pure Appl. Math.*, vol. 278, Chapman & Hall/CRC, 2006.
18. B. Girard, *An asymptotically tight bound for the Davenport constant*, *Journal de l'Ecole polytechnique - Mathématiques* **5**(2018) 605-611.
19. D. Han and H. Zhang, *On zero-sum subsequences of prescribed length*, *Int. J. Number Theory* **14**(2018) 167-191.
20. X. He, *Zero-sum subsequences of length kq over finite abelian p -groups*, *Discrete Math.* **339**(2016), no.1, 399-407.
21. H. Harborth, *Ein Extremalproblem für Gitterpunkte*, *J. Reine Angew. Math.* **262**(1973) 356-360.
22. S. Kubertin, *Zero-sums of length kq in \mathbb{Z}_q^d* , *Acta Arith.* **116**(2005) 145-152.
23. A. Sidorenko, *Extremal problems on the hypercube and the codegree Turán density of complete r -graphs*, arXiv:1710.08228.
24. A. Sidorenko, *On generalized Erdős-Ginzburg-Ziv constants for \mathbb{Z}_2^d* , arXiv:1808.06555.
25. G. Tenenbaum, *Introduction to analytic and probabilistic number theory*, *Cambridge Studies in Advanced Mathematics*, 46, Cambridge University Press, 1995.

DEPARTMENT OF MATHEMATICS, SOUTHWEST JIAOTONG UNIVERSITY, CHENGDU 610000, P.R.
CHINA

E-mail address: han-qingfeng@163.com

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING
100190, P.R. CHINA

E-mail address: zhanghanbin@amss.ac.cn