

On TQFT representations of mapping class groups with boundary

Greg Kuperberg; Shuang Ming

Abstract

We study the TQFT mapping class group representations for surfaces with boundary associated with the $SU(2)$ gauge group, or equivalently the quantum group $U_q(\mathfrak{sl}(2))$. We show that at a prime root of unity, these representations are all irreducible. We also examine braid group representations for transcendental values of the quantum parameter, where we show that the image of every mapping class group is Zariski dense.

1 Introduction

Given a integer $p \geq 3$, A a primitive $4p$ -th root of unity, we can associate a vector space to a surface with its boundary colored by nonnegative integers less than $p - 1$. The mapping class group of the surface act on the vector space projectively. These representations are called $SU(2)$ -TQFT representations and have many good properties [4] and it is used to study mapping class groups and provide interesting examples in both geometric topology [11] and quantum algebra [9].

In this paper, we are studying the image of mapping class groups in their $SU(2)$ -TQFT representations. Usually we understand this question in the following steps: We start with irreducibility, so we can restrict the image into blocks. Then we check the (in)finiteness of the image. If the image is finite then we have a finite quotient (e.g. Weil representation). Otherwise, we study its closure under Zariski topology or Analytic topology.

The irreducibility problem is first studied by Roberts [16]. Roberts proved when p is an odd prime, the representations are irreducible for closed surfaces. After that, Korinman [12] studied non-prime case: He showed that the representations are irreducible when p is the product of two distinct odd primes and when p is square of an odd prime, and for some other p , decomposable examples have been given. In [7], they first studied the cases that are with boundary. They proved when p is an odd prime, the representations of any surfaces with one boundary component is irreducible. Recently, Koberda and Santharoubane [10] considered the cases that are with boundaries, and they proved that for any p , if one of the boundaries is colored by 1, the representation can be shown irreducible.

The (in)finiteness problem is first studied by Funar [6]. Funar showed that when $p \neq 2, 3, 4, 6, 10$, the image of B_3 in $PSL(V_{0,3;1,1,1,3})$ is infinite. It implies for all representations containing this vector space as a tensor-factor of a subspace, the images of the mapping class groups will be infinite. After that, in [13], Korinman gives a criterion for (in)finiteness, which can be checked by hand for surfaces with low genus and small number of boundary components. Especially, Korinman gave the computation for the case p is odd prime, $g = 1$ and $b = 1$.

By the results of Funar and Korinman, in most cases, the image of the representation is infinite. The closure problem is studied by Freedman, Larsen, Wang [5][15] and the first author [14]. In [5] and [14], The Jones representation of braided group is proved dense in $PSU(N)$ for p not equal to 2, 3, 4, 6, 10. In [14], the first author also proved when A is generic, the image of the braid group is Zariski dense in $PSL(N)$. After that, In [15], Larsen and Wang showed when $p > 3$ is an odd prime. The $SO(3)$ -TQFT representations have dense image when genus $g \geq 2$.

It worth noting that in our setting, Jones representations of B_n can be seen as a special case of $SU(2)$ -TQFT representations of $MCG(\Sigma_{0,n+1})$. By taking the surface to be a sphere with $(n + 1)$ circle boundaries, and n of its boundary components are colored by 1.

In this paper, we study the case that surfaces are with nonempty boundary. Let $V_{g,b;\vec{c}}$ be the representation associated to the surface of genus g and b boundary components colored by \vec{c} (the precise definition will be given in the next section). We proved the following irreducibility theorem.

Theorem 1.1. $V_{g,b;\vec{c}}$ is irreducible at odd prime level under the action of $MCG(\Sigma_{g,b})$ for all admissible color \vec{c} .

When A is not a root of unity, the TQFT is not well defined because the dimension of $V_{g,b;\vec{c}}$ will be infinite when $g \geq 1$. However, when $g = 0$, this representation is well-defined because the color set is bounded by the half of the

total colors on the boundary. We have the following denseness theorem which generalize an earlier result [14] of the first author.

Theorem 1.2. *Let A be a transcendental complex number. The image of \widetilde{PB}_n is Zariski dense in the algebraic group $SL(V_{0,n;\vec{c}})$.*

The structure of this paper is the following. In section 2, we discuss the background for $SU(2)$ -TQFT representations. In section 3 to 6, we prove the irreducibility theorem. In section 7, we develop a set of tools for studying group homomorphisms. In section 8, we prove the Zariski denseness theorem.

2 Background

In this section, we give a short introduction to the background of $SU(2)$ -TQFT representations. One can find full explanation in [2].

2.1 The space $V_{p,g,b;\vec{c}}$

Given an integer $p \geq 3$, let $A \in \mathbb{C}$ be a primitive $4p$ -th root of unity. Using the Kauffman skein relation(see picture below), we can associate a link $L \in S^3$ an complex number $\langle L \rangle_p$.

$$\left(\begin{array}{c} \diagup \\ \diagdown \end{array} \right) = -A \left(\begin{array}{c} \diagdown \\ \diagup \end{array} \right) \quad ; \quad \left(\begin{array}{c} \frown \\ \smile \end{array} \right) = -A^{-1} \left(\begin{array}{c} \smile \\ \frown \end{array} \right) \quad ; \quad \bigcirc = -A^2 - A^{-2}.$$

Choose a handlebody of genus g and let C_g be the vector space spanned by the isotopy classes of framed links together with empty links. We fix two of such genus g handlebody H_g^1, H_g^2 and a glueing:

$$H_g^1 \bigcup_{\partial H_g^1 \rightarrow \partial H_g^2} H_g^2 \cong S^3.$$

This gives us a bilinear form

$$(\cdot, \cdot)_{g,p}^H : C_g \times C_g \rightarrow \mathbb{C}$$

by composing with the evaluation $\langle \cdot \rangle_p$ in S^3 . We define the space $V_{p,g}$ as C_g mod out by the kernel of $\langle \cdot, \cdot \rangle$. It was proved in [2] that the spaces $V_{p,g}$ are finite-dimensional, and a set of basis can be given as following.

We choose a trivalent graph Γ such that the handlebody H_g retracts to Γ , and color the edges from the set $\{0, 1, \dots, p-2\}$. We say a coloring is *admissible* if at each trivalent vertex, the color of the adjacent edges (i, j, k) satisfies:

- i) $|i - j| \leq k \leq i + j$,
- ii) $i + j + k \leq 2p - 4$,
- iii) $i + j + k = 0 \pmod{2}$

A representative of such a colored graph in C_g is given by Jones-Wenzl's idempotent. At each edges colored by c , we put c parallel string with an idempotent f_c . At vertices colored by (i, j, k) , we connect the projectors with the link below.

$$\begin{array}{c} \begin{array}{c} | \\ \diagup \quad \diagdown \\ j \quad k \end{array} \quad i \\ = \quad \begin{array}{c} | \\ \diagup \quad \diagdown \\ j \quad k \end{array} \quad i \\ \begin{array}{c} \boxed{f_i} \\ \diagup \quad \diagdown \\ \boxed{f_j} \quad \boxed{f_k} \\ \diagdown \quad \diagup \\ \end{array} \end{array}$$

This set of basis can be generalize to the surfaces with non-empty boundary. Let $V_{p,g,b;\vec{c}}$ be the vector space generated by the coloring of a fixed connected uni-trivalent graph $\Gamma_{g,b}$ satisfying the following three conditions.

- 1) The uni-trivalent graph $\Gamma_{g,b}$ is the retraction of H_g and have b univalent vertexes.
- 2) The colors of the edges connected to the univalent vertexes are \vec{c} .
- 3) At each trivalent vertexes, the coloring are admissible.

For example, $V_{p,4;1,1,3,3}$ is spanned by the following set of basis.

$$V_{0,4;1,1,3,3} = \left\langle \left(\begin{array}{c} 1 \\ \diagdown \quad \diagup \\ \quad 2 \\ \diagup \quad \diagdown \\ 1 \end{array} ; \begin{array}{c} 3 \\ \diagdown \quad \diagup \\ \quad 3 \\ \diagup \quad \diagdown \\ 1 \end{array} \right) \quad \left(\begin{array}{c} 3 \\ \diagdown \quad \diagup \\ \quad 3 \end{array} \right) \right\rangle$$

Remark 2.1. *The vector spaces we defined are closely related to the representation category of quantum group $SU(2)_q$. (i, j, k) is an admissible triple is equivalent to that the vector space $\text{Inv}(V_i \otimes V_j \otimes V_k)$ is nonempty, where V_i are Verlindre modules, e.g $V_{p,4;1,1,3,3} \cong \text{Inv}(V_1 \otimes V_1 \otimes V_3 \otimes V_3)$ as vector spaces.*

By the point of remark above, given a punctured surface $\Sigma_{g,b}$ and p , we say a coloring \vec{c} is admissible if the dimension of $V_{p,g,b;\vec{c}}$ is positive.

2.2 Mapping class group

Let S be the surface with or without boundary. The mapping class group of S is defined to be

$$\text{MCG}(S) = \text{Diff}(S; \partial S) / \{\text{isotopy}\}.$$

In this section, we define an action of $\text{MCG}(\Sigma_{g,b})$ on vector space $V_{p,g,b;\vec{c}}$.

We first view $V_{p,g,b;\vec{c}}$ as vector space spanned by the skeins in handlebody H_g , with strands and clasps on the boundary if $b \neq 0$. Mapping class groups are generated by Dehn twists. We define the mapping class group action by defining actions of Dehn twists.

Here we introduce a new color Ω , which is defined to be

$$\Omega = \sqrt{2/p} \sin(\pi/p) \sum_{i=0}^{p-2} (-1)^i [i+1] \phi_i,$$

where ϕ_i is the strand colored by i .

Consider a single closed curve γ on $\Sigma_{g,b}$. The action of the Dehn twist D_γ on the skein space is by adding a curve γ colored by Ω with (-1) framing along the boundary of H_g . This action turns out to be projective.

In particular, if γ is the boundary of a disc perpendicular to a strand colored by i , then $D_\gamma v = (-1)^i A^{i(i+2)} v$ under some central extension where v denotes the vector representing the skein.

The following theorem allows us to restrict the (projective)representations of $\text{MCG}(S)$ to $\text{MCG}(S' \subset S)$

Theorem 2.2. [3][The inclusion homomorphism] *Let S be a closed subsurface of a surface S' . Assume that S is not homeomorphic to a closed annulus and no component of $S'-S$ is an open disk. Let $\eta : \text{MCG}(S) \rightarrow \text{MCG}(S')$ be the induced map. Let $\alpha_1, \dots, \alpha_m$ denote the boundary components of S that bound once-punctured disks in $S' - S$ and let $\{\beta_1, \gamma_1\}, \dots, \{\beta_n, \gamma_n\}$ denote the pairs of boundary components of S that bound annuli in $S' - S$. Then the kernel of η is the free abelian group:*

$$\ker(\eta) = \langle T_{\alpha_1}, \dots, T_{\alpha_m}, T_{\beta_1} T_{\gamma_1}^{-1}, \dots, T_{\beta_n} T_{\gamma_n}^{-1} \rangle$$

In particular, if no connected component of $S'-S$ is an open annulus, an open disk, or an open once marked disk, then η is injective.

2.3 Notations and conventions

p is a fixed prime number throughout section 3 to section 6. The index p in the vector space $V_{p,g,b;\vec{c}}$ will be hid for notational simplicity.

Let $[i]$ be the i -th quantum integer, i.e.

$$[i] = \frac{A^{2i} - A^{-2i}}{A^2 - A^{-2}}$$

3 A criterion for irreducibility

Lemma 3.1. [14] Consider a Hermitian vector space X over some field F is a (projective) representation of some group G , and G_1, G_2 are two subgroups. Let $X = \bigoplus_I V_i$ (resp. $\{X = \bigoplus_J W_j\}$) be an irreducible multiplicity free decomposition under the action of some central extension \tilde{G}_1 (resp. \tilde{G}_2) of G_1 (resp. G_2). Define a graph $C(X, G_1, G_2)$ on the set of irreducible summands, with an edge connecting V_j and W_k if there exists some element $v \in V_j$ and $w \in W_k$ such that $[v, w] \neq 0$.

If the graph is connected, then X is irreducible as a representation of G .

Note that decompositions in Lemma 4 do not depend on the choice of central extensions. Thus, when proving irreducibility, G and \tilde{G} will not be distinguished. As discussed in [16], central extension does not affect the irreducibility.

With the setting of Lemma 3.1, each W_j (resp. V_i) is connected with some V_i (resp. W_j) since W_j (resp. V_i) is not $\{0\}$ as a set. Thus, to prove irreducibility, we just need to prove one side of this bipartite graph is connected. We formulate it in the following corollary.

Corollary 3.2. With the assumptions in Lemma 4, we can conclude irreducibility with one of the following conditions on the graph.

- a) for all $i, j \in I, V_i, V_j$ are connected by some path.
- b) there exist $i \in I$ such that V_i is connected to W_j for all $j \in J$.

4 Two base cases

In this section, we discuss two base cases, where the surfaces are sphere with 4 punctures and one-holed torus.

4.1 Sphere with 4 punctures


Lemma 4.1. $V_{0,4;a,b,c,d}$ is irreducible under the action of PB_4 for all admissible a, b, c, d . By admissible, we mean the vector space associated has positive dimension.

Proof. The Dehn twist around γ_1 and γ_2 gives 2 othogonal decompositions:

$$V_{0,4;a,b,c,d} = \bigoplus_{i=\max\{|a-b|, |c-d|\}; i \equiv a-b \pmod{2}}^{\min\{a+b, c+d, 2p-4-a-b, 2p-4-c-d\}} V_i,$$







$$V_{0,4;a,b,c,d} = \bigoplus_{j=\max\{|a-d|, |b-c|\}; i \equiv a-d \pmod{2}}^{\min\{a+d, c+b, 2p-4-a-d, 2p-4-c-b\}} W_j,$$

where V_i (resp. W_j) are spanned by the single vector v_i (resp. w_j).



When p is an odd prime, $(-1)^i A^{i(i+2)}$ are different for different i . Thus, both decompositions are multiplicity free because they have different eigenvalues under the action of D_{γ_t} ($t = 1, 2$).

By lemma 3.1, the irreducibility of the representation of PB_4 is equivalent to the connectivity of the graph $C(V_{0,4;a,b,c,d}, \langle D_{\gamma_1} \rangle, \langle D_{\gamma_2} \rangle)$. Before we proceed, we give the change of basis formula [1] for $\{v_i\}$ and $\{w_j\}$ below.

$$v_i = \sum_j \left\langle \begin{array}{ccc} a & b & i \\ c & d & j \end{array} \right\rangle w_j = \sum_j \frac{\text{Diagram 1} \cdot \text{Diagram 2}}{\text{Diagram 3} \cdot \text{Diagram 4}} w_j$$







where we follow the conventions in [1]:

$$\langle a, b, c \rangle = \text{Diagram 1}; \quad \left\langle \begin{array}{ccc} a & b & i \\ c & d & j \end{array} \right\rangle = \text{Diagram 2}$$

V_i and W_k are connected is equivalent to the span of v_i in $\{w_j\}$ has non-zero coefficient on w_k . According to Corollary 3.2 case (a), Our strategy is to find some i such that V_i is connected to W_j for all j .

Without loss of generality, we assume $a - b \geq |c - d|$ (otherwise, we rotate the symbols), write v_{a-b} as summation of w_j :

$$v_{a-b} = \sum_j \left\langle \begin{array}{ccc} a & b & a-b \\ c & d & j \end{array} \right\rangle w_j \quad (1)$$

If j is admissible, $\langle b, c, j \rangle$ and $\langle a, d, j \rangle$ are nonzero, so we just need to check the tetrahedron symbols. Theorem 2 of [1] gives an explicit formula for the tetrahedron symbol. As in [1], Let $m_1 = (a + b + i)/2$, $m_2 = (a + d + j)/2$, $m_3 = (b + c + j)/2$, $m_4 = (i + d + c)/2$; $n_1 = (a + b + c + d)/2$, $n_2 = (b + i + d + j)/2$, $n_3 = (a + i + c + j)/2$.

$$\left\langle \begin{array}{ccc} a & b & i \\ c & d & j \end{array} \right\rangle = \sum_{z=\max m_s}^{\min n_t} \frac{\prod_{s,t} [n_s - m_t]!}{[a]![b]![c]![d]![i]![j]!} \frac{(-1)^z [z+1]!}{\prod_s [n_s - z]! \prod_t [z - m_t]!} \quad (2)$$

In general, the tetrahedron symbol is a summation over z for all $\max_t n_t \leq z \leq \min_s m_s$. In our case, we only have one summand because $\max_t n_t = z = \min_s m_s$. which is

$$\left\langle \begin{array}{ccc} a & b & a-b \\ c & d & j \end{array} \right\rangle = \frac{\prod_{s,t} [n_s - m_t]!}{[a]![b]![c]![d]![a-b]![j]!} \frac{(-1)^z [z+1]!}{\prod_s [n_s - z]! \prod_t [z - m_t]!} \quad (3)$$

Then we need to check the q -factorials in the above formula are less than p to make sure all factors are nonzero. Note that $2(n_s - m_t)$ can be realized as summation of two labels of an admissible triple subtracting the other one, which is always less than $2(p - 2)$. Thus $n_s - m_t \leq p - 2$, which implies $[n_s - m_t]! \neq 0$; z is half of the sum of labels of an admissible triple, so $z \leq p - 2$, $[z + 1]! \neq 0$.


The above computation showed that V_{a-b} are connected to W_j for all j . By Corollary 3.2(a), $V_{0,4;a,b,c,d}$ is irreducible under the action of PB_4 . □

4.2 One-holed torus


This case has been studied by G. Patrick and G. Masbaum [8]. They proved irreducibility of $V_{g,1;2a}$ for any g when p is an odd prime. In our case, we just need $g = 1$ to start the induction. For completeness, we put a more elementary proof here, and we would like to thank Julien Korinman [13] for teaching us the proof.

Lemma 4.2. *Let p be an odd prime, $1 \leq a \leq \frac{p-3}{2}$, $V_{1,1;2a}$ is irreducible under the action of $MCG(\Sigma_{1,1})$.*

Proof. In [7], their computation showed that the Hopf pairing (see figure below) \langle, \rangle of $V_{1,1;2a}$ is a nondegenerate bilinear form.



Let v_j be the lollipop basis shown below, and $w_i = D_\beta^i v_0$ be the vector derived from applying Dehn twist along β i times to v_0 .



We compute the Hopf pairing of w_i and v_j , where $0 \leq i, j \leq p - a - 2$:

$$\langle w_i, v_j \rangle = \langle D_\beta^i v_0, v_j \rangle = \langle v_0, D_\gamma^i v_j \rangle = (-1)^{ij} A^{i(j+a)(2+j+a)} \langle v_0, v_j \rangle$$

When p is an odd prime, $(-1)^{ij} A^{i(j+a)(2+j+a)}$ are different complex numbers, so matrix $V = \{(-1)^{ij} A^{i(j+a)(2+j+a)}\}_{ij}$ is a Vandermonde matrix, thus invertible. $\{\langle w_i, v_j \rangle\}_{ij} = V \cdot \text{diag}\{\langle v_0, v_j \rangle\}$. $\langle v_0, v_j \rangle \neq 0$ (c. f. [7] Page 100), so the product of the two matrices is invertible, which implies w_i spans $V_{1,1;2a}$.


Consider the bipartite graph $C(V_{1,1;2a}, \langle D_\beta \rangle, \langle D_\gamma \rangle)$. $\langle v_0 \rangle$ is invariant under the action of D_γ . The decomposition of $V_{1,1;2a}$ under the action of $\langle D_\gamma \rangle$ is multiplicity free. The argument above showed that v_0 have component in all eigenspaces of D_β and each eigenspaces are 1-dimensional. That is, $\langle v_0 \rangle$ is connected to all eigenspaces of $\langle D_\beta \rangle$, and all the eigenspaces are multiplicity free due to eigenvalue test. By corollary 3.2(a), $V_{1,1;2a}$ is irreducible. \square

5 The induction

In this section, we will develop the induction steps. The idea is the following: We can decompose the representation by restricting it to a mapping class group of a subsurface $S' \subset S$. Usually $S - S'$ is a cylinder $\alpha \times I$. The decomposition depends on the choice of α on S . Given two different such decompositions, we study the connectivity of the bipartite graph described in Lemma 3.1 to conclude the irreducibility.

The following three lemmas provide us the tools for the induction on genus g and the number of boundaries b .


Lemma 5.1. *Let p be an odd prime and $g \geq 1$, Suppose $V_{g,1,c}$ and $V_{g-1,2;i,j}$ are irreducible as $MCG(\Sigma_{g,1})$ and $MCG(\Sigma_{g-1,2})$ representation respectively for all admissible c, i, j , then $V_{g,2;a,b}$ is an irreducible representation of $MCG(\Sigma_{g,2})$ for all a, b .*



Proof. Let S be a representative of surface $\Sigma_{g,b}$ as above. We restrict $V_{g,2;a,b}$ to the action of the image of $\text{MCG}(S - \alpha \times I) \rightarrow \text{MCG}(S)$ and the image of $\text{MCG}(S - (\beta \cup \gamma) \times I) \rightarrow \text{MCG}(S)$ respectively. According to Theorem 3, the kernels factor through the corresponding representations. Notice $\text{MCG}(S - \alpha \times I) \cong \text{MCG}(\Sigma_{0,3}) \times \text{MCG}(\Sigma_{g,1})$ and $\text{MCG}(S - (\beta \cup \gamma) \times I) \cong \text{MCG}(\Sigma_{0,4}) \times \text{MCG}(\Sigma_{g-1,2})$. Thus, with the assumptions in this lemma, $V_{g,2;a,b}$ has the following two irreducible decompositions accordingly.

$$V_{g,2;a,b} = \bigoplus_c A_c = \bigoplus_c V_{g,1;c} \otimes V_{0,3;c,a,b}; V_{g,2;a,b} = \bigoplus_{i,j} B_{i,j} = \bigoplus_{i,j} V_{g-1,2;i,j} \otimes V_{0,4;i,j,a,b},$$


where A_c and $B_{i,j}$ are invariant spaces of D_α and $D_\beta \times D_\gamma$ respectively. They are spanned by the graphs shown below.



When p is odd prime, these invariant spaces have different eigenvalues for the Dehn twists, so the decompositions are multiplicity free for all central extensions.

According to Lemma 3.1, we just need to prove the bipartite graph $C(V_{g,2;a,b}, \text{MCG}(S - \alpha \times I), \text{MCG}(S - (\beta \cup \gamma) \times I))$ is connected.


Note that if (c, i, j) is an admissible triple, A_c and $B_{i,j}$ are connected because the following element will be in the intersection of A_c and $B_{i,j}$.



Then notice for all c , $(c, \frac{p-3}{2}, \frac{p-3}{2})$ will be an admissible triple, so all A_c are connected to $B_{\frac{p-3}{2}, \frac{p-3}{2}}$. $B_{\frac{p-3}{2}, \frac{p-3}{2}}$ is of positive dimension for all g . Thus, $V_{g,2;a,b}$ is an irreducible $\text{MCG}(\Sigma_{g,2})$ representation by corollary 3.2. \square

If all boundaries are colored by 0, we should consider it as the closed surface. This case not only shows up in the question itself, but also contribute to the induction. Although closed surface case have been proved by J. Roberts in [16], we still put this case in our induction for completeness.


Lemma 5.2. *Let p be an odd prime and $g \geq 2$. Assume $V_{g-1,2;i,i}$ is irreducible under the action of $\text{MCG}(\Sigma_{g-1,2})$ for all i . Then $V_{g,0}$ is irreducible under the action of $\text{MCG}(\Sigma_g)$.*



Proof. Let S be a representative of $\Sigma_{g,0}$ as above. We restrict $V_{g,0}$ to the representation of subgroups of the image of $\text{MCG}(S - \alpha \times I)$ and the image of $\text{MCG}(S - \beta \times I)$ in $\text{MCG}(S)$ respectively. According to Theorem 2.2, the kernels factor through the corresponding representations. Notice that both $\text{MCG}(S - \alpha \times I)$ and $\text{MCG}(S - \beta \times I)$ are isomorphic to $\text{MCG}(\Sigma_{g-1,2})$. Thus, with the assumptions in this lemma, $V_{g,0}$ has the following two irreducible decompositions accordingly.

$$V_{g,0} \cong \oplus_i A_i = V_{g-1,2;i,i}; V_{g,0} \cong \oplus_j B_j = V_{g-1,2;j,j}$$

Where A_i and B_j are invariant spaces of D_α and D_β respectively. They are spanned by the graphs shown below:




When p is odd prime, these invariant spaces have different eigenvalues under the action of D_α and D_β respectively, so the decompositions are multiplicity free for all central extension of $\text{MCG}(\Sigma_g)$. Notice the following element is in the intersection of A_i and B_j :



Argument above showed that the bipartite graph $\langle V_{g,0}; \text{MCG}(S - \alpha \times I), \text{MCG}(S - \beta \times I) \rangle$ is complete. Thus, the representation is irreducible by Lemma 3.1. \square

Before we introduce the next lemma, we define \prec to be the lexicographical order on pair (g, b) .

Lemma 5.3. *Let p be an odd prime and $(g, b) \notin \{(0, 1), (0, 2), (0, 3), (0, 4), (1, 1), (1, 2)\}$ and $b \geq 0$. Consider vector space $V_{g,b;\vec{c}}$ with $b \geq 1$ and one of the boundary is colored by $a \neq p - 2$. If $V_{g',b';\vec{c}}$ is irreducible under the action of $\text{MCG}(\Sigma_{g',b'})$ for all $(g', b') \prec (g, b)$ and any color \vec{c} , $V_{g,b;\vec{c}}$ is irreducible under the action of $\text{MCG}(\Sigma_{g,b})$.*




Proof. With the numerical conditions on (g, b) described in the lemma, we can find $(g_1, b_1) \prec (g, b)$ and $(g_2, b_2) \prec (g, b)$ satisfying $g = g_1 + g_2$ and $b = b_1 + b_2 - 1$. By restricting to the group actions of $\text{MCG}(S - \alpha \times I)$ and $\text{MCG}(S - \beta \times I)$, we can decompose the representation $V_{g,b;\vec{c}} = V_{g_1+g_2, b_1+b_2+1; \vec{c}_1, \vec{c}_2, a}$ in following two ways:

$$A_i = V_{g_1, b_1+1; \vec{c}_1, i} \otimes V_{g_2, b_2+2; \vec{c}_2, i, a}$$


$$B_j = V_{g_1, b_1+2; \vec{c}_1, a, j} \otimes V_{g_2, b_2+1; \vec{c}_2, j}$$

Where A_i and B_j are invariant spaces of D_α and D_β respectively. They are spanned by the graphs shown below:



where i (resp. j) run through all colors such that both of their tensor factor have positive dimension. That is, if both g_1 and g_2 are positive, i and j run through all colors satisfying the parity condition. If g_1 or g_2 is zero, then we have extra inequality constraints for i and j , but these constraints always reduce to an interval.

By the assumption, $V = \oplus_i A_i$ (resp. $V = \oplus_j B_j$) is an irreducible and multiplicity free (by checking eigenvalues of the Dehn twist around the circle in the graph) decomposition under the group action of $\text{MCG}(\Sigma_{g_1, b_1+1}) \times \text{MCG}(\Sigma_{g_2, b_2+2})$ (resp. $\text{MCG}(\Sigma_{g_1, b_1+2}) \times \text{MCG}(\Sigma_{g_2, b_2+1})$), and we noticed that if (i, j, a) is an admissible triple, A_i and B_j are connected because the following element is in the intersection:



Now we prove the following claim: Suppose $i \geq i'$ and there exist j, j' such that (i, j, a) and (i', j', a) are admissible triples, then there exist some j'' such that (i, j'', a) and $(i-2, j'', a)$ are admissible triples.

If $(i-2, j, a)$ is admissible, we are done. Otherwise, one of the inequality conditions must fail, so we have either $i+j = a$ or $j-i = a$:

Case 1 $i+j = a$: $i' + j' \geq a$ and $i' < i$ implies $j' > j$, so B_{j+2} is not $\{0\}$. Let $j'' = j+2$, we check $(i-2) + j'' = a$, $|(i-2) - j''| = |i-4 + j| \leq |i-4| + j$. Since $i > i', i \geq 2$, so $|i-4| + j \leq i+j = a$. Thus $(i-2, j+2, a)$ is an admissible triple.

Case 2 $j-i = a$: $j' - i' \leq a$ and $i' < i$ implies $j' < j$, so B_{j-2} is not $\{0\}$. Let $j'' = j-2$, we check $|(i-2) - j''| = a$, $(i-2) + j'' = j-i+2i-4 \geq j-i = a$. Thus $(i-2, j-2, a)$ is an admissible triple.

We take i' to be the least i such that $A_i \neq 0$. The claim said $B_{j''}$ is connected to A_i and A_{i-2} . This gives us paths connect all pairs of adjacent A'_i 's. Together with corollary 3.2(b), we proved $V_{g_1+g_2, b_1+b_2+1; \vec{c}_1, \vec{c}_2, a}$ is irreducible. \square

6 Proof of the theorem 1.1

In this section, we prove the theorem 1.1.

Proof. We are going to do induction on the lexicographical order \prec of pairs (g, b) . We just need to show that for all $V_{g,b;\vec{c}}$ where $(g, b) \notin \{(0, 1), (0, 2), (0, 3), (0, 4), (1, 1), (1, 2)\}$, The irreducibility of $V_{g,b;\vec{c}}$ is implied by some $V_{g',b';\vec{c}'}$ satisfying $(g', b') \prec (g, b)$.

We first consider the case that one of the boundary is colored by $p-2$. Because summation on colors on the boundary should be an even number, so $b \geq 2$. Pick another boundary that colored by i . We have the following isomorphism of $\text{MCG}(\Sigma_{g,b-1})$ representations:

$$V_{g,b;\vec{c}} = V_{0,3;p-2,i,p-2-i} \otimes V_{g,b-1;\vec{c},p-2-i \setminus \{p-2,i\}}$$

By this, we lower the number of boundary components by 1.

If $b \geq 1$ and none of the boundary is colored by $p-2$, and $(g, b) \notin \{(0, 1), (0, 2), (0, 3), (0, 4), (1, 1), (1, 2)\}$, Lemma 5.3 implies it is enough to show $V_{g',b';\vec{c}'}$ are irreducible for all $(g', b') \prec (g, b)$.

If $b = 0$ and $g \leq 2$, Lemma 5.2 implies it is enough to show $V_{g,2;i,i}$ is irreducible.

If $(g, b) = (1, 2)$, Lemma 5.1 implies it is enough to show $V_{1,1;a}$ is irreducible.

For Base cases:

- $(g, b) = (0, 1), (0, 2), (0, 3)$. The vector spaces are either 0 or 1-dimensional.
- $(g, b) = (1, 0)$. This is Weil representation, and it is irreducible.
- $(g, b) = (0, 4)$ and $(1, 1)$. We proved them in section 4.

The above induction proves $V_{g,b;\vec{c}}$ is irreducible for all g, b and all coloring \vec{c} . \square

7 Tools for prove denseness

In the section, we introduce some tools to study images of group homomorphisms that originate with Goursat's Lemma.

In generic q case, we no longer have the Hermitian form on the skein spaces. Thus, we rephrase lemma 3.1.

Lemma 7.1. [14] Consider a vector space X over some field F is a (projective) representation of some group G , and G_1, G_2 are two subgroups. Let $X = \bigoplus_I V_i$ (resp. $\{X = \bigoplus_J W_j\}$) be an irreducible multiplicity free decomposition under the action of some central extension \tilde{G}_1 (resp. \tilde{G}_2) of G_1 (resp. G_2). Define a directed graph $C(X, G_1, G_2)$ on the set of irreducible summands, with an edge from V_j to W_k if there exists some element $v \in V_j$ has nonzero component in W_k .

If the graph is strongly connected, then X is irreducible as a representation of G .

The following theorem is often called non-commutative Chinese remainder theorem, that locally surjectivity implies global surjectivity.

Theorem 7.2. *Suppose that each of G_1, G_2, \dots, G_l is a minimal simple Lie group or a non-abelian finite simple group, and suppose that*

$$H \subset G = G_1 \times G_2 \times \dots \times G_l$$

is a closed subgroup that surjects onto each factor G_k . Then H is a diagonal subgroup of G .

In our case, group H is the Zariski closure of the mapping class group and G_i are complex simple Lie groups. Thus, we have a corollary for our case.

Corollary 7.3. *If*

$$f : H \rightarrow PSL(W_1) \times PSL(W_2) \times \dots \times PSL(W_n)$$

is surjective when restrict to $H \rightarrow G_i$ for all i , and non of pairs W_i and W_j are isomorphic or dual to each other, then f is surjective.

Lastly, we need the following surjectivity theorem:

Theorem 7.4. *Let V be a finite-dimensional complex representation of a connected Lie group G , Let $H \subset G$ be a closed, connected subgroup, and let*

$$V|_H \cong \bigoplus_{k=1}^n W_k$$

be the decomposition of the restricted representation. Suppose that:

1. V is G -irreducible.
2. At most one of W_i is of dimension 2 and at most one of W_i is of dimension 1.
3. For every $j \neq k$, the summands W_j and W_k are neither isomorphic nor dual as projective representation of H .
4. For each j , H surjects onto $PSL(W_j)$.

Then G surjects onto $PSL(V)$.

Proof. Let L be the Lie algebra of G , $K \subset L$ be the corresponding Lie algebra of H . Then we have the splitting

$$\mathfrak{gl}(V) = \mathfrak{sl}(V) \oplus \mathbb{C}$$

The non-commutative Chinese remainder theorem for Lie algebra gives us that H is jointly surjective:

$$H \twoheadrightarrow \bigoplus_k \mathfrak{sl}(W_k)$$

Meanwhile we have the partial decomposition

$$\mathfrak{sl}(V) = \bigoplus_{j \neq k} W_j \otimes W_k^* \oplus \bigoplus_k \mathfrak{sl}(W_k) \oplus \mathbb{C}^{n-1}$$

Notice that if for some i , $\dim W_i > 2$, then $W_i \not\cong W_i^*$ as $\mathfrak{sl}(W_i)$ representation. Together with the 3rd assumption in the theorem, $W_i \otimes W_j^*$ and $W_j \otimes W_i^*$ are both unique in the decomposition in the sense that they are not isomorphic to each other and all other summands as a representation of H . That is, The off-diagonal blocks are multiplicity free if and only if at most one of W_i satisfying $\dim W_i \leq 2$. In the rest of the proof, we discuss it case by case. Without lose of generality, we assume the $\dim W_i \leq \dim W_j$ if $i \leq j$.

Case 1 $\dim W_2 \geq 3$.

As we discussed above, the off diagonal summands are multiplicity free in this case. so the image of L in $\mathfrak{sl}(V)$ contains a subset of the off-diagonal blocks $W_j \otimes W_k^*$ and some subspace of \mathbb{C}^{n-1} . We can make a directed graph Γ with a directed edge $k \rightarrow j$ for every off-diagonal block $V_j \otimes V_k^*$ which is in the image of L . We claim Γ is complete.

First, we prove Γ is strongly connected. Assuming otherwise, Γ would have a strongly connected component C with no outward edges. In this case $\bigoplus_{k \in C} W_k$ would be a non-trivial subrepresentation of L . G is connected, so $\bigoplus_{k \in C} W_k$ is also a subrepresentation of G , contradicting the hypothesis that V is irreducible.

Second, we prove that Γ is transitively closed. Suppose that $i \rightarrow k \rightarrow j$ is a path of length two using three distinct vertices. Choose two operators

$$\begin{aligned} Y &\in \text{Hom}(W_i, W_k) \cong W_k \otimes W_i^* \\ X &\in \text{Hom}(W_k, W_j) \cong W_j \otimes W_k^* \end{aligned}$$

whose product

$$XY \in \text{Hom}(W_i, W_j) \cong W_j \otimes W_i^*$$

is non-zero. Then $[X, Y]$ is a non-zero element in $W_j \otimes W_i^*$. The image of L thus contains some elements of $W_j \otimes W_i^*$, and therefore it contains all of them and Γ is transitively closed.

If Γ is both strongly connected and transitively close, then it is the complete directed graph. In the final step, choose some basis of V that refines the decomposition of $V|_H$. In this basis, the image of L contains all off-diagonal elementary matrices, so there commutators gives us all of $\mathfrak{sl}(V)$

Case 2 $n \geq 3$ and $\dim W_2 = 2, \dim W_1 = 1$.

Define $V' = \bigoplus_{k=2}^n W_k$. The identical argument as in case 1 shows that $\mathfrak{sl}(V')$ is in the image of L . Thus we have decomposition

$$\mathfrak{sl}(V) = \mathfrak{sl}(V') \oplus V' \oplus V'^* \oplus \mathbb{C}$$

as representation of $\mathfrak{sl}(V')$. The decomposition is multiplicity free since $\dim V' \geq 3$. V is irreducible as L representation, so The image of L contains both V and V^* . The commutators will give us all of $\mathfrak{sl}(V)$.

Case 3 $n = 2$ and $\dim W_2 = 2, \dim W_1 = 1$.

In this case, the V is a 3-dimensional representation. we list all the Lie subalgebra of $\mathfrak{sl}(V)$. Only $\mathfrak{sl}(V)$ itself makes V irreducible. □

Before we get to the last section, we introduce a lemma conjectured by the first author and supported by David Speyer. It is useful for analyzing weight diagrams.

Lemma 7.5 (Speyer). *Let $D(\lambda)$ be the set of weights in the weight diagram of the irreducible representation $V(\lambda)$ of a simple Lie algebra \mathfrak{g} . Let $S(\lambda) = \lambda - D(\lambda)$ as a set of vectors in \mathfrak{h}^* . The indecomposable elements of $S(\lambda)$ are roots of \mathfrak{g} .*

Proof. Let R be the set of positive roots of \mathfrak{g} . $C = S(\lambda) \cap R$, $C' = R - C$ and v_λ be a vector in $V(\lambda)$ have weight λ . Notice $U(\mathfrak{n}_-)(v_\lambda) = V(\lambda)$. By PBW theorem, $U(\mathfrak{n}_-)$ is spanned by monomials of form $f_{c_1} f_{c_2} \dots f_{c_m} f_{c'_1} f_{c'_2} \dots f_{c'_n} v_\lambda$ where $c_i \in C$ and $c'_j \in C'$. Since $\lambda - c'_j \notin D(\lambda)$, we have $f_{c'_j} v_\lambda = 0$ if $c'_j \neq 0$, so $V(\lambda)$ is spanned by $f_{c_1} f_{c_2} \dots f_{c_m} v_\lambda$. That is, for all weight $\lambda' \in D(\lambda)$ such that $\lambda - \lambda'$ indecomposable in $S(\lambda)$. λ' is either λ or $v_{\lambda'} = f_{c_i} v_\lambda$. This proves $\lambda - \lambda_i = c_i$ is a positive root. □

8 Proof of the theorem 1.2

The proof is by induction on number of boundaries. The induction begins at $b = 4$.

Lemma 8.1. *Let A be a transcendental complex number. \widetilde{PB}_4 is dense in the algebraic group $SL(V_{0,4;a,b,c,d})$ if $\dim V_{0,4;a,b,c,d} \geq 2$*

Proof. • The representation is irreducible.

We adapt the notations of section 4.1 and claim the graph $C(V_{0,4;a,b,c,d}, \langle D_{\gamma_1} \rangle, \langle D_{\gamma_2} \rangle)$ is complete.

We still have the change of basis formula (1) in section 4.1. To prove the v_i have a non-zero component in W_j , need to show coefficient

$$\left\langle \begin{array}{ccc} a & b & i \\ c & d & j \end{array} \right\rangle = \sum_{z=\max m_s}^{\min n_t} \frac{\prod_{s,t} [n_s - m_t]!}{[a]![b]![c]![d]![i]![j]!} \frac{(-1)^z [z+1]!}{\prod_s [n_s - z]! \prod_t [z - m_t]!} \neq 0. \quad (4)$$

Notice that when consider it as a rational function of A , the degree is different among its summands. Thus, the leading term only appeared in one of the summands. Since A transcendental, it is not a root of any rational function, so the coefficient is not zero. We proved the graph is complete and the representation is irreducible.

- The image is infinite.

Consider two consecutive colors v_i and v_{i+2} . Let λ_i and λ_{i+1} be the eigenvalues of D_{γ_1} respectively. $\lambda_{i+1}/\lambda_i = -A^{4i+8}$. A is not a root of unity. Thus, the image is infinite.

- The closure of the image is $SL(V_{0,4;a,b,c,d})$.

Assume the Zariski closure of the image is a semisimple lie algebra G . the action of D_{γ_1} have the following eigenvalue set under some central extension:

$$X = \{A^{i^2} | i \in [\max\{|a-b|, |c-d|\}, \min\{|a+b|, |c+d|\}], i \equiv a-b \pmod{2}\}$$

Let's assume i start at i_0 and end at i_n . There exist an element $a \in \mathfrak{g} = Lie(G)$, the eigenvalues for a acting on $V_{0,4;a,b,c,d}$ is $\lambda_0 = C + i_0^2, \lambda_1 = C + (i_0 + 2)^2, \dots, \lambda_n = C + (i_n)^2$ for some constant C . We claim \mathfrak{g} is \mathfrak{sl}_{n+1} .

Notice in the weight space \mathfrak{h}^* , λ_0 is the highest weight under some choice of simple roots. $\lambda_0 - \lambda_1$ satisfying the following properties:

1. $\lambda_0 - \lambda_1$ is not repetitive, that is, $\lambda_0 - \lambda_1 \neq \lambda_k - \lambda_j$ for any $(k, j) \neq (0, 1)$.
2. $\lambda_0 - \lambda_1$ is indecomposable, that is $\lambda_0 - \lambda_1$ can not be written as positive integral linear combination of $\lambda_0 - \lambda_k$ for $k \neq 1$.

By Speyer's lemma, $\lambda_0 - \lambda_1$ is a root. Consider the action of $W_{\lambda_0 - \lambda_1}$ on the weight diagram. It send λ_k to λ_k plus(or minus) some copy of $\lambda_0 - \lambda_1$. Because $\lambda_0 - \lambda_1$ is not repetitive, $W_{\lambda_0 - \lambda_1}$ interchanges λ_0 and λ_1 and fix all other λ_k .

Then we prove for all k , $\lambda_0 - \lambda_k$ is indecomposable. Assume otherwise, say $\lambda_0 - \lambda_k = \sum_t \alpha_t (\lambda_0 - \lambda_t)$. the decomposition is perserved by the Weyl group action. We have $\lambda_1 - \lambda_k = \sum_t \alpha_t (\lambda_1 - \lambda_t)$. These two equation together implies $\sum_t \alpha_t = 1$. Thus $\lambda_0 - \lambda_k$ is indecomposable for all k .

Use Speyer's lemma again, we know all $\lambda_0 - \lambda_k$ are roots. Consider the action of $W_{\lambda_0 - \lambda_k}$. It interchanges λ_0 and λ_k . Otherwise, $\lambda_0 - W_{\lambda_0 - \lambda_k}(\lambda_0)$ will be multiple of $\lambda_0 - \lambda_k$. The action also have to fix all other λ_i . Assume otherwise, it send λ_i to λ_j . then $\lambda_0 - \lambda_j = \lambda_0 - \lambda_i + (\lambda_i - \lambda_j)$ which is equal to $\lambda_0 - \lambda_i$ plus multiple of $\lambda_0 - \lambda_k$, so the indecomposable condition is contradicted.

$W_{\lambda_0 - \lambda_k}$ generate S_{n+1} . The lie algebra \mathfrak{g} have an $(n+1)$ -dimensional irreducible representation and has S_{n+1} as a quotient of a subgroup of its Weyl group. \mathfrak{g} have to be \mathfrak{sl}_{n+1} . □


To apply theorem 7.4, we need the following proposition about irreducibility.

Proposition 8.2. *For A any transcendental number, $V_{0,b;\bar{c}}$ is irreducible as a representation of PB_b .*

Proof. The proof is identical to lemma 5.3. Notice that we find a element in the intersection of V_i and W_j . In transcendental case, this means we have directed edges of both direction. □

The following lemma gives a criterion that for some specific decomposition, the components is not isomorphic nor dual to each other.

Lemma 8.3. *Let A be a transcendental number, and c_1, c_2, \dots, c_n be a sequence of fixed non-decreasing natural numbers and $n \geq 3$. Then there is no pair of elements in $\{V_{0,n+1;a,c_1,\dots,c_n} \mid a \leq 2c_1\}$ are isomorphic or dual to each other.*



Proof. Consider the Dehn twist around the curve γ that bounds the first two boundaries. The eigenvalue set will be $E_a = \{A^{i^2} \mid \max\{c_n - \sum_{k=2}^{n-1} c_k, |a - c_1|\} \leq i \leq \min\{a + c_1, \sum_{k=2}^n c_k\}\}$ up to some central extension. For different a , we claim either the cardinality of the eigenvalue set will be different or the set of ratios of eigenvalues are different. Set of ratios of eigenvalues is an invariant of central extensions, so the claim implies the lemma.

Suppose $a, a' \leq c_1$ such that E_a and $E_{a'}$ have the same cardinality. Then the interval for i will be of the same length. To make the set of ratios of eigenvalues the same, the interval should start and end at the same place. Since we assumed $a, a' \leq 2c_1 \leq 2c_2$, so at least one of the boundary of the interval is determined by a or a' . Thus we proved the claim. \square

Now we can give the proof of theorem 1.2 by induction:

Proof. The case number of boundary components $b = 4$ is proved in lemma 8.1.

Suppose the Zariski closure of the image of PB_i in $PSL(V_{0,i;\vec{c}})$ is surjective for all $i \leq n$ and coloring \vec{c} . we prove PB_{n+1} has a dense image in $PSL(V_{0,n+1;\vec{c}'})$ for any \vec{c}' .

Let G be the Zariski closure of PB_{n+1} . G contains elements of infinite order, so G must be of positive dimension. G is generated by 1-dimensional subgroups that densely generated by Dehn twists, so G is connected.

Without loss of generality, assume $\vec{c}' = (c_0, c_1, c_2, \dots, c_n)$ such that $c_i \leq c_j$ if $i \leq j$. Then we can have a decomposition of $V_{0,n+1;\vec{c}'}$ by restricting to a subgroup $H = MCG(S') \cong PB_n$.

$$V_{0,n+1;\vec{c}'} = \bigoplus_{i=c_1-c_0}^{c_0+c_1} V_{0,n;i,c_2,c_3,\dots,c_n}$$

According to lemma 8.3, all summands are not isomorphic nor dual to each other.

To apply theorem 7.4, we prove the dimension of the summands satisfying (2) of theorem 7.4.

If the number of boundary component $b \geq 4$, the dimension of the vector spaces $V_{0,b;\vec{c}}$ have dimension 1 if and only if one of the colors is equal to the summation of colors on all other boundaries. In our case, $b \geq 5$, $n \geq 4$, $i \leq c_0 + c_1$. $\dim V_{0,n;i,c_2,c_3,\dots,c_n} = 1$ if and only if $i = c_n - c_{n-1} - \dots - c_2$, so it happens at most once.

Next, we consider the 2-dimensional summands. We fix a uni-trivalent tree Γ with n boundary vertex colored by \vec{c} . If $\dim V_{0,n;\vec{c}} = 2$, then we have 2 admissible coloring for Γ . Fix an edge e that have different colors a and $a+2$ in the two different admissible colorings. Cut the tree at e , the tree Γ split to Γ_1 and Γ_2 , and coloring e by a and $a+2$ both give unique admissible coloring for Γ_1 and Γ_2 . This implies Γ_1 and Γ_2 can only have 3 boundary vertices and $n = 4$. When $n = 4$, we can check by hand that at most for only one $i \leq c_0 + c_1$, $\dim V_{0,n;i,c_2,c_3,\dots,c_n} = 2$.

By theorem 7.4, G surjects onto $PSL(V_{0,n+1;\vec{c}'})$. \square

References

- [1] Christian Blanchet, Nathan Habegger, Gregor Masbaum, and Pierre Vogel. Three-manifold invariants derived from the Kauffman bracket. *Topology*, 31(4):685–699, 1992.
- [2] Christian Blanchet, Nathan Habegger, Gregor Masbaum, and Pierre Vogel. Topological quantum field theories derived from the kauffman bracket. *Topology*, 34(4):883–927, 1995.
- [3] Benson Farb and Dan Margalit. *A primer on mapping class groups*. Princeton University Press, 2011.

- [4] Michael Freedman and Vyacheslav Krushkal. On the asymptotics of quantum $su(2)$ representations of mapping class groups. In *Forum Mathematicum*, volume 18, pages 293–304. Walter de Gruyter, 2006.
- [5] Michael H Freedman, Michael J Larsen, and Zhenghan Wang. The two-eigenvalue problem and density of jones representation of braid groups. *Communications in mathematical physics*, 228(1):177–199, 2002.
- [6] Louis Funar. On the TQFT representations of the mapping class groups. *Pacific J. Math.*, 188(2):251–274, 1999.
- [7] Patrick Gilmer and Gregor Masbaum. Integral topological quantum field theory for a one-holed torus. *Pacific journal of mathematics*, 252(1):93–112, 2011.
- [8] Patrick M Gilmer and Gregor Masbaum. Irreducible factors of modular representations of mapping class groups arising in integral tqft. *arXiv preprint arXiv:1110.5666*, 2011.
- [9] Patrick M Gilmer and Gregor Masbaum. An application of tqft to modular representation theory. *Inventiones mathematicae*, 210(2):501–530, 2017.
- [10] Thomas Koberda and Ramanujan Santharoubane. Irreducibility of quantum representations of mapping class groups with boundary. *arXiv:1701.08901*.
- [11] Thomas Koberda and Ramanujan Santharoubane. Quotients of surface groups and homology of finite covers via quantum representations. *Inventiones mathematicae*, 206(2):269–292, 2016.
- [12] Julien Korinman. Decomposition of the weil representations at even levels into irreducible factors. *arXiv:1310.0390*.
- [13] Julien Korinman. On the (in)finiteness of the image of reshetikhin-turaev representations. *arXiv:1412.2671*.
- [14] Greg Kuperberg. Denseness and zariski denseness of jones braid representations. *Geometry & Topology*, 15(1):11–39, 2011.
- [15] Michael Larsen and Zhenghan Wang. Density of the $SO(3)$ TQFT representation of mapping class groups. *Comm. Math. Phys.*, 260(3):641–658, 2005.
- [16] Justin Roberts. Irreducibility of some quantum representations of mapping class groups. *Journal of Knot Theory and Its Ramifications*, 10(05):763–767, 2001.