

**CRITICAL EXPONENT FOR THE SEMILINEAR WAVE
EQUATIONS WITH A DAMPING INCREASING IN THE FAR
FIELD**

KENJI NISHIHARA, MOTOHIRO SOBAJIMA, AND YUTA WAKASUGI

ABSTRACT. We consider the Cauchy problem of the semilinear wave equation with a damping term

$$\begin{cases} u_{tt} - \Delta u + c(t, x)u_t = |u|^p, & (t, x) \in (0, \infty) \times \mathbb{R}^N, \\ u(0, x) = \varepsilon u_0(x), \quad u_t(0, x) = \varepsilon u_1(x), & x \in \mathbb{R}^N, \end{cases}$$

where $p > 1$ and the coefficient of the damping term has the form

$$c(t, x) = a_0(1 + |x|^2)^{-\alpha/2}(1 + t)^{-\beta}$$

with some $a_0 > 0$, $\alpha < 0$, $\beta \in (-1, 1]$. In particular, we mainly consider the cases

$$\alpha < 0, \beta = 0 \quad \text{or} \quad \alpha < 0, \beta = 1,$$

which imply $\alpha + \beta < 1$, namely, the damping is spatially increasing and effective. Our aim is to prove that the critical exponent is given by

$$p = 1 + \frac{2}{N - \alpha}.$$

This shows that the critical exponent is the same as that of the corresponding parabolic equation

$$c(t, x)v_t - \Delta v = |v|^p.$$

The global existence part is proved by a weighted energy estimates with an exponential-type weight function and a special case of the Caffarelli-Kohn-Nirenberg inequality. The blow-up part is proved by a test-function method introduced by Ikeda and Sobajima [15]. We also give an upper estimate of the lifespan.

1. INTRODUCTION

We consider the Cauchy problem of the semilinear wave equation with a damping term

$$(1.1) \quad \begin{cases} u_{tt} - \Delta u + c(t, x)u_t = |u|^p, & (t, x) \in (0, \infty) \times \mathbb{R}^N, \\ u(0, x) = \varepsilon u_0(x), \quad u_t(0, x) = \varepsilon u_1(x), & x \in \mathbb{R}^N, \end{cases}$$

where $N \geq 1$, $u = u(t, x)$ is a real-valued unknown function, ε is a small positive parameter, u_0, u_1 are given initial data, $p > 1$, and the coefficient of the damping term has the form

$$(1.2) \quad c(t, x) = a(x)b(t) = a_0 \langle x \rangle^{-\alpha} (1 + t)^{-\beta},$$

Key words and phrases. semilinear damped wave equation; time and space dependent damping; critical exponent; lifespan.

2010 Mathematics Subject Classification. 35L15; 35A01, 35B44

where $\langle x \rangle := \sqrt{1 + |x|^2}$, $a(x) = a_0 \langle x \rangle^{-\alpha}$, $b(t) = (1+t)^{-\beta}$, with some $a_0 > 0$, $\alpha < 2$ and $\beta > -1$. In particular, in this paper we mainly consider the cases

$$(1.3) \quad \alpha < 0, \beta = 0 \quad \text{or} \quad \alpha < 0, \beta = 1.$$

We assume that the initial data satisfies

$$(1.4) \quad u_0 \in H^1(\mathbb{R}^N), \quad u_1 \in L^2(\mathbb{R}^N), \quad \text{supp}(u_0, u_1) \subset \{x \in \mathbb{R}^N; |x| \leq R_0\}$$

with some $R_0 > 0$.

Our aim is to determine the critical exponent p_c . Here, the meaning of the *critical exponent* is the following: if $p > p_c$, then for any (u_0, u_1) satisfying (1.4), there exists a unique global solution for sufficiently small ε ; if $p \leq p_c$, then there exists (u_0, u_1) satisfying (1.4), the local solution blows up in finite time for any small ε . In this paper, we will show that under the conditions (1.2) and (1.3), the critical exponent is determined by

$$p_c = 1 + \frac{2}{N - \alpha}.$$

Comparing with previous studies we will explain below, our novelties are to determine the critical exponent for spatially increasing damping, and to give the blow-up of solutions for the damping depending on the time and space variables.

The Cauchy problem of the linear damped wave equation

$$(1.5) \quad \begin{cases} u_{tt} - \Delta u + c(t, x)u_t = 0, & (t, x) \in (0, \infty) \times \mathbb{R}^N, \\ u(0, x) = \varepsilon u_0(x), \quad u_t(0, x) = \varepsilon u_1(x), & x \in \mathbb{R}^N, \end{cases}$$

with a damping coefficient $c(t, x) = a(x)b(t) = a_0 \langle x \rangle^{-\alpha} (1+t)^{-\beta}$ has been studied for a long time. Roughly speaking, it is known that if the damping is sufficiently strong, in other words, effective, then the solution behaves like that of the corresponding parabolic equation $c(t, x)u_t - \Delta u = 0$ (diffusion phenomenon). On the other hand, if the damping is sufficiently weak, in other words, non-effective, then the solution behaves like that of the wave equation without damping (scattering).

It is known that the classical damping $\alpha = \beta = 0$ is included in the effective case, and the diffusion phenomenon was studied by [40, 13, 45, 26, 75, 46, 39, 12, 44, 2, 54, 41].

On the other hand, Yamazaki [74] and Wirth [70, 71, 72, 73] considered time-dependent damping $\alpha = 0, \beta \in \mathbb{R}$, and classified the behavior of the solution in the following way: (i) Scattering: if $\beta > 1$, then the solution behaves like that of the wave equation without damping; (ii) Scale-invariant weak damping: if $\beta = 1$, then the asymptotic behavior of the solution depends on a_0 ; (iii) Effective: if $-1 \leq \beta < 1$, then the solution behaves like that of the corresponding parabolic equation; (iv) Overdamping: if $\beta < -1$, then the solution does not decay to zero in general.

The space-dependent damping $\alpha \in \mathbb{R}, \beta = 0$ was also studied by [53, 42, 21, 60, 25, 65, 55, 56, 57, 58], and similarly to the above, the behavior of the solution was classified in the following way: (i) Scattering: if $\alpha > 1$, then the solution behaves like that of the wave equation without damping; (ii) Scale-invariant weak damping: if $\alpha = 1$, then the asymptotic behavior of the solution depends on a_0 ; (iii) Effective: if $\alpha < 1$, then the solution behaves like that of the corresponding parabolic equation. We note that in the space-dependent case, the overdamping phenomenon does not occur.

In a similar approach to the space-dependent case, these results are partially extended to the space-time dependent damping $\alpha, \beta \in \mathbb{R}$. Mochizuki and Nakazawa

[43] proved that the case $\alpha, \beta \geq 0, \alpha + \beta > 1$ belongs to the scattering case. For $0 \leq \alpha < 1, -1 < \beta < 1, 0 < \alpha + \beta < 1$, by [27, 28], energy estimates of solutions were obtained and they indicate the solution has diffusion phenomenon.

Based on the studies on the linear problem, recently, the Cauchy problem of the semilinear damped wave equation (1.1) has been intensively studied. In particular, if the damping is effective, we expect that the critical exponent is the same as that of the corresponding parabolic problem. Indeed, when $\alpha = \beta = 0$, it was shown by [34, 59, 76, 30] that the critical exponent is given by $p_c = p_F(N) = 1 + \frac{2}{N}$, which is called the Fujita exponent named after the pioneering work by [8].

For the time-dependent damping case, namely, $\alpha = 0$, it was revealed by [37] that the critical exponent remains $p_c = p_F(N)$ when $\alpha = 0, \beta \in (-1, 1)$ (see [67, 9, 18] for the case $\beta = -1$). When $\beta > 1$, Lai and Takamura [31] and Wakasa and Yordanov [63] showed the small data blow-up for the sub-Strauss or Strauss exponent $1 < p < \infty$ ($N = 1$); $1 < p \leq p_S(N) = \frac{N+1+\sqrt{N^2+10N-7}}{2(N-1)}$ ($N \geq 2$). Recently, Liu and Wang [38] gave the global existence result for $p > p_S(N)$ with $N = 3, 4$, while the cases $N = 2$ and $N \geq 5$ remain open. On the other hand, in the scale-invariant case $\beta = 1$, the situation becomes more complicated. First, if a_0 is sufficiently large, we expect that the critical exponent coincides with $p_F(N)$. Indeed, by [4, 3, 65] it is proved that $p_c = p_F(N)$ holds for $a_0 \geq \frac{5}{3}$ ($N = 1$), 3 ($N = 2$), $N + 2$ ($N \geq 3$). On the other hand, when $a_0 = 2$, D'Abbico, Lucente and Reissig [7] showed $p_c = \max\{p_F(N), p_S(N + 2)\}$ for $N \leq 3$ (see [5, 49] for higher dimensional cases). This implies that the critical exponent depends on a_0 . For $a_0 \neq 2$, the best known result is by Ikeda and Sobajima [17]. They obtained the small data blow-up for $N \geq 1, 0 < a_0 < \frac{N^2+N+2}{N+2}$ and $1 < p \leq p_S(N + a_0)$ (see also [32, 61]). Also, by [20] it is proved that the critical exponent is given by $p_c = 1$ when $\beta < -1$, namely, the small data global existence holds for any $p > 1$.

On the other hand, for the space-dependent damping case, namely, $\beta = 0$, Ikehata, Todorova and Yordanov [24] proved if $0 \leq \alpha < 1$, then the critical exponent is given by $p_c = 1 + \frac{2}{N-\alpha}$. However, there is no result in the case $\alpha < 0$, where the damping coefficient is unbounded with respect to the space variable, while we can expect that the critical exponent is still given by $p_c = 1 + \frac{2}{N-\alpha}$ in view of the result of linear problem [57]. When $\alpha = 1$, the equation has scale-invariance and the critical exponent seems to change depending on a_0 . Indeed, Ikeda and Sobajima [16] proved the small data blow-up for $N \geq 3, 0 \leq a_0 < \frac{(N-1)^2}{N+1}, \frac{N}{N-1} < p \leq p_S(N + a_0)$, while the global existence part remains open.

In contrast, there were only few results on time and space dependent cases. By [64], the small data global existence was proved for $\alpha, \beta \geq 0, \alpha + \beta < 1$ and $p > 1 + \frac{2}{N-\alpha}$. Khader [29] also proved the small data global existence for $0 \leq \alpha < 1, -1 < \beta < 1, 0 < \alpha + \beta < 1$, and $p > 1 + \frac{4(\beta+1)}{2(N-\alpha)(\beta+1)-\beta(2-\alpha)}$. However, there are no results on the small data blow-up for subcritical or critical case.

We also refer the reader to [34, 47, 19, 33, 14, 15, 62] for studies on estimates of lifespan of blow-up solutions.

Summarizing the previous studies, we can conjecture the following for the critical exponent of the problem (1.1) with the condition (1.2).

Conjecture

- (i) For $\alpha < 2, \beta > -1$ with $\alpha + \beta < 1$, the critical exponent is $p_c = 1 + \frac{2}{N-\alpha}$.
- (ii) For $\alpha, \beta \in \mathbb{R}$ with $\alpha + \beta = 1$, the equation has scale-invariance and the critical

exponent will depend on a_0 .

(iii) For $\alpha, \beta \in \mathbb{R}$ with $\alpha + \beta > 1$, the critical exponent is given by the Strauss number $p_c = p_S(N)$.

In this paper, we present some partial answers which support the Conjecture (i). In particular, we completely give the critical exponent in the case $\alpha < 0, \beta = 0$.

Before going to our main results, we mention the existence of the local solution.

Proposition 1.1 (Existence of the local solution). *Let $N \geq 1$ and let $c(t, x)$ has the form (1.2) with some $a_0 > 0, \alpha, \beta \in \mathbb{R}$. We assume that the initial data satisfy (1.4) with some $R_0 > 0$. If p satisfies*

$$(1.6) \quad 1 < p < \infty \quad (N = 1, 2), \quad 1 < p \leq \frac{N}{N-2} \quad (N \geq 3),$$

then for any $\varepsilon > 0$, there exists a time $T > 0$ such that the Cauchy problem (1.1) admits a unique solution

$$(1.7) \quad u \in C([0, T]; H^1(\mathbb{R}^N)) \cap C^1([0, T]; L^2(\mathbb{R}^N))$$

satisfying

$$(1.8) \quad \text{supp } u(t, \cdot) \subset \{x \in \mathbb{R}^N; |x| \leq R_0 + t\}.$$

Moreover, with the notion of the lifespan

$$(1.9) \quad T(\varepsilon) := \{T \in (0, \infty]; \text{ there exists a unique solution } u \text{ in the class (1.7)}\},$$

we have the following blow-up alternative: if $T(\varepsilon) < \infty$, then

$$(1.10) \quad \lim_{t \rightarrow T(\varepsilon) - 0} \|(u, u_t)(t)\|_{H^1 \times L^2} = \infty$$

holds.

The proof is given by a standard energy estimate and the contraction mapping principle (see [23, Proposition 2.1]).

Our first main results are the small data global existence in the supercritical case.

Theorem 1.2 (Small data global existence for $\beta \in (-1, 1)$ in the supercritical case). *Let $N \geq 1$ and let $c(t, x)$ has the form (1.2) with some $a_0 > 0, \alpha < 0$ and $\beta \in (-1, 1)$. We assume that the initial data satisfy (1.4) with some $R_0 > 0$. If p satisfies*

$$(1.11) \quad 1 + \frac{2}{N-\alpha} < p < \infty \quad (N = 1, 2), \quad 1 + \frac{2}{N-\alpha} < p \leq \frac{N}{N-2} \quad (N \geq 3),$$

then there exists a constant $\varepsilon_0 > 0$ depending on $N, p, a_0, \alpha, \beta, u_0, u_1, R_0$ such that for any $\varepsilon \in (0, \varepsilon_0]$, the Cauchy problem (1.1) admits a unique global solution

$$u \in C([0, \infty); H^1(\mathbb{R}^N)) \cap C^1([0, \infty); L^2(\mathbb{R}^N)).$$

Theorem 1.3 (Small data global existence for $\beta = 1$ in the supercritical case). *Let $N \geq 1$ and let $c(t, x)$ has the form (1.2) with some $a_0 > 0, \alpha < 0$ and $\beta = 1$. We assume that the initial data satisfy (1.4) with some $R_0 > 0$. If p satisfies (1.11), then there exist constants $a_* > 0$ depending on p and $\varepsilon_0 > 0$ depending on $N, p, a_0, \alpha, u_0, u_1, R_0$ such that if $a_0 \geq a_*$ and $\varepsilon \in (0, \varepsilon_0]$, then the Cauchy problem (1.1) admits a unique global solution*

$$u \in C([0, \infty); H^1(\mathbb{R}^N)) \cap C^1([0, \infty); L^2(\mathbb{R}^N)).$$

The second main result is the finite time blow-up of the solution in the subcritical and the critical cases when α and β satisfy (1.3). Moreover, we give the sharp upper estimates of the lifespan.

Theorem 1.4 (Blow-up in the subcritical or the critical case). *Let $N \geq 1$, and let $c(t, x)$ has the form (1.2) with some $a_0 > 0$, α and β satisfying (1.3). Moreover, we assume that p satisfies*

$$(1.12) \quad 1 < p \leq 1 + \frac{2}{N - \alpha},$$

and the initial data satisfy (1.4) with some $R_0 > 0$ and

$$(1.13) \quad \int_{\mathbb{R}^N} (u_1(x) + (a_0 \langle x \rangle^{-\alpha} - \beta)u_0(x)) dx > 0.$$

Then, there exist constants $\varepsilon_1 > 0$ and $C > 0$ depending on $N, p, a_0, \alpha, u_0, u_1, R_0$ such that for any $\varepsilon \in (0, \varepsilon_1]$, the lifespan of the local solution is estimated as

$$(1.14) \quad T(\varepsilon) \leq \begin{cases} C\varepsilon^{-\frac{2-\alpha}{2(1+\beta)}(\frac{1}{p-1} - \frac{N-\alpha}{2})^{-1}} & 1 < p < 1 + \frac{2}{N-\alpha}, \\ \exp(C\varepsilon^{-(p-1)}) & p = 1 + \frac{2}{N-\alpha}. \end{cases}$$

Remark 1.1. (i) By Theorems 1.2, and 1.4 we conclude that if the damping term is given by (1.2) with some $a_0 > 0$, α, β satisfying $\alpha < 0, \beta = 0$, then the critical exponent of the Cauchy problem (1.1) is determined by

$$p_c = 1 + \frac{2}{N - \alpha}.$$

When $\alpha < 0, \beta = 1$, by Theorems 1.3 and 1.4, the critical exponent is almost determined as the above one, in the sense that if $1 < p \leq 1 + \frac{2}{N-\alpha}$, then the small data blow-up occurs (for all $a_0 > 0$), and if $p > 1 + \frac{2}{N-\alpha}$, then the small data global existence holds provided that a_0 is sufficiently large (depending on p).

(ii) When the damping is effective, namely, $\alpha + \beta < 1$, if we do not impose the condition (1.3), the blow-up of solutions in the subcritical or the critical case $1 < p \leq 1 + \frac{2}{N-\alpha}$ is still an open problem.

We shall comment on the method of proof and construction of the paper. Theorem 1.2 is proved in Section 2 by weighted energy method with a weight function having the form $e^{\psi(t,x)}$ with an appropriate function $\psi(t, x)$ (see Definition 2.1). Such a weight function was developed by [21, 60, 48, 56]. Making use of this weight, we can estimate the weighted energy of the solution by the sum of the initial energy and the nonlinear terms (see Lemma 2.4). Then, to control the nonlinear terms, we apply the Caffarelli–Kohn–Nirenberg inequality, which is suitable with the energy including polynomially increasing coefficient $a(x)$.

Theorem 1.3 can be also proved in the same strategy. To avoid proceeding the similar computations as before, we emphasize the difference from the proof of Theorem 1.2 by giving only the outline of proof in Appendix B.

For Theorem 1.4, in Section 3, we apply the so-called test function method developed by [76, 37, 4, 15]. Multiplying the equation by $(1+t)$, we can transform the linear part of the equation (1.1) into divergence form. This is a simple but crucial idea in the proof. Then, we further multiply a test function scaled by a large parameter $R \in (0, T(\varepsilon))$, and apply the integration by parts, which gives a certain estimate including the parameter R , the initial data, and the nonlinear term. Finally, letting R to $T(\varepsilon)$, we have the estimate of the lifespan.

2. SMALL DATA GLOBAL EXISTENCE IN THE SUPERCRITICAL CASE

To keep the paper readable length, we will give the detailed proof only for the case $\alpha < 0$ and $\beta = 0$, namely, $c(t, x) = a(x)$, and for the other cases we will give an outline of the proof in Appendix B.

2.1. Construction of a weight function. We first prepare a suitable weight function, which will be used for weighted energy estimates of the solution. In the previous works [21, 24, 48, 37, 64] for the case $\alpha \geq 0$, the so-called Ikehata-Todorova-Yordanov type weight function

$$(2.1) \quad e^{\psi(t,x)} \quad \text{with} \quad \psi(t,x) = \mu \frac{\langle x \rangle^{2-\alpha}}{1+t}, \quad \mu > 0$$

was used. The function $\psi(t, x)$ has the properties

$$(2.2) \quad -\psi_t(t, x)a(x) = (2 + \delta)|\nabla\psi(t, x)|^2,$$

$$(2.3) \quad \Delta\psi(t, x) \geq \left(\frac{N - \alpha}{2(2 - \alpha)} - \delta \right) \frac{a(x)}{1+t}$$

with some small $\delta > 0$, and these properties are essential for the weighted energy estimates. However, if $\alpha < 0$, then the above weight function $\psi(t, x)$ does not satisfy the estimate (2.3).

Therefore, following the idea of [56], we modify the weight function (2.1) as follows.

Definition 2.1. Let $\delta \in (0, 1)$, $\mu = \frac{a_0}{(2-\alpha)^2(2+\delta)}$, and $A_0 > 0$. Let $R_\delta > 0$ be a sufficiently large constant depending on $\delta > 0$, and we take a cut-off function $\eta_{R_\delta} \in C_0^\infty(\mathbb{R}^N)$ such that

$$\eta_{R_\delta}(x) = 1 \quad (|x| \leq R_\delta), \quad \eta_{R_\delta}(x) = 0 \quad (|x| \geq 2R_\delta).$$

Let \mathcal{N} be the Newton potential, that is,

$$\mathcal{N}(x) = \begin{cases} \frac{|x|}{2} & (N = 1), \\ \frac{1}{2\pi} \log \frac{1}{|x|} & (N = 2), \\ \frac{\Gamma(\frac{N}{2} + 1)}{N(N-2)\pi^{N/2}} |x|^{2-N} & (N \geq 3). \end{cases}$$

We define

$$\psi(t, x) = \frac{\mu}{1+t} \left\{ \langle x \rangle^{2-\alpha} + A_0 - \mathcal{N} * (\alpha(2-\alpha)\langle x \rangle^{-2-\alpha}\eta_{R_\delta}(x)) \right\}.$$

Lemma 2.2. For any $\delta \in (0, 1)$, there exist constants $R_\delta > 0$ and $A_0 > 0$ such that the function $\psi(t, x)$ defined by Definition 2.1 satisfies

$$(2.4) \quad -\psi_t(t, x)a(x) \geq (2 + \delta_1)|\nabla\psi(t, x)|^2,$$

$$(2.5) \quad \Delta\psi(t, x) \geq \left(\frac{N - \alpha}{2(2 - \alpha)} - \delta_2 \right) \frac{a(x)}{1+t},$$

where $\delta_1 = \frac{2}{3}\delta$ and $\delta_2 = \frac{N-\alpha}{2(2-\alpha)}\delta$.

Proof. We calculate

$$\begin{aligned}\Delta\psi(t, x) &= \mu(N - \alpha)(2 - \alpha)\frac{\langle x \rangle^{-\alpha}}{1 + t} + \mu\alpha(2 - \alpha)\frac{\langle x \rangle^{-2-\alpha}}{1 + t}(1 - \eta_{R_\delta}(x)) \\ &\geq \left(\frac{N - \alpha}{2(2 - \alpha)} - \delta_2\right)\frac{a(x)}{1 + t}\end{aligned}$$

for sufficiently large $R_\delta > 0$. Thus, we have (2.5). Next, we prove (2.4). We compute

$$\begin{aligned}& -\psi_t(t, x)a(x) \\ &= \frac{1}{1 + t}\psi(t, x)a(x) \\ &= \frac{a_0}{(2 - \alpha)^2(2 + \delta)}\frac{a(x)}{(1 + t)^2}\left\{\langle x \rangle^{2-\alpha} + A_0 - \mathcal{N} * (\alpha(2 - \alpha)\langle x \rangle^{-2-\alpha}\eta_{R_\delta}(x))\right\}.\end{aligned}$$

On the other hand, we have

$$\begin{aligned}& (2 + \delta_1)|\nabla\psi(t, x)|^2 \\ &= (2 + \delta_1)\frac{a_0^2}{(2 - \alpha)^2(2 + \delta)^2}\frac{\langle x \rangle^{-2\alpha}|x|^2}{(1 + t)^2} \\ &\quad - 2(2 + \delta_1)\frac{a_0^2}{(2 - \alpha)^3(2 + \delta)^2}\frac{\langle x \rangle^{-\alpha}x}{(1 + t)^2} \cdot \nabla[\mathcal{N} * (\alpha(2 - \alpha)\langle x \rangle^{-2-\alpha}\eta_{R_\delta}(x))] \\ &\quad + (2 + \delta_1)\frac{a_0^2}{(2 - \alpha)^4(2 + \delta)^2}\frac{1}{(1 + t)^2}|\nabla\mathcal{N} * (\alpha(2 - \alpha)\langle x \rangle^{-2-\alpha}\eta_{R_\delta}(x))|^2.\end{aligned}$$

We easily obtain

$$\begin{aligned}|\mathcal{N} * (\alpha(2 - \alpha)\langle x \rangle^{-2-\alpha}\eta_{R_\delta}(x))| &\leq C \begin{cases} \langle x \rangle^{2-N} & (N = 1, N \geq 3), \\ \log(2 + |x|) & (N = 2), \end{cases} \\ |\nabla\mathcal{N} * (\alpha(2 - \alpha)\langle x \rangle^{-2-\alpha}\eta_{R_\delta}(x))| &\leq C\langle x \rangle^{1-N}.\end{aligned}$$

Now, we take a constant $\delta_3 > 0$ so that $1 - \delta_3 \geq \frac{2+\delta_1}{2+\delta}$ holds. Then, we have

$$(1 - \delta_3)\frac{a_0}{(2 - \alpha)^2(2 + \delta)}\frac{\langle x \rangle^{2-\alpha}a(x)}{(1 + t)^2} \geq (2 + \delta_1)\frac{a_0^2}{(2 - \alpha)^2(2 + \delta)^2}\frac{\langle x \rangle^{-2\alpha}|x|^2}{(1 + t)^2}.$$

Finally, we take a sufficiently large constant $A_0 > 0$ so that

$$\begin{aligned}& \frac{a_0}{(2 - \alpha)^2(2 + \delta)}\frac{\delta_3\langle x \rangle^{2-\alpha}a(x) + A_0a(x)}{(1 + t)^2} \\ &\geq \frac{a_0}{(2 - \alpha)^2(2 + \delta)}\frac{a(x)}{(1 + t)^2}|\mathcal{N} * (\alpha(2 - \alpha)\eta_{R_\delta}(x)\langle x \rangle^{-2-\alpha})| \\ &\quad + \left|2(2 + \delta_1)\frac{a_0^2}{(2 - \alpha)^3(2 + \delta)^2}\frac{\langle x \rangle^{-\alpha}x}{(1 + t)^2} \cdot \nabla[\mathcal{N} * (\alpha(2 - \alpha)\eta_{R_\delta}(x)\langle x \rangle^{-2-\alpha})]\right| \\ &\quad + (2 + \delta_1)\frac{a_0^2}{(2 - \alpha)^4(2 + \delta)^2}\frac{1}{(1 + t)^2}|\nabla\mathcal{N} * (\alpha(2 - \alpha)\eta_{R_\delta}(x)\langle x \rangle^{-2-\alpha})|^2.\end{aligned}$$

Consequently, we have (2.4). \square

2.2. Weighted energy estimates. Using the function $\psi(t, x)$ constructed in Definition 2.1, we prove the following weighted energy estimates for solutions to (1.1). Continuing from the previous subsection, we consider the case $\alpha < 0$ and $\beta = 0$. Let $\delta_0 \in (0, \frac{N-\alpha}{2-\alpha})$. We define

$$(2.6) \quad M(t) := \sup_{0 \leq \tau < t} \left\{ (1 + \tau)^{\frac{N-\alpha}{2-\alpha} + 1 - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} (u_t^2 + |\nabla u|^2) dx \right. \\ \left. + (1 + \tau)^{\frac{N-\alpha}{2-\alpha} - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} a(x) u^2 dx \right\}.$$

By the blow-up alternative in Proposition 1.1, Theorem 1.2 is obtained from the following a priori estimate.

Proposition 2.3. *Under the assumptions of Theorem 1.2, there exist $\delta_0 \in (0, \frac{N-\alpha}{2-\alpha})$, and constants $t_0 = t_0(a_0, R_0, \delta_0) \geq 1$, $C = C(N, \alpha, a_0, p, R_0, \delta_0, t_0) > 0$ such that the solution u constructed in Proposition 1.1 satisfies the a priori estimate*

$$M(t) \leq CM(0) + CM(t)^{(p+1)/2}$$

for $t \in [0, T(\varepsilon))$, where the function $\psi(t, x)$ is defined in Definition 2.1 with $\delta = \frac{2-\alpha}{2(N-\alpha)}\delta_0$.

Remark 2.1. *The restriction $p \leq \frac{N}{N-2}$ ($N \geq 3$) is due to the local existence (Proposition 1.1), and we can obtain the above a priori estimate for $1 + \frac{2}{N-\alpha} < p \leq \frac{N+2}{N-2}$.*

To prove Proposition 2.3, we first prove the following energy estimate. After that, in the next subsection, we give the nonlinear estimates and complete the proof of Proposition 2.3.

Lemma 2.4. *Under the assumptions of Proposition 2.3, for any $\delta_0 \in (0, \frac{N-\alpha}{2-\alpha})$, there exist constants $t_0 = t_0(N, \alpha, a_0, R_0, \delta_0) \geq 1$, and $C = C(N, \alpha, a_0, p, R_0, \delta_0, t_0) > 0$ such that the solution u constructed in Proposition 1.1 satisfies*

$$(t_0 + t)^{\frac{N-\alpha}{2-\alpha} + 1 - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} (u_t^2 + |\nabla u|^2) dx + (t_0 + t)^{\frac{N-\alpha}{2-\alpha} - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} a(x) u^2 dx \\ \leq C \int_{\mathbb{R}^N} (u_1^2 + |\nabla u_0|^2 + u_0^2 + |u_0|^{p+1}) dx \\ + C(t_0 + t)^{\frac{N-\alpha}{2-\alpha} + 1 - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} |F(u)| dx \\ + C \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha} - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} |F(u)| dx d\tau \\ + C \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha} + 1 - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} (-\psi_t) |F(u)| dx d\tau,$$

where $F(u) = \int_0^u |v|^p dv$, and the function $\psi(t, x)$ is defined in Definition 2.1 with $\delta = \frac{2-\alpha}{2(N-\alpha)}\delta_0$.

Proof. Multiplying the equation of (1.1) by $e^{2\psi}u_t$, we have

$$\begin{aligned} & \frac{\partial}{\partial t} \left[\frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) \right] - \nabla \cdot (e^{2\psi} u_t \nabla u) \\ & + e^{2\psi} \left(a(x) - \frac{|\nabla \psi|^2}{-\psi_t} - \psi_t \right) u_t^2 + \frac{e^{2\psi}}{-\psi_t} |\psi_t \nabla u - u_t \nabla \psi|^2 \\ & = \frac{\partial}{\partial t} [e^{2\psi} F(u)] + 2e^{2\psi} (-\psi_t) F(u). \end{aligned}$$

By (2.4), the last term of the left-hand side satisfies

$$\frac{e^{2\psi}}{-\psi_t} |\psi_t \nabla u - u_t \nabla \psi|^2 \geq e^{2\psi} \left(\frac{1}{5} (-\psi_t) |\nabla u|^2 - \frac{a(x)}{4(2+\delta_1)} u_t^2 \right).$$

This and using again (2.4) imply

$$\begin{aligned} (2.7) \quad & \frac{\partial}{\partial t} \left[\frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) \right] - \nabla \cdot (e^{2\psi} u_t \nabla u) \\ & + e^{2\psi} \left\{ \left(\frac{a(x)}{4} - \psi_t \right) u_t^2 + \frac{-\psi_t}{5} |\nabla u|^2 \right\} \\ & \leq \frac{\partial}{\partial t} [e^{2\psi} F(u)] + 2e^{2\psi} (-\psi_t) F(u). \end{aligned}$$

On the other hand, multiplying the equation of (1.1) by $e^{2\psi}u$, we have

$$\begin{aligned} & \frac{\partial}{\partial t} \left[e^{2\psi} \left(uu_t + \frac{a(x)}{2} u^2 \right) \right] - \nabla \cdot (e^{2\psi} u \nabla u) \\ & + e^{2\psi} \{ |\nabla u|^2 - \psi_t a(x) u^2 + 2u \nabla \psi \cdot \nabla u - 2\psi_t uu_t - u_t^2 \} \\ & = e^{2\psi} |u|^p u. \end{aligned}$$

By noting

$$2e^{2\psi} u \nabla \psi \cdot \nabla u = 4e^{2\psi} u \nabla \psi \cdot \nabla u - \nabla \cdot (e^{2\psi} u^2 \nabla \psi) + 2e^{2\psi} |\nabla \psi|^2 u^2 + e^{2\psi} (\Delta \psi) u^2,$$

we see that

$$\begin{aligned} (2.8) \quad & \frac{\partial}{\partial t} \left[e^{2\psi} \left(uu_t + \frac{a(x)}{2} u^2 \right) \right] - \Delta \left(\frac{e^{2\psi}}{2} u^2 \right) \\ & + e^{2\psi} \{ |\nabla u|^2 + 4u \nabla \psi \cdot \nabla u + ((-\psi_t) a(x) + 2|\nabla \psi|^2) u^2 \} \\ & + e^{2\psi} (\Delta \psi) u^2 - e^{2\psi} (2\psi_t uu_t + u_t^2) \\ & = e^{2\psi} |u|^p u. \end{aligned}$$

By (2.4) and the Schwarz inequality, we estimate

$$\begin{aligned} & |\nabla u|^2 + 4u \nabla \psi \cdot \nabla u + ((-\psi_t) a(x) + 2|\nabla \psi|^2) u^2 \\ & \geq |\nabla u|^2 + 4u \nabla \psi \cdot \nabla u + (4 + \delta_4) |\nabla \psi|^2 u^2 + \frac{\delta}{4} (-\psi_t) a(x) u^2 \\ & \geq \delta_5 |\nabla u|^2 + \frac{\delta}{4} (-\psi_t) a(x) u^2, \end{aligned}$$

where $\delta_4 = \delta_1 - \frac{\delta}{2} - \frac{\delta\delta_1}{4}$ and $\delta_5 = \frac{\delta_4}{4+\delta_4}$. Here, we remark that $\delta \in (0, 1)$ and $\delta_1 = \frac{2}{3}\delta$ ensure $\delta_4 > 0$. Also, (2.5) implies

$$e^{2\psi}(\Delta\psi)u^2 \geq \left(\frac{N-\alpha}{2(2-\alpha)} - \delta_2\right) e^{2\psi} \frac{a(x)}{1+t} u^2.$$

Moreover, it follows from the Schwarz inequality that

$$|2\psi_t uu_t| \leq \frac{8}{\delta a(x)} (-\psi_t) u_t^2 + \frac{\delta}{8} a(x) (-\psi_t) u^2.$$

Plugging these into (2.8), we conclude

$$\begin{aligned} & \frac{\partial}{\partial t} \left[e^{2\psi} \left(uu_t + \frac{a(x)}{2} u^2 \right) \right] - \Delta \left(\frac{e^{2\psi}}{2} u^2 \right) \\ & + e^{2\psi} \left(\delta_5 |\nabla u|^2 + \frac{\delta}{8} (-\psi_t) a(x) u^2 \right) + \left(\frac{N-\alpha}{2(2-\alpha)} - \delta_2 \right) e^{2\psi} \frac{a(x)}{1+t} u^2 \\ & - e^{2\psi} \left(1 + \frac{8(-\psi_t)}{\delta a(x)} \right) u_t^2 \\ & \leq e^{2\psi} |u|^p u. \end{aligned}$$

Integrating the above over \mathbb{R}^N and multiplying it by $(t_0 + t)^{\frac{N-\alpha}{2-\alpha} - \delta_0}$, we have

$$\begin{aligned} (2.9) \quad & \frac{d}{dt} \left[(t_0 + t)^{\frac{N-\alpha}{2-\alpha} - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(uu_t + \frac{a(x)}{2} u^2 \right) dx \right] \\ & - \left(\frac{N-\alpha}{2-\alpha} - \delta_0 \right) (t_0 + t)^{\frac{N-\alpha}{2-\alpha} - 1 - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(uu_t + \frac{a(x)}{2} u^2 \right) dx \\ & + (t_0 + t)^{\frac{N-\alpha}{2-\alpha} - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(\delta_5 |\nabla u|^2 + \frac{\delta}{8} (-\psi_t) a(x) u^2 \right) dx \\ & + \left(\frac{N-\alpha}{2(2-\alpha)} - \delta_2 \right) (t_0 + t)^{\frac{N-\alpha}{2-\alpha} - 1 - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} a(x) u^2 dx \\ & - (t_0 + t)^{\frac{N-\alpha}{2-\alpha} - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(1 + \frac{8(-\psi_t)}{\delta a(x)} \right) u_t^2 dx \\ & \leq (t_0 + t)^{\frac{N-\alpha}{2-\alpha} - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} |u|^p u dx. \end{aligned}$$

Since $\delta = \frac{2-\alpha}{2(N-\alpha)}\delta_0$ and $\delta_2 = \frac{N-\alpha}{2(2-\alpha)}\delta$, we see that $4\delta_2 = \delta_0$ holds and hence, we compute

$$\begin{aligned} & - \left(\frac{N-\alpha}{2-\alpha} - \delta_0 \right) (t_0 + t)^{\frac{N-\alpha}{2-\alpha} - 1 - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} \frac{a(x)}{2} u^2 dx \\ & + \left(\frac{N-\alpha}{2(2-\alpha)} - \delta_2 \right) (t_0 + t)^{\frac{N-\alpha}{2-\alpha} - 1 - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} a(x) u^2 dx \\ & = \delta_2 (t_0 + t)^{\frac{N-\alpha}{2-\alpha} - 1 - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} a(x) u^2 dx. \end{aligned}$$

Moreover, the Schawarz inequality implies

$$\begin{aligned}
& \left(\frac{N-\alpha}{2-\alpha} - \delta_0 \right) (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} u u_t \, dx \\
& \leq \delta_6 (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} a(x) u^2 \, dx + C(\delta_6) (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \frac{1}{a(x)} u_t^2 \, dx \\
& \leq \delta_6 (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} a(x) u^2 \, dx + \frac{C(\delta_6)}{t_0} (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \frac{1}{a(x)} u_t^2 \, dx
\end{aligned}$$

where $\delta_6 = \frac{\delta_2}{2}$ and $C(\delta_6) = \frac{1}{4\delta_6} \left(\frac{N-\alpha}{2-\alpha} - \delta_0 \right)^2$. Applying these estimates to (2.9), we deduce

$$\begin{aligned}
& \frac{d}{dt} \left[(t_0 + t)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(u u_t + \frac{a(x)}{2} u^2 \right) \, dx \right] \\
& + (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(\delta_5 |\nabla u|^2 + \frac{\delta}{8} (-\psi_t) a(x) u^2 \right) \, dx \\
& + \delta_6 (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} a(x) u^2 \, dx \\
& - (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(1 + \frac{C(\delta_6)}{t_0 a(x)} + \frac{8(-\psi_t)}{\delta a(x)} \right) u_t^2 \, dx \\
& \leq (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} |u|^p u \, dx.
\end{aligned}$$

Integrating it over $[0, t]$, we conclude

$$\begin{aligned}
(2.10) \quad & (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(u u_t + \frac{a(x)}{2} u^2 \right) \, dx \\
& + \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(\delta_5 |\nabla u|^2 + \frac{\delta}{8} (-\psi_t) a(x) u^2 \right) \, dx d\tau \\
& + \delta_6 \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha}-1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} a(x) u^2 \, dx d\tau \\
& - \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(1 + \frac{C(\delta_6)}{t_0 a(x)} + \frac{8(-\psi_t)}{\delta a(x)} \right) u_t^2 \, dx d\tau \\
& \leq C(t_0) \int_{\mathbb{R}^N} e^{2\psi(0,x)} (u_0 u_1 + \frac{a(x)}{2} u_0^2) \, dx + \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} |u|^p u \, dx d\tau.
\end{aligned}$$

On the other hand, integrating (2.7) over \mathbb{R}^N , we have

$$\begin{aligned}
& \frac{d}{dt} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) \, dx \\
& + \int_{\mathbb{R}^N} e^{2\psi} \left\{ \left(\frac{a(x)}{4} - \psi_t \right) u_t^2 + \frac{-\psi_t}{5} |\nabla u|^2 \right\} \, dx \\
& \leq \frac{d}{dt} \int_{\mathbb{R}^N} e^{2\psi} F(u) \, dx + 2 \int_{\mathbb{R}^N} e^{2\psi} (-\psi_t) F(u) \, dx.
\end{aligned}$$

We multiply it by $(t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0}$ to obtain

$$\begin{aligned}
& \frac{d}{dt} \left[(t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) dx \right] \\
& - \left\{ \frac{N-\alpha}{2-\alpha} + 1 - \delta_0 \right\} (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) dx \\
& + (t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left\{ \left(\frac{a(x)}{4} - \psi_t \right) u_t^2 + \frac{-\psi_t}{5} |\nabla u|^2 \right\} dx \\
& \leq \frac{d}{dt} \left[(t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} F(u) dx \right] \\
& - \left\{ \frac{N-\alpha}{2-\alpha} + 1 - \delta_0 \right\} (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} F(u) dx \\
& + 2(t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} (-\psi_t) F(u) dx.
\end{aligned}$$

Noting that

$$\left\{ \frac{N-\alpha}{2-\alpha} + 1 - \delta_0 \right\} (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \leq (t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \frac{a(x)}{8}$$

holds if we choose $t_0 \geq 1$ so that $t_0 \geq \frac{8}{a_0} \left(\frac{N-\alpha}{2-\alpha} + 1 - \delta_0 \right)$, we calculate

$$\begin{aligned}
& \frac{d}{dt} \left[(t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) dx \right] \\
& + (t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left\{ \left(\frac{a(x)}{8} - \psi_t \right) u_t^2 + \frac{-\psi_t}{5} |\nabla u|^2 \right\} dx \\
& - \left\{ \frac{N-\alpha}{2-\alpha} + 1 - \delta_0 \right\} (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} |\nabla u|^2 dx \\
& \leq \frac{d}{dt} \left[(t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} F(u) dx \right] \\
& - \left\{ \frac{N-\alpha}{2-\alpha} + 1 - \delta_0 \right\} (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} F(u) dx \\
& + 2(t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} (-\psi_t) F(u) dx.
\end{aligned}$$

Integrating it over $[0, t]$, we conclude

$$\begin{aligned}
(2.11) \quad & (t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) dx \\
& + \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left\{ \left(\frac{a(x)}{8} - \psi_t \right) u_t^2 + \frac{-\psi_t}{5} |\nabla u|^2 \right\} dx d\tau \\
& - \left\{ \frac{N-\alpha}{2-\alpha} + 1 - \delta_0 \right\} \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} |\nabla u|^2 dx d\tau \\
& \leq C(t_0) \int_{\mathbb{R}^N} e^{2\psi(0,x)} (u_1^2 + |\nabla u_0|^2 + |u_0|^{p+1}) dx \\
& + (t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} F(u) dx \\
& - \left\{ \frac{N-\alpha}{2-\alpha} + 1 - \delta_0 \right\} \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} F(u) dx d\tau \\
& + 2 \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} (-\psi_t) F(u) dx d\tau.
\end{aligned}$$

Finally, we combine (2.10) and (2.11). We take $\nu > 0$ so that

$$\nu \left\{ \frac{N-\alpha}{2-\alpha} + 1 - \delta_0 \right\} \leq \frac{\delta_5}{2}$$

holds and, we choose t_0 sufficiently large so that

$$\begin{aligned}
\frac{1}{t_0} \left(1 + \frac{C(\delta_6)}{t_0} \right) & \leq \frac{\nu}{16} a(x), \\
\frac{8}{\delta t_0 a(x)} & \leq \frac{\nu}{2},
\end{aligned}$$

Computing (2.10) $+\nu \cdot$ (2.11), we conclude

$$\begin{aligned}
& \nu(t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) dx + (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(uu_t + \frac{a(x)}{2} u^2 \right) dx \\
& + \nu \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left\{ \left(\frac{a(x)}{16} + \frac{-\psi_t}{2} \right) u_t^2 + \frac{-\psi_t}{5} |\nabla u|^2 \right\} dx d\tau \\
& + \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(\frac{\delta_5}{2} |\nabla u|^2 + \frac{\delta}{8} (-\psi_t) a(x) u^2 \right) dx d\tau \\
& + \delta_6 \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha}-1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} a(x) u^2 dx d\tau \\
& \leq C(t_0) \int_{\mathbb{R}^N} e^{2\psi(0,x)} (u_1^2 + |\nabla u_0|^2 + |u_0|^{p+1} + a(x) u_0^2) dx \\
& + \nu(t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} |F(u)| dx \\
& + C(N, \alpha, \delta_0, \nu) \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} |F(u)| dx d\tau \\
& + C(\nu) \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} (-\psi_t) |F(u)| dx d\tau.
\end{aligned}$$

Finally, noting that

$$\begin{aligned} & \nu(t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) dx + (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(uu_t + \frac{a(x)}{2} u^2 \right) dx \\ & \geq \frac{\nu}{2} (t_0 + t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) dx + \frac{1}{4} (t_0 + t)^{\frac{N-\alpha}{2-\alpha}-\delta_0} \int_{\mathbb{R}^N} e^{2\psi} a(x) u^2 dx \end{aligned}$$

if $t_0 \geq \frac{2}{\nu a_0}$, we have the assertion of Lemma 2.4. \square

Remark 2.2. *In the above proof, we have determined the positive constants δ, δ_j ($j = 1, \dots, 6$), ν, t_0 in the following way. First, for given $\delta_0 \in (0, \frac{N-\alpha}{2-\alpha})$, we define $\delta := \frac{2-\alpha}{2(N-\alpha)} \delta_0$. Then, we choose δ_j ($j = 1, \dots, 6$) as $\delta_1 := \frac{2}{3} \delta$, $\delta_2 := \frac{N-\alpha}{2(2-\alpha)} \delta$, δ_3 such that $1 - \delta_3 \geq \frac{2+\delta_1}{2+\delta}$, $\delta_4 := \delta_1 - \frac{\delta}{2} - \frac{\delta \delta_1}{4}$, $\delta_5 := \frac{\delta_4}{4+\delta_4}$, $\delta_6 := \frac{\delta_2}{2}$. Note that $\delta = O(\delta_0)$ and $\delta_j = O(\delta_0)$ ($j = 1, \dots, 6$) as $\delta_0 \rightarrow 0+$. After that, we take ν as $\nu(\frac{N-\alpha}{2-\alpha} + 1 - \delta_0) \leq \frac{\delta_5}{2}$. Then, we choose $t_0 \geq 1$ so that*

$$\begin{aligned} t_0 & \geq \frac{8}{a_0} \left(\frac{N-\alpha}{2-\alpha} + 1 - \delta_0 \right), \quad \frac{1}{t_0} \left(1 + \frac{C(\delta_6)}{t_0} \right) \leq \frac{\nu}{16} a(x), \\ \frac{8}{\delta t_0 a(x)} & \leq \frac{\nu}{2}, \quad t_0 \geq \frac{2}{\nu a_0} \end{aligned}$$

hold for any $x \in \mathbb{R}^N$. These observations will be useful when we discuss the case $\alpha < 0, \beta = 1$ (see Appendix B).

Finally, we also remark that δ_0 will be determined depending on p in the nonlinear estimates discussed in the next subsection.

2.3. Nonlinear estimates and proof of Proposition 2.3. In this subsection, we give the nonlinear estimates for the right-hand side of Lemma 2.4 and complete the proof of Proposition 2.3. We first recall the following special case of Gagliardo–Nirenberg inequality.

Lemma 2.5. (Gagliardo–Nirenberg inequality, see [10, Section 6.1.1]) *For $1 \leq p < \infty$ ($N = 1, 2$), $1 \leq p \leq \frac{N+2}{N-2}$ ($N \geq 3$), there exists a constant $C > 0$ such that for any $u \in H^1(\mathbb{R}^N)$, we have*

$$\|u\|_{L^{p+1}} \leq C \|\nabla u\|_{L^2}^\theta \|u\|_{L^2}^{1-\theta},$$

where $\theta = \frac{N(p-1)}{2(p+1)} \in [0, 1]$.

Besides the above lemma, we also use the following special case of Caffarelli–Kohn–Nirenberg inequality.

Lemma 2.6. (Caffarelli–Kohn–Nirenberg inequality, see [1]) *For $k = 0, 1, 2, \dots$, there exist $C(k, N) > 0$ such that for $u \in H^1(\mathbb{R}^N)$ with compact support, we have*

$$\|u\|_{L^2} \leq C(k, N) \|\nabla u\|_{L^2}^{1-1/2^k} \| |x|^{2^k-1} u \|_{L^2}^{1/2^k}.$$

We give a short proof of this lemma in Appendix.

Based on the above lemma, we first prepare the following.

Lemma 2.7. *Under the assumptions of Proposition 2.3, for any integer k satisfying $2^k - 1 \geq -\frac{\alpha}{2}$, there exists a constant $C(\alpha, p, k, N) > 0$ such that for any $u \in H^1(\mathbb{R}^N)$*

with compact support, we have

$$\begin{aligned} \|e^{\frac{2\psi}{p+1}}u\|_{L^2} &\leq C(\alpha, p, k, N) \left((1+t)^{-\frac{1}{2}} \|e^\psi \sqrt{a}u\|_{L^2} + \|e^\psi \nabla u\|_{L^2} \right)^{1-1/2^k} \\ &\quad \times \left((1+t)^{(2^k-1+\frac{\alpha}{2})/(2-\alpha)} \|e^\psi \sqrt{a}u\|_{L^2} \right)^{1/2^k}. \end{aligned}$$

Proof. By Lemma 2.6, we have

$$(2.12) \quad \|e^{\frac{2\psi}{p+1}}u\|_{L^2} \leq C(k, N) \|\nabla(e^{\frac{2\psi}{p+1}}u)\|_{L^2}^{1-1/2^k} \|\langle x \rangle^{2^k-1} e^{\frac{2\psi}{p+1}}u\|_{L^2}^{1/2^k}$$

with k satisfying $2^k - 1 \geq -\frac{\alpha}{2}$. We estimate

$$\begin{aligned} \langle x \rangle^{2^k-1} e^{\frac{2\psi}{p+1}} &= \langle x \rangle^{-\frac{\alpha}{2}} \langle x \rangle^{2^k-1+\frac{\alpha}{2}} e^{\frac{2\psi}{p+1}} \\ &= \langle x \rangle^{-\frac{\alpha}{2}} \left(\frac{\langle x \rangle^{2-\alpha}}{1+t} \right)^{(2^k-1+\frac{\alpha}{2})/(2-\alpha)} e^{\frac{2\psi}{p+1}} \cdot (1+t)^{(2^k-1+\frac{\alpha}{2})/(2-\alpha)} \\ &\leq C(\alpha, p, k, N) (1+t)^{(2^k-1+\frac{\alpha}{2})/(2-\alpha)} e^{\psi(t,x)} \sqrt{a(x)}. \end{aligned}$$

On the other hand, we compute

$$\nabla(e^{\frac{2\psi}{p+1}}u) = e^{\frac{2\psi}{p+1}} \left(\frac{2}{p+1} (\nabla\psi)u + \nabla u \right)$$

and

$$\begin{aligned} |\nabla\psi| e^{\frac{2\psi}{p+1}} &\leq C \frac{\langle x \rangle^{1-\alpha}}{1+t} e^{\frac{2\psi}{p+1}} \\ &\leq C \langle x \rangle^{-\frac{\alpha}{2}} \left(\frac{\langle x \rangle^{2-\alpha}}{1+t} \right)^{(1-\frac{\alpha}{2})/(2-\alpha)} e^{\frac{2\psi}{p+1}} \cdot (1+t)^{(1-\frac{\alpha}{2})/(2-\alpha)-1} \\ &\leq C (1+t)^{-\frac{1}{2}} e^{\psi(t,x)} \sqrt{a(x)}. \end{aligned}$$

Plugging these estimates into (2.12), we have the desired estimate. \square

Combining Lemmas 2.5 and 2.7, we obtain the following interpolation estimate.

Lemma 2.8. *Under the assumptions of Proposition 2.3, for any integer k satisfying $2^k - 1 \geq -\frac{\alpha}{2}$, there exists a constant $C(\alpha, p, k, N) > 0$ such that for any $u \in H^1(\mathbb{R}^N)$ with compact support, we have*

$$\begin{aligned} \|e^{\frac{2\psi}{p+1}}u\|_{L^{p+1}} &\leq C(\alpha, p, k, N) \left((1+t)^{-\frac{1}{2}} \|e^\psi \sqrt{a}u\|_{L^2} + \|e^\psi \nabla u\|_{L^2} \right)^{\theta+(1-1/2^k)(1-\theta)} \\ &\quad \times \left((1+t)^{(2^k-1+\frac{\alpha}{2})/(2-\alpha)} \|e^\psi \sqrt{a}u\|_{L^2} \right)^{(1-\theta)/2^k}, \end{aligned}$$

where $\theta = \frac{N(p-1)}{2(p+1)} \in [0, 1]$.

Proof. By Lemmas 2.5 and 2.7, we estimate

$$\begin{aligned}
\|e^{\frac{2\psi}{p+1}}u\|_{L^{p+1}} &\leq C\|\nabla(e^{\frac{2\psi}{p+1}}u)\|_{L^2}^\theta\|e^{\frac{2\psi}{p+1}}u\|_{L^2}^{1-\theta} \\
&\leq C\left((1+t)^{-\frac{1}{2}}\|e^\psi\sqrt{a}u\|_{L^2} + \|e^\psi\nabla u\|_{L^2}\right)^\theta\|e^{\frac{2\psi}{p+1}}u\|_{L^2}^{1-\theta} \\
&\leq C\left((1+t)^{-\frac{1}{2}}\|e^\psi\sqrt{a}u\|_{L^2} + \|e^\psi\nabla u\|_{L^2}\right)^\theta \\
&\quad \times \left((1+t)^{-\frac{1}{2}}\|e^\psi\sqrt{a}u\|_{L^2} + \|e^\psi\nabla u\|_{L^2}\right)^{(1-1/2^k)(1-\theta)} \\
&\quad \times \left((1+t)^{(2^k-1+\frac{\alpha}{2})/(2-\alpha)}\|e^\psi\sqrt{a}u\|_{L^2}\right)^{(1/2^k)(1-\theta)} \\
&= C\left((1+t)^{-\frac{1}{2}}\|e^\psi\sqrt{a}u\|_{L^2} + \|e^\psi\nabla u\|_{L^2}\right)^{\theta+(1-1/2^k)(1-\theta)} \\
&\quad \times \left((1+t)^{(2^k-1+\frac{\alpha}{2})/(2-\alpha)}\|e^\psi\sqrt{a}u\|_{L^2}\right)^{(1-\theta)/2^k},
\end{aligned}$$

which completes the proof. \square

Now we are in a position to estimate the nonlinearities

$$\begin{aligned}
&C(t_0+t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0}\int_{\mathbb{R}^N}e^{2\psi}|F(u)|dx \\
&+ C\int_0^t(t_0+\tau)^{\frac{N-\alpha}{2-\alpha}-\delta_0}\int_{\mathbb{R}^N}e^{2\psi}|F(u)|dxd\tau \\
&+ C\int_0^t(t_0+\tau)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0}\int_{\mathbb{R}^N}e^{2\psi}(-\psi_t)|F(u)|dxd\tau \\
&=: N_1 + N_2 + N_3
\end{aligned}$$

in the right-hand side of Lemma 2.4. We first consider N_1 . Applying Lemma 2.8 and using the definition of $M(t)$ (see (2.6)), we deduce

$$\begin{aligned}
N_1 &\leq C(t_0+t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0}\|e^{\frac{2\psi}{p+1}}u\|_{L^{p+1}}^{p+1} \\
&\leq C(t_0+t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0}\left((1+t)^{-\frac{1}{2}}\|e^\psi\sqrt{a}u\|_{L^2} + \|e^\psi\nabla u\|_{L^2}\right)^{(\theta+(1-1/2^k)(1-\theta))(p+1)} \\
&\quad \times \left((1+t)^{(2^k-1+\frac{\alpha}{2})/(2-\alpha)}\|e^\psi\sqrt{a}u\|_{L^2}\right)^{(1-\theta)(p+1)/2^k} \\
&\leq C(t_0+t)^{\frac{N-\alpha}{2-\alpha}+1-\delta_0}(1+t)^{-\frac{1}{2}(\frac{N-\alpha}{2-\alpha}+1-\delta_0)(\theta+(1-1/2^k)(1-\theta))(p+1)} \\
&\quad \times (1+t)^{[(2^k-1+\frac{\alpha}{2})/(2-\alpha)-\frac{1}{2}(\frac{N-\alpha}{2-\alpha}-\delta_0)](1-\theta)(p+1)/2^k}M(t)^{\frac{p+1}{2}}.
\end{aligned}$$

By a straightforward calculation, we can see that the condition

$$\begin{aligned}
&\frac{N-\alpha}{2-\alpha}+1-\delta_0-\frac{1}{2}\left(\frac{N-\alpha}{2-\alpha}+1-\delta_0\right)\left(\theta+\left(1-\frac{1}{2^k}\right)(1-\theta)\right)(p+1) \\
&+ \left[\left(2^k-1+\frac{\alpha}{2}\right)\frac{1}{2-\alpha}-\frac{1}{2}\left(\frac{N-\alpha}{2-\alpha}-\delta_0\right)\right](1-\theta)(p+1)\frac{1}{2^k} < 0
\end{aligned}$$

if and only if

$$(2.13) \quad p > 1 + \frac{2}{N-\alpha-(2-\alpha)\delta_0/2}.$$

Noting $p > 1 + \frac{2}{N-\alpha}$ and $\delta_0 < \frac{2}{2-\alpha}(N - \alpha - \frac{2}{p-1})$, we see have (2.13) holds and hence,

$$(2.14) \quad N_1 \leq CM(t)^{\frac{p+1}{2}}.$$

We can obtain the same estimate as (2.14) for N_2 in the same way. Finally, for N_3 , noting

$$(-\psi_t)e^{2\psi} \leq C \frac{\langle x \rangle^{2-\alpha}}{(1+t)^2} e^{2\psi} \leq C(1+t)^{-1} e^{\frac{p+3}{2}\psi},$$

we have

$$N_3 \leq C \int_0^t (t_0 + \tau)^{\frac{N-\alpha}{2-\alpha} - \delta_0} \int_{\mathbb{R}^N} e^{\frac{p+3}{2}\psi} |F(u)| dx d\tau.$$

We can apply the same argument to the right-hand side and obtain the same estimate as (2.14) for N_3 . This completes the proof of Proposition 2.3.

3. BLOW-UP AND THE SHARP UPPER ESTIMATES OF THE LIFESPAN IN THE SUBCRITICAL AND THE CRITICAL CASE

In this section, we give a proof of Theorem 1.4 for the case $\alpha < 0, \beta = 1$. We can also prove Theorem 1.4 for the case $\alpha < 0, \beta = 0$ in the same argument with a slight modification (see Remark 3.1 below). The proof is based on the test-function method developed by Ikeda and Sobajima [15].

First, we remark that if $T(\varepsilon) \leq R_0$, then the assertion of Theorem 1.4 is obvious, provided that $\varepsilon \leq 1$. Thus, we may assume that $T(\varepsilon) > R_0$. Let $\eta = \eta(s)$ be a test function such that

$$\eta(s) = \begin{cases} 1 & \text{if } s \leq \frac{1}{2}, \\ \text{decreasing} & \text{if } \frac{1}{2} < s < 1, \\ 0 & \text{if } s \geq 1. \end{cases}$$

Let $R \in [R_0, T(\varepsilon))$ a parameter and, we define

$$\psi_R(t, x) = \left[\eta \left(\frac{|x|^{2-\alpha} + t^2}{R^2} \right) \right]^{2p'}.$$

We also define

$$\eta^*(s) = \begin{cases} 0 & \text{if } s \leq \frac{1}{2}, \\ \eta(s) & \text{if } s > \frac{1}{2}, \end{cases} \quad \psi_R^*(t, x) = \left[\eta^* \left(\frac{|x|^{2-\alpha} + t^2}{R^2} \right) \right]^{2p'}.$$

Then, we have

$$\begin{aligned} |\partial_t \psi_R(t, x)| &\leq CR^{-2}(1+t) [\psi_R^*(t, x)]^{1-\frac{1}{2p'}}, \\ |\partial_t^2 \psi_R(t, x)| &\leq CR^{-2} [\psi_R^*(t, x)]^{\frac{1}{p}}, \\ |\Delta \psi_R(t, x)| &\leq CR^{-2} \langle x \rangle^{-\alpha} [\psi_R^*(t, x)]^{\frac{1}{p}} \end{aligned}$$

and

$$\langle x \rangle^{-\alpha} \leq CR^{-\frac{2\alpha}{2-\alpha}} \quad \text{on } \text{supp } \psi_R, \text{ for } R \geq R_0.$$

Finally, we define

$$\Psi_R(t, x) = (1+t)\psi_R(t, x).$$

Multiplying the equation (1.1) by Ψ_R and integrating it over \mathbb{R}^N , we have

$$\int_{\mathbb{R}^N} |u|^p \Psi_R dx = \int_{\mathbb{R}^N} \left(u_{tt} - \Delta u + \frac{a(x)}{1+t} u_t \right) \Psi_R dx.$$

By integration by parts, we calculate

$$\begin{aligned} \int_{\mathbb{R}^N} |u|^p \Psi_R dx &= \frac{d}{dt} \int_{\mathbb{R}^N} \left(u_t \Psi_R - u \partial_t \Psi_R + \frac{a(x)}{1+t} u \Psi_R \right) dx \\ &\quad + \int_{\mathbb{R}^N} u \left(\partial_t^2 \Psi_R - \Delta \Psi_R - \partial_t \left(\frac{a(x)}{1+t} \Psi_R \right) \right) dx \\ &= \frac{d}{dt} \int_{\mathbb{R}^N} \left(u_t \Psi_R - u \partial_t \Psi_R + \frac{a(x)}{1+t} u \Psi_R \right) dx \\ &\quad + \int_{\mathbb{R}^N} u \left(2\partial_t \psi_R + (1+t) \partial_t^2 \psi_R + (1+t) \Delta \psi_R - a(x) \partial_t \psi_R \right) dx \\ &\leq \frac{d}{dt} \int_{\mathbb{R}^N} \left(u_t \Psi_R - u \partial_t \Psi_R + \frac{a(x)}{1+t} u \Psi_R \right) dx \\ &\quad + CR^{-2} \int_{\mathbb{R}^N} |u| (1 + \langle x \rangle^{-\alpha}) (1+t) [\psi_R^*]^{\frac{1}{p}} dx \\ &\leq \frac{d}{dt} \int_{\mathbb{R}^N} \left(u_t \Psi_R - u \partial_t \Psi_R + \frac{a(x)}{1+t} u \Psi_R \right) dx \\ &\quad + CR^{-\frac{4}{2-\alpha}} \int_{\mathbb{R}^N} |u| (1+t) [\psi_R^*]^{\frac{1}{p}} dx. \end{aligned}$$

Integrating it over $[0, R]$ and applying the Hölder inequality, we have

(3.1)

$$\begin{aligned} &\varepsilon \int_{\mathbb{R}^N} (u_1(x) + (a(x) - 1)u_0(x)) dx + \iint_{\mathbb{R}^N \times (0, R)} |u|^p (1+t) \psi_R dx dt \\ &\leq CR^{-\frac{4}{2-\alpha}} \left(\iint_{\mathbb{R}^N \times (0, R)} |u|^p (1+t) \psi_R^* dx dt \right)^{\frac{1}{p}} \left(\iint_{|x|^{2-\alpha} + t^2 \leq R^2} (1+t) dx dt \right)^{\frac{1}{p'}} \\ &\leq CR^{-\frac{4}{2-\alpha} \left(\frac{1}{p-1} - \frac{N-\alpha}{2} \right) \frac{1}{p'}} \left(\iint_{\mathbb{R}^N \times (0, R)} |u|^p (1+t) \psi_R^* dx dt \right)^{\frac{1}{p}}. \end{aligned}$$

In the subcritical case $1 < p < 1 + \frac{2}{N-\alpha}$, from $\psi_R^* \leq \psi_R$ and the Young inequality, we obtain

$$\varepsilon \int_{\mathbb{R}^N} (u_1(x) + (a(x) - 1)u_0(x)) dx \leq CR^{-\frac{4}{2-\alpha} \left(\frac{1}{p-1} - \frac{N-\alpha}{2} \right)}.$$

By the assumption of Theorem 1.4, the left-hand side is bounded from below by $C\varepsilon$. Therefore, we have

$$R \leq C\varepsilon^{-\frac{2-\alpha}{4} \left(\frac{1}{p-1} - \frac{N-\alpha}{2} \right)^{-1}}.$$

Since R is arbitrary in $[1, T(\varepsilon))$, we obtain the desired estimate for $T(\varepsilon)$.

In the critical case $p = 1 + \frac{2}{N-\alpha}$, we define

$$Y(\rho) = \int_0^\rho \left(\iint_{\mathbb{R}^N \times (0, R)} |u|^p (1+t) \psi_R^* dx dt \right) R^{-1} dR.$$

We note that the changing variable $s = \frac{\sqrt{|x|^{2-\alpha} + t^2}}{R}$ implies

$$\begin{aligned} Y(\rho) &\leq \iint_{\mathbb{R}^N \times (0, \rho)} |u|^p (1+t) \left(\int_0^\rho \left[\eta^* \left(\frac{|x|^{2-\alpha} + t^2}{R^2} \right) \right]^{2p'} R^{-1} dR \right) dx dt \\ &= \iint_{\mathbb{R}^N \times (0, \rho)} |u|^p (1+t) \left(\int_{\frac{\sqrt{|x|^{2-\alpha} + t^2}}{\rho}}^\infty [\eta^*(s^2)]^{2p'} s^{-1} ds \right) dx dt. \end{aligned}$$

Here, noting $\text{supp } \eta^* \subset [\frac{1}{2}, 1]$, $\eta^* \leq \eta$ and η is decreasing, we estimate

$$\int_{\frac{\sqrt{|x|^{2-\alpha} + t^2}}{\rho}}^\infty [\eta^*(s^2)]^{2p'} s^{-1} ds \leq \log 2 \left[\eta \left(\frac{|x|^{2-\alpha} + t^2}{\rho^2} \right) \right]^{2p'}.$$

Hence, we obtain

$$Y(\rho) \leq \log 2 \iint_{\mathbb{R}^N \times (0, \rho)} |u|^p (1+t) \psi_\rho dx dt.$$

Combining this with (3.1), we have the differential inequality of $Y(R)$

$$\left(\varepsilon \int_{\mathbb{R}^N} (u_1(x) + (a(x) - 1)u_0(x)) dx + Y(R) \right)^p \leq CRY'(R)$$

for $R \geq 1$. Noting $Y(1) = 0$ and solving the above, we conclude

$$\log R \leq C \left(\varepsilon \int_{\mathbb{R}^N} (u_1(x) + (a(x) - 1)u_0(x)) dx \right)^{-(p-1)}.$$

Since R is arbitrary in $[1, T(\varepsilon))$, we obtain the desired estimate for $T(\varepsilon)$.

Remark 3.1. In the case $\alpha < 0, \beta = 0$, we modify the definition of $\psi_R(t, x)$ and $\psi_R^*(t, x)$ by

$$\psi_R(t, x) = \left[\eta \left(\frac{|x|^{2-\alpha} + t}{R} \right) \right]^{2p'}, \quad \psi_R^*(t, x) = \left[\eta^* \left(\frac{|x|^{2-\alpha} + t}{R} \right) \right]^{2p'},$$

and we use ψ_R itself instead of Ψ_R . Then, corresponding to (3.1), we can prove

$$\begin{aligned} &\varepsilon \int_{\mathbb{R}^N} (u_1(x) + a(x)u_0(x)) dx + \iint_{\mathbb{R}^N \times (0, R)} |u|^p \psi_R dx dt \\ &\leq CR^{-\frac{2}{2-\alpha}(\frac{1}{p-1} - \frac{N-\alpha}{2})\frac{1}{p'}} \left(\iint_{\mathbb{R}^N \times (0, R)} |u|^p \psi_R^* dx dt \right)^{\frac{1}{p}}. \end{aligned}$$

From this, one has the assertion of Theorem 1.4 for $\alpha < 0, \beta = 0$ in the same manner.

APPENDIX A. SHORT PROOF OF A SPECIAL CASE OF THE CAFFARELLI-KOHN-NIRENBERG INEQUALITY

We give a short proof of Lemma 2.6. For any $\gamma \geq 0$, by the integration by parts, we compute

$$\begin{aligned} \int_{\mathbb{R}^N} \nabla \cdot (|x|^\gamma x) u^2 dx &= -2 \int_{\mathbb{R}^N} |x|^\gamma (x \cdot \nabla u) u dx \\ &\leq 2 \|\nabla u\|_{L^2} \| |x|^{\gamma+1} u \|_{L^2}, \end{aligned}$$

which leads to

$$(A.1) \quad \| |x|^{\gamma/2} u \|_{L^2}^2 \leq C(\gamma, N) \|\nabla u\|_{L^2} \| |x|^{\gamma+1} u \|_{L^2}.$$

Taking $\gamma = 0$ and $\gamma = 2$, we have

$$(A.2) \quad \|u\|_{L^2}^2 \leq C(N) \|\nabla u\|_{L^2} \| |x| u \|_{L^2}$$

and

$$(A.3) \quad \| |x| u \|_{L^2}^2 \leq C(\gamma, N) \|\nabla u\|_{L^2} \| |x|^3 u \|_{L^2},$$

respectively. The inequality (A.2) gives the assertion for $k = 1$. Next, combining (A.2) and (A.3), we deduce

$$(A.4) \quad \|u\|_{L^2}^2 \leq C(N) \|\nabla u\|_{L^2}^{3/4} \| |x|^3 u \|_{L^2}^{1/4},$$

which gives the assertion for $k = 2$. Furthermore, taking $\gamma = 6$ in (A.1), we have

$$\| |x|^3 u \|_{L^2}^2 \leq C(N) \|\nabla u\|_{L^2} \| |x|^7 u \|_{L^2}.$$

This and (A.4) imply

$$\|u\|_{L^2} \leq C(N) \|\nabla u\|_{L^2}^{7/8} \| |x|^7 u \|_{L^2}^{1/8},$$

which proves the assertion for $k = 3$. Repeating this argument, we obtain the desired estimate.

APPENDIX B. SKETCH OF A PRIORI ESTIMATE IN CASE OF TIME AND SPACE DEPENDENT COEFFICIENT DAMPING

Under Proposition 1.1, we here sketch the proof of the following proposition, corresponding to Proposition 2.3, assuming (i) $\alpha < 0$, $-1 < \beta < 1$ or (ii) $\alpha < 0$, $\beta = 1$ with $a_0 \gg 1$. Then we can reach to Theorems 1.2–1.3, using usual procedure of the energy method.

Proposition B.1. *Under the assumptions of Theorems 1.2–1.3, there exist $\delta_0 \in (0, \frac{(N-\alpha)(1+\beta)}{2-\alpha})$, and constants $t_0 = t_0(a_0, R_0, \delta_0) \geq 1$, $C = C(N, \alpha, \beta, a_0, R_0, \delta_0, t_0)$ such that the solution u constructed in Proposition 1.1 satisfies the a priori estimate*

$$M^{(\beta)}(t) \leq C M^{(\beta)}(0) + C M^{(\beta)}(t)^{\frac{p+1}{2}}$$

for $t \in [0, T(\varepsilon))$, where $\psi^{(\beta)}$ and $M^{(\beta)}(t)$ are defined in (B.1) and (B.2) with $\delta = \frac{2-\alpha}{2(N-\alpha)(1+\beta)} \delta_0$ below, respectively.

First, the function $\psi^{(\beta)}$ is defined by

$$(B.1) \quad \psi^{(\beta)}(t, x) = \frac{\mu}{B(t)} \left\{ \langle x \rangle^{2-\alpha} + A_0 - N * (\alpha(2-\alpha) \langle x \rangle^{-2-\alpha} \eta_{R_\delta}(x)) \right\},$$

where $B(t) = \frac{1}{1+\beta} + \int_0^t \frac{d\tau}{b(\tau)} = \frac{1}{1+\beta} (1+t)^{1+\beta}$. Since $\psi^{(0)}$ satisfies (2.4)–(2.5) in Lemma 2.2, and since $\psi^{(\beta)} = \frac{1+t}{B(t)} \psi^{(0)}$ and $(\frac{1}{B(t)})' = -\frac{(1+t)^\beta}{B(t)^2}$,

$$\begin{aligned} -c(t, x) \psi_t^{(\beta)} &= -c(t, x) \frac{(1+t)^\beta}{B(t)^2} (1+t)^2 \psi_t^{(0)} \geq -\left(\frac{1+t}{B(t)}\right)^2 a(x) \psi_t^{(0)} \\ &\geq \left(\frac{1+t}{B(t)}\right)^2 (2 + \delta_1^{(0)}) |\nabla \psi^{(0)}|^2 = (2 + \delta_1^{(0)}) |\nabla \psi^{(\beta)}|^2 \end{aligned}$$

and

$$\begin{aligned}\Delta\psi^{(\beta)} &= \frac{1+t}{B(t)}\Delta\psi^{(0)} \geq \frac{1+\beta}{(1+t)^\beta} \left(\frac{N-\alpha}{2(2-\alpha)} - \delta_2^{(0)} \right) \frac{a(x)}{1+t} \\ &= \left(\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} - \delta_2^{(\beta)} \right) \frac{c(t,x)}{1+t}.\end{aligned}$$

Here $\delta_1^{(\beta)} = \frac{2}{3}\delta$, $\delta_2^{(\beta)} = \frac{(N-\alpha)(1+\beta)}{2(2-\alpha)}\delta$. Hence, samely as in Lemma 2.2, it holds that

$$\begin{aligned}-\psi_t(t,x)c(t,x) &\geq (2 + \delta_1^{(\beta)})|\nabla\psi(t,x)|^2, \\ \Delta\psi(t,x) &\geq \left(\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} - \delta_2^{(\beta)} \right) \frac{c(t,x)}{1+t}.\end{aligned}$$

Next, since the number $\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)}$ suggests the decay rate, for the weighted energy estimate we define $M^{(\beta)}$ by

$$(B.2) \quad M^{(\beta)}(t) = \sup_{0 < \tau < t} \left\{ (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + \beta + 1 - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} (u_t^2 + |\nabla u|^2) dx \right. \\ \left. + (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} a(x) u^2 dx \right\},$$

where t_0 is suitably large number and $\delta_0 \geq 4\delta_2$.

Here and after, we abbreviate the suffix (β) for $\psi^{(\beta)}$, $\delta^{(\beta)}$ and $M^{(\beta)}$. Multiplying (1.1) by $e^{2\psi}u_t$ and $e^{2\psi}u$, we respectively have

$$(B.3) \quad \begin{aligned}\frac{\partial}{\partial t} \left[\frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) \right] - \nabla \cdot (e^{2\psi}u_t \nabla u) + e^{2\psi} \left\{ \left(\frac{1}{4}c(t,x) - \psi_t \right) u_t^2 + \frac{-\psi_t}{5} |\nabla u|^2 \right\} \\ \leq \frac{\partial}{\partial t} [e^{2\psi}F(u)] + 2e^{2\psi}(-\psi_t)F(u), \quad \text{where } F(u) = \frac{1}{p+1}|u|^p u,\end{aligned}$$

and

$$(B.4) \quad \begin{aligned}\frac{\partial}{\partial t} \left[e^{2\psi} \left(uu_t + \frac{c(t,x)}{2} u^2 \right) \right] - \Delta \left(\frac{e^{2\psi}}{2} u^2 \right) + e^{2\psi} \left(\delta_5 |\nabla u|^2 + \frac{\delta}{8} (-\psi_t) c(t,x) u^2 \right) \\ + \left(\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + \frac{\beta}{2} - \delta_2 \right) e^{2\psi} \frac{c(t,x)}{1+t} u^2 - e^{2\psi} \left(1 + \frac{8(-\psi_t)}{\delta c(t,x)} \right) u_t^2 \\ \leq e^{2\psi} |u|^p u,\end{aligned}$$

since $c(t,x) \frac{\partial}{\partial t} \frac{u^2}{2} = \frac{\partial}{\partial t} \frac{c(t,x)}{2} u^2 + \frac{\beta c(t,x)}{2(1+t)} u^2$, which is only different from the case $\beta = 0$, though $a(x)$ is changed to $c(t,x)$. So, note that $\delta_i (i \geq 3)$ is the same as one in Section 2 (cf. Remark 2.2). Integrating (B.3) over \mathbb{R}^N and multiplying it by

$(t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0}$, we have

$$\begin{aligned}
& \text{(B.5)} \\
& \frac{d}{dt} \left[(t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) dx \right] \\
& - \left\{ \frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0 \right\} (t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + \beta - \delta_0} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) dx \\
& + (t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left\{ \frac{1}{4} c(t, x) - \psi_t \right\} u_t^2 + \frac{-\psi_t}{5} |\nabla u|^2 \Big\} dx \\
& \leq \frac{d}{dt} \left[(t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} F(u) dx \right] \\
& - \left\{ \frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0 \right\} (t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + \beta - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} F(u) dx \\
& + 2(t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} (-\psi_t) F(u) dx.
\end{aligned}$$

Since

$$\text{(B.6)} \quad (t_0 + t)c(t, x) = (t_0 + t)a_0 \langle x \rangle^{-\alpha} (1+t)^{-\beta} \geq a_0 t_0^{1-\beta_+} \gg 1,$$

if (i) or (ii) holds, the term $\int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} u_t^2$ in the second term in (B.5) is absorbed into $\int_{\mathbb{R}^N} e^{2\psi} \frac{1}{4} c(t, x) u_t^2 dx$, where $\beta_+ = 0$ ($\beta \leq 0$), $\beta_+ = \beta$ ($0 < \beta \leq 1$). Hence, integration of (B.5) over $[0, t]$ yields

$$\begin{aligned}
& \text{(B.7)} \\
& (t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) dx \\
& + \int_0^t (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left\{ \left(\frac{1}{8} c(\tau, x) - \psi_t \right) u_t^2 + \frac{-\psi_t}{5} |\nabla u|^2 \right\} dx d\tau \\
& - \left\{ \frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0 \right\} \int_0^t (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} |\nabla u|^2 dx d\tau \\
& \leq C(t_0) \int_{\mathbb{R}^N} e^{2\psi(0, x)} (u_1^2 + |\nabla u_0|^2 + |u_0|^{p+1}) dx \\
& + (t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} F(u) dx \\
& - \left\{ \frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0 \right\} \int_0^t (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + \beta - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} F(u) dx d\tau \\
& + 2 \int_0^t (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} (-\psi_t) F(u) dx d\tau.
\end{aligned}$$

Similarly, integrating (B.4) over $\mathbb{R}^N \times [0, t]$, we have

(B.8)

$$\begin{aligned}
& (t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + \beta - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(uu_t + \frac{c(t, x)}{2} u^2 \right) dx \\
& + \int_0^t (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + \beta - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(\delta_5 |\nabla u|^2 + \frac{\delta}{8} (-\psi_t) c(\tau, x) u^2 \right) dx d\tau \\
& + \delta_6 \int_0^t (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + \beta - 1 - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} c(\tau, x) u^2 dx d\tau \\
& - \left(1 + \frac{C(\delta_6)}{a_0 t_0^{1-\beta_+}} \right) \int_0^t (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + \beta - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} u_t^2 dx d\tau \\
& - \frac{8}{\delta} \int_0^t (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + \beta - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} \frac{-\psi_t}{c(\tau, x)} u_t^2 dx d\tau \\
& \leq C(\alpha, \beta, t_0) \int_{\mathbb{R}^N} e^{2\psi(0, x)} \left(u_0 u_1 + \frac{c(0, x)}{2} u_0^2 \right) dx \\
& + \int_0^t (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + \beta - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} |u|^p u dx d\tau,
\end{aligned}$$

where we used, by (B.6),

$$\begin{aligned}
\int_{\mathbb{R}^N} e^{2\psi} uu_t dx & \leq \delta_6 \int_{\mathbb{R}^N} e^{2\psi} c(\tau, x) u^2 dx + (t_0 + \tau) \cdot \frac{C(\delta_0)}{(t_0 + \tau)c(\tau, x)} \int_{\mathbb{R}^N} e^{2\psi} u_t^2 dx \\
& \leq \delta_6 \int_{\mathbb{R}^N} e^{2\psi} c(\tau, x) u^2 dx + (t_0 + \tau) \cdot \frac{C(\delta_6)}{a_0 t_0^{1-\beta_+}} \int_{\mathbb{R}^N} e^{2\psi} u_t^2 dx.
\end{aligned}$$

Now, we add (B.7) to $\nu \cdot$ (B.8) ($0 < \nu \ll 1$) and cover the bad terms. In fact, fix ν small as $\nu \cdot \left\{ \frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0 \right\} \leq \frac{\delta_5}{2}$, and then take t_0 or a_0 large in case of (i) or (ii) as

$$1 + \frac{C(\delta_6)}{a_0 t_0^{1-\beta_+}} \leq \frac{\nu}{16} (t_0 + \tau) c(\tau, x) \quad \text{and} \quad \frac{8}{\delta} \cdot \frac{1}{c(\tau, x)} \leq \frac{\nu}{2} (t_0 + \tau),$$

that is, by (B.6),

$$1 + \frac{C(\delta_6)}{a_0 t_0^{1-\beta_+}} \leq \frac{\nu}{16} a_0 t_0^{1-\beta_+} \leq \frac{\nu}{16} (t_0 + \tau) c(\tau, x) \quad \text{and} \quad \frac{8}{\delta} \leq \frac{\nu}{2} a_0 t_0^{1-\beta_+} \leq \frac{\nu}{2} (t_0 + \tau) c(\tau, x).$$

Thus, the bad terms in the left hand sides in (B.7)–(B.8) are absorbed to good ones and the following desired inequality holds:

$$\begin{aligned}
& \nu(t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \int_{\mathbb{R}^N} \frac{e^{2\psi}}{2} (u_t^2 + |\nabla u|^2) dx \\
& + (t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + \beta - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} (uu_t + \frac{c(t, x)}{2} u^2) dx \\
& + \nu \int_0^t (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left\{ \left(\frac{1}{16} c(\tau, x) - \frac{1}{2} \psi_t \right) u_t^2 + \frac{-\psi_t}{5} |\nabla u|^2 \right\} dx d\tau \\
& + \int_0^t (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + \beta - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} \left(\frac{\delta_6}{2} |\nabla u|^2 + \frac{\delta}{8} (-\psi_t) c(\tau, x) u^2 \right) dx d\tau \\
& \leq C(t_0) \int_{\mathbb{R}^N} e^{2\psi(0, x)} \left(u_1^2 + |\nabla u_0|^2 + |u_0|^{p+1} + \frac{c(0, x)}{2} u_0^2 \right) dx \\
& + \nu(t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} |F(u)| dx \\
& + C(t_0) \int_0^t (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + \beta - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} |u|^{p+1} dx d\tau \\
& + 2\nu \int_0^t (t_0 + \tau)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} (-\psi_t) |F(u)| dx d\tau.
\end{aligned}$$

For the semilinear terms, we estimate, for an example,

$$\begin{aligned}
N_1 & := \nu(t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \int_{\mathbb{R}^N} e^{2\psi} |F(u)| dx \\
& = \nu(t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \|e^{\frac{2\psi}{p+1}} u\|_{L^{p+1}}^{p+1}.
\end{aligned}$$

By the Gagliardo-Nirenberg inequality

$$\begin{aligned}
\|e^{\frac{2\psi}{p+1}} u\|_{L^{p+1}} & \leq C \|\nabla(e^{\frac{2\psi}{p+1}} u)\|_{L^2}^\theta \|e^{\frac{2\psi}{p+1}} u\|_{L^2}^{1-\theta}, \quad \text{where } \theta = \frac{N(p-1)}{2(p+1)}, \\
& \leq C (\|e^{\frac{2\psi}{p+1}} |\nabla \psi| u\|_{L^2} + \|e^{\frac{2\psi}{p+1}} \nabla u\|_{L^2})^\theta \|e^{\frac{2\psi}{p+1}} u\|_{L^2}^{1-\theta}.
\end{aligned}$$

Since $\psi = \psi^{(\beta)}$ has the t -dependent coefficient $\frac{\mu}{B(t)}$, instead of $\frac{\mu}{1+t}$ for $\psi = \psi^{(0)}$ in Section 2, by the Caffarelli–Korn–Nirenberg inequality, samely as in Lemmas 2.7–2.8, we have

$$\begin{aligned}
& \|e^{\frac{2\psi}{p+1}} u\|_{L^{p+1}} \\
& \leq C(B(t))^{-\frac{1}{2}} \|e^\psi \sqrt{a} u\|_{L^2} + \|e^\psi \nabla u\|_{L^2}^{\theta + (1 - \frac{1}{2k})(1-\theta)} \times B(t)^{\frac{2k}{2-\alpha} - \frac{1}{2}} \|e^\psi \sqrt{a} u\|_{L^2}^{\frac{1}{2k}(1-\theta)}.
\end{aligned}$$

Hence

$$\begin{aligned}
N_1 & \leq C(t_0 + t)^{\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0} \times \left[(t_0 + t)^{-\frac{1}{2} \left(\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} + (1+\beta) - \delta_0 \right) (\theta + (1 - \frac{1}{2k})(1-\theta))} \right. \\
& \quad \left. \cdot (t_0 + t)^{\{(1+\beta)(\frac{2k}{2-\alpha} - \frac{1}{2} - \frac{1}{2}(\frac{(N-\alpha)(1+\beta)}{2(2-\alpha)} - \delta_0))\} \frac{1}{2k}(1-\theta)} \right]^{p+1} \times M(t)^{\frac{p+1}{2}}.
\end{aligned}$$

The exponent of $(t_0 + t)$ is

$$(1 + \beta) \left[\frac{N - \alpha}{2 - \alpha} + 1 - \frac{\delta_0}{1 + \beta} + (p + 1) \left\{ -\frac{1}{2} \left(\frac{N - \alpha}{2 - \alpha} + 1 - \frac{\delta_0}{1 + \beta} \right) \left(\theta + \left(1 - \frac{1}{2^k} \right) (1 - \theta) \right) \right. \right. \\ \left. \left. + \left(\left(\frac{2^k}{2 - \alpha} - \frac{1}{2} \right) - \frac{1}{2} \left(\frac{N - \alpha}{2 - \alpha} - \frac{\delta_0}{1 + \beta} \right) \right) \frac{1 - \theta}{2^k} \right\} \right].$$

Samely as (2.13), this is negative if and only if

$$p > 1 + \frac{2}{N - \alpha - (2 - \alpha)\delta_0/2(1 + \beta)},$$

because of $1 + \beta > 0$. The other semilinear terms are estimated in a similar fashion to the above. Thus, taking $\delta_0 > 0$ small, we obtain the desired estimate on $M(t) = M^{(\beta)}(t)$ if $p > 1 + \frac{2}{N - \alpha}$, which completes Proposition B.1 for $\beta \neq 0$.

ACKNOWLEDGEMENT

This work was supported by JSPS KAKENHI Grant Number JP18K134450 and JP16K17625.

REFERENCES

- [1] L. CAFFARELLI, R. KOHN, L. NIRENBERG, *First order interpolation inequalities with weights*, Compositio Math. **53** (1984), 259–275.
- [2] R. CHILL, A. HARAUX, *An optimal estimate for the difference of solutions of two abstract evolution equations*, J. Differential Equations **193** (2003), 385–395.
- [3] M. D’ABBICCO, *The threshold of effective damping for semilinear wave equations*, Math. Methods Appl. Sci. **38** (2015), 1032–1045.
- [4] M. D’ABBICCO, S. LUCENTE, *A modified test function method for damped wave equations*, Adv. Nonlinear Stud. **13** (2013), 867–892.
- [5] M. D’ABBICCO, S. LUCENTE, *NLWE with a special scale invariant damping in odd space dimension*, in: Discrete Contin. Dyn. Syst. 2015, Dynamical systems, differential equations and applications. 10th AIMS Conference. Suppl., 312–319.
- [6] M. D’ABBICCO, S. LUCENTE, M. REISSIG, *Semi-Linear wave equations with effective damping*, Chin. Ann. Math., Ser. B **34** (2013), 345–380.
- [7] M. D’ABBICCO, S. LUCENTE, M. REISSIG, *A shift in the Strauss exponent for semilinear wave equations with a not effective damping*, J. Differential Equations **259** (2015), 5040–5073.
- [8] H. FUJITA, *On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$* , J. Fac. Sci. Univ. Tokyo Sec. I, **13** (1966), 109–124.
- [9] K. FUJIWARA, M. IKEDA, Y. WAKASUGI, *Estimates of lifespan and blow-up rates for the wave equation with a time-dependent damping and a power-type nonlinearity*, to appear in Funkcial. Ekvac., arXiv:1609.01035v2.
- [10] M.-H. GIGA, Y. GIGA, J. SAAL, *Nonlinear Partial Differential Equations*, Progress in Nonlinear Differential Equations and their Applications, **79**, Birkhäuser, Boston, MA, 2010.
- [11] N. HAYASHI, E. I. KAIKINA, P. I. NAUMKIN, *Damped wave equation with super critical nonlinearities*, Differential Integral Equations **17** (2004), 637–652.
- [12] T. HOSONO, T. OGAWA, *Large time behavior and L^p - L^q estimate of solutions of 2-dimensional nonlinear damped wave equations*, J. Differential Equations **203** (2004), 82–118.
- [13] L. HSIAO, T. -P. LIU, *Convergence to nonlinear diffusion waves for solutions of a system of hyperbolic conservation laws with damping*, Comm. Math. Phys. **43** (1992), 599–605.
- [14] M. IKEDA, T. INUI, *The sharp estimate of the lifespan for the semilinear wave equation with time-dependent damping*, arXiv:1707.03950v1.
- [15] M. IKEDA, M. SOBAJIMA, *Upper bound for lifespan of solutions to certain semilinear parabolic, dispersive and hyperbolic equations via a unified test function method*, arXiv:1710.06780v1.
- [16] M. IKEDA, M. SOBAJIMA, *Life-span of blowup solutions to semilinear wave equation with space-dependent critical damping*, arXiv:1709.04401v1.

- [17] M. IKEDA, M. SOBAJIMA, *Life-span of solutions to semilinear wave equation with time-dependent critical damping for specially localized initial data*, to appear in *Mathematische Annalen*, arXiv:1709.04406v1.
- [18] M. IKEDA, M. SOBAJIMA, Y. WAKASUGI, *Sharp lifespan estimates of blowup solutions to semilinear wave equations with time-dependent effective damping*, arXiv:1808.06189v1.
- [19] M. IKEDA, Y. WAKASUGI, *A note on the lifespan of solutions to the semilinear damped wave equation*, *Proc. Amer. Math. Soc.* **143** (2015), 163–171.
- [20] M. IKEDA, Y. WAKASUGI, *Global well-posedness for the semilinear wave equation with time dependent damping in the overdamping case*, to appear in *Proc. Amer. Math. Soc.*
- [21] R. IKEHATA, *Some remarks on the wave equation with potential type damping coefficients*, *Int. J. Pure Appl. Math.* **21** (2005), 19–24.
- [22] R. IKEHATA, H. TAKEDA, *Uniform Energy Decay for Wave Equations with Unbounded Damping Coefficients*, arXiv:1706.03942v1.
- [23] R. IKEHATA, K. TANIZAWA, *Global existence of solutions for semilinear damped wave equations in \mathbb{R}^N with noncompactly supported initial data*, *Nonlinear Anal.* **61** (2005), 1189–1208.
- [24] R. IKEHATA, G. TODOROVA, B. YORDANOV, *Critical exponent for semilinear wave equations with space-dependent potential*, *Funkcialaj Ekvacioj* **52** (2009), 411–435.
- [25] R. IKEHATA, G. TODOROVA, B. YORDANOV, *Optimal decay rate of the energy for wave equations with critical potential*, *J. Math. Soc. Japan* **65** (2013), 183–236.
- [26] G. KARCH, *Selfsimilar profiles in large time asymptotics of solutions to damped wave equations*, *Studia Math.* **143** (2000), 175–197.
- [27] J. S. KENIGSON, J. J. KENIGSON, *Energy decay estimates for the dissipative wave equation with space-time dependent potential*, *Math. Meth. Appl. Sci.* **34** (2011), 48–62.
- [28] M. KHADER, *Nonlinear dissipative wave equations with spacetime dependent potential*, *Nonlinear Anal.* **74** (2011), 3945–3963.
- [29] M. KHADER, *Global existence for the dissipative wave equations with space-time dependent potential*, *Nonlinear Anal.* **81** (2013), 87–100.
- [30] M. KIRANE, M. QAFSAOUI, *Fujita’s exponent for a semilinear wave equation with linear damping*, *Adv. Nonlinear Stud.* **2** (2002), 41–49.
- [31] N. A. LAI, H. TAKAMURA, *Blow-up for semilinear damped wave equations with sub-Strauss exponent in the scattering case*, *Nonlinear Anal.* **168** (2018), 222–237.
- [32] N. A. LAI, H. TAKAMURA, K. WAKASA, *Blow-up for semilinear wave equations with the scale invariant damping and super-Fujita exponent*, *J. Differential Equations* **263** (2017), 5377–5394.
- [33] N. A. LAI, Y. ZHOU, *The sharp lifespan estimate for semilinear damped wave equation with Fujita critical power in higher dimensions*, to appear in *J. Math. Pure Appl.*
- [34] T.-T. LI, Y. ZHOU, *Breakdown of solutions to $\square u + u_t = |u|^{1+\alpha}$* , *Discrete Contin. Dynam. Syst.* **1** (1995), 503–520.
- [35] J. LIN, K. NISHIHARA, J. ZHAI, *L^2 -estimates of solutions for damped wave equations with space-time dependent damping term*, *J. Differential Equations* **248** (2010), 403–422.
- [36] J. LIN, K. NISHIHARA, J. ZHAI, *Decay property of solutions for damped wave equations with space-time dependent damping term*, *J. Math. Anal. Appl.* **374** (2011) 602–614.
- [37] J. LIN, K. NISHIHARA, J. ZHAI, *Critical exponent for the semilinear wave equation with time-dependent damping*, *Discrete Contin. Dyn. Syst.* **32** (2012), 4307–4320.
- [38] M. LIU, C. WANG, *Global existence for semilinear damped wave equations in relation with the Strauss conjecture*, arXiv1807.05908v1.
- [39] P. MARCATI, K. NISHIHARA, *The L^p - L^q estimates of solutions to one-dimensional damped wave equations and their application to the compressible flow through porous media*, *J. Differential Equations* **191** (2003), 445–469.
- [40] A. MATSUMURA, *On the asymptotic behavior of solutions of semi-linear wave equations*, *Publ. Res. Inst. Math. Sci.* **12** (1976), 169–189.
- [41] H. MICHIIHISA, *L^2 -asymptotic profiles of solutions to linear damped wave equations*, arXiv1710.04870v1.
- [42] K. MOCHIZUKI, *Scattering theory for wave equations with dissipative terms*, *Publ. Res. Inst. Math. Sci.* **12** (1976), 383–390.
- [43] K. MOCHIZUKI, H. NAKAZAWA, *Energy decay and asymptotic behavior of solutions to the wave equations with linear dissipation*, *Publ. RIMS, Kyoto Univ.* **32** (1996), 401–414.

- [44] T. NARAZAKI, L^p - L^q estimates for damped wave equations and their applications to semilinear problem, *J. Math. Soc. Japan* **56** (2004), 585–626.
- [45] K. NISHIHARA, *Asymptotic behavior of solutions of quasilinear hyperbolic equations with linear damping*, *J. Differential Equations* **137** (1997), 384–395.
- [46] K. NISHIHARA, L^p – L^q estimates of solutions to the damped wave equation in 3-dimensional space and their application, *Math. Z.* **244** (2003), 631–649.
- [47] K. NISHIHARA, L^p - L^q estimates for the 3-D damped wave equation and their application to the semilinear problem, *Seminar Notes of Math. Sci.* **6**, Ibaraki Univ., (2003), 69–83.
- [48] K. NISHIHARA, *Decay properties for the damped wave equation with space dependent potential and absorbed semilinear term*, *Comm. Partial Differential Equations* **35** (2010), 1402–1418.
- [49] A. PALMIERI, *A global existence result for a semilinear wave equation with scale-invariant damping and mass in even space dimension*, arXiv:1804.03978v1.
- [50] P. RADU, G. TODOROVA, B. YORDANOV, *Higher order energy decay rates for damped wave equations with variable coefficients*, *Discrete Contin. Dyn. Syst. Ser. S.* **2** (2009), 609–629.
- [51] P. RADU, G. TODOROVA, B. YORDANOV, *Decay estimates for wave equations with variable coefficients*, *Trans. Amer. Math. Soc.* **362** (2010), 2279–2299.
- [52] P. RADU, G. TODOROVA, B. YORDANOV, *The generalized diffusion phenomenon and applications*, *SIAM J. Math. Anal.* **48** (2016), 174–203.
- [53] J. RAUCH, M. TAYLOR, *Exponential decay of solutions to hyperbolic equations in bounded domains*, *Indiana Univ. Math. J.* **24** (1974), 79–86.
- [54] S. SAKATA, Y. WAKASUGI *Movement of time-delayed hot spots in Euclidean space*, *Math. Z.* **285** (2017), 1007–1040.
- [55] M. SOBAJIMA, Y. WAKASUGI, *Diffusion phenomena for the wave equation with space-dependent damping in an exterior domain*, *J. Differential Equations* **261** (2016), 5690–5718.
- [56] M. SOBAJIMA, Y. WAKASUGI, *Remarks on an elliptic problem arising in weighted energy estimates for wave equations with space-dependent damping term in an exterior domain*, *AIMS Mathematics* **2** (2017), 1–15.
- [57] M. SOBAJIMA, Y. WAKASUGI, *Diffusion phenomena for the wave equation with space-dependent damping term growing at infinity*, *Advances in Differential Equations* **23** (2018), 581–614.
- [58] M. SOBAJIMA, Y. WAKASUGI, *Weighted energy estimates for wave equation with space-dependent damping term for slowly decaying initial data*, to appear in *Commun. Contemp. Math.*
- [59] G. TODOROVA, B. YORDANOV, *Critical exponent for a nonlinear wave equation with damping*, *J. Differential Equations* **174** (2001), 464–489.
- [60] G. TODOROVA, B. YORDANOV, *Weighted L^2 -estimates for dissipative wave equations with variable coefficients*, *J. Differential Equations* **246** (2009), 4497–4518.
- [61] Z. TU, J. LIN, *A note on the blowup of scale invariant damping wave equation with sub-Strauss exponent*, arXiv:1709.00866v2.
- [62] K. WAKASA, *The Lifespan of solutions to semilinear damped wave equations in one space dimension*, *Commun. Pure Appl. Anal.* **15** (2016), 1265–1283.
- [63] K. WAKASA, B. YORDANOV, *On the blow-up for critical semilinear wave equation with damping in the scattering case*, arXiv:1807.06164v1.
- [64] Y. WAKASUGI, *Small data global existence for the semilinear wave equation with space-time dependent damping*, *J. Math. Anal. Appl.* **393** (2012), 66–79.
- [65] Y. WAKASUGI, *Critical exponent for the semilinear wave equation with scale invariant damping*, in: *Fourier Analysis* (M. Ruzhansky and V. Turunen, eds.), *Trends in Mathematics*, Birkhäuser, Basel, 2014, 375–390.
- [66] Y. WAKASUGI, *On diffusion phenomena for the linear wave equation with space-dependent damping*, *J. Hyp. Diff. Eq.* **11** (2014), 795–819.
- [67] Y. WAKASUGI, *Scaling variables and asymptotic profiles for the semilinear damped wave equation with variable coefficients*, *J. Math. Anal. Appl.* **447** (2017), 452–487.
- [68] Y. WAKASUGI, *Blow-up of solutions to the one-dimensional semilinear wave equation with damping depending on time and space variables*, *Discrete Contin. Dyn. Syst.* **34** (2014), 3831–3846.
- [69] Y. WAKASUGI, *On diffusion phenomena for the linear wave equation with space-dependent damping*, *J. Hyp. Diff. Eq.* **11** (2014), 795–819.

- [70] J. WIRTH, *Solution representations for a wave equation with weak dissipation*, Math. Meth. Appl. Sci. **27** (2004), 101–124.
- [71] J. WIRTH, *Wave equations with time-dependent dissipation I. Non-effective dissipation*, J. Differential Equations **222** (2006), 487–514.
- [72] J. WIRTH, *Wave equations with time-dependent dissipation II. Effective dissipation*, J. Differential Equations **232** (2007), 74–103.
- [73] J. WIRTH, *Scattering and modified scattering for abstract wave equations with time-dependent dissipation*, Adv. Differential Equations **12** (2007), 1115–1133.
- [74] T. YAMAZAKI, *Asymptotic behavior for abstract wave equations with decaying dissipation*, Adv. Differential Equations **11** (2006), 419–456.
- [75] H. YANG, A. MILANI, *On the diffusion phenomenon of quasilinear hyperbolic waves*, Bull. Sci. Math. **124** (2000), 415–433.
- [76] QI S. ZHANG, *A blow-up result for a nonlinear wave equation with damping: the critical case*, C. R. Acad. Sci. Paris Sér. I Math. **333** (2001), 109–114.

(K. Nishihara) PROFESSOR EMERITUS, WASEDA UNIVERSITY, TOKYO 169-8050, JAPAN
E-mail address: `kenji@waseda.jp`

(M. Sobajima) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY, TOKYO UNIVERSITY OF SCIENCE, 2641 YAMAZAKI, NODA-SHI, CHIBA, 278-8510, JAPAN
E-mail address: `msobajima1984@gmail.com`

(Y. Wakasugi) DEPARTMENT OF ENGINEERING FOR PRODUCTION AND ENVIRONMENT, GRADUATE SCHOOL OF SCIENCE AND ENGINEERING, EHIME UNIVERSITY, 3 BUNKYO-CHO, MATSUYAMA, EHIME, 790-8577, JAPAN
E-mail address: `wakasugi.yuta.vi@ehime-u.ac.jp`