

Vieta Matrix and its Determinant

Ufuk KAYA^a

^a*Department of Mathematics, Faculty of Arts and Sciences, Bitlis Eren University,
Bitlis, Turkey*

Abstract

In this paper, we define a matrix which we call Vieta matrix and calculate its determinant:

$$\begin{pmatrix} 1 & 1 & \cdots & 1 \\ a_2 + a_3 + \cdots + a_n & a_1 + a_3 + \cdots + a_n & \cdots & a_1 + a_2 + \cdots + a_{n-1} \\ a_2a_3 + a_2a_4 + \cdots + a_{n-1}a_n & a_1a_3 + a_1a_4 + \cdots + a_{n-1}a_n & \cdots & a_1a_2 + a_1a_3 + \cdots + a_{n-2}a_{n-1} \\ \vdots & \vdots & \ddots & \vdots \\ a_2a_3 \cdots a_{n-1}a_n & a_1a_3 \cdots a_{n-1}a_n & \cdots & a_1a_2 \cdots a_{n-2}a_{n-1} \end{pmatrix}$$

Keywords: Vieta's formula, matrix, determinant

2010 MSC: 15A15

1. Introduction

The determinant of a square matrix can be calculated by Leibnitz formula or Laplace formula. Especially, the determinant of a 3×3 matrix can be calculated by rule of Sarrus. Generally, the determinants of 3×3 matrices can be easily calculated, but 4×4 and more are so difficult. In some cases, the determinants of $n \times n$ matrices can be formulated, such as Vandermonde

URL: mat-ufuk@hotmail.com (Ufuk KAYA)

matrix:

$$\begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ a_1 & a_2 & a_3 & \cdots & a_n \\ a_1^2 & a_2^2 & a_3^2 & \cdots & a_n^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_1^{n-1} & a_2^{n-1} & a_3^{n-1} & \cdots & a_n^{n-1} \end{pmatrix}.$$

The determinant of Vandermonde matrix is

$$\prod_{k>i} (a_k - a_i).$$

In this paper, we define a matrix which we call Vieta matrix:

$$\begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ a_2 + a_3 + \cdots + a_n & a_1 + a_3 + \cdots + a_n & a_1 + a_2 + \cdots + a_n & \cdots & a_1 + a_2 + \cdots + a_{n-1} \\ a_2a_3 + a_2a_4 + \cdots + a_{n-1}a_n & a_1a_3 + a_1a_4 + \cdots + a_{n-1}a_n & a_1a_2 + a_1a_4 + \cdots + a_{n-1}a_n & \cdots & a_1a_2 + a_1a_3 + \cdots + a_{n-2}a_{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_2a_3 \cdots a_{n-1}a_n & a_1a_3 \cdots a_{n-1}a_n & a_1a_2 \cdots a_{n-1}a_n & \cdots & a_1a_2 \cdots a_{n-2}a_{n-1} \end{pmatrix} \quad (1)$$

and we calculate its determinant. Also, we give some applications of it.

2. The Determinant of Vieta Matrix

Definition 1. We state that the matrix given by the formula (1) is the Vieta matrix of the complex numbers a_1, a_2, \dots, a_n . It is equal to the following

$$\begin{pmatrix} 1 & 1 \\ a_2 & a_1 \end{pmatrix}$$

for $n = 2$,

$$\begin{pmatrix} 1 & 1 & 1 \\ a_2 + a_3 & a_1 + a_3 & a_1 + a_2 \\ a_2a_3 & a_1a_3 & a_1a_2 \end{pmatrix}$$

for $n = 3$ and

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ a_2 + a_3 + a_4 & a_1 + a_3 + a_4 & a_1 + a_2 + a_4 & a_1 + a_2 + a_3 \\ a_2a_3 + a_2a_4 + a_3a_4 & a_1a_3 + a_1a_4 + a_3a_4 & a_1a_2 + a_1a_4 + a_2a_4 & a_1a_2 + a_1a_3 + a_2a_3 \\ a_2a_3a_4 & a_1a_3a_4 & a_1a_2a_4 & a_1a_2a_3 \end{pmatrix}$$

for $n = 4$.

One can easily calculate that the determinants of the above matrices are

$$a_1 - a_2,$$

$$(a_1 - a_2)(a_1 - a_3)(a_2 - a_3),$$

and

$$(a_1 - a_2)(a_1 - a_3)(a_1 - a_4)(a_2 - a_3)(a_2 - a_4)(a_3 - a_4)$$

respectively. Now we prove that the determinant of the Vieta matrix is

$$\prod_{1 \leq i < k \leq n} (a_i - a_k) \quad (2)$$

in the general case by the following theorem:

Theorem 1. The determinant of the Vieta matrix (1) is equal to the product in the formula (2).

Proof. It is obvious that if at least two numbers of a_1, a_2, \dots, a_n are equal to 0, then the last row of the matrix (1) is the zero vector, and so, the determinant is 0. Assume that the numbers a_1, a_2, \dots, a_n include at most one zero. We now use the induction method. For $n = 2$, the determinant of the Vieta matrix

$$\begin{pmatrix} 1 & 1 \\ a_2 & a_1 \end{pmatrix}$$

is $a_1 - a_2$. We assume that the assertion is valid for n , i.e. the determinant

3. Some Applications of Vieta Matrix

3.1. Wronskian of some polynomials

Let $a_1, a_2, \dots, a_n \in \mathbb{C}$. Consider the polynomials

$$\begin{aligned} p_1(x) &= (x - a_2)(x - a_3) \dots (x - a_n), \\ p_2(x) &= (x - a_1)(x - a_3) \dots (x - a_n), \\ &\vdots \\ p_n(x) &= (x - a_1)(x - a_2) \dots (x - a_{n-1}). \end{aligned}$$

We now find the Wronskian of these polynomials, i.e. calculate the following

$$W[p_1, p_2, \dots, p_n](x) = \begin{vmatrix} p_1(x) & p_2(x) & \dots & p_n(x) \\ p_1'(x) & p_2'(x) & \dots & p_n'(x) \\ \vdots & \vdots & \ddots & \vdots \\ p_1^{(n-1)}(x) & p_2^{(n-1)}(x) & \dots & p_n^{(n-1)}(x) \end{vmatrix}. \quad (5)$$

Note that the polynomials p_1, p_2, \dots, p_n can be rewritten as follows:

$$\begin{aligned} p_1(x) &= x^{n-1} - (a_2 + a_3 + \dots + a_n)x^{n-2} + (a_2a_3 + a_2a_4 + \dots + a_{n-1}a_n)x^{n-3} + \dots + (-1)^{n-1}a_2a_3 \dots a_n, \\ p_2(x) &= x^{n-1} - (a_1 + a_3 + \dots + a_n)x^{n-2} + (a_1a_3 + a_1a_4 + \dots + a_{n-1}a_n)x^{n-3} + \dots + (-1)^{n-1}a_1a_3 \dots a_n, \\ &\vdots \\ p_n(x) &= x^{n-1} - (a_1 + a_2 + \dots + a_{n-1})x^{n-2} + (a_1a_2 + a_1a_3 + \dots + a_{n-2}a_{n-1})x^{n-3} + \dots + (-1)^{n-1}a_1a_2 \dots a_{n-1}. \end{aligned} \quad (6)$$

Furthermore, each of the functions p_1, p_2, \dots, p_n is a solution of the linear differential equation $y^n = 0$. Then, the Wronskian $W[p_1, p_2, \dots, p_n](x)$ is independent of the variable x by Abel's identity, see [1]. So, it is equal to the value $W[p_1, p_2, \dots, p_n](0)$, i.e.

$$W[p_1, p_2, \dots, p_n](x) = W[p_1, p_2, \dots, p_n](0). \quad (7)$$

By the relations (5), (6), (7) and, Theorem 1 we have

$$W[p_1, p_2, \dots, p_n](x) =$$

$$\begin{aligned}
& \begin{vmatrix} (-1)^{n-1} a_2 a_3 \dots a_{n-1} a_n & (-1)^{n-1} a_1 a_3 \dots a_{n-1} a_n & \dots & (-1)^{n-1} a_1 a_2 \dots a_{n-2} a_{n-1} \\ \vdots & \vdots & \ddots & \vdots \\ - (n-2)! (a_2 + a_3 + \dots + a_n) & - (n-2)! (a_1 + a_3 + \dots + a_n) & \dots & - (n-2)! (a_1 + a_2 + \dots + a_{n-1}) \\ (n-1)! & (n-1)! & \dots & (n-1)! \end{vmatrix} \\
&= \prod_{k=0}^{n-1} k! \begin{vmatrix} 1 & 1 & \dots & 1 \\ a_2 + a_3 + \dots + a_n & a_1 + a_3 + \dots + a_n & \dots & a_1 + a_2 + \dots + a_{n-1} \\ \vdots & \vdots & \ddots & \vdots \\ a_2 a_3 \dots a_{n-1} a_n & a_1 a_3 \dots a_{n-1} a_n & \dots & a_1 a_2 \dots a_{n-2} a_{n-1} \end{vmatrix} \\
&= \prod_{k=0}^{n-1} k! \cdot \prod_{1 \leq i < k \leq n} (a_i - a_k).
\end{aligned}$$

3.2. Jacobian of some functions

Consider the functions

$$\begin{aligned}
f_1(x_1, x_2, \dots, x_n) &= x_1 + x_2 + \dots + x_n, \\
f_2(x_1, x_2, \dots, x_n) &= x_1 x_2 + x_1 x_3 + \dots + x_{n-1} x_n, \\
&\vdots \\
f_n(x_1, x_2, \dots, x_n) &= x_1 x_2 \dots x_n.
\end{aligned}$$

If we calculate the Jacobian of the functions f_1, f_2, \dots, f_n by using Theorem 1, we have

$$\begin{aligned}
& \begin{vmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} & \dots & \frac{\partial f_1}{\partial x_n} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} & \dots & \frac{\partial f_2}{\partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial x_1} & \frac{\partial f_n}{\partial x_2} & \dots & \frac{\partial f_n}{\partial x_n} \end{vmatrix} \\
&= \begin{vmatrix} 1 & 1 & \dots & 1 \\ x_2 + x_3 + \dots + x_n & x_1 + x_3 + \dots + x_n & \dots & x_1 + x_2 + \dots + x_{n-1} \\ \vdots & \vdots & \ddots & \vdots \\ x_2 x_3 \dots x_{n-1} x_n & x_1 x_3 \dots x_{n-1} x_n & \dots & x_1 x_2 \dots x_{n-2} x_{n-1} \end{vmatrix}
\end{aligned}$$

$$= \prod_{1 \leq i < k \leq n} (x_i - x_k).$$

References

- [1] Boyce WE, DiPrima RC. Elementary differential equations and boundary value problems. New York (NY): Wiley; 1986.