

# Representing the Smallest Ideal and its closure in the semigroup $(\beta N, \cdot)$ as equivalence classes under the extension of divisibility relations

Salahdeen Khalifa

Department of Mathematics and Computer Science  
University of Missouri- St. Louis  
St.Louis, MO 63121 U.S.A  
e-mail: skkf57@mail.umsl.edu

July 13, 2021

## Abstract

In this paper we will prove that all the elements in the smallest ideal  $K(\beta N)$  in the semigroup of the Stone Cech compactification  $(\beta N, \cdot)$  of the discrete semigroup of natural numbers  $N$  under multiplication constitute a single equivalence class under the relation  $=_m$  ( $x =_m y$  if  $x|_m y$  and  $y|_m x$ ). And all the elements in the closure of the smallest ideal  $Cl(K(\beta N))$  constitute a single equivalence class under the relation  $=_{\sim}$  ( $x =_{\sim} y$  if  $x|\tilde{y}$  and  $y|\tilde{x}$ )

## 1 Introduction

we consider the discrete space  $N$  of natural numbers and its Stone-Cech compactification  $\beta N = \{x : x \text{ is ultrafilter on } N\}$  with topology that is introduced by a base  $\mathcal{B} = \{\bar{A} : A \subseteq N\}$  where  $\bar{A} = \{x \in \beta N : A \in x\}$  is a clopen subset of  $\beta N$ . Then according to ([1] Chapter 4 (4.1)) we can extend the usual multiplication  $(\cdot)$  which is defined on the semigroup  $N$  to an operation that is defined on  $\beta N$  as follows:  $A \in x \cdot y$  iff  $\{n \in N : A/n \in y\} \in x$  where  $A/n = \{m \in N : mn \in A\}$ .  $\beta N$  contains the set of natural numbers  $N$

as principal ultrafilters which are defined on  $N$ , and the set of non-principal ultrafilters are denoted by  $N^* = \beta N - N$ .  $|\beta N| = 2^c$  where  $c = |R|$ .  
 $\mu = \{A \subseteq N : A = A \uparrow\}$  is called the collection of **upward closed** subsets of  $N$  where  $A \uparrow = \{n \in N : \exists a \in A, a|n\}$  and  $\nu = \{A \subseteq N : A = A \downarrow\}$  is called the collection of **downward closed** subsets of  $N$  where  $A \downarrow = \{n \in N : \exists a \in A, n|a\}$ .  $\mu$  and  $\nu$  have the finite intersection property and if  $A \in \mu$  then  $N - A \in \nu$ .

**Lemma 1.1.** (1) ([1] Theorem 4.15 ) Let  $x, y \in \beta N$  and let  $A \subseteq N$ . Then  $A \in x.y$  if and only if there exists  $B \in x$  and an indexed family  $(C_n)_{n \in B}$  in  $y$  such that  $\bigcup_{n \in B} nC_n \subseteq A$ .

(2) Let  $x, y \in \beta N$  and let  $A, B \subseteq N$ . Then

(b) If  $A \in x, B \in y$ , then  $AB \in xy$

(c)  $\overline{A.B} \subseteq \overline{AB}$  for any  $A, B \subseteq N$

## 2 Ideals

A subset  $L$  of  $\beta N$  is a left ideal if  $\beta N.L \subseteq L$ , and  $L$  is a minimal left ideal if for any left ideal  $L_1$  we have  $L \subset L_1$ . A subset  $R$  of  $\beta N$  is a right ideal if  $R.\beta N \subseteq R$ , and  $R$  is a minimal right ideal if for any right ideal  $R_1$  we have  $R \subset R_1$ . A subset  $J$  of  $\beta N$  is an ideal if  $J$  left (right) ideal, and  $J$  is a minimal ideal if it is a minimal left (right) ideal.

Each set of the form  $\beta Nx = \{px : p \in \beta N\}$ , where  $x \in \beta N$  is called principal left ideal generated by  $x$ , each set of the form  $x\beta N = \{xp : p \in \beta N\}$  where  $x \in \beta N$  is called principal right ideal generated by  $x$ , and each set of the form  $\beta Nx\beta N = \{pxq : p, q \in \beta N\}$  where  $x \in \beta N$  is called principal ideal generated by  $x$ . Each minimal left ideal  $L$  is principal:  $L = \beta Nx$  for every  $x \in L$ , each minimal right ideal  $R$  is principal:  $R = x\beta N$  for every  $x \in R$ , and each minimal ideal  $J$  is principal:  $J = \beta Nx\beta N$  for every  $x \in J$

**Theorem 2.1.** (a) ([1] Theorem 2.7 (d)) If  $L$  is a left ideal and  $R$  is a right ideal then  $L \cap R \neq \phi$

(b) ([1] Corollary 2.6, Theorem 2.7(a)) Every left (right) ideal in  $\beta N$  contains a minimal left (right) ideal.

**Theorem 2.2.** ([1] Theorem 6.30, Corollary 6.41) There are  $2^c$  disjoint minimal left (right) ideals in  $\beta N$  and each of them contains  $2^c$  elements.

**Theorem 2.3.** (a) ([1], Theorem 1.51)  $\beta N$  has the smallest ideal denoted by  $K(\beta N)$  that is contained in all other ideals in  $\beta N$  .  
(b) ([1], Theorem 1.64)

$$\begin{aligned} K(\beta N) &= \cup\{L : L \text{ is a minimal left ideal of } \beta N\} \\ &= \cup\{R : R \text{ is a minimal right ideal of } \beta N\} \end{aligned}$$

### 3 Divisibility relations on $\beta N$

We can extend the usual divisibility relation  $|$  that is defined on  $N$  to a divisibility relation on  $\beta N$  in many ways. If  $x, y \in \beta N$  then the most common extension divisibility relations of  $|$  are the following :  $y$  is **left-divisible** by  $x, x|_l y$  if there is  $z \in \beta N$  such that  $y = zx$  .  $y$  is **right-divisible** by  $x, x|_r y$ , if there is  $z \in \beta N$  such that  $y = xz$  .  $y$  is **mid-divisible** by  $x, x|_m y$  , if there are  $z, w \in \beta N$  such that  $y = zxw$  .  $y$  is  $\tilde{|}$  -**divisible** by  $x, x\tilde{|}y$  if for all  $A \subseteq N, A \in x$  implies  $|A = \{n \in N : \exists a \in A, a|n\} \in y$ . When  $x = n \in N$  , we write  $n|y, (y = nz, z \in \beta N)$  .

**Lemma 3.1.** (a) ([3])  $|_l, |_r \subset |_m \subset \tilde{|}$   
(b) ([3]) All the relations  $|_l, |_r, |_m$  and  $\tilde{|}$  are preorders (reflexive and transitive), but none of them is symmetric and antisymmetric  
Since  $|_l, |_r, |_m$  and  $\tilde{|}$  are not antisymmetric then we can introduce another relations on  $\beta N$  that will be symmetric such as :  $x =_l y$  if  $x|_l y$  and  $y|_l x, x =_r y$  if  $x|_r y$  and  $y|_r x, x =_m y$  if  $x|_m y$  and  $y|_m x$  and  $x =_{\sim} y$  if  $x\tilde{|}y$  and  $y\tilde{|}x$

**Lemma 3.2.** ([4] Lemma 1.4) The following conditions are equivalent:

- (a)  $x\tilde{|}y$
- (b)  $x \cap \mu \subset y \cap \mu$
- (c)  $y \cap \nu \subset x \cap \nu$

**Lemma 3.3.** (a)  $=_l, =_r, =_m$  and  $\tilde{|}$  are equivalence relations.

(b) ([4] Lemma 1.5 (b)) For each  $x \in \beta N$  the sets  $x \uparrow_{\sim} = \{y \in \beta N : x\tilde{|}y\}$  ,  $x \downarrow_{\sim} = \{y \in \beta N : y\tilde{|}x\}$  and  $[x]_{\sim}$  are closed subsets of  $\beta N$  .

### 4 Levels of ultrafilters

**Definition 4.1.** ([5] Definition 2.4) Let  $P$  be the set of prime numbers in  $N$ , and let

$$L_0 = \{1\}$$

$$L_1 = \{a : a \in P\}$$

$$L_2 = \{a_1 a_2 : a_1, a_2 \in P\}$$

·  
·  
·

$L_n = \{a_1 a_2 \dots a_n : a_1, a_2, \dots, a_n \in P\}$ . Then, the ultrafilters  $x$  is called on "finite level" if it is in exactly on one of the following sets :  $\bar{L}_i, i = 0, 1, 2, \dots$  where  $\bar{L}_i = \{y \in \beta N : L_i \in y\}$

**Lemma 4.2.** ([2] Lemma 3.1(c))  $\bigcup_{i=0}^{\infty} \bar{L}_i \neq \overline{\bigcup_{i=0}^{\infty} L_i}$

By the lemma(4.2) there are ultrafilters that are not on finite levels, these ultrafilters are given by the following definition

**Definition 4.3.** ( [2] Definition 3.2 ) The ultrafilters that belong to

$\mathcal{I} = \bigcap_{i=0}^{\infty} \overline{N - L_i}$  are called the "ultrafilters that are not on finite levels".

**Theorem 4.4.** ([2] Theorem 3.7)

(a) If  $x, y \in \beta N$  and  $x, y \in \mathcal{I}$  . Then  $xy \in \mathcal{I}$  and  $yx \in \mathcal{I}$

(b) If  $x, y \in \beta N$  and  $x \in \mathcal{I}, y \notin \mathcal{I}$  . Then  $xy \in \mathcal{I}$  and  $yx \in \mathcal{I}$

## 5 Smallest ideal $K(\beta N)$ in the semigroup $(\beta N, \cdot)$

**Lemma 5.1.** (a)  $\mathcal{I}$  is a left ideal in the semigroup  $(\beta N, \cdot)$

(b)  $\mathcal{I}$  is a right ideal in the semigroup  $(\beta N, \cdot)$

(c)  $\mathcal{I}$  is an ideal in the semigroup  $(\beta N, \cdot)$

*Proof.* By (Theorem 4.4) ■

**Lemma 5.2.** (a)  $\mathcal{I}$  is not a minimal left ideal in  $(\beta N, \cdot)$

(b)  $\mathcal{I}$  is not a minimal right ideal in  $(\beta N, \cdot)$

(c)  $\mathcal{I}$  is not a minimal ideal in  $(\beta N, \cdot)$

*Proof.* (a) Let  $L = \{2x : x \in \mathcal{I}\}$  , then  $L$  is a left ideal and  $L \subset \mathcal{I}, L \neq \mathcal{I}$  [ by ( [1], Lemma 6.28 )  $2x \neq nx$  for any  $n \in N, n \neq 2$  ] . Thus  $\mathcal{I}$  is not a minimal left ideal.

(b) Similar to (a)

(c) By (a) and (b) ■

**Corollary 5.3.** *The smallest ideal  $K(\beta N)$  of the semigroup  $(\beta N, \cdot)$  exists and is a subset of  $\mathcal{I}$  i.e  $K(\beta N) \subset \mathcal{I}$*

*Proof.* By (Theorem 2.2)  $\beta N$  has  $2^c$  minimal left (right) ideals and by ([1 Lemma 1-45) for any minimal left ideal  $L$  and minimal right ideal  $R$  we have  $L \subseteq \mathcal{I}$  and  $R \subseteq \mathcal{I}$ , and by (Lemma 5.2) since  $\mathcal{I}$  is not a minimal ideal, then  $L \subset \mathcal{I}$  and  $R \subset \mathcal{I}$ . Then, by (Theorem 2.3 (a))  $K(\beta N)$  exists and

$$\begin{aligned} K(\beta N) &= \cup\{L : L \text{ is a minimal left ideal of } \beta N\} \\ &= \cup\{R : R \text{ is a minimal right ideal of } \beta N\} \end{aligned}$$

Thus  $K(\beta N) \subset \mathcal{I}$  ■

**Lemma 5.4.** (a) *If  $L$  is a minimal left ideal and  $x \in L$ . Then  $x|_l y$  if and only if  $y \in L$*

(b) *If  $R$  is a minimal right ideal and  $x \in R$ , Then  $x|_r y$  if and only if  $y \in R$*

(c) *If  $J$  is a minimal ideal and  $x \in J$ . then  $x|_m y$  if and only if  $y \in J$*

*Proof.* (a) ( $\Rightarrow$ ) Let  $L$  be a minimal left ideal. Then  $L = \beta N x$  for any  $x \in L$ . Let  $x|_l y$ , then  $y = zx$  for some  $z \in \beta N$ . Thus  $y \in \beta N x$  so  $y \in L$ .

( $\Leftarrow$ ) If  $y \in L$ , then  $y \in \beta N x$  for any  $x \in L$ , so  $y = zx$  for some  $z \in \beta N$ . Thus  $x|_l y$ .

(b) ( $\Rightarrow$ ) Let  $R$  be a minimal right ideal. Then  $R = x\beta N$  for any  $x \in R$ . Let  $x|_r y$ , then  $y = xz$  for some  $z \in \beta N$ . Thus  $y \in x\beta N$ , so  $y \in R$ .

( $\Leftarrow$ ) If  $y \in R$  then  $y \in x\beta N$  for any  $x \in R$ , so  $y = xz$  for some  $z \in \beta N$ . Thus  $x|_r y$ .

(c) ( $\Rightarrow$ ) Let  $J$  be a minimal ideal. Then  $J = \beta N x \beta N$  for any  $x \in J$ . Let  $x|_m y$ , then  $y = zxw$  for some  $z, w \in \beta N$ . Thus  $y \in \beta N x \beta N$  so  $y \in J$ .

( $\Leftarrow$ ) If  $y \in J$ , then  $y \in \beta N x \beta N$  for any  $x \in J$ , so  $y = zxw$  for some  $z, w \in \beta N$ . Thus  $x|_m y$  ■

**Theorem 5.5.** *Let  $(\beta N, =_l)$  where  $x =_l y$  if  $x|_l y$  and  $y|_l x$ . Then*

(a) *If  $x \in K(\beta N)$ , then  $[x]_l = L$ ,  $L$  is a minimal left ideal,  $x \in L$ , and  $[x]_l$  has  $2^c$  elements.*

(b)  *$K(\beta N)$  is partitioned by  $2^c$  equivalence classes  $[x]_l$ .*

*Proof.* (a) Let  $x \in K(\beta N)$ , then by (Theorem 2.2) there exists only one minimal left ideal  $L$  such that  $x \in L$ . Since  $L$  is a minimal left ideal, then  $L$  is principal, so  $L = \beta N x$ . Also for any  $y \in L$  we have  $L = \beta N y$ . Then  $y \in \beta N x$ , and there exist  $z \in \beta N$  such that  $y = zx$ . Thus  $x|_l y$ . Also

$x \in \beta N y$ , and there exist  $w \in \beta N$  such that  $x = wy$ . Thus  $y|_l x$ . Therefore  $x =_l y$  for any  $y \in L$ . If  $p \in \beta N, p \notin L = \beta N x$ , by (Lemma 5.4(a)) we have  $x \neq_l p$ . So  $[x]_l = L$ . Since  $L$  has  $2^c$  elements, then  $[x]_l$  has  $2^c$  elements.  
(b) Since  $K(\beta N)$  has  $2^c$  disjoint minimal left ideals, and by (a) each of them is  $[x]_l$  equivalence class. Then  $K(\beta N)$  is partitioned by  $2^c$  equivalence classes  $[x]_l$ . ■

**Theorem 5.6.** *Let  $(\beta N, =_r)$  such that  $x =_r y$  if  $x|_r y$  and  $y|_r x$ . Then*  
(a) *If  $x \in K(\beta N)$ , then  $[x]_r = R, R$  is a minimal right ideal,  $x \in R$ , and  $[x]_r$  has  $2^c$  elements.*  
(b)  *$K(\beta N)$  is partitioned by  $2^c$  equivalence classes  $[x]_R$ .*

*Proof.* Similar to (Theorem 5.5) ■

**Theorem 5.7.** *Let  $(\beta N, =_m)$  where  $x =_m y$  if  $x|_m y$  and  $y|_m x$ . Then*  
(a) *If  $x \in K(\beta N)$ , then  $[x]_m = K(\beta N)$ , and  $[x]_m$  has  $2^c$  elements.*  
(b)  *$=_m$  has one equivalence class for any  $x \in K(\beta N)$ .*

*Proof.* (a) By (Theorem 2.1 (a)) for any minimal left ideal  $L$  and any minimal right ideal  $R$  we have  $L \cap R \neq \phi$ . So there exist  $x$  such  $x \in L$  and  $x \in R$ . If  $x \in L$ , then for any  $y \in L$  we have  $y \in \beta N x$ , and  $x \in \beta N y$ , so  $x|_l y$  and  $y|_l x, x|_m y$  and  $y|_m x$ . Thus  $x =_m y$  for any  $y \in L$ . If  $x \in R$ , then for any  $z \in R$  we have  $z \in x\beta N$  and  $x \in z\beta N$ , so  $x|_R z$  and  $z|_R x, x|_m z$  and  $z|_m x$ . Thus  $x =_m z$  for any  $z \in R$ . If  $w \in \beta N, w \notin L$  and  $w \notin R$ , then  $w \notin \beta N x \beta N$ , and by (Lemma 5.4 (c)) we have  $w \neq_m x$ . So  $[x]_m = K(\beta N)$ . Since  $K(\beta N)$  has  $2^c$  elements, so  $[x]_m$  has  $2^c$  elements.  
(b) By (a) we have one equivalence class such that  $|[x]_m| = 2^c$  for any  $x \in K(\beta N)$ . ■

## 6 Closure of $K(\beta N)$ in the semigroup $(\beta N, \cdot)$

**Definition 6.1.** Consider the semigroup  $(N, \cdot)$ . Then

- (a)([1] Definition 4.45) A set  $A \subseteq N$  is "thick" if the family  $\{A/n : n \in N\}$  has the finite intersection property.  
(b)([1] Definition 4.38(a)) A set  $A \subseteq N$  is "syndetic" if there is a finite subset  $E$  of  $N$  ( $E \in \mathcal{P}_f(N)$ ) such that  $\bigcup_{n \in E} A/n = N$ .  
(c)([1] Definition 4.38(b)) A set  $A \subseteq N$  is piecewise syndetic if there is a

finite subset  $E$  of  $N$  ( $E \in \mathcal{P}_f(N)$ ) such that the set  $\bigcup_{n \in E} A/n$  is thick in  $N$  i.e  $\{(\bigcup_{n \in E} A/n)/m : m \in N\}$  has the finite intersection property.

**Theorem 6.2.** *Let  $A \subseteq N$  . Then*

(a)([1] Theorem 4.48(a))  *$A$  is thick if and only if there is a left ideal of  $\beta N$  contained in  $\bar{A}$  .*

(b)([1] Theorem 4.48(b))  *$A$  is syndetic if and only if for every left ideal  $L$  of  $\beta N$ ,  $L \cap \bar{A} \neq \phi$  .*

(c)([1] Theorem 4.40)  *$A$  is piecewise syndetic if and only if  $\bar{A} \cap K(\beta N) \neq \phi$ .*

**Lemma 6.3.** ([1] 4.46)

(a) *Every thick set is piecewise syndetic .*

(b) *Every syndetic is piecewise syndetic.*

**Example 6.4.** (1)  $N - L_i$  is thick for all  $i = 0, 1, \dots$  since for any minimal left ideal  $L$  of  $\beta N$  we have  $L \subseteq \overline{N - L_i}$  . Thus by (Theorem 6.2(a))  $N - L_i$  is thick.

(2)  $N - L_i$  is syndetic for all  $i = 0, 1, \dots$  . Since  $\overline{N - L_i} \cap L \neq \phi$  for every left ideal  $L$  of  $\beta N$  . Then by (Theorem 6.2(b))  $N - L_i$  is syndetic.

(3)  $N - L_i$  is piecewise syndetic.

**Lemma 6.5.** (a)  *$A \subseteq N$  is syndetic if and only if  $N - A$  is not thick*

(b)  *$A \subseteq N$  is thick if and only if  $N - A$  is not syndetic*

**Lemma 6.6.** *Let  $\mu$  be the set of all upward closed subsets of  $N$  and  $\nu$  be the set of all downward closed subsets of  $N$ . Then*

(a) *Any  $A \in \mu$  is thick*

(b) *Any  $A \in \nu$  is not thick*

(c) *Any  $A \in \mu$  is syndetic*

(d) *Any  $A \in \nu$  is not syndetic*

*Proof.* (a) Let  $A \in \mu$ .

$A/2 = \{m \in N : 2m \in A\}$ , and  $A/2 \uparrow = \{n \in N : \exists a \in A/2, a|n\}$ .

Let  $n \in A/2 \uparrow$ , so  $n = ra$  for some  $r \in N$  and  $a \in A/2$ , so  $2a \in A$ . Since  $A \in \mu$  then  $r(2a) \in A$  ( $2a \mid r(2a)$ ), so  $2(ra) \in A$ , and  $ra \in A/2$ . Thus  $n \in A/2$ , and since  $A/2 \subseteq A/2 \uparrow$ . So  $A/2 = A/2 \uparrow$  i.e  $A/2 \in \mu$

In general. Let  $A \in \mu$ .

$A/n = \{m \in N : mn \in A\}$ , and  $A/n \uparrow = \{r \in N : \exists a \in A/n, a|r\}$ .

Let  $r \in A/n \uparrow$ , so  $r = ta$  for some  $t \in N$  and  $a \in A/n$ , so  $na \in A$ . Since

$A$  is upward closed, then  $t(na) \in A (na|t(na))$  , so  $n(ta) \in A$  , and  $ta \in A/n$  . Thus  $r \in A/n$ , and since  $A/n \subseteq A/n \uparrow$  , then  $A/n = A/n \uparrow$  , so  $A/n \in \mu$  . Therefore  $\{A/n : n \in N\} \subseteq \mu$  . And since  $\mu$  has *F.I.P* , so  $\{A/n : n \in N\}$  has *F.I.P* , i.e  $A$  is thick

(b) Let  $A \in \nu, A/m = \{r \in N : rm \in A\}$ . If  $m \notin A$  then  $A/m = \{r \in N : rm \in A\} = \phi$  , because if  $rm \in A$  , and since  $m|rm$  , then  $m \in A$  a contradiction . So, if  $A \in \nu$  , then  $\{A/n : n \in N\}$  does not have *F.I.P* . Therefore  $A$  is not thick

(c) Let  $A \in \mu$  , we have  $N - A \in \nu$  and by (b)  $N - A$  is not thick . Thus by (Lemma 6.5(a))  $A$  is syndetic .

(d) Let  $A \in \nu$  , then we have  $N - A \in \mu$  , and by (a)  $N - A$  is thick . Thus by (Lemma 6.5 (b))  $A$  is not syndetic . ■

**Definition 6.7.** The set of ultrafilters which are generated by the set of upward closed subsets of  $N$  is denoted by  $\hat{\mu}$  and it is given by  $\hat{\mu} = \bigcap_{A \in \mu} \bar{A}$

i.e  $\hat{\mu} = \{x \in \beta N : x \in \bar{A}, A \in \mu\}$ .

By (Lemma 3.2) the set of ultrafilters which belong to  $\hat{\mu}$  are divided by all ultrafilters of  $\beta N$  under  $\bar{}$ - divisibility

**Lemma 6.8.** (a) For any ultrafilter  $x \in \beta N$  such that  $x \in \hat{\mu}, L_i \notin x (x \notin \bar{L}_i)$  for all  $i = 0, 1, 2, \dots$

(b)  $x \in \mathcal{I}$  for any  $x \in \hat{\mu} (\hat{\mu} \subset \mathcal{I})$

(c) There are  $2^c$  ultrafilters  $x$  such that  $x \in \mathcal{I}, x \notin \hat{\mu}$

*Proof.* (a) Let  $x \in \beta N, x \in \hat{\mu}$  . First we need to prove that any subset  $A \subseteq N, A \in x$  where  $x \in \hat{\mu}$  is syndetic. For any element  $A \in x$  we have three cases:  $A \in \mu$  or  $A$  is a finite intersection of elements of  $\mu$  or  $A$  contains some elements from  $\mu$  .

(1) If  $A \in \mu$  then by (Lemma6.6(c))  $A$  is syndetic

(2) If  $A$  is finite intersection of elements of  $\mu$  , then  $A \in \mu$  and  $A$  is syndetic.

(3) If  $B \subset A, B \in \mu$  , since  $B$  is syndetic , then by (Theorem 6.2 (b) )  $L \cap \bar{B} \neq \phi$  for any left ideal  $L$  of  $\beta N$  , so  $L \cap \bar{A} \neq \phi$  . Thus  $A$  is syndetic .

We know, by (Example 6-4)  $L_i$  is not syndetic for any  $L_i, i = 0, 1, 2, \dots$ . Therefore  $L_i \notin x (x \in \bar{L}_i)$

(b) Let  $x \in \beta N$  , and  $x \in \hat{\mu}$  by (a)  $L_i \notin x$  for all  $i = 0, 1, \dots$ , so  $N - L_i \in x, x \in \overline{N - L_i}$ . Therefore  $x \in \bigcap_{i=0}^{\infty} \overline{N - L_i}$  , Thus  $x \in \mathcal{I}$  and  $\hat{\mu} \subset \mathcal{I}$

(c) Let  $A = \{n_1, n_2, n_3, \dots\}$  such that  $n_i \in L_i, i \geq 1, n_i$  are odd number,

so by ([6] Theorem 3.3 )  $\bar{A}$  contains  $2^c$  ultrafilters  $x$  and since there is no nonprincipal ultrafilters  $y$  such that  $y \in \bar{A}$  and  $y \in \bar{L}_i$  for all  $i = 1, \dots$ . Thus  $x \in \mathcal{I}$  . And let  $B = \{2, 4, 6, \dots\}$  , then  $B \in \mu$  and  $B \notin x$  for any  $x \in \bar{A}$  ( $A \cap B = \phi$ ). Thus  $x \notin \hat{\mu}$  for any  $x \in \bar{A}$ . ■

**Lemma 6.9.** (a)  $\hat{\mu}$  is a left ideal in the semigroup  $(\beta N, \cdot)$

(b)  $\hat{\mu}$  is a right ideal in the semigroup  $(\beta N, \cdot)$

(c)  $\hat{\mu}$  is an ideal in the semigroup  $(\beta N, \cdot)$  .

(d)  $K(\beta N) \subset \hat{\mu}$

(e)  $|\hat{\mu}| = 2^c$

*Proof.* (a) Let  $x \in \hat{\mu}$  and  $y \in \beta N$  . For any  $A \in \mu$  we have  $A \in x$  , and for any  $B \in y$  by ( Lemma 1.1 (2) ) we have  $yx \in \bar{B}\bar{A} \subseteq \overline{AB}$  , so  $AB \in yx$ . Since  $A \in \mu$  , then for any  $ba \in BA$  we have  $ba \in A$  , so  $BA \subseteq A$  . Thus  $A \in yx$  for any  $A \in \mu$  . Therefore  $yx \in \hat{\mu}$  .

(b) Similar to (a)

(c) By (a) and (b)

(d) By (c)

(e) Since  $|K(\beta N)| = 2^c$  , then by (d) we have  $|\hat{\mu}| = 2^c$ . ■

**Lemma 6.10.** (a) Let  $(\beta N, =_{\sim})$  where  $x =_{\sim} y$  if  $x \tilde{|} y$  and  $y \tilde{|} x$  . Then  $[x]_{\sim} = \hat{\mu}$  for any  $x \in \hat{\mu}$

(b)  $K(\beta N) \neq \hat{\mu}$

*Proof.* (a) Let  $x \in \hat{\mu}$  , then for any  $y \in \hat{\mu}$  we have  $x \cap \mu \subseteq y$  and  $y \cap \mu \subseteq x$ , so by (Lemma 3.2 ) we have  $x \tilde{|} y$  and  $y \tilde{|} x$  . So  $x =_{\sim} y$  for any  $y \in \hat{\mu}$  . If  $y \notin \hat{\mu}$ , then  $x \cap \mu \not\subseteq y$  . So  $x \not\tilde{|} y$ . Therefore  $[x]_{\sim} = \hat{\mu}$  for any  $x \in \hat{\mu}$

(b) Since for any  $x \in k(\beta N)$ , by (Theorem 5.7 (a)) we have  $[x]_m = K(\beta N)$ , and by (a) we have  $[x]_{\sim} = \hat{\mu}$  , and since  $|_m \subset \tilde{|}$  , then  $[x]_m \subset [x]_{\sim}$  . Thus  $K(\beta N) \neq \hat{\mu}$  ■

**Corollary 6.11.**  $\hat{\mu}$  is closed subset of  $\beta N$ .

*Proof.* By (Lemma 3.3 (b))  $[x]_{\sim}$  is closed and compact for any  $x \in \beta N$  . Then by (Lemma 6.10)  $\hat{\mu}$  is closed subset of  $\beta N$ . ■

**Lemma 6.12.** Any subset  $A \subseteq N$  such that  $A \in x$  where  $x \in \hat{\mu}$  is piecewise syndetic

*Proof.* Let  $A \subseteq N$  such that  $A \in x, x \in \hat{\mu}$  . Analogues to proof of ( Lemma 6.8(a) ) we have  $A$  is syndetic , and by (Lemma 6-3)  $A$  is piecewise syndetic. ■

**Theorem 6.13.** (a) If  $x \in \hat{\mu}$ . Then  $x \in Cl(K(\beta N))$ .  
 (b)  $Cl(K(\beta N)) = \hat{\mu}$ .  
 (c)  $=_{\sim}$  has one equivalence class for any  $x \in Cl(K(\beta N))$ .

*Proof.* Let  $x \in \hat{\mu}$  and let  $U$  be any open subset of  $\beta N$  such that  $x \in U$ . So there exist a basic open subset  $\bar{A}$  of  $\beta N$  such that  $x \in \bar{A} \subset U$ , and since  $A \in x$  then by (Lemma 6.12)  $A$  is piecewise syndetic. So by (Theorem 6-2) we have  $U \cap K(\beta N) \neq \emptyset$ . Thus  $x \in Cl(K(\beta N))$ .

(b) By (Lemma 6.9 (d)) we have  $K(\beta N) \subseteq \hat{\mu}$ , so  $Cl(K(\beta N)) \subseteq CL(\hat{\mu})$  and by (Corollary 6.11) we have  $Cl(K(\beta N)) \subseteq \hat{\mu}$ . Since by (a)  $\hat{\mu} \subseteq Cl(K(\beta N))$ . Thus  $Cl(K(\beta N)) = \hat{\mu}$ .

(c) By (b) and ( Lemma 6.10(a) ) ■

## References

- [1] N. Hindman, D. Strauss: Algebra in the Stone-Cech compactification, theory and applications. 2nd revised and extended, De Gruyter, 2012.
- [2] S. Khalifa. Levels of ultrafilters with extension divisibilities, Math.Logic, submitted.
- [3] B. Sobot, Divisibility in the Stone-Cech compactification, Rep. Math. Logic 50(2015), 53-66.
- [4] B. Sobot, Divisibility orders in the Stone-Cech compactification, Math.Logic(2016).
- [5] B. Sobot,  $\tilde{}$ -divisibility of ultrafilters, Math.Logic, submitted.
- [6] R. Walker, The Cech-Stone compactification. Springer, 1974