A Schur-Toponogov theorem in Riemannian geometry & a new proof of Toponogov's theorem in Alexandrov geometry

Yusheng Wang¹

Abstract. In the paper, we give a Schur-Toponogov theorem in Riemannian geometry, which not only generalizes Schur's and Toponogov's theorem but also indicates their relation. Inspired by its proof, we also supply a new proof of Toponogov's theorem (in the large) in Alexandrov geometry.

Key words. Schur's theorem, Toponogov's theorem, Alexandrov geometry Mathematics Subject Classification (2000): 53-C20.

0 Introduction

In Rimannian geometry, a kind of elegant geometry is on distance comparison by curvature; for instance, Shcur's theorem ([Ho]). In the present paper, we denote by [pq] a minimal geodesic between p and q in a Riemannian manifold or Alexandrov space.

Theorem 0.1 (A. Shcur) Let $\gamma(s)|_{[0,L]} \subset \mathbb{E}^n$ (n-dimensional Euclidean space) and $\tilde{\gamma}(s)|_{[0,L]} \subset \mathbb{E}^2$ be two arc-length parameterized C^2 -curves, and let $\kappa(s), \tilde{\kappa}(s)$ be their absolute curvature respectively. If $\kappa(s) \leq \tilde{\kappa}(s)$ for all s, and if $\tilde{\gamma}(s)|_{[0,L]} \cup [\tilde{\gamma}(0)\tilde{\gamma}(L)]$ is a convex curve, then the distance

 $|\gamma(0)\gamma(L)| \ge |\tilde{\gamma}(0)\tilde{\gamma}(L)|;$

and if equality holds, then $\gamma(s)|_{[0,L]}$ is equal to $\tilde{\gamma}(s)|_{[0,L]}$ up to an isometry of \mathbb{E}^n .

Recall that $\kappa(s) = |D_{\dot{\gamma}(s)}\dot{\gamma}(s)|$, where $\dot{\gamma}(s)$ denotes the tangent vector of $\gamma(s)$ and D is the canonical connection of \mathbb{E}^n . Theorem 0.1 also holds for piecewise C^2 -curves ([Ho], cf. [Su]), i.e., $\gamma(s)$ and $\tilde{\gamma}(s)$ are not differential at only a finite number of points, and at such a point ' $\kappa(s) \leq \tilde{\kappa}(s)$ ' means that $|\dot{\gamma}_{-}(s)\dot{\gamma}_{+}(s)| \leq |\dot{\tilde{\gamma}}_{-}(s)\dot{\tilde{\gamma}}_{+}(s)|$ (where $\dot{\gamma}_{\pm}(s)$ denotes the right and left tangent vectors).

We now consider another significant result on distance comparison—Toponogov's theorem ([BGP], [Pe], [GM]). In the paper, we always denote by \mathbb{S}_k^n the complete and simply connected *n*-dimensional Riemannian manifold with constant sectional curvature k. For a given [pq] in a Riemannian manifold (or Alexandrov space), we denote by \uparrow_p^q the direction from p to q of [pq] (in a Riemannian manifold, \uparrow_p^q is just the unit tangent vector of [pq] at p).

¹Supported by NSFC 11471039.

Theorem 0.2 (Toponogov) Let M^n be a complete Riemannian manifold with $\sec_M \ge k$. *k.* Let $p \in M$ and $[qr] \subset M$, and let $\tilde{p} \in \mathbb{S}^2_k$ and $[\tilde{q}\tilde{r}] \subset \mathbb{S}^2_k$ such that $|\tilde{p}\tilde{q}| = |pq|$, $|\tilde{q}\tilde{r}| = |qr|$, and $|pq| + |pr| + |qr| < \frac{2\pi}{\sqrt{k}}$ for k > 0. Then the following holds: (0.2.1) If $|\tilde{p}\tilde{r}| = |pr|$, then for $s \in [qr]$ and $\tilde{s} \in [\tilde{q}\tilde{r}]$ with $|qs| = |\tilde{q}\tilde{s}|$,

$$|ps| \ge |\tilde{p}\tilde{s}|. \tag{0.1}$$

(0.2.2) If $|\uparrow_q^p\uparrow_q^r| = |\uparrow_{\tilde{q}}^{\tilde{p}}\uparrow_{\tilde{q}}^{\tilde{r}}|$ for some [pq], then for $s \in [qr]$ and $\tilde{s} \in [\tilde{q}\tilde{r}]$ with $|qs| = |\tilde{q}\tilde{s}|$,

$$|ps| \le |\tilde{p}\tilde{s}|. \tag{0.2}$$

(0.2.3) If the equality in (0.1) (resp. (0.2)) holds for some $s \in [qr]^{\circ 2}$ (resp. $s \in [qr] \setminus \{q\}$), then there is [pq] and [pr] (resp. [ps]) which together with [qr] (resp. [pq] and [qs]) bounds a convex surface which can be isometrically embedded into \mathbb{S}_k^2 .

Remark 0.3 (0.3.1) In Theorem 0.2, if k > 0, then $|pq| + |pr| + |qr| \le \frac{2\pi}{\sqrt{k}}$, and equality holds if and only if M^n is isometric to \mathbb{S}^n_k with p, q, r lying in a great circle (cf. [BGP]). (0.3.2) It is not hard to see that (0.2.1) is equivalent to (0.2.2), and (0.2.2) is equivalent to that, in (0.2.1), $|\uparrow_q^p\uparrow_q^r| \ge |\uparrow_{\tilde{q}}^{\tilde{p}}\uparrow_{\tilde{q}}^{\tilde{r}}|$ for any [pq].

(0.3.3) Theorem 0.2 has a corresponding version for $\sec_M \leq k$ with inverse inequalities, but it requires that $[qr] \subset M \setminus C(p)$, where C(p) denotes the cut locus of p (cf. [Me]). If $[qr] \subset M \setminus C(p)$, the corresponding ' $|\uparrow_q^p \uparrow_q^r| \leq |\uparrow_{\tilde{q}}^{\tilde{p}} \uparrow_{\tilde{q}}^{\tilde{r}}|$ ' (see (0.3.2)) is an immediate corollary of Rauch's first comparison theorem (but this way does not work if $\sec_M \geq k$).

We now state our main result, a Schur-Toponogov theorem, which unifies and generalizes Theorem 0.1, 0.2 and (0.3.3).

Theorem A Let M be a complete Riemannian manifold with $\sec_M \leq k$ (resp. $\sec_M \geq k$) and let $\gamma(s)|_{[0,L]} \subset M$ and $\tilde{\gamma}(s)|_{[0,L]} \subset \mathbb{S}^2_k$ be two arc-length parameterized C^2 -curves. Suppose that the absolute curvature of $\gamma(s)$ and $\tilde{\gamma}(s)$, $\kappa(s)$ and $\tilde{\kappa}(s)$, satisfy that $\kappa(s) \leq \tilde{\kappa}(s)$ (resp. $\kappa(s) \geq \tilde{\kappa}(s)$) for all s. Then the following holds:

(A1) If $\tilde{\gamma}(s)|_{[0,L]} \cup [\tilde{\gamma}(0)\tilde{\gamma}(L)]$ is a convex curve and $\gamma(s)|_{[0,L]} \subset M \setminus C(\gamma(0))$ (resp. if $\gamma(s)|_{[0,L]}$ is convex to $\gamma(0)$), then

$$|\gamma(0)\gamma(L)| \ge (resp. \le) |\tilde{\gamma}(0)\tilde{\gamma}(L)|; \tag{0.3}$$

and if equality holds and $|\gamma(0)\gamma(L)| + L < \frac{2\pi}{\sqrt{k}}$ for k > 0, then (resp. then there is $[\gamma(0)\gamma(s)]$ such that) $\bigcup_{s\in[0,L]}[\gamma(0)\gamma(s)]$ with induced metric is isometric to $\bigcup_{s\in[0,L]}[\tilde{\gamma}(0)\tilde{\gamma}(s)]$. (A2) Let $p \in M, \tilde{p} \in \mathbb{S}^2_k$ with $|p\gamma(0)| = |\tilde{p}\tilde{\gamma}(0)|$, and let $|p\gamma(0)| + |p\gamma(L)| + L < \frac{2\pi}{\sqrt{k}}$ for k > 0. If $[\tilde{p}\tilde{\gamma}(0)] \cup \tilde{\gamma}(s)|_{[0,L]} \cup [\tilde{p}\tilde{\gamma}(L)]$ is a convex curve and $\gamma(s)|_{[0,L]} \subset M \setminus C(p)$ (resp. if $\gamma(s)|_{[0,L]}$ is convex to p), then the following holds:

² $[qr]^{\circ}$ denotes the interior part of [qr].

(A2.1) If $|p\gamma(L)| = |\tilde{p}\tilde{\gamma}(L)|$, then for $s \in (0, L)$

$$|p\gamma(s)| \le (resp. \ge) |\tilde{p}\tilde{\gamma}(s)|. \tag{0.4}$$

(A2.2) If (resp. If for some $[p\gamma(0)]$) $|\uparrow_{\gamma(0)}^{p} \dot{\gamma}(0)| = |\uparrow_{\tilde{\gamma}(0)}^{\tilde{p}} \dot{\tilde{\gamma}}(0)|$, then for $s \in (0, L]$

$$|p\gamma(s)| \ge (resp. \le) |\tilde{p}\tilde{\gamma}(s)|. \tag{0.5}$$

(A2.3) If the equality in (0.4) or (0.5) holds for some $s \in (0, L)$ or $s_0 \in (0, L]$ respectively, then (resp. then there is $[p\gamma(s)]$ such that) $\bigcup_{s\in[0,L]} [p\gamma(s)]$ or $\bigcup_{s\in[0,s_0]} [p\gamma(s)]$ with induced metric is isometric to $\bigcup_{s\in[0,L]} [\tilde{p}\tilde{\gamma}(s)]$ or $\bigcup_{s\in[0,s_0]} [\tilde{p}\tilde{\gamma}(s)]$ respectively.

In Theorem A, $\kappa(s)$ (the absolute curvature of $\gamma(s)$) is still defined to be $|D_{\dot{\gamma}(s)}\dot{\gamma}(s)|$, where D is the Levi-Civita connection of M. And the definition of a convex curve in \mathbb{S}_k^2 is the same as in \mathbb{E}^2 , while the notion $\gamma(s)|_{[0,L]}$ is convex to p (or $\gamma(0)$)' will be introduced in Section 2 (see Definition 2.1).

Remark 0.4 (0.4.1) From our proof, one can see that (A2.1) is equivalent to (A2.2), and (A2.2) is equivalent to that, in (A2.1), $|\uparrow_{\gamma(0)}^{p} \dot{\gamma}(0)| \leq (\text{resp.} \geq) |\uparrow_{\tilde{\gamma}(0)}^{\tilde{p}} \dot{\tilde{\gamma}}(0)|$ for any $[p\gamma(0)]$ (cf. (0.3.2)).

(0.4.2) From our proof, (A1) can be viewed as a corollary of (A2), but we formulate (A1) and (A2) in a theorem because the proof of (A2.3) requires (0.3).

Remark 0.5 (0.5.1) It is clear that (A1) of Theorem A includes Theorem 0.1. And (A2) for $\sec_M \geq k$ includes Theorem 0.2 (see Remark 2.4, where a key point is that if $\gamma(s)|_{[0,L]}$ is a minimal geodesic (so is $\tilde{\gamma}(s)|_{[0,L]}$), then it is convex to p automatically). We would like to point out that Theorem 0.1 and 0.2 are not used in our proof; in other words, we supply a new proof for Theorem 0.1 and 0.2 in this paper.

(0.5.2) One can find counterexamples for Theorem A without the condition $\gamma(s)|_{[0,L]} \subset M \setminus C(\gamma(0))$ or $\gamma(s)|_{[0,L]} \subset M \setminus C(p)$ (resp. $|\gamma(0)\gamma(L)| + L < \frac{2\pi}{\sqrt{k}}$ or $|p\gamma(0)| + |p\gamma(L)| + L < \frac{2\pi}{\sqrt{k}}$ for k > 0) when M is a cylinder (resp. \mathbb{S}_k^2).

(0.5.3) Similar to Theorem 0.1, Theorem A also holds for piecewise C^2 -curves $\gamma(s)$ and $\tilde{\gamma}(s)$ (in fact, our proof works for such general case), and at each non-differential point ' $\kappa(s) \leq (\geq) \tilde{\kappa}(s)$ ' means ' $|\dot{\gamma}_{-}(s)\dot{\gamma}_{+}(s)| \leq (\geq) |\dot{\tilde{\gamma}}_{-}(s)\dot{\tilde{\gamma}}_{+}(s)|$ '.

Toponogov's theorem is not only a powerful tool in Riemannian geometry, but also the base of Alexandrov geometry. An Alexandrov space X with curvature $\geq k$ is roughly defined to be a locally compact length space ³ on which Theorem 0.2 without the rigidity part (0.2.3) holds locally ([BGP]). In fact, if X is complete, then Theorem 0.2 holds globally on X, for which without the rigidity part there are three proofs so far ([BGP], [P1], [Sh]; refer to [GM] for a proof for the rigidity part). The proof in [BGP]

³If an Alexandrov space X is locally compact, then for any $p \in X$ there is a neighborhood U of p such that between any two points in U there is a shortest path (geodesic). In [BGP], the original definition of Alexandrov spaces is for locally complete spaces.

can be viewed as a version corresponding to the original definition of Alexandrov spaces with curvature $\geq k$ in [BGP] (which also adapts to locally complete spaces), while the proofs in [Pl] and [Sh] can be viewed corresponding to (0.2.2) in Theorem 0.2. Inspired by our proof of Theorem A (especially Lemma 1.2), we supply a new proof which is a version corresponding to (0.2.1) in Theorem 0.2, i.e. we will prove:

Theorem B Let X be a complete Alexandrov space with curvature $\geq k$. Let $p \in X$ and $[qr] \subset X$, and let $\tilde{p} \in \mathbb{S}_k^2$ and $[\tilde{q}\tilde{r}] \subset \mathbb{S}_k^2$ such that $|\tilde{p}\tilde{q}| = |pq|, |\tilde{p}\tilde{r}| = |pr|, |\tilde{q}\tilde{r}| = |qr|$. Suppose that $|pq| + |pr| + |qr| < \frac{2\pi}{\sqrt{k}}$ for k > 0. Then for any $s \in [qr]$ and $\tilde{s} \in [\tilde{q}\tilde{r}]$ with $|qs| = |\tilde{q}\tilde{s}|$, we have that $|ps| \geq |\tilde{p}\tilde{s}|$.

Similar to (0.3.1), if k > 0 in Theorem B, then one can show that $|pq| + |pr| + |qr| \le \frac{2\pi}{\sqrt{k}}$ ([BGP]).

1 Proof of Theorem A for the case where $sec_M \leq k$

1.1 Preparations

a. Closed convex curves in \mathbb{S}_k^2

Let $\tilde{\gamma}(s)|_{[0,L]}$ be an arc-length parameterized C^2 -curve in \mathbb{S}^2_k , and $p \in \mathbb{S}^2_k$. If $[\tilde{p}\tilde{\gamma}(0)] \cup \tilde{\gamma}(s)|_{[0,L]} \cup [\tilde{p}\tilde{\gamma}(L)]$ is a convex curve, then for any $\tilde{\gamma}(s)$ and $\tilde{q} \in [\tilde{p}\tilde{\gamma}(0)] \cup \tilde{\gamma}(s)|_{[0,L]} \cup [\tilde{p}\tilde{\gamma}(L)]$,

the angle between $D_{\dot{\tilde{\gamma}}(s)}\dot{\tilde{\gamma}}(s)$ and $[\tilde{\gamma}(s)\tilde{q}]$ at $\tilde{\gamma}(s)$ is not bigger than $\frac{\pi}{2}$. (1.1)

Moreover,

$$|\tilde{p}\tilde{\gamma}(0)| + |\tilde{p}\tilde{\gamma}(L)| + L \leq \frac{2\pi}{\sqrt{k}}$$
 for $k > 0$, and
'=' holds iff $[\tilde{p}\tilde{\gamma}(0)] \cup \tilde{\gamma}(s)|_{[0,L]} \cup [\tilde{p}\tilde{\gamma}(L)]$ is the union of two half great circles. (1.2)

In order to see (1.2), it suffices to show that a closed and convex piecewise minimal geodesic $\Gamma \subset \mathbb{S}_k^2$ (i.e., a convex polygon) is of length $\leq \frac{2\pi}{\sqrt{k}}$, and equality holds if and only if Γ is the union of two half great circles. Note that Γ is the boundary of the intersection of a finite number of half spheres. Then by induction on the number of these half spheres, one can see the wanted property of Γ .

b. Index forms of Jacobi fields

Let M be a complete Riemannian manifold, and let $c(t)|_{[a,b]} \subset M$ be a normal geodesic. Recall that for a normal ⁴ Jacobi field X(t) along c(t), the index form

$$I_a^b(X,X) \triangleq \int_a^b \left(|\dot{X}(t)|^2 - \langle R_{\dot{c}(t)X(t)}\dot{c}(t), X(t) \rangle \right) dt.$$

Let $p \in M$ and $\gamma(s)|_{(-\epsilon,\epsilon)} \subset M$ be an arc-length parameterized C^2 -curve. If $\gamma(0) \notin C(p)$, then there is a Jacobi field U(t) along the normal geodesic $c(t) \triangleq [p\gamma(0)]$ with U(0) = 0 and $U(|p\gamma(0)|) = \dot{\gamma}(0)$ such that

$$\frac{d|p\gamma(s)|}{ds}|_{s=0} = -\cos|\dot{\gamma}(0)\uparrow^p_{\gamma(0)}|$$

⁴Here, 'normal' means that $X(t) \perp \dot{c}(t)$

and

$$\frac{d^2|p\gamma(s)|}{d^2s}|_{s=0} = \left\langle D_{U(t)}U(t), \dot{c}(t) \right\rangle|_0^{|p\gamma(0)|} + I_0^{|p\gamma(0)|}(U^{\perp}, U^{\perp}), \tag{1.3}$$

where $U^{\perp}(t)$ is the projection of U(t) to the orthogonal space of $\dot{c}(t)$ in $T_{c(t)}M$.

c. Index form comparison

On index forms, we have the following comparison result (cf. [Wu], [CE]), which plays an essential role in proving Rauch's first comparison theorem.

Lemma 1.1 Let M and \tilde{M} be two complete n-dimensional Riemannian manifolds, and let $c(t)|_{[0,\ell]} \subset M$ and $\tilde{c}(t)|_{[0,\ell]} \subset \tilde{M}$ be two normal minimal geodesics. Let $K_{\max}(t)$ (resp. $\tilde{K}_{\min}(t)$) be the maximum (resp. the minimum) of sectional curvatures at c(t) (resp. $\tilde{c}(t)$). And let J(t) and $\tilde{J}(t)$ be normal Jacobi fields along c(t) and $\tilde{c}(t)$ respectively. If $K_{\max}(t) \leq \tilde{K}_{\min}(t), J(0) = \tilde{J}(0) = 0$ and $|J(\ell)| = |\tilde{J}(\ell)|$, then

$$I_0^\ell(J,J) \ge I_0^\ell(\tilde{J},\tilde{J}).$$

Moreover, if equality holds and if $\tilde{M} = \mathbb{S}_k^n$, then $K(\dot{c}(t), J(t)) \equiv k$ (where $K(\dot{c}(t), J(t))$) is the sectional curvature of the plane spanned by $\dot{c}(t)$ and J(t)).

d. Two easy facts on distance comparison

Lemma 1.2 Let M and \tilde{M} be two complete Riemannian manifolds, let $p \in M$ and $\tilde{p} \in \tilde{M}$, and let $c(t)|_{[0,\ell]} \subset M$ and $\tilde{c}(t)|_{[0,\ell]} \subset \tilde{M}$ be two c^1 -curves such that $|pc(0)| = |\tilde{p}\tilde{c}(0)|$ and $|pc(\ell)| = |\tilde{p}\tilde{c}(\ell)|$. If $|pc(t)| - |\tilde{p}\tilde{c}(t)|$ attains a local minimum at $t_0 \in (0,\ell)$ and there is a unique minimal geodesic between \tilde{p} and $\tilde{c}(t_0)$, then

$$|\dot{c}(t_0)\uparrow^p_{c(t_0)}| = |\dot{\tilde{c}}(t_0)\uparrow^p_{\tilde{c}(t_0)}|.$$

Proof. This is an almost immediate corollary of the first variation formula. \Box

Lemma 1.3 Let $[p_1p_2] \subset \mathbb{S}_k^2$ with $|p_1p_2| < \frac{\pi}{\sqrt{\kappa}}$ for $\kappa > 0$, and let $q \in [p_1p_2]^\circ$ and $c(t)|_{[0,\ell]} \subset \mathbb{S}_k^2$ be a normal minimal geodesic with c(0) = q and $\ell \leq \frac{\pi}{\sqrt{\kappa}}$ for $\kappa > 0$. Then $|p_1c(t)| + |p_2c(t)|$ is strictly increasing with respect to t. In particular, if $|qp_i|$ (i = 1 or 2) is sufficiently small compared to $|qc(t_0)|$, then $|p_ic(t)| < |p_ic(t_0)|$ for all $t \in [0, t_0)$.

Proof. It is not hard to see that the lemma follows from the Law of Cosine. \Box

1.2 Proof of Theorem A for the case where $\sec_M \leq k$

In this subsection, we always assume that $\sec_M \leq k$. And we remark that the following proof goes through for piecewise C^2 -curves $\gamma(s)$ and $\tilde{\gamma}(s)$ as mentioned in (0.5.3).

Step 1. To prove (A2.1).

(This step is the essential part of the whole proof for Theorem A.)

Observe that $|p\gamma(0)| + |p\gamma(L)| + L < \frac{2\pi}{\sqrt{k}}$ for k > 0 implies that $|p\gamma(s)| < \frac{\pi}{\sqrt{k}}$ for all s. It follows that for any $\bar{s} \in [0, L]$ there is a C^2 -curve $\tilde{\gamma}_{\bar{s}}(s)|_{[0,L]} \subset \mathbb{S}^2_k$, equal to $\tilde{\gamma}(s)|_{[0,L]}$ up to an isometry of \mathbb{S}^2_k , such that

$$|p\gamma(\bar{s})| = |\tilde{p}\tilde{\gamma}_{\bar{s}}(\bar{s})| \text{ and } |\uparrow^{p}_{\gamma(\bar{s})}\dot{\gamma}(\bar{s})| = |\uparrow^{\tilde{p}}_{\tilde{\gamma}_{\bar{s}}(\bar{s})}\dot{\tilde{\gamma}}_{\bar{s}}(\bar{s})|.$$
(1.4)

Since $\gamma(s)|_{[0,L]} \subset M \setminus C(p)$, $\gamma(s)|_{[0,L]}$ (and $\tilde{\gamma}_{\bar{s}}(s)|_{[0,L]}$ similarly) determines a Jacobi field $U_s(t)|_{[0,|p\gamma(s)|]}$ along $\beta_s(t)|_{[0,|p\gamma(s)|]} \triangleq [p\gamma(s)]$ such that $U_s(|p\gamma(s)|) = \dot{\gamma}(s)$. Then putting (1.1), (1.4), Lemma 1.1 and (1.3) together, we can conclude that

$$\frac{d^2 |p\gamma(s)|}{d^2 s}\Big|_{s=\bar{s}} \ge \frac{d^2 |\tilde{p}\tilde{\gamma}_{\bar{s}}(s)|}{d^2 s}\Big|_{s=\bar{s}};$$
(1.5)

and equality holds if and only if

either
$$|\uparrow_{\gamma(\bar{s})}^{p} \dot{\gamma}(\bar{s})| = 0 \text{ or } \pi,$$
 (1.6)

or
$$\kappa(\bar{s}) = \tilde{\kappa}(\bar{s})$$
 and the angle between $D_{\dot{\gamma}(\bar{s})}\dot{\gamma}(\bar{s})$ and $\uparrow^p_{\gamma(\bar{s})}$ is equal to

that between
$$D_{\hat{\gamma}_{\bar{s}}(\bar{s})} \dot{\tilde{\gamma}}_{\bar{s}}(\bar{s})$$
 and $\uparrow_{\hat{\gamma}_{\bar{s}}(\bar{s})}^{\tilde{p}}$. (1.7)

Moreover, when (1.7) occurs, $D_{\dot{\gamma}(\bar{s})}\dot{\gamma}(\bar{s})$ lies in the plane spanned by $\dot{\gamma}(\bar{s})$ and $\uparrow_{\gamma(\bar{s})}^{p}$ in $T_{\gamma(\bar{s})}M$, and the sectional curvature of the plane spanned by $\dot{\beta}_{\bar{s}}(t)$ and $U_{\bar{s}}(t)$ in $T_{\beta_{\bar{s}}(t)}M$

$$K(\bar{s},t) = \kappa \text{ for all } t \in [0, |p\gamma(\bar{s})|]$$
(1.8).

Then we will finish Step 1 through the following three substeps.

Substep 1. To give a proof for the case where [0, L] is one kind of "best" intervals, i.e.: the ' \geq ' in (1.5) is '>' for all \bar{s} , and for any $s_1 \neq s_2 \in [0, L]$ there is a C^2 -curve $\tilde{\gamma}_{s_1,s_2}(s)|_{[0,L]} \subset \mathbb{S}^2_k$, which is equal to $\tilde{\gamma}(s)|_{[0,L]}$ up to an isometry of \mathbb{S}^2_k , such that

$$|p\gamma(s_i)| = |\tilde{p}\tilde{\gamma}_{s_1,s_2}(s_i)| \tag{1.9}$$

and

$$[\tilde{p}\tilde{\gamma}_{s_1,s_2}(s_1)] \cup \tilde{\gamma}_{s_1,s_2}(s)|_{[s_1,s_2]} \cup [\tilde{p}\tilde{\gamma}_{s_1,s_2}(s_2)] \text{ is convex.}$$
(1.10)

(For example, [0, L] will be such a "best" interval if $\tilde{\gamma}(s)|_{[0,L]}$ (and so $\gamma(s)|_{[0,L]}$) is a minimal geodesic.)

We now assume that (A2.1) is not true for such a "best" case. Our strategy is to find some $[a, b] \subset [0, L]$ with b - a sufficiently small such that

$$|\uparrow_{\gamma(a)}^{p} \dot{\gamma}(a)| > |\uparrow_{\tilde{\gamma}_{a,b}(a)}^{\tilde{p}} \dot{\tilde{\gamma}}_{a,b}(a)| \text{ and } \frac{d^{2}|p\gamma(s)|}{d^{2}s}|_{[a,b]} > \frac{d^{2}|\tilde{p}\tilde{\gamma}_{a,b}(s)|}{d^{2}s}|_{[a,b]},$$
(1.11)

which contradicts to $|p\gamma(b)| = |\tilde{p}\tilde{\gamma}_{a,b}(b)|$ ' (see (1.9)).

By the assumption right above, we can let $s_0 \in (0, L)$ such that

$$|p\gamma(s_0)| - |\tilde{p}\tilde{\gamma}(s_0)| = \max_{s \in (0,L)} \{|p\gamma(s)| - |\tilde{p}\tilde{\gamma}(s)|\} > 0.$$

By Lemma 1.2,

$$|\uparrow_{\gamma(s_0)}^p \dot{\gamma}(s_0)| = |\uparrow_{\tilde{\gamma}(s_0)}^{\tilde{p}} \dot{\tilde{\gamma}}(s_0)|.$$
(1.12)

Note that we can prolong $[\tilde{\gamma}(s_0)\tilde{p}]$ to $[\tilde{\gamma}(s_0)\tilde{p}']$ such that $|\tilde{\gamma}(s_0)\tilde{p}'| = |\gamma(s_0)p|$. Note that $[\tilde{\gamma}(0)\tilde{\gamma}(L)] \cap [\tilde{\gamma}(s_0)\tilde{p}] \neq \emptyset$ due to the convexity of $[\tilde{p}\tilde{\gamma}(0)] \cup \tilde{\gamma}(s)|_{[0,L]} \cup [\tilde{p}\tilde{\gamma}(L)]$. Then by Lemma 1.3, at least one of $|\tilde{p}'\tilde{\gamma}(0)| > |p\gamma(0)|$ ' and $|\tilde{p}'\tilde{\gamma}(L)| > |p\gamma(L)|$ ' holds; moreover,

$$|\tilde{p}'\tilde{\gamma}(0)| > |p\gamma(0)| \text{ (resp. } |\tilde{p}'\tilde{\gamma}(L)| > |p\gamma(L)|) \text{ if } s_0 \text{ (resp. } L - s_0)$$

is sufficiently small. (1.13)

We now assume that $|\tilde{p}'\tilde{\gamma}(L)| > |p\gamma(L)|$, and consider the curve $\tilde{\gamma}_{s_0,L}(s)|_{[0,L]}$ (due to that [0,L] is "best"). $|\tilde{p}'\tilde{\gamma}(L)| > |p\gamma(L)|$ ' implies that $|\uparrow_{\tilde{\gamma}(s_0)}^{\tilde{p}} \dot{\tilde{\gamma}}(s_0)| > |\uparrow_{\tilde{\gamma}_{s_0,L}(s_0)}^{\tilde{p}} \dot{\tilde{\gamma}}_{s_0,L}(s_0)|$, so by (1.12)

$$|\uparrow_{\gamma(s_0)}^{p} \dot{\gamma}(s_0)| > |\uparrow_{\tilde{\gamma}_{s_0,L}(s_0)}^{\tilde{p}} \dot{\tilde{\gamma}}_{s_0,L}(s_0)|.$$
(1.14)

By (1.14) and the first variation formula, it is easy to see that there is $s'_0 \in (s_0, L)$ such that $|p\gamma(s'_0)| - |\tilde{p}\tilde{\gamma}_{s_0,L}(s'_0)| = \max_{s \in (s_0,L)}\{|p\gamma(s)| - |\tilde{p}\tilde{\gamma}_{s_0,L}(s)|\} > 0$. And consequently, we can set $[s_1, s'_1] \triangleq [s_0, s'_0]$ or $[s'_0, L]$ such that

$$|\uparrow_{\gamma(s_1)}^p \dot{\gamma}(s_1)| > |\uparrow_{\tilde{\gamma}_{s_1,s_1'}(s_1)}^{\tilde{p}} \dot{\tilde{\gamma}}_{s_1,s_1'}(s_1)| \text{ or } |\uparrow_{\gamma(s_1')}^p \dot{\gamma}(s_1')| < |\uparrow_{\tilde{\gamma}_{s_1,s_1'}(s_1')}^{\tilde{p}} \dot{\tilde{\gamma}}_{s_1,s_1'}(s_1')|.$$

By repeating the above process, we can get $[s_1, s'_1] \supseteq \cdots \supseteq [s_i, s'_i] \supseteq \cdots$ such that

$$|\uparrow_{\gamma(s_{i})}^{p} \dot{\gamma}(s_{i})| > |\uparrow_{\tilde{\gamma}_{s_{i}},s_{i}'}^{\tilde{p}}(s_{i})} \dot{\tilde{\gamma}}_{s_{i},s_{i}'}(s_{i})| \text{ or } |\uparrow_{\gamma(s_{i}')}^{p} \dot{\gamma}(s_{i}')| < |\uparrow_{\tilde{\gamma}_{s_{i}},s_{i}'}^{\tilde{p}}(s_{i}') \dot{\tilde{\gamma}}_{s_{i},s_{i}'}(s_{i}')|.$$
(1.15)

Moreover, by (1.13) we can select s_i and s'_i such that they converge to some \bar{s} as $i \to \infty$, and then we put $[a, b] \triangleq [s_i, s'_i]$ for a sufficiently large i.

We now need to check that [a, b] satisfies (1.11). Note that by (1.15) we can have the first inequality of (1.11). As for the second one, since we have assumed that the ' \geq ' in (1.5) is '>', it suffices to show that

$$\tilde{\gamma}_{s_i,s_i'}(s)|_{[0,L]}$$
 converges to $\tilde{\gamma}_{\bar{s}}(s)|_{[0,L]}$ as $i \to \infty$ (1.16)

(note that $\tilde{\gamma}_{s_i,s'_i}(s)|_{[0,L]}$ and $\tilde{\gamma}_{\bar{s}}(s)|_{[0,L]}$ are all equal to $\tilde{\gamma}(s)|_{[0,L]}$ up to isometries of \mathbb{S}^2_k). Observe that the smaller $|s_i - s'_i|$ is the smaller $\left||\uparrow^p_{\gamma(s_i)}\dot{\gamma}(s_i)| - |\uparrow^{\tilde{p}}_{\tilde{\gamma}_{s_i,s'_i}(s_i)}\dot{\tilde{\gamma}}_{s_i,s'_i}(s_i)|\right|$ is because both $\gamma(s)$ and $\tilde{\gamma}(s)$ are C^2 -curves, and thus (1.16) follows.

Substep 2. To give a proof for the case where [0, L] is another "best" interval, namely, the ' \geq ' in (1.5) is '=' for all \bar{s} and (1.6) does not occur at any \bar{s} .

Since $\gamma(s)|_{[0,L]} \subset M \setminus C(p)$ and $|p\gamma(s)||_{[0,L]} < \frac{\pi}{\sqrt{k}}$, we can draw a C^2 -curve $\bar{\tilde{\gamma}}(s)|_{[0,L]}$ in \mathbb{S}_k^2 with s being arc-length parameter such that $|p\gamma(s)| = |\tilde{p}\bar{\tilde{\gamma}}(s)|$ for all $s \in [0, L]$. Consider the corresponding (1.3) to $|\tilde{p}\bar{\tilde{\gamma}}(s)|$ together with (1.7) and (1.8), and notice that (1.6) does not occur at any \bar{s} in this substep. It has to hold that $\bar{\tilde{\gamma}}(s)|_{[0,L]}$ is equal to $\tilde{\gamma}(s)|_{[0,L]}$ up to an isometry of \mathbb{S}_k^2 , and thus $|p\gamma(s)| = |\tilde{p}\tilde{\gamma}(s)|$ for all $s \in [0, L]$.

Substep 3. To give a proof for general cases.

We first observe that (A2.1) holds if (1.6) occurs at all $\bar{s} \in [0, L]$ (which implies that (1.5) is an equality). In fact, in such a situation, $\gamma(s)|_{[0,L]}$ has to be a minimal geodesic with $[p\gamma(1)] = [p\gamma(0)] \cup \gamma(s)|_{[0,L]}$ or $[p\gamma(0)] = [p\gamma(1)] \cup \gamma(s)|_{[0,L]}$, and $\tilde{\gamma}(s)|_{[0,L]}$ has the same phenomenon.

We now assume that $W \triangleq \{\bar{s} \in [0, L] | (1.6) \text{ occurs at } \bar{s}\}$ is not equal to [0, L]. Note that if (A2.1) is not true, in order to get a contradiction we can assume that

$$|p\gamma(s)| > |\tilde{p}\tilde{\gamma}(s)| \text{ for all } s \in (0, L),$$
(1.17)

which implies that

$$|\uparrow_{\gamma(0)}^{p} \dot{\gamma}(0)| \ge |\uparrow_{\tilde{\gamma}(0)}^{\tilde{p}} \dot{\tilde{\gamma}}(0)| \text{ (and } |\uparrow_{\gamma(L)}^{p} \dot{\gamma}(L)| \le |\uparrow_{\tilde{\gamma}(L)}^{\tilde{p}} \dot{\tilde{\gamma}}(L)|).$$
(1.18)

In the following, we will derive a contradiction under (1.18) (with (1.17)).

Note that W is a closed subset of [0, L], so $W^c \triangleq [0, L] \setminus W$ is a union of some intervals. Due to the C^2 -property of $\gamma(s)|_{[0,L]}$ and $\tilde{\gamma}(s)|_{[0,L]}$, for any $\bar{s} \in W^c$ (i.e., $|\uparrow_{\gamma(\bar{s})}^p \dot{\gamma}(\bar{s})| \neq 0$ or π), there is an interval $I \subset W^c$ containing \bar{s} such that for any $s_1, s_2 \in I$ there exists $\tilde{\gamma}_{s_1,s_2}(s)|_{[0,L]}$ (equal to $\tilde{\gamma}(s)|_{[0,L]}$ up to an isometry of \mathbb{S}^2_k) satisfying (1.9) and (1.10). Then by Substep 1 and 2, W^c must contain a *nice* interval (s_1, s_2) . Here, we call (s_1, s_2) a nice interval if $\tilde{\gamma}_{s_1,s_2}(s)|_{[0,L]}$ satisfying (1.9) and (1.10) exists and $|p\gamma(s)| \leq |\tilde{p}\tilde{\gamma}_{s_1,s_2}(s)|$ for any $s \in (s_1, s_2)$.

We now can take a maximal nice interval $(s_{11}, s_{21}) \subset (0, L)$ (i.e. any $(c, d) \supseteq (s_{11}, s_{21})$ is not nice), which may contain the point in W. It is clear that

$$|\uparrow_{\gamma(s_{11})}^{p} \dot{\gamma}(s_{11})| \leq |\uparrow_{\tilde{\gamma}_{s_{11},s_{21}}(s_{11})}^{\tilde{p}} \dot{\tilde{\gamma}}_{s_{11},s_{21}}(s_{11})|, |\uparrow_{\gamma(s_{21})}^{p} \dot{\gamma}(s_{21})| \geq |\uparrow_{\tilde{\gamma}_{s_{11},s_{21}}(s_{21})}^{\tilde{p}} \dot{\tilde{\gamma}}_{s_{11},s_{21}}(s_{21})|.$$

$$(1.19)$$

By (1.17), $[s_{11}, s_{21}] \neq [0, L]$, so outside of $[s_{11}, s_{21}]$ it is possible that there exists a maximal nice interval. Note that there are at most $\{(s_{1j}, s_{2j})\}_{j=1}^{\infty}$ such that $(s_{1(j+1)}, s_{2(j+1)})$ is a maximal nice interval outside of $\bigcup_{l=1}^{j} [s_{1j}, s_{2j}]$ and

$$W \cup \bigcup_{j} [s_{1j}, s_{2j}] = [0, L]$$
, and any $s \notin W$ lies in some $[s_{1j}, s_{2j})$ and $(s_{1j'}, s_{2j'}]$. (1.20)

In order to complete Substep 3, we introduce the following function

$$d(s) \triangleq \begin{cases} |\tilde{p}\tilde{\gamma}_{s_{1j},s_{2j}}(L)|, & s \in (s_{1j},s_{2j}) \\ |\tilde{p}\tilde{\gamma}_s(L)|, & s \in [0,L] \setminus \bigcup_j (s_{1j},s_{2j}) \end{cases}$$

(for $\tilde{\gamma}_s$ refer to the beginning of Step 1). Note that $d(L) = |\tilde{p}\tilde{\gamma}(L)|$, and (1.18) implies that $d(0) \ge |\tilde{p}\tilde{\gamma}(L)| \ (= d(L))$. However, we claim that

$$d(s) \ge d(0) \text{ for all } s \text{ with } d(L) > d(0), \tag{1.21}$$

a contradiction (and thus Substep 3 is finished).

We now need only to verify (1.21) (in fact, the proof implies that $d(s)|_{[0,L]}$ is nondecreasing). We first show that $d(s) \ge d(0)$ for all s. Let A be the collection of such $\bar{s} \in [0, L]$ that $d(s) \geq d(0)$ on $[0, \bar{s}]$, and that for $s' \in (s_{1j}, s_{2j}) \cap [0, \bar{s}]$ (resp. $s' \in [0, \bar{s}] \setminus \bigcup_j (s_{1j}, s_{2j})) [\tilde{p}\tilde{\gamma}_{s_{1j}, s_{2j}}(s')] \cup \tilde{\gamma}_{s_{1j}, s_{2j}}(s)|_{[s', L]} \cup [\tilde{p}\tilde{\gamma}_{s_{1j}, s_{2j}}(L)]$ (resp. $[\tilde{p}\tilde{\gamma}_{s'}(s')] \cup \tilde{\gamma}_{s'}(s)|_{[s', L]} \cup [\tilde{p}\tilde{\gamma}_{s'}(L)]$) is convex. In fact, we can prove that A is a non-empty, open and closed subset of [0, L], i.e. A = [0, L]. Note that $A \neq \emptyset$ because $0 \in A$ by (1.18). As for 'open and closed', it suffices to show that

if
$$\bar{s} \in A$$
 and $\bar{s} \neq L$, then there is a $\delta > 0$ such that $[\bar{s}, \bar{s} + \delta) \subset A$ (1.22)

and

if
$$(\bar{s} - \delta', \bar{s}) \subset A$$
 for some $\delta' > 0$, then $\bar{s} \in A$. (1.23)

Case 1: If \bar{s} lies in some (s_{1j}, s_{2j}) , then (1.22) and (1.23) hold automatically.

Case 2: If \bar{s} is equal to some s_{1j} (resp. s_{2j}), then the first (resp. the second) inequality of corresponding (1.19) (together with the convexity of $\tilde{\gamma}(s)|_{[\bar{s},L]}$) implies (1.22) (resp. (1.23)). Moreover, if $\bar{s} = s_{1j} = s_{2j'}$, then the maximality of each (s_{1j}, s_{2j}) implies that at least one inequality of corresponding (1.19) is strict, and thus

$$d(s) \ge d(\bar{s}) > d(s') \text{ or } d(s) > d(\bar{s}) \ge d(s'), \ \forall \ s \in (s_{1j}, s_{2j}), s' \in (s_{1j'}, s_{2j'}).$$
(1.24)

(Note that if $[0, L] = \bigcup_{i} [s_{1i}, s_{2i}]$, then Case 1 and 2 right above implies (1.21).)

By (1.20), the residual case is:

Case 3: For (1.22) (resp. (1.23)), $\bar{s} \in W \setminus \bigcup_j (s_{1j}, s_{2j})$ and $\bar{s} \neq s_{1j}$ (resp. $\bar{s} \neq s_{2j}$) for any j. In this case, due to the similarity, we just give a proof for (1.22). We first observe that $|\uparrow_{\gamma(\bar{s})}^p \dot{\gamma}(\bar{s})| = \pi$. Otherwise, $|\uparrow_{\gamma(\bar{s})}^p \dot{\gamma}(\bar{s})| = 0$ (see (1.6)), so the convexity of $[\tilde{p}\tilde{\gamma}_{\bar{s}}(\bar{s})] \cup \tilde{\gamma}_{\bar{s}}(s)|_{[\bar{s},L]} \cup [\tilde{p}\tilde{\gamma}_{\bar{s}}(L)]$ (note that $\bar{s} \in A \setminus \bigcup_j (s_{1j}, s_{2j})$) implies that $\tilde{\gamma}_{\bar{s}}(s)|_{[\bar{s},L]}$ is a minimal geodesic ($\subseteq [\tilde{\gamma}_{\bar{s}}(\bar{s})\tilde{p}]$), so is $\gamma(s)|_{[\bar{s},L]}$ ($\subseteq [\gamma(\bar{s})p]$); which implies that \bar{s} has to be L (otherwise, (\bar{s}, L) belongs to some (s_{1j}, s_{2j})). Due to $|\uparrow_{\gamma(\bar{s})}^p \dot{\gamma}(\bar{s})| = \pi$, it is not hard to see that there is a small $\delta > 0$ such that one of the following holds: Subcase 1: $[\bar{s}, \bar{s} + \delta) \subset W \setminus \bigcup_i (s_{1j}, s_{2j})$;

Subcase 2: For any $\delta' \leq \delta$, $(\bar{s}, \bar{s} + \delta')$ contains an infinite number of (s_{1j}, s_{2j}) .

In Subcase 1, $\gamma(s)|_{[\bar{s},\bar{s}+\delta)}$ is a minimal geodesic (belonging to $[p\gamma(\bar{s}+\delta)]$), but $\tilde{\gamma}_{\bar{s}}(s)|_{[\bar{s},\bar{s}+\delta')}$ with $\delta' \leq \delta$ is not (otherwise, $(\bar{s},\bar{s}+\delta')$ belongs to some (s_{1j},s_{2j})). Then for any $\hat{s} \in (\bar{s},\bar{s}+\delta)$, we can rotate $\tilde{\gamma}_{\hat{s}}(s)|_{[0,L]}$ around $\tilde{\gamma}_{\hat{s}}(\hat{s})$ to $\tilde{\gamma}^*_{\hat{s}}(s)|_{[0,L]}$ such that $\tilde{\gamma}^*_{\hat{s}}(\bar{s}) \in [\tilde{p}\tilde{\gamma}_{\hat{s}}(\hat{s})]$. Note that $|\tilde{p}\tilde{\gamma}^*_{\hat{s}}(\bar{s})| > |\tilde{p}\tilde{\gamma}_{\bar{s}}(\bar{s})|$. By comparing $\tilde{\gamma}_{\hat{s}}(s)|_{[0,L]}$, $\tilde{\gamma}^*_{\hat{s}}(s)|_{[0,L]}$ and $\tilde{\gamma}_{\hat{s}}(s)|_{[0,L]}$ (which are all isometric to the convex $\tilde{\gamma}(s)|_{[0,L]}$), it is not hard to see that

$$d(\hat{s}) = |\tilde{p}\tilde{\gamma}_{\hat{s}}(L)| > |\tilde{p}\tilde{\gamma}^*_{\hat{s}}(L)| > |\tilde{p}\tilde{\gamma}_{\bar{s}}(L)| = d(\bar{s})$$

$$(1.25)$$

and $[\tilde{p}\tilde{\gamma}_{\hat{s}}(\hat{s})] \cup \tilde{\gamma}_{\hat{s}}(s)|_{[\hat{s},L]} \cup [\tilde{p}\tilde{\gamma}_{\hat{s}}(L)]$ is convex. It follows that $[\bar{s}, \bar{s} + \delta) \subset A$.

In Subcase 2, an ideal model is that $(\bar{s}, \bar{s} + \delta)$ belongs to a union of many $[s_{1j}, s_{2j}]$. In this case, by a similar reason for Case 1 and 2, we can conclude that $[\bar{s}, \bar{s} + \delta) \subset A$ and (1.24) implies that

$$d(\bar{s}+\delta) > d(\bar{s}). \tag{1.26}$$

In other cases, similarly, we only need to show that $\hat{s} \in A$, where $\hat{s} \in (\bar{s}, \bar{s} + \delta) \cap W \setminus \bigcup_{i}(s_{1j}, s_{2j})$ and $\hat{s} \neq s_{2j}$ for any j. Moreover, we can assume that

for any
$$\hat{s}' \in [\bar{s}, \hat{s}), \, (\hat{s}', \hat{s})$$
 is not a nice interval; (1.27)

otherwise, since $\bar{s}, \hat{s} \notin \bigcup_j (s_{1j}, s_{2j})$, we can reselect the maximal nice intervals in (\bar{s}, \hat{s}) such that \hat{s} is just some s_{2j} . Note that when δ is sufficiently small, $|\uparrow_{\gamma(\hat{s})}^p \dot{\gamma}(\hat{s})| = \pi$ because $|\uparrow_{\gamma(\bar{s})}^p \dot{\gamma}(\bar{s})| = \pi$ and $\hat{s} \in W$. Similarly, we can rotate $\tilde{\gamma}_{\hat{s}}(s)|_{[0,L]}$ around $\tilde{\gamma}_{\hat{s}}(\hat{s})$ to $\tilde{\gamma}_{\hat{s}}^*(s)|_{[0,L]}$ such that $\tilde{\gamma}_{\hat{s}}^*(\bar{s}) \in [\tilde{p}\tilde{\gamma}_{\hat{s}}(\hat{s})]$. Then similar to Subcase 1, we can conclude that $|\bar{s}, \bar{s} + \delta) \subset A$ and the corresponding (1.25) holds if we can show that $|\tilde{p}\tilde{\gamma}_{\hat{s}}^*(\bar{s})| > |\tilde{p}\tilde{\gamma}_{\bar{s}}(\bar{s})|$. In fact, if $|\tilde{p}\tilde{\gamma}_{\hat{s}}^*(\bar{s})| \leq |\tilde{p}\tilde{\gamma}_{\bar{s}}(\bar{s})|$, then $\tilde{\gamma}_{\hat{s}}^*(s)|_{[0,L]}$ has to be equal to some $\tilde{\gamma}_{\hat{s}',\hat{s}}(s)|_{[0,L]}$ with $\hat{s}' \in [\bar{s}, \hat{s})$ and (\hat{s}', \hat{s}) is a nice interval, which contradicts (1.27).

Note that we have proven $d(s)|_{[0,L]} \ge d(0)$, and the proof (especially (1.24-26)) implies d(L) > d(0), i.e. (1.21) holds (so Substep 3 (and thus Step 1) is completed).

Step 2. To prove (A2.2), i.e. $|p\gamma(s)| \ge |\tilde{p}\tilde{\gamma}(s)| \forall s \in (0, L]$; and if equality holds for some $s_0 \in (0, L]$, then $|p\gamma(s)| = |\tilde{p}\tilde{\gamma}(s)|$ for all $s \in [0, s_0]$.

We first give an observation: there is a $\delta > 0$ such that $|p\gamma(s)| \ge |\tilde{p}\tilde{\gamma}(s)|$ for all $s \in (0, \delta)$. In fact, due to $|\uparrow_{\gamma(0)}^{p} \dot{\gamma}(0)| = |\uparrow_{\tilde{\gamma}(0)}^{\tilde{p}} \dot{\tilde{\gamma}}(0)|$ (the condition in (A2.2)), for sufficiently small $\delta > 0$ if $|p\gamma(\delta)| < |\tilde{p}\tilde{\gamma}(\delta)|$, then $\tilde{\gamma}_{0,\delta}(s)|_{[0,L]}$ satisfying (1.9) and (1.10) exists and $|\uparrow_{\gamma(0)}^{p} \dot{\gamma}(0)| > |\uparrow_{\tilde{\gamma}_{0,\delta}(0)}^{\tilde{p}} \dot{\tilde{\gamma}}_{0,\delta}(0)|$. However, by applying (A2.1) on $\gamma(s)|_{[0,\delta]}$ and $\tilde{\gamma}_{0,\delta}(s)|_{[0,\delta]}$, it has to hold that $|\uparrow_{\gamma(0)}^{p} \dot{\gamma}(0)| \le |\uparrow_{\tilde{\gamma}_{0,\delta}(0)}^{\tilde{p}} \dot{\tilde{\gamma}}_{0,\delta}(0)|$, a contradiction.

Due to the observation right above, we can let $s_0 \neq 0$ be the maximal s such that $|p\gamma(s)| \geq |\tilde{p}\tilde{\gamma}(s)|$ for all $s \in [0, s_0]$. In order to prove (A2.2), we need to show that $s_0 = L$. If it is not true, then $|p\gamma(s_0)| = |\tilde{p}\tilde{\gamma}(s_0)|$. Then by applying (A2.1) on $\gamma(s)|_{[0,s_0]}$ and $\tilde{\gamma}(s)|_{[0,s_0]}$, we have that $|p\gamma(s)| \leq |\tilde{p}\tilde{\gamma}(s)|$ for all $s \in [0, s_0]$. It follows that $|p\gamma(s)| = |\tilde{p}\tilde{\gamma}(s)|$ for all $s \in [0, s_0]$, which implies that $|\uparrow_{\gamma(s_0)}^p \dot{\gamma}(s_0)| = |\uparrow_{\tilde{\gamma}(s_0)}^{\tilde{p}} \dot{\tilde{\gamma}}(s_0)|$. Thereby, similar to the observation right above, there is a $\delta > 0$ such that $|p\gamma(s)| \geq |\tilde{p}\tilde{\gamma}(s)|$ for all $s \in (s_0, s_0 + \delta)$, which contradicts the maximality of s_0 (i.e., $s_0 = L$).

Note that the proof for $s_0 = L'$ implies that if $|p\gamma(s_0)| = |\tilde{p}\tilde{\gamma}(s_0)|$ with $s_0 \in (0, L]$, then $|p\gamma(s)| = |\tilde{p}\tilde{\gamma}(s)|$ for all $s \in [0, s_0]$.

Step 3. To prove (0.3) in (A1), i.e. $|\gamma(0)\gamma(L)| \ge |\tilde{\gamma}(0)\tilde{\gamma}(L)|$.

We will derive a contradiction by assuming $|\gamma(0)\gamma(L)| < |\tilde{\gamma}(0)\tilde{\gamma}(L)|$.

We first claim that $|\gamma(0)\gamma(s)| < |\tilde{\gamma}(0)\tilde{\gamma}(s)|$ for all $s \in (0, L)$. If it is not true, then there is $s_0 \in (0, L)$ such that $|\gamma(0)\gamma(s_0)| = |\tilde{\gamma}(0)\tilde{\gamma}(s_0)|$. Note that if k > 0, by $(1.2), \, |\gamma(0)\gamma(L)| < |\tilde{\gamma}(0)\tilde{\gamma}(L)|$ ' implies that $|\gamma(0)\gamma(L)| + L < \frac{2\pi}{\sqrt{k}}$, and thus $|\gamma(0)\gamma(s_1)| + |\gamma(0)\gamma(s_1)| + |\gamma(0)\gamma(s_1)| + |s_2 - s_1| < \frac{2\pi}{\sqrt{k}}$ for all $s_1, s_2 \in [0, L]$. Hence, we can apply (A2.1) on $\gamma(s)|_{[0,s_0]}$ and $\tilde{\gamma}(s)|_{[0,s_0]}$ (with $p = \gamma(0)$ and $\tilde{p} = \tilde{\gamma}(0)$) to conclude that $|\gamma(0)\gamma(s)| \le |\tilde{\gamma}(0)\tilde{\gamma}(s)|$ for all $s \in (0, s_0)$. In fact, this implies that $|\gamma(0)\gamma(s)| \le |\tilde{\gamma}(0)\tilde{\gamma}(s)|$ for all $s \in (0, L)$. Hence, by Lemma 1.2 it holds that $|\uparrow_{\gamma(s_0)}^{\gamma(0)} \dot{\gamma}(s_0)| = |\uparrow_{\tilde{\gamma}(s_0)}^{\tilde{\gamma}(0)} \dot{\tilde{\gamma}}(s_0)|$. Then we can apply (A2.2) on $\gamma(s)|_{[s_0,L]}$ and $\tilde{\gamma}(s)|_{[s_0,L]}$ (with $p = \gamma(0)$ and $\tilde{p} = \tilde{\gamma}(0)$) to conclude that $|\gamma(0)\gamma(L)| \ge |\tilde{\gamma}(0)\tilde{\gamma}(L)|$, which contradicts $|\gamma(0)\gamma(L)| < |\tilde{\gamma}(0)\tilde{\gamma}(L)|$ '. By the claim right above, we can select a sufficiently small $s_1 \in (0, L)$ such that $|\gamma(0)\gamma(s_1)| - |\tilde{\gamma}(0)\tilde{\gamma}(s_1)| = \min_{s \in [0,s_1]} \{|\gamma(0)\gamma(s)| - |\tilde{\gamma}(0)\tilde{\gamma}(s)|\}$, which implies that

$$\left|\uparrow_{\gamma(s_1)}^{\gamma(0)}\left(-\dot{\gamma}(s_1)\right)\right| \ge \left|\uparrow_{\tilde{\gamma}(s_1)}^{\tilde{\gamma}(0)}\left(-\dot{\tilde{\gamma}}(s_1)\right)\right|.$$
(1.28)

On the other hand, since s_1 is sufficiently small, there is $q \in M$ such that

$$|q\gamma(s_1)| = |\gamma(0)\gamma(s_1)|$$
 and $|q\gamma(0)| + |q\gamma(s_1)| = |\tilde{\gamma}(0)\tilde{\gamma}(s_1)|,$ (1.29)

and that $|\uparrow_{\gamma(s_1)}^q(-\dot{\gamma}(s_1))| = |\uparrow_{\gamma(s_1)}^q\uparrow_{\gamma(s_1)}^{\gamma(0)}| + |\uparrow_{\gamma(s_1)}^{\gamma(0)}(-\dot{\gamma}(s_1))|$, which together with (1.28) implies that

$$\left|\uparrow_{\gamma(s_1)}^q \left(-\dot{\gamma}(s_1)\right)\right| > \left|\uparrow_{\tilde{\gamma}(s_1)}^{\tilde{\gamma}(0)} \left(-\dot{\tilde{\gamma}}(s_1)\right)\right|.$$
(1.30)

Let $\tilde{q} \in [\tilde{\gamma}(0)\tilde{\gamma}(s_1)]$ such that $|\tilde{q}\tilde{\gamma}(s_1)| = |q\gamma(s_1)| (= |\gamma(0)\gamma(s_1)|)$, which enables us to apply (A2.1) here to conclude that $|q\gamma(s)| \le |\tilde{q}\tilde{\gamma}(s)|$ for all $s \in [0, s_1]$. However, by (1.30) it is clear that $|q\gamma(s)| > |\tilde{q}\tilde{\gamma}(s)|$ for s less than and close to s_1 , a contradiction.

Step 4. To prove (A2.3) and the rigidity part of (A1).

We first prove the rigidity part for (A2.1), i.e. if $|p\gamma(s_0)| = |\tilde{p}\tilde{\gamma}(s_0)|$ for some $s_0 \in (0, L)$ in (A2.1), then $\bigcup_{s \in [0, L]} [p\gamma(s)]$ with induced metric is isometric to $\bigcup_{s \in [0, L]} [\tilde{p}\tilde{\gamma}(s)]$. Note that the proof will be done once one show that for any sufficiently small interval $[s_1, s_2] \subset [0, L], \bigcup_{s \in [s_1, s_2]} [p\gamma(s)]$ is convex and is isometric to $\bigcup_{s \in [s_1, s_2]} [\tilde{p}\tilde{\gamma}(s)]$.

Note that if $[s_1, s_2]$ is sufficiently small, we can assume that $\bigcup_{s \in [s_1, s_2]} [p\gamma(s)]$ lies in $M \setminus C(q)$ for any $q \in \bigcup_{s \in [s_1, s_2]} [p\gamma(s)]$. Then it suffices to verify that

$$|a_1a_2| = |\tilde{a}_1\tilde{a}_2| \text{ for any } a_i \in [p\gamma(s_i)] \text{ and } \tilde{a}_i \in [\tilde{p}\tilde{\gamma}(s_i)] \text{ with } |pa_i| = |\tilde{p}\tilde{a}_i|, \qquad (1.31)$$

and

$$[a_1a_2] \subset \bigcup_{s \in [s_1, s_2]} [p\gamma(s)]. \tag{1.32}$$

We first show that (1.31) is true if $a_2 = \gamma(s_2)$, i.e.

$$|a_1\gamma(s_2)| = |\tilde{a}_1\tilde{\gamma}(s_2)|.$$
(1.33)

From Step 1, we know that $|p\gamma(s)| \leq |\tilde{p}\tilde{\gamma}(s)|$ for all $s \in (0, L)$, so by Lemma 1.2

$$\left|\uparrow_{\gamma(s_0)}^p \dot{\gamma}(s_0)\right| = \left|\uparrow_{\tilde{\gamma}(s_0)}^{\tilde{p}} \dot{\tilde{\gamma}}(s_0)\right|.$$

Then by the rigidity part of Step 2, we have that

$$|p\gamma(s)| = |\tilde{p}\tilde{\gamma}(s)| \text{ for all } s \in [0, L],$$
(1.34)

which in turn implies that

$$\left|\uparrow_{\gamma(s)}^{p} \dot{\gamma}(s)\right| = \left|\uparrow_{\tilde{\gamma}(s)}^{\tilde{p}} \dot{\tilde{\gamma}}(s)\right| \text{ for all } s \in [0, L].$$

$$(1.35)$$

By (1.34) and (1.35), we can apply (A2.1) on {the curve $\gamma(s)|_{[s_1,s_2]} \cup [p\gamma(s_1)], \gamma(s_2)$ } and {the curve $\tilde{\gamma}(s)|_{[s_1,s_2]} \cup [\tilde{p}\tilde{\gamma}(s_1)], \tilde{\gamma}(s_2)$ } to conclude that $|\gamma(s_1)\gamma(s_2)| \leq |\tilde{\gamma}(s_1)\tilde{\gamma}(s_2)|$. On the other hand, by Step 3 we know that $|\gamma(s_1)\gamma(s_2)| \geq |\tilde{\gamma}(s_1)\tilde{\gamma}(s_2)|$, and thus $|\gamma(s_1)\gamma(s_2)| = |\tilde{\gamma}(s_1)\tilde{\gamma}(s_2)|$. Then (1.33) follows (just as (1.34)), by which we can similarly apply (A2.1) on {the curve $\gamma(s)|_{[s_1,s_2]} \cup [p\gamma(s_2)], a_1$ } and {the curve $\tilde{\gamma}(s)|_{[s_1,s_2]} \cup [\tilde{p}\tilde{\gamma}(s_2)], \tilde{a}_1$ } to conclude (1.31).

For (1.32), we let $q \in [a_1a_2]$ and $\tilde{q} \in [\tilde{a}_1\tilde{a}_2]$ with $|qa_1| = |\tilde{q}\tilde{a}_1|$, and assume that $\tilde{q} \in [\tilde{p}\tilde{\gamma}(\bar{s})]$ with $\bar{s} \in [s_1, s_2]$. Note that (1.31) implies that $|\uparrow_{a_1}^{\gamma(s_1)}\uparrow_{a_1}^{a_2}| =$ $|\uparrow_{\tilde{a}_1}^{\tilde{\gamma}(s_1)}\uparrow_{\tilde{a}_1}^{\tilde{a}_2}|$. This together with (1.35) enables us to apply the rigidity part of Step 2 on {the curve $[a_1\gamma(s_1)] \cup \gamma(s)|_{[s_1,s_2]} \cup [a_2\gamma(s_2)], q$ } and {the curve $[\tilde{a}_1\tilde{\gamma}(s_1)] \cup \tilde{\gamma}(s)|_{[s_1,s_2]} \cup$ $[\tilde{a}_2\tilde{\gamma}(s_2)], \tilde{q}$ }, and {the curve $[a_1p] \cup [a_2p], q$ } and {the curve $[\tilde{a}_1\tilde{p}] \cup [\tilde{a}_2\tilde{p}], \tilde{q}$ } respectively to conclude that q lies in $[p\gamma(\bar{s})]$. That is, (1.32) follows.

The rigidity part for (A2.2) can be seen similarly because the corresponding (1.34) and (1.35) to $[0, s_0]$ hold (note that by the rigidity part of Step 2, if the equality in (0.5) holds for some $s_0 \in (0, L]$, then $|p\gamma(s)| = |\tilde{p}\tilde{\gamma}(s)|$ for all $s \in [0, s_0]$).

Similarly, the rigidity part of (A1) can be verified once we show that $|\gamma(0)\gamma(s)| = |\tilde{\gamma}(0)\tilde{\gamma}(s)|$ for all $s \in [0, L]$. In fact, due to $|\gamma(0)\gamma(L)| = |\tilde{\gamma}(0)\tilde{\gamma}(L)|'$, we can apply (A2.1) on {the curve $\gamma(s)|_{[0,L]}, \gamma(0)$ } and {the curve $\tilde{\gamma}(s)|_{[0,L]}, \tilde{\gamma}(0)$ } to conclude that $|\gamma(0)\gamma(s)| \leq |\tilde{\gamma}(0)\tilde{\gamma}(s)|$ for all $s \in [0, L]$. On the other hand, by Step 3 we can see that $|\gamma(0)\gamma(s)| \geq |\tilde{\gamma}(0)\tilde{\gamma}(s)|$, so it follows that $|\gamma(0)\gamma(s)| = |\tilde{\gamma}(0)\tilde{\gamma}(s)|$.

2 Proof of Theorem A for the case where $\sec_M \ge k$

In this section, we always assume that $\sec_M \ge k$.

We first introduce the notion $\gamma(s)|_{[0,L]}$ is convex to p' in Theorem A. Let $\gamma(s)|_{[0,L]} \subset M$ be an arc-length parameterized C^2 -curve. Note that given $p \in M$, $|p\gamma(s)||_{[0,L]}$ is a 1-Lipschitz function (maybe not a C^1 -function). Then for $k \leq 0$ and if $|p\gamma(s)||_{[0,L]} < \frac{\pi}{\sqrt{k}}$ for k > 0, given $\bar{p} \in \mathbb{S}^2_k$, there is a unique (up to a rotation) arc-length parameterized Lipschitz-curve $\bar{\gamma}(s)|_{[0,L]}$ in \mathbb{S}^2_k such that

$$|p\gamma(s)| = |\bar{p}\bar{\gamma}(s)| \text{ for all } s \in [0, L],$$
(2.1)

and $[\bar{p}\bar{\gamma}(s)]$ turns clockwise as s increases ([PP], in which $\bar{\gamma}(s)|_{[0,L]}$ is called a development of $\gamma(s)|_{[0,L]}$ with respect to p). If k > 0 and $|p\gamma(s)| = \frac{2\pi}{\sqrt{k}}$ for some $s \in [0, L]$, by the Maximum Diameter Theorem ([CE]), we know that M is isometric to \mathbb{S}_k^n ; and thus there is an arc-length parameterized C^2 -curve $\bar{\gamma}(s)|_{[0,L]}$ such that (2.1) holds.

Definition 2.1 The $\gamma(s)|_{[0,L]}$ right above is said to be *convex to* p if (2.1.1) for all $s \in [0, L]$, there is $[p\gamma(s)]$ such that the angle between $D_{\dot{\gamma}(s)}\dot{\gamma}(s)$ and $\uparrow_{\gamma(s)}^{p}$ is less than or equal to $\frac{\pi}{2}$, and $\{\dot{\gamma}(s), D_{\dot{\gamma}(s)}\dot{\gamma}(s), \uparrow_{\gamma(s)}^{p}\}$ lies in a plane of $T_{\gamma(s)}M^{-5}$. (2.1.2) $[\bar{p}\bar{\gamma}(0)] \cup \bar{\gamma}(s)|_{[0,L]} \cup [\bar{p}\bar{\gamma}(L)]$ is a convex curve in \mathbb{S}^{2}_{k} .

⁵ If $M = \mathbb{S}_k^n$, (2.1.1) implies that p and $\gamma(s)|_{[0,L]}$ fall in an $\mathbb{S}_k^2 \subset \mathbb{S}_k^n$.

Remark 2.2 (2.2.1) In Definition 2.1, if $\gamma(s)|_{[0,L]}$ is a piecewise C^2 -curve, then at non-differential points, $\dot{\gamma}(s)$ shall be replaced by $\dot{\gamma}_{\pm}(s)$ and it shall be added that

$$|(-\dot{\gamma}_{-}(s))\uparrow_{\gamma(s)}^{p}| + |\dot{\gamma}_{+}(s)\uparrow_{\gamma(s)}^{p}| = |(-\dot{\gamma}_{-}(s))\dot{\gamma}_{+}(s)|.$$

(2.2.2) If k > 0 and $|p\gamma(s)| = \frac{2\pi}{\sqrt{k}}$ for some $s \in [0, L]$ (so M is isometric to \mathbb{S}_k^n), then by (1.2), (2.1.2) implies that each of $[p\gamma(0)] \cup \gamma(s)|_{[0,L]} \cup [p\gamma(L)]$ and $[\bar{p}\bar{\gamma}(0)] \cup \bar{\gamma}(s)|_{[0,L]} \cup [\bar{p}\bar{\gamma}(L)]$ is a union of two half great circles.

Since Theorem A for $\sec_M \ge k$ is almost an immediate corollary of the proof in Section 1, we only supply a rough proof for it.

Proof of Theorem A for $sec_M \ge k$.

Since $\gamma(s)|_{[0,L]}$ is convex to $\gamma(0)$ and p in (A1) and (A2), for the $\bar{\gamma}(s)|_{[0,L]}$ and \bar{p} corresponding to $\gamma(s)|_{[0,L]}$ and $p, \bar{\gamma}(s)|_{[0,L]} \cup [\bar{\gamma}(0)\bar{\gamma}(L)]$ and $[\bar{p}\bar{\gamma}(0)] \cup \bar{\gamma}(s)|_{[0,L]} \cup [\bar{p}\bar{\gamma}(L)]$ are convex curves in \mathbb{S}_k^2 (by (2.1.2)). We observe that the proof of Theorem A for sec_ $M \leq k$ in Section 1 applies to $\tilde{\gamma}(s)|_{[0,L]}$ and $\bar{\gamma}(s)|_{[0,L]}$ (which both lie in \mathbb{S}_k^2) by replacing $\gamma(s)|_{[0,L]}$ and $\tilde{\gamma}(s)|_{[0,L]}$ (in Section 1) by $\tilde{\gamma}(s)|_{[0,L]}$ and $\bar{\gamma}(s)|_{[0,L]}$ respectively. A problem here is that $\bar{\gamma}(s)|_{[0,L]}$ is only a 1-Lipschitz curve. However, via (1.3), (2.1.1) implies that the corresponding (1.5) still holds (in the support sense), and then one can check that the whole proof in Section 1 still works here. This together with (2.1) implies that (0.3-5) follows. Moreover, if equalities hold in (0.3-5), then $\tilde{\gamma}(s)|_{[0,L]}$ has to be equal to $\bar{\gamma}(s)|_{[0,L]}$ up to an isometry of \mathbb{S}_k^2 , which implies $\bar{\gamma}(s)|_{[0,L]}$ is also a C^2 -curve. It then is not hard to see the rigidity part of (A1) and (A2.3).

We will end this section by showing that Definition 2.1 is not so artificial through the following two remarks (especially Remark 2.4) on it.

Remark 2.3 (2.1.1) implies that $\bar{\gamma}(s) \subset \mathbb{S}^2_k$ satisfies that

$$\forall s \in [0, L], \exists \delta > 0 \text{ s.t. } [\bar{p}\bar{\gamma}(s-\delta)] \cup \bar{\gamma}(s)|_{[s-\delta, s+\delta]} \cup [\bar{p}\bar{\gamma}(s+\delta)] \text{ is convex}$$
(2.2)

(here if s = 0 (resp. L), then $s - \delta$ (resp. $s + \delta$) should be 0 (resp. L)). Recall that for $p \in M$ (sec_M $\geq k$) and an arc-length parameterized minimal geodesic $\sigma(s)|_{[a,b]} \subset M$, if

we let
$$f(s) = \rho_k(|p\sigma(s)|)$$
 with $\rho_k(x) = \begin{cases} \frac{1}{k}(1 - \cos(\sqrt{kx})), & k > 0\\ \frac{x^2}{2}, & k = 0, \text{ then}\\ \frac{1}{k}(1 - \cosh(\sqrt{-kx})), & k < 0 \end{cases}$

$$f''(s) \le 1 - kf(s)$$
 for all $s \in [a, b]$ (in the support sense, [Pe]). (2.3)

Thereby, for each $s_0 \in [0, L]$, by comparing $\gamma(s)|_{[0,L]}$ and the minimal geodesic tangent to $\gamma(s)|_{[0,L]}$ at $\gamma(s_0)$, via (1.3) one can see that (2.1.1) implies that (2.3) holds for \bar{p} and $\bar{\gamma}(s)|_{[0,L]}$. On the other hand, according to [PP], (2.2) is equivalent to (2.3) for \bar{p} and $\bar{\gamma}(s)|_{[0,L]}$ ($\subset \mathbb{S}^2_k$). It then follows that (2.1.1) implies (2.2). **Remark 2.4** (2.4.1) If $\gamma(s)|_{[0,L]}$ is a minimal geodesic in Definition 2.1, then $\gamma(s)|_{[0,L]}$ is convex to p automatically. It is clear that $\gamma(s)|_{[0,L]}$ satisfies (2.1.1), so it suffices to verify (2.1.2). We only need to consider the case where $\gamma(0) \notin [p\gamma(1)]$ and $\gamma(1) \notin [p\gamma(0)]$. In such a situation, for the \bar{p} and $\bar{\gamma}(s)|_{[0,L]}$ corresponding to p and $\gamma(s)|_{[0,L]}$, $s < |\bar{p}\bar{\gamma}(0)| + |\bar{p}\bar{\gamma}(s)|$ and thus $|\uparrow_{\bar{p}}^{\bar{\gamma}(0)}\uparrow_{\bar{p}}^{\bar{\gamma}(s)}| < \pi$ for all $s \in [0, L]$. This together with (2.2) (see Remark 2.3) implies (2.1.2).

(2.4.2) In (A2) of Theorem A for the case where $\sec_M \geq k$, if $\gamma(s)|_{[0,L]}$ is a minimal geodesic, then $\tilde{\gamma}(s)|_{[0,L]}$ has to be a minimal geodesic too by ' $\kappa(s) \geq \tilde{\kappa}(s)$ '. Then due to (2.4.1), (A2.1) and (A2.2) imply (0.2.1) and (0.2.2) respectively. Moreover, note that $[p\gamma(0)]$ and $[p\gamma(L)]$ are also convex to $\gamma(L)$ and $\gamma(0)$ respectively. Hence, if the equality of (0.4) holds for some $s \in (0, L)$, then by (A2.3) and by applying (A2) of Theorem A on $\{[p\gamma(0)], \gamma(L)\}$ and $\{[\tilde{p}\tilde{\gamma}(0)], \tilde{\gamma}(L)\}$ (and $\{[p\gamma(L)], \gamma(0)\}$ and $\{[\tilde{p}\tilde{\gamma}(L)], \tilde{\gamma}(0)\}$), it is not hard to conclude (0.2.3). That is, (A2) of Theorem A includes Theorem 0.2.

3 Proof of Theorem B

3.1 Preparations

Note that a proof of Theorem B has to depend only on its local version, i.e. the definition of Alexandrov spaces with lower curvature bound, which enables us to define the angle between two minimal geodesics starting from a common point ([BGP]).

In this section, X always denotes a complete Alexandrov space with curvature $\geq k$. Given $[pq], [pr] \subset X$, we denote by $\angle qpr$ the angle between them (i.e. $|\uparrow_p^q\uparrow_p^r|$). And we denote by $\triangle qpr$, a triangle, the union of three minimal geodesics $[pq], [pr], [qr] \subset X$.

From the definition of angles, one can see the following easy properties ([BGP]).

Lemma 3.1 Let $[pq], [rr'] \subset X$ with $r \in [pq]^{\circ}$. Then $\angle prr' + \angle qrr' = \pi$.

Lemma 3.2 Let $\{[p_iq_i]\}_{i=1}^{\infty}$ and $\{[p_ir_i]\}_{i=1}^{\infty}$ be two sequences of minimal geodesics in X. If $[p_iq_i] \rightarrow [pq]$ and $[p_ir_i] \rightarrow [pr]$ as $i \rightarrow \infty$, then $\angle qpr \leq \liminf \angle q_ip_ir_i$.

Furthermore, we have the following easy observation.

Lemma 3.3 Let $\{[p_iq_i]\}_{i=1}^{\infty}, \{[p_ir_i]\}_{i=1}^{\infty} \subset X$, and let $p \in X$. If $|p_ip| \to 0$, $|p_ir_i| \to 0$ and $|p_iq_i| \to a > 0$ as $i \to \infty$, then we have that

$$|q_i r_i| \le |q_i p_i| - |p_i r_i| \cos \angle q_i p_i r_i + o(|p_i r_i|).$$
(3.1)

Proof. By Toponogov's theorem around p, there is $\bar{q}_i \in [p_i q_i]$ near p such that

$$|\bar{q}_i r_i| \le |\bar{q}_i p_i| - |p_i r_i| \cos \angle q_i p_i r_i + o(|p_i r_i|).$$

Then (3.1) follows from that $|q_i r_i| \leq |q_i \bar{q}_i| + |\bar{q}_i r_i|$.

In Lemma 3.3, if $p_i = p$, $q_i = q$, and $[p_i r_i]$ lie in a minimal geodesic, then we can select [pq] such that (3.1) is an equality (the first variation formula, [BGP]). We will end this subsection by Alxandrov's lemma ([BGP]), a basic tool in Alexandrov geometry.

Lemma 3.4 Let $\triangle pqr$, $\triangle pqs$, $\triangle abc \subset \mathbb{S}_k^2$, where $\triangle pqr$ and $\triangle pqs$ are joined to each other in an exterior way along [pq], such that |ab| = |pr|, |ac| = |ps|, |bc| = |qr| + |qs|, and $|ab| + |ac| + |bc| < \frac{2\pi}{\sqrt{k}}$ if k > 0. Then $\angle pqr + \angle pqs \leq \pi$ (resp. $\geq \pi$) if and only if $\angle prq \geq \angle abc$ and $\angle psq \geq \angle acb$ (resp. $\angle prq \leq \angle abc$ and $\angle psq \leq \angle acb$).

3.2 Proof of Theorem B

Due to the similarity, we only give a proof for the case where k = 0.

Assume that Theorem B is not true, i.e. there is $s \in [qr]$ and $\tilde{s} \in [\tilde{q}\tilde{r}]$ with $|qs| = |\tilde{q}\tilde{s}|$ such that $|ps| < |\tilde{p}\tilde{s}|$. To get a contradiction, roughly, we will construct a sufficiently small triangle which does not satisfy Toponogov's theorem. The process is completed through the following three steps, where for $\triangle abc \subset X$ we denote by $\tilde{\angle}abc$ the corresponding angle of its comparison triangle ⁶ in \mathbb{S}_k^2 .

Step 1. To find $r' \in [qr]^{\circ}$ such that, for any [pr'], $\angle pr'q < \tilde{\angle}pr'q$ or $\angle pr'r < \tilde{\angle}pr'r$.

By Step 1, there is a $\triangle p_1 q_1 r_1$ (with $p_1, q_1, r_1 \in \triangle pqr$) such that

$$\angle p_1 q_1 r_1 < \angle p_1 q_1 r_1. \tag{3.2}$$

Let $peri(p_1q_1r_1)$ and $(p_1q_1r_1)_{min}$ denote the perimeter (i.e. $|p_1q_1| + |p_1r_1| + |q_1r_1|)$ and the length of the minimal side of $\Delta p_1q_1r_1$ respectively.

Step 2. To find a $\triangle p_2 q_2 r_2$ with $p_2, q_2, r_2 \in \triangle p_1 q_1 r_1$ such that $\operatorname{peri}(p_2 q_2 r_2) < \operatorname{peri}(p_1 q_1 r_1)$, $\angle p_2 q_2 r_2 < \tilde{\angle} p_2 q_2 r_2$, and either

(3.3) $\cos \angle p_2 q_2 r_2 - \cos \tilde{\angle} p_2 q_2 r_2 \ge \cos \angle p_1 q_1 r_1 - \cos \tilde{\angle} p_1 q_1 r_1$, or

(3.4) the following holds:

(3.4.1) $\operatorname{peri}(p_2q_2r_2) < \operatorname{peri}(p_1q_1r_1) - \frac{1}{2}(p_1q_1r_1)_{\min},$

(3.4.2) if $(p_2q_2r_2)_{\min} < (p_1q_1r_1)_{\min}$, then $\operatorname{peri}(p_2q_2r_2) < 4.5(p_1q_1r_1)_{\min}$.

Step 3. To repeat Step 2 to get $\{ \Delta p_i q_i r_i \}_{i=1}^{\infty}$ each of which satisfies $\operatorname{peri}(p_i q_i r_i) < \operatorname{peri}(p_{i-1}q_{i-1}r_{i-1}), \angle p_i q_i r_i < \tilde{\angle} p_i q_i r_i$, and the corresponding (3.3) or (3.4).

Note that any subsequence of $\{ \Delta p_i q_i r_i \}_{i=1}^{\infty}$ contains a converging subsequence if $\{ \Delta p_i q_i r_i \}_{i=1}^{\infty}$ lie in a compact domain of X. It is obviously true if X is compact. Anyway, due to our method of locating $\Delta p_i q_i r_i$, it turns out to be true even when X is not compact (see Remark 3.5 below).

Thereby, if $\operatorname{peri}(p_iq_ir_i) \to 0$ as $i \to \infty$, i.e. $\Delta p_iq_ir_i$ converge to a point passing to a subsequence, then $\langle \angle p_iq_ir_i < \widetilde{\angle}p_iq_ir_i \rangle$ contradicts the local Toponogov's theorem (cf. (0.3.2)). In fact, either all $(p_iq_ir_i)_{\min}$ have a positive lower bound, or $(p_iq_ir_i)_{\min} \to 0$ passing to a subsequence; and thus if there is an infinite number of $\Delta p_iq_ir_i$ satisfying (3.4), it is easy to see that $\operatorname{peri}(p_iq_ir_i) \to 0$.

Then we can assume that $\operatorname{peri}(p_i q_i r_i)$ converges to a positive number as $i \to \infty$, and meanwhile the corresponding (3.3) occurs for all $i \ge 2$. In this case, passing to a subsequence, $\Delta p_i q_i r_i$ converge to a minimal geodesic or a triangle as $i \to \infty$. For the

⁶We usually call $\triangle \tilde{a}\tilde{b}\tilde{c} \subset \mathbb{S}_k^2$ the comparison triangle of $\triangle abc$ if $|\tilde{a}\tilde{b}| = |ab|, |\tilde{a}\tilde{c}| = |ac|$ and $|\tilde{b}\tilde{c}| = |bc|$.

former case, it is not hard to see that (3.3) for each *i* together with Lemma 3.3 implies a contradiction. For the latter case, it is clear that (3.3) for each *i* and Lemma 3.2 imply that $\Delta p_i q_i r_i$ converge to a triangle $\Delta \bar{p} \bar{q} \bar{r}$ with

$$\cos \angle \bar{p}\bar{q}\bar{r} - \cos \tilde{\angle} \bar{p}\bar{q}\bar{r} \ge \cos \angle p_1 q_1 r_1 - \cos \tilde{\angle} p_1 q_1 r_1;$$

and thus we can repeat such a process on $\Delta \bar{p}\bar{q}\bar{r}$ until a contradiction is gotten.

In the rest of the proof, we need only to show how we accomplish Step 1 and 2.

On Step 1:

Let $r' \in [qr]$ and $\tilde{r}' \in [\tilde{q}\tilde{r}]$ with $|qr'| = |\tilde{q}\tilde{r}'|$ such that

$$|pr'| - |\tilde{p}\tilde{r}'| = \min_{s \in [qr], \tilde{s} \in [\tilde{q}\tilde{r}], |qs| = |\tilde{q}\tilde{s}|} \{ |ps| - |\tilde{p}\tilde{s}| \} < 0.$$

Then by Lemma 3.1 and 3.3 we can conclude that, similar to Lemma 1.2,

$$\angle qr'p = \angle \tilde{q}\tilde{r}'\tilde{p} \text{ and } \angle rr'p = \angle \tilde{r}\tilde{r}'\tilde{p}.$$

On the other hand, since $|pr'| < |\tilde{p}\tilde{r}'|$, by Lemma 1.3 it is easy to see that

$$\angle \tilde{q}\tilde{r}'\tilde{p} < \tilde{\angle}qr'p \text{ or } \angle \tilde{r}\tilde{r}'\tilde{p} < \tilde{\angle}rr'p$$

It therefore follows that

$$\angle qr'p < \tilde{\angle}qr'p$$
 or $\angle rr'p < \tilde{\angle}rr'p$.

On Step 2:

We can finish Step 2 according to the following two cases.

Case 1: One of $[p_1q_1]$ and $[q_1r_1]$, say $[p_1q_1]$, is the minimal side of $\Delta p_1q_1r_1$.

We first notice that by (3.2) and Lemma 3.3 there is $t \in [q_1r_1]$ and $\tilde{t} \in [\tilde{q}_1\tilde{r}_1]$ with $|\tilde{t}\tilde{r}_1| = |tr_1|$ such that $|p_1t| = |\tilde{p}_1\tilde{t}|$ and for any $t' \in [q_1t]^\circ$ and $\tilde{t}' \in [\tilde{q}_1\tilde{t}]$ with $|\tilde{t}'\tilde{t}| = |t't|$

$$|p_1 t'| < |\tilde{p}_1 \tilde{t}'|. \tag{3.5}$$

If $t \neq r_1$, then peri $(p_1q_1t) < \text{peri}(p_1q_1r_1)$ and $\cos \angle p_1q_1t - \cos \angle p_1q_1t = \cos \angle p_1q_1r_1 - \cos \angle p_1q_1r_1$. $\cos \angle p_1q_1r_1$. I.e., it suffices to let $(p_2, q_2, r_2) = (p_1, q_1, t)$.

We now assume that $t = r_1$, and take the point $s \in [q_1r_1]$ with $|sr_1| = \frac{1}{2}|p_1q_1|$. It is clear that both $\operatorname{peri}(p_1q_1s)$ and $\operatorname{peri}(p_1r_1s)$ are less than $\operatorname{peri}(p_1q_1r_1)$. If $\angle p_1q_1s < \tilde{\angle}p_1q_1s$ or $\angle p_1sq_1 < \tilde{\angle}p_1sq_1$, then we can let $(p_2,q_2,r_2) \triangleq (p_1,q_1,s)$ or (p_1,s,q_1) respectively which satisfies (3.4) (due to (3.5), it is easy to see that $\operatorname{peri}(p_1q_1s) < \operatorname{peri}(p_1q_1r_1) - \frac{1}{2}|p_1q_1|$, and if $(p_1q_1s)_{\min} < |p_1q_1|$ then $\operatorname{peri}(p_1q_1s) < 4|p_1q_1|$). If $\angle p_1q_1s \ge \tilde{\angle}p_1q_1s$ and $\angle p_1sq_1 \ge \tilde{\angle}p_1sq_1$, then we claim that $(p_2,q_2,r_2) = (p_1,s,r_1)$ which satisfies (3.3). First of all, note that $\angle p_1sr_1 = \pi - \angle p_1sq_1$ and $\angle p_1q_1r_1 = \angle p_1q_1s$. Then by Lemma 3.4, " $\angle p_1q_1s \ge \tilde{\angle}p_1q_1s$, $\angle p_1sq_1 \ge \tilde{\angle}p_1sq_1$ and $\angle p_1q_1r_1 < \tilde{\angle}p_1q_1r_1$ " implies that $\angle p_1sr_1 < \tilde{\angle}p_1sr_1$, and thus one can check that

$$\frac{\cos \angle p_1 s r_1 - \cos \angle p_1 s r_1}{\cos \angle p_1 q_1 r_1 - \cos \widetilde{\angle} p_1 q_1 r_1} \ge \frac{\cos(\pi - \angle p_1 s q_1) - \cos \angle p_1 s r_1}{\cos \widetilde{\angle} p_1 q_1 s - \cos \widetilde{\angle} p_1 q_1 r_1} = \frac{|p_1 q_1| \cdot |q_1 r_1|}{|p_1 s| \cdot |s r_1|} \overset{\text{by } (3.5)}{>} 1.$$

Case 2: $[p_1r_1]$ is the strictly minimal side of $\triangle p_1q_1r_1$.

Without loss of generality, we assume that $|p_1r_1| < |p_1q_1| \leq |q_1r_1|$. By the same reason as in Case 1, we can assume that (3.5) holds for all $t' \in [q_1r_1]^\circ$ and $\tilde{t}' \in [\tilde{q}_1\tilde{r}_1]$ with $|\tilde{t}'\tilde{r}_1| = |t'r_1|$. Then we can conclude that there is $s \in [q_1r_1]$ such that either $|sr_1| = |p_1r_1|$ and $|p_1r_1| \leq |p_1s| < \sqrt{2}|p_1r_1|$, or $|sr_1| > |p_1r_1|$ and $|p_1s| = |p_1r_1|$. It follows that $\operatorname{peri}(p_1q_1s) < \operatorname{peri}(p_1q_1r_1) - \frac{1}{2}|p_1r_1|$, and $\operatorname{peri}(p_1sq_1) < 4.5|p_1r_1|$ if $|q_1s| < |p_1r_1|$. Moreover, similar to the proof for Case 1, we can let $(p_2, q_2, r_2) = (p_1, q_1, s)$ or (p_1, s, q_1) which satisfies (3.4), or let $(p_2, q_2, r_2) = (p_1, s, r_1)$ which satisfies (3.3).

Remark 3.5 Let $\{ \Delta p_i q_i r_i \}_{i=1}^{\infty}$ be triangles in the right above proof, and define $d_{ij} \triangleq \max\{|p_j\{p_i, q_i, r_i\}|, |q_j\{p_i, q_i, r_i\}|, |r_j\{p_i, q_i, r_i\}|\}$ for $j > i \ge 1$. We can show that $\{d_{1j}\}_{j=2}^{\infty}$ have an upper bound, so $\{ \Delta p_i q_i r_i \}_{i=1}^{\infty}$ lie in a compact domain of X even when X is not compact. Note that by the choice of $\Delta p_i q_i r_i$, it is possible that

$$d_{i(i+1)} \le \operatorname{peri}(p_i q_i r_i) - \operatorname{peri}(p_{i+1} q_{i+1} r_{i+1});$$
(3.6)

and thus it is clear that if (3.6) holds for all i with $1 \le i \le j - 1$, then

$$d_{1j} \le \sum_{i=1}^{j-1} d_{i(i+1)} \le \sum_{i=1}^{j-1} \left(\operatorname{peri}(p_i q_i r_i) - \operatorname{peri}(p_{i+1} q_{i+1} r_{i+1}) \right) < \operatorname{peri}(p_1 q_1 r_1).$$

Unfortunately, (3.6) might not be true, and even there is no constant $c \in (0, 1)$ such that $c \cdot d_{i(i+1)} \leq \operatorname{peri}(p_i q_i r_i) - \operatorname{peri}(p_{i+1} q_{i+1} r_{i+1})$ for all *i*. However, by analyzing the process of locating $\Delta p_i q_i r_i$ in Step 2 and 3 case by case, we can conclude that, for all *j*, there is a constant $c \in (0, 1)$ and $1 = i_0 < i_1 < \cdots < i_k = j$ such that

$$d_{1j} \le \sum_{l=0}^{k-1} d_{i_l i_{l+1}} \le \frac{1}{c} \cdot \sum_{l=0}^{k-1} \left(\operatorname{peri}(p_{i_l} q_{i_l} r_{i_l}) - \operatorname{peri}(p_{i_{l+1}} q_{i_{l+1}} r_{i_{l+1}}) \right) < \frac{1}{c} \cdot \operatorname{peri}(p_1 q_1 r_1)$$

(where it might hold that $c \cdot d_{i_l i_{l+1}} > \text{peri}(p_{i_l} q_{i_l} r_{i_l}) - \text{peri}(p_{i_{l+1}} q_{i_{l+1}} r_{i_{l+1}})$ for some *l*).

References

- [BGP] Yu. Burago, M. Gromov and G. Perel'man, A.D. Alexandrov spaces with curvature bounded below, Uspeckhi Mat. Nank 47:2 (1992): 3-51.
- [CE] J. Cheeger and D. Ebin, Comparison theorems in Riemannian geometry, New York, 1975.
- [GM] K. Grove and S. Markvorsen, New Extremal Problems for the Riemannian Recognition Program via Alexandrov Geometry, J. of AMS, 8:1 (1995): 1-28.
- [Ho] H. Hopf, Differential Geometry in the Large, Springer-Verlag, 1983.
- [Me] W. Meyer, Toponogov's Theorem and Applications.

- [PP] G. Perel'man and A. Petrunin, Quasigeodesics and Gradient Curves in Alexandrov Spaces, 1994.
- [Pe] P. Peterson, Riemannian Geometry, GTM 171, Springer-Verlag New York, 1998.
- [Pl] C. Plaut, Spaces of Wald-Berestovskii curvature bounded below. J. Geom. Anal. 6 (1996): 113-134.
- [Su] J. Sullivan, Curves of Finite Total Curvature, Discrete Diff. Geom. (OWS, volume 38) (2008): 137-161.
- [Sh] K. Shiohama, An introduction to the geometry of Alexandrov spaces, 1992.
- [Wu] H. Wu, Introduction to Riemannian geometry (in Chinese), Peking University Press, 1989.

School of Mathematical Sciences (and Lab. math. Com. Sys.), Beijing Normal University, Beijing, 100875 P.R.C. e-mail: wyusheng@bnu.edu.cn