# The one dimensional semi-classical Bogoliubov-de Gennes Hamiltonian with PT symmetry: generalized Bohr-Sommerfeld quantization rules

A Ifa

Université Tunis El-Manar, Tunis, Tunisia

#### M Rouleux

Aix-Marseille Univ, Université de Toulon, CNRS, CPT, Marseille, France E-mail: abdelwaheb.ifa@fsm.rnu.tn; rouleux@univ-tln.fr

Abstract. We present a method for computing first order asymptotics of semiclassical spectra for 1-D Bogoliubov-de Gennes (BdG) Hamiltonian from Supraconductivity, which models the electron/hole scattering through two SNS junctions. This involves: 1) reducing the system to Weber equation near the branching point at the junctions; 2) constructing local sections of the fibre bundle of microlocal solutions; 3) normalizing these solutions for the "flux norm" associated to the microlocal Wronskians; 4) finding the relative monodromy matrices in the gauge group that leaves invariant the flux norm; 5) from this we deduce Bohr-Sommerfeld (BS) quantization rules that hold precisely when the fibre bundle of microlocal solutions (depending on the energy parameter E) has trivial holonomy. Such a semi-classical treatement reveals interesting continuous symetries related to monodromy. Details will appear elsewhere.

#### 1. Bogoliubov-de Gennes Hamiltonian

BdG Hamiltonian describes the dynamics of a pair of quasi-particles electron/hole in the Theory of Supraconductivity [2]. We consider a narrow metallic 1-D wire (Normal Metal N) connected to Supraconducting bulks S through a SNS junction, and compute the excitation spectrum in the normal contact region as a function of gate voltage, when electronic levels transform into phase sensitive Andreev levels. The wire, or lead, is identified with a 1-D structure, the interval  $x \in [-L, L]$  (case of a perfect junction) or  $x \in [-L + \ell/2, L - \ell/2]$  ("dirty junction"), where  $\ell \ll L$ . The reference energy in the lead is Fermi level  $E_F$ . The pair electron/hole is acted upon by two kinds of potentials:

(1) the "order parameter"  $\Delta(x)$  times a phase function  $e^{i\phi(x)/2}$ , which is the potential due to Cooper pairs in the supraconducting bulk. This potential, subject to self-consistency relations, is priori unknown. Namely, inside S,  $\Delta(x)e^{i\phi(x)/2}$  is a solution of Ginzburg-Landau (or Pitaevskiy) equations, and shows typically a vortex profile (in 2-D). In BdG Hamiltonian it is assumed, however, that  $\Delta(x)e^{i\phi(x)/2}$  is an "effective" potential. Inside N, superconducting gap  $\Delta(x) \equiv 0$ : quasi-particles live in the "clean metal". For  $|x| \geq L + \ell$ ,  $\Delta(x) = \Delta_0 > 0$ .

We assume that the phase function  $\phi(x)$  is constant near the junction, and gauge the interaction by  $\phi_{-} = -\phi_{+} = -\phi$  in the superconducting banks, so that  $\phi(x) = \operatorname{sgn}(x)\phi$ . We

assume further that this equality holds everywhere: since  $\Delta(x) = 0$  inside N, the discontinuity of  $x \mapsto \phi(x)$  is irrelevant.

(2) a smooth chemical potential  $\mu(x)$ : typically  $\mu(x)$  is flat in N and drops smoothly to the band bottom in the superconducting banks S. In our model we assume again  $\mu(x)$  to be constant in the superconducting bank, i.e.  $\mu(x) = \mu_0$  when  $|x| \ge L + \ell$ . Andreev currents at energy E occur only if  $\mu(x) \ge E$  in [-L, L].

The case of a perfect junction ( $\Delta$  "hard-wall potential") has been considered in [5], see also [4] for a SFS junction, and makes use scattering matrix techniques. In this work, justifying semiclassical techniques as in [8] (also in the multi-dimensional case) we rather consider an imperfect (or "dirty") junction:  $\Delta(x)e^{i\phi(x)/2}$  is a smooth function. In a neighborhood of [-L, L], say  $x \in [-L - \ell, L + \ell]$ , the system is described at the classical level by BdG Hamiltonian

$$\mathcal{P}(x,\xi) = \begin{pmatrix} \xi^2 - \mu(x) & \Delta(x)e^{i\phi(x)/2} \\ \Delta(x)e^{-i\phi(x)/2} & -\xi^2 + \mu(x) \end{pmatrix}$$
(1)

The energy surface:  $\Sigma_E = \{\det(\mathcal{P} - E) = -(\xi^2 - \mu(x))^2 - \Delta(x)^2 + E^2 = 0\} = \Lambda_E^{\leq} \cup \Lambda_E^{\geq}$  splits into 2 branches separated in momentum space, so consists of two microlocal wells. Interaction between these wells gives the imaginary parts of the resonances for the electron/hole scattering, and will be ignored in this paper. Because of smoothness of  $x \mapsto \Delta(x)$ , the reflections occur inside [-L, L], we denote by  $(\pm x_E, \xi_E) \in \Lambda_E^{\geq}$ , the one-parameter family of "branching points" defined by  $\Delta(\pm x_E) = E$  with  $x_E$  near  $x_0 \in [L - \frac{\ell}{2}, L + \frac{\ell}{2}], \Delta(x_0) > 0$ . We do not consider the problem of "clustering" of eigenvalues as  $E \to 0 = E_F$  (Fermi level). In the "hard wall potential" limit for x near  $x_0$ , the potential  $\Delta(x)$  can be safely approximated by a linear function such that  $\Delta(x_0) = E_0$ , and  $\mu(x)$  by a constant  $\mu$ . So near  $x_0$  we assume that

$$\phi(x) = \phi, \quad \mu(x) = \mu > E, \quad \Delta(x) = E + \alpha(x - x_E)$$

for large  $\alpha > 0$ . Condition  $a_E = (x_E, \xi_E) \in \Sigma_E$  gives  $\xi_E^2 = \mu > E$ ,  $\Delta(x_E) = E$ .

The physical mechanism goes roughly as follows (see [5] for a detailed exposition): An electron  $e^-$  moving in the metallic lead, say, to the right, with energy  $0 < E \leq \Delta$  below the gap and kinetic energy  $K_+(x) = \mu(x) + \sqrt{E^2 - \Delta(x)^2}$  is reflected back as a hole  $e^+$  from the supraconductor, injecting a Cooper pair into the superconducting contact. The hole has kinetic energy  $K_-(x) = \mu(x) - \sqrt{E^2 - \Delta(x)^2}$ , and a momentum of the same sign as this of the electron. When  $\inf_{[-L,L]} K_-(x) > 0$  it bounces along the lead to the left and picks up a Cooper pair in the supraconductor, transforming again to the original electron state, a process known as Andreev reflection. This works also the other way in  $\Lambda_E^<$ , since Hamiltonian system conserves both charge and energy. Actually, the hole can propagate throughout the lead only if  $\inf_{[-L,L]} \mu(x) \geq E$ . Otherwise, it is reflected from the potential  $\mu(x)$  in the junction, and Andreev levels are quenched at higher energies, i.e. transform into localized electronic states.

For a rescaled "Planck constant" h so that  $h \ll \ell$ , we consider Weyl h-quantization of BdG Hamiltonian  $\mathcal{P}(x, hD_x)$  on  $L^2(I) \otimes \mathbb{C}^2$ ,  $I = [-(L+\ell), L+\ell]$ , which is self-adjoint when imposing Dirichlet boundary conditions at  $\partial I$ . Phase-sensitive Andreev states carry supercurrents that turn out to be proportional to the  $\phi$ -derivative of the eigen-energies of  $\mathcal{P}(x, hD_x)$ .

We have 
$$\sigma^y \mathcal{P}(\phi) \sigma^y = -\mathcal{P}(-\phi)$$
, with  $\sigma^y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ , accounting for "negative energies". We

shall assume here E > 0. When potentials are even functions (typical for metals),  $\mathcal{P}(x, hD_x)$  verifies PT symmetry  ${}^{\vee}\mathcal{IP}(x, hD_x) = \mathcal{P}(x, hD_x)\mathcal{I}^{\vee}$  which is essential for our approach to work.

At least formally, since BdG is only defined locally near N, removing boundary conditions leads to "resonances" (i.e. metastable states or quasi-particles with a finite life-time). Thus for simplicity we have assumed that (1), together with its semi-classical quantization, describes the system not only in *I*, but on the whole real line, provided  $h \ll \ell \ll L$ . Thus  $\mathcal{P}(x, hD_x)$  extends to  $L^2(\mathbf{R}) \otimes \mathbf{C}^2$ , Our general goal is to give a precise mathematical meaning to these "resonances". Here we content to compute their real parts through Bohr-Sommerfeld quantization rules.

# 2. Monodromy operator, scattering matrix: an outlook

a) Schrödinger operator on the real line.

We first recall from [1] basic facts for a 1-D Schrödinger operator with a compactly supported potential V. The generalized wave-functions u with energy  $E = k^2 > 0$  satisfy

$$-h^{2}u''(x) + V(x)u(x) = Eu(x)$$
(2)

and outside supp V,

$$-h^2 u''(x) = k^2 u(x) \tag{3}$$

defines the state space  $\mathcal{Z} \approx \mathbb{C}^2$  of the "free particle", spanned by  $f_1(x) = e^{ikx/h}$ ,  $f_2(x) = e^{-ikx/h}$ . The monodromy operator  $M(k) : f_1 + Bf_2 \mapsto Af_1$  is such that

$$M(k) = \begin{pmatrix} 1/\overline{A} & -\overline{B}/\overline{A} \\ -B/A & 1/A \end{pmatrix} \in \mathrm{SU}(1,1)$$

In particular,  $|A|^2 + |B|^2 = 1$ . We call  $|A|^2$  the transmission coefficient and  $|B|^2$  the reflection coefficient. Along with the passage from the left to the right of the support of V, consider the passage from the right to the left. The corresponding solution v of (2) is  $e^{-ikx/h} + B_2e^{ikx/h}$  to the right of suppV, and  $A_2e^{-ikx/h}$  to the left. The scattering matrix is defined as

$$S(k) = \begin{pmatrix} A & B \\ -\overline{B}A/\overline{A} & A \end{pmatrix} \in \mathrm{U}(2)$$

S(k) remains unitary and symmetric for complex values of k. Resonances of (2) are then defined as  $E = k^2 \in \mathbb{C}$ , where k is a pole of S, and physical resonances those with Imk > 0. Thus E is a resonance iff the solution of (3) is purely outgoing as  $x \to +\infty$  and  $x \to -\infty$ . The poles coincide with the poles of meromorphic extension of the resolvent  $(P - k^2)^{-1}$  from the physical half-plane ImE < 0 to the second sheet ImE > 0.

# b) Monodromy matrix for BdG equation: heuristics.

Now we discuss BdG equation  $(\mathcal{P}(x,hD_x) - E)U = 0$  for large |x|, i.e. (within our approximation above) when  $|x| \ge L + \ell$ , so  $\Delta(x) = \Delta_0$ ,  $\mu(x) = \mu_0 > E$ . Solutions are of the form

$$U(x;h) = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} e^{ikx/h} \\ e^{i\ell x/h} \end{pmatrix}$$

 $\mu_0 + E \pm i\Delta_0 \in \{k^2, \ell^2\}$ , so eigenfrequencies are  $(\pm k, \pm \overline{k})$ ,  $k = \sqrt{\mu_0 + E + i\Delta_0}$ , and the corresponding solutions as follows:

Let  $\phi(x) = \operatorname{sgn}(x)\phi$ ,  $\mathcal{Z}$  be the 2-D complex line bundle spanned by  $F_1^{\pm}(x) = \binom{e^{i\phi(x)/2}}{-i}e^{\pm ikx/h}$ (associated with the scattering process  $e^+ \to e^-$ ), and  $\overline{\mathcal{Z}}$  the 2-D complex line bundle spanned by  $F_2^{\pm}(x) = \binom{e^{i\phi(x)/2}}{i}e^{\pm i\overline{k}x/h}$  (associated with the scattering process  $e^- \to e^+$ ).

The space of solutions of exponential type for BdG is  $\mathcal{Z} \oplus \overline{\mathcal{Z}}$ , and  $\mathcal{Z}, \overline{\mathcal{Z}}$  are orthogonal for the usual pointwise Hermitian product in  $\mathbb{C}^2$ . Declare that  $E \in \mathbb{C}$  is a  $\mathcal{Z}$ -resonance iff the  $\mathcal{Z}$ -component of the wave function solving BdG equation is outgoing and evanescent ("physical solution") at infinity, i.e.

$$U(x,h) = A\binom{e^{i\phi/2}}{-i}e^{ikx/h}, x \to +\infty$$
$$U(x,h) = B\binom{e^{-i\phi/2}}{-i}e^{-ikx/h}, x \to -\infty$$

Similarly we say that E is a  $\overline{Z}$ -resonance iff the  $\overline{Z}$ -component of the wave function is outgoing (and evanescent) at infinity, i.e.

$$U(x,h) = A\binom{e^{i\phi/2}}{i}e^{-i\overline{k}x/h}, x \to +\infty$$
$$U(x,h) = B\binom{e^{-i\phi/2}}{i}e^{i\overline{k}x/h}, x \to -\infty$$

So for both sets of resonances, the corresponding solution is simultaneously decaying, and outgoing at  $\pm \infty$ . These sets of resonances need not coincide (although they come up in pairs), but their real parts are given by Bohr-Sommerfeld quantization rules. Namely, define the monodromy operator  $M^{\mathbb{Z}}(k)$  acting on  $\mathbb{Z}$  according to the formula

$$\binom{e^{-i\phi/2}}{-i}e^{ikx/h} + B\binom{e^{-i\phi/2}}{-i}e^{-ikx/h} \mapsto A\binom{e^{i\phi/2}}{-i}e^{ikx/h}$$

and similarly for  $M^{\overline{Z}}(k)$ . It is plausible to expect that  $M^{\mathcal{Z}}(k), M^{\overline{Z}}(k) \in U(1,1)$ , and that the corresponding scattering matrices  $S^{\mathcal{Z}}(k), S^{\overline{Z}}(k)$  have a meromorphic extension to the complex plane, their poles defining the resonances  $E^{\mathcal{Z}}$  and  $E^{\overline{Z}}$ . Actually, we shall construct "relative monodromy operators" in the "classically allowed region". In particular the relative monodromy operators are in U(1,1) for some specific Lorenzian form which is constructed below.

#### 3. Bohr-Sommerfeld quantization rules

In this work, we content to determine the real parts of the resonances, extending to this setting the method of positive commutators elaborated in [12], [9] and [10]. Imaginary parts may be determined as in [11]. We obtain Bohr-Sommerfeld quantization rules for the quasi-particle, alternating even and odd quantum numbers associated with the electron and the hole. In the sequel we will sketch a proof of the following result:

**Theorem 1**: Let  $\int_{-x_0}^{x_0} \eta^{\rho}(y;h) dy$  be the semi-classical actions (see Proposition 8 below)  $\rho = 1$  for the electron,  $\rho = -1$  for the hole. Bohr-Sommerfeld quantization conditions near  $E_0$  are given at first order by:

$$\oint_{\gamma_E} \eta^{\rho}(y;h) \, dy - h\phi + h\pi + \mathcal{O}(h^2) = 2\pi nh; \ n \in \mathbf{Z}$$

with even (resp. odd) quantum numbers n for the electron (resp. the hole). Here  $\oint_{\gamma_E}$  denotes integral over the loop  $\gamma_E$  obtained by gluing together  $\Lambda_E^>$  and  $\Lambda_E^<$ , if we ignore tunneling in momentum space.

### 4. Microlocal solutions in Fourier representation near the branching points

# a) Reduction of the system.

In *h*-Fourier representation,  $\mathcal{F}_h u(\xi) = (2\pi h)^{-1/2} \int e^{-ix\xi/h} u(x) dx$  the local Hamiltonian near  $a = a_E = (x_E, \xi_E)$ ,  $\mathcal{P}^a$  takes the form :

$$\mathcal{P}^{a}(-hD_{\xi},\xi) = \begin{pmatrix} \xi^{2} - \mu & e^{i\phi/2}(E - \alpha hD_{\xi} - \alpha x_{E}) \\ e^{-i\phi/2}(E - \alpha hD_{\xi} - \alpha x_{E}) & -\xi^{2} + \mu \end{pmatrix}$$
(4)

By PT symmetry  $\mathcal{P}^{a'} = \mathcal{I}\mathcal{P}^{a}\mathcal{I}$  near  $a' = a'_{E} = (-x_{E}, \xi_{E})$ . Solving the system  $\mathcal{P}^{a}(-hD_{\xi}, \xi)\widehat{U} = 0$ ,  $\widehat{U} = \begin{pmatrix}\widehat{\varphi}_{1}\\\widehat{\varphi}_{2}\end{pmatrix}$  gives second order ODE for  $u(\xi) = \exp[-i\int^{\xi}g(s)ds/h]\widehat{\varphi}_{2}(\xi)$ ,

$$P^{a}(-hD_{\xi},\xi,h)u(\xi) = \frac{E^{2}}{\alpha^{2}}u(\xi)$$
(5)

$$P^{a}(-hD_{\xi},\xi,h) = (hD_{\xi})^{2} + \alpha^{-2}(\xi^{2} - \mu)^{2} + h^{2}(\xi^{2} - \mu - E)^{-2}(2\xi^{2} + \mu + E)$$

After *E*-dependent scalings  $\beta = \sqrt{\alpha}(2\xi_E)^{-3/2} > 0$ ,  $E_1 = (2\xi_E)^{-2}E$ ,  $\xi = 2\xi_E\beta\omega\xi' + \xi_E$ ,  $\omega = \pm 1$  ( $\xi'$  is "local momentum") we obtain  $P^a_{\omega}(-hD_{\xi'},\xi',h)u_{\omega}(\xi') = \left(\frac{E_1}{\beta}\right)^2 u_{\omega}(\xi')$ , where

$$P^{a}_{\omega}(-hD_{\xi'},\xi';h) = (-hD_{\xi'})^{2} + (\xi' + \beta\omega\xi'^{2})^{2} + h^{2}\beta^{2}f(\omega\beta\xi')$$

is an anharmonic Schrödinger operator. The lower order term  $f(z) = (2z^2 + 2z + \frac{3}{4} + E_1)(z^2 + z - E_1)^{-2}$  has a pole on  $\Lambda_E^>$  where the linear approximation of  $\Delta(x)$  breaks down. The linear approximation only holds for small  $\xi'$ . Consider the map

$$\iota^{a}: \sum_{\omega=\pm 1} \operatorname{Ker}_{h}(P_{\omega}^{a} - \left(\frac{E_{1}\omega}{\beta}\right)^{2}) \to \operatorname{Ker}_{h}(\mathcal{P}^{a} - E)$$
(6)

where  $\operatorname{Ker}_h$  denotes the microlocal kernel. The index  $\omega$  is to be chosen carefully with the complex germ of solutions having the right decay beyond the branching points  $\pm x_E$ . We shall endow the RHS of (6) with a Lorenzian structure and "diagonalize"  $\iota^a$  in some orthogonal subspaces.

# b) The normal form of Helffer-Sjöstrand

When  $E_1 < \frac{1}{4}$ , we take  $P^a_{\omega}$  microlocally to its normal form, namely:

**Proposition 2** [9]: There exists an analytic diffeomorphism  $t \mapsto F_0(t)$  defined in a neighborhood of 0,  $F_0(0) = 0$ , with inverse  $G_0$ , and a real analytic phase function  $\phi_\beta(\xi',\theta)$ , defined in a neighborhood of (0,0), of the form  $\phi_\beta(\xi',\theta) = \xi'\theta + g_\beta(\xi',\theta), g_\beta(\xi',\theta) = \mathcal{O}(|\xi',\theta|^3)$ , parametrizing the canonical transformation  $\kappa_\beta : (\partial_\theta \phi_\beta, \theta) \mapsto (\xi', \partial_{\xi'} \phi_\beta)$ , such that  $F_0 \circ p_\beta \circ \kappa_\beta = p_0$ . At the semi-classical level, there is a (formally) unitary FIO operator A defined microlocally near (0,0)

$$Av(\xi',h) = (2\pi h)^{-1} \int \int e^{i\varphi(\xi',\eta,\theta)/h} c(\xi',\eta,\theta,h) e^{ib(\xi',\eta,\theta,h)} v(\eta,h) \, d\eta d\theta$$

and a real valued analytic symbol

$$F(t,\beta,h) = F_0(t,\beta) + hF_1(t,\beta) + h^2F_2(t,\beta) + \cdots$$

with  $F_1(t,\beta) = -\frac{1}{2}$  such that

$$A^*F(P_{\omega},\beta,h)A = P_0(\eta,hD_{\eta}) = \frac{1}{2}((hD_{\eta})^2 + \eta^2 - h), \quad A^*A \equiv \mathrm{Id}$$

The function  $F_0$ , taking the period T(E) of Hamilton vector flow for  $P^a_{\omega}$  at energy  $(E_1/\beta)^2$  to  $2\pi$ , involves an elliptic integral, which requires sometimes the use of formal calculus.

#### c) Weber equation and parabolic cylinder functions

Weber equation  $P_0 v = \nu h v$ , through change of variables  $\eta = (h/2)^{1/2} \zeta$ ,  $\tilde{v}(\zeta) = v(\eta)$  scales to

$$-\widetilde{v}'' + \frac{1}{4}\zeta^2 \widetilde{v} = \left(\nu + \frac{1}{2}\right)\widetilde{v}$$

Fundamental solutions express as parabolic cylinder functions  $D_{\nu}$ , entire in **C**. The systems  $(D_{\nu}(\pm\zeta), D_{-\nu-1}(\pm i\zeta))$  are fundamental solutions for any choice of  $\pm$ . Integral representations give asymptotic solutions of  $(P_0 - \nu h)u(\eta) = 0$  by stationary phase for real  $\nu$ ,  ${E'}_1^2 = 2\beta^2 F(\beta^{-2}E_1^2, \beta, h) = 2\beta^2(\nu+1)h$ .

$$D_{\nu}(\varepsilon(h/2)^{-1/2}\eta) = \frac{\Gamma(\nu+1)}{-2i\pi\sqrt{h}}h^{E^2/4h} \int_{\infty}^{(0^+)} \exp\left[i\Phi_{\varepsilon}^{\nu}(s;\eta)/h\right] ds$$
$$D_{-\nu-1}(i\varepsilon(h/2)^{-1/2}\eta) = \frac{\Gamma(-\nu)}{2i\pi}h^{-E^2/4h} \int_{\infty}^{(0^+)} \exp\left[i\Phi_{\varepsilon}^{-\nu-1}(s;\eta)/h\right] \frac{ds}{s}$$

with  $\varepsilon = \pm 1$ ,  $E = \sqrt{2(\nu + 1)h}$ , see [13]. This normalization is called Whittaker normalization. Classically forbidden regions  $|\eta| > E$  lie on Stokes lines, classically allowed region  $|\eta| < E$  in between, and 3 Stokes lines stem from each "turning point"  $\eta = \pm E$ .

#### d) Microlocal solutions.

We apply asymptotic stationary phase to  $AD_j$ ,  $j \in \{\nu, -\nu - 1\}$ . With  $h' = \beta^2 h$  as a "rescaled" Planck constant, we get:

**Proposition 3:** In Fourier representation, the image  $K_h^a(E) = \operatorname{Ker}_h(\mathcal{P}^a(-hD_{\xi},\xi)-E)$  of  $\iota^a$  is a 2-D vector space spanned by the spinors  $\widehat{U}_{\varepsilon,\omega}^j = \left(\widehat{\varphi}_2^{j}\right)_{\varepsilon,\omega}^j$ ,  $(j,\varepsilon,\omega) \in \{\nu, -\nu - 1\} \times \{-1,1\}^2$ , of the form:

$$\begin{split} \widehat{U}_{\varepsilon,\omega}^{\nu} &= C_{h'}^{\nu} \sum_{\theta_{\omega} = \pm \widehat{\theta}_{\omega}(\xi_{1})} {e^{i\phi/2} (\xi^{2} - \mu - E)^{-1/2} X_{\varepsilon,\omega}^{\nu} \choose |\widetilde{a}_{\varepsilon,\omega}^{\nu}|} \exp[i(\Phi_{\varepsilon,\omega}^{\nu} + h' R_{\omega}^{\nu})/h'] + \mathcal{O}(h') \\ \widehat{U}_{\varepsilon,\omega}^{-\nu-1} &= C_{h'}^{-\nu-1} \sum_{\theta_{\omega} = \pm \widehat{\theta}_{\omega}(\xi_{1})} \varepsilon \operatorname{sgn}(\theta_{\omega}) {e^{i\phi/2} (\xi^{2} - \mu - E)^{-1/2} X_{\varepsilon,\omega}^{-\nu-1} \choose (\xi^{2} - \mu - E)^{1/2}} \\ &|\widetilde{a}_{\varepsilon,\omega}^{-\nu-1}| \exp[i(\Phi_{\varepsilon,\omega}^{-\nu-1} + h' R_{\omega}^{-\nu-1})/h'] + \mathcal{O}(h') \end{split}$$

Here  $\hat{\theta}_{\omega}(\xi_1)$  is a critical point (from stationary phase),  $\Phi^j_{\varepsilon,\omega} + h'R^j_{\omega}$ ) the h'-dependent phase functions, and  $X^j_{\varepsilon,\omega}$ ,  $|\tilde{a}^j_{\varepsilon,\omega}|$  some positive amplitudes. Spinors  $U^j_{\varepsilon,\omega}$  verify the symmetry  $^{\dagger}\hat{U}^j_{-\varepsilon,-\omega} = \hat{U}^j_{\varepsilon,\omega}$  for the "local time" reversal operator  $^{\dagger}u(\xi_1) = u(-\xi_1)$ , and the constants  $C^j_{h'}$  (from Whittaker normalization of  $D_{\nu}$ ,  $D_{-\nu-1}$ ) are related by  $C^{\nu}_{h'}C^{-\nu-1}_{h'} = \left((2\sqrt{h'})^3\pi^2\sin\pi\nu\right)^{-1}$ .

#### 5. Normalization

# a) The microlocal Wronskian.

We extend to BdG Hamiltonian the classical "positive commutator method" using conservation of some quantity called a "quantum flux" ([12], [9], [11], [10]).

**Definition 4:** Let  $\mathcal{P}$  be (formally) self-adjoint, and  $U^a, V^a \in K_h(E)$  be supported on  $\Lambda_E^>$ . We call the sesquilinear form  $\mathcal{W}^a_\rho(U^a, V^a) = \left(\frac{i}{\hbar}[\mathcal{P}, \chi^a]_\rho U^a | V^a\right) = \left(\frac{i}{\hbar}[\mathcal{P}, \chi^a]_\rho \widehat{U}^a | \widehat{V}^a\right)$  the microlocal Wronskian of  $(U^a, \overline{V^a})$  in  $\omega^a_\rho$ . Here  $\frac{i}{\hbar}[\mathcal{P}, \chi^a]_\rho$  denotes the part of the commutator supported microlocally on  $\omega^a_\rho$  (a small neighborhood of supp $[\mathcal{P}, \chi^a] \cap \Lambda_E$  near  $\rho$ ).

A crucial property of the microlocal Wronskian is to be invariant by Fourier transformation:  $\mathcal{W}^a_{\rho}(U^a, V^a) = \mathcal{W}^a_{\rho}(\hat{U}^a, \hat{V}^a)$ . The relation  $\mathcal{W}^a_+(U^a, V^a) + \mathcal{W}^a_-(U^a, V^a) = 0$  doesn't readily follow as in the scalar case [10], the microlocal solutions being neither smooth in spatial of Fourier representation near the branching point, but from a careful inspection, involving also formal calculus. This is used essentially in Propositions 5 and 8 below. Choosing  $\varepsilon, \omega$  such that  $\varepsilon \omega = 1$  we define a Lorenzian metric  $\mathcal{W}_{\rho}$  on the space of microlocal solutions near a. In the basis  $\widehat{U}_{\varepsilon,\omega}^{j}, j \in \{\nu, -\nu - 1\}$  we have, up to a constant factor:

$$\rho \mathcal{W}_{\rho} = \begin{pmatrix} |C_{h'}^{\nu}|^2 \mathcal{O}(h') & C_{h'}^{\nu} \overline{C_{h'}^{-\nu-1}} \exp[-i\pi E_1'^2/4h'] (1 + \mathcal{O}(h')) \\ \overline{C_{h'}^{\nu}} \overline{C_{h'}^{-\nu-1}} \exp[i\pi E_1'^2/4h'] (1 + \mathcal{O}(h')) & |C_{h'}^{-\nu-1}|^2 \mathcal{O}(h') \end{pmatrix}$$

Changing Whittaker normalization for the  $D_{\nu}, D_{-\nu-1}$  functions, and the microlocal solutions by some constant phase factors, we can reduce to  $\rho \mathcal{W}_{\rho} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \mathcal{O}(h')$ , and prove:

**Proposition 5:** Under PT symmetry above the microlocal Wronskians  $\mathcal{W}^a_{\rho}$  endow  $K^a_h(E)$  (mod h') with a Lorenzian form  $\mathcal{W}^a = \frac{1}{2}(\mathcal{W}^a_+ - \mathcal{W}^a_-)$ . The same holds at a', and the corresponding structures on  $K^a_h \times K^{a*}_h$  and  $K^{a'}_h \times K^{a'*}_h$  are anti-isomorphic. The group of automorphisms preserving  $\mathcal{W}^a$  and  $\mathcal{W}^{a'}$  mod  $\mathcal{O}(h')$  is therefore U(1,1).

# 6. Spinors in the spatial representation

We compute  $U_{\varepsilon,\omega}^{a,j}, U_{\varepsilon,\omega}^{a',j}$  in spatial representation, then extend along the branches  $\rho = \pm 1$  of  $\Lambda_E^>$  with WKB solutions.

# a) Spinors near the branching points.

Near a, a' we apply inverse *h*-Fourier transform and get:

Proposition 6: Up to a constant phase factor

$$\begin{aligned} U_{\varepsilon,\omega}^{\nu}(x,h) &= 2\omega\beta\xi_E e^{ix\xi_E/h} \sum_{\rho=\pm} {e^{i\phi/2}(\xi^2 - \mu - E)^{-1/2}X_{\varepsilon,\omega}^{\nu}} |a_{\varepsilon,\omega}^{\nu}||_{\theta_1 = \theta_{\omega}(\xi_1), \xi_1 = \xi_{\omega}^{\rho}(x)} \\ &\times \left(\frac{L_{\omega}^{\rho}(x)}{i}\right)^{-1/2} \exp[i\left(\Psi_{\varepsilon,\omega}^{\nu,\rho}(x) + h'R_{\varepsilon,\omega}^{\nu,\rho}(x)\right)/h'](1 + \mathcal{O}(h')) \\ U_{\varepsilon,\omega}^{-\nu-1}(x,h) &= 2\omega\beta\xi_E e^{ix\xi_E/h} \sum_{\rho=\pm} \varepsilon \operatorname{sgn}(\theta_1) {e^{i\phi/2}(\xi^2 - \mu - E)^{-1/2}X_{\varepsilon,\omega}^{-\nu-1}} |\tilde{a}_{\varepsilon,\omega}^{-\nu-1}||_{\theta_1 = \theta_{\omega}(\xi_1), \xi_1 = \xi_{\omega}^{\rho}(x)} \\ &\times \left(\frac{L_{\omega}^{\rho}(x)}{i}\right)^{-1/2} \exp[i\left(\Psi_{\varepsilon,\omega}^{\nu,\rho}(x) + h'R_{\varepsilon,\omega}^{\nu,\rho}(x)\right)/h'](1 + \mathcal{O}(h')) \end{aligned}$$

Here  $(L_{\omega}^{\rho}(x))^{-1/2}$  is a real density (singular at  $x = x_E$ ), and  $\rho$  labels the branch of the Lagrangian manifold. The phases  $\Psi_{\varepsilon,\omega}^{j,\rho}(x) + h' R_{\varepsilon,\omega}^{j,\rho}(x), j \in \{\nu, -\nu - 1\}$  differ only by a constant.

# b) WKB spinors away from the branching points

The Lagrangian manifold  $\Lambda_E^>$  consists of 2 branches  $\Lambda_E^{>,\rho}$  (or simply  $\rho$ )  $\rho = \pm 1$  so that  $\rho = +1$  belongs to the electronic state ( $\xi_1 > 0$  in the local coordinates near a above), resp.  $\rho = -1$  to the hole state ( $\xi_1 < 0$ ). These states mix up when  $\Delta(x) \neq 0$ , but we can sort them out semiclassically, outside a, a'. Call the vector space of  $\mathbf{C}^2$  generated by  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$  the space of (pure) electronic states, or electronic spinors, and this by  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$  the space of (pure) hole states, or hole spinors.

The principal symbol  $\mathcal{P}(x,\xi)$  has eigenvalues  $\lambda_{\rho} = \rho \lambda(x,\xi) = \rho \sqrt{\Delta(x)^2 + (\xi^2 - \mu(x))^2}$ . By diagonalizing, we obtain a line bundle  $\Lambda_E^{\rho}$  with fiber

$$Y_{\rho}(x,\xi) = (\Delta^2 + (-\xi^2 + \mu + \rho\sqrt{\Delta^2 + (\xi^2 - \mu)^2})^2)^{-1/2} \begin{pmatrix} \Delta e^{i\phi/2} \\ -\xi^2 + \mu + \rho\sqrt{\Delta^2 + (\xi^2 - \mu)^2} \end{pmatrix}^2 = (\Delta^2 + (-\xi^2 + \mu + \rho\sqrt{\Delta^2 + (\xi^2 - \mu)^2})^2)^{-1/2} \begin{pmatrix} \Delta e^{i\phi/2} \\ -\xi^2 + \mu + \rho\sqrt{\Delta^2 + (\xi^2 - \mu)^2} \end{pmatrix}^2 = (\Delta^2 + (-\xi^2 + \mu + \rho\sqrt{\Delta^2 + (\xi^2 - \mu)^2})^2)^{-1/2} \begin{pmatrix} \Delta e^{i\phi/2} \\ -\xi^2 + \mu + \rho\sqrt{\Delta^2 + (\xi^2 - \mu)^2} \end{pmatrix}^2 \end{pmatrix}^2$$

Looking at the electronic state, we choose  $\rho = +1$  so that  $\lambda_{\rho}(x_{\rho}, \xi_{\rho}) - E = 0$ , while  $\lambda_{-\rho}(x_{\rho}, \xi_{\rho}) - E$  is elliptic. and similarly when looking at the hole state.

**Proposition 7** The microlocal kernel  $\operatorname{Ker}_h(\mathcal{P} - E)$  on  $\Lambda_E^{>,\rho}$  is one-dimensional space spanned by

$$W^{\rho}(x,h) = e^{iS_{\rho}(x,h)/h} \left( w_0^{\rho}(x,h) Y_{\rho}(x,\partial_x S_{\rho}) + \mathcal{O}(h) \right) = e^{iS_{\rho}(x,h)/h} \widetilde{W}^{\rho}(x,h)$$

where  $w_0^{\rho}(x)|dx|^{1/2}$  is a smooth half-density. By the uniqueness property of WKB solutions along simple bicharacteristics, the h (or h')-dependent phase function  $S_{\rho}(x,h)$  should coincide, up to a constant (in a punctured neighborhood of a) with either one of  $\Psi_{\varepsilon,\omega}^{j,\rho}(x) + h' R_{\varepsilon,\omega}^{j,\rho}(x)$ above,  $j \in \{\nu, -\nu - 1\}$ , and similarly for the half-densities.

### 7. Relative monodromy matrices

Now we look for connexion formulas. For each  $\varepsilon, \omega, \rho = \pm 1, j \in \{\nu, -\nu - 1\}$ , the normalized microlocal solutions  $U^{a',j,\rho}_{\varepsilon,\omega}$  are related to the extension  $U^{a,k,\rho}_{-\varepsilon,-\omega,\text{ext}}$  of the normalized microlocal solutions  $U^{a,k,\rho}_{\varepsilon,\omega}$  along the bicharacteristics by a monodromy matrix

$$\mathcal{M}^{a,a',\rho} = \begin{pmatrix} d^{\rho}_{11} & d^{\rho}_{12} \\ d^{\rho}_{21} & d^{\rho}_{22} \end{pmatrix} \in U(1,1)$$

(defined at least mod  $\mathcal{O}(h')$ ) which we call a *relative monodromy matrix*. Since there is a pair of particles, the symmetry between the  $\mathcal{M}^{a,a',\rho}$  and  $\mathcal{M}^{a',a,\rho}$  is order 4;  $\mathcal{M}^{a',a,\rho} \in U(1,1)$  is obtained by extending from the left to the right, and applying symmetry

$$\rho \mathcal{M}^{a',a,\rho} = \mathcal{I}(\mathcal{M}^{a,a',\rho})^{-1} \mathcal{I} =, \quad \rho = \pm 1$$
(7)

where  $\mathcal{I}$  denotes complex conjugation. We compute the coefficients  $d_{ij} = d_{ij}^{\rho}$ . Considering behavior of  $U_{e,\omega}^{a',j,\rho}$  in the classically forbidden region (according to scattering process  $e^+ \to e^$ or  $e^- \to e^+$ ) we obtain

$$\mathcal{M}^{a,a',\rho} = \begin{pmatrix} 0 & d_{12}^{\rho} \\ d_{21}^{\rho} & 0 \end{pmatrix}, \quad \overline{d_{12}^{\rho}} \, d_{21}^{\rho} = 1$$

Note that if we do not look too closely at the relevant complex branches, as is the case when computing BS, it makes no difference to choose instead  $\mathcal{M}^{a,a',\rho} = \begin{pmatrix} d_{11}^{\rho} & 0\\ 0 & d_{22}^{\rho} \end{pmatrix}$ , with  $\overline{d_{11}}^{\rho} d_{22}^{\rho} = 1$ .

As in [12], [9], [11], [10], the argument consists now in extending microlocal solutions obtained above from a to a', and computing the resulting semi-classical action. So take first  $U_1$  equal to  $U_1^a = U_{\varepsilon,\omega}^{\nu,a}$  near a, extend it along to a' along the bicharacteristics  $\rho = \pm 1$  by WKB. Evaluating on  $\rho$  near a' we find  $U_1^{a',\rho} = U_{\varepsilon,\omega,\text{ext}}^{\nu,a,\rho} = d_{21}^{\rho}U_{\varepsilon,\omega}^{-\nu-1,a',\rho}$ . Similarly, take  $U_2$  starting at a' and with  $-\nu - 1$  instead of  $\nu$ , we get  $U_1^{a,\rho} = U_{\varepsilon,\omega,\text{ext}}^{-\nu-1,a',\rho} = e_{12}^{\rho}U_{\varepsilon,\omega}^{\nu,a,\rho}$ , where  $e_{12}^{\rho} = \rho(d_{21}^{\rho})^{-1}$  is the matrix element of  $\mathcal{M}^{a',a,\rho}$  given in (7). We compute  $d_{21}^{\rho}$  in two different ways and compare the result.

(1) Using time-reversal and PT symmetries in the microlocal Wronskians, we get

$$\begin{pmatrix} \frac{i}{h} \left[ \mathcal{P}^{a'}, \chi^{a'} \right]_{\rho} U_1 | U_{\varepsilon,\omega}^{\nu} \rangle = d_{21}^{\rho} \left( \frac{i}{h} \left[ \mathcal{P}^{a'}, \chi^{a'} \right]_{\rho} U_{\varepsilon,\omega}^{-\nu-1} | U_{\varepsilon,\omega}^{\nu} \right) = \\ = d_{21}^{\rho} \mathcal{W}_{\rho}^{a'} \left( U_{\varepsilon,\omega}^{-\nu-1}, U_{\varepsilon,\omega}^{\nu} \right) = d_{21}^{\rho} \mathcal{W}_{\rho}^{a'} \left( \widehat{U}_{\varepsilon,\omega}^{-\nu-1}, \widehat{U}_{\varepsilon,\omega}^{\nu} \right) = \\ = -d_{21}^{\rho} \overline{\mathcal{W}_{\rho}^{a}} \left( \widehat{U}_{-\varepsilon,-\omega}^{-\nu-1}, \widehat{U}_{-\varepsilon,-\omega}^{\nu} \right) = -d_{21}^{\rho} \mathcal{W}_{\rho}^{a} \left( \widehat{U}_{-\varepsilon,-\omega}^{\nu}, \widehat{U}_{-\varepsilon,-\omega}^{-\nu-1} \right) = -d_{21}^{\rho} \end{aligned}$$

(2) Using the extensions described in Proposition 7. Near a' we have  $U_{1,\text{ext}}^{\rho} = e^{i\phi/2}W^{\rho}(x,h) =$  $d_{21}^{\rho}U_{\varepsilon,\omega}^{-\nu-1,a',\rho}$  (by solving transport equation along  $\rho$  the amplitude picks up the phase factor  $e^{i\phi/2}$ ), so we need to compute  $\left(\frac{i}{h}\left[\mathcal{P}^{a'},\chi^{a'}\right]_{\rho}W^{\rho}(x,h)|U_{\varepsilon,\omega}^{\nu}\right)$ . The amplitude  $W^{\rho}(x,h)$  is actually defined up to a real, constant factor  $C^{\rho}$ .

**Proposition 8:** Let  $\widetilde{\Psi}_{\varepsilon,\omega}^{\nu,a',\rho}(x) = x \xi_E + \frac{(2\xi_E)^3}{\alpha} \Psi_{\varepsilon,\omega}^{\nu,a',\rho}(x)$ . We have

$$\left(\frac{i}{h}\left[\mathcal{P}^{a'},\chi^{a'}\right]_{\rho}W^{\rho}|U^{\nu,a',\rho}_{\varepsilon,\omega}\right) = 2\,\widetilde{C}^{\rho}\,e^{i\pi/4}\int\exp\left[i\left(\widetilde{S}_{\rho}(x;h)/h\right]\beta(x,h)\,(\chi^{a'}_{1})'(x)\,dx\right]$$
(8)

where the amplitude  $\beta(x, h)$ , real mod  $\mathcal{O}(h)$ , is computed from the WKB solutions in Proposition 7, and

$$\widetilde{S}_{\rho}(x,h) = S_{\rho}(x;h) - \left(x\xi_E + \widetilde{\Psi}_{\varepsilon,\omega}^{\nu,a',\rho}(x) - h R_{-\omega}^{\nu} \left(\theta_{-\omega}(\xi_{-\omega}^{\rho}(-x))\right)\right) = \frac{(2\xi_E)^3}{\alpha} \Psi_{-\varepsilon,-\omega}^{\nu,a,\rho}(x_0) - \int_{-x_0}^{x_0} \eta^{\rho}(y;h) \, dy + h R_{-\omega}^{\nu} \left(\theta_{-\omega}(0)\right)$$

Moreover,  $\beta(x, h)$  is also independent of x, so that, comparing the former expression (1) and (8) for a suitable choice of  $C^{\rho}$ , we get

$$d_{21}^{\rho} = -e^{i\tau^{\rho}(h)/h} \int (\chi_1^{a'})'(x) \, dx = e^{i\tau^{\rho}(h)/h} \tag{9}$$

Here  $\tau^{\rho}(h) = h \frac{\phi}{2} + h \frac{\pi}{2} - \int_{-x_0}^{x_0} \eta^{\rho}(y;h) dy + \text{Const.}$ , where Const. is evaluated at the boundaries  $x = \pm x_E$ , and depends only on  $E'_1$ . It will eventually disappear from the final formula, by adding to BS the contribution of the lower branch  $\Lambda_E^{<,\rho}$ . Note that  $\int_{-x_0}^{x_0} \eta^{\rho}(y;h) dy$ ,  $\eta^{\rho}(y;h)$  being the derivative of the h'-depending phase function, is the semi-classical action.

8. Bohr-Sommerfeld quantization rules We set  $F_{\varepsilon,\omega}^{j,a,\rho} = \frac{i}{h} [\mathcal{P}^a, \chi^a]_{\rho} U_{\varepsilon,\omega}^{j,a,\rho}$ , and similarly with a'. The set  $\{G_{\varepsilon,\omega}^{j,\flat} = F_{\varepsilon,\omega}^{j,\flat,+} - F_{\varepsilon,\omega}^{j,\flat,-} : j \in \{\nu, -\nu - 1\}, \flat \in \{a, a'\}\}$  (or their *h*-Fourier transform) can be interpreted as a basis of the microlocal co-kernel of  $\mathcal{P}$  near a, a'. Following [10], we introduce Gram matrix  $\mathcal{G}^{\rho}$  of vectors  $\widehat{U}_1^{\rho}$ 

and  $\widehat{U}_{2}^{\rho}$  in this basis, namely  $\mathcal{G} = \begin{pmatrix} (\widehat{U}_{1}|\widehat{G}_{\varepsilon,\omega}^{-\nu-1,a}) & (\widehat{U}_{2}|\widehat{G}_{\varepsilon,\omega}^{-\nu-1,a}) \\ (\widehat{U}_{1}|\widehat{G}_{\varepsilon,\omega}^{\nu,a'}) & (\widehat{U}_{2}|\widehat{G}_{\varepsilon,\omega}^{\nu,a'}) \end{pmatrix}$ . Using symmetries we get

$$\mathcal{G} = \mathcal{G}^{\rho} = 2 \begin{pmatrix} 1 & e_{12}^{\rho} \\ -d_{21}^{\rho} & -1 \end{pmatrix}$$

The condition  $det(\mathcal{G}^{(\rho)}) = 0$  means that  $U_1$  is collinear to  $U_2$ , i.e. there is a global section of  $\operatorname{Ker}_{h}(\mathcal{P}-E)$ . Recall  $e_{12}^{\rho} = \rho(\overline{d_{21}^{\rho}})^{-1}$ ; for  $\rho = +1$  (electronic state) we get  $\operatorname{Im} d_{21}^{+} = 0$ , that is  $\sin\left(\frac{\tau^{(+)}(h)}{h}\right) = 0$ . We eventually obtain BS by "surgery": namely (ignoring tunneling) we cut and paste the half-bicharacteristic  $\Lambda_E^{>,+}$  in the upper-half plane  $\xi > 0$  with its symmetric part  $\Lambda_E^{<,-}$  in  $\xi < 0$  and add together the contributions. By symmetry, the constant term Const. in  $\tau^{+}(h)$  drops out, while the other terms  $h \frac{\phi}{2} + h \frac{\pi}{2} - \int_{-x_0}^{x_0} \eta^{\rho}(y;h) dy$  add up, which yields BS for the electronic state. We argue similarly for the hole state. This eventually gives Theorem 1.

Acknowledgements: We thank Timur Tudorovskiy for having introduced us to the problem. This work has been partially supported by the grant PRC CNRS/RFBR 2017-2019 No.1556 "Multidimensional semi-classical problems of Condensed Matter Physics and Quantum Dynamics".

- [1] Arnold V 1983 Geometrical methods in the theory of ordinary differential equations (Springer, Berlin)
- [2] Bardeen J, Cooper L and Schriefer J 1959 Phys. Rev. 108(5) 1175
- Bensouissi A, M'hadbi N and Rouleux M 2011 Proc. "Days of Diffraction 2011" (Saint-Petersburg) (IEEE 101109/DD.2011.6094362) 39
- [4] Cayssol J and Montambaux G 2004 Phys. Rev. B 70 224520
- [5] Chtchelkatchev N, Lesovik G and Blatter G 2000 Phys. Rev. B 62(5) 3559
- [6] de Gennes P G 1966 Superconductivity of Metals and Alloys (Benjamin, New York)
- [7] Gérard C and Sigal I M 1992 Comm. Math. Phys. 145 281
- [8] Duncan K P and Györffy B L 2002 Annals of. Phys. 298 273
- [9] Helffer B and Sjöstrand J 1989 Soc. Math. de France, Mémoire 39 117(4)
- [10] Ifa A, Louati H and Rouleux M 2018 J. Math. Sci. Univ. Tokyo 25 1
- [11] Rouleux M 1999 Tunneling effects for h-Pseudodifferential Operators, Feshbach resonances and the Born-Oppenheimer approximation (Adv. Part. Diff. Eq. 16) ed M Demuth, E Schrohe et al. (Wiley VCH, Berlin)
- [12] Sjöstrand J 1990 Density of states oscillations for magnetic Schrödinger operators Proc. Diff. Eq. Math. Phys. (Univ. Alabama, Birmingham) ed Bennewitz
- [13] Whittaker E T and Watson G N 1980 A Course of Modern Analysis (Cambridge Univ.Press)