

GLOBAL CHANG'S CONJECTURE AND SINGULAR CARDINALS

MONROE ESKEW AND YAIR HAYUT

ABSTRACT. We investigate the possibilities of global versions of Chang's Conjecture that involve singular cardinals. We show some ZFC limitations on such principles, and prove relative to large cardinals that Chang's Conjecture can consistently hold between all pairs of limit cardinals below $\aleph_{\omega\omega}$.

1. INTRODUCTION

The Löwenheim-Skolem theorem asserts that for every pair of infinite cardinals $\kappa > \mu$ and every structure \mathfrak{A} on κ in a countable language, there is a substructure $\mathfrak{B} \subseteq \mathfrak{A}$ of size μ . “Chang's Conjecture” is a type of principle strengthening this theorem to assert similar relationships between sequences of cardinals. For example $(\kappa_1, \kappa_0) \rightarrow (\mu_1, \mu_0)$ says that for every structure \mathfrak{A} on κ_1 in a countable language, there is a substructure \mathfrak{B} of size μ_1 such that $|\mathfrak{B} \cap \kappa_0| = \mu_0$. The following basic observation puts some constraints on this type of principle:

Proposition 1. *Suppose $\kappa, \lambda \leq \delta$ and $\kappa^\lambda \geq \delta$. Then there is a structure \mathfrak{A} on δ such that for every $\mathfrak{B} \prec \mathfrak{A}$,*

$$|\mathfrak{B} \cap \kappa|^{|\mathfrak{B} \cap \lambda|} \geq |\mathfrak{B} \cap \delta|.$$

Corollary 2. *If $(\kappa_1, \kappa_0) \rightarrow (\mu_1, \mu_0)$, $\nu \leq \kappa_0$, and $\kappa_0' \geq \kappa_1$, then $\mu_0^{\min(\mu_0, \nu)} \geq \mu_1$.*

From this, we immediately see that under GCH, $(\kappa^+, \kappa) \rightarrow (\mu^+, \mu)$ can only occur when $\text{cf}(\kappa) \geq \text{cf}(\mu)$. (The consistency of contrary cases is unknown.) This inspires the following bold conjecture:

Definition (Global Chang's Conjecture). *We say that the Global Chang's Conjecture holds if for all infinite cardinals $\mu < \kappa$ with $\text{cf}(\mu) \leq \text{cf}(\kappa)$, $(\kappa^+, \kappa) \rightarrow (\mu^+, \mu)$.*

In the paper [5], we showed, assuming the consistency of a huge cardinal, that there is a model of ZFC + GCH in which $(\kappa^+, \kappa) \rightarrow (\mu^+, \mu)$ holds whenever κ is regular and $\mu < \kappa$ is infinite. Surprisingly, the full Global Chang's Conjecture is inconsistent (even without assuming GCH), as we show in Theorem 9. Indeed, there is a tension between instances of Chang's Conjecture at successors of singular cardinals and at double successors of singulars.

Next, we investigate other forms of Global Chang's Conjecture:

Definition (Singular Global Chang's Conjecture). *We say that the Singular Global Chang's Conjecture holds if for all infinite $\mu < \kappa$ of the same cofinality, $(\kappa^+, \kappa) \rightarrow (\mu^+, \mu)$.*

Obtaining the Singular Global Chang's Conjecture seems to be hard. We present here a partial result, showing that there is a model in which the Singular Global Chang's Conjecture holds for cardinals below $\aleph_{\omega\omega}$.

The paper is organized as follows. In Section 2 we discuss some relationships between Chang's Conjecture and PCF-theoretic scales, and derive some ZFC limitations on the simultaneous occurrence of some instances of Chang's Conjecture. In

Section 3, we introduce the technology for obtaining $(\aleph_{\alpha+1}, \aleph_\alpha) \rightarrow (\aleph_{\beta+1}, \aleph_\beta)$ for various choices of α and β of countable cofinality. In Section 4 we construct a model in which $(\aleph_{\alpha+1}, \aleph_\alpha) \rightarrow (\aleph_{\beta+1}, \aleph_\beta)$ holds for all limit ordinals $0 \leq \beta < \alpha < \omega^\omega$. In Section 5, we show the consistency of $(\aleph_{\alpha+1}, \aleph_\alpha) \rightarrow (\aleph_{\beta+1}, \aleph_\beta)$ holding for a fixed β but for α ranging over a longer interval. We conclude with some open questions.

2. LIMITATIONS ON GLOBAL CHANG'S CONJECTURE

A useful strengthening of Chang's Conjecture is the following, introduced by Shelah [19]:

Definition. We say $(\kappa_1, \kappa_0) \rightarrow_\nu (\mu_1, \mu_0)$ if for all structures \mathfrak{A} on κ_1 in a countable language, there is a substructure \mathfrak{B} such that $|\mathfrak{B}| = \mu_1$, $|\mathfrak{B} \cap \kappa_0| = \mu_0$, and $\nu \subseteq \mathfrak{B}$.

Note that nothing more is asserted by adding the subscript ν when $\nu < \omega_1$. These versions of Chang's Conjecture are robust under mild forcing:

Lemma 3. Suppose $(\kappa_1, \kappa_0) \rightarrow_\nu (\mu_1, \mu_0)$ and \mathbb{P} is a ν^+ -c.c. partial order. Then $\Vdash_{\mathbb{P}} (\kappa_1, \kappa_0) \rightarrow_\nu (\mu_1, \mu_0)$.

Of particular interest is the case $\nu = \mu_0$. The following lemma gives a stepping-up of the Chang's Conjecture if the distance between the cardinals considered is not too great, or enough GCH holds relatively close to the upper end. A proof is contained in [7, Section 2.2.1].

Lemma 4. Suppose $(\kappa_1, \kappa_0) \rightarrow_\nu (\mu_1, \mu_0)$.

- (1) If $\kappa_0 = \mu_0^{+\nu}$, then $(\kappa_1, \kappa_0) \rightarrow_{\mu_0} (\mu_1, \mu_0)$.
- (2) If $\lambda \leq \mu_0$ and there is $\kappa \leq \kappa_0$ such that $\kappa_0 = \kappa^{+\nu}$ and $\kappa^\lambda \leq \kappa_0$, then $(\kappa_1, \kappa_0) \rightarrow_\lambda (\mu_1, \mu_0)$.

When the hypotheses of the above lemma cannot be applied, some GCH at the lower end allows a similar conclusion in a special case.

Lemma 5. Suppose $\mu^{<\nu} = \mu$, and $(\kappa^+, \kappa) \rightarrow (\mu^+, \mu)$. Then $(\kappa^+, \kappa) \rightarrow_\nu (\mu^+, \mu)$.

Proof. If $\kappa^\nu = \kappa$, then the conclusion follows from (2) of Lemma 4. Otherwise, let \mathfrak{A} be a structure on κ^+ which is isomorphic to a transitive elementary substructure of $(H_{\kappa^{++}}, \in, \triangleleft, \mu, \nu)$, where \triangleleft is a well-order of $H_{\kappa^{++}}$. It is easy to see that the conclusion of Proposition 1 applies to \mathfrak{A} with respect to the cardinals κ, ν, κ^+ . If $\mathfrak{B} \prec \mathfrak{A}$ witnesses Chang's Conjecture, then $|\mathfrak{B} \cap \kappa|^{|\mathfrak{B} \cap \nu|} = \mu^{|\mathfrak{B} \cap \nu|} \geq |\mathfrak{B} \cap \kappa^+| = \mu^+$. Thus $|\mathfrak{B} \cap \nu| = \nu$.

Let $\delta \in \mathfrak{B} \cap \nu$. Corollary 2 implies that $\kappa^\delta = \kappa$. Let $\langle f_\alpha : \alpha < \kappa \rangle \in \mathfrak{B}$ list all functions from δ to κ . Let $\mathfrak{B}' = \text{Hull}^{\mathfrak{A}}(\mathfrak{B} \cup \delta)$. If $\beta \in \kappa \cap \mathfrak{B}'$, then there is function $f \in {}^\delta \kappa \cap \mathfrak{B}$ and $\gamma < \delta$ such that $\beta = f(\gamma)$. Thus $\mathfrak{B}' \cap \kappa = \{f_\alpha(\gamma) : \alpha \in \mathfrak{B} \cap \kappa \text{ and } \gamma < \delta\}$, which has size μ . Now let $\mathfrak{C} = \text{Hull}^{\mathfrak{A}}(\mathfrak{B} \cup \nu)$. Since \mathfrak{B} is cofinal in ν , $\mathfrak{C} = \bigcup \{\text{Hull}^{\mathfrak{A}}(\mathfrak{B} \cup \delta) : \delta \in \mathfrak{B} \cap \nu\}$, so $|\mathfrak{C} \cap \kappa| = \mu$. \square

Versions of Chang's Conjecture involving singular cardinals have a strong influence on the combinatorics in their neighborhood, even without cardinal arithmetic assumptions. Recall that if κ is singular, a *scale for κ* is a collection of functions $\langle f_\alpha : \alpha < \kappa^+ \rangle$ contained in some product $\prod_{i < \text{cf}(\kappa)} \kappa_i$, where $\langle \kappa_i : i < \text{cf}(\kappa) \rangle$ is an increasing and cofinal sequence of regular cardinals below κ , such that the functions f_α are increasing and cofinal in the partial order of the product where we put $f < g$ when $|\{i : f(i) \geq g(i)\}| < \text{cf}(\kappa)$. It is easy to construct scales under the assumption $2^\kappa = \kappa^+$, but Shelah proved in ZFC that scales exist for all singular cardinals (see [1]).

A scale $\langle f_\alpha : \alpha < \kappa^+ \rangle$ is *good at α* when there is a sequence $\vec{g} = \langle g_i : i < \text{cf}(\alpha) \rangle$ and $j_\star < \text{cf}(\kappa)$, such that for all $j \geq j_\star$, $\langle g_i(j) \mid i < \text{cf}(\alpha) \rangle$ is increasing and \vec{g} and

$\langle f_\beta : \beta < \alpha \rangle$ are interleaved (i.e., cofinal in each other). A scale is *bad at* α when it is not good at α . A scale is *better at* α if there is a club $C \subseteq \alpha$ such that for all $\beta \in C$ there is $j < \text{cf}(\kappa)$ such that $f_\gamma(i) < f_\beta(i)$ for $i \geq j$ and $\gamma \in C \cap \beta$. Note that if $\text{cf}(\alpha) > \text{cf}(\kappa)$, then being better at α implies being good at α . A scale is simply called *good* (or *better*) if it is good (or better) at every α such that $\text{cf}(\alpha) > \text{cf}(\kappa)$. The key connection with Chang's Conjecture is the following (see [9] or [19]):

Lemma 6. *If $(\kappa^+, \kappa) \rightarrow_{\text{cf}(\kappa)} (\mu^+, \mu)$ and $\mu \geq \text{cf}(\kappa)$, then there is no good scale for κ . Moreover, every scale $\langle f_\alpha : \alpha < \kappa^+ \rangle$ for κ is bad at stationarily many α of cofinality μ^+ .*

We now show that the full Global Chang's Conjecture is inconsistent with ZFC.

Lemma 7. *Suppose κ is regular, $\mu < \kappa$ is singular, and $(\kappa^+, \kappa) \rightarrow (\mu^+, \mu)$. Then μ carries a better scale. Moreover, if $\text{cf} \kappa = \omega$ then \square_κ^* holds.*

Proof. Let us start with a general observation, following [8, Theorem 2.15].

Claim 8. *Let $\mu < \kappa = \text{cf}(\kappa)$ be cardinals. Let θ be a regular cardinal above κ^+ . If H is the transitive collapse of some elementary substructure of H_θ of size κ^+ containing κ^+ , and $M \prec H$ is such that $|M \cap \kappa^+| = \mu^+$ and $|M \cap \kappa| = \mu$, then $\text{cf}(\sup(M \cap \kappa)) = \text{cf}(\mu)$.*

Proof. Fix in such an H a sequence $\langle x_\alpha : \alpha < \kappa^+ \rangle$ of “strongly almost disjoint” unbounded subsets of κ . That is, for every $\alpha < \kappa^+$, there is a sequence $\langle \gamma_\beta^\alpha : \beta < \alpha \rangle \in H$ of ordinals below κ such that $\langle x_\beta \setminus \gamma_\beta^\alpha : \beta < \alpha \rangle$ is pairwise disjoint. This principle, due to Shelah, is called ADS_κ and it holds for κ regular (see [3] and [20]).

Let $M \prec H$ be as above. Let $f : \mu \rightarrow M \cap \kappa$ be a bijection. If $\text{cf}(\sup(M \cap \kappa)) \neq \text{cf}(\mu)$, then for each $\alpha < M \cap \kappa^+$ there is $\delta_\alpha < \mu$ such that $f[\delta_\alpha] \cap x_\alpha$ is cofinal in $M \cap \kappa$. Since $|M \cap \kappa^+| = \mu^+$, there is a set $Y \subseteq M \cap \kappa^+$ of size μ^+ and a fixed $\delta < \mu$ such that $\delta_\alpha = \delta$ for all $\alpha \in Y$. Let $\zeta \in M \cap \kappa^+$ be large enough so that $|Y \cap \zeta| = \mu$. Note that $\langle \gamma_\beta^\zeta : \beta < \zeta \rangle \in M$ and thus for every $\beta \in M \cap \zeta$, $\gamma_\beta^\zeta \in M \cap \kappa$.

For $\beta \in Y \cap \zeta$, let $y_\beta = f[\delta] \cap x_\beta \setminus \gamma_\beta^\zeta$. Then $\{f^{-1}[y_\beta] : \beta \in Y \cap \zeta\}$ is a collection of μ many pairwise disjoint subsets of δ , which is impossible. \square

Let us return to the proof of the lemma.

By a theorem of Shelah [19], κ carries a “partial weak square”, a weak square sequence that misses only cofinality κ . That is, there is a sequence $\langle \mathcal{C}_\alpha : \alpha < \kappa^+ \rangle$ such that whenever $\omega \leq \text{cf}(\alpha) < \kappa$, then \mathcal{C}_α is a nonempty collection of size $\leq \kappa$ such that each $C \in \mathcal{C}_\alpha$ is a club subset of α of size $< \kappa$, and if $C \in \mathcal{C}_\alpha$ and $\beta \in \lim C$, then $C \cap \beta \in \mathcal{C}_\beta$.

Let $M \prec H$ be as above, with $\vec{C} \in M$ a partial weak square at κ . Let $\pi : M \rightarrow N$ be the transitive collapse. Let $\vec{D} = \pi(\vec{C})$. Since $\text{ot}(M \cap \kappa^+) = \mu^+$ and $|M \cap \mathcal{C}_\alpha| \leq \mu$ for each $\alpha \in M \cap \kappa^+$, \vec{D} is a sequence $\langle \mathcal{D}_\alpha : \alpha < \mu^+ \rangle$, such that each \mathcal{D}_α has size $\leq \mu$, if $D \in \mathcal{D}_\alpha$ and $\beta \in \lim D$, then $D \cap \beta \in \mathcal{D}_\beta$, and \mathcal{D}_α is nonempty whenever α is a limit ordinal such that $\text{cf}(\pi^{-1}(\alpha)) < \kappa$. If α is such that $\text{cf}(\pi^{-1}(\alpha)) = \kappa$, then there is an increasing cofinal map $f : \kappa \rightarrow \pi^{-1}(\alpha)$ in M , which implies that $\text{cf}(\alpha) = \text{cf}(\mu)$. Therefore, \mathcal{D}_α is nonempty whenever $\text{cf}(\alpha) > \text{cf}(\mu)$. Furthermore, if $D \in \mathcal{D}_\alpha$, then $\text{ot}(D) < \pi(\kappa)$.

Next, we modify \vec{D} to a sequence $\vec{\mathcal{E}}$ with the same properties except that $|C| < \mu$ whenever $C \in \mathcal{E}_\alpha$ and $\alpha < \mu^+$. It is easy to show by induction that for each $\eta < \mu^+$, there is a “short square” of length η —a coherent sequence of clubs $\langle E_\alpha : \alpha < \eta \rangle$ such that $|E_\alpha| < \mu$ for each $\alpha < \eta$. Fix such a sequence $\langle E_\alpha : \alpha < \pi(\kappa) \rangle$. For each $\alpha < \mu^+$, let $\mathcal{E}_\alpha = \{\{\beta \in D : \text{ot}(D \cap \beta) \in E_{\text{ot}(D)}\} : D \in \mathcal{D}_\alpha\}$. Clearly each element of each \mathcal{E}_α has size $< \mu$. If $C \in \mathcal{E}_\alpha$ and $\beta \in \lim C$, then there is $D \in \mathcal{D}_\alpha$ such

that $\beta \in \lim D$ and $C = \{\beta \in D : \text{ot}(D \cap \beta) \in E_{\text{ot}(D)}\}$, and thus $D \cap \beta \in \mathcal{D}_\beta$ and $\text{ot}(D \cap \beta) \in \lim E_{\text{ot}(D)}$, so $C \cap \beta \in \mathcal{E}_\beta$.

Note that $\vec{\mathcal{E}}$ is a partial weak square, avoiding only ordinals of cofinality $\leq \omega$. Thus if $\text{cf } \kappa = \omega$, one can easily obtain a weak square sequence by completing the missing points in $\vec{\mathcal{E}}$.

Fix a scale for μ , $\langle f_\alpha : \alpha < \mu^+ \rangle \subseteq \prod_{i < \text{cf}(\mu)} \mu_i$. Let us inductively construct a better scale $\langle g_\alpha : \alpha < \mu^+ \rangle$ as follows. Let $g_0 = f_0$. If \mathcal{E}_α is empty, let $g_\alpha = f_\gamma$, where $\gamma \geq \alpha$ and f_γ eventually dominates g_β for each $\beta < \alpha$. If \mathcal{E}_α is nonempty, first, for all $C \in \mathcal{E}_\alpha$, define

$$g_C(i) = \begin{cases} \sup\{g_\beta(i) + 1 : \beta \in C\} & \text{if } \mu_i > |C|, \\ 0 & \text{otherwise.} \end{cases}$$

Then let $g_\alpha = f_\gamma$, where $\gamma \geq \alpha$ and f_γ eventually dominates g_β for each $\beta < \alpha$ and g_C for each $C \in \mathcal{E}_\alpha$.

Clearly $\langle g_\alpha : \alpha < \mu^+ \rangle$ is a scale. To check betterness, if $\text{cf}(\alpha) > \text{cf}(\mu)$, let $C \in \mathcal{E}_\alpha$. If $\beta \in \lim C$, then $C \cap \beta \in \mathcal{E}_\beta$. There is $i < \text{cf}(\mu)$ such that $g_{C \cap \beta}(j) > g_\gamma(j)$ for $i < j < \text{cf}(\mu)$ and $\gamma \in C \cap \beta$. Thus if C' is the set of limit points of some $C \in \mathcal{E}_\alpha$, then for all $\beta \in C'$ there is $i < \text{cf}(\mu)$ such that $g_\beta(j) > g_\gamma(j)$ for $i < j < \text{cf}(\mu)$ and $\gamma \in C' \cap \beta$. \square

Combining the above with Lemmas 5 and 6, we have:

Theorem 9. *Suppose κ is singular, $\lambda > \kappa$ is regular, $(\lambda^+, \lambda^+) \rightarrow (\kappa^+, \kappa)$, and $\text{cf}(\kappa) \leq \mu < \kappa$. Then $(\kappa^+, \kappa) \not\rightarrow_{\text{cf}(\kappa)} (\mu^+, \mu)$. Thus if $\mu^{<\text{cf}(\kappa)} = \mu$, then $(\kappa^+, \kappa) \not\rightarrow (\mu^+, \mu)$.*

Corollary 10. *$[\aleph_0, \aleph_\omega]$ is the maximal initial interval of cardinals on which the Global Chang's Conjecture can hold.*

The negative direction follows from Theorem 9 and the positive direction is proven in [5, Section 5].

It seems to be unknown whether $(\kappa^+, \kappa) \rightarrow (\mu^+, \mu)$ is equivalent to $(\kappa^+, \kappa) \rightarrow_\mu (\mu^+, \mu)$ for regular μ . However, further analysis of scales allows us to rule out some instances of Chang's Conjecture in ZFC, and to show that these two notions are not in general equivalent for singular μ , even under GCH. The authors are grateful to Chris Lambie-Hanson for showing us how to prove the following:

Theorem 11. *Suppose κ is a singular cardinal and $\vec{f} = \langle f_\alpha : \alpha < \kappa^+ \rangle$ is a scale for κ . There is a club $C \subseteq \kappa^+$ such that for all regular cardinals μ, ν such that $\text{cf}(\kappa) < \mu < \mu^{+3} \leq \nu < \mu^{+\text{cf}(\kappa)} \leq \kappa$, \vec{f} is good at every $\alpha \in C$ of cofinality ν .*

Proof. Suppose $\text{cf}(\kappa) < \mu < \mu^{+3} \leq \nu < \mu^{+\text{cf}(\kappa)} \leq \kappa$. By [1, Theorem 2.21], there is a club $C_{\mu, \nu} \subseteq \kappa^+$ such that for every $\alpha \in C_{\mu, \nu}$ of cofinality ν , $\langle f_\beta : \beta < \alpha \rangle$ has an exact upper bound g such that $\text{cf}(g(i)) \geq \mu$ for all i . g being an exact upper bound means that g is an upper bound to $\langle f_\beta : \beta < \alpha \rangle$, and for every $h < g$, there is $\beta < \alpha$ such that $h < f_\beta$.

The arguments for Lemmas 6–8 of [15] show that $\text{cf}(g(i)) = \nu$ on a cobounded set of $i < \text{cf}(\kappa)$, which implies \vec{f} is good at α . For the reader's convenience: Let $\langle \alpha_j : j < \nu \rangle$ be cofinal in α . We cannot have that $\text{cf}(g(i)) > \nu$ for all i in an unbounded set $X \subseteq \text{cf}(\kappa)$. For then there would be an $i^* < \text{cf}(\kappa)$, an unbounded $Y \subseteq \nu$, and an $h < g$ such that $f_{\alpha_j}(i) < h(i) < g(i)$ for $i \in X \setminus i^*$ and $j \in Y$, contradicting that g is an exact upper bound. Thus there is some $\nu' \in [\mu, \nu]$ and an unbounded $X \subseteq \text{cf}(\kappa)$ such that $\text{cf}(g(i)) = \nu'$ for all $i \in X$. Let $\langle g_k : k < \nu' \rangle$ be a pointwise increasing sequence such that $\sup_{k < \nu'} g_k(i) = g(i)$ for all $i \in X$. Since g is an exact upper bound, for each $k < \nu'$, there is $j < \nu$ such that $g_k \upharpoonright X < f_{\alpha_j} \upharpoonright X$.

Also, for each $j < \nu$, there is $i^* < \text{cf}(\kappa)$ such that $f_{\alpha_j}(i) < g(i)$ for $i \in X \setminus i^*$, and thus some $k < \nu'$ such that $f_{\alpha_j} \upharpoonright X < g_k \upharpoonright X$. This implies $\nu' = \nu$.

Finally, we can take the intersection of all the $C_{\nu, \mu}$ for regular $\nu, \mu < \kappa^+$ to get the desired club C . \square

Therefore, if κ is singular, $(\kappa^+, \kappa) \twoheadrightarrow_{\text{cf}(\kappa)} (\mu^+, \mu)$ fails whenever $\text{cf}(\kappa)^{+3} \leq \mu < \text{cf}(\kappa)^{+\text{cf}(\kappa)}$. However, it is possible that the version of Chang's Conjecture holds when we drop the subscript " $\text{cf}(\kappa)$ " on the arrow:

Proposition 12. *Suppose there is a 3-huge cardinal. Then there are singular cardinals $\lambda < \delta$ such that $\text{cf}(\delta) < \lambda < \text{cf}(\delta)^{+\text{cf}(\delta)}$ and $(\delta^+, \delta) \twoheadrightarrow (\lambda^+, \lambda)$.*

Proof. Let $j : V \rightarrow M$ have critical point κ , with $M^{j^3(\kappa)} \subseteq M$. Let $\delta = j^2(\kappa)^{+j(\kappa)}$ and let $\lambda = j(\kappa)^{+\kappa}$. Let \mathfrak{A} be any structure on δ^+ . In M , $j[\mathfrak{A}] \prec j(\mathfrak{A})$, and we have that $|j[\mathfrak{A}]| = \delta^+$ and $|j[\mathfrak{A}] \cap j(\delta)| = \delta$. Reflecting through j , we have that there is $\mathfrak{B} \prec \mathfrak{A}$ such that $|\mathfrak{B}| = j(\kappa)^{+\kappa+1} = \lambda^+$ and $|\mathfrak{B} \cap \delta| = j(\kappa)^{+\kappa} = \lambda$. \square

3. CHANG'S CONJECTURE BETWEEN SUCCESSORS OF VARIOUS SINGULARS

Recall that a partial order is (κ, λ) -distributive if forcing with it adds no functions from κ to λ . The following lemma is a mild generalization of a lemma that was proved in [5].

Lemma 13. *Let $\gamma < \kappa$ be such that $\kappa^{+\gamma}$ is a strong limit cardinal and κ is $\kappa^{+\gamma+1}$ -supercompact, as witnessed by an embedding $j : V \rightarrow M$. If \mathcal{U} is the ultrafilter on κ derived from j , then there is $A \in \mathcal{U}$ such that for every $\alpha < \beta$ in $A \cup \{\kappa\}$ and every iteration $\mathbb{P} * \dot{\mathbb{Q}}$ of size $< \beta^{+\gamma}$, such that \mathbb{P} is $\alpha^{+\gamma+1}$ -Knaster and $\Vdash_{\mathbb{P}} \dot{\mathbb{Q}}$ is $(\alpha^{+\gamma+1}, \alpha^{+\gamma+1})$ -distributive,*

$$\Vdash_{\mathbb{P} * \dot{\mathbb{Q}}} (\beta^{+\gamma+1}, \beta^{+\gamma}) \twoheadrightarrow_{\alpha^{+\gamma}} (\alpha^{+\gamma+1}, \alpha^{+\gamma}).$$

Proof. We show that for a set $A \in \mathcal{U}$, for every $\alpha \in A$ and every iteration $\mathbb{P} * \dot{\mathbb{Q}}$ satisfying the hypothesis for $\beta = \kappa$ forces $(\kappa^{+\gamma+1}, \kappa^{+\gamma}) \twoheadrightarrow_{\alpha^{+\gamma}} (\alpha^{+\gamma+1}, \alpha^{+\gamma})$. Then standard reflection arguments yield the desired conclusion. By Lemma 4, it suffices to prove that for all $\alpha \in A$, every such $\mathbb{P} * \dot{\mathbb{Q}}$ forces $(\kappa^{+\gamma+1}, \kappa^{+\gamma}) \twoheadrightarrow_{\gamma} (\alpha^{+\gamma+1}, \alpha^{+\gamma})$, since by the assumptions that $\kappa^{+\gamma}$ is a strong limit and $|\mathbb{P} * \dot{\mathbb{Q}}| < \kappa^{+\gamma}$, it is forced that for some $\lambda \in [\kappa, \kappa^{+\gamma})$, $\lambda^\kappa < \kappa^{+\gamma}$, so we may increase the subscript to $\alpha^{+\gamma}$. If the claim fails, then on a set $B \in \mathcal{U}$, for every $\alpha \in B$, there is an iteration $\mathbb{P}_\alpha * \dot{\mathbb{Q}}_\alpha$ and a name for a function $\dot{f}_\alpha : (\kappa^{+\gamma+1})^{<\omega} \rightarrow \kappa^{+\gamma}$ such that it is forced that for every $X \subseteq \kappa^{+\gamma+1}$ of size $\alpha^{+\gamma+1}$ with $\gamma \subseteq X$, the closure of X under \dot{f}_α contains $\alpha^{+\gamma+1}$ many ordinals below $\kappa^{+\gamma+1}$. We may assume that \dot{f}_α is closed under compositions.

In M , let $\mathbb{P} * \dot{\mathbb{Q}} = j(\langle \mathbb{P}_\alpha * \dot{\mathbb{Q}}_\alpha : \alpha < \kappa \rangle)(\kappa)$ and let $\dot{f} = j(\langle \dot{f}_\alpha : \alpha < \kappa \rangle)(\kappa)$. Let $X = j[\kappa^{+\gamma+1}]$. Note that X is a subset of $j(\kappa^{+\gamma+1})$ containing γ and of size $\kappa^{+\gamma+1}$. By hypothesis, $\Vdash_{\mathbb{P} * \dot{\mathbb{Q}}}^M |\dot{f}[X^{<\omega}]| = \kappa^{+\gamma+1}$. Since $j(\kappa^{+\gamma})$ is singular, it is forced that there is a sequence $\langle \dot{b}_\alpha : \alpha < \kappa^{+\gamma+1} \rangle \subseteq X$ such that $\langle \dot{f}(\dot{b}_\alpha) : \alpha < \kappa^{+\gamma+1} \rangle$ is a strictly increasing sequence of ordinals below $j(\kappa^{+\xi})$, for some $\xi < \gamma$. Let $\nu < \gamma$ and $(p_0, \dot{q}_0) \in \mathbb{P} * \dot{\mathbb{Q}}$ be such that $|\mathbb{P} * \dot{\mathbb{Q}}| < j(\kappa^{+\nu})$ and $(p_0, \dot{q}_0) \Vdash \dot{f}(\dot{b}_\alpha) < j(\kappa^{+\nu})$ for all $\alpha < \kappa^{+\gamma+1}$.

Since $\dot{\mathbb{Q}}$ adds no subsets to X , there is a \mathbb{P} -name \dot{Y} and a condition $(p_1, \dot{q}_1) \leq (p_0, \dot{q}_0)$ such that $(p_1, \dot{q}_1) \Vdash \langle \dot{b}_\alpha : \alpha < \kappa^{+\gamma+1} \rangle = \dot{Y}$. Next, for each $\alpha < \kappa^{+\gamma+1}$, find $r_\alpha \leq p_1$ and $a_\alpha \in (\kappa^{+\gamma+1})^{<\omega}$ such that $r_\alpha \Vdash_{\mathbb{P}} j(\check{a}_\alpha) = \dot{Y}(\alpha)$. Since \mathbb{P} is $\kappa^{+\gamma+1}$ -Knaster, there is $Z \subseteq \kappa^{+\gamma+1}$ of size $\kappa^{+\gamma+1}$ such that r_α and r_β are compatible for $\alpha, \beta \in Z$. Therefore, for $\alpha < \beta$ in Z , there is $r \in \mathbb{P}$ such that $(r, \dot{q}_1) \Vdash \dot{f}(j(\check{a}_\alpha)) < \dot{f}(j(\check{a}_\beta)) < j(\kappa^{+\nu})$.

Reflecting these statements to V , we have that for $\alpha < \beta$ in Z , there are $\gamma < \kappa$ and $(p, \dot{q}) \in \mathbb{P}_\gamma * \dot{\mathbb{Q}}_\gamma$ such that $|\mathbb{P}_\gamma * \dot{\mathbb{Q}}_\gamma| < \kappa^{+\nu}$ and $(p, \dot{q}) \Vdash_{\mathbb{P}_\gamma * \dot{\mathbb{Q}}_\gamma}^V \dot{f}_\gamma(\dot{a}_\alpha) < \dot{f}_\gamma(\dot{a}_\beta) < \kappa^{+\nu}$. This defines a coloring of $[\kappa^{+\gamma+1}]^2$ in $\kappa^{+\nu}$ colors. Since $\kappa^{+\gamma}$ is a strong limit, the Erdős-Rado Theorem implies that there is a set $H \subseteq Z$ of size $\kappa^{+\nu+1}$ such that all pairs in $[H]^2$ get the same color. Thus we have a fixed η and a fixed $(p, \dot{q}) \in \mathbb{P}_\eta * \dot{\mathbb{Q}}_\eta$ such that $(p, \dot{q}) \Vdash \dot{f}_\eta(a_\alpha) < \dot{f}_\eta(a_\beta) < \kappa^{+\nu}$ for $\alpha < \beta$ in H . This is a contradiction. \square

Corollary 14. *If there is a $(+\omega+1)$ -supercompact cardinal, then there is a forcing extension in which $(\aleph_{\alpha+1}, \aleph_\alpha) \rightarrow (\aleph_{\beta+1}, \aleph_\beta)$ holds for all limit ordinals $0 \leq \beta < \alpha < \omega^2$.*

Proof. Let κ be $\kappa^{+\omega+1}$ -supercompact, and let $A \subseteq \kappa$ be given by Lemma 13. Let $\langle \alpha_i : i < \omega \rangle$ enumerate the first ω elements of A . Let

$$\mathbb{P} = \text{Col}(\omega, \alpha_0^{+\omega}) \times \prod_{n < \omega} \text{Col}(\alpha_n^{+\omega+2}, \alpha_{n+1}).$$

Clearly, \mathbb{P} forces that $\alpha_n^{+\omega} = \aleph_{\omega \cdot n}$ for all n . For a fixed n , we can factor \mathbb{P} as $\mathbb{Q}_0 \times \text{Col}(\alpha_n^{+\omega+2}, \alpha_{n+1}) \times \mathbb{Q}_1$. The middle term forces $(\alpha_{n+1}^{+\omega+1}, \alpha_{n+1}^{+\omega}) \rightarrow_{\alpha_n^{+\omega}} (\alpha_n^{+\omega+1}, \alpha_n^{+\omega})$. Since \mathbb{Q}_1 is $\alpha_{n+1}^{+\omega+2}$ -distributive after forcing with the middle term, the instance of Chang's Conjecture is preserved. Then, forcing with \mathbb{Q}_0 preserves this instance by Lemma 3. Since Chang's Conjecture is transitive, i.e. $(\kappa_1, \kappa_0) \rightarrow (\mu_1, \mu_0)$ and $(\mu_1, \mu_0) \rightarrow (\nu_1, \nu_0)$ implies $(\kappa_1, \kappa_0) \rightarrow (\nu_1, \nu_0)$, the conclusion follows. \square

The limitation of our argument so far is that we only get Chang's Conjecture between successors of singulars for which there are tail-end sequences of cardinals below that are order-isomorphic. We will overcome this with a forcing that collapses singular cardinals to onto others of different types while preserving their successors and the desired instances of Chang's Conjecture.

Theorem 15. *Assume GCH. Suppose $\alpha < \beta$ are countable limit ordinals and κ is $\kappa^{+\beta+1}$ -supercompact. Then there is a forcing extension in which $(\aleph_{\beta+1}, \aleph_\beta) \rightarrow (\aleph_{\alpha+1}, \aleph_\alpha)$.*

The proof breaks into cases depending on the “tail types” of α and β . For ordinals $\alpha \geq \beta$, let $\alpha - \beta$ be the unique γ such that $\alpha = \beta + \gamma$. For an ordinal α , let $\tau(\alpha)$ (the tail of α) be $\min_{\beta < \alpha} (\alpha - \beta)$. Let $\iota(\alpha)$ be the least β such that $\alpha = \beta + \tau(\alpha)$. An ordinal α is indecomposable iff $\alpha = \tau(\alpha)$, and all tails are indecomposable.

Case 1: $\tau(\alpha) = \tau(\beta) = \gamma$, or $\alpha = 0$. Note that $\iota(\beta) \geq \alpha$, and let $\delta = \iota(\beta) - \alpha$. Let $A \subseteq \kappa$ be given by Lemma 13 (with respect to γ). Let $\zeta < \eta$ be in A , and force with $\text{Col}(\zeta^{+\gamma+\delta+2}, \eta)$, so that the ordertype of the set of cardinals between $\zeta^{+\gamma}$ and $\eta^{+\gamma}$ becomes $\delta + \gamma$. By Lemma 13, we have $(\eta^{+\gamma+1}, \eta^{+\gamma}) \rightarrow_{\zeta^{+\gamma}} (\zeta^{+\gamma+1}, \zeta^{+\gamma})$. If $\alpha = 0$, force with $\text{Col}(\omega, \zeta^{+\gamma})$, and if $\alpha > 0$, force with $\text{Col}(\aleph_{\iota(\alpha)+1}, \zeta)$. In both cases, Chang's Conjecture is preserved, and we get $|\zeta^{+\gamma}| = \aleph_\alpha$ and $\eta^{+\gamma} = \aleph_{\alpha+\delta+\gamma} = \aleph_\beta$.

For the other cases, we will use a variation on the Gitik-Sharon forcing [11], which singularizes a large cardinal while collapsing a singular cardinal above it.

The following definition is standard (see [10]).

Definition. *A structure $\langle \mathbb{P}, \leq, \leq^* \rangle$ is a Prikry-type forcing when \leq and \leq^* are partial orders of \mathbb{P} (called extension and direct extension respectively), with $p \leq^* q \Rightarrow p \leq q$, and such that whenever σ is a statement in the forcing language of $\langle \mathbb{P}, \leq \rangle$ and $p \in \mathbb{P}$, then there is $q \leq^* p$ deciding σ . Such a forcing is called weakly κ -closed for a cardinal κ if $\langle \mathbb{P}, \leq^* \rangle$ is κ -closed.*

It is easy to see that if \mathbb{P} is of Prikry type and weakly κ^+ -closed, then it is (κ, κ) -distributive.

Suppose $\gamma < \delta$ are limit ordinals of countable cofinality, and $\vec{\gamma} = \langle \gamma_i : 1 \leq i < \omega \rangle$, $\vec{\delta} = \langle \delta_i : 1 \leq i < \omega \rangle$ are sequences such that:

- (1) $\vec{\gamma}$ is strictly increasing with $\sup_i \gamma_i = \gamma$.
- (2) $\vec{\delta}$ is nondecreasing with $\gamma \leq \delta_1$ and $\sum_i \delta_i = \delta$.

Suppose $\kappa > \delta$ is $\kappa^{+\gamma_n}$ -supercompact for each $n \geq 1$, and $\mu < \kappa$ is regular. For $1 \leq n < \omega$, let U_n be a κ -complete normal measure on $\mathcal{P}_\kappa(\kappa^{+\gamma_n})$, and let $j_n : V \rightarrow M_n \cong \text{Ult}(V, U_n)$ be the ultrapower embedding. By the closure of the ultrapowers and GCH, we may choose an M_n -generic $K_n \subseteq \text{Col}(\kappa^{+\delta_n+2}, j_n(\kappa))^{M_n}$. Let $\vec{U} = \langle U_n : n < \omega \rangle$ and $\vec{K} = \langle K_n : n < \omega \rangle$.

With these choices made, we may define the forcing $\mathbb{P}(\mu, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K})$, which will have the following properties:

- The forcing is of Prikry type, weakly μ -closed, and $\kappa^{+\gamma}$ -centered (and thus has the $\kappa^{+\gamma+1}$ -c.c.).
- κ is forced to become $\mu^{+\delta}$.
- $(\kappa^{+\gamma})^V$ is collapsed to κ .

Conditions in $\mathbb{P}(\mu, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K})$ are sequences

$$\langle f_0, x_1, f_1, \dots, x_n, f_n, F_{n+1}, F_{n+2}, \dots \rangle,$$

where:

- (1) For $1 \leq i \leq n$, $x_i \in \mathcal{P}_\kappa(\kappa^{+\gamma_i})$, and $\kappa_i := x_i \cap \kappa$ is inaccessible.
- (2) For $1 \leq i < n$, $\kappa_{i+1} > |x_i|$.
- (3) $f_0 \in \text{Col}(\mu, \kappa)$, and $\text{ran}(f_0) \subseteq \kappa_1$ if x_1 is defined.
- (4) For $1 \leq i < n$, $f_i \in \text{Col}(\kappa_i^{+\delta_i+2}, \kappa_{i+1})$.
- (5) $f_n \in \text{Col}(\kappa_n^{+\delta_n+2}, \kappa)$.
- (6) For $i > n$, $\text{dom } F_i \in U_i$.
- (7) For $i > n$ and $x \in \text{dom } F_i$, $\kappa_x := x \cap \kappa$ is an inaccessible cardinal greater than $|x_n| + \sup(\text{ran } f_n)$.
- (8) For $i > n$ and $x \in \text{dom } F_i$, $F_i(x) \in \text{Col}(\kappa_x^{+\delta_i+2}, \kappa)$.
- (9) For $i > n$, $[F_i]_{U_i} \in K_i$.

Suppose $p = \langle f_0, \dots, x_n, f_n, F_{n+1}, \dots \rangle$ and $q = \langle f'_0, \dots, x'_m, f'_m, F'_{m+1}, \dots \rangle$. We say $q \leq p$ when:

- (1) $m \geq n$.
- (2) $f'_i \supseteq f_i$ for $i \leq n$, and $x_i = x'_i$ for $1 \leq i \leq n$.
- (3) For $n < i \leq m$, $x'_i \in \text{dom } F_i$ and $f'_i \supseteq F_i(x'_i)$.
- (4) For $i > m$, $\text{dom } F'_i \subseteq \text{dom } F_i$, and $F'_i(x) \supseteq F_i(x)$ for $x \in \text{dom } F'_i$.

For p as above, let $\text{stem}(p) = \langle f_0, \dots, x_n, f_n \rangle$, and say the *length* of p is n . (The stem of length-0 conditions is of the form $\langle f_0 \rangle$.)

Lemma 16. *Suppose $\mu, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K}$ are as above, and $p = \langle f_0, x_1, \dots, x_n, f_n \rangle \frown \vec{F}$ is a condition of length $n > 0$. Then $\mathbb{P}(\mu, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K}) \restriction p$ is canonically isomorphic to*

$$\text{Col}(\mu, \kappa_1) \restriction f_0 \times \dots \times \text{Col}(\kappa_{n-1}^{+\delta_{n-1}+2}, \kappa_n) \restriction f_{n-1} \times \mathbb{P}(\kappa_n^{+\delta_n+2}, \vec{\gamma}', \vec{\delta}', \vec{U}', \vec{K}') \restriction \langle f_n \rangle \frown \vec{F}',$$

where for each sequence $s \in \{\vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K}, \vec{F}\}$, s' is the sequence such that $s'(m) = s(n+m)$ for $m \geq 1$.

We say $q \leq^* p$ when $q \leq p$ and they have the same length. If $q \leq p$ and $\text{stem}(p)$ is an initial segment of $\text{stem}(q)$, we say q is an *end-extension* of p , or $q \leq p$. Given a sequence $\vec{F} = \langle F_i : 1 \leq i < \omega \rangle$ such that $\langle \emptyset \rangle \frown \vec{F}$ is a condition of length 0, and

another condition $p = \text{stem}(p) \smallfrown \langle H_i : n < i < \omega \rangle$, define

$$p \wedge \vec{F} := \text{stem}(p) \smallfrown \langle \{ \langle x, F_i(x) \cup H_i(x) \rangle : x \in \text{dom } F_i \cap \text{dom } H_i \\ \text{and } F_i(x) \cup H_i(x) \text{ is a function} \} : n < i < \omega \rangle.$$

Note that $p \wedge \vec{F}$ is both \preceq and \leq^* p , but $p \wedge \vec{F}$ is not necessarily $\leq \langle \emptyset \rangle \smallfrown \vec{F}$. For a given stem s and sequence \vec{F} as above, we define $s \wedge \vec{F} = p \wedge \vec{F}$, where p is the weakest condition with stem s .

It is easy to see that $\mathbb{P}(\mu, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K})$ is $\kappa^{+\gamma}$ -centered, and a density argument shows that it forces all cardinals in $[\kappa, \kappa^{+\gamma}]$ to have countable cofinality. The fact that not more damage is done than intended is a consequence of the Prikry Property, which follows from a more basic combinatorial property. If \mathbb{P} is a partial order and $c : \mathbb{P} \rightarrow \{0, 1, 2\}$, we say c is a *decisive coloring* if whenever $c(p) > 0$ and $q \leq p$, then $c(q) = c(p)$.

Lemma 17. *Let c be a decisive coloring of $\mathbb{P}(\mu, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K})$.*

- (1) *There is a sequence \vec{F} such that for every condition p , every two $r, r' \preceq p \wedge \vec{F}$ of the same length have the same color.*
- (2) *For every condition p , there is $q \leq^* p$ such that every two $r, r' \leq q$ of the same length have the same color.*

Proof. Let $\mathbb{P} = \mathbb{P}(\mu, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K})$. For (1), we prove the following claim by induction: For all $n < \omega$ and all decisive colorings of the conditions of length n , there is \vec{F} such that for all $m \leq n$ and every condition p of length m , every two $r, r' \preceq p \wedge \vec{F}$ of length n have the same color. Suppose $n = 0$ and c is such a coloring. For every $s \in \text{Col}(\mu, \kappa)$, choose if possible some \vec{F}_s such that $c(\langle s \rangle \smallfrown \vec{F}_s) > 0$. Using the closure of the higher collapses and diagonal intersections, we may select a single sequence \vec{F} such that $\langle s \rangle \wedge \vec{F} \leq \langle s \rangle \smallfrown \vec{F}_s$ for all s . By decisiveness, \vec{F} witnesses the claim for $n = 0$.

Suppose the claim is true for $n - 1$. Let c be any decisive coloring of the conditions of length n . Using the closure of $\text{Col}(\kappa^{+\delta_n+2}, j_{U_n}(\kappa))^{M_n}$, the genericity of K_n , and the decisiveness of $j_{U_n}(c)$, we can find a function $f^* \in K_n$ such that for every stem s of length $n - 1$, if there are some g and F such that $g \supseteq f^*$ and $s \smallfrown \langle j_{U_n}[\kappa^{+\gamma_n}], g \rangle \smallfrown \vec{F}$ has color > 0 , then $s \smallfrown \langle j_{U_n}[\kappa^{+\gamma_n}], f^* \rangle \smallfrown \vec{F}$ already has this color. If F_n represents f^* , then for all stems s of length $n - 1$, there is $A_s \in U_n$ and a color $c_s < 3$ such that for all $x \in A_s$, either there is $\vec{F}^{s,x} = \langle F_k^{s,x} : n + 1 \leq k < \omega \rangle$ such that $s \smallfrown \langle x, F_n(x) \rangle \smallfrown \vec{F}^{s,x}$ has color $c_s > 0$, or for all $x \in A_s$ and all $g \supseteq F_n(x)$, any p of length n with stem $s \smallfrown \langle x, g \rangle$ has color 0. Let A be the diagonal intersection of the sets A_s . Using the directed-closure of the filters K_k and diagonal intersections, we may select a single sequence \vec{F} that plays the role of $\vec{F}^{s,x}$ for all s and x . Putting $\vec{F}' = \langle F_n \restriction A \rangle \smallfrown \vec{F}$, we have that for any condition p of length $n - 1$, all $q \preceq p \wedge \vec{F}'$ of length n have the same color. This defines a decisive coloring c' of the conditions of length $n - 1$ of the form $p \wedge \vec{F}'$, by coloring them whatever color an arbitrary length- n extension receives. By induction, there is \vec{F}'' such that for every $m \leq n - 1$, for every condition p of length m , every $q \preceq p \wedge \vec{F}''$ of length $n - 1$ receives the same color under c' . This means that every such $p \wedge \vec{F}''$ receives the same color under c when end-extended to a condition of length n .

To finish the argument for (1), let c be a decisive coloring of \mathbb{P} . We have for each n a sequence \vec{F}_n such that the restriction of c to conditions of length n satisfies the inductive claim. Using the countable closure of the filters K_m , we can find the desired \vec{F} by taking a lower bound to all the conditions of the form $\langle \emptyset \rangle \smallfrown \vec{F}_n$.

For (2), let \vec{F} be given by (1) and let $p \in \mathbb{P}$. If there is $s \leq \text{stem}(p)$ such that some end-extension of $s \smallfrown \vec{F}$ has color > 0 , then pick such an s which achieves such

a color c^* by end-extending to length n , where n is as small as possible. Then every $r, r' \leq s \cap \vec{F}$ have color 0 if their length is $< n$, and color c^* otherwise. \square

Corollary 18. $\langle \mathbb{P}(\mu, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K}), \leq, \leq^* \rangle$ is a Prikry-type forcing.

Proof. If σ is a sentence in the forcing language of $\mathbb{P}(\mu, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K})$, then we color a condition 0 if it does not decide σ , 1 if it forces σ , and 2 if it forces $\neg\sigma$. This is decisive, so for every p , there is $q \leq^* p$ such that all extensions of q of the same length have the same color. If q does not decide σ , then there are $r, r' \leq q$ of the same length forcing opposite decisions about σ , contradicting the property of q . \square

Case 2 (of Theorem 15): $\tau(\alpha) > \tau(\beta) = \gamma$. Again, we have $\iota(\beta) \geq \alpha$, so let $\xi = \iota(\beta) - \alpha$. Let $A \subseteq \kappa$ be given by Lemma 13 (with respect to γ). Find $\nu < \mu$ in A such that ν is $\nu^{+\gamma+1}$ -supercompact. Let $G \subseteq \text{Col}(\nu^{+\gamma+\xi+2}, \mu)$ be generic over V . In $V[G]$, $(\mu^{+\gamma+1}, \mu^{+\gamma}) \twoheadrightarrow_{\nu^{+\gamma}} (\nu^{+\gamma+1}, \nu^{+\gamma})$ holds, and ν is still $\nu^{+\gamma+1}$ -supercompact. Let $\vec{\gamma} = \langle \gamma_i : 1 \leq i < \omega \rangle$ be an increasing sequence converging to γ . Since $\tau(\alpha) > \gamma$, we may find a nondecreasing sequence $\vec{\alpha} = \langle \alpha_i : 1 \leq i < \omega \rangle$ such that $\gamma \leq \alpha_1$ and $\sum_i \alpha_i = \alpha$.

Since ν is $\nu^{+\gamma+1}$ -supercompact, we can construct \vec{U} and \vec{K} as above according to the sequences $\vec{\gamma}, \vec{\alpha}$. Let $H \subseteq \mathbb{P}(\omega, \vec{\gamma}, \vec{\alpha}, \vec{U}, \vec{K})$ be generic over $V[G]$. Since this forcing is $\nu^{+\gamma+1}$ -c.c., Chang's Conjecture is preserved. In the extension, $\nu = \aleph_\alpha$, $(\nu^{+\gamma+1})^{V[G]} = (\nu^+)^{V[G][H]}$, and $\mu^{+\gamma} = \aleph_{\alpha+\xi+\gamma} = \aleph_\beta$.

The third case requires a more detailed analysis of the Gitik-Sharon forcing. Suppose $\mathbb{P}(\mu, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K})$ is built as above, around a sufficiently supercompact κ . Associated to a generic filter G are sequences $\langle x_n : 1 \leq n < \omega \rangle$, and $\langle C_n : n < \omega \rangle$ determined by the stems of all conditions in G , where C_0 is generic for $\text{Col}(\mu, \kappa_1)$, and for $n \geq 1$, C_n is generic for $\text{Col}(\kappa_n^{+\delta_n+2}, \kappa_{n+1})$ and $x_n \in \mathcal{P}_\kappa(\kappa^{+\gamma_n})$. From this sequence, we can recover G by taking all conditions $\langle f_0, x_1, f_1, \dots, x_n, f_n, F_{n+1}, \dots \rangle$ such that:

- (1) $\langle x_i : 1 \leq i \leq n \rangle$ is an initial segment of $\langle x_i : 1 \leq i < \omega \rangle$.
- (2) For $i \leq n$, $f_i \in C_i$.
- (3) For $i > n$, $x_i \in \text{dom } F_i$, and $F_i(x_i) \in C_i$.

The collection of such conditions is a filter containing G , so it must equal G by the maximality of generic filters.

Lemma 19. Let V be a model of set theory, and let $\langle \mathbb{P}_i, \kappa_i, G_i : i < n \rangle$ be such that:

- (1) $\langle \kappa_i : i < n \rangle$ is an increasing sequence of regular cardinals in V .
- (2) For each i , \mathbb{P}_i is a partial order in V that is (κ_i, κ_i) -distributive and of size $\leq \kappa_{i+1}$.
- (3) For each i , G_i is \mathbb{P}_i -generic over V .

Then $\prod_{i < n} G_i$ is $\prod_{i < n} \mathbb{P}_i$ -generic over V .

Proof. We show this by induction on $m \leq n$. Suppose that $\prod_{i < m} G_i$ is $\prod_{i < m} \mathbb{P}_i$ -generic over V . Since \mathbb{P}_m is (κ_m, κ_m) -distributive, forcing with it adds no antichains to $\prod_{i < m} \mathbb{P}_i$. Thus $\prod_{i < m} G_i$ is $\prod_{i < m} \mathbb{P}_i$ -generic over $V[G_m]$, and so $\prod_{i \leq m} G_i$ is $\prod_{i \leq m} \mathbb{P}_i$ -generic over V . \square

Lemma 20. (\vec{x}, \vec{C}) generates a generic for $\mathbb{P}(\mu, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K})$ over V iff the following hold:

- (1) For every sequence $\vec{F} = \langle F_n : 1 \leq n < \omega \rangle$ such that $\langle \emptyset \rangle \cap \vec{F}$ is a condition of length 0, there is m such that for all $n \geq m$, $x_n \in \text{dom } F_n$ and $F_n(x_n) \in C_n$.

- (2) C_0 is generic for $\text{Col}(\mu, \kappa_1)$, and C_n is generic for $\text{Col}(\kappa_n^{+\delta_n+2}, \kappa_{n+1})$ for all $n > 0$.

Proof. The forward direction is clear. For the reverse direction, let $D \in V$ be a dense open subset of $\mathbb{P} = \mathbb{P}(\mu, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K})$, and let G be the filter generated by (\vec{x}, \vec{C}) . Let $c : \mathbb{P} \rightarrow 2$ be defined by $c(p) = 0$ if $p \notin D$ and $c(p) = 1$ otherwise. This is decisive, so let \vec{F} be given by Lemma 17. Let m be given by (1).

Consider the condition $p = \langle \emptyset, x_1, \emptyset, \dots, x_{m-1}, \emptyset \rangle \cap \langle F_i : m \leq i < \omega \rangle$. Let $D' = \{q \in D : q \leq p\}$. D' projects to a dense subset of $\text{Col}(\mu, \kappa_1) \times \text{Col}(\kappa_1^{+\delta_1+2}, \kappa_2) \times \dots \times \text{Col}(\kappa_{m-1}^{+\delta_{m-1}+2}, \kappa_m)$. By (2) and Lemma 19, there is a sequence $\langle f_i : i < m \rangle$ that is in the projection of D' intersected with $C_0 \times \dots \times C_{m-1}$. Thus there is some condition of the form

$$\langle f_0, x_1, f_1, \dots, x_{m-1}, f_{m-1}, y_m, f_m, \dots, y_n, f_n, F'_{n+1}, \dots \rangle$$

that is in D' . But by the homogeneity property of \vec{F} , we also have that

$$\langle x_0, f_0, \dots, x_{m-1}, f_{m-1}, x_m, F_m(x_m), \dots, x_n, F_n(x_n), F_{n+1}, \dots \rangle \in D.$$

Therefore, $D \cap G \neq \emptyset$. \square

Case 3 (of Theorem 15): $0 < \tau(\alpha) = \gamma < \tau(\beta)$. Let $\delta = \beta - \iota(\alpha)$. We can find a nondecreasing sequence $\vec{\delta} = \langle \delta_i : 1 \leq i < \omega \rangle$ such that $\delta_1 \geq \gamma$ and $\sum_i \delta_i = \delta$. Let $\vec{\gamma} = \langle \gamma_i : 1 \leq i < \omega \rangle$ be an increasing sequence converging to γ . Let j be an embedding witnessing that κ is $\kappa^{+\gamma+1}$ -supercompact, and let $A \subseteq \kappa$ be given by Lemma 13 (with respect to γ). For each $n \geq 1$, let U_n be a κ -complete normal measure on $\mathcal{P}_\kappa(\kappa^{+\gamma_n})$ derived from j , so that A is in the projection of each U_n to κ . Let $\mu = \aleph_{\iota(\alpha)+1}$, and let us force with $\mathbb{P} = \mathbb{P}(\mu, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K})$ for where \vec{K} is a sequence of filters as in the construction.

Let p_0 be a condition of length 0 forcing every Prikry point to be in A . Let $p_1 \leq p_0$ be a condition of length 1 deciding the statement $\sigma := “(\kappa^+, \kappa) \twoheadrightarrow (\mu^{+\gamma+1}, \mu^{+\gamma})”$. We claim $p_1 \Vdash \sigma$.

Let us define an iteration of ultrapowers. Let $N_1 = V$. Given a commuting system of elementary embeddings $j_{m,m'} : N_m \rightarrow N_{m'}$ for $1 \leq m \leq m' \leq n$, let $j_{n,n+1} : N_n \rightarrow \text{Ult}(N_n, j_{0,n}(U_{n+1})) = N_{n+1}$ be the ultrapower embedding, and let $j_{m,n+1} = j_{n,n+1} \circ j_{m,n}$ for $1 \leq m < n$. For $n < \omega$, let $j_{n,\omega} : N_n \rightarrow N_\omega$ be the direct limit embedding. N_ω is well-founded, and thus can be identified with a transitive class, because of the following generalization of a well-known theorem of Gaifman (see [22]).

Fact 21. *If \mathcal{E} is a set of countably complete ultrafilters, and $j_{\alpha,\beta} : N_\alpha \rightarrow N_\beta$, $\alpha < \beta \leq \theta$, is a system of elementary embeddings defined by taking at each $\alpha < \theta$ the ultrapower map $j_{\alpha,\alpha+1} : N_\alpha \rightarrow \text{Ult}(N_\alpha, U) = N_{\alpha+1}$ for some $U \in j_{0,\alpha}(\mathcal{E})$, and taking direct limits at limit stages, then each N_α is well-founded.*

Let $\text{stem}(p_1) = \langle f_0, x_1, f_1 \rangle$, and let $C_0 \times C_1 \subseteq \text{Col}(\mu, \kappa_1) \times \text{Col}(\kappa_1^{+\delta_1+2}, \kappa)$ be a filter that contains $\langle f_0, f_1 \rangle$ and is generic over V . For $n > 1$, let $y_n = j_{n-1,n}[j_{1,n-1}(\kappa^{+\gamma_n})]$, and let $x_n = j_{n,\omega}(y_n)$, and let $C_n = j_{1,n-1}(K_n)$.

Claim 22. $\langle x_n : 1 \leq n < \omega \rangle$ and $\langle C_n : n < \omega \rangle$ together generate a generic filter for $j_{1,\omega}(\mathbb{P})$ over N_ω .

Proof. We need to verify the two conditions of Lemma 20. For (1), suppose $\vec{F} = \langle F_n : 1 \leq n < \omega \rangle$ is such that $\langle \emptyset \rangle \cap \vec{F} \in j_{1,\omega}(\mathbb{P})$ is a condition of length 0. Let $m < \omega$ be such that $\vec{F} = j_{m,\omega}(\vec{F}')$ for some \vec{F}' . For $n \geq m$, $\text{dom } j_{m,n}(F'_{n+1}) \in j_{1,n}(U_{n+1})$, and $N_n \models [j_{m,n}(F'_{n+1})]_{j_{1,n}(U_{n+1})} \in j_{1,n}(K_{n+1})$. Thus for $n \geq m$, $y_{n+1} \in \text{dom } j_{m,n+1}(F'_{n+1})$, and $f_{n+1} := j_{m,n+1}(F'_{n+1})(y_{n+1}) \in C_{n+1}$. Note that

f_{n+1} is an object of rank $< j_{1,n+1}(\kappa) = \text{crit}(j_{n+1,\omega})$. Thus for $n > m$, $x_n \in \text{dom } F_n$ and $f_n = F_n(x_n) \in C_n$.

To verify (2), note that for each $n > 1$, $N_{n-1} \models j_{1,n-1}(K_n)$ is generic for $\text{Col}(j_{1,n-1}(\kappa^{+\delta_n+2}), j_{1,n}(\kappa))$ over N_n . It is also generic over the submodel N_ω . Note also for each $n > 1$, $\kappa_n := x_n \cap j_{1,\omega}(\kappa) = j_{1,n-1}(\kappa)$. \square

Let G be the generated filter for $j_{1,\omega}(\mathbb{P})$. Note that $j_{1,\omega}(p_1) \in G$. We claim that $N_\omega[G]$ is closed under κ -sequences from $V[C_0 \times C_1]$. Since $C_0 \times C_1$ is generic for a forcing of size κ , it suffices to show that $N_\omega[\langle x_n : 2 \leq n < \omega \rangle]$ is closed under κ -sequences from V , an idea due to Bukovsky [2] and independently to Dehornoy [4]. This follows from the fact that every element of N_ω is of the form $j_{1,\omega}(f)(x_2, \dots, x_n)$ for some function $f \in V$ and some $n < \omega$. Let $\langle f_\alpha : \alpha < \kappa \rangle$ be a sequence of functions in V , such that for each α , there is n_α such that $\text{dom } f_\alpha = \mathcal{P}_\kappa(\kappa^{+\gamma_2}) \times \dots \times \mathcal{P}_\kappa(\kappa^{+\gamma_{n_\alpha}})$. Then $\langle j_{1,\omega}(f_\alpha)(x_2, \dots, x_{n_\alpha}) : \alpha < \kappa \rangle$ can be computed from $j_{1,\omega}(\langle f_\alpha : \alpha < \kappa \rangle)$ and $\langle x_n : 2 \leq n < \omega \rangle$.

For all $\alpha < j_{1,\omega}(\kappa)$, there are $n < \omega$ and $\beta < j_{1,n}(\kappa)$ such that $\alpha = j_{n,\omega}(\beta)$, and $\alpha = \beta$ since $\text{crit}(j_{n,\omega}) = j_{1,n}(\kappa)$. By GCH and the nature of the measures, for $2 \leq n < \omega$, $\kappa^{+\gamma_n} < j_{1,n}(\kappa) < \kappa^{+\gamma}$. Therefore, $j_{1,\omega}(\kappa) = \kappa^{+\gamma}$. Furthermore, an easy counting argument shows that $j_{1,\omega}(\kappa^{+\gamma+1}) = \kappa^{+\gamma+1}$.

By Lemma 13, $V[C_0 \times C_1] \models (\kappa^{+\gamma+1}, \kappa^{+\gamma}) \twoheadrightarrow (\mu^{+\gamma+1}, \mu^{+\gamma})$. Let $\mathfrak{A} \in N_\omega[G]$ be an algebra on $\kappa^{+\gamma+1} = (j_{1,\omega}(\kappa))^+{}^{N_\omega[G]}$. In $V[C_0 \times C_1]$, there is $\mathfrak{B} \prec \mathfrak{A}$ of size $\mu^{+\gamma+1}$ such that $|\mathfrak{B} \cap \kappa^{+\gamma}| = \mu^{+\gamma}$. By the closure of $N_\omega[G]$, $\mathfrak{B} \in N_\omega[G]$. This shows that $N_\omega[G]$ satisfies the desired instance of Chang's Conjecture, and thus by elementarity that p_1 forces $(\kappa^+, \kappa) \twoheadrightarrow (\mu^{+\gamma+1}, \mu^{+\gamma})$. This completes the proof of Theorem 15.

Corollary 23. *Suppose $\mathbb{P} = \mathbb{P}(\mu, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K})$ is as above. Then there is a condition $p \in \mathbb{P}$ of length 0 that forces*

$$\begin{aligned} (\mu^{+\delta+1}, \mu^{+\delta}) &\twoheadrightarrow (\mu^{+\sum_1^n \delta_i + \gamma + 1}, \mu^{+\sum_1^n \delta_i + \gamma}) \twoheadrightarrow \\ &(\mu^{+\sum_1^m \delta_i + \gamma + 1}, \mu^{+\sum_1^m \delta_i + \gamma}) \twoheadrightarrow (\mu^{+\gamma+1}, \mu^{+\gamma}) \end{aligned}$$

for $1 \leq m < n < \omega$.

Proof. Note that it is forced that $\mu^{+\gamma} = \kappa_1^{+\gamma}$, and for each $n \geq 1$, $\kappa_n^{+\delta_n + \gamma} = \kappa_{n+1}^{+\gamma} = \mu^{+\sum_1^n \delta_i + \gamma}$. Let p be a condition of length 0 that forces all Prikry points to be in the set A given by Lemma 13. Fix $1 \leq m < n < \omega$, and let $q \leq p$ be a condition of length n . By Lemma 16, $\mathbb{P} \restriction q$ is isomorphic to a restriction of

$$\text{Col}(\mu, \kappa_1) \times \dots \times \text{Col}(\kappa_{n-1}^{+\delta_{n-1}+2}, \kappa_n) \times \mathbb{P}(\kappa_n^{+\delta_n+2}, \vec{\gamma}', \vec{\delta}', \vec{U}', \vec{K}'),$$

where s' denotes the shift of a sequence s by n . By Lemma 13, this product forces $(\kappa_n^{+\gamma+1}, \kappa_n^{+\gamma}) \twoheadrightarrow (\kappa_m^{+\gamma+1}, \kappa_m^{+\gamma}) \twoheadrightarrow (\mu^{+\gamma+1}, \mu^{+\gamma})$. The last two terms of the product are isomorphic to a restriction of $\mathbb{P}(\kappa_{n-1}^{+\delta_{n-1}+2}, \vec{\gamma}'', \vec{\delta}'', \vec{U}'', \vec{K}'')$ to a condition of length 1, where s'' denotes the shift of the original sequences s by $n-1$. By the argument for Case 3 of Theorem 15, this forces $(\kappa_n^{+\sum_1^\infty \delta_i + 1}, \kappa_n^{+\sum_1^\infty \delta_i}) \twoheadrightarrow (\kappa_n^{+\gamma+1}, \kappa_n^{+\gamma})$. \square

Our methods are not limited to getting $(\aleph_{\beta+1}, \aleph_\beta) \twoheadrightarrow (\aleph_{\alpha+1}, \aleph_\alpha)$ where α and β are countable. For example, if we opt not to interleave collapses in the Gitik-Sharon forcing, we obtain:

Porism 24. *Let $\alpha \geq \omega$ be a countable limit ordinal, and let κ be a $\kappa^{\alpha+1}$ -supercompact cardinal. Then there is a generic extension in which $(\lambda^+, \lambda) \twoheadrightarrow (\aleph_{\alpha+1}, \aleph_\alpha)$, and another in which $(\lambda^{\alpha+1}, \lambda^{\alpha+1}) \twoheadrightarrow (\lambda^+, \lambda)$, where in both cases $\text{cf}(\lambda) = \omega$ and $\aleph_\lambda = \lambda$.*

4. SINGULAR GLOBAL CHANG'S CONJECTURE BELOW \aleph_{ω^ω}

In this section we will prove the following theorem:

Theorem 25. *If there is a model of ZFC with a cardinal δ which is $\delta^{+\omega+1}$ -supercompact and Woodin for supercompactness, then there is a model in which $(\aleph_{\alpha+1}, \aleph_\alpha) \twoheadrightarrow (\aleph_{\beta+1}, \aleph_\beta)$ holds for all limit $\beta < \alpha < \omega^\omega$ (including $\beta = 0$).*

We will need the stronger large cardinal assumption so that we can use a Gitik-Sharon-style forcing over a model where Chang's Conjecture already holds quite frequently.

Definition. *A cardinal δ is called Woodin for supercompactness when for every $A \subseteq \delta$ there is $\kappa < \delta$ such that for all $\lambda \in [\kappa, \delta)$, there is a normal κ -complete ultrafilter U on $\mathcal{P}_\kappa(\lambda)$ such that $j_U(A) \cap \lambda = A \cap \lambda$.*

Like Woodin cardinals, Woodin for supercompactness cardinals need not be even weakly compact, but they have higher consistency strength than supercompact cardinals. Every almost-huge cardinal is Woodin for supercompactness. Woodin for supercompact cardinals are the same as Vopěnka cardinals (see [17]).

Lemma 26. *Suppose GCH and δ is $\delta^{+\omega+1}$ -supercompact and Woodin for supercompactness. Then there is a model of ZFC in which GCH holds, there is a supercompact cardinal, and for all $\alpha < \beta$,*

$$(\beta^{+\omega+1}, \beta^{+\omega}) \twoheadrightarrow (\alpha^{+\omega+1}, \alpha^{+\omega}).$$

Furthermore, such instances of Chang's Conjecture are preserved by forcing over this model with any $(\alpha^{+\omega+1}, \alpha^{+\omega+1})$ -distributive forcing of size $< \beta^{+\omega}$.

Proof. Let $A \subseteq \delta$ be given by Lemma 13. Let $\langle \alpha_i : i < \delta \rangle$ enumerate the closure of A . Force with the following Easton support iteration $\langle \mathbb{P}_i, \dot{\mathbb{Q}}_i : i \leq \delta, j < \delta \rangle$:

- (1) $\mathbb{Q}_0 = \text{Col}(\omega, \alpha_0^{+\omega}) * \text{Col}(\alpha_0^{+\omega+2}, \alpha_1)$.
- (2) If $i > 0$ and $\alpha_i \in A$, $\Vdash_i \dot{\mathbb{Q}}_i = \text{Col}(\alpha_i^{+\omega+2}, \alpha_{i+1})$.
- (3) If $i > 0$ and $\alpha_i \notin A$, $\Vdash_i \dot{\mathbb{Q}}_i = \text{Col}(\alpha_i^+, \alpha_{i+1})$.

It is easy to see that this iteration forces that for all infinite $\alpha < \delta$,

$$(\alpha^{+\omega})^{V^{\mathbb{P}_\delta}} = (\beta^{+\omega})^V,$$

for some $\beta \in A$. By standard arguments, δ remains inaccessible in $V^{\mathbb{P}_\delta}$.

Suppose that in $V^{\mathbb{P}_\delta}$, $\alpha < \alpha^{+\omega} < \beta < \delta$, and let $i < j$ be such that

$$(\alpha^{+\omega})^{V^{\mathbb{P}_\delta}} = (\alpha_i^{+\omega})^V \text{ and } (\beta^{+\omega})^{V^{\mathbb{P}_\delta}} = (\alpha_j^{+\omega})^V.$$

Then \mathbb{P}_δ factors as $\mathbb{P}_i * \mathbb{P}_j / \mathbb{P}_i * \mathbb{P}_\delta / \mathbb{P}_j$, where $|\mathbb{P}_i| \leq \alpha_i^{+\omega}$, $\mathbb{P}_j / \mathbb{P}_i$ is forced to be $\alpha_i^{+\omega+2}$ -closed and of size $\leq \alpha_j$, and $\mathbb{P}_\delta / \mathbb{P}_j$ is forced to be $\alpha_j^{+\omega+2}$ -closed.

Suppose \mathbb{Q} is an $(\alpha_i^{\omega+1}, \alpha_i^{\omega+1})$ -distributive forcing of size $< \alpha_j^{+\omega}$ in $V^{\mathbb{P}_\delta}$. Then $\mathbb{Q} \in V^{\mathbb{P}_j}$. Since \mathbb{P}_i forces that $\mathbb{P}_j / \mathbb{P}_i * \mathbb{Q}$ is $(\alpha_i^{+\omega+1}, \alpha_i^{+\omega+1})$ -distributive, Lemma 13 implies that $\mathbb{P}_j * \mathbb{Q}$ forces $(\alpha_j^{+\omega+1}, \alpha_j^{+\omega}) \twoheadrightarrow (\alpha_i^{+\omega+1}, \alpha_i^{+\omega})$. This is preserved by $\mathbb{P}_\delta / \mathbb{P}_j$, which remains $(\alpha_j^{+\omega+1}, \infty)$ -distributive after forcing with \mathbb{Q} .

Finally, we need to find a supercompact. In V , let $\kappa < \delta$ be given by Woodin for supercompactness with respect to A . Let $\lambda > \kappa$ be an inaccessible limit point of A . Let U be a normal κ -complete ultrafilter on $\mathcal{P}_\kappa(\lambda)$ such that $j_U(A) \cap \lambda = A \cap \lambda$. We have that $j_U(\mathbb{P}_\kappa) = \mathbb{P}_\kappa * \mathbb{P}_\lambda / \mathbb{P}_\kappa * \mathbb{Q}$, for some \mathbb{Q} that is forced to be λ^+ -closed. Let $G_\delta \subseteq \mathbb{P}_\delta$ be generic, and let $G_\lambda = G_\delta \restriction \mathbb{P}_\lambda$. By GCH, $j_U(\kappa) < \lambda^{++}$ and $j_U(\lambda^{++}) = \lambda^{++}$, so we may build $H \subseteq \mathbb{Q}$ in $V[G_\lambda]$ that is generic over $M[G_\lambda]$. Thus we can extend the embedding to $j : V[G_\kappa] \rightarrow M[G_\lambda * H]$. Since $M[G_\lambda * H]$ is λ -closed in $V[G_\lambda]$ and $\mathbb{P}_\lambda / G_\kappa$ is κ -directed-closed, there is $p \in j_U(\mathbb{P}_\lambda) / (G_\lambda * H)$ below

$j[G_\lambda/G_\kappa]$. Since $|\mathbb{P}_\lambda| = \lambda$ and $j_U(\lambda^+) < \lambda^{++}$, we can build $K \subseteq j_U(\mathbb{P}_\lambda)/(G_\lambda * H)$ below p in $V[G_\lambda]$ that is generic over $M[G_\lambda * H]$. Thus we can extend the embedding to $j : V[G_\lambda] \rightarrow M[G_\lambda * H * K]$. This shows that κ is λ -supercompact in $V[G_\lambda]$, a property that is preserved by $\mathbb{P}_\delta/G_\lambda$. Thus, $V_\delta[G_\delta] \models$ "There is a supercompact cardinal." \square

Let us work in a model satisfying the conclusion of the above lemma. We define by induction on $1 \leq n \leq \omega$ the class of "order- n Gitik-Sharon forcings" (abbreviated by GS_n). Formally, we fix a large enough regular θ and define these inductively as subsets of H_θ , but it will be clear that choice of θ is irrelevant, and for $\theta < \theta'$, $\text{GS}_n^{H_\theta} = \text{GS}_n^{H_{\theta'}} \cap H_\theta$. Each order- n forcing will add a club of ordertype ω^n to a large cardinal κ , consisting of former inaccessibles, while preserving that κ is a cardinal, collapsing $\kappa^{+\omega \cdot n}$ to κ , and preserving larger cardinals.

GS_1 is the collection of forcings of the form $\mathbb{P}(\mu, \vec{\omega}, \vec{\omega}^2, \vec{U}, \vec{K})$, as defined in the previous section, where $\vec{\omega}$ is the identity sequence $\langle 1, 2, 3, \dots \rangle$, and $\vec{\omega}^2$ is the constant sequence $\langle \omega, \omega, \omega, \dots \rangle$. We will define partial orders in GS_n using sequences $\langle U_\alpha, K_\alpha : \alpha < \omega \cdot n \rangle$ such that:

- (1) There is a $\kappa > \omega$ such that each U_α is an ultrafilter with completeness κ (to be called the *critical point* of the sequence).
- (2) For $n < \omega$, U_n is a normal ultrafilter on $\mathcal{P}_\kappa(\kappa^{+n})$ and for $\omega \leq \alpha < \omega \cdot n$, U_α is a normal ultrafilter on $\mathcal{P}_\kappa(H_{\kappa+\alpha})$.
- (3) For $\alpha < \omega \cdot n$, if $j_\alpha : V \rightarrow M_\alpha$ is the ultrapower embedding from U_α , then K_α is $\text{Col}(\kappa^{+\alpha+\omega+2}, j_\alpha(\kappa))^{M_\alpha}$ -generic over M_α .

A partial order $\mathbb{P} \in \text{GS}_n$ will be determined by the choice of a sequence $d = \langle U_\alpha, K_\alpha : \alpha < \omega \cdot n \rangle$ as above and a regular cardinal $\mu < \text{crit}(d)$. Suppose $n > 1$ and that we have defined GS_m for $m < n$, and we have a function defined on pairs $(\mu, d) \in H_\theta$ that outputs a partial order $\mathbb{P}(\mu, d) \in \text{GS}_m$ whenever d is a sequence of length $\omega \cdot m$ as above and $\mu < \text{crit}(d)$ is regular.

Let $d = \langle U_\alpha, K_\alpha : \alpha < \omega \cdot n \rangle$ be as above and let $\mu < \text{crit}(d)$ be regular. Conditions in $\mathbb{P}(\mu, d) \in \text{GS}_n$ take the form:

$$p = \langle f_0, e_1, (x_1, a_1), f_1, e_2, (x_2, a_2), f_2, \dots, e_l, (x_l, a_l), f_l, \vec{F} \rangle.$$

The *stem* of p is the initial segment obtained by removing \vec{F} . The *length* of p as above is l . We require:

- (1) For $1 \leq i \leq l$:
 - (a) $|x_i| < \kappa$, $x_i \prec H_{\kappa+\omega \cdot (n-1)+i}$, $\kappa_i := x_i \cap \kappa$ is inaccessible, the transitive collapse of x_i is $H_{\kappa_i+\omega \cdot (n-1)+i}$, and $\langle U_\alpha, K_\alpha : \alpha < \omega \cdot (n-1) \rangle \in x_i$.
 - (b) If $\pi : x_i \rightarrow H$ is the transitive collapse map, put $\pi(\langle U_\alpha, K_\alpha : \alpha < \omega \cdot (n-1) \rangle) := \langle u_\alpha^i, k_\alpha^i : \alpha < \omega \cdot (n-1) \rangle := d_i$. We require that a_i is a sequence of functions $\langle b_\alpha^i : \alpha < \omega \cdot (n-1) \rangle$ such that $\text{dom}(b_\alpha^i) \in u_\alpha^i$ and $[b_\alpha^i]_{u_\alpha^i} \in k_\alpha^i$.
- (2) $f_0 \in \text{Col}(\mu, \kappa)$, and if x_0 is defined, then $\langle f_0 \rangle^\frown e_1^\frown a_1 \in \mathbb{P}(\mu, d_1)$.
- (3) For $1 \leq i < l$, $x_i \in x_{i+1}$, and $\langle f_i \rangle^\frown e_{i+1}^\frown a_{i+1} \in \mathbb{P}(\kappa_i^{+\omega \cdot n+2}, d_{i+1})$.
- (4) $f_l \in \text{Col}(\kappa_l^{+\omega \cdot n+2}, \kappa)$.
- (5) \vec{F} is a sequence of functions $\langle F_\alpha : \alpha < \omega \cdot n \rangle$ such that for each α , $\text{dom } F_\alpha \in U_\alpha$ and $[F_\alpha]_{U_\alpha} \in K_\alpha$.

Suppose we have two conditions

$$p = \langle f_0, e_1, (x_1, a_1), f_1, \dots, e_l, (x_l, a_l), f_l, \vec{F} \rangle;$$

$$q = \langle f'_0, e'_1, (x'_1, a'_1), f'_1, \dots, e'_m, (x'_m, a'_m), f'_m, \vec{F}' \rangle.$$

We put $q \leq p$ when:

- (1) $m \geq l$, and for $1 \leq i \leq l$, $x_i = x'_i$.
- (2) For $i \leq l$, $f'_i \supseteq f_i$.
- (3) For $1 \leq i \leq l$, $\langle f'_{i-1} \rangle \frown e'_i \frown a'_i \leq \langle f_{i-1} \rangle \frown e_i \frown a_i$ in the relevant partial order from GS_{n-1} .
- (4) For $l < i \leq m$, $x'_i \in \text{dom } F_{\omega \cdot (n-1) + i}$ and $f'_i \supseteq F_{\omega \cdot (n-1) + i}(x_i)$.
- (5) $\vec{F} \restriction \omega \cdot (n-1) \in x$ if $x = x'_i$ for $l < i \leq m$, or if $x \in \text{dom } F'_{\omega \cdot (n-1) + k}$ for $k > m$.
- (6) Put $f_k = F_{\omega \cdot (n-1) + k}(x'_k)$ for $l < k < m$. If $l < i \leq m$ and $\pi : x_i \rightarrow H$ is the transitive collapse map, then $\langle f'_{i-1} \rangle \frown e'_i \frown a'_i \leq \langle f_{i-1} \rangle \frown \pi(\vec{F} \restriction \omega \cdot (n-1))$.
- (7) For each $\alpha < \omega \cdot n$, $\text{dom } F'_\alpha \subseteq \text{dom } F_\alpha$, and for each $x \in \text{dom } F'_\alpha$, $F'_\alpha(x) \supseteq F_\alpha(x)$.

Finally, we may define the order- ω forcings which generically stack the order- n forcings for finite n . Everything looks quite similar, except now our sequences of functions \vec{F} have length ω^2 , and stems of length $n > 0$ look like stems of length-1 conditions from forcings in GS_{n+1} .

We define some notions to describe the conditions in our forcings. A *type-1 sequence* is a natural number. For $n > 1$, a *type- n sequence* is a finite sequence of type- $(n-1)$ sequences. We can define inductively a partial order on these sequences. For a type-1 sequence, this is just the usual linear order. If $s = \langle t_1, \dots, t_l \rangle$ and $s' = \langle t'_1, \dots, t'_m \rangle$ are of type- n , then we say $s' \geq s$ when $m \geq l$ and $t'_i \geq t_i$ for $1 \leq i \leq m$. It is easy to see by induction that this ordering is upward-directed.

If $p \in \mathbb{P} \in \text{GS}_1$, then by the *shape* of p we mean its length. If $s = \langle t_1, \dots, t_l \rangle$ is a type- n sequence, and

$$p = \langle f_0, e_1, (x_1, a_1), f_1, \dots, e_l, (x_l, a_l), f_l, \vec{F} \rangle \in \mathbb{P} \in \text{GS}_n,$$

then we say, inductively, that the stem of p has *shape* s if each $\langle f_{i-1} \rangle \frown e_i \frown a_i$ has shape t_i . If $s = \langle t_1, \dots, t_l \rangle$ is such that each t_i is a type- i sequence, and $p \in \mathbb{P} \in \text{GS}_\omega$ takes the same form as above, then we say p has shape s if each $\langle f_{i-1} \rangle \frown e_i \frown a_i$ has shape t_i . Note that if $q \leq p$, then the shape of q is greater or equal to the shape of p in the ordering on sequences. Since the shape of a condition only depends on its stem, we will also speak of the shapes of stems and their subsequences.

Suppose $\mathbb{P} \in \text{GS}_n$ for $n \leq \omega$. For conditions $p, q \in \mathbb{P}$, we say $p \leq^* q$ if $p \leq q$ and they have the same shape. If $p \leq q$ and $\text{stem}(p)$ is an initial segment of $\text{stem}(q)$, then we say $p \preceq q$. We have an operation $p \wedge \vec{F}$ defined similarly as before.

Lemma 27. *Suppose $\mathbb{P}(\mu, d) \in \text{GS}_n$, $\mu > \omega$, and $c : \mathbb{P} \rightarrow 3$ is a decisive coloring.*

- (1) *There is a sequence \vec{F} such that for every condition p , every two $r, r' \preceq p \wedge \vec{F}$ of the same shape have the same color.*
- (2) *For every condition p , there is $q \leq^* p$ such that every two $r, r' \preceq q$ of the same shape have the same color.*

Proof. The case $n = 1$ was proven in Lemma 17. Assume $n > 1$ and the lemma holds for GS_m , $m < n$. Let $\mathbb{P}(\mu, d) \in \text{GS}_n$, with $\text{crit}(d) = \kappa$ and $\omega < \mu < \kappa$. Like before, we prove (1) by showing the following claim by induction: For all $l < \omega$ and all decisive colorings of the conditions of length l , there is \vec{F}_l such that for all $m \leq l$ and every condition p of length m , every two $r, r' \preceq p \wedge \vec{F}_l$ of the same shape and of length l have the same color. This suffices, since we can find \vec{F} that is a lower bound to the countably many \vec{F}_l . Suppose $l = 0$ and c is such a coloring. For every $s \in \text{Col}(\mu, \kappa)$, choose if possible some \vec{F}_s such that $c(\langle s \rangle \frown \vec{F}_s) > 0$. Using the closure of the collapses and diagonal intersections, we may select a single sequence \vec{F} such that $\langle s \rangle \wedge \vec{F} \leq \langle s \rangle \frown \vec{F}_s$ for all s .

Suppose the claim is true for $m < l$. Let c be a decisive coloring of the conditions of length l . For each stem $s = \langle f_0^s, \dots, (x_{l-1}^s, a_{l-1}^s), f_{l-1}^s \rangle$ of length $l-1$, and each

candidate (x, a) for the last node in a one-step extension containing s , we can define a coloring $c_{s,x}$ on conditions of the form $\langle f_{l-1}^s \rangle \frown e \frown a \in \mathbb{P}(\kappa_{l-1}^{+\omega \cdot n+2}, d_x)$ as follows. First, as in the proof of Lemma 17, we find a sequence $\vec{F} = \langle F_\alpha : \alpha < \omega \cdot n \rangle$ such that for each stem s , each $x \in \text{dom } F_{\omega \cdot (n-1)+l}$, and each choice of e and a such that there are f and \vec{H} such that $s \frown \langle e, (x, a), f \rangle \frown \vec{H}$ is a condition below $s \frown \vec{F}$ with color > 0 , then already $s \frown \langle e, (x, a), F_{\omega \cdot (n-1)+l}(x) \rangle \frown \vec{F}$ has this color. We can then define $c_{s,x}(\langle f_{l-1}^s \rangle \frown e \frown a) = c(s \frown \langle e, (x, a), F_{\omega \cdot (n-1)+l}(x) \rangle \frown \vec{F})$. By the induction hypothesis on the order of Gitik-Sharon forcing, for each such s, x , there is a choice of $a_{s,x}$ such that $c_{s,x}(\langle f_{l-1}^s \rangle \frown e \frown a_{s,x})$ depends only on the shape of e , for conditions below $\langle f_{l-1}^s \rangle \frown a_{s,x}$. For each x , we can use diagonal intersections to select a sequence a_x such that for all s , $\langle f_{l-1}^s \rangle \wedge a_x \leq \langle f_{l-1}^s \rangle \frown a_{s,x}$.

In the ultrapower by $U = U_{\omega \cdot (n-1)+l}$, the function $x \mapsto a_x$ represents a sequence of functions \vec{G} strengthening $\vec{F} \restriction \omega \cdot (n-1) = \pi(j_U(\vec{F} \restriction \omega \cdot (n-1)))$, where π is the transitive collapse of $j_U[H_{\kappa+\omega \cdot (n-1)+l}]$. Let \vec{F}' be \vec{F} with the initial segment below $\omega \cdot (n-1)$ replaced by \vec{G} . Thus we have for each stem s of length $l-1$, a set $A_s \in U_{\omega \cdot (n-1)+l}$ such that for all $x \in A_s$, $a_x = \pi_x(F' \restriction \omega \cdot (n-1))$, and the color of $s \frown \langle e, (x, a_x), F'_{\omega \cdot (n-1)+l}(x) \rangle \frown \vec{F}'$ depends only on the shape of e , if $\langle f_{l-1}^s \rangle \frown e \frown a_x \leq \langle f_{l-1}^s \rangle \frown a_x$. Let A^* be the diagonal intersection of the A_s , and let \vec{F}'' be F' restricted to A^* at coordinate $\omega \cdot (n-1) + l$.

Now for any condition p of shape $\langle t_1, \dots, t_{l-1} \rangle$, the color under c of any $q \leq p \wedge \vec{F}''$ of shape $\langle t_1, \dots, t_{l-1}, t \rangle$ depends only on t . So for each type- $(n-1)$ sequence t , let c_t color the length- $(l-1)$ conditions accordingly. Note that each c_t inherits decisiveness from c . By the induction hypothesis, for each t , there is a sequence \vec{F}_t such that for all $m < l-1$ and all p of length m , every $q \leq p \wedge \vec{F}_t$ of length $l-1$ has a color under c_t depending only on the shape of q . If \vec{F}''' is a lower bound to the countably many sequences \vec{F}_t , then \vec{F}''' satisfies the inductive claim for l . This concludes the argument for (1).

To show (2), let us assume inductively that it holds for GS_m , $m < n$. Let $\mathbb{P} \in \text{GS}_n$, let $c : \mathbb{P} \rightarrow 3$ be decisive, and let \vec{F} be given by (1). Let $p \in \mathbb{P}$, with $\text{stem}(p) = \langle f_0, \dots, f_{l-1}, e_l, (x_l, a_l), f_l \rangle$. For every end-extension $q = \text{stem}(p) \frown s \frown \vec{F}$ of $p \wedge \vec{F}$, the color of q depends only on the shape of s . Using the closure of $\text{Col}(\kappa_l^{+\omega \cdot n+2}, \kappa)$, we can find $f'_l \supseteq f_l$ such that for every strengthening s of the initial segment of $\text{stem}(p)$ before f_l , and every type- n sequence t , if there is $f \supseteq f'_l$ such that some s' of shape t with $s \frown \langle f \rangle \frown s' \frown \vec{F} \leq p \wedge \vec{F}$ has color > 0 , then already $s \frown \langle f'_l \rangle \frown s' \frown \vec{F}$ has this color.

Now for each type- n sequence t , and each strengthening s of $\text{stem}(p)$ before f_{l-1} , we have a coloring $c_{s,t}$ of the conditions $\langle f \rangle \frown e \frown a \leq \langle f_{l-1} \rangle \frown e_l \frown a_l$ according to the color under c of $s \frown \langle f, e, (x_l, a), f'_l \rangle \frown s' \frown \vec{F}$, where s' is anything of shape t , such that the resulting condition is below $p \wedge \vec{F}$. Using the inductive hypothesis and the weak closure of $\mathbb{P}(\kappa_{l-1}^{+\omega \cdot n+2}, d_l)$, we find $\langle f'_{l-1} \rangle \frown e'_l \frown a'_l \leq^* \langle f_{l-1} \rangle \frown e_l \frown a_l$ such that any two extensions of the former of the same shape have the same color under every $c_{s,t}$. As a result, we have that for any s strengthening $\text{stem}(p)$ before f_{l-1} , for any two r, r' of the same shape below $s \frown \langle f'_{l-1}, e'_l, (x_l, a'_l), f'_l \rangle \frown \vec{F}$, for which s is an initial segment of both, $c(r) = c(r')$. We continue this process in the same fashion down the stem of p , in a total of l steps, so that at step $k \leq l$, we find $\langle f'_{l-k-1} \rangle \frown e'_{l-k} \frown a'_{l-k} \leq^* \langle f_{l-k-1} \rangle \frown e_{l-k} \frown a_{l-k}$, such that for every strengthening s of the initial segment of $\text{stem}(p)$ before f_{l-k-1} , any two conditions r, r' of the same shape, with s as an initial segment, and below $s \frown \langle f'_{l-k-1}, e'_{l-k}, (x_{l-k}, a'_{l-k}), f'_{l-k}, \dots, (x_l, a'_l), f'_l, \vec{F} \rangle$, we have $c(r) = c(r')$. Eventually we reach the desired condition $q \leq^* p$.

The inductive argument for GS_ω is entirely similar. \square

Corollary 28. *If $\mathbb{P}(\mu, d) \in \text{GS}_n$, $1 \leq n \leq \omega < \mu$, then $\langle \mathbb{P}(\mu, d), \leq, \leq^* \rangle$ is a Prikry-type forcing. Furthermore, if p_0 is a condition of length $m > 0$, then $\mathbb{P}(\mu, d) \restriction p_0$ is canonically isomorphic to*

$$\mathbb{P}(\mu, d_1) \restriction \langle f_0 \rangle \frown e_1 \frown a_1 \times \cdots \times \mathbb{P}(\kappa_{m-1}^{+\omega \cdot n+2}, d_m) \restriction \langle f_{m-1} \rangle \frown e_m \frown a_m \times \mathbb{Q},$$

where \mathbb{Q} is a weakly $\kappa_m^{+\omega \cdot m+2}$ -closed Prikry-type forcing.

Proof. Let σ be a sentence in the forcing language, and color conditions 0 if they do not decide σ , 1 if they force σ , and 2 if they force $\neg\sigma$. Let $p \in \mathbb{P}(\mu, d)$, and let $q \leq^* p$ be such that any two extensions of q of the same shape have the same color. If q does not decide σ , then by the fact that the ordering on sequences is upward-directed, we can find $r, r' \leq q$ of the same shape that force opposite decisions about σ , a contradiction.

For the second claim, the map is the obvious one, where the elements of \mathbb{Q} are the tail-ends beyond place m , of conditions below p_0 . Let us write $\mathbb{P}(\mu, d) \restriction p_0$ as $\mathbb{R} \times \mathbb{Q}$. From any decisive coloring c of the conditions in \mathbb{Q} , we can define a decisive coloring c' of $\mathbb{R} \times \mathbb{Q}$ by setting $c'(r, q) = c(q)$. Given any $q \in \mathbb{Q}$, we can find $p \leq^* (1, q)$ such that any two $p', p'' \leq p$ of the same shape have the same color under c' . This means that any two $q', q'' \leq q$ of the same shape have the same color under c . Then we apply the argument of the previous paragraph. \square

If $\mathbb{P}(\mu, d) \in \text{GS}_n$ for $1 < n < \omega$, with $\text{crit}(d) = \kappa$, and $G \subseteq \mathbb{P}(\mu, d)$ is generic over V , then we have a sequence $\langle x_i, G_i : 1 \leq i < \omega \rangle$ such that:

- (1) Each x_i is a typical point in $\mathcal{P}_\kappa(H_{\kappa^{+\omega \cdot (n-1)+i}})$.
- (2) $\langle x_i : 1 \leq i < \omega \rangle$ is \in - and \subseteq -increasing, with $\bigcup_i x_i = H_{\kappa^{+\omega \cdot n}}$.
- (3) G_1 is $\mathbb{P}(\mu, d_1)$ -generic, and for $i > 1$, G_i is $\mathbb{P}(\kappa_{i-1}^{+\omega \cdot n+2}, d_i)$ -generic, where κ_i and d_i are as in the definition of GS_n .

From $\langle x_i, G_i : 1 \leq i < \omega \rangle$, we can recover G as the collection of all conditions $\langle f_0, e_1, (x_1, a_1), f_1, \dots, e_l, (x_l, a_l), f_l, \vec{F} \rangle$ such that:

- (1) $\langle x_1, \dots, x_l \rangle$ is an initial segment of $\langle x_i : 1 \leq i < \omega \rangle$.
- (2) For $i > l$, $F \restriction \omega \cdot (n-1) \in x_i \in \text{dom } F_{\omega \cdot n+i}$.
- (3) For $1 \leq i \leq l$, $\langle f_{i-1} \rangle \frown e_i \frown a_i \in G_i$.
- (4) Putting $f_i = F_{\omega \cdot n+i}(x_i)$ for $i > l$, $\langle f_{i-1} \rangle \frown \vec{F} \restriction \omega \cdot (n-1) \in G_i$.

We need the following characterization of genericity, proof of which is essentially the same as for Lemma 20:

Lemma 29. *Suppose $d = \langle U_\alpha, K_\alpha : \alpha < \omega \cdot n \rangle$ and $\mathbb{P}(\mu, d) \in \text{GS}_n$, with $\omega < \mu < \text{crit}(d) = \kappa$. Suppose in some outer model $W \supseteq V$, there is a sequence $\langle x_i, G_i : 1 \leq i < \omega \rangle$ as above.*

Then this sequence generates a V -generic filter G for $\mathbb{P}(\mu, d)$ iff for every sequence $\vec{F} = \langle F_\alpha : \alpha < \omega \cdot n \rangle$ such that $\langle \emptyset \rangle \frown \vec{F}$ is a condition, there is $m < \omega$ such that for all $k \geq m$, $\vec{F} \restriction \omega \cdot (n-1) \in x_k \in \text{dom } F_{\omega \cdot (n-1)+k}$, and

$$\langle F_{\omega \cdot (n-1)+k}(x_k) \rangle \frown \pi_{k+1}(\vec{F} \restriction \omega \cdot (n-1)) \in G_{k+1},$$

where π_{k+1} is the transitive collapse of x_{k+1} .

To prove the main theorem, we will show by induction that, in a model satisfying the conclusion of Lemma 26, if $\mu = \nu^{+\omega \cdot k+2}$ and $\mathbb{P}(\mu, d) \in \text{GS}_n$, for $1 \leq k, n < \omega$, then $\mathbb{P}(\mu, d)$ forces that $(\nu^{+\alpha+1}, \nu^{+\alpha}) \twoheadrightarrow (\nu^{+\beta+1}, \nu^{+\beta})$ holds for all limit ordinals $\omega \leq \beta < \alpha \leq \omega^{n+1}$. Note that we include the case $\nu = 0$ so that the lower pair may be (\aleph_1, \aleph_0) .

For the base case, suppose $\mu = \nu^{+\omega \cdot k+2}$, for $1 \leq k < \omega$, and $\mathbb{P}(\mu, d) \in \text{GS}_1$, with $\text{crit}(d) = \kappa$. By Lemma 16 and the preservation claim of Lemma 26, we have that in $V^{\mathbb{P}(\mu, d)}$, $(\nu^{+\omega \cdot i+1}, \nu^{+\omega \cdot i}) \rightarrow (\nu^{+\omega \cdot j+1}, \nu^{+\omega \cdot j})$ holds for all $1 \leq j < i < \omega$. Using again the fact that for $\alpha < \kappa$, $(\kappa^{+\omega+1}, \kappa^{+\omega}) \rightarrow (\alpha^{+\omega+1}, \alpha^{+\omega})$ is indestructible by any $\alpha^{+\omega+2}$ -closed forcing of size κ , the iterated ultrapower construction in the previous section shows that $\mathbb{P}(\mu, d)$ also forces $(\kappa^+, \kappa) \rightarrow (\nu^{+\omega \cdot i+1}, \nu^{+\omega \cdot i})$ for $1 \leq i < \omega$.

Assuming that the inductive claim holds for n , let us first argue for the weaker claim that if $\mu = \nu^{+\omega \cdot k+2}$, for $1 \leq k < \omega$, and $\mathbb{P}(\mu, d) \in \text{GS}_{n+1}$, then $\mathbb{P}(\mu, d)$ forces $(\nu^{+\alpha+1}, \nu^{+\alpha}) \rightarrow (\nu^{+\beta+1}, \nu^{+\beta})$ to hold for all limit ordinals $\omega \leq \beta < \alpha < \omega^{n+2}$ (where the last inequality is strict). A generic $G \subseteq \mathbb{P}(\mu, d)$ introduces a Prikry sequence of generics for GS_n forcings, $\langle G_i : 1 \leq i < \omega \rangle$, where G_1 is generic for $\mathbb{P}(\mu, d_1)$, and for $i \geq 2$, G_i is generic for $\mathbb{P}(\kappa_{i-1}^{+\omega \cdot (n+1)+2}, d_i)$. In $V[G_1]$, $\kappa_1 = \nu^{+\omega^{n+1}}$, its successor is $(\kappa_1^{+\omega \cdot n+1})^V$, and we have $(\nu^{+\alpha+1}, \nu^{+\alpha}) \rightarrow (\nu^{+\beta+1}, \nu^{+\beta})$ for all limit ordinals $\omega \leq \beta < \alpha \leq \omega^{n+1}$. This is preserved by adjoining $\langle G_j : 2 \leq j < \omega \rangle$, which adds no subsets of $(\kappa_1^{+\omega \cdot n+1})^V$. For $i > 1$, we have that in $V[G_i]$,

$$(\kappa_{i-1}^{+\omega \cdot n+\alpha+1}, \kappa_{i-1}^{+\omega \cdot n+\alpha}) \rightarrow (\kappa_{i-1}^{+\omega \cdot n+\beta+1}, \kappa_{i-1}^{+\omega \cdot n+\beta}),$$

holds for all limit ordinals $0 \leq \beta < \alpha \leq \omega^{n+1}$. For each such i , these instances of Chang's Conjecture are preserved by adjoining $\langle G_j : i < j < \omega \rangle$, which adds no subsets of $(\kappa_i^{+\omega \cdot n+1})^V$, the $(\omega^{n+1} + 1)^{\text{st}}$ cardinal above κ_{i-1} in the extension, and also by adjoining $G_1 \times \cdots \times G_{i-1}$, which is generic for a $\kappa_{i-1}^{+\omega \cdot n}$ -centered forcing. By the transitivity of Chang's Conjecture, we can combine finitely many instances to bridge the different intervals that lie between adjacent Prikry points, and get the weaker conclusion for $n + 1$.

The hard part is to improve the final inequality to allow $\alpha = \omega^{n+2}$. If the critical point of d as above is κ , then by applying transitivity again, it suffices to show that the extension satisfies $(\kappa^+, \kappa) \rightarrow (\kappa_i^+, \kappa_i)$ for infinitely many i . Towards this, we generalize Claim 22 and produce an iterated ultrapower for which we can find a generic filter for (the image of) a forcing $\mathbb{P} \in \text{GS}_{n+1}$.

Claim 30. *Suppose $1 \leq n < \omega$, W is a model of ZFC, and $\mathbb{P}(\mu, d) \in \text{GS}_n^W$, with $\text{crit}(d) = \kappa$. Suppose $p \in \mathbb{P}(\mu, d)$ is a condition of length l , $p = \langle f_0, \dots, f_l \rangle \cap \vec{F}$. If $l > 0$, let $\nu = \kappa_l^{+\omega \cdot n+2}$ and let \mathbb{R} be such that $\mathbb{P}(\mu, d) \restriction p \cong \mathbb{R} \times \mathbb{Q}$, as in Corollary 28. Otherwise let $\nu = \mu$ and let \mathbb{R} be trivial.*

There is an elementary embedding $j: W \rightarrow W'$, where W' is transitive, $\text{crit } j = \kappa$, $j(\kappa) = \kappa^{+\omega \cdot n}$, and $\kappa^{+\omega \cdot n+1}$ is a fixed point of j . If there is a W -generic filter $H \subseteq \mathbb{R} \times \text{Col}(\nu, \kappa)$, then there is a W' -generic filter $G \subseteq j(\mathbb{P}(\mu, d))$ in $W[H]$ such that $j(p) \in G$. Moreover, $W[H]$ and $W'[G]$ have the same κ -sequences of ordinals.

Proof. First, let us introduce a temporary notation in order to describe generic filters for $\mathbb{P}(\mu, d)$. Every ordinal $\alpha < \omega^\omega$, can be represented using Cantor Normal Form as a sum

$$\alpha = \omega^m \cdot k_m + \cdots + \omega \cdot k_1 + k_0,$$

where $k_i < \omega$ for all i . For $\alpha \neq 0$, let $n_*(\alpha) = \min\{r \mid k_r \neq 0\}$ and let $m_*(\alpha) = k_{n_*(\alpha)}$.

A generic $G \subseteq \mathbb{P}(\mu, d)$ can be unraveled into a sequence $\langle x_\alpha : 1 \leq \alpha < \omega^n \rangle \subseteq \mathcal{P}_\kappa(H_{\kappa^{+\omega \cdot n}})$ and filters $\langle C_\alpha : \alpha < \omega^n \rangle$, from which we can recover G . If $\rho_\alpha = x_\alpha \cap \kappa$, then the ρ_α are increasing, continuous, and cofinal in κ . C_0 is generic for $\text{Col}(\mu, \rho_1)$, and for $\alpha \geq 1$, C_α is generic for $\text{Col}(\rho_\alpha^{+\omega \cdot n_*(\alpha)+\omega+2}, \rho_{\alpha+1})$. If $\beta < \alpha$ and $n_*(\beta) \leq n_*(\alpha)$, then $x_\beta \in x_\alpha$.

Let us note that by unraveling the criteria for being in the filters associated to the sequences, we can recover G in the following way. Let $\vec{F} = \langle F_\alpha : \alpha < \omega \cdot n \rangle$ be a sequence of functions. For each $\alpha < \omega^n$, define a finite sequence $\langle F_\alpha^0, \dots, F_\alpha^{n-n_*(\alpha)-1} \rangle$

by putting $F_\alpha^0 = F_{n_*(\alpha)+m_*(\alpha)}$, and for $0 < k < n - n_*(\alpha)$, $F_\alpha^k = \pi(F_\alpha^{k-1})$, where π is the transitive collapse of $x_{\alpha+\omega^{n-k}}$, if that object is in $\text{dom } \pi$. Put $F'_\alpha = F_\alpha^{n-n_*(\alpha)-1}$. Then we have $\langle \emptyset \rangle \cap \vec{F} \in G$ iff for all $\alpha < \omega^n$, F'_α is defined, $x_\alpha \in \text{dom } F'_\alpha$, and $F'_\alpha(x_\alpha) \in C_\alpha$.

Given a sequence of ultrafilters and guiding generics d , let us construct an iterated ultrapower and a sequence $\langle x_\alpha, C_\beta \mid 1 \leq \alpha < \omega^n, \beta < \omega^n \rangle$ as above. We will assume, by induction on n (simultaneously for all models of ZFC, all sequences d and all generics H) that this process provides a generic filter for the limit ultrapower.

Let μ, d, H, W be as hypothesized, and let $d = \langle U_\alpha, K_\alpha \mid \alpha < \omega \cdot n \rangle$. Let us define by induction on $\omega \cdot l < \alpha \leq \omega^n$, a model N_α and elementary embeddings $j_{\beta, \alpha} : N_\beta \rightarrow N_\alpha$. The choice of the measures which are applied at each step resembles the iterated ultrapower for obtaining a Radin generic filter (see [16]).

Let $\alpha_0 = \omega \cdot l + 1$, and let $N_{\alpha_0} = W$. For limit ordinals α , let N_α be the direct limit of the system $\langle N_\beta, j_{\beta, \gamma} \mid \beta < \gamma < \alpha \rangle$ and let $j_{\beta, \alpha}$ be the corresponding limit embeddings. For $\alpha = \beta + 1$, let $j_{\beta, \alpha} : N_\beta \rightarrow N_\alpha \cong \text{Ult}(N_\beta, j_{\alpha_0, \beta}(U_{\omega \cdot n_*(\beta)+m_*(\beta)}))$, and let $j_{\gamma, \alpha} = j_{\beta, \alpha} \circ j_{\gamma, \beta}$ for $\gamma < \beta$. By Fact 21, N_{ω^n} is well-founded. By counting arguments similar to those in the previous section, we can show that $j_{\alpha_0, \omega^n}(\kappa) = \kappa^{+\omega \cdot n}$, and $\kappa^{+\omega \cdot n+1}$ is fixed.

Let us define a sequence of filters $\langle C_i \mid i < \omega^n \rangle$ and a sequence of sets $\langle x_i \mid 1 \leq i < \omega^n \rangle$. For $i \leq \omega^{n-1} \cdot l$ we extract C_i and x_i from the W -generic filter H .

Let us define the Prikry points for $\alpha > \omega^{n-1} \cdot l$. Let $X_\alpha = \kappa^{+m_*(\alpha)}$ if $n_*(\alpha) = 0$, and $X_\alpha = H_{\kappa^{+\omega \cdot n_*(\alpha)+m_*(\alpha)}}$ otherwise. Let $y_\alpha = j_{\alpha, \omega^n}[j_{\alpha_0, \alpha}(X_\alpha)]$. Note that $y_\alpha = j_{\alpha+1, \omega^n}(j_{\alpha, \alpha+1}[j_{\alpha_0, \alpha}(X_\alpha)])$, and in particular it is in N_{ω^n} . In other words, we take y_α to be the seed of the measure $j_{\alpha_0, \alpha}(U_{n_*(\alpha)+m_*(\alpha)})$, pushed by the map $j_{\alpha+1, \omega^n}$ to the limit model N_{ω^n} . Since the critical point of the elementary map $j_{\alpha+1, \omega^n}$ is above the cardinality of y_α , it acts pointwise.

If $n_*(\alpha) = n - 1$, let $x_\alpha = y_\alpha$. Otherwise, let π be the Mostowski collapse of $y_{\alpha+\omega^{n_*(\alpha)+1}}$ and let $x_\alpha = \pi(y_\alpha)$. Let $C_\alpha = j_{\alpha_0, \alpha}(K_{\omega \cdot n_*(\alpha)+m_*(\alpha)})$. Let us verify that the obtained filter satisfies the requirements of Lemma 29.

Let $m > l$. Let G_m be the filter for the forcing $\mathbb{P}(\rho_{\omega^{n-1} \cdot (m-1)}^{+\omega \cdot n+2}, d_m)^{N_{\omega^{n-1} \cdot m}}$, where $d_m = j_{\alpha_0, \omega^{n-1} \cdot m}(d) \restriction \omega \cdot (n-1)$, which is derived from the sequences $\langle x_\alpha \mid \omega^{n-1} \cdot (m-1) < \alpha < \omega^{n-1} \cdot m \rangle$ and $\langle C_\alpha \mid \omega^{n-1} \cdot (m-1) \leq \alpha < \omega^{n-1} \cdot m \rangle$. Let us assume, by induction, that G_m is an $N_{\omega^{n-1} \cdot m}$ -generic filter. Note that

$$\mathbb{P}(\rho_{\omega^{n-1} \cdot (m-1)}^{+\omega \cdot n+2}, d_m)^{N_{\omega^{n-1} \cdot m}} = \mathbb{P}(\rho_{\omega^{n-1} \cdot (m-1)}^{+\omega \cdot n+2}, d_m)^{N_{\omega^n}},$$

and that G_m is also N_{ω^n} -generic. For $m \leq l$, G_m is derived from the W -generic filter H , and thus it is clearly N_{ω^n} -generic.

Let $z_i = x_{\omega^{n-1} \cdot i}$ for $1 \leq i < \omega$. Let us check that for every sequence $\vec{F} = \langle F_i \mid i < \omega \cdot n \rangle \in N_{\omega^n}$ there is some k such that for all $m > k$, $\vec{F} \restriction \omega \cdot (n-1) \in z_m \in \text{dom } F_{\omega \cdot (n-1)+m}$, and $\langle F_{\omega \cdot (n-1)+m}(z_m) \rangle \cap \pi_{m+1}(\vec{F} \restriction \omega \cdot (n-1)) \in G_{m+1}$. Let us show that for $\alpha_0 \leq \alpha < \omega^n$, if $\vec{F} \in N_\alpha$ is a sequence of functions such that $\langle \emptyset \rangle \cap \vec{F}$ is a condition in $j_{\alpha_0, \alpha}(\mathbb{P}(\mu, d))$, then for every $\beta > \alpha$,

$$\begin{aligned} j_{\alpha, \omega^n}(\vec{F} \restriction \omega \cdot n_*(\beta)) &\in y_\beta \in \text{dom } j_{\alpha, \omega^n}(F_{\omega \cdot n_*(\beta)+m_*(\beta)}), \\ \text{and } j_{\alpha, \omega^n}(F_{\omega \cdot n_*(\beta)+m_*(\beta)})(y_\beta) &\in C_\beta. \end{aligned}$$

The relation $j_{\alpha, \omega^n}(\vec{F} \restriction \omega \cdot n_*(\beta)) \in y_\beta$ holds simply because $\vec{F} \restriction \omega \cdot n_*(\beta) \in (H_{j_{\alpha_0, \alpha}(\kappa)^{+\omega \cdot n_*(\beta)+1}})^{N_\alpha}$. The other claims are true since $\bar{y}_\beta := j_{\beta+1, \omega^n}^{-1}(y_\beta)$ is the seed of the measure $j_{\alpha_0, \beta}(U_{\omega \cdot n_*(\beta)+m_*(\beta)})$ and the domain of $j_{\alpha, \beta}(F_{\omega \cdot n_*(\beta)+m_*(\beta)})$ is large with respect to this measure. Moreover, this function represents an element

of $j_{\alpha_0, \beta}(K_{\omega \cdot n_*(\beta) + m_*(\beta)})$. But

$$\begin{aligned} j_{\alpha, \beta+1}(F_{\omega \cdot n_*(\beta) + m_*(\beta)})(\bar{y}_\beta) &= \\ j_{\beta, \beta+1}(j_{\alpha, \beta}(F_{\omega \cdot n_*(\beta) + m_*(\beta)})(\bar{y}_\beta)) &= \\ [j_{\alpha, \beta}(F_{\omega \cdot n_*(\beta) + m_*(\beta)})]_{j_{\alpha_0, \beta}(U_{\omega \cdot n_*(\beta) + m_*(\beta)})} &\in j_{\alpha_0, \beta}(K_{\omega \cdot n_*(\beta) + m_*(\beta)}) = C_\beta. \end{aligned}$$

Note that for $\omega^{n-1} \cdot l < \alpha < \omega^n$, the sequence $\langle y_\alpha, y_{\alpha + \omega^{n_*(\alpha)+1}}, \dots, y_{\alpha + \omega^{n-1}} \rangle$ is both \in - and \subseteq -increasing. Thus to compute x_α , we get the same result by taking the image of y_α under the transitive collapse $y_{\alpha + \omega^{n_*(\alpha)+1}}$, as by first collapsing $y_{\alpha + \omega^{n-1}}$, then collapsing the image of $y_{\alpha + \omega^{n-2}}$, etc., until we take the image of y_α under $n - n_*(\alpha) - 1$ successive collapses. The point is that the latter process parallels exactly the sequence of collapses applied to a sequence of functions \vec{F} to determine whether $\langle \emptyset \rangle \cap \vec{F}$ is in the filter generated from the sequences $\langle x_\alpha, C_\beta : 1 \leq \alpha < \omega^n, \beta < \omega^n \rangle$.

Hence, if

$$\begin{aligned} j_{\alpha, \omega^n}(\vec{F} \upharpoonright \omega \cdot n_*(\beta)) &\in y_\beta \in \text{dom } j_{\alpha, \omega^n}(F_{\omega \cdot n_*(\beta) + m_*(\beta)}), \\ \text{and } j_{\alpha, \omega^n}(F_{\omega \cdot n_*(\beta) + m_*(\beta)})(y_\beta) &\in C_\beta, \end{aligned}$$

then $j_{\alpha, \omega^n}(\vec{F})'_\beta \in x_\beta \in \text{dom } j_{\alpha, \omega^n}(\vec{F})'_\beta$, and $j_{\alpha, \omega^n}(\vec{F})'_\beta(x_\beta) \in C_\beta$. So if $\vec{F} \in N_\alpha$, the genericity criteria holds for $j_{\alpha, \omega^n}(\vec{F})$ for the cofinal segment above α . Since N_{ω^n} is a direct limit, the generated filter G is generic.

We would like to claim now that $N_{\omega^n}[G]$ has the same κ -sequences as $W[H]$. Indeed, since the forcing that introduces H has cardinality κ , any sequence of ordinals in $W[H]$ has a name of cardinality κ and thus can be coded using a sequence of ordinals of length κ from W .

Let $\langle \xi_i \mid i < \kappa \rangle$ be a sequence of ordinals in W . In N_{ω^n} , for every ordinal there is a representing function f_i , and a finite sequence $s_i \subseteq \langle y_\alpha : \omega \cdot l < \alpha < \omega^n \rangle$, such that $j_{\alpha_0, \omega^n}(f_i)(s_i) = \xi_i$. By our choices of x_i and C_i , the sequence $\langle y_\alpha : \omega \cdot l < \alpha < \omega^n \rangle$ can be computed from the generic filter G . Since $j_{\alpha_0, \omega^n}(\langle f_i \mid i < \kappa \rangle)$ and $j_{\alpha_0, \omega^n}(\langle s_i \mid i < \kappa \rangle)$ are in N_{ω^n} , and since $\langle y_\alpha \mid \omega^{n-1} \cdot l < \alpha < \omega^n \rangle \in N_{\omega^n}[G]$ we conclude that $\langle \xi_i \mid i < \kappa \rangle \in N_{\omega^n}[G]$. \square

Let us return to the proof of the theorem. Recall that, assuming the inductive claim holds for GS_n , we must only show that for every $\mathbb{P}(\mu, d) \in \text{GS}_{n+1}$ with $\text{crit}(d) = \kappa$, it is forced that $(\kappa^+, \kappa) \rightarrow (\kappa_i^+, \kappa_i)$ holds for infinitely many i . Let p be a condition of length l , let $H \subseteq \mathbb{R} \times \text{Col}(\nu, \kappa)$ be as in Claim 30, with H generic over V . Note that $V[H] \models |(\kappa_l^{+\omega \cdot n})^V| = \kappa_l$, and $(\kappa_l^{+\omega \cdot n+1})^V = \kappa_l^+$. By Lemma 26, $(\kappa^{+\omega \cdot (n+1)+1}, \kappa^{+\omega \cdot (n+1)}) \rightarrow (\kappa_l^+, \kappa_l)$ holds in $V[H]$. Let $j : V \rightarrow M$ and G be given by Claim 30, with $j(p) \in G$.

Let $\mathfrak{A} \in M[G]$ be any structure on $j(\kappa^{+\omega \cdot (n+1)+1}) = \kappa^{+\omega \cdot (n+1)+1} = (j(\kappa)^+)^{M[G]}$. By Chang's Conjecture in $V[H]$, there is a $\mathfrak{B} \prec \mathfrak{A}$ of size $\kappa_l^{+\omega \cdot n+1}$ such that $|\mathfrak{B} \cap \kappa^{+\omega \cdot (n+1)}| = |\mathfrak{B} \cap j(\kappa)| = \kappa_l^{+\omega \cdot n}$. By the closure of $M[G]$, $\mathfrak{B} \in M[G]$, and thus $M[G] \models (j(\kappa)^+, j(\kappa)) \rightarrow (\kappa_l^+, \kappa_l)$. By elementarity, the desired conclusion follows.

5. CHANG'S CONJECTURE WITH THE SAME TARGET

In this section we will discuss two restricted versions of the Singular Global Chang's Conjecture.

Theorem 31. *Suppose that κ is ν^+ -supercompact, where $\text{cf } \nu = \kappa^+$ and ν is a limit of measurable cardinals, and α_* is a countable ordinal. Then there is a generic extension in which*

$$(\mu^+, \mu) \rightarrow (\omega_1, \omega),$$

for all $\mu < \aleph_{\alpha_*}$.

Theorem 32. *Suppose there are two supercompact cardinals and $\alpha_\star > 0$ is a countable limit ordinal. Then there is a generic extension in which*

$$(\mu^+, \mu) \twoheadrightarrow (\aleph_{\alpha_\star+1}, \aleph_{\alpha_\star}),$$

for all singular μ , $\aleph_{\alpha_\star} < \mu < \aleph_{\omega_1}$.

Proof. The proof of both theorems follows closely the ideas from [12], which in turn are motivated by the forcing arguments from [15].

For Theorem 31, let us assume that κ is Laver-indestructible (with respect to κ -directed closed forcing notions of cardinality $\leq \nu^+$) and that GCH holds above κ . If this is not the case, we can always force it using Laver forcing [14]. Let $\langle \zeta_\beta \mid \beta < \kappa^+ \rangle$ be a continuous increasing sequence with $\sup \zeta_\beta = \nu$, $\zeta_0 = \kappa$, and $\zeta_{\beta+1}$ measurable for each $\beta < \kappa^+$.

For every $\alpha < \kappa^+$ of countable cofinality, let us pick a cofinal ω -sequence of ordinals $s_\alpha: \omega \rightarrow \alpha$. Let us assume that for each α , s_α is increasing, $s_\alpha(0) = 0$, and $s(n)$ is a successor ordinal for $n > 0$.

Let us consider the forcing

$$\mathbb{C}_\alpha = \prod_{n < \omega} \mathbb{E}(\zeta_{s_\alpha(n)}, \zeta_{s_\alpha(n+1)}) \times \text{Col}(\zeta_\alpha^+, \nu^+),$$

where $\mathbb{E}(\mu, \delta)$ is the Easton-support product of $\text{Col}(\mu, \eta)$ over all $\eta < \delta$. The product in the definition of \mathbb{C}_α is taken with full support.

For each $\alpha < \kappa^+$ of countable cofinality, after forcing with \mathbb{C}_α ,

$$(\zeta_\alpha^+)^V = (\kappa^{+\omega+1})^{V^{\mathbb{C}_\alpha}}.$$

By the arguments of [5] related to Lemma 13, there is $\rho_\alpha < \kappa$ such that

$$V^{\mathbb{C}_\alpha} \models (\kappa^{+\omega+1}, \kappa^{+\omega}) \twoheadrightarrow (\rho_\alpha^+, \rho_\alpha),$$

and remains true after forcing with $\mathbb{D}_\alpha = \text{Col}(\omega, \rho_\alpha) * \dot{\text{Col}}(\rho_\alpha^+, < \kappa)$. In fact, $(\zeta_\alpha^+, \zeta_\alpha) \twoheadrightarrow (\rho_\alpha^+, \rho_\alpha)$ must already hold in V , by the distributivity of \mathbb{C}_α .

Since the forcing \mathbb{C}_α is weakly homogeneous, the value of ρ_α depends only on α and does not depend on the generic filter for \mathbb{C}_α . Therefore, the function $\alpha \rightarrow \rho_\alpha$ is in V , and it has the property that

$$1 \Vdash_{\mathbb{D}_\alpha \times \mathbb{C}_\alpha} (\kappa^{+\omega+1}, \kappa^{+\omega}) \twoheadrightarrow (\check{\rho}_\alpha^+, \check{\rho}_\alpha).$$

By the κ^+ -completeness of NS_{κ^+} , there is a stationary set $S \subseteq \kappa^+$ and a cardinal $\rho_\star < \kappa$ such that for all $\alpha \in S$, $\rho_\alpha = \rho_\star$. Let \mathbb{D} be the common value of \mathbb{D}_α for $\alpha \in S$. There is $n_0 < \omega$ such that for every club $C \subseteq \kappa^+$, $\{s_\alpha(n_0) : \alpha \in C \cap S\}$ is unbounded. By Fodor's Lemma, we may assume $s_\alpha \restriction n_0$ is constant on S .

Let us define a partial order \mathbb{P} that searches for a “thread” of the sequences s_α for $\alpha \in S$. A condition $t \in \mathbb{P}$ is a continuous increasing function from a countable successor ordinal γ into κ^+ , such that $\text{ran } t \subseteq S \cup \bigcup_{\alpha < \kappa^+} \text{ran } s_\alpha$, and for every limit ordinal $\beta < \gamma$, $\text{ran } s_{t(\beta)} \subseteq \text{ran } t$. As in [12], we have:

Claim 33. *For every $t \in \mathbb{P}$, every $\gamma < \omega_1$, and every $\xi < \kappa^+$, there is a stronger condition $t' \supseteq t$ with $\gamma \subseteq \text{dom } t'$ and $s_\beta(n_0) > \xi$ for limit $\beta \in \text{dom } t' \setminus \text{dom } t$.*

In particular, we can find a thread of any countable length. Let t be a thread of length α_\star . Define a sequence $s: \alpha_\star \rightarrow \nu$ as follows. If β is an infinite limit ordinal, then $s(\beta) = \zeta_{t(\beta)}^+$, and otherwise $s(\beta) = \zeta_{t(\beta)}$. Consider the forcing:

$$\mathbb{C} = \prod_{\beta < \alpha_\star} \mathbb{E}(s(\beta), s(\beta+1)).$$

First let us claim that in the generic extension by $\mathbb{D} \times \mathbb{C}$, we have $(\aleph_{\beta+1}, \aleph_\beta) \twoheadrightarrow (\aleph_1, \aleph_0)$ for limit $\beta < \alpha_\star$. As in [12], the projection properties of the Levy collapse,

together with the fact that $\text{ran } s_\beta \subseteq \text{ran } t$ for limit $\beta < \alpha_*$, imply that for each limit $\beta < \alpha_*$, there is a projection $\pi_\beta : \mathbb{C}_\beta \rightarrow \mathbb{C}$. If \mathfrak{A} is a structure on ζ_β^+ in $V^{\mathbb{D} \times \mathbb{C}}$, then in $V^{\mathbb{D} \times \mathbb{C}_\beta}$, there is an elementary $\mathfrak{B} \prec \mathfrak{A}$ such that $|\mathfrak{B}| = \rho_\star^+ = \aleph_1$, and $|\mathfrak{B} \cap \zeta_\beta| = |\rho_\star| = \aleph_0$. Since the quotient forcing adds no sets of ordinals of size $< \kappa = \aleph_2$, the instance of Chang's Conjecture holds in $V^{\mathbb{D} \times \mathbb{C}}$.

To obtain the result for successors below α_* , we consider instead the forcing $\mathbb{D} * \dot{\mathbb{C}}$, where $\dot{\mathbb{C}}$ is the forcing with the same definition as \mathbb{C} , but constructed in $V^{\mathbb{D}}$ rather than V . By [21], there is a projection from $\mathbb{D} \times \mathbb{C}$ to $\mathbb{D} * \dot{\mathbb{C}}$ that is the identity on \mathbb{D} . By the same argument as above, the relevant instances of Chang's Conjecture at limit ordinals also hold in $V^{\mathbb{D} * \dot{\mathbb{C}}}$.

Suppose $\beta < \alpha_*$ is zero or a successor ordinal. For convenience, let $\zeta_{-1} = \rho_\star^+$. Since ζ_β is measurable, in the extension by

$$\mathbb{D} * \prod_{\gamma < \beta} \dot{\mathbb{E}}(s(\gamma), s(\gamma + 1)),$$

there is a normal ideal I on ζ_β such that $\mathcal{P}(\zeta_\beta)/I$ contains a countably closed dense set—in particular the boolean algebra is a proper forcing. By [18], the following version of Strong Chang's Conjecture holds in this model: If M is a countable elementary submodel of $H_{\zeta_\beta^+}$ then there is an elementary $M' \supseteq M$ such that $M \cap \zeta_{\beta-1} = M' \cap \zeta_{\beta-1}$ and $M \cap \zeta_\beta \neq M' \cap \zeta_\beta$. By Lemma 15 of [5], $(\zeta_\beta, \zeta_{\beta-1}) \twoheadrightarrow (\aleph_1, \aleph_0)$ is preserved by the formerly ζ_β -closed quotient $\prod_{\beta \leq \gamma < \alpha_*} \dot{\mathbb{E}}(s(\gamma), s(\gamma + 1))$.

Finally, let us show Theorem 32. Let $\kappa_0 < \kappa$ be supercompact, and let $\alpha_* > 0$ be a fixed countable limit ordinal. First force Martin's Maximum (MM) while turing κ_0 into \aleph_2 [6]. MM is indestructible under \aleph_2 -directed-closed forcing [13]. Then force with Laver's forcing, which is \aleph_2 -directed-closed, to force that κ is indestructibly supercompact and GCH holds above κ .

Next we need, for large enough $\mu < \kappa$, a forcing \mathbb{D}_μ that turns κ into \aleph_{α_*+3} while preserving ω_1 and satisfying the hypotheses of Lemma 13. If $\tau(\alpha_*) = \omega$, let $\mathbb{D}_\mu = \text{Col}(\aleph_{\iota(\alpha_*)+1}, \mu) \times \text{Col}(\mu^{+\omega+2}, < \kappa)$. If $\tau(\alpha_*) > \omega$, let $\vec{\gamma}$ be the identity sequence converging to ω , and let $\vec{\delta}$ be a nondecreasing sequence summing to $\tau(\alpha_*)$, with $\delta_1 \geq \omega$. Let $\mathbb{D}_\mu = \mathbb{P}(\aleph_{\iota(\alpha_*)+1}, \vec{\gamma}, \vec{\delta}, \vec{U}, \vec{K}) \times \text{Col}(\mu^{+\omega+2}, < \kappa)$, where \vec{U} and \vec{K} are ω -sequences such that U_n is a normal μ -complete ultrafilter on $\mathcal{P}_\mu(\mu^{+n})$, and K_n is sufficiently generic filter, as in Section 3.

For each $\alpha < \kappa^+$ of countable cofinality, choose a cofinal increasing sequence $s_\alpha : \omega \rightarrow \alpha$ with $s_\alpha(0) = \kappa$ and $s_\alpha(n)$ is a double successor ordinal for $n > 0$. For each $\alpha < \kappa^+$ of countable cofinality, define

$$\mathbb{C}_\alpha = \prod_{n < \omega} \text{Col}(\kappa^{+s_\alpha(n)}, \kappa^{+s_\alpha(n+1)-1}) \times \text{Col}(\kappa^{+\alpha+2}, \kappa^{+\kappa^++1}).$$

For each α , there is $\mu_\alpha < \kappa$ such that

$$1 \Vdash_{\mathbb{D}_{\mu_\alpha} \times \mathbb{C}_\alpha} (\kappa^{+\omega+1}, \kappa^{+\omega}) \twoheadrightarrow (\aleph_{\alpha_*+1}, \aleph_{\alpha_*}).$$

As above, let $S \subseteq \kappa^+$ be a stationary set of countable cofinality ordinals such that μ_α has the same value for all $\alpha \in S$, and that the threading forcing \mathbb{P} satisfies Claim 33. Let $\mathbb{D} = \mathbb{D}_{\mu_\alpha}$ for any $\alpha \in S$. We now claim that \mathbb{P} preserves stationary subsets of ω_1 . To see this, fix some stationary set $A \subseteq \omega_1$ and some condition $t_0 \in \mathbb{P}$ let \dot{C} be a \mathbb{P} -name for a club subset of ω_1 , and let

$$M \prec (H_{\kappa^{++}}, \in, \langle s_\alpha : \alpha < \kappa^+ \rangle, S, \mathbb{P}, p_0, A, \dot{C})$$

be such that $M \cap \kappa^+ = \delta \in S$. Let $N \prec M$ be countable with $\sup(N \cap \kappa^+) = \delta$, $\text{ran } s_\delta \subseteq N$, and $N \cap \omega_1 \in A$. Let $\langle D_n : n < \omega \rangle$ enumerate the dense subsets of \mathbb{P} in N . Using Claim 33, we can build a sequence $t_0 \geq t_1 \geq t_2 \geq \dots$ such that for

$n > 0$, $t_n \in D_n$ and $s_\delta(n) \in \text{ran } t_n$. If $t = \bigcup_n t_n \cup \{\delta\}$, then t is an (N, \mathbb{P}) -master condition, and so it forces $A \cap \dot{C} \neq \emptyset$.

Applying MM, we find a thread t of length ω_1 . Let $s : \omega_1 \rightarrow \kappa^+$ be such that $s(\alpha) = t(\alpha) + 2$ for limit $\alpha > 0$ and $s(\alpha) = t(\alpha)$ otherwise. Let us consider the forcing

$$\mathbb{C} = \prod_{\alpha < \omega_1} \text{Col}(\kappa^{+s(\alpha)}, \kappa^{+s(\alpha+1)-1}).$$

For every $\beta \in S$, there is a projection from \mathbb{C}_β to \mathbb{C} . Therefore, since the quotient adds no sets of ordinals of size $< \kappa$, $\mathbb{D} \times \mathbb{C}$ forces the desired conclusion. \square

6. OPEN PROBLEMS

The construction in Section 4 is limited to instances of Chang's Conjecture between successors of singular cardinals below \aleph_{ω^ω} . In order to push this mechanism forwards, one needs to start with a model in which there is a cardinal κ which is $\kappa^{+\alpha+1}$ -supercompact and Chang's Conjecture holds between any pair of singular cardinals in the interval $[\kappa, \kappa^{+\alpha}]$. Since our method to produce an interval with such properties with limits of limit cardinals includes Prikry forcing, it cannot preserve supercompactness.

Question 1. *Is it consistent relative to large cardinals that $(\mu^+, \mu) \rightarrow (\nu^+, \nu)$ holds whenever μ and ν have countable cofinality?*

The known limitations on Global Chang's Conjecture do not seem to rule out the consistency of a strengthening of Theorem 31 to a global statement:

Question 2. *Is it consistent relative to large cardinals that $(\kappa^+, \kappa) \rightarrow (\omega_1, \omega)$ holds for all infinite cardinals κ ?*

REFERENCES

- [1] Uri Abraham and Menachem Magidor, *Cardinal arithmetic*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 1149–1227. MR 2768693
- [2] Lev Bukovský, *Iterated ultrapower and Prikry's forcing*, Comment. Math. Univ. Carolinae **18** (1977), no. 1, 77–85. MR 0446978
- [3] James Cummings, Matthew Foreman, and Menachem Magidor, *Squares, scales and stationary reflection*, J. Math. Log. **1** (2001), no. 1, 35–98. MR 1838355 (2003a:03068)
- [4] Patrick Dehornoy, *Iterated ultrapowers and Prikry forcing*, Ann. Math. Logic **15** (1978), no. 2, 109–160. MR 514228
- [5] Monroe Eskew and Yair Hayut, *On the consistency of local and global versions of chang's conjecture*, Transactions of the American Mathematical Society **370** (2018), no. 4, 2879–2905.
- [6] M. Foreman, M. Magidor, and S. Shelah, *Martin's maximum, saturated ideals, and nonregular ultrafilters. I*, Ann. of Math. (2) **127** (1988), no. 1, 1–47. MR 924672
- [7] Matthew Foreman, *Smoke and mirrors: combinatorial properties of small cardinals equiconsistent with huge cardinals*, Adv. Math. **222** (2009), no. 2, 565–595. MR 2538021
- [8] Matthew Foreman and Menachem Magidor, *Large cardinals and definable counterexamples to the continuum hypothesis*, Annals of Pure and Applied Logic **76** (1995), no. 1, 47–97.
- [9] Matthew Foreman and Menachem Magidor, *A very weak square principle*, J. Symb. Log. **62** (1997), no. 1, 175–196.
- [10] Moti Gitik, *Prikry-type forcings*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 1351–1447. MR 2768695
- [11] Moti Gitik and Assaf Sharon, *On SCH and the approachability property*, Proc. Amer. Math. Soc. **136** (2008), no. 1, 311–320. MR 2350418
- [12] Mohammad Golshani and Yair Hayut, *The tree property on a countable segment of successors of singular cardinals*, Fund. Math. **240** (2018), no. 2, 199–204. MR 3720924
- [13] Paul Larson, *Separating stationary reflection principles*, J. Symbolic Logic **65** (2000), no. 1, 247–258. MR 1782117
- [14] Richard Laver, *Making the supercompactness of κ indestructible under κ -directed closed forcing*, Israel J. Math. **29** (1978), no. 4, 385–388. MR 0472529
- [15] Menachem Magidor and Saharon Shelah, *When does almost free imply free? (For groups, transversals, etc.)*, J. Amer. Math. Soc. **7** (1994), no. 4, 769–830. MR 1249391

- [16] William J. Mitchell, *Beginning inner model theory*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 1449–1495. MR 2768696
- [17] Norman Lewis Perlmutter, *The large cardinals between supercompact and almost-huge*, Arch. Math. Logic **54** (2015), no. 3-4, 257–289. MR 3324126
- [18] Hiroshi Sakai, *Semiproper ideals*, Fund. Math. **186** (2005), no. 3, 251–267. MR 2191239
- [19] Saharon Shelah, *Reflecting stationary sets and successors of singular cardinals*, Arch. Math. Logic **31** (1991), no. 1, 25–53.
- [20] ———, *Proper and improper forcing*, second ed., Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1998. MR 1623206
- [21] Masahiro Shioya, *The Easton collapse and a saturated filter*, RIMS Kokyuroku **1754** (2011), 108–114.
- [22] John Steel, *An introduction to iterated ultrapowers*, Forcing, iterated ultrapowers, and Turing degrees (Chitat Chong et al., eds.), Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap., vol. 29, World Sci. Publ., Hackensack, NJ, 2016, pp. 123–174.

(Monroe Eskew) KURT GÖDEL RESEARCH CENTER, UNIVERSITY OF VIENNA, 25 WÄHRINGER STRASSE, 1090 VIENNA, AUSTRIA

E-mail address: `monroe.eskew@univie.ac.at`

(Yair Hayut) DEPARTMENT OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM 9190401, ISRAEL

E-mail address: `yair.hayut@mail.huji.ac.il`