KIRILLOV–RESHETIKHIN CRYSTALS $B^{7,s}$ FOR TYPE $E_7^{(1)}$

REKHA BISWAL AND TRAVIS SCRIMSHAW

ABSTRACT. We construct a combinatorial crystal structure on the Kirillov–Reshetikhin crystal $B^{7,s}$ in type $E_7^{(1)}$, where 7 is the unique node in the orbit of 0 in the affine Dynkin diagram. We then describe the combinatorial *R*-matrix $R: B^{7,s} \otimes B^{7,s'} \to B^{7,s'} \otimes B^{7,s}$.

1. INTRODUCTION

An important class of finite-dimensional representations for affine Lie algebras are the *Kirillov-Reshetikhin (KR) modules*, which are characterized by their Drinfel'd polynomials [4, 5]. We denote a KR module by $W^{r,s}$, where r is a node of the classical Dynkin diagram and s is a positive integer. KR modules have been well-studied and have many interesting properties. For example, their characters (resp. *q*-characters) are solutions of the Q-system (resp. T-system) [13] (see also [21] and references therein). Moreover, graded (Demazure-type) characters of tensor products of single-column KR modules are (nonsymmetric) Macdonald polynomials at t = 0 for untwisted affine types [23–25].

One significant aspect of a KR module $W^{r,s}$ is that it (conjecturally) admits a crystal base [11, 12] despite not being a highest weight module. The corresponding crystal of $W^{r,s}$ is called a *Kirillov–Reshetikhin (KR) crystal* and denoted by $B^{r,s}$. KR crystals have been shown to exist in all nonexceptional types in [33], types $G_2^{(1)}$ and $D_4^{(3)}$ in [31], for a number of nodes in exceptional types [1, 32], and for r being in the orbit of or adjacent to 0 in all affine types from the general theory [16, 17]. An open problem is to determine a uniform model for KR crystals. This has been achieved for $B^{r,1}$ by using Kashiwara's construction of projecting an extremal level-zero module/crystal [20]. This was done explicitly by Naito and Sagaki using Lakshmibai–Seshadri (LS) paths [28–30]. The construction of Kashiwara was also shown to partially extend to general $B^{r,s}$ in nonexceptional affine types (conjecturally in all affine types) [26]. In contrast, the models in [7, 15, 18, 37] are all type-dependent, but are given for $B^{r,s}$ for all s.

KR crystals are connected with mathematical physics. For instance, tensor products of KR modules are used to describe certain vertex models and are related with Heisenberg spin chains by the X = M conjecture of [11, 12]. The X = M conjecture implies a fermionic formula for the graded characters of a tensor product of KR crystals; see [34, 36] for recent progress. Furthermore, tensor products of KR crystals describe the dynamics of soliton cellular automata, a generalization of the Takehashi–Satsuma box-ball system (which is an ultradiscrete version of the Korteweg–de Vries (KdV) equation). We refer the reader to [14, 27] for more details. Another important (conjectural) property of KR crystals is that

²⁰¹⁰ Mathematics Subject Classification. 05E10, 17B37.

Key words and phrases. Kirillov–Reshetikhin crystal, crystal, crystal basis, affine Lie algebra.

R.B. was partially supported by the NSERC discovery grant of her Postdoc supervisor Michael Lau at Université Laval. T.S. was partially supported by the Australian Research Council grant DP170102648.

they are perfect [8, 16-18, 37], a technical condition that allows highest weight crystals to be modeled using a semi-infinite tensor product known as the Kyoto path model [17].

In this note, we give a combinatorial model for the KR crystal $B^{7,s}$ in type $E_7^{(1)}$, where 7 is the unique node in the orbit of 0 in the Dynkin diagram (see Figure 1 below). We achieve this by considering the (Levi) decomposition of the classical (type E_7) highest weight crystal $B(s\overline{\omega}_7)$ into A_6 highest weight crystals, which is multiplicity free. From this, we reconstruct the A_7 decomposition of $B^{7,s}$ since $B^{7,s} \cong B(s\overline{\omega}_7)$ as E_7 crystals and the decomposition of A_7 highest weight crystals into A_6 crystals is multiplicity free. We note that the KR crystal $B^{7,s}$ exists since 7 is a minuscule node, so $B^{7,s}$ is irreducible as a classical crystal. The novelty of our approach is doing a further Levi decomposition and reconstructing the affine action to a type A_7 crystal rather than through the classical decomposition. We then given an explicit description of the combinatorial R-matrix $R: B^{7,s} \otimes B^{7,s'} \to B^{7,s'} \otimes B^{7,s}$. We note that the local energy function is given by [36, Thm. 7.5].

As a potential application of our results, the combinatorial *R*-matrix allows us to study soliton cellular automata of $B^{7,1}$ using different techniques from [26]. Moreover, our results could potentially be used to show that $B^{7,s}$ is a perfect crystal of level *s*.

This paper is organized as follows. In Section 2, we give the necessary background. In Section 3, we give our main results. In Section 4, we give a conjecture about the decomposition of $B^{1,s}$ into A_7 crystals in an effort to prove [15, Conj. 3.26].

Acknowledgments. The authors thank the referee for useful comments on our manuscript.

2. Background

Let \mathfrak{g} be an affine Kac–Moody Lie algebra with index set I, Cartan matrix $(A_{ij})_{i,j\in I}$, simple roots $(\alpha_i)_{i\in I}$, fundamental weights $(\omega_i)_{i\in I}$, and simple coroots $(\alpha_i^{\vee})_{i\in I}$. Let $U_q(\mathfrak{g})$ denote the corresponding (Drinfel'd–Jimbo) quantum group, and we will be using $U'_q(\mathfrak{g}) := U_q([\mathfrak{g},\mathfrak{g}])$, which has weight lattice $P = \sum_{i\in I} \mathbb{Z}\omega_i$. Let P^+ denote the positive weight lattice. Let Qbe the root lattice with Q^+ being the positive root lattice. We denote the canonical pairing $\langle , \rangle \colon P^{\vee} \times P \to \mathbb{Z}$, which is given by $\langle \alpha_i^{\vee}, \alpha_j \rangle = A_{ij}$.

Recall that $lev(\lambda) := \langle c, \lambda \rangle$ is the level of the weight λ , where c is the canonical central element of \mathfrak{g} . In particular, for \mathfrak{g} of type $E_7^{(1)}$, we have

$$\begin{aligned}
 & \text{lev}(\omega_0) = 1, & \text{lev}(\omega_1) = 2, & \text{lev}(\omega_2) = 2, & \text{lev}(\omega_3) = 3, \\
 & \text{lev}(\omega_4) = 4, & \text{lev}(\omega_5) = 3, & \text{lev}(\omega_6) = 2, & \text{lev}(\omega_7) = 1.
 \end{aligned}$$
(2.1)

We denote the dominant weights of level ℓ by P_{ℓ}^+ .

Let \mathfrak{g}_0 denote the canonical simple Lie algebra given by the index set $I_0 = I \setminus \{0\}$, and $U_q(\mathfrak{g}_0)$ the corresponding quantum group. Let P_0 and Q_0 be the weight and root lattice of \mathfrak{g}_0 , and let $\overline{\omega}_i$ be the natural projection of the fundamental weight ω_i onto P_0 . Let W_0 be the Weyl group of \mathfrak{g}_0 .

2.1. Crystals. An *abstract* $U_q(\mathfrak{g})$ -*crystal* is a set B endowed with *crystal operators* $e_i, f_i: B \to B \sqcup \{0\}$, for $i \in I$, and *weight function* wt: $B \to P$ that satisfy the following conditions:

- (1) $\varphi_i(b) = \varepsilon_i(b) + \langle \alpha_i^{\vee}, \operatorname{wt}(b) \rangle$, for all $b \in B$ and $i \in I$,
- (2) $f_i b = b'$ if and only if $b = e_i b'$, for $b, b' \in B$ and $i \in I$,
- (3) wt($f_i b$) = wt(b) α_i if $f_i b \neq 0$;




FIGURE 1. The Dynkin diagram of type $E_7^{(1)}$.

where the statistics $\varepsilon_i, \varphi_i \colon B \to \mathbb{Z}_{\geq 0}$ are defined by

$$\varepsilon_i(b) := \max\{k \mid e_i^k b \neq 0\}, \qquad \qquad \varphi_i(b) := \max\{k \mid f_i^k b \neq 0\}.$$

Remark 2.1. The definition of an abstract crystal given in this paper is sometimes called a *regular* or *seminormal* abstract crystal in the literature. See, *e.g.*, [2] for the more general definition.

Using the axioms, we identify B with an I-edge colored weighted directed graph whose vertices are B and having an *i*-colored edge $b \xrightarrow{i} b'$ if and only if $f_i b = b'$. Therefore, we can depict an entire *i*-string through an element $b \in B$ diagrammatically by

$$e_i^{\varepsilon_i(b)}b \xrightarrow{i} \cdots \xrightarrow{i} e_i^2 b \xrightarrow{i} e_i b \xrightarrow{i} b \xrightarrow{i} f_i b \xrightarrow{i} f_i^2 b \xrightarrow{i} \cdots \xrightarrow{i} f_i^{\varphi_i(b)} b.$$

Let $J \subseteq I$. An element $b \in B$ is *J*-highest (resp. lowest) weight if $e_i b = 0$ (resp. $f_i b = 0$) for all $i \in J$. When J = I, we simply say b is highest (resp. lowest) weight.

For abstract $U_q(\mathfrak{g})$ -crystals B_1, B_2, \ldots, B_L , the action of the crystal operators on the *tensor* product $B_L \otimes \cdots \otimes B_2 \otimes B_1$, which equals the Cartesian product $B_L \times \cdots \times B_2 \times B_1$ as sets, can be defined by the signature rule. Let $b := b_L \otimes \cdots \otimes b_2 \otimes b_1 \in B$, and for $i \in I$, we write

$$\underbrace{-\cdots}_{\varphi_i(b_L)} \underbrace{+\cdots+}_{\varepsilon_i(b_L)} \cdots \underbrace{-\cdots}_{\varphi_i(b_1)} \underbrace{+\cdots+}_{\varepsilon_i(b_1)}.$$

Then by successively deleting consecutive +--pairs (in that order), we obtain a sequence

$$\operatorname{sgn}_i(b) := \underbrace{-\cdots -}_{\varphi_i(b)} \underbrace{+\cdots +}_{\varepsilon_i(b)},$$

called the *reduced signature*. If there does not exist a + (resp. -) in $\operatorname{sgn}_i(b)$, then $e_i b = 0$ (resp. $f_i b = 0$). Otherwise, suppose $1 \leq j_- \leq j_+ \leq L$ are such that b_{j_-} contributes the rightmost - in $\operatorname{sgn}_i(b)$ and b_{j_+} contributes the leftmost + in $\operatorname{sgn}_i(b)$. Then, we have

$$e_ib := b_L \otimes \cdots \otimes b_{j_++1} \otimes e_i b_{j_+} \otimes b_{j_+-1} \otimes \cdots \otimes b_1,$$

$$f_ib := b_L \otimes \cdots \otimes b_{j_-+1} \otimes f_i b_{j_-} \otimes b_{j_--1} \otimes \cdots \otimes b_1.$$

Remark 2.2. Our tensor product convention follows [2], which is opposite to that of Kashiwara [19].

Let B_1 and B_2 be two abstract $U_q(\mathfrak{g})$ -crystals. A crystal morphism $\psi: B_1 \to B_2$ is a map $B_1 \sqcup \{0\} \to B_2 \sqcup \{0\}$ with $\psi(0) = 0$, such that the following properties hold for all $b \in B_1$ and $i \in I$:

- (1) if $\psi(b) \in B_2$, then wt $(\psi(b)) = wt(b)$, $\varepsilon_i(\psi(b)) = \varepsilon_i(b)$, and $\varphi_i(\psi(b)) = \varphi_i(b)$;
- (2) we have $\psi(e_i b) = e_i \psi(b)$ if $\psi(e_i b) \neq 0$ and $e_i \psi(b) \neq 0$;

(3) we have $\psi(f_i b) = f_i \psi(b)$ if $\psi(f_i b) \neq 0$ and $f_i \psi(b) \neq 0$.

An *embedding* (resp. *isomorphism*) is a crystal morphism such that the induced map $B_1 \sqcup \{0\} \to B_2 \sqcup \{0\}$ is an embedding (resp. bijection).

An abstract crystal B is a $U_q(\mathfrak{g})$ -crystal if B is the crystal basis of some $U_q(\mathfrak{g})$ -module. Kashiwara [19] has shown that the irreducible highest weight module $V(\lambda)$, for $\lambda \in P^+$, admits a crystal basis denoted $B(\lambda)$. The highest weight crystal $B(\lambda)$ is generated by a unique highest weight element u_{λ} that satisfies $\operatorname{wt}(u_{\lambda}) = \lambda$.

2.2. Minuscule crystals. We say a highest weight $U_q(\mathfrak{g}_0)$ -crystal $B(\lambda)$ is minuscule if W_0 acts transitively on $B(\lambda)$. In other words, there exists a bijection between $B(\lambda)$ and W_0^{λ} , the set of minimal length coset representatives of $W_0/\operatorname{stab}_{W_0}(\lambda)$ (recall $\operatorname{stab}_{W_0}(\lambda) := \{w \in W_0 \mid w\lambda = \lambda\}$ is the stabilizer of λ and a parabolic subgroup of W_0). Indeed, consider a minimal length coset representative $w \in W_0^{\lambda}$ with a reduced expression $s_{i_1}s_{i_2}\cdots s_{i_k}$, then the corresponding element is $u_{w\lambda} := f_{i_1}f_{i_2}\cdots f_{i_k}u_{\lambda}$. We note that the element $u_{w\lambda}$ is independent of the choice of reduced expression.

For a minuscule representation $B(\lambda)$, we can characterize the elements in $B(s\lambda)$ as follows. Recall that since $B(\lambda)$ is a highest weight crystal, it can be considered as the Hasse diagram of a poset with u_{λ} being the smallest element.

Proposition 2.3 ([36, Prop. 7.29]). Let $B(\lambda)$ be a minuscule representation. The crystal $B(s\lambda)$ is isomorphic to the set of semistandard tableaux whose shape is a single row of length s

$$T = \begin{bmatrix} x_1 & x_2 & \cdots & x_s \end{bmatrix},$$

with $x_1 \leq x_2 \leq \cdots \leq x_s$ in $B(\lambda)$, and the crystal structure is given by considering T as the element $x_1 \otimes x_2 \otimes \cdots \otimes x_s \in B(\lambda)^{\otimes s}$.

2.3. Type A_n crystals. In this section, we consider the Lie algebra of type A_n , which is \mathfrak{sl}_{n+1} . We denote the fundamental weights of type A_n by $\{\eta_i \mid 1 \leq i \leq n\}$. Recall that we have a natural bijection between P^+ and partitions of length at most n by η_i corresponding to a column of height i. Let $B(\eta_1)$ denote the crystal of the vector representation of \mathfrak{sl}_{n+1} :

$$1 \xrightarrow{1} 2 \xrightarrow{2} 3 \xrightarrow{3} \cdots \xrightarrow{n-1} n \xrightarrow{n} n+1$$

Furthermore, the crystal $B(\eta)$ can be described by *semistandard Young tableaux (SSYT)*, written in English convention, whose entries are at most n+1. The crystal structure is given by embedding a SSYT $T \in B(\eta)$ into $B(\eta_1)^{\otimes |\eta|}$ by the reverse Far-Eastern reading word: reading bottom-to-top and left-to-right.

Next, we recall the Levi branching rule $\mathfrak{sl}_{n+1} \searrow \mathfrak{sl}_n$ given at the level of crystals. In terms of the crystal graph, we simply remove all *n*-colored edges. For the SSYT, this amounts to fixing all n + 1's that appear. Since any n + 1 must be the bottom entry of every column and the largest entry in a given row, we obtain the following statement (which is well-known to experts).

Proposition 2.4. As $U_q(\mathfrak{sl}_n)$ -crystals, we have

$$B(\eta) \cong \bigoplus_{\substack{\mu\\4}} B(\mu),$$

where the sum is taken over all μ such that η/μ is a horizontal strip (i.e. a skew partition that does not contain a vertical domino).

Note that the decomposition of Proposition 2.4 is *multiplicity-free*.

We recall that there exists a natural order 2 diagram automorphism σ on type A_n crystals given by $i^{\sigma} = n + 1 - i$. Indeed, we define an automorphism of the weight lattice, which we abuse notation and also denote by $\sigma: P \to P$, by $\eta_i \mapsto \eta_{i^{\sigma}} = \eta_{n+1-i}$; in particular, we note that $\eta \mapsto -w_0\eta$. Next, we define a map also denoted by $\sigma: B(\eta) \mapsto B(-w_0\eta)$ given by

$$\sigma(f_{i_1}\cdots f_{i_k}u_\eta) = f_{i_1^\sigma}\cdots f_{i_k^\sigma}u_{-w_0\eta}$$

We recall from [35] that $\sigma = \lor \circ \ast$, where \lor denotes the contragredient dual map and \ast is the the Lusztig involution (which equals the Schützenberger involution [22]).

In the sequel, we require the A_7 Levi subalgebra \mathfrak{g}_2 of the type $E_7^{(1)}$ affine Lie algebra \mathfrak{g} given by the index set $I_2 := I \setminus \{2\}$. Therefore, the fundamental weights correspond by

$$\eta_1 = \omega_7, \quad \eta_2 = \omega_6, \quad \eta_3 = \omega_5, \quad \eta_4 = \omega_4, \quad \eta_5 = \omega_3, \quad \eta_6 = \omega_1, \quad \eta_7 = \omega_0.$$

We let $I_{0,2} := J \setminus \{0, 2\}$ be index set of the the A_6 Levi subalgebra of \mathfrak{g}_0 (of type E_7).

3. Results

In this section, we give our main results.

We follow [15, Fig. 3] and label an element $b \in B(\overline{\omega}_7)$ as a word in I_0 and $\overline{I}_0 := \{\overline{1}, \ldots, \overline{7}\}$, where for $i \in b$ such that $i \in I_0$ (resp. $i \in \overline{I}_0$), we have $\varphi_i(b) = 1$ (resp. $\varepsilon_i(b) = 1$). See Figure 2 for the crystal $B(\overline{\omega}_7)$ with this labeling convention.


3.1. Multiplicity freeness of $B(s\overline{\omega}_7)$. We first prove that when we decompose the E_7 crystals $B(s\overline{\omega}_7)$ into the Levi subalgebra of type A_6 , we obtain a multiplicity free decomposition.

The $I_{0,2}$ -highest weight elements of $B(s\overline{\omega}_7)$ are single row tableaux of size k whose entries consist of

-

$$x_1 = 7,$$
 $x_2 = \overline{65},$ $x_3 = \overline{423},$ $x_4 = 12,$ $x_5 = \overline{124}$ $x_6 = \overline{26},$ $x_7 = \overline{21}.$

This can be seen by a direct computation using Proposition 2.3, the crystal graph $B(\overline{\omega}_7)$, and the signature rule. Note that in the crystal graph of $B(\overline{\omega}_7)$, we have



and so, nearly all of the elements x_i are comparable except x_4 with $x_{4'}$.

We recall from [15, Def. 3.10] that a *(reduced) composition graph* $G_k(\overline{\omega})$ is essentially the smallest acyclic digraph with loops whose vertices are elements of $B(\overline{\omega})$ such that for any s > 0 and every $(I_0 \setminus \{k\})$ -highest weight element $b_1 \otimes \cdots \otimes b_s \in B(s\overline{\omega}) \subseteq B(\overline{\omega})^{\otimes s}$, the elements (b_1, \ldots, b_s) occurs as a subsequence of a directed path in $G_k(\overline{\omega})$. We remark by reversing the arrows and adding loops to every vertex in (3.1), we obtain the composition graph $G_1(\overline{\omega}_7)$.




FIGURE 2. The crystal $B(\overline{\omega}_7)$ in type E_7 .

Note that for a semistandard tableau $T \in B(s\overline{\omega}_7)$, we have

$$\operatorname{wt}(T) = \sum_{i \in I} (a_i - a_{\overline{\imath}}) \overline{\omega}_i,$$

where a_i (resp. $a_{\overline{i}}$) equals the number if *i*'s (resp. \overline{i} 's) that appear in *T*. Therefore, from (3.1) and the signature rule, we have the following.

Lemma 3.1. Let m_i denote the number of occurrences of x_i in $T \in B(s\overline{\omega}_7)$. Then T is a $I_{0,2}$ -highest weight element if and only if entries in T consist of $\{x_1, x_2, x_3, x_4, x_{4'}, x_5, x_6, x_7\}$ and

$$m_2 \le m_6,$$
 $m_3 \le m_5,$ $m_4 + m_5 \le m_7,$ $\min(m_4, m_{4'}) = 0,$

with
$$\sum_{i} m_{i} = s$$
. Moreover, if $T \in B(s\overline{\omega}_{7})$ is a $I_{0,2}$ -highest weight element, then
wt $(T) = (m_{7} - m_{4} - m_{5})\overline{\omega}_{1} + (m_{3} + m_{4} - m_{4'} - m_{5} - m_{6} - m_{7})\overline{\omega}_{2} + (m_{3} + m_{4'})\overline{\omega}_{3}$
 $+ (m_{5} - m_{3})\overline{\omega}_{4} + m_{2}\overline{\omega}_{5} + (m_{6} - m_{2})\overline{\omega}_{6} + m_{1}\overline{\omega}_{7}.$
(3.2)

We note that the condition $\min(m_4, m_{4'}) = 0$ is precisely the fact that x_4 and $x_{4'}$ cannot simultaneously appear in an element of $B(s\overline{\omega}_7)$.

Proposition 3.2. The decomposition of $B(s\overline{\omega}_7)$ into type A_6 crystals is given by

$$B(s\overline{\omega}_7) \cong \bigoplus_{\mu} B(\mu),$$

where

$$\mu = (m_7 - m_4 - m_5)\eta_6 + (m_3 + m_{4'})\eta_5 + (m_5 - m_3)\eta_4 + m_2\eta_3 + (m_6 - m_2)\eta_2 + m_1\eta_3$$

such that m_1, \ldots, m_7 satisfy

$$m_2 \le m_6,$$
 $m_3 \le m_5,$ $m_4 + m_5 \le m_7,$
 $\min(m_4, m_{4'}) = 0,$ $s = m_1 + m_2 + m_3 + m_4 + m_{4'} + m_5 + m_6 + m_7$

Moreover, this decomposition is multiplicity free.

Proof. The first claim follows immediately from Lemma 3.1 and relabeling the fundamental weights. For the second claim, consider a weight $\mu = \sum_{i=1}^{6} a_i \eta_i$ such that $B(\mu)$ appears in the A_6 decomposition of $B(s\overline{\omega}_7)$. Thus, we have

$$\begin{split} m_1 &= a_1, & m_2 &= a_3, \\ m_6 &= a_2 + m_2 &= a_2 + a_3, & m_3 &= a_5 - m_{4'}, \\ m_5 &= a_4 + m_3 &= a_4 + a_5 - m_{4'}, & m_7 &= a_6 + m_4 + m_5 &= a_4 + a_5 + a_6 + m_4 - m_{4'}. \end{split}$$

Since $\min(m_4, m_{4'}) = 0$ with $m_4, m_{4'} \ge 0$, there exists a unique m_4 and $m_{4'}$ such that $m_4 - m_{4'} = C$ for any constant C. Next, we have

$$s = m_1 + m_2 + m_3 + m_4 + m_{4'} + m_5 + m_6 + m_7$$

= $a_1 + a_3 + (a_5 - m_{4'}) + m_4 + m_{4'} + (a_4 + a_5 - m_{4'}) + (a_2 + a_3) + (a_4 + a_5 + a_6 + m_4 - m_{4'})$
= $a_1 + a_2 + 2a_3 + 2a_4 + 3a_5 + a_6 + 2(m_4 - m_{4'}).$

Hence, we have

$$m_4 - m_{4'} = \frac{1}{2}(s - a_1 - a_2 - 2a_3 - 2a_4 - 3a_5 - a_6).$$

Therefore, there is a unique m_1, \ldots, m_7 that yields the weight μ .

3.2. Reconstructing the A_7 crystals. In this section, we continue to use the notation of Proposition 3.2.

Lemma 3.3. Let $T \in B^{7,s}$ be a $I_{0,2}$ -highest weight element. Then

$$-\langle \alpha_0^{\vee}, \operatorname{wt}(T) \rangle = m_1 + m_2 + m_3 + m_{4'}$$

Proof. This follows from Equation (3.2), that all elements in $B^{7,s}$ are of level 0, and (2.1).

Since $\langle \alpha_0^{\vee}, \operatorname{wt}(T) \rangle \leq 0$ for all $I_{0,2}$ -highest weight elements $T \in B^{7,s}$, we know that any potential I_2 -highest weight element must have $m_1 = m_2 = m_3 = m_{4'} = 0$. Hence, the possible I_2 -dominant weights are

$$\mu = (m_7 - m_4 - m_5)\eta_6 + m_5\eta_4 + m_6\eta_2$$

 $(i.e., \langle \alpha_i^{\lor}, \mu \rangle \ge 0$ for all $i \in I_2$) such that

 $m_4 + m_5 \le m_7, \qquad m_4 + m_5 + m_6 + m_7 = s.$

Because we can only remove horizontal strips for the branching rule from $A_7 \searrow A_6$ (Proposition 2.4), each of such I_2 -dominant weights μ must correspond to a I_2 -highest weight component $B(\mu)$. Hence, we obtain the following.

Proposition 3.4. The decomposition of $B(s\overline{\omega}_7)$ into type A_7 crystals is given by

$$B(s\overline{\omega}_7) \cong \bigoplus_{\mu} B(\mu),$$

where

$$\mu = (m_7 - m_4 - m_5)\eta_6 + m_5\eta_4 + m_6\eta_2$$

such that m_4, m_5, m_6, m_7 satisfy

$$m_4 + m_5 \le m_7, \qquad s = m_4 + m_5 + m_6 + m_7$$

Moreover, Proposition 2.4 states that for any element T expressed as an A_6 tableau, we add in a horizontal strip to T with every entry an 8 such that every column has even height to obtain the representation as an A_7 tableau. In particular, for a $I_{0,2}$ -highest weight element b, we have

$$\varepsilon_0(b) = m_1 + m_2 + m_3 + m_{4'}, \qquad \qquad \varphi_0(b) = 0$$

Now we combine this to form a combinatorial crystal structure on $B^{7,s}$ by extending the E_7 crystal structure on $B(s\overline{\omega}_7)$ as follows. Let $\overline{\psi}: B(s\overline{\omega}_7) \to \bigoplus_{\overline{\mu}} B(\overline{\mu})$ be the $I_{0,2}$ -crystal (*i.e.*, type A_6) isomorphism given by Proposition 3.2. From Proposition 3.4 and Proposition 2.4, we can uniquely extend the image of $\overline{\psi}$ to highest weight crystals of type A_7 . Therefore, we define

$$e_0 := \overline{\psi}^{-1} \circ e_7^A \circ \overline{\psi}, \qquad \qquad f_0 := \overline{\psi}^{-1} \circ f_7^A \circ \overline{\psi},$$

where e_7^A and f_7^A are the crystal operators from this extended type A_7 crystal. Let $\mathcal{B}^{7,s}$ denote the corresponding crystal.

In order to show this is the combinatorial structure of KR crystal, we need the following uniqueness theorem. The proof is similar to [15, Thm. 3.15] with $K = I_{0,2}$ and using Proposition 3.2 instead of [15, Lemma 3.12].

Theorem 3.5. Let \mathcal{B} and \mathcal{B}' be two affine type $E_7^{(1)}$ crystals such that there exists a I_2 -crystal (i.e., type A_7) isomorphism and I_0 -crystal (i.e., type E_7) isomorphism

$$\Psi_{I_2}\colon \mathcal{B}|_{I_2} \to \mathcal{B}'|_{I_2} \cong \bigoplus_{\mu} B(\mu) \qquad \qquad \Psi_{I_0}\colon \mathcal{B}|_{I_0} \to \mathcal{B}'|_{I_0} \cong B(s\overline{\omega}_7),$$

where the direct sum is over μ given in Proposition 3.4. Then, we have $\Psi_J(b) = \Psi_{I_0}(b)$ for all $b \in \mathcal{B}$. Moreover, there exists an I-crystal isomorphism $\Psi \colon \mathcal{B} \to \mathcal{B}'$.

Corollary 3.6. We have $\mathcal{B}^{7,s} \cong B^{7,s}$.

Proof. We have that $B^{7,s} \cong B(s\overline{\omega}_7)$ as I_0 -crystals since the corresponding KR module is irreducible as a $U_q(\mathfrak{g}_0)$ -module [3]. Hence, the KR crystal $B^{7,s}$ exists by [16, 17]. We can then decompose the crystal $B(s\overline{\omega}_7)$ into $I_{0,2}$ -crystals according to Proposition 3.2, and so the I_2 -crystal decomposition of $B^{7,s}$ is given by Proposition 3.4. Hence, we have isomorphisms between $\mathcal{B}^{7,s}$ and $B^{7,s}$ as I_0 -crystals and I_2 -crystals because these decompositions are multiplicity free.

We have verified that $B^{7,s}$ is a perfect crystal of level s for all $s \leq 4$ using the implementation of $B^{7,s}$ in SAGEMATH [6] done by the second author.

3.3. Combinatorial isomorphism. Let $\langle b \rangle$ denote the $I_{0,2}$ -subcrystal generated by an element b. Consider the subcrystals

$$\begin{array}{ll} \mathcal{C}_1 = \langle x_1 \rangle, & \mathcal{C}_2 = \langle x_6 \rangle, & \mathcal{C}_3 = \langle x_2 \otimes x_6 \rangle, \\ \mathcal{C}_4 = \langle x_5 \otimes x_7 \rangle, & \mathcal{C}_5 = \langle x_3 \otimes x_5 \otimes x_7 \rangle, & \mathcal{C}_6 = \langle x_7 \rangle. \end{array}$$

It is clear that we have $I_{0,2}$ -crystal isomorphisms $\psi_i \colon B(\eta_i) \to C_i$ for all i. (Note that we could have $C_7 = \langle x_4 \otimes x_7 \rangle$, which would correspond to m_4 .) Let $C_{5'} = \langle x_{4'} \rangle$, and we also have a $I_{0,2}$ -crystal isomorphism $\psi_{5'} \colon B(\eta_5) \to C_{5'}$.

Therefore, we can construct a bijection between SSYT for $B(\mu)$ with the A_6 component

$$\mathcal{C}_6^{\otimes a_6} \otimes \mathcal{C}_{5'}^{\otimes m_{4'}} \otimes \mathcal{C}_5^{\otimes a_5 - m_{4'}} \otimes \mathcal{C}_4^{\otimes a_4} \otimes \mathcal{C}_3^{\otimes a_3} \otimes \mathcal{C}_2^{\otimes a_2} \otimes \mathcal{C}_1^{\otimes a_1}$$

by applying ψ_i on each column of height *i*, with possibly $\psi_{5'}$ on columns of height 5, of a SSYT (of type A_6). Note that the result is "out-of-order" in that it does not result is $x_{i_1} \otimes \cdots \otimes x_{i_\ell}$ with $i_1 \leq \cdots \leq i_\ell$ (*i.e.*, a semistandard tableaux given by Proposition 2.3). Thus it would remain to "sort" these elements to obtain an honest component of $B(s\overline{\omega}_7)$. However, we will instead do the reverse process: we refine the isomorphisms ψ_i^{-1} to apply them to elements of $B(\overline{\omega}_7)$, which we combine to apply to an element of $B(s\overline{\omega}_7)$.

Using the tensor product rule, we obtain the following defining crystal isomorphism with a tensor product of A_6 columns:

$$x_1 \mapsto 1, x_2 \mapsto 3, x_3 \mapsto 5, x_4 \mapsto 7, x_{4'} \mapsto \frac{1}{2}, x_5 \otimes x_7 \mapsto \frac{1}{2}, x_6 \mapsto 1, x_7 \mapsto \frac{1}{2}, x_7 \mapsto \frac{1}$$

We need one additional step to decouple $x_5 \otimes x_7$, which we do by

$$x_5 \mapsto \frac{1}{2}{\frac{3}{4}}.$$

We could have arrived at this refined isomorphism by using the A_6 decomposition of $B(\overline{\omega}_7)$. Now using our refined isomorphism and jeu-de-taquin, we obtain a crystal isomorphism Ψ from elements of $B(s\overline{\omega}_7)$ to a direct sum of A_6 SSYT given by Proposition 3.2. Recall that jeu-de-taquin is a crystal isomorphism. Indeed, performing jeu-de-taquin (see, e.g., [10]) on a highest weight element of the tensor product of single column SSYT, we obtain the SSYT

m_1			$m_{4'}$	$m_{4'}$ n			m_6	m_6		m_7
$m_{4'}$ m_5					m_6		m_7			
m_2	$m_{4'}$	m_5			m_7					
$m_{4'}$	m_5			m ₇						
m_3 m_3		$m_{4'}$	m ₇							
m_7										
m_4 m_5										

where row *i* is filled with only *i*. Note that we have included an entry from both m_4 and $m_{4'}$ for illustrative purposes, but both cannot appear. Augmenting the result with a horizontal strip of 8's to obtain all columns of even height (which agrees with Proposition 3.2 after ignoring the columns of height 8), we have

m_1			$m_{4'}$		m_5			m_6		m_7	
$m_{4'}$	$m_{4'}$ m_5			m_6		m_7					m_1
m_2	$m_{4'}$ m			m_7							
$m_{4'}$	$m_{4'}$ m_5			m ₇				m_2			
m_3 r			$m_{4'}$	m_7							
m ₇				$m_3 + m_{4'}$							
m_4 m_5											
$m_4 + m_5$											

Note that the 8's in the even rows can overlap any of the blocks depending on m_1 , m_2 , and $m_3 + m_{4'}$ compares with $m_{4'} + m_5 + m_6 + m_7$, $m_{4'} + m_5 + m_7$, and m_7 respectively.

3.4. **Diagram automorphism.** We can construct the type E_6 crystal decomposition as follows. Let $I_{0,7} := I_0 \setminus \{7\}$. We construct the $I_{0,7}$ -highest weight elements by considering semistandard tableaux in $B(s\overline{\omega}_7)$ consisting of the elements of

$$7 \longleftarrow \overline{7}6 \longleftarrow \overline{7}1 \longleftarrow \overline{7}.$$

The computation is similar to (3.1). Similarly, by reversing the arrows and adding loops at every vertex, we obtain the composition graph $G_7(\overline{\omega}_7)$. Thus, the $I_{0,7}$ -highest weight elements in $B(s\overline{\omega}_7)$ are given by

$$7^{\otimes m_7} \otimes \overline{7}6^{\otimes m_{\overline{7}6}} \otimes \overline{7}1^{\otimes m_{\overline{7}1}} \otimes \overline{7}^{\otimes m_{\overline{7}}} \tag{3.3}$$

(recall that we naturally identify this with a semistandard tableau) with $m_7 + m_{\overline{76}} + m_{\overline{71}} + m_{\overline{7}} = s$.

Proposition 3.7. Let m_b be the number of occurrences of b in a single row tableau $R \in B(s\overline{\omega}_7)$. Then R is an $I_{0,7}$ -highest weight element. Moreover, as E_6 crystals, we have

$$B(s\overline{\omega}_7) \cong \bigoplus_{m_{\overline{7}6} + m_{\overline{7}1} + m_{\overline{7}} \le s} B(m_{\overline{7}6}\overline{\omega}_6 + m_{\overline{7}1}\overline{\omega}_1).$$

Note that the decomposition into E_6 crystals from Proposition 3.7 is not multiplicity free; indeed, the multiplicity of $B(m_{\overline{7}6}\overline{\omega}_6 + m_{\overline{7}1}\overline{\omega}_1)$ is $s - m_{\overline{7}6} - m_{\overline{7}1}$ (the number of distinct values $m_{\overline{7}}$ can take). However, we can distinguish each of the $I_{0,7}$ -highest weight elements by using the extra information (in addition to the weight) of $\langle \operatorname{wt}(b), \alpha_7^{\vee} \rangle = m_7 - m_{\overline{7}1} - m_{\overline{7}6} - m_{\overline{7}}$. Furthermore, by the level-zero condition, we have $\langle \operatorname{wt}(b), \alpha_0^{\vee} \rangle = m_{\overline{7}} - m_{\overline{7}} - m_{\overline{7}6} - m_{\overline{7}1}$. By also using $m_7 + m_{\overline{7}6} + m_{\overline{7}1} + m_{\overline{7}} = s$, we thus obtain the following.

Proposition 3.8. The map $\Phi: B^{7,s} \to B^{7,s}$ given by

$$\Phi\left(7^{\otimes a} \otimes \overline{7}6^{\otimes b} \otimes \overline{7}1^{\otimes c} \otimes \overline{7}^{\otimes d}\right) = 7^{\otimes d} \otimes \overline{7}6^{\otimes c} \otimes \overline{7}1^{\otimes b} \otimes \overline{7}^{\otimes a}$$

and extended as a twisted $I_{0,7}$ -crystal morphism is a twisted $I_{0,7}$ -crystal isomorphism. Moreover Φ is an twisted crystal involution that is induced from the order 2 diagram automorphism of $E_7^{(1)}$.

By restricting to $I_{0,2}$ -highest weight elements, we have that $\Phi = \psi^{-1} \circ \sigma \circ \psi$, where ψ is the isomorphism from Proposition 3.4. This can be seen by equating the weight and $\langle wt(b), \alpha_7^{\vee} \rangle$. Thus, Φ when restricted to the A_7 crystal of $B^{7,s}$ is the order 2 diagram automorphism of A_7 .

As a consequence, we have that $B^{7,s}$ satisfies [9, Assumption 1], and so we have the following by [9, Thm. 4.7]. (We refer the reader to [9] for the precise definitions.)

Corollary 3.9. The tensor product $B = (B^{7,s})^{\otimes m} \otimes \{u_{s\omega_0}\}$ is isomorphic to a Demazure subcrystal $B_w(s\omega_{\tau(0)})$, where $t_{\mu} = w\tau$ in the extended affine Weyl group with $\mu = -m\omega_7$. Moreover, B and $(B^{7,s})^{\otimes m}$ are connected.

3.5. Combinatorial *R*-matrix. We give an explicit description of the combinatorial *R*matrix $R: B^{7,s} \otimes B^{7,s'} \to B^{7,s'} \otimes B^{7,s}$ by noting the classical decomposition of $B^{7,s} \otimes B^{7,s'}$ is multiplicity free as E_7 crystals. We may assume $s \leq s'$ without loss of generality as the combinatorial *R*-matrix is an involution. Thus it is sufficient to consider the $I_{0,7}$ -highest weight elements of $B^{7,s}$, which is given by (3.3). Therefore, the I_0 -highest weight elements have weight

$$(m_{\overline{7}} - m_{\overline{7}} - m_{\overline{7}6} - m_{\overline{7}1} - s')\omega_0 + m_{\overline{7}1}\omega_1 + m_{\overline{7}6}\omega_6 + (s' + m_{\overline{7}} - m_{\overline{7}6} - m_{\overline{7}1} - m_{\overline{7}})\omega_7.$$

The fact that the decomposition into E_7 crystals is multiplicity free is exactly the same reasoning as in the proof of Proposition 3.8. Since the combinatorial *R*-matrix must map classical components to classical components, we have the following.

Theorem 3.10. For $s \leq s'$, the combinatorial *R*-matrix $R: B^{7,s} \otimes B^{7,s'} \mapsto B^{7,s'} \otimes B^{7,s}$ is defined by

$$(7^{\otimes m_7} \otimes \overline{7}6^{\otimes m_{\overline{7}6}} \otimes \overline{7}1^{\otimes m_{\overline{7}1}} \otimes \overline{7}^{\otimes m_{\overline{7}}}) \otimes 7^{\otimes s'} \mapsto (7^{\otimes (s'-s)+m_7} \otimes \overline{7}6^{\otimes m_{\overline{7}6}} \otimes \overline{7}1^{\otimes m_{\overline{7}1}} \otimes \overline{7}^{\otimes m_{\overline{7}}}) \otimes 7^{\otimes s}$$

and extended as an I_0 -crystal isomorphism.

4. Conjectures for $B^{1,s}$

We conclude with a conjectural decomposition of $B^{1,s}$ into A_7 crystals. Recall that A_7 has a natural diagram symmetry σ that respects the $E_7^{(1)}$ diagram symmetry. Thus, proving this conjecture using E_7 crystal decomposition of $B^{1,s} \cong \bigoplus_{k=0}^s B(k\overline{\omega}_1)$ could possibly lead to a proof of [15, Conj. 3.26].

Conjecture 4.1. Let $a, b, c, d \in \mathbb{Z}_{>0}$ such that $a + 2b + 3c + d \leq s$. Then we have

$$B^{1,s} \cong \bigoplus B(a(\eta_1 + \eta_7) + b(\eta_2 + \eta_6) + c(\eta_3 + \eta_5) + d\eta_4)^{\oplus m_{a,b,c,d}},$$

as A_7 crystals, where the multiplicities are $m_{a,b,c,d} = m_{d,s-a-2b-3c}$, where

$$m_{d,s'} = \sum_{i=M}^{d+1} \left\lceil \frac{i}{2} \right\rceil$$

with $M = \max(d + 1 - (s' - d), 0)$.

We compute the multiplicity of η_4 in $B^{1,s}$ using SAGEMATH [6]:

```
sage: for s in range(10):
....: [sum(ceil(i/2) for i in range(max(0,2*d+1-s),d+1+1))
....: for d in range(s+1)]
[1]
[1, 1]
[1, 2, 2]
[1, 2, 3, 2]
[1, 2, 4, 4, 3]
[1, 2, 4, 5, 5, 3]
[1, 2, 4, 6, 7, 6, 4]
[1, 2, 4, 6, 8, 8, 7, 4]
[1, 2, 4, 6, 9, 10, 10, 8, 5]
[1, 2, 4, 6, 9, 11, 12, 11, 9, 5]
```

We compute the decomposition of $B^{7,s}$ into A_7 crystals using SAGEMATH:

```
sage: def compute_branching(s):
          A7 = WeylCharacterRing(['A',7], style="coroots")
. . . . :
          E7 = WeylCharacterRing(['E',7], style="coroots")
. . . . :
          La = E7.fundamental_weights()
. . . . :
          chi = sum(E7(k*La[1]) for k in range(s+1))
          return chi.branch(A7, rule="extended")
sage: compute_branching(1)
A7(0,0,0,0,0,0,0) + A7(0,0,0,1,0,0,0) + A7(1,0,0,0,0,0,1)
sage: compute_branching(2)
2*A7(0,0,0,0,0,0,0) + 2*A7(0,0,0,1,0,0,0) + A7(0,0,0,2,0,0,0)
+ A7(0,1,0,0,0,1,0) + A7(1,0,0,0,0,0,1) + A7(1,0,0,1,0,0,1)
+ A7(2,0,0,0,0,0,2)
sage: compute_branching(3)
2*A7(0,0,0,0,0,0,0) + 3*A7(0,0,0,1,0,0,0) + 2*A7(0,0,0,2,0,0,0)
+ A7(0,0,1,0,1,0,0) + A7(0,1,0,0,0,1,0) + 2*A7(1,0,0,0,0,0,1)
+ 2*A7(1,0,0,1,0,0,1) + A7(0,1,0,1,0,1,0) + A7(0,0,0,3,0,0,0)
+ A7(1,0,0,2,0,0,1) + A7(1,1,0,0,0,1,1) + A7(2,0,0,0,0,0,2)
+ A7(2,0,0,1,0,0,2) + A7(3,0,0,0,0,0,3)
```

We have verified Conjecture 4.1 for $s \leq 9$ by using a heavily optimized version of the branching rule code in SAGEMATH [6].

One way to construct the A_7 highest weight elements would be to use the A_6 highest weight elements, which we can compute from the composition graph in Figure 3 (equivalently Figure 4). From there, we will want the elements invariant (as a set) under the diagram automorphism from [15] as highest weight elements of weight η must map to a highest weight element of weight $-w_0\eta$ under the A_7 diagram automorphism σ .

As a step towards proving Conjecture 4.1, we show the analog of [15, Prop. 2.13] that characterizes the elements in $B(2\overline{\omega}_1) \subseteq B(\overline{\omega}_1)^{\otimes 2}$.

Proposition 4.2. We have

 $(b_1 \otimes c_1) \otimes (b_2 \otimes c_2) \in B(2\overline{\omega}_1) \subseteq B(\overline{\omega}_1)^{\otimes 2} \subseteq B(\overline{\omega}_7)^{\otimes 4}$

if and only if $b_1 \leq b_2$ and $c_1 \leq c_2$ (the comparisons are in $B(\overline{\omega}_7)$) and $(b_1 \otimes c_1) < (b_2 \otimes c_2)$ (the comparison is in $B(\overline{\omega}_1)$).

Proof. This is a finite computation that can be done, e.g., by using the following SAGE-MATH [6] code:

```
sage: L = crystals.Letters(['E',7])
sage: x = L.highest_weight_vector().f_string([7,6,5,4,2,3,4,5,6,7])
sage: A = tensor([x,L.highest_weight_vector()]).subcrystal()
sage: S = tensor([A.module_generators[0].value,
                  A.module_generators[0].value]).subcrystal()
. . . . :
sage: all(P.le(x.value[0][1], x.value[1][1]) for x in S) # Check the b_i condition
True
sage: all(P.le(x.value[0][0], x.value[1][0]) for x in S) # Check the c_i condition
True
sage: P = Poset(L.digraph())
sage: PA = Poset(A.digraph())
sage: data = [[x.value[0], x.value[1]] for x in S]
sage: T = tensor([A, A])
sage: all((not PA.le(A(x[0].value), A(x[1].value))) or x[0].value == x[1] value
          for x in T if P.le(x[0].value[0], x[1].value[0])
. . . . :
                    and P.le(x[0].value[1], x[1].value[1])
. . . . :
                     and [x[0].value, x[1].value] not in data)
. . . . :
True
```

APPENDIX A. SAGEMATH CODE FOR COMPOSITION GRAPHS

To compute the composition graph in Figure 3, we first do some setup:

```
sage: L = crystals.Letters(['E',7])
sage: x = L.highest_weight_vector().f_string([7,6,5,4,2,3,4,5,6,7])
sage: A = tensor([x,L.highest_weight_vector()]).subcrystal()
sage: TA = tensor([A.module_generators[0], A.module_generators[0]]).subcrystal()
sage: _ = A.list()
```

To compute the composition graph $G_2(\overline{\omega}_1)$, we run the following functions:

```
def check_le(x, y):
    return tensor([x,y]) in TA

def composition_graph(J):
    I = A.index_set()
    ImJ = sorted(set(I) - set(J))
    G = DiGraph([[b for b in A if b.is_highest_weight(ImJ)], check_le])
    num_verts = 0
    while num_verts != G.num_verts():
```

FIGURE 3. The composition graph $G_2(\overline{\omega}_1)$ to compute the $(I_0 \setminus \{2\})$ -highest weight elements. We have suppressed the loops that occur at every node except $\overline{32} \otimes \overline{23}$.

```
num_verts = G.num_verts()
    verts = set(G.vertices())
    for b in A:
        ep = set([i for i in I if b.epsilon(i) > 0])
        Jplus = set(list(J) + [i for i in ImJ
                                if any(bp.phi(i) > 0 and check_le(b,bp
                                       for bp in verts)
                               ]
                   )
        if ep.issubset(Jplus):
            G.add_vertex(b)
            for bp in verts:
                if check_le(b, bp):
                    G.add_edge(b, bp)
loops = G.loops()
G = G.transitive_reduction()
for l in loops:
    G.add_edge(*1)
G.set_latex_options(format='dot2tex')
return G
```

References

- Rekha Biswal and Travis Scrimshaw. Existence of Kirillov-Reshetikhin crystals for multiplicity free nodes. Publ. Res. Inst. Math. Sci., 56(4), 2020.
- [2] Daniel Bump and Anne Schilling. Crystal bases. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2017. Representations and combinatorics.




FIGURE 4. The composition graphs of Figure 3 with every node written in "compact form," where a k adds 1 to $\varphi_k(b)$ and \overline{k} adds 1 to $\varepsilon_k(b)$. Recall that the only vertex that does not have a loop is $\overline{22}$.

- [3] Vyjayanthi Chari. On the fermionic formula and the Kirillov-Reshetikhin conjecture. Int. Math. Res. Not. IMRN, (12):629-654, 2001.
- [4] Vyjayanthi Chari and Andrew Pressley. Quantum affine algebras and their representations. In *Representations of groups (Banff, AB, 1994)*, volume 16 of *CMS Conf. Proc.*, pages 59–78. Amer. Math. Soc., Providence, RI, 1995.
- [5] Vyjayanthi Chari and Andrew Pressley. Twisted quantum affine algebras. Comm. Math. Phys., 196(2):461–476, 1998.
- [6] The Sage Developers. Sage Mathematics Software (Version 9.2). The Sage Development Team, 2020. http://www.sagemath.org.
- [7] Ghislain Fourier, Masato Okado, and Anne Schilling. Kirillov-Reshetikhin crystals for nonexceptional types. Adv. Math., 222(3):1080–1116, 2009.
- [8] Ghislain Fourier, Masato Okado, and Anne Schilling. Perfectness of Kirillov-Reshetikhin crystals for nonexceptional types. *Contemp. Math.*, 506:127–143, 2010.
- [9] Ghislain Fourier, Anne Schilling, and Mark Shimozono. Demazure structure inside Kirillov-Reshetikhin crystals. J. Algebra, 309(1):386–404, 2007.
- [10] William Fulton. Young tableaux, volume 35 of London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 1997. With applications to representation theory and geometry.
- [11] G. Hatayama, A. Kuniba, M. Okado, T. Takagi, and Y. Yamada. Remarks on fermionic formula. In Recent developments in quantum affine algebras and related topics (Raleigh, NC, 1998), volume 248 of Contemp. Math., pages 243–291. Amer. Math. Soc., Providence, RI, 1999.
- [12] Goro Hatayama, Atsuo Kuniba, Masato Okado, Taichiro Takagi, and Zengo Tsuboi. Paths, crystals and fermionic formulae. In *MathPhys odyssey*, 2001, volume 23 of *Prog. Math. Phys.*, pages 205–272. Birkhäuser Boston, Boston, MA, 2002.
- [13] David Hernandez. Kirillov-Reshetikhin conjecture: the general case. Int. Math. Res. Not. IMRN, (1):149–193, 2010.
- [14] Rei Inoue, Atsuo Kuniba, and Taichiro Takagi. Integrable structure of box-ball systems: crystal, Bethe ansatz, ultradiscretization and tropical geometry. J. Phys. A, 45(7):073001, 64, 2012.

- [15] Brant Jones and Anne Schilling. Affine structures and a tableau model for E_6 crystals. J. Algebra, 324(9):2512-2542, 2010.
- [16] Seok-Jin Kang, Masaki Kashiwara, Kailash C. Misra, Tetsuji Miwa, Toshiki Nakashima, and Atsushi Nakayashiki. Affine crystals and vertex models. In *Infinite analysis, Part A, B (Kyoto, 1991)*, volume 16 of Adv. Ser. Math. Phys., pages 449–484. World Sci. Publ., River Edge, NJ, 1992.
- [17] Seok-Jin Kang, Masaki Kashiwara, Kailash C. Misra, Tetsuji Miwa, Toshiki Nakashima, and Atsushi Nakayashiki. Perfect crystals of quantum affine Lie algebras. Duke Math. J., 68(3):499–607, 1992.
- [18] M. Kashiwara, K. C. Misra, M. Okado, and D. Yamada. Perfect crystals for $U_q(D_4^{(3)})$. J. Algebra, 317(1):392-423, 2007.
- [19] Masaki Kashiwara. On crystal bases of the q-analogue of universal enveloping algebras. Duke Math. J., 63(2):465–516, 1991.
- [20] Masaki Kashiwara. On level-zero representations of quantized affine algebras. Duke Math. J., 112(1):117– 175, 2002.
- [21] Atsuo Kuniba, Tomoki Nakanishi, and Junji Suzuki. T-systems and Y-systems in integrable systems. J. Phys. A, 44(10):103001, 146, 2011.
- [22] Cristian Lenart. Combinatorial aspects of the K-theory of Grassmannians. Ann. Comb., 4(1):67–82, 2000.
- [23] Cristian Lenart, Satoshi Naito, Daisuke Sagaki, Anne Schilling, and Mark Shimozono. A uniform model for Kirillov-Reshetikhin crystals I: Lifting the parabolic quantum Bruhat graph. Int. Math. Res. Not. IMRN, (7):1848–1901, 2015.
- [24] Cristian Lenart, Satoshi Naito, Daisuke Sagaki, Anne Schilling, and Mark Shimozono. A uniform model for Kirillov-Reshetikhin crystals II. Alcove model, path model, and P = X. Int. Math. Res. Not. IMRN, 2016.
- [25] Cristian Lenart, Satoshi Naito, Daisuke Sagaki, Anne Schilling, and Mark Shimozono. A uniform model for Kirillov-Reshetikhin crystals III: Nonsymmetric Macdonald polynomials at t = 0 and Demazure characters. *Transform. Groups*, pages 1–39, 2017.
- [26] Cristian Lenart and Travis Scrimshaw. On higher level Kirillov–Reshetikhin crystals, Demazure crystals, and related uniform models. J. Algebra, 539:285–304, 2019.
- [27] Xuan Liu and Travis Scrimshaw. A uniform approach to soliton cellular automata using rigged configurations. Ann. Henri Poincaré, 20(4):1175–1215, 2019.
- [28] Satoshi Naito and Daisuke Sagaki. Path model for a level-zero extremal weight module over a quantum affine algebra. Int. Math. Res. Not., (32):1731–1754, 2003.
- [29] Satoshi Naito and Daisuke Sagaki. Path model for a level-zero extremal weight module over a quantum affine algebra. II. Adv. Math., 200(1):102–124, 2006.
- [30] Satoshi Naito and Daisuke Sagaki. Crystal structure on the set of Lakshmibai-Seshadri paths of an arbitrary level-zero shape. Proc. Lond. Math. Soc. (3), 96(3):582–622, 2008.
- [31] Katsuyuki Naoi. Existence of Kirillov-Reshetikhin crystals of type $G_2^{(1)}$ and $D_4^{(3)}$. J. Algebra, 512:47–65, 2018.
- [32] Katsuyuki Naoi and Travis Scrimshaw. Existence of Kirillov-Reshetikhin crystals for near adjoint nodes in exceptional types. J. Pure Appl. Algebra, 225(5):106593, 2021.
- [33] Masato Okado and Anne Schilling. Existence of Kirillov-Reshetikhin crystals for nonexceptional types. Represent. Theory, 12:186–207, 2008.
- [34] Masato Okado, Anne Schilling, and Travis Scrimshaw. Rigged configuration bijection and proof of the X = M conjecture for nonexceptional affine types. J. Algebra, 516:1–37, 2018.
- [35] Anne Schilling and Mark Shimozono. X = M for symmetric powers. J. Algebra, 295(2):562–610, 2006.
- [36] Travis Scrimshaw. Uniform description of the rigged configuration bijection. Selecta Math. (N.S.), 26(3):article 42, 2020.
- [37] Shigenori Yamane. Perfect crystals of $U_q(G_2^{(1)})$. J. Algebra, 210(2):440–486, 1998.

(R. Biswal) MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, 53111 BONN, GERMANY *Current address*: University of Edinburgh, Old College, South Bridge, Edinburgh EH8 9YL, United Kingdom

Email address: rbiswal@ed.ac.uk *URL*: https://rekhabiswal.github.io/

(T. Scrimshaw) School of Mathematics and Physics, The University of Queensland, St. Lucia, QLD 4072, Australia

Email address: tcscrims@gmail.com

URL: https://people.smp.uq.edu.au/TravisScrimshaw/