

KY FAN INEQUALITIES FOR BIVARIATE MEANS

ALFRED WITKOWSKI

ABSTRACT. In this note we give sufficient conditions for bivariate, homogeneous, symmetric means M and N to satisfy Ky Fan inequalities

$$\frac{M}{M'} < \frac{N}{N'} \quad \text{and} \quad \frac{1}{M} - \frac{1}{M'} < \frac{1}{N} - \frac{1}{N'}.$$

1. INTRODUCTION

Denote by G and A the geometric and arithmetic means of nonnegative numbers x_i , $i = 1, \dots, n$

$$G(x_1, \dots, x_n) = \prod_{i=1}^n x_i^{1/n}, \quad A(x_1, \dots, x_n) = \frac{1}{n} \sum_{i=1}^n x_i.$$

The classical Ky Fan inequality states that if $0 < x_i \leq 1/2$, then

$$\frac{G(x_1, \dots, x_n)}{G(1-x_1, \dots, 1-x_n)} \leq \frac{A(x_1, \dots, x_n)}{A(1-x_1, \dots, 1-x_n)}.$$

This result has been published in 1961 and generalized in many directions. For more information see [1].

This note is inspired by the result of Neuman and Sándor [2] where we find the following sequence of inequalities

$$(1) \quad \frac{G}{G'} \leq \frac{L}{L'} \leq \frac{P}{P'} \leq \frac{A}{A'} \leq \frac{NS}{NS'} \leq \frac{T}{T'},$$

(here L, P, NS, T stand for the logarithmic, first Seiffert, Neuman-Sándor and second Seiffert means of arguments $0 < x, y \leq 1/2$, and prime denotes the same mean with arguments $1-x$ and $1-y$).

The aim of this note is to establish the conditions for two symmetric, homogeneous means M, N under which the Ky Fan inequality

$$\frac{M(x, y)}{M(1-x, 1-y)} \leq \frac{N(x, y)}{N(1-x, 1-y)}$$

holds.

Date: March 5, 2019.

1991 Mathematics Subject Classification. 26D60, 26E20.

Key words and phrases. Ky Fan inequality, mean, homogeneity.

2. DEFINITION AND NOTATION

A function $M : \mathbb{R}_+^2 \rightarrow \mathbb{R}$ is called a *mean* if for all $x, y > 0$ holds

$$\min(x, y) \leq M(x, y) \leq \max(x, y).$$

Mean is *symmetric* if for all $M(x, y) = M(y, x)$ for all $x, y > 0$ and *homogeneous* if for all $\lambda > 0$

$$M(\lambda x, \lambda y) = \lambda M(x, y).$$

A function $m : (0, 1) \rightarrow \mathbb{R}$ satisfying

$$\frac{z}{1+z} \leq m(z) \leq \frac{z}{1-z}$$

is called a *Seiffert function*.

Let us recall a result from [7]

Theorem 2.1. *The formula*

$$M(x, y) = \frac{|x - y|}{2m\left(\frac{|x - y|}{x + y}\right)}$$

establishes a one-to-one correspondence between the set of symmetric, homogeneous means and the set of Seiffert functions.

If there is no risk of ambiguity we shall skip the argument of means. For $0 < x, y \leq 1/2$ and an arbitrary mean M we shall denote $M(1 - x, 1 - y)$ by M' .

3. KY FAN INEQUALITIES

Our first theorem shows that the Ky Fan inequality is a consequence of monotonicity of the ratio of Seiffert means.

Theorem 3.1. *Let M, N be two symmetric, homogeneous means and m, n be their Seiffert functions. If the function $\frac{n}{m}$ is decreasing, then for all $0 < x, y \leq 1/2$ the Ky Fan inequality*

$$\frac{M(x, y)}{M(1 - x, 1 - y)} \leq \frac{N(x, y)}{N(1 - x, 1 - y)}$$

holds.

Proof. Denote $x' = 1 - x, y' = 1 - y$. Clearly $|x - y| = |x' - y'|$ and $x + y \leq x' + y'$, thus $\frac{|x - y|}{x + y} \geq \frac{|x' - y'|}{x' + y'}$. By Theorem 2.1 we obtain

$$\frac{M(x, y)}{N(x, y)} = \frac{n}{m} \left(\frac{|x - y|}{x + y} \right) < \frac{n}{m} \left(\frac{|x' - y'|}{x' + y'} \right) = \frac{M(x', y')}{N(x', y')},$$

which concludes the proof. \square

Sometimes it is more convenient to investigate the ratio of means instead of their Seiffert functions.

Corollary 3.2. *In the assumptions of Theorem 3.1 if the function $q(t) = \frac{M(1, t)}{N(1, t)}$ increases in $(0, 1)$, then for all $0 < x, y \leq 1/2$ the Ky Fan inequality*

$$\frac{M(x, y)}{M(1 - x, 1 - y)} \leq \frac{N(x, y)}{N(1 - x, 1 - y)}$$

holds.

Proof. Let us recall the formula connecting the mean and its Seiffert function ([7])

$$(2) \quad m(z) = \frac{z}{M(1+z, 1-z)}.$$

This gives:

$$\frac{n}{m}(z) = \frac{M(1+z, 1-z)}{N(1+z, 1-z)} = \frac{M\left(1, \frac{1-z}{1+z}\right)}{N\left(1, \frac{1-z}{1+z}\right)} = \frac{M(1, s)}{N(1, s)} = q(s).$$

Here $s = \frac{1-z}{1+z}$ decreases from 1 to 0 as z travels in the opposite direction, so n/m decreases if and only if q increases. \square

Let us illustrate our results with some examples:

Example 3.1. This result was proved by Chan, Goldberg and Gonek in [3]: if

$$A_r(x, y) = \begin{cases} \left(\frac{x^r + y^r}{2}\right)^{1/r} & r \neq 0, \\ \sqrt{xy} & r = 0 \end{cases}$$

is a power mean of order r , then for $r < s$ holds

$$(3) \quad \frac{A_r}{A'_r} \leq \frac{A_s}{A'_s}.$$

Indeed, for $0 < t < 1$ and $rs \neq 0$

$$\operatorname{sgn} \frac{d}{dt} \log \frac{A_r(1, t)}{A_s(1, t)} = \operatorname{sgn} \frac{t^r - t^s}{t(t^r + 1)(t^s + 1)} = \operatorname{sgn}(s - r),$$

else if $s = 0$, then

$$\operatorname{sgn} \frac{d}{dt} \log \frac{A_r(1, t)}{A_0(1, t)} = \operatorname{sgn} \frac{t^r - 1}{t(2t^r + 1)} = -\operatorname{sgn} r,$$

so the inequality (3) is true by Corollary 3.2.

Example 3.2. The last term in the chain of inequalities (1) contains the second Seiffert mean

$$T(x, y) = \frac{|x - y|}{2 \arctan \frac{|x - y|}{x + y}}.$$

Let $Q(x, y) = \sqrt{\frac{x^2 + y^2}{2}}$ be the quadratic (or root mean square).

We have $t(z) = \arctan z$ and $q(z) = \frac{z}{\sqrt{1+z^2}}$ and

$$\frac{d}{dz} \frac{t}{q}(z) = \frac{z - \arctan z}{z^2 \sqrt{1+z^2}} > 0,$$

so by Theorem 3.1 we can extend the chain (1)

$$\frac{T}{T'} < \frac{Q}{Q'}.$$

Example 3.3. Consider now the Heronian mean

$$\operatorname{He}(x, y) = \frac{x + \sqrt{xy} + y}{3}.$$

For $0 < t < 1$ we have

$$q(t) = \frac{\text{He}(1, t)}{\text{A}_{2/3}(1, t)} = \frac{2\sqrt{2}}{3} \frac{t + \sqrt{t} + 1}{(t^{2/3} + 1)^{3/2}}$$

and

$$\frac{dq}{dt}(t) = \frac{\sqrt{2}}{3} \frac{(1 - t^{1/6})^3(1 + t^{1/6})}{(1 + t^{2/3})^{5/2}t^{1/2}} > 0.$$

We also see that the quotient

$$\begin{aligned} \frac{\text{He}(1, t)}{\text{A}_{1/2}(1, t)} &= \frac{4}{3} \frac{t + \sqrt{t} + 1}{t + 2\sqrt{t} + 1} = \frac{4}{3} \left[1 - \frac{\sqrt{t}}{t + 2\sqrt{t} + 1} \right] \\ &= \frac{4}{3} \left[1 - \frac{1}{\sqrt{t} + 2 + \frac{1}{\sqrt{t}}} \right] \end{aligned}$$

decreases in $(0, 1)$. Therefore by Corollary 3.2 we have

$$\frac{\text{A}_{1/2}}{\text{A}'_{1/2}} < \frac{\text{He}}{\text{He}'} < \frac{\text{A}_{2/3}}{\text{A}'_{2/3}}.$$

Example 3.4. We know ([5]) that the logarithmic mean $\text{L}(x, y) = \frac{x-y}{\log x - \log y}$ satisfies the inequality $\text{L} < \text{A}_{1/3}$. Consider

$$q(t) = \frac{\text{A}_{1/3}(1, t)}{\text{L}(1, t)} = \frac{(t^{1/3} + 1)^3}{8(t-1)} \log t.$$

Its derivative equals

$$\frac{dq}{dt}(t) = \frac{(t^{1/3} + 1)^2 [t^{4/3} + t - t^{1/3} - 1 - (t + t^{1/3}) \log t]}{8t(t-1)^2}.$$

To evaluate the sign of the expression in square brackets substitute $t = s^3$ and calculate its Taylor series at $s = 1$

$$\begin{aligned} t^{4/3} + t - t^{1/3} - 1 - (t + t^{1/3}) \log t &= s^4 + s^3 - s - 1 - 3(s^2 + 1)s \log s \\ &= - \sum_{n=5}^{\infty} \frac{n^2 - 5n + 12}{n(n-1)(n-2)(n-3)} (1-s)^n < 0, \end{aligned}$$

which proves the inequality

$$\frac{\text{L}}{\text{L}'} < \frac{\text{A}_{1/3}}{\text{A}'_{1/3}}.$$

Example 3.5. There is another mean similar to the Seiffert means that lies between the arithmetic and the first Seiffert mean:

$$\text{P}(x, y) = \frac{|x-y|}{2 \arcsin\left(\frac{|x-y|}{x+y}\right)} \leq \frac{|x-y|}{2 \sinh\left(\frac{|x-y|}{x+y}\right)} = \text{S}_{\sinh}(x, y) \leq \frac{x+y}{2} = \text{A}(x, y).$$

Their respective Seiffert functions are \arcsin , \sinh and id . The hyperbolic sine is convex for positive arguments, therefore $\frac{\sinh z}{z}$ increases as a divided difference of a convex function. So by Theorem 3.1 we conclude

$$\frac{\text{S}_{\sinh}}{\text{S}'_{\sinh}} \leq \frac{\text{A}}{\text{A}'}$$

Example 3.6. It is known [4] that $A_{1/2} < P$. We have $a_{1/2}(z) = \frac{2z}{1+\sqrt{1-z^2}}$. To check the monotonicity of $p/a_{1/2}$ we substitute $z = \sin t$ to get $\frac{t}{2\sin t}(1 + \cos t)$. Its derivative $\frac{t-\sin t}{2\cos t-2}$ is negative, so

$$\frac{A_{1/2}}{A'_{1/2}} < \frac{P}{P'}.$$

Example 3.7. The tangent is also a Seiffert function. It satisfies the inequality $\tan z < \operatorname{artanh} z$ for $0 < z < 1$, so their means satisfy

$$L(x, y) = \frac{|x-y|}{2 \operatorname{artanh}\left(\frac{|x-y|}{x+y}\right)} < \frac{|x-y|}{2 \tan\left(\frac{|x-y|}{x+y}\right)} = S_{\tan}(x, y).$$

Let us investigate the quotient of the Seiffert functions:

$$\frac{d}{dz} \frac{\operatorname{artanh} z}{\tan z} = \frac{\frac{1}{2} \sin 2z - (1-z^2) \operatorname{artanh} z}{(1-z^2) \sin^2 z}$$

and using Taylor expansion we obtain

$$\begin{aligned} \frac{1}{2} \sin 2z - (1-z^2) \operatorname{artanh} z &= \frac{1}{2} \sum_{n=0}^{\infty} \frac{(-1)^n (2z)^{2n+1}}{(2n+1)!} - (1-z^2) \sum_{n=0}^{\infty} \frac{z^{2n+1}}{2n+1} \\ &= \sum_{n=1}^{\infty} \frac{1}{(2n+1)(2n-1)} \left[2 + \frac{(-1)^n 4^n (2n-1)}{(2n)!} \right] z^{2n+1} > 0, \end{aligned}$$

because all the coefficients in square brackets are nonnegative. Thus by Theorem 3.1

$$\frac{L}{L'} < \frac{S_{\tan}}{S'_{\tan}}.$$

4. HARMONIC KY FAN INEQUALITIES

In [6] the authors derive the inequalities of type

$$\frac{1}{A} - \frac{1}{A'} < \frac{1}{P} - \frac{1}{P'}$$

from the inequalities $P/P' < A/A'$ using some elementary algebraic transformation. Here we show an approach that bases on the concept of Seiffert functions.

Suppose M and N are two symmetric and homogeneous means. Using Theorem 2.1 the inequality

$$\frac{1}{M} - \frac{1}{M'} \leq \frac{1}{N} - \frac{1}{N'}$$

can be written as

$$m\left(\frac{|x-y|}{x+y}\right) - n\left(\frac{|x-y|}{x+y}\right) \leq m\left(\frac{|x-y|}{2-x-y}\right) - n\left(\frac{|x-y|}{2-x-y}\right).$$

Since $2-x-y > x+y$ for $0 < x, y < 1/2$ we can formulate the following result.

Theorem 4.1. *Suppose the means M and N are symmetric and homogeneous and m and n are their Seiffert means. If the function $m-n$ decreases, then the Ky Fan inequality*

$$\frac{1}{M} - \frac{1}{M'} \leq \frac{1}{N} - \frac{1}{N'}$$

holds.

Taking into account the formula (2) and setting $s = (1+z)/(1-z)$ we get the following

Corollary 4.2. *In the assumptions of Theorem 4.1 if the function*

$$g(s) = (s-1) \left(\frac{1}{M(s,1)} - \frac{1}{N(s,1)} \right)$$

decreases for $s > 1$, then the inequality

$$\frac{1}{M} - \frac{1}{M'} \leq \frac{1}{N} - \frac{1}{N'}$$

holds.

Example 4.1. Consider the chain of inequalities between Seiffert functions (see [7, Lemma 3.1]) valid for $0 < z < 1$

$$z > \operatorname{arsinh} z > \sin z > \arctan z > \tanh z.$$

The mean corresponding to the second function is called Neuman-Sándor mean — NS and the fourth one generates the second Seiffert mean — T.

$$\text{Since } \cosh^2 z - 1 = (\cosh z + 1)(\cosh z - 1) > 2 \cdot \frac{z^2}{2}$$

$$\frac{d}{dz}(\arctan z - \tanh z) = \frac{\cosh^2 z - 1 - z^2}{(1+z^2)\cosh^2 z} > 0.$$

Inequality $\cos z > 1 - z^2/2$ leads to

$$\frac{d}{dz}(\sin z - \arctan z) = \frac{(1+z^2)\cos z - 1}{1+z^2} > \frac{(1+z^2)(1-z^2/2) - 1}{1+z^2} = \frac{z^2(1-z^2)}{2(1+z^2)} > 0.$$

On the other hand $\cos z < 1 - z^2/2 + z^4/6$, so

$$\begin{aligned} \frac{d}{dz}(\operatorname{arsinh} z - \sin z) &= \frac{1 - \sqrt{1+z^2}\cos z}{\sqrt{1+z^2}} > \frac{1 - \sqrt{(1+z^2)\left(1 - \frac{z^2}{2} + \frac{z^4}{6}\right)^2}}{\sqrt{1+z^2}} \\ &= \frac{1 - \sqrt{1 - \frac{5}{12}z^4(1-z^2) - \frac{1}{36}z^8(5-z^2)}}{\sqrt{1+z^2}} > 0. \end{aligned}$$

And finally

$$\frac{d}{dz}(z - \operatorname{arsinh} z) = 1 - \frac{1}{\sqrt{1+z^2}} > 0.$$

Therefore by Theorem 4.1

$$\frac{1}{S_{\tanh}} - \frac{1}{S'_{\tanh}} \leq \frac{1}{T} - \frac{1}{T'} \leq \frac{1}{S_{\sin}} - \frac{1}{S'_{\sin}} \leq \frac{1}{NS} - \frac{1}{NS'} \leq \frac{1}{A} - \frac{1}{A'}.$$

Example 4.2. On the other side of the arithmetic mean there are two chains of inequalities for Seiffert means involving sine and tangent (see [7, Lemma 3.2]):

$$(4) \quad z < \sinh z < \left\{ \begin{array}{c} \tan z \\ \arcsin z \end{array} \right\} < \operatorname{artanh} z.$$

The two Seiffert functions in curly brackets are not comparable, arcsine defines the first Seiffert mean and inverse hyperbolic tangent corresponds to the logarithmic mean.

The difference $\sinh z - z$ increases, because this is the gap between a convex function and its supporting line.

To show that $\frac{d}{dz}(\tan z - \sinh z) = \frac{1 - \cos^2 z \cosh z}{\cos^2 z}$ is positive note that

$$1 - \cos^2 z \cosh z > 1 - \cos z \cosh z =: f(z).$$

The function f vanishes at $z = 0$ and

$$f'(z) = \cos z \cosh z (\tan z - \tanh z) > 0,$$

because $\tan z > z > \tanh z$. Thus f' is positive, and so is f .

The difference between inverse hyperbolic tangent and tangent also increases, since

$$\frac{d}{dz}(\operatorname{artanh} z - \tan z)' = \frac{1}{1 - z^2} - \frac{1}{\cos^2 z} = \frac{z^2 - \sin^2 z}{(1 - z^2) \cos^2 z} > 0.$$

To deal with the lower chain of inequalities (4) note that

$$(5) \quad \begin{aligned} \cosh z &= 1 + \frac{z^2}{2!} + \frac{z^4}{4!} + \frac{z^6}{6!} + \dots \\ &< 1 + \frac{z^2}{2} + \frac{z^4}{12} \left(\frac{1}{2} + \frac{1}{4} + \dots \right) = 1 + \frac{z^2}{2} + \frac{z^4}{12}. \end{aligned}$$

Now

$$\frac{d}{dz}(\arcsin z - \sinh z) = \frac{1 - \sqrt{1 - z^2} \cosh z}{\sqrt{1 - z^2}} > 0$$

because using (5)

$$\begin{aligned} 1 - \sqrt{1 - z^2} \cosh z &> 1 - \sqrt{(1 - z^2) \left(1 + \frac{z^2}{2} + \frac{z^4}{12} \right)^2} \\ &= 1 - \sqrt{1 - \frac{7z^4}{12} - \frac{z^6}{3} - \frac{11z^8}{144} - \frac{z^{10}}{144}} > 0. \end{aligned}$$

Comparison of the last pair is simple

$$\frac{d}{dz}(\operatorname{artanh} z - \arcsin z) = \frac{1}{1 - z^2} - \frac{1}{\sqrt{1 - z^2}} > 0.$$

Now we can use Theorem 4.1 to write the chain of inequalities

$$\frac{1}{A} - \frac{1}{A'} < \frac{1}{S_{\sinh}} - \frac{1}{S'_{\sinh}} < \left\{ \frac{1}{S_{\tan}} - \frac{1}{S'_{\tan}} \right\} < \frac{1}{L} - \frac{1}{L'}.$$

NOTE: In the proof of Theorems 3.1 we use monotonicity of $\frac{m}{n}$ to obtain inequality between $\frac{m}{n} \left(\frac{y-x}{x+y} \right)$ and $\frac{m}{n} \left(\frac{y-x}{2-x-y} \right)$. It is natural to ask whether the inverse is true, i.e. if for all $0 < x < y < 1/2$ the inequality

$$(6) \quad f \left(\frac{y-x}{x+y} \right) > f \left(\frac{y-x}{2-x-y} \right)$$

holds then f needs to be monotone. The counterexample has been produced by *tometomek91* - user of the mathematical portal matematyka.pl.

It is clear that $\frac{y-x}{x+y} > \frac{y-x}{2-x-y}$. Moreover $\sup_{0 < x < y < 1/2} \frac{y-x}{2-x-y} = \frac{1}{3}$. Therefore every function f satisfying the two conditions:

- f is increasing on $(0, 1/3)$
- $f(x) \geq \lim_{t \nearrow 1/3} f(t)$ for $x \in [1/3, 1]$

satisfies (6).

5. ACKNOWLEDGEMENT

The author wishes to thank Janusz Matkowski and Monika Nowicka for improvement suggestions and efficient bug and typo hunting.

6. REFERENCES

REFERENCES

- [1] H. Alzer. The inequality of Ky Fan and related results. *Acta Applicandae Mathematica*, 38(3):305–354, Mar 1995.
- [2] E. Neuman and J. Sándor. On the Schwab-Borchardt mean. *Math. Pannon.*, 14(2):253–266, 2003.
- [3] F. Chan, D. Goldberg, and S. Gonek. On extensions of an inequality among means. *Proc. Amer. Math. Soc.*, 42:202–207 (1974), 1973.
- [4] A.A. Jagers. Solution of problem 887. *Nieuw Arch. Wisk. (4)*, 12:230–231, 1994.
- [5] Tung Po Lin. The power mean and the logarithmic mean. *Amer. Math. Monthly*, 81:879–883, 1974.
- [6] E. Neuman and J. Sándor. On the Schwab-Borchardt mean. II. *Math. Pannon.*, 17(1):49–59, 2006.
- [7] A. Witkowski. On Seiffert-like means. *J. Math. Inequal.*, 9(4):1071–1092, 2015.
E-mail address: `alfred.witkowski@utp.edu.pl`