

MICHAEL HIPPKKE

Sonneberg Observatory, Sternwartestr. 32, 96515 Sonneberg, Germany

ABSTRACT

In recent publications, Borra (2013); Borra & Trottier (2016); Borra (2017) claimed the discovery of ultra-short (10^{-12} s) optical pulses originating from stars and galaxies, asserted to be sent by extraterrestrial intelligence. I show that these signals are not astrophysical or instrumental in nature, but originate from the non-random spacings of spectral absorption lines. They can be shown to arise in their clearest form in synthetic solar spectra, as these do not suffer from noise.

1. INTRODUCTION

Over the last years, Borra (2010a,b, 2012) developed and publicized a novel method, dubbed Spectral Fourier Transform (SFT), to detect and characterize ultra-short pulses from spectra. It appears that the method has gradually evolved from work by Delisle & St-Arnaud (1970); Mandel & Wolf (1976); Chin et al. (1992). It is important to appreciate the effort to build such a new technique, a tool which may prove valuable in the future.

As a first application, SFTs were used to examine astronomical spectra from the SDSS survey of galaxies (Borra 2013), stars (Borra & Trottier 2016), and exotic objects (Borra 2017). Astrophysical pulse detections were claimed with repetition times $\rho = 1.64 \times 10^{-12}$ s (Borra & Trottier 2016, stars) and $\rho = 1.09 \times 10^{-13}$ s (Borra 2013, galaxies). The authors speculate that the factor of ~ 15 between these repetition times is caused by different originators: Extraterrestrial intelligence (section 7.2 in Borra & Trottier 2016) versus black holes. Recent re-observations of 3 candidate stars with different equipment have failed to reproduce the signals (Isaacson et al. 2018).

The Berkeley SETI Research Center team (2016) speculated that the signals may be artifacts from instrumental optics, fringing, movement of the telescope, variations in observing conditions, the process of wavelength calibration, or “inconsistencies in the manufacture of detectors (...) of high resolution spectrographs”. This is a valid concern, as a myriad of CCD artifacts exist (De-sai et al. 2016) and more are discovered regularly (e.g., Boone et al. 2018).

In this note, I show that the signals are caused by the non-random spacing of spectral absorption lines (Fowler 1924). They can be shown to occur in their clearest form in synthetic spectra, as these do not suffer from noise. They can also be shown to arise in high signal-to-noise ratio spectra of our sun and many other sources.

2. METHOD: SPECTRAL FOURIER TRANSFORMS

Most commonly, Fourier transforms are applied to time series data, to identify periodic signals. Equally, spectra can be Fourier transformed to yield periodic signals (Chin et al. 1992). A seminal summary of the method is given by Borra (2012).

A unique feature of SFTs is their sensitivity to ultra-short periods, $\rho_{\min} = \lambda_{\min}/c \approx 10^{-15}$ s for $\lambda_{\min} = 300$ nm optical pulses, which is 6 orders of magnitude shorter than time-resolved (10^{-9} s) photometry using photomultipliers (Abbasi et al. 2010). The period of the longest detectable pulses depends on λ_{\min} and the spectral resolution R . For SDSS spectra ($R = 4,000$ at a wavelength $380 < \lambda < 920$ nm, Yanny et al. 2009; Eisenstein et al. 2011) we get $\rho_{\max} = R \rho_{\min} \sim 4 \times 10^{-12}$ s.

SFTs are not sensitive to the actual pulse length Δt_{\min} , but to the duration between pulses (ρ), so that $1/\rho$ is the pulse repetition rate. As the pulse lengths are unresolved in time, it may well be that $\Delta t_{\min} \ll \rho$, exacerbating the atmospheric effects of group delay dispersion. Atmospheric models indicate that turbulence smears picosecond pulses, so that it may well be impossible to use SFT-based methods on the ground. Then, the pulses could not be of astrophysical origin. Without picosecond photon counters, however, this is impossible to verify in practice (Hippke 2018).

Astronomical spectra are typically sampled at equal wavelength intervals, $\lambda = c/f$ with $\Delta\lambda = \text{const.}$ For a search of signals with constant temporal periods, these spectra need to be converted to equal frequency intervals $f = c/\lambda$ where $\Delta f = \text{const.}$ Afterwards, the overlaying blackbody-like flux shape needs to be subtracted out in order to perform a period search. It is adequate to use a sliding median with a boxcar window size of a few percent. Typically, sizes of 1–5 % give very similar results, with only a slight variation in the relative amplitudes of the peaks in the resulting Spectral Fourier transform. The SFT then produces a spectrum with power for signals with a constant temporal period, e.g., repeating laser pulses.

As the SFT method is not intuitive and requires careful implementation, I provide an open-source Python

arXiv:1901.00523v1 [astro-ph.IM] 2 Jan 2019




Figure 1. (a): Spectral Fourier Transform of the synthetic solar spectrum by Kurucz (2005a,b). A sliding median with a boxcar width of 3% (red line) is fitted to the spectral flux (black line). (b): After subtracting the median from the flux, a Fourier transformation is calculated (black). The highest peak coincides with one of the claimed pulse spacings (dashed red). (c): The median-subtracted flux (black) is shown together with the best-fit sine derived from the FFT. (d): A zoom to show the fit of the sine to the data.

code¹ which reads spectra, performs the transformation, and creates Figure 1.

3. RESULTS: THE ORIGIN OF THE SIGNALS

3.1. Synthetic spectrum

To determine whether the signals are inherent to the data reduction method, I check the synthetic solar spectrum by Kurucz (2005a,b) where the telluric lines have been removed. It has the advantage that it is essentially noise-free and without instrumental issues. I down-sample and crop the resolution from $R \approx 10^6$ between

¹ <http://github.com/hippke/pulses>. In the case of evenly spaced data without gaps, as is the case here, the Fourier transformation is equivalent to the Lomb-Scargle periodogram (Lomb 1976; Scargle 1981; VanderPlas 2018). The code uses the periodogram provided by Astropy (Astropy Collaboration et al. 2013; Price-Whelan et al. 2018), but delivers identical results when swapped for SciPy’s (Jones et al. 2001) FFT implementation.

300–1000 nm to match the $R = 4,000$ coverage between 380–920 nm of the SDSS survey, where the signals have originally been found. Following the recipe described in the previous section, I perform the spectral Fourier transform and show the result in Figure 1.

The highest spectral peak in the SFT corresponds exactly to one of the two claimed pulses at $\rho = 1.09 \times 10^{-13}$ s. The chance for this to happen by coincidence is 0.05%. As there are no artificial laser pulses in a synthetic solar spectrum, it appears highly likely that such signals commonly arise as part of the processing method.

The plot of the best-fitting sine found in the period search shows that its mean is slightly below the mean of the normalized spectrum, so that the peaks of the sine fit out the flux near its nominal values, and the bottoms of the sine fit out (more or less) the absorption lines.

3.2. Stellar spectra




Figure 2. (a): SFT of a solar spectrum and (b): a F9V star (TYC 2041-872-1) from SDSS found by [Borra & Trottier \(2016\)](#) to exhibit the signals. In both cases, a prominent peak is visible in the spectrum near one of the claimed periods.

Additionally, it can be shown that the signals arise in many high signal-to-noise ratio spectra of luminous bodies with absorption lines, and quite often with the highest, or one of the highest peaks, at exactly the same $\rho = 1.09 \times 10^{-13}$ s. Figure 2 (a) shows an SFT made from a high signal-to-noise solar spectrum². Here, the $\rho = 1.09 \times 10^{-13}$ s signal corresponds to the second highest peak when performing the same reduction as for the synthetic spectrum. The other claimed pulse spacing with $\rho = 1.64 \times 10^{-12}$ s coincides with a peak in the periodogram which is also significantly above the noise floor.

As a crosscheck, I have also re-reduced the SDSS spectrum of one of the stars for which [Borra & Trottier \(2016\)](#) found a signal (SDSS J160133.35+273355.2, Figure 2 (b)). This star was also re-observed using the optical Levy Spectrometer at the Automated Planet Finder (APF) without a detection of the signal ([Isaacson et al. 2018](#)). The APF spectrum with $R \approx 100,000$ was not downsampled to the $R = 4,000$ of the SDSS survey. Processing variations and different noise levels plausibly explain a non-detection of the signal.

Stars of different stellar type and metallicity exhibit different absorption lines. The power of the corresponding pulse spacings ρ in the periodogram will show similar variations. Further exploration would likely not yield additional insights and may be taken as an exercise in numerology.

3.3. Galaxy spectra

² Taken with the solar spectrometer installed by the University of Liège at the International Scientific Station of the Jungfraujoch. Available at http://bass2000.obspm.fr/solar_spect.php

[Borra \(2013\)](#) reports a correlation of signals with redshift and argues that this excludes instrumental and data processing effects as a source. This is incorrect: the integrated flux of galaxies originates from starlight, and thus comprises the same spectral line spacings, although stacked and redshifted.

As an independent validation, I have randomly selected 12,000 galactic spectra ($\sim 10\%$) from the 6df survey ([Jones et al. 2004](#)), DR3 ([Jones et al. 2009](#)). Data were taken with the Anglo-Australian Observatory’s UK Schmidt Telescope at a wavelength range $400 < \lambda < 750$ nm at $R \sim 1,000$, similar to SDSS, but with completely independent hardware, and placed in a different location.

Each galactic spectrum was converted to SFT using the method described in section 2 and in [Borra \(2012\)](#). Redshifts were taken from [Jones et al. \(2010\)](#) obtained through cross-correlation with template absorption-line spectra. SFTs were summed in redshift bins $\Delta z = 10^{-4}$, typically 10–100 SFTs per bin.

The signal at 1.09×10^{-13} s is clearly visible in the resulting Figure 3 (as 109×10^{-15} s for $z \rightarrow 0$). Additional features, undetected by [Borra \(2013\)](#), are also visible in the stack. While the vast majority of individual spectra show no significant ($\text{SNR} > 6$) peaks, it is apparent that all (or a large fraction of) galaxies contribute to these patterns. It appears that, very commonly, signals are overwhelmed by noise, but can be recovered by stacking many spectra. By comparing Figures 2 and 3, it is clear that a whole set of frequencies is detected in many stars and galaxies. This comb-like structure corresponds to a set of correlated distances of absorption lines (in units of wavelength) in the stellar and galactic spectra.

The statistical correlation of the stacked signals with redshift has been tested using linear regressions to the five strongest features for $0 < z < 0.2$ with a t-test.




Figure 3. SFT stack of 12,000 galaxies from the 6df survey. The stacked periodograms run from the bottom to the top in each redshift bin. The correlation of the peaks with redshift can be traced at least out to $z \sim 0.2$ until noise dominates. Consequently, the peaks appear at constant periods in the rest frame of the galaxies.

These yield positive slopes at $> 6\sigma$ confidence in each case, for a total significance (over noise) in excess of 30σ confidence. This is a clear verification that the features in question are not instrumental in nature.

4. DISCUSSION AND CONCLUSION

The discovery of artificial signals from outside our solar system may well require the discovery of new techniques, such as SFTs, or yet unknown tools. For this and

other goals, the development of new methods will prove invaluable. My refutation of this first SFT application does not mean that the method should be abandoned; instead it should be refined. Peaks caused by the non-random spacing of absorption lines can be flagged in the future.

A natural application for the SFT method could be the template-free estimation of galactic redshift from spectra. A follow-up work could analyze the feasibility of this approach. False-positives may arise; I have found several such cases. One is SDSS J130155.84+083631.6 with a SNR > 6 signal near 1.09×10^{-13} s, which was classified as a galaxy with redshift $z = 0.9703971$ by Jones et al. (2010) through cross-correlation with template absorption-line spectra but is really a carbon star (Green 2013). The outstanding advantage of SFTs is that they do not require dedicated SETI observations, but can be performed automatically with many existing spectroscopic data originally acquired for other purposes.

The signals in question can be found in galactic spectra from the 6df survey, as well as the SDSS survey, and show a correlation with redshift. Therefore, they can not be instrumental artifacts. However, their origin is not astrophysical. They can be found in SFTs of synthetic spectra, making them a processing effect from the non-random spacings of spectral absorption lines.

Acknowledgments I am thankful to Ermanno Borra, Brian Lacki, and Howard Isaacson for useful discussions. I thank the referee for their suggestion to examine the solar spectrum, which has led to the results described in this manuscript.

REFERENCES (0)

- Abbasi, R., Abdou, Y., Abu-Zayyad, T., et al. 2010, *Nuclear Instruments and Methods in Physics Research A*, 618, 139
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33
- Berkeley SETI Research Center team. 2016, Breakthrough Listen at UC Berkeley to conduct follow up observations of reported anomalous spectral features in solar type stars, https://seti.berkeley.edu/bl_sdss_seti_2016.pdf
- Boone, K., Aldering, G., Copin, Y., et al. 2018, *PASP*, 130, 064504
- Borra, E. F. 2010a, *ApJ*, 715, 589
- . 2010b, *A&A*, 511, L6
- . 2012, *AJ*, 144, 181
- . 2013, *ApJ*, 774, 142
- . 2017, *Journal of Astrophysics and Astronomy*, 38, 23
- Borra, E. F., & Trottier, E. 2016, *PASP*, 128, 114201
- Chin, S. L., François, V., Watson, J. M., & Delisle, C. 1992, *ApOpt*, 31, 3383
- Delisle, C., & St-Arnaud, J.-M. 1970, *Canadian Journal of Physics*, 48, 1214
- Desai, S., Mohr, J. J., Bertin, E., Kümmel, M., & Wetzstein, M. 2016, *Astronomy and Computing*, 16, 67
- Eisenstein, D. J., Weinberg, D. H., Agol, E., et al. 2011, *AJ*, 142, 72
- Fowler, A. 1924, *JRASC*, 18, 373
- Green, P. 2013, *ApJ*, 765, 12
- Hippke, M. 2018, *Journal of Astrophysics and Astronomy*, 39, 74
- Isaacson, H., Siemion, A. P. V., Marcy, G. W., et al. 2018, arXiv e-prints, arXiv:1812.02258
- Jones, D. H., Saunders, W., Colless, M., et al. 2004, *MNRAS*, 355, 747

- Jones, D. H., Read, M. A., Saunders, W., et al. 2009, [MNRAS](#), **399**, 683
- . 2010, VizieR Online Data Catalog, J/MNRAS/399/683
- Jones, E., Oliphant, T., Peterson, P., et al. 2001, SciPy: Open source scientific tools for Python, [Online; accessed 27 Nov 2018]
- Kurucz, R. L. 2005a, Memorie della Societa Astronomica Italiana Supplementi, **8**, 189
- . 2005b, Memorie della Societa Astronomica Italiana Supplementi, **8**, 73
- Lomb, N. R. 1976, [Ap&SS](#), **39**, 447
- Mandel, L., & Wolf, E. 1976, [JOSA](#), **66**, 529
- Price-Whelan, A. M., Sipócz, B. M., Günther, H. M., et al. 2018, [AJ](#), **156**, 123
- Scargle, J. D. 1981, [The Astrophysical Journal Supplement Series](#), **45**, 1
- VanderPlas, J. T. 2018, [The Astrophysical Journal Supplement Series](#), **236**, 16
- Yanny, B., Rockosi, C., Newberg, H. J., et al. 2009, [AJ](#), **137**, 4377