

MINIMAL HYPERSURFACES WITH ARBITRARILY LARGE AREA

OTIS CHODOSH AND CHRISTOS MANTOULIDIS

ABSTRACT. For $3 \leq n \leq 7$, we prove that a bumpy closed Riemannian n -manifold contains a sequence of connected embedded closed minimal surfaces with unbounded area.

1. INTRODUCTION

The goal of this note is to prove the following result:

Theorem 1.1. *For $3 \leq n \leq 7$, suppose that (M^n, g) is a connected, closed Riemannian n -manifold with a bumpy metric. Then, there is a sequence $\Sigma_j \subset (M^n, g)$ of connected, embedded, closed minimal surfaces with $\text{area}(\Sigma_j) \rightarrow \infty$.*

We recall here the following standard definition:

Definition 1.2. *We say that a metric g on a Riemannian manifold M^n is bumpy if there is no immersed closed minimal hypersurface Σ^{n-1} with a non-trivial Jacobi field.*

By work of White [Whi91, Whi17], bumpy metrics are generic in the sense of Baire category. Here, “generic” will always mean in the Baire category sense.

The key quantifier in Theorem 1.1 is *connected*. Indeed, thanks to the resolution of Marques–Neves’s multiplicity-one conjecture by the authors [CM18] for $n = 3$ and recently by Zhou [Zho19] for $3 \leq n \leq 7$, if one does not require that the Σ_j are connected, then Theorem 1.1 would be an immediate consequence of the following statement concerning the existence of minimal surfaces Σ_p realizing the p -widths (note that if (M^n, g) does not satisfy the Frankel property, then it is not clear that the number of connected components of Σ_p is uniformly bounded):

Theorem 1.3 ($n = 3$ [CM18], $3 \leq n \leq 7$ [Zho19, MN19]). *For $3 \leq n \leq 7$, suppose that (M^n, g) is a closed Riemannian n -manifold with a bumpy metric. Then, there is a constant $C = C(M, g) > 0$ so that for each positive integer p , there is a smooth embedded closed minimal surface Σ_p so that*

- *each component of Σ_p is two-sided,*
- *the area of Σ_p satisfies $C^{-1}p^{\frac{1}{n}} \leq \text{area}_g(\Sigma_p) \leq Cp^{\frac{1}{n}}$, and*
- *the index of Σ_p is satisfies $\text{ind}(\Sigma_p) = p$.*

We emphasize that the p -widths were introduced by Gromov, Guth, and Marques–Neves [Gro03, Gut09, MN17] in the Almgren–Pitts setting (as considered in the work of Zhou [Zho19]) and were understood in the Allen–Cahn setting (as considered in the work of the authors [CM18]) by Gaspar–Guaraco [Gua18, GG18a].

Recently, there has been exciting progress on the existence and behavior of min-max minimal hypersurfaces in Riemannian manifolds; besides those works already mentioned, we point out

that Liokumovich–Marques–Neves [LMN18] have proven a Weyl law for the p -widths. This a key component of the proof by Irie–Marques–Neves that minimal surfaces are generically dense [IMN18]. These results were extended to the Allen–Cahn setting by Gaspar–Guaraco [GG18b]. Marques–Neves–Song have proven that generically there is an equidistributed set of minimal hypersurfaces [MNS19]. Song has proven Yau’s conjecture on infinitely many minimal surfaces in all cases [Son18]. Finally, we note that Zhou–Zhu’s min-max theory for prescribed mean curvature [ZZ18] was instrumental in the proof of Zhou’s multiplicity-one result.

Theorem 1.1 is a consequence of the following statement:

Theorem 1.4. *For $3 \leq n \leq 7$, suppose that (M^n, g) is a connected Riemannian n -manifold with a bumpy metric. Then, either:*

- (1) *there exists a sequence of connected closed embedded stable minimal hypersurfaces $\Sigma_j \subset (M^n, g)$ with $\text{area}_g(\Sigma_j) \rightarrow \infty$, or*
- (2) *the hypersurfaces Σ_p from Theorem 1.3 have at least one connected component Σ'_p with $\text{area}_g(\Sigma'_p) \geq Cp^{\frac{1}{n}}$ for some $C = C(M, g) > 0$ independent of p .*

We note that for (M^3, g) with a bumpy metric of positive scalar curvature, the first case never occurs by [CKM17, Car17] and moreover we can conclude that $\text{ind}(\Sigma'_p) \rightarrow \infty$. On the other hand, Colding–Minicozzi have shown that any 3-manifold admits a (bumpy) metric in which the first alternative occurs [CM00].

We further note that in [CM18] we have shown how to use the monotonicity formula to find a component Σ''_p of Σ_p with $\text{genus}(\Sigma''_p) \geq C^{-1} \text{ind}(\Sigma''_p) \geq p^{\frac{2}{3}}$ (the genus bound follows from the work of Ejiri–Micallef [EM08]). The same argument applies in higher dimensions to find $\Sigma''_p \subset (M^n, g)$ with $\text{ind}(\Sigma''_p) \geq Cp^{1-\frac{1}{n}}$. It is not clear to us, however, that Σ'_p and Σ''_p can be taken to be the same.

1.1. Acknowledgments. O.C. was supported in part by an NSF grant DMS-1811059. He is grateful to Simon Brendle and Davi Maximo for several interesting conversations concerning the behavior of the Σ_p surfaces. C.M. would like to thank Yevgeny Liokumovich for interesting conversations on Almgren–Pitts theory. Both authors would like to thank André Neves for a helpful discussion about this work. This work originated during the authors’ visit to the International Centre for Mathematical Sciences (ICMS) in Edinburgh during the “Geometric Analysis Workshop” during the summer of 2018, which they would like to acknowledge for its support.

2. PROOF OF THEOREM 1.4

Lemma 2.1. *For $3 \leq n \leq 7$, consider (M^n, g) a closed Riemannian n -manifold with a bumpy metric. Then for $A_0 > 0$, there are finitely many closed embedded stable minimal hypersurfaces with area at most A_0 .*

Proof. This follows from curvature estimates [Sch83, SSY75, SS81] and the proof of [CM00] (see also the proof of [Sha17]). Namely, if there is a sequence Σ_j of stable embedded minimal hypersurfaces with uniformly bounded area, then they have uniformly bounded curvature. Thus, after passing to a subsequence, they converge smoothly (possibly with multiplicity) to a stable minimal hypersurface Σ_∞ . Passing to a double cover if Σ_∞ is one-sided, we can thus find—for j sufficiently large—graphs $u_j : \Sigma_\infty \rightarrow \mathbf{R}$ so that the exponential normal graph of u_j is some component of Σ_j .

We have that $u_j \rightarrow 0$ in $C^\infty(\Sigma_\infty)$. After normalizing by $\|u_j\|_{C^{2,\alpha}(\Sigma_\infty)}$, it is standard to pass to a further subsequence to find a non-trivial Jacobi field on (a cover of) Σ_∞ . This contradicts the bumpyness of (M^n, g) .

Alternatively, the implicit function theorem guarantees that any strictly stable minimal surface has a (strictly) mean convex tubular neighborhood (as before, passing to a double cover if the surface is one-sided). No minimal surface (besides the original one) can be completely contained in this neighborhood by the maximum principle. However, if there were infinitely many embedded stable minimal surfaces with bounded area, then curvature estimates and area bounds would allow one to find a sequence of such surfaces converging (possibly with multiplicity) to a limiting such surface. This cannot occur by considering the aforementioned mean convex neighborhood. \square

Lemma 2.2. *For $3 \leq n \leq 7$, consider (M^n, g) a connected closed Riemannian n -manifold. Assume there are at most N stable minimal surfaces in (M^n, g) . Then M contains at most $\max\{2N, 1\}$ disjoint closed embedded unstable minimal hypersurfaces.*

Proof of Lemma 2.2. Suppose there are $Q \geq 2$ disjoint unstable embedded minimal hypersurfaces $\Sigma_1, \dots, \Sigma_Q$. Consider

$$\tilde{M} := M \setminus \bigcup_{i=1}^Q \Sigma_i.$$

We can give \tilde{M} the structure of a compact Riemannian manifold with boundary $\partial\tilde{M}$ equal to the Σ_i (or their two-sided double cover if they were one-sided originally). Note that each component of $\partial\tilde{M}$ is unstable (for the components that were originally two-sided, this is clear; for the one-sided surfaces this follows by lifting an unstable variation to the double cover). Write $\tilde{M} = \tilde{M}_1 \cup \tilde{M}_2$, where \tilde{M}_1 is the union of the connected components of \tilde{M} with exactly one boundary component and \tilde{M}_2 is the union of components of \tilde{M} with at least 2 boundary components.

We claim that $\partial\tilde{M}_2$ has at least Q components. Indeed, consider some Σ_i . Suppose first that Σ_i is two sided and separating. Consider Ω_\pm the components of \tilde{M} containing Σ_i in their boundary. Suppose that $\Omega_- \cup \Omega_+ \subset \tilde{M}_1$. In this case, it is clear that $Q = 1$ and Σ_i was the only unstable surface in the family, contradicting our assumption that $Q \geq 2$. Thus, at least one of the Ω_\pm is contained in \tilde{M}_2 . This determines at least one element of $\partial\tilde{M}_2$. Now, suppose that Σ_i is two-sided but not separating. It is clear that any component of $\partial\tilde{M}$ associated to Σ_i cannot bound a component of \tilde{M}_1 (otherwise Σ_i would be separating). Finally, if Σ_i was one-sided and some component of $\partial\tilde{M}$ bounded a component of \tilde{M}_1 , then as before, we would have $Q = 1$, a contradiction.

Given this, Lemma 2.2 is a consequence of the following claim:

Claim. For $3 \leq n \leq 7$, consider (\check{M}^n, g) a connected compact Riemannian manifold with $J \geq 2$ boundary components that are all unstable minimal hypersurfaces. Then, there are at least $J/2$ closed embedded stable minimal surfaces in the interior of \check{M} .

Proof of the claim. Write the components of $\partial\check{M}$ as $\Gamma_1, \dots, \Gamma_J$. Because $J \geq 2$,

$$[\Gamma_i] \neq 0 \in H_{n-1}(\check{M}; \mathbf{Z}_2).$$

Find $\Gamma'_1 \in [\Gamma_1]$ minimizing area in the homology class. Because each component of the boundary is an unstable minimal surface, Γ'_1 is contained in the interior of \check{M} . There is thus an open set $\check{\Omega}_1 \subset \check{M}$ so that $\partial\check{\Omega}_1 = \Gamma_1 \cup \Gamma'_1$. Set $\check{M}_1 := \check{M} \setminus \check{\Omega}_1$. This manifold has $J - 1$ unstable boundary components and at least 1 stable boundary component. For Γ_2 one of the unstable boundary

components minimize area in homology to get Γ'_2 . Each component of Γ'_2 is contained in the interior of \check{M}_1 or coincides with one of the stable components of $\partial\check{M}_1$. In either case, we can repeat this process J times. Consider the disjoint stable minimal surfaces $\Gamma'_1, \dots, \Gamma'_J$ in \check{M} . We also note that, by construction, the $\check{\Omega}_1, \dots, \check{\Omega}_J$ are all disjoint.

We would like to estimate the number of connected components of $\cup_{j=1}^J \Gamma'_j$ (note that it is not necessarily true that the Γ'_j are pairwise disjoint, but at any point of intersection of Γ'_j and Γ'_k , the two surfaces locally agree, by construction). Consider a connected component $\Gamma \subset \cup_{j=1}^J \Gamma'_j$. There is some $a \in \{1, \dots, J\}$ so that $\Gamma \subset \partial\check{\Omega}_a$. Moreover, there is at most one other index $b \in \{1, \dots, J\} \setminus \{a\}$ so that $\Gamma \subset \partial\check{\Omega}_b$ (this holds because the $\check{\Omega}_j$ are pairwise disjoint). Finally, for any Γ_j , we can find such a component Γ (possibly multiple such components). Choosing such a component Γ corresponding to each Γ_j , we have just seen that a component can appear at most twice in this process. Thus we have found at least $J/2$ components, proving the claim. \square

This completes the proof of Lemma 2.2. \square

Proof of Theorem 1.4. If the first possibility fails, then there is a uniform area bound for embedded stable minimal hypersurfaces. Thus, Lemma 2.1 implies that there are finitely many stable embedded minimal hypersurfaces in (M^n, g) . Call this number N . Applying Lemma 2.2, we find that there are at most $N + \max\{2N, 1\}$ pairwise disjoint embedded minimal surfaces in (M, g) . Applied to Σ_p , there is at least one component Σ'_p with $\text{area}(\Sigma'_p) \geq Cp^{\frac{1}{n}}$. \square

REFERENCES

- [Car17] Alessandro Carlotto. Generic finiteness of minimal surfaces with bounded Morse index. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 17(3):1153–1171, 2017.
- [CKM17] Otis Chodosh, Daniel Ketover, and Davi Maximo. Minimal hypersurfaces with bounded index. *Invent. Math.*, 209(3):617–664, 2017.
- [CM00] Tobias H. Colding and William P. Minicozzi, II. Embedded minimal surfaces without area bounds in 3-manifolds. In *Geometry and topology: Aarhus (1998)*, volume 258 of *Contemp. Math.*, pages 107–120. Amer. Math. Soc., Providence, RI, 2000.
- [CM18] Otis Chodosh and Christos Mantoulidis. Minimal surfaces and the Allen–Cahn equation on 3-manifolds: index, multiplicity, and curvature estimates. <https://arxiv.org/abs/1803.02716>, 2018.
- [EM08] Norio Ejiri and Mario Micalef. Comparison between second variation of area and second variation of energy of a minimal surface. *Adv. Calc. Var.*, 1(3):223–239, 2008.
- [GG18a] Pedro Gaspar and Marco A. M. Guaraco. The Allen-Cahn equation on closed manifolds. *Calc. Var. Partial Differential Equations*, 57(4):Art. 101, 42, 2018.
- [GG18b] Marco A. M. Guaraco and Pedro Gaspar. The Weyl Law for the phase transition spectrum and density of limit interfaces. *To appear in Geom. Funct. Anal.*, <https://arxiv.org/abs/1804.04243>, 2018.
- [Gro03] M. Gromov. Isoperimetry of waists and concentration of maps. *Geom. Funct. Anal.*, 13(1):178–215, 2003.
- [Gua18] Marco A. M. Guaraco. Min–max for phase transitions and the existence of embedded minimal hypersurfaces. *J. Differential Geom.*, 108(1):91–133, 2018.
- [Gut09] Larry Guth. Minimax problems related to cup powers and Steenrod squares. *Geom. Funct. Anal.*, 18(6):1917–1987, 2009.
- [IMN18] Kei Irie, Fernando Marques, and André Neves. Density of minimal hypersurfaces for generic metrics. *Ann. of Math. (2)*, 187(3):963–972, 2018.
- [LMN18] Yevgeny Liokumovich, Fernando Marques, and André Neves. Weyl law for the volume spectrum. *Ann. of Math. (2)*, 187(3):933–961, 2018.
- [MN17] Fernando C. Marques and André Neves. Existence of infinitely many minimal hypersurfaces in positive Ricci curvature. *Invent. Math.*, 209(2):577–616, 2017.
- [MN19] Fernando C. Marques and André Neves. Morse index of multiplicity one min-max minimal hypersurfaces. <https://arxiv.org/abs/1803.04273>, 2019.

- [MNS19] Fernando C. Marques, André Neves, and Antoine Song. Equidistribution of minimal hypersurfaces for generic metrics. *To appear in Invent. Math.*, <https://arxiv.org/abs/1712.06238>, 2019.
- [Sch83] Richard Schoen. Estimates for stable minimal surfaces in three-dimensional manifolds. In *Seminar on minimal submanifolds*, volume 103 of *Ann. of Math. Stud.*, pages 111–126. Princeton Univ. Press, Princeton, NJ, 1983.
- [Sha17] Ben Sharp. Compactness of minimal hypersurfaces with bounded index. *J. Differential Geom.*, 106(2):317–339, 2017.
- [Son18] Antoine Song. Existence of infinitely many minimal hypersurfaces in closed manifolds. <https://arxiv.org/abs/1806.08816>, 2018.
- [SS81] Richard Schoen and Leon Simon. Regularity of stable minimal hypersurfaces. *Comm. Pure Appl. Math.*, 34(6):741–797, 1981.
- [SSY75] Richard Schoen, Leon Simon, and Shing-Tung Yau. Curvature estimates for minimal hypersurfaces. *Acta Math.*, 134(3-4):275–288, 1975.
- [Whi91] Brian White. The space of minimal submanifolds for varying Riemannian metrics. *Indiana Univ. Math. J.*, 40(1):161–200, 1991.
- [Whi17] Brian White. On the bumpy metrics theorem for minimal submanifolds. *Amer. J. Math.*, 139(4):1149–1155, 2017.
- [Zho19] Xin Zhou. On the multiplicity one conjecture in min-max theory. <https://arxiv.org/abs/1901.01173>, 2019.
- [ZZ18] Xin Zhou and Jonathan J. Zhu. Existence of hypersurfaces with prescribed mean curvature I - generic min-max. <https://arxiv.org/abs/1808.03527>, 2018.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08544

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON, NJ 08540

E-mail address: ochodosh@math.princeton.edu

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139

E-mail address: c.mantoulidis@mit.edu