# A MULTI-YEAR SEARCH FOR TRANSITS OF PROXIMA CENTAURI. II: NO EVIDENCE FOR TRANSIT EVENTS WITH PERIODS BETWEEN 1–30 DAYS

DAX L. FELIZ<sup>1,2,3</sup>, DAVID L. BLANK<sup>4</sup>, KAREN A. COLLINS<sup>5</sup>, GRAEME L. WHITE<sup>4</sup>, KEIVAN G. STASSUN<sup>1,2</sup>, IVAN A. CURTIS<sup>6</sup>, RHODES HART<sup>4</sup>, JOHN F. KIELKOPF<sup>7</sup>, PETER NELSON<sup>8</sup>, HOWARD RELLES<sup>5</sup>, CHRISTOPHER STOCKDALE<sup>9</sup>, BANDUPRIYA JAYAWARDENE<sup>4</sup>, PAUL SHANKLAND<sup>10</sup>, DANIEL E. REICHART<sup>11</sup>, JOSHUA B. HAISLIP<sup>11</sup>, AND VLADIMIR V. KOUPRIANOV<sup>11</sup>

Draft version May 18, 2022

## ABSTRACT

Using a global network of small telescopes, we have obtained light curves of Proxima Centauri at 329 observation epochs from 2006 to 2017. The planet Proxima b discovered by Anglada-Escudé et al. (2016) with an orbital period of 11.186 d has an *a priori* transit probability of  $\sim 1.5\%$ ; if it transits, the predicted transit depth is about 5 millimagnitudes. In Blank et al. (2018), we analyzed 96 of our light curves that overlapped with predicted transit ephemerides from previously published tentative transit detections, and found no evidence in our data that would corroborate claims of transits with a period of 11.186 d. Here we broaden our analysis, using 262 high-quality light curves from our data set to search for any periodic transit-like events over a range of periods from 1 to 30 d. We also inject a series of simulated planet transits and find that our data are sufficiently sensitive to have detected transits of 5 millimagnitude depth, with recoverability ranging from  $\sim 100\%$  for orbital period of 1 d to  $\sim 20\%$  for orbital period of 20 d for the parameter spaces tested. The recoverability specifically at the 11.186 d period of Proxima b is  $\sim 40\%$  for the parameter spaces tested. While we are able to rule out virtually all transits of other planets at periods shorter than 5 d and depths greater than 5 millimagnitudes, we cannot rule out the possibility that transits of Proxima b are present but escaped detection. However, we are able to rule out virtually all transits of other planets at periods shorter than 5 d and depths greater than 3 millimagnitudes.

Subject headings: planetary systems - stars: individual (Proxima Centauri) - techniques: photometric

## 1. INTRODUCTION

The discovery of Proxima Centauri b (Proxima b, hereafter) via the radial velocity (RV) technique by Anglada-Escudé et al. (2016) was a landmark event in exoplanet studies. We now know that orbiting in the habitable zone (Kopparapu et al. 2013) of the star nearest to our Sun is a planet that is likely to be rocky (Brugger et al. 2016; Kane et al. 2017; Bixel & Apai 2017) and possibly habitable (Ribas et al. 2016; Barnes et al. 2016; Meadows et al. 2016; Turbet et al. 2016; Boutle et al. 2017). We report here further results from our transit search of Proxima Centauri from 2006 to 2017 (Blank et al. 2018, Paper I hereafter) which was motivated by the possibility that such planets may exist, and that they could be found using sub-meter size telescopes with commercial grade CCD cameras.

In the 11 years of this photometric campaign, we collected light curves at 329 epochs. Of these 329 light curves, 262 passed various quality tests (detailed in Paper I), 96 of

<sup>1</sup> Department of Physics and Astronomy, Vanderbilt University, NashviÎle, TN 37235, UŠA

Department of Physics, Fisk University, 1000 17th Avenue North, Nashville, TN 37208, ÚSA

Corresponding Author, dax.feliz@vanderbilt.edu

<sup>4</sup> Centre for Astronomy, University of Southern Queensland, Australia <sup>5</sup> Harvard-Smithsonian Center for Astrophysics, Cambridge, MA

02138, USA ICO, Adelaide, South Australia

<sup>7</sup> Department of Physics and Astronomy, University of Louisville, Louisville, KY 40292, ÚSA

Ellinbank Observatory, Victoria, Australia

9 Hazelwood Observatory, Churchill, Victoria, Australia

<sup>10</sup> Lawrence Berkeley National Laboratory and Dept. of Physics, University of California Berkeley, Berkeley CA 94720-7300, USA

<sup>11</sup> Department of Physics and Astronomy, University of North Carolina at Chapel Hill, Campus Box 3255, Chapel Hill, NC 27599, USA

which overlapped with the previously published ephemerides of Anglada-Escudé et al. 2016, Kipping et al. 2017, Liu et al. 2017 and Li et al. 2017. A search for transits corresponding to Proxima b at these ephemerides is reported in Paper I. No convincing transit event attributable to Proxima b was detected in this subset of light curves.

In this work, we proceed to search all 262 quality light curves systematically for a planet of any period in the period range 1.01 to 30.5 days. In Section 2 we summarize the data collection, drawing reference to Paper I. We also describe our strategy for determining a period range to conduct our planet search. In Section 3 we describe our methods of analysis for algorithmically searching for periodic transit events and tests for statistical significance and sensitivity to detect transit events. Section 4 contains our results and the tests of sensitivity needed to place limits on any possible detection. We discuss our findings in Section 5.

# 2. OBSERVATIONS AND DATA COMPLETENESS

# 2.1. Summary of Observations and Data Reduction done in Blank et al. (2018)

The observations that make up our data set of 329 light curves came from the world-wide robotic telescope network Skynet (Reichart et al. 2005), the Real Astronomy Experience (RAE) robotic telescope (Fadavi et al. 2006) located in in Bickley, Western Australia, and from several participating observatories from the Kilodegree Extremely Little Telescope (KELT; Pepper et al. 2007, 2012) Follow-Up Network (KELT-FUN; Collins et al. 2018). More details about the participating instruments are in Table 2 in Paper I.

Our data reduction techniques are described in Section 3.2 of Paper I. To briefly summarize, in order to minimize the effects of long term stellar variability and differential chromatic airmass we performed a linear detrend so that the final mean flux value was 1.0. To remove obvious flares that are predicted to occur ~ 63 times per day (Davenport et al. 2016) and photometric outliers, we performed an iterative  $3\sigma$  clipping.

# 2.2. Geometric Transit Probability of Proxima b-Like Planets For A Given Period

To estimate the geometric transit probability of Proxima, we assumed a planet mass of 1.27  $M_{\oplus}$  and a planet radius of  $\sqrt{\delta}R_*$ , where  $\delta \sim 5$  millimagnitudes (or mmag) as reported by Anglada-Escudé et al. 2016 and then calculated the probability of a planet transiting as the fraction of the area of the celestial sphere that is swept out by the shadow of the planet during one orbital period (Borucki & Summers 1984; Winn 2011):

Transit Probability = 
$$\left(\frac{R_* + R_p}{a}\right) \left(\frac{1 + esinw}{1 - e^2}\right)$$
 (1)

Applying Kepler's Third Law, assuming an eccentricity e=0 and that  $R_p \ll R_*$  gives

$$\frac{R_*}{a} = R_* \left(\frac{G(M_p + M_*)P^2}{4\pi^2}\right)^{-1/3} \tag{2}$$

Adopting a stellar mass 0.1221  $M_{\odot}\,$  and stellar radius 0.1542  $R_{\odot}\,$  (Kervella et al. 2017), the geometric probability of transit detection for orbital periods from 0.01 to 365 days is shown in the top panel of Figure 1. It should be noted that this geometric estimation is based on the planet density model assumed by Anglada-Escudé et al. 2016 and that all transits are assumed to be across the face of the host star. Scaled curves of this type will have varied results from different assumptions of the density of the planet and from grazing transits.

### 2.3. Phase Coverage of Photometric Observations

To estimate the phase coverage of our data, we phase folded our data for each day in the period range of 0.01 - 365 days and then binned our data into 5 minute bins and calculated the inverse variance weighted means for each bin.

$$\hat{y} = \frac{\sum_{i} y_i / \sigma_i^2}{\sum_{i} 1 / \sigma_i^2} \tag{3}$$

We then define phase coverage as the number of finite values in our phase folded data bins divided by the total number of values in our phase folded data bins. The result of this procedure is shown in the bottom panel of Figure 1 where it is clear that the phase coverage falls below  $\sim 75\%$  for periods longer than 30 days. Both panels of Figure 1 show the increasing difficulty of detection of a transit event for longer periods. For period longer than 30 days, both the low sensitivity of detection and the poor phase coverage imply a low probability of detecting a transit, and we have decided to limit our search to the period region 1.01 to 30.5 days. A lower orbital period limit of 1.01 days was chosen to avoid potential aliases due to diurnal and sidereal day sampling.

# 3. METHODOLOGY

#### 3.1. Box-fitting Least-Squares Algorithm

The box-fitting least-squares (BLS) algorithm (Kovács et al. 2002) searches for periodic decreases in star brightness of a photometric time series. The BLS algorithm models



FIG. 1.— Top: The Geometric Transit Probability (GTP) of Proxima Centauri. At the orbital period of 11.186 days determined by the radial velocity discovery of Anglada-Escudé et al. (2016) the transit probability is  $\sim 1.5\%$  based on their assumed planet density model. Bottom: Phase Coverage of the 329 individual light curves phase folded and binned into 5 minute bins, as described in Section 2.3. The colored solid lines mark the periods where phase coverage is 100%, 75%, 50% and 25%. The phase coverage of our data drops below  $\sim 75\%$  around 30 days.

transit events as simple step functions and identifies transiting planet candidates by phase folding light curves to trial frequencies and searching a grid of transit epochs and durations at each frequency, and then picking the parameters that maximize the transit depth significance with a least-squares optimization. We use the VARTOOLS software (Hartman & Bakos 2016) to produce the BLS power spectrum shown in Figure 2. To determine the fractional transit length (transit duration divided by orbital period) q, we estimated a minimum value  $q_{min} = 0.017$  to account for transit of at least 25 minutes duration for a minimum orbital period of 1.01 days, and  $q_{max} = 0.1$  for events of ~ 3 days for a maximum orbital period of 30.5 days. With  $q_{min}$ , a range of desired frequencies  $(\frac{1}{P_{min}}, \frac{1}{P_{max}})$ , and the total duration of our observations (T) , we can roughly estimate the number of frequencies required for our BLS search:

$$N_{freg} = 4T \times (f_{max} - f_{min})/q_{min} \tag{4}$$

All parameters used in the BLS search are listed in Table 1.

#### 3.2. False Alarm Probability

To determine the statistical significance of peaks in the BLS power spectrum, we define a false alarm probability (FAP) to be the probability of a peak having equal strength by random chance or due to the cadence of our sampling. This is done by randomly rearranging the detrended fluxes and error information while keeping the time stamps fixed, reapplying the BLS search and recording the BLS outputs for the top peak of each iteration. This random permutation is then repeated 1,000 times. The 0.1% FAP is the highest peak out of 1,000 permutations, the 1% FAP is the 10th highest peak out of 1,000 permutations. In Section 4.2, we apply different variations of this definition of FAP.

### 3.3. Transit Injections

To test the sensitivity of the BLS algorithm in recovering transit-like events, we injected fake transits into our data and ran the BLS algorithm on the injected data sets. We simulated these fake transits using a given transit depth  $\delta$ , and orbital

 TABLE 1

 PARAMETERS USED IN BLS PERIODIC SEARCHES

$q_{mi}$	n	$q_{max}$	$P_{min}$	$P_{max}$	$N_{freq}$	$N_{bin}$	$d\!f$
0.01	7	0.1	1.01	30.5	920,492	120	$1.05\times 10^{-6}$

periods P, along with the stellar mass, stellar radius, the orbital eccentricity e, orbital inclination i and the argument of periastron w (all transit model parameters are listed in Table 2). With each permutation of our transit model parameters, we simulated a total of 550 Mandel-Agol transit models (Mandel & Agol 2002) using the Python package PyTransit, (Parviainen 2015), which in addition to the parameters listed in Table 2, also uses quadratic limb darkening coefficients. Using the EXOFAST<sup>1</sup>(Eastman et al. 2013) website to interpolate quadratic limb darkening coefficients from the limb darkening tables in Claret & Bloemen (2011), we were able to obtain  $u \sim [0.425, 0.298]$  for Proxima. After model creation, we then use BLS to test our ability to successfully detect our injected transit events, as described in Section 4.3.

#### 4. RESULTS

## 4.1. BLS Power Spectrum

Using the parameters from Table 1, we applied the VAR-TOOLS (Hartman & Bakos 2016) BLS algorithm on our combined and detrended 262 observations from the Skynet, KELT-FUN and RAE telescopes. We ran our BLS search from 1.01 to 30.5 days and found that all peaks in the power spectrum within orbital periods of 1.01 - 30.5 days fall below the majority of the calculated FAP and FAP(P) thresholds, as described in Section 4.2. The top peak of the BLS power spectrum corresponds to an orbital period of  $\sim 1.808$  days, which lies above the 10% FAP(P) threshold. As an example of the transit-like events detected by BLS in our data, Figure 3 displays the phase folded light curve of our data, using the orbital period and transit center time reported by the BLS algorithm. Figure 9 in the Appendix section, shows the 32 individual light curves that contribute to this signal. Although there is some evidence for a transit-like event in Figure 3, the existence of such an event is not supported by analysis of the individual light curves.

We note that there is no significant power in the BLS power spectrum at the orbital period determined by the radial velocity of Proxima b (Anglada-Escudé et al. 2016) which is consistent with our failure to find transits in Paper I. In section 4.2.3, we describe a methodology to detect low power peaks like those near the 11.186 day RV period that are displayed in the inset panel of Figure 2.

# 4.2. Testing Statistical Significance of BLS Power Spectra 4.2.1. FAP

In order for each power spectrum of the randomly permuted data sets to be on comparable scales, we normalized the power spectra using a modified version of the Spectroscopic Signal to Noise used in Hartman & Bakos (2016):

$$S/N(f) = \frac{SR(f)_{random} - SR(f)}{\sigma_{\overline{SR}}}$$
(5)

<sup>1</sup> http://astroutils.astronomy.ohio-state.edu/exofast/limbdark.shtml

where we use the average and standard deviations of the BLS reported Signal Residue, SR(f), of the power spectrum in Figure 2 with the SR(f) of the randomized data sets. Based on our definition of FAP in Section 3.2, we find FAP thresholds for the 0.1% FAP occurs at a S/N of ~ 7.61, 1% FAP at ~ 6.96 and the 10% FAP at ~ 6.35 as shown as horizontal black lines in Figure 2.

# 4.2.2. FAP As A Function of Period

To better assess the validity of peaks below the 0.1%, 1% and 10% FAP values, we divided  $N_{freq}$  into 20 period ranges, starting from 1.01 days and used Equation 4 to calculate the bounds of each subsequent period range so that each range contained the same number of frequencies.

$$P[i] = \frac{P[i-1]}{1 - P[i-1] \times (q_{min}/4T) \times (N_{free}/20)} \forall i \in \{1, ..., 20\}$$
(6)

These period ranges are displayed in Figure 4. For each period range, we then follow a similar process as our FAP procedure, where we randomly shuffle our data with fixed time stamps and then run a BLS search on the shuffled data. Within each period range, we then record information from the highest peak in the resulting BLS power spectra and repeat the process a total of 1,000 times. From these 1,000 permutations, we calculate the FAP thresholds within each period range which we refer to hereafter as FAP(P), which is shown as the red line-connected dots in Figure 2.

#### 4.2.3. Robust Estimation Of The Mean S/N

As a visually intuitive alternative to identifying potentially significant, low power, peaks like those that are near the RV orbital period of 11.186 days, we utilized the Python package StatsModels<sup>2</sup> module for Huber's robust estimator of scale and location (Huber 1981) to estimate the mean and standard deviation of the S/N of the power spectrum. By fixing the orbital period on these 12 low power peaks near the 11.186 day period, we then ran a BLS search to obtain parameters for transit center time, transit duration and transit depth. In Figure 5, we use these parameters to phase fold our data around these 12 orbital periods corresponding to peaks near the 11.186 day RV period, in addition to the 11.186 day RV period itself. We then carefully and critically examined curves all light curves that contribute to each of these 12 peaks. We find that on average, these peaks correspond to transit depths of  $\sim 0.73$ mmag and there are no consistent light curve events that display these periodic decreases in flux.

To apply this method as an additional transit detection criteria in Section 4.3, we calculated the robust estimations of mean and standard deviation of the transit injected power spectra S/N within each of the 20 defined period ranges described in Section 4.2.2 and shown in Figure 4. Using these robust statistics provides a better estimation for the location of the mean and standard deviation of S/N without rejecting outliers.

<sup>2</sup> https://www.statsmodels.org/stable/index.html

 TABLE 2

 Parameters used in Transit injection Analysis

Parameter	Value / Model	Citation
Stellar Radius	$0.1542~\mathrm{R}_{\odot}$	Kervella et al. (2017)
Stellar Mass	$0.1221~\mathrm{M}_{\odot}$	Kervella et al. (2017)
Effective Temperature	3042 K	Ségransan et al. (2003)
log g	5.20	Ségransan et al. (2003)
[Fe/H]	0.21	Schlaufman & Laughlin (2010)
Transit Depth $\delta$ (mmag)	1.0, 2.0, 3.0, 4.0, 5.0, 6.0, 7.5, 10.0, 15.0, 20.0	This work
Orbital Period (days)	1.1, 2.1, 3.1, 5.1, 7.6, 10.1, 11.186, 15.1, 20.1, 25.1, 30.1	This work
Orbital Phase	-0.4, -0.2, 0.0, 0.2, 0.4	This work
Planetary Radius	$\sqrt{\delta}R_*$	This work
Planetary Mass	$M_{\oplus}R_p/R_{\oplus}$	This work
Eccentricity	0.0	This work
Inclination	$\pi/2$	This work
Argument of Periastron	$\pi/2$	This work
Quadratic Limb darkening Coefficients	(0.425, 0.298)	This work, determined with EXOFAST (Eastman et al. 2013)



FIG. 2.— We present a power spectrum from the VARTOOLS BLS transit search algorithm. The black vertical solid lines represent the orbital periods corresponding to the top 4 peaks of the power spectrum. The horizontal black lines correspond to the 0.1%, 1%, and 10% FAP as described in Section 3.2. The red line-connected dots are the FAP(P) thresholds calculated in 20 period ranges with equal  $N_{freq}$  as described in Section 4.2.2. The green, orange, cyan and brown line-connected dots represent the robust estimations for the mean and mean plus the  $3\sigma$ ,  $5\sigma$  and  $7\sigma$  of the S/N in each period range as described in Section 4.2.3. The inset figure is a close up of peaks in the power spectrum that are near the 11.186 day RV period and are also above the mean +  $5\sigma$  and mean +  $7\sigma$  lines, marked with black dots.



FIG. 3.— From the power spectrum in Figure 2, we took the orbital period corresponding to the top peak and phase folded our combined 262 light curves. The grey dots are the unbinned data and the red dots are binned with 5 minute bin widths. The BLS model of this peak, shown as the blue line, has a transit depth and duration of  $\sim 5.28$  mmag and 43.4 minutes. We do not believe this to be a real transit event as shown in Figure 9 and explained in the Appendix section.



FIG. 4.— An illustration of the 20 period ranges used to define FAP(P). The edges of the ranges are calculated with Equation 6. Each colored shaded region is a different period range where each range has an even amount of frequencies,  $N_{freq}/20$ .

## 4.3. Transit Injection Recovery

To test the sensitivity of the BLS algorithm's transit detection ability on our data set, we injected 550 transit models into

5



FIG. 5.— From the inset panel of Figure 2, we have identified 12 peaks that have S/N values larger than the robust mean +  $5\sigma$  of the power spectrum's S/N within the corresponding period range. We then conducted a BLS search using a fixed period for each of these 12 peaks, in addition to the 11.186 day RV period, and phase folded our data with the corresponding transit center time output by the VARTOOLS BLS algorithm. In each panel of this figure, we have the phase folded light curves with the unbinned data as grey points, the binned data with 5 minute bins as red points and the BLS model as the blue lines. The black points are data from the UT April 11 2007 RAE light curve which is also in shown Figures 9 and discussed in the Appendix section. In each case, the transit depths reported by BLS are on the order of 1 mmag or smaller which is reflected by the relatively weak peak strength as seen in Figure 2.

our data as described in Section 3.3. We then ran the BLS algorithm using identical input parameters as in Table 1 and recorded the resulting power spectra and BLS reported transit parameters. In this analysis, a successfully recovered transit injection is defined as a peak in the power spectrum that is within  $\pm 1\%$  of the injected transit model's orbital period and has a BLS Power above a threshold. We use the FAP, FAP(P) and robust estimations of the mean plus standard deviation of the S/N as three separate thresholds to our ability to detect varying peak strengths in the transit injected power spectra.

We also considered harmonics and sub-harmonics (1/3, 1/2, 2 and 3 times) of the injected transit model periods in our detection criteria. In Figure 6, we apply our detection criteria to the 550 transit injections and perform BLS searches to recover the injected transits across multiple orbital phases. The numbers of the color bars shown in Figures 6, 7 and 8 represent the

number of orbital phases where a detection by BLS occurred, ranging from 0 (for no detections in any of the phases tested) to 5 (detections in all phases tested) for the FAP, FAP(P) and robust mean plus standard deviation thresholds, respectively.

As expected, the number of successfully recovered transit injections decrease for orbital periods beyond 15 days where we lose about 10% of our phase coverage as shown in Figure 1. For the injected transits with Proxima b's orbital period of 11.186 days and a transit depth of 5 mmag (highlighted by red and cyan colored boxes in Figures 6, 7 and 8) and higher, we successfully recover transit injections in at least 2 of the 5 orbital phases with and without the requirement of FAP, FAP(P) or robust mean plus standard deviation thresholds, respectively.

# 4.4. Constraints on Transiting Planet Properties

As a simple exercise, we also estimated which combinations of transit depth and orbital period could be detected by the Doppler semi-amplitude of Proxima b ( $\sim 1.4$  m/s) as reported by Anglada-Escudé et al. (2016). For a given planet radius ( $R_p = \sqrt{\delta}R_*$ ), we can define the planet mass as:

$$M_p = \left(\frac{\rho_p}{\rho_{\oplus}}\right) \left(\frac{R_p}{R_{\oplus}}\right)^3 M_{\oplus} \tag{7}$$

We assume a circular orbit and the planet density to be Earthlike ( $\rho_p/\rho_{\oplus} = 1$ ). By keeping the Doppler semi-amplitude (K) fixed,

$$K = \frac{2\pi \, a \sin i}{P\sqrt{1-e^2}} \tag{8}$$

applying Kepler's Third Law, and assuming a circular orbit (e=0), we can solve for the orbital period:

$$P = 2\pi G \left(\frac{M_p sin(i)}{K}\right)^3 \left(\frac{M_\odot}{M_*}\right)^2 \left(\frac{1}{1-e^2}\right)^{3/2} \quad (9)$$

In Figures 6, 7 and 8, we display this RV model over our transit injection recovery plots.

The RV model roughly follows our transit detection recoverability up to an orbital period  $\sim 25$  days and transit depths as low as  $\sim 6$  mmag. Each orbital period and transit depth cell in Figures 6, 7 and 8 that are above the line correspond to the transit depth and orbital period combinations that should be detectable in the RV data.

If Proxima b does transit and is not denser than Earth, then the planet would appear in a cell above the RV model; however our light curve data rule out any transit above the curve for orbital periods shorter than  $\sim 15$  days. For other rocky planets (with Earth-like density) that might exist in the system, they must be below the RV model in Figures 6, 7 and 8 or else their RV signatures would presumably have been detectable. While our light curve data does not generally probe the regions under the RV model, we are able to rule out detectable transit events for orbital periods  $\lesssim 5$  days and transit depths  $\gtrsim 3$  mmag.

# 5. DISCUSSION

Our BLS analysis confirms our work in Paper I that Proxima b does not transit its host star in our light curves and we find no evidence that Proxima b transits its host star with a depth greater than 3 mmag with any other orbital period in the period range of 1 to 30 days. In Paper I, we describe the selection criteria based on the amount of scatter in individual light curves that led to 262 of our 329 light curves to be included in our overall analysis. As shown in Figure 1 of Blank et al. (2018), the median standard deviation of individual unbinned light curves after the detrended and vetting processes is  $\sim 0.52\%$  which is similar to the expected 0.5% transit depth for Proxima b. For the minimum mass of 1.27  $M_\oplus$  estimated by Anglada-Escudé et al. (2016), smaller radii of Proxima b would translate to higher planet densities.

Brugger et al. (2016) determine the ranges of planet mass and radius of Proxima b to be (1.1 -1.46)  $M_{\oplus}$  and (0.94 -1.40)  $R_{\oplus}$ . In the case of the 1.1  $M_{\oplus}$ , 0.94  $R_{\oplus}$  model of Proxima b, 65% of the planets mass is located in the core and the remaining 35% as part of the mantle. In the 1.46  $M_{\oplus}$ , 1.4  $R_{\oplus}$  case, the corresponding composition is 50% of the planets mass being in the form of water and the remaining 50% in the mantle. We estimate a planet radius and mass of  $\sim 0.94$   $R_\oplus$  and  $\sim 1.1~M_\oplus$  corresponds to a transit depth of about 3 mmag and a planet density  $1.32~\rho_\oplus$  while a planet of radius 1.4  $R_\oplus$  and mass 1.46  $M_\oplus$  corresponds to a transit depth of about 7 mmag and a planet density of  $\sim 0.5~\rho_\oplus$ .

To determine our lower limit for detectable planet densities, we have extended our exercise Section 4.4 to also fix the planet mass to 1.27  $M_{\oplus}$  in addition to the fixed Doppler semi-amplitude of 1.4 m/s. Using Equation 9 gives an orbital period ~13.09 days. The lowest recovered injected transit depth near that orbital period in Figures 6 and 7 is ~ 3 mmag which corresponds to a planet radius ~0.92 R<sub> $\oplus$ </sub>. This results in a planet density of ~1.63  $\rho_{\oplus}$  which is below the minimum density of ~2.07  $\rho_{\oplus}$  estimated from the lower bounds of the probabilistically constrained result from Bixel & Apai (2017) for their rocky planet model.

Lovd et al. (2018) discusses the mechanisms of flares potentially inducing photoionization heating of the upper atmosphere of planets orbiting very near their host stars. Through the authors' work, they determined that the extreme ultraviolet radiation during flare events can be intense enough to drive hydrodynamic escape. Howard et al. (2018) detected a super flare where Proxima's optical flux increased by 68 times in an hour long event. Through the two years of observations with the Evryscope telescope, they observed 23 other large flare events and determined that super flares of this scale occur roughly 5 times per year, and that this level of repeated flaring may be sufficient enough to reduce the ozone of an Earth-like atmosphere by 90% within five years and complete depletion may occur within several hundred thousand years. In the event of Proxima b's planetary radius being smaller due to a different planet density or decreasing due to flare driven atmospheric loss, our data should have a sensitivity to transit depths up to  $\sim 0.5\%$ , supported by the 5 minute binned RMS of our data being  $\sim 0.26\%$ .

There are several areas for improvement within our data reduction pipeline. It is likely that our data have residual correlated noise after our detrending process and there are a many ways to approach modeling the noise (e.g. Gaussian Processes). In addition to scatter due to the intrinsic variability of the star, Proxima is also a well known flare star (Shapley 1951; Walker 1981). There are numerous flares within our data set that can be quantified (Jayawardene et al., in preparation) and modelled to subtract the events from our light curves to obtain a flatter baseline more suitable for transit detection. Another alternative to estimate the FAP is with Bayesian statistics rather than our Frequentist approach which assumes pure white noise.

#### 6. CONCLUSION

In this work, we present an analysis of 262 photometric observations of Proxima Centauri where we search for periodic transit-like events. The light curves have been cleaned and detrended as described in Paper I, and then fed into the box-fitting least squares (BLS) period finding algorithm. To estimate the statistical significance of the peaks in the BLS power spectrum, we estimated 0.1%, 1% and 10% false alarm probability (FAP) thresholds. We also determined FAP as a function of orbital period (FAP(P)) by calculating the FAP thresholds for 20 period ranges, each with an equal number of frequencies to be searched for by the BLS algorithm. To explore peaks in the BLS power spectra that have low power, we used Huber's robust estimator of scale and location to estimate the mean and standard deviation of the power spectrum



FIG. 6.— Applying the detection criteria described in Section 3.3 to our 550 transit injected data sets, we present a color map of transit detections that occurred in the orbital phases: -0.4, -0.2, 0, 0.2, 0.4. The upper left figure represents recovered transit injections within  $\pm 1\%$  of their injected periods but considering no FAP threshold. The upper right figure represents recovered transit injections within  $\pm 1\%$  of their injected periods and over the 10% FAP thresholds. Similarly the lower left and right figures represent recovered transit injections within  $\pm 1\%$  of their injected periods and above the 1%, 0.1% FAP thresholds, respectively.



FIG. 7.— Similarly to Figure 6, we altered our detection criteria to use the FAP(P) function described in Section 3.3. The upper left figure represents recovered transit injections within  $\pm 1\%$  of their injected periods but considering no FAP(P) threshold. The upper right figure represents recovered transit injections within  $\pm 1\%$  of their injected period and over the 10% FAP(P) thresholds. Similarly the lower left and right figures represent recovered transit injections within  $\pm 1\%$  of their injected period and over the 10% FAP(P) thresholds, respectively.



FIG. 8.— Similarly to Figure 6, we altered our detection criteria to use the robust estimation of mean,  $\sigma$  of the BLS power spectrum as described in Section 3.3. The upper left figure represents recovered transit injections that are within  $\pm 1\%$  of their injected periods and are over the robust mean of the injected power spectra. The upper right figure represents recovered transit injections within  $\pm 1\%$  of their injected periods and over the mean +  $3\sigma$  line. Similarly the lower left and right figures represent recovered transit injections within  $\pm 1\%$  of their injected periods and above the mean +  $3\sigma$  line. Similarly the lower left and right figures represent recovered transit injections within  $\pm 1\%$  of their injected periods and above the mean +  $5\sigma$  and mean +  $7\sigma$  lines, respectively.

in each of the 20 period ranges. The majority of the high-

est peaks of the BLS power spectrum fall below the FAP and

FAP(P) thresholds. We note that there is no significant power in the power spectrum near the orbital period of Proxima b (Anglada-Escudé et al. 2016); however we have identified 12 peaks that are above the robust mean plus 5 times the standard deviation. We then phase folded those 12 peaks and examined individual light curves that contribute to those periods. We conclude that these 12 peaks are unlikely to be caused by transit events. To test our sensitivity for detecting transit-like events, we injected 550 fake transits with parameters differing in transit depth, orbital phase and orbital period. We were able to detect most injections at transit depths 5 mmag and greater up to an orbital period of 11.186 days across multiple orbital phases. Overall, we were unable to confirm the existence of transits of Proxima b.

- Anglada-Escudé, G., Amado, P. J., Barnes, J., et al. 2016, Nature, 536, 437 Barnes, R., Deitrick, R., Luger, R., et al. 2016, ArXiv e-prints,
- arXiv:1608.06919
- Bixel, A., & Apai, D. 2017, ApJ, 836, L31
- Blank, D. L., Feliz, D., Collins, K. A., et al. 2018, ArXiv e-prints, arXiv:1802.04284
- Borucki, W. J., & Summers, A. L. 1984, Icarus, 58, 121 Boutle, I. A., Mayne, N. J., Drummond, B., et al. 2017, A&A, 601, A120 Brugger, B., Mousis, O., Deleuil, M., & Lunine, J. I. 2016, ApJ, 831, L16
- Claret, A., & Bloemen, S. 2011, A&A, 529, A75
- Collins, K. A., Collins, K. I., Pepper, J., et al. 2018, ArXiv e-prints, arXiv:1803.01869
- Davenport, J. R. A., Kipping, D. M., Sasselov, D., Matthews, J. M., & Cameron, C. 2016, ApJ, 829, L31 Eastman, J., Gaudi, B. S., & Agol, E. 2013, PASP, 125, 83
- Fadavi, M., Verveer, A., Aymon, J., et al. 2006, Astronomische Nachrichten, 327.811
- Hartman, J. D., & Bakos, G. Á. 2016, Astronomy and Computing, 17, 1 Howard, W. S., Tilley, M. A., Corbett, H., et al. 2018, ApJ, 860, L30 Huber, P. 1981, Robust Statistics (Wiley)

- Kane, S. R., Gelino, D. M., & Turnbull, M. C. 2017, AJ, 153, 52 Kervella, P., Thévenin, F., & Lovis, C. 2017, A&A, 598, L7 Kipping, D. M., Cameron, C., Hartman, J. D., et al. 2017, AJ, 153, 93
- Kopparapu, R. K., Ramirez, R., Kasting, J. F., et al. 2013, ApJ, 765, 131

In our upcoming Paper III (D.L. Feliz et al., in preparation), we intend to model the numerous flares in our data as well as the correlated noise to reduce the scatter in our light curves and conduct a more thorough period finding search.

The authors would also like to thank Joshua Pepper, Scott Gaudi, David Latham, Mike Lund, and Robert Siverd for their thoughtful discussions and input regarding our data reduction and analysis. Additionally, the authors would like to once again thank the science teams of RAE, Skynet and KELT-FUN for their contributions to our survey. Dax Feliz would also like to acknowledge support from the Fisk-Vanderbilt Masters-to-PhD Bridge Program.

#### REFERENCES

- Kovács, G., Zucker, S., & Mazeh, T. 2002, A&A, 391, 369
- Li, Y., Stefansson, G., Robertson, P., et al. 2017, ArXiv e-prints, arXiv:1712.04483
- Liu, H.-G., Jiang, P., Huang, X., et al. 2017, ArXiv e-prints, arXiv:1711.07018

- Loyd, R. O. P., France, K., Youngblood, A., et al. 2018, ApJ, 867, 71 Mandel, K., & Agol, E. 2002, ApJ, 580, L171 Meadows, V. S., Arney, G. N., Schwieterman, E. W., et al. 2016, ArXiv e-prints, arXiv:1608.08620
- Parviainen, H. 2015, MNRAS, 450, 3233
- Pepper, J., Kuhn, R. B., Siverd, R., James, D., & Stassun, K. 2012.
- Publications of the Astronomical Society of the Pacific, 124, 230 Pepper, J., Pogge, R. W., DePoy, D. L., et al. 2007, Publications of the Astronomical Society of the Pacific, 119, 923
- Reichart, D., Nysewander, M., Moran, J., et al. 2005, Nuovo Cimento C Geophysics Space Physics C, 28, 767 Ribas, I., Bolmont, E., Selsis, F., et al. 2016, A&A, 596, A111 Schlaufman, K. C., & Laughlin, G. 2010, A&A, 519, A105

- Ségransan, D., Kervella, P., Forveille, T., & Queloz, D. 2003, A&A, 397, L5
- Shapley, H. 1951, Proceedings of the National Academy of Science, 37, 15
- Turbet, M., Leconte, J., Selsis, F., et al. 2016, A&A, 596, A112
- Walker, A. R. 1981, MNRAS, 195, 1029
- Winn, J. 2011, Exoplanets, edited by S. Seager. Tucson, AZ: University of Arizona Press, 2011, 526 pp. ISBN 978-0-8165-2945-2, 55

# APPENDIX

To verify whether or not the top peak of the BLS power spectrum is due to a transit-like event, we phase folded our 262 light curves around the orbital period corresponding to the top peak which is  $\sim 1.808$  days. We then examined a subset of 32 light curves that contribute data points to within  $\pm 3$  hours of the transit center time reported by the BLS algorithm,  $T_C$ =2453880.516604185 BJD<sub>TDB</sub>. In Figure 9, the individual light curves are then binned with 5 minute bins and are vertically offset from one another. In Figure 11 of Paper I, we modeled the best fit transit model of the Skynet Prompt 2 UT 2014 May 14 light curve where we deemed it unlikely to be caused by a transiting exoplanet. The Skynet RCOP light curve from UT 2014 August 2 has a decrease in flux near phase 0 relative to  $T_C$  but this is due to the remaining points after our iterative  $3\sigma$  of a flare event that is followed by another smaller pair of flare events. The KELT-FUN Ivan Curtis Observatory light curves from UT 2017 March 7 and UT 2017 March 18 display decreases in flux near phase 0 in relation to  $T_C$ . In the UT 2017 March 7 light curve, the decrease at the end of the light curve are suspected to be due to atmospheric fluctuation but remained in our quality checks in Paper I. The UT March 18 2017 has a relatively high amount of scatter compared to the rest of our data but also passed our quality check due to the transit-like feature shown. In Figure 5, we highlighted data from the UT April 11 2007 RAE light curve which displays a ~20 mmag dip which we've chosen to include in our ensemble of light curves. We conclude that the that top peak of the BLS power spectrum is unlikely to be due to a transiting exoplanet as there is no consistent transit events in all other light curves near phase 0 relative to  $T_C$ .



FIG. 9.— These light curves correspond to the contributions in phase of Figure 3. Each light curve is phase folded around the orbital period and transit center time from the highest peak from the VARTOOLS BLS algorithm. The orbital period for this peak is 1.808 days and has a transit depth of 5.28 mmag. We vertically separated each light curves by an constant and alternated their colors for easier distinction between observation. Each light curve is binned at 5 minute intervals and contributes at least one data point within  $\pm 1$  hour of the transit center time.