

Attractors of Trees of Maps and of Sequences of Maps between Spaces with Applications to Subdivision

Nira Dyn, David Levin and Peter Massopust

Abstract. An extension of the Banach fixed point theorem for a sequence of maps on a complete metric space (X, d) has been presented in a previous paper. It has been shown that backward trajectories of maps $X \rightarrow X$ converge under mild conditions and that they can generate new types of attractors such as scale dependent fractals. Here we present two generalisations of this result and some potential applications. First, we study the structure of an infinite tree of maps $X \rightarrow X$ and discuss convergence to a unique “attractor” of the tree. We also consider “staircase” sequences of maps, that is, we consider a countable sequence of metric spaces $\{(X_i, d_i)\}$ and an associated countable sequence of maps $\{T_i\}$, $T_i : X_i \rightarrow X_{i-1}$. We examine conditions for the convergence of backward trajectories of the $\{T_i\}$ to a unique attractor. An example of such trees of maps are trees of function systems leading to the construction of fractals which are both scale dependent and location dependent. The staircase structure facilitates linking all types of linear subdivision schemes to attractors of function systems.

Mathematics Subject Classification (2010). Primary 47H10; Secondary 28A80, 41A30, 54E50.

Keywords. Fractals, subdivision schemes, fixed points, attractors, function systems.

1. Introduction and Preliminaries

In a recent paper [14], the authors investigated the relation between non-stationary subdivision and sequences of function systems (SFSS). This study introduced the notion of backward trajectories on a metric space and a related version of the Banach fixed point theorem for sequences of maps. With this theory, limits of non-stationary subdivision schemes with masks of fixed size are related to attractors of SFSS.

The present paper is motivated by an attempt to relate more general types of subdivision processes to SFSs. In order to explain these goals we start with a short introduction to the field of subdivision schemes.

1.1. Some Notions of Subdivision Schemes

Subdivision schemes play an important role in Computer-Aided Geometric Design (CAGD) and in wavelets theory [4, 9]. Here, we only consider univariate scalar binary subdivision schemes.

Given a set of control points $p^0 = \{p_j^0 \in \mathbb{R}^m : j \in \mathbb{Z}\}$ at level 0, a stationary binary subdivision scheme recursively defines new sets of points $p^k = \{p_j^k : j \in \mathbb{Z}\}$ at level $k \geq 1$ by the refinement rule

$$p_i^{k+1} = \sum_{j \in \mathbb{Z}} a_{i-2j} p_j^k, \quad k \geq 0, \quad (1.1)$$

or in short form,

$$\bar{p}^{k+1} = S_a \bar{p}^k, \quad k \geq 0, \quad (1.2)$$

The refinement is manifested by relating each point p_j^k at level k of the subdivision process to the binary parametric value $j2^{-k}$. The set of real coefficients $a = \{a_j : j \in \mathbb{Z}\}$ that determines the refinement rule is called the mask of the scheme. We assume that the support of the mask, $\sigma(a) := \{j \in \mathbb{Z} : a_j \neq 0\}$, is finite. In (1.2), S_a is a bi-infinite two-slanted matrix with entries $(S_a)_{i,j} = a_{i-2j}$, and \bar{p}^k is an infinite matrix whose rows are the points at level k , $(\bar{p}^k)_i = p_i^k \in \mathbb{R}^m$.

A non-stationary binary subdivision scheme is defined formally as

$$\bar{p}^{k+1} = S_{a^{[k]}} \bar{p}^k, \quad k \geq 0, \quad (1.3)$$

where the refinement rule at refinement level k is of the form

$$p_i^{k+1} = \sum_{j \in \mathbb{Z}^s} a_{i-2j}^{[k]} p_j^k, \quad i \in \mathbb{Z}, \quad (1.4)$$

i.e., in a non-stationary scheme, the mask $a^{[k]} := \{a_j^{[k]} : j \in \mathbb{Z}\}$ depends on the refinement level k .

The classical definition of a convergent subdivision scheme is the following.

Definition 1.1 (C^ν -Convergent Subdivision). A subdivision scheme is termed C^ν -convergent, $\nu \geq 0$, if for any initial data p^0 there exists a C^ν function $f : \mathbb{R} \rightarrow \mathbb{R}^m$ such that

$$\lim_{k \rightarrow \infty} \sup_{i \in \mathbb{Z}} |p_i^k - f(2^{-k}i)| = 0, \quad (1.5)$$

and for some initial data $f \neq 0$.

Remark 1.2.

The limit curve of a C^0 -convergent subdivision applied to initial data p^0 is denoted by $p^\infty = S_a^\infty p^0$. The function f in Definition 1.1 specifies a parametrization of the limit curve.

The analysis of subdivision schemes aims at studying the smoothness properties of the limit function f . For further reading, see [9].

A weaker type of convergence is obtained by using a set distance approach.

Definition 1.3 (*h-Convergent Subdivision*). A subdivision scheme is termed *h-convergent* if for arbitrary initial data p^0 there exists a set $p^\infty \subset \mathbb{R}^m$, such that

$$\lim_{k \rightarrow \infty} h(p^k, p^\infty) = 0, \quad (1.6)$$

where h is the Euclidian Hausdorff metric on \mathbb{R}^m . The set p^∞ is termed the *h-limit* of the subdivision scheme.

The relationship between curves and surfaces generated by stationary subdivision algorithms and self-similar fractals generated by iterated function systems (IFSs) was first presented in [18]. The work in [14] establishes a relation between non-stationary subdivision with a mask of fixed support size and sequences of function systems (SFSSs). The present paper is motivated by our goal to find the relationship between subdivision and fractals for an extended families of subdivision schemes such as non-stationary schemes with growing mask size and non-uniform schemes. It turns out that two new structures of sequences of maps are needed here. The first involves an infinite binary tree of maps on a metric space and the second a sequence of metric spaces $\{(X_i, d_i)\}_{i \in \mathbb{N}}$ together with a “staircase” type sequence of maps $\{T_i\}_{i \in \mathbb{N}}$, $T_i : X_{i+1} \rightarrow X_i$. For both structures, we extend the Banach fixed point theorem.

In Section 2, we study the structure of an infinite tree of maps (TMs). We consider convergence of backward trajectories along paths in the tree and introduce the notion of attractor of a tree. Staircase maps are analyzed in Section 3 and in Sections 4 and 5 we demonstrate the application of staircase trajectories to SFSSs related to non-stationary subdivision procedures, in particular that which generates the up-function [9]. In Section 6, staircase maps and trees of maps are applied to the analysis of linear non-uniform subdivision schemes.

The results in the present work are built upon the results in [14] which we briefly review below.

1.2. Sequences of Maps on $\{X, d\}$ and their Trajectories

Let (X, d) be a complete metric space. For a map $f : X \rightarrow X$, we define the Lipschitz constant associated with f by

$$\text{Lip}(f) = \sup_{x, y \in X, x \neq y} \frac{d(f(x), f(y))}{d(x, y)}.$$

A map f is said to be Lipschitz if $\text{Lip}(f) < +\infty$ and a contraction if $\text{Lip}(f) < 1$.

Now, consider a sequence of continuous maps $\{T_i\}_{i \in \mathbb{N}}$, $T_i : X \rightarrow X$.

Definition 1.4 (Forward and Backward Trajectories). The forward and backward trajectories of $\{T_i\}_{i \in \mathbb{N}}$ in X , starting from $x \in X$, are

1. $\Phi_k(x) = T_k \circ T_{k-1} \circ \cdots \circ T_1(x)$, $k \in \mathbb{N}$, respectively,
2. $\Psi_k(x) = T_1 \circ T_2 \circ \cdots \circ T_k(x)$, $k \in \mathbb{N}$.

In [14], the convergence of both types of trajectories is studied and the results are applied to the case where the maps are iterated function systems (IFSs) generating fractals. Such fractals are then related to limits of non-stationary subdivision procedures generating curves. The following definition is presented in [14] and is used in this paper.

Definition 1.5. Two sequences $\{x_i\}_{i \in \mathbb{N}}$ and $\{y_i\}_{i \in \mathbb{N}}$ in a complete metric space (X, d) are said to be asymptotically equivalent if $d(x_i, y_i) \rightarrow 0$ as $i \rightarrow \infty$. We denote this relation by

$$\{x_i\} \sim \{y_i\}. \quad (1.7)$$

Obviously \sim is an equivalence relation.

With this definition, we can formulate an important property of these trajectories.

Proposition 1.6 (Asymptotic Equivalence of Trajectories). *Let $\{T_i\}_{i \in \mathbb{N}}$ be a sequence of transformations on X where each T_i is a Lipschitz map with Lipschitz constant s_i . If $\lim_{k \rightarrow \infty} \prod_{i=1}^k s_i = 0$ then for any $x, y \in X$,*

$$\begin{aligned} \{\Phi_k(x)\} &\sim \{\Phi_k(y)\}, \\ \{\Psi_k(x)\} &\sim \{\Psi_k(y)\}. \end{aligned} \quad (1.8)$$

To present the results in [14] concerning convergence of trajectories, we need the following definition.

Definition 1.7 (Invariant Domain of $\{T_i\}$). We call $C \subseteq X$ an invariant domain of a sequence of transformations $\{T_i\}_{i \in \mathbb{N}}$ if

$$\forall i \in \mathbb{N}, \forall x \in C : T_i(x) \in C. \quad (1.9)$$

For the convergence of forward trajectories $\{\Phi_k(x)\}$, it is assumed in [14] that the sequence of maps converges to a limit map: $\lim_{i \rightarrow \infty} T_i = T$.

Proposition 1.8 (Convergence of Forward Trajectories). *Let $\{T_i\}_{i \in \mathbb{N}}$ be a sequence of transformations on X with a common compact invariant domain C . Let $T : C \rightarrow C$ be a Lipschitz map with Lipschitz constant $\mu < 1$. If*

$$\lim_{i \rightarrow \infty} \sup_{x \in C} d(T_i(x), T(x)) = 0, \quad (1.10)$$

then for any $x \in C$ the forward trajectory $\{\Phi_k(x)\}_{k \in \mathbb{N}}$ converges to the fixed point p of T , $p = Tp$, namely,

$$\lim_{k \rightarrow \infty} d(\Phi_k(x), p) = 0. \quad (1.11)$$

For the convergence of the backward trajectories $\{\Psi_k(x)\}$ the following result is presented in [14].

Proposition 1.9 (Convergence of Backward Trajectories). *Let $\{T_i\}_{i \in \mathbb{N}}$ be a sequence of transformations on X with a common compact invariant domain C . Assume that each T_i is a Lipschitz map with Lipschitz constant s_i . If $\sum_{k=1}^{\infty} \prod_{i=1}^k s_i < \infty$, then the backward trajectories $\{\Psi_k(x)\}_{k \in \mathbb{N}}$ converge for all points $x \in C$ to a unique point in C .*

Proposition 1.9 is an example of an extension of the Banach fixed point theorem. Namely, for a given infinite iterative process, it states conditions that guarantee the existence of a basin of attraction from which the iterative process converges to a unique attractor. In this paper we present such extensions for several types of infinite iterative processes.

2. Trees of Maps (TMs)

Having in mind non-uniform subdivision schemes and possible applications to the generation of fractals which are non-uniform in space, we introduce the structure of a tree of maps. For expository purposes, we present the idea for the case of binary trees of maps.

Let us denote by $B^{[k]}$ the set of binary codes of length k from the alphabet $\{1, 2\}$,

$$B^{[k]} := \{\eta_k : \eta_k = (i_1 i_2 \dots i_{k-1} i_k), i_j \in \{1, 2\}\} = \{1, 2\}^k. \quad (2.1)$$

We denote the set of infinite binary codes by $B^{[\infty]}$,

$$B^{[\infty]} := \{\eta : \eta = (i_1 i_2 \dots i_k \dots), i_j \in \{1, 2\}\} = \{1, 2\}^{\mathbb{N}}. \quad (2.2)$$

In the following, we define operations on $B^{[k]}$ and on $B^{[\infty]}$.

Definition 2.1 (Prolongation and Truncation Operators).

For a finite code $\eta_k = (i_1 i_2 \dots i_{k-1} i_k) \in B^{[k]}$, we define the prolongation operator $\pi_j : B^{[k]} \rightarrow B^{[k+1]}$ by

$$\pi_j \eta_k := \eta_k j := (i_1 i_2 \dots i_{k-1} i_k j), \quad j = 1, 2.$$

The ℓ -truncation operator $\tau_\ell : B^{[k]} \rightarrow B^{[\ell]}$ for $1 \leq \ell < k$, is defined by


$$\tau_\ell \eta_k := (i_1 i_2 \dots i_\ell) := \eta_\ell.$$

The last definition applies also to $\eta \in B^{[\infty]}$.

Now, consider an infinite collection of continuous functions (maps) from X to itself,

$$\{f_{\eta_k} : \eta_k \in B^{[k]}, k \in \mathbb{Z}_+\}.$$

A function f_{η_k} is also denoted by $f_{i_1 i_2 \dots i_{k-1} i_k}$. We arrange the maps in a tree structure, as depicted below,



where the functions corresponding to the two branches of the same code $f_{\eta_{k1}}$ and $f_{\eta_{k2}}$ are different.

2.1. Attractor of a Tree of Maps

In the following, we define the notion of the attractor of a tree of maps. We present sufficient conditions for the convergence of an infinite process on the tree to a unique attractor $U_{TM} \subset C \subset X$ where C is a common invariant domain of all the maps $\{f_{\eta_k} : \eta_k \in B^{[k]}, k \in \mathbb{N}\}$ in the tree.

Definition 2.2 (Paths in a Tree). Each infinite code $\eta \in B^{[\infty]}$ defines a path in the tree which is an infinite sequence of codes $P_\eta = \{\eta_1, \eta_2, \dots, \eta_k, \dots\}$ such that $\eta_\ell = \tau_\ell \eta$ for $\ell \in \mathbb{N}$.

A path P_η defines a sequence of maps from X to itself, $\{f_{\eta_k}\}_{k \in \mathbb{N}}$. Therefore, the proof of the next proposition is a direct consequence of Proposition 1.9 for the convergence of backward trajectories.

Proposition 2.3 (Convergence along a Path P_η). *Let P_η be a path in a tree T of maps and let $\{f_{\eta_k}\}_{\eta_k \in P_\eta}$ be the sequence of maps along this path with a common compact invariant domain C_{P_η} and with associated Lipschitz constants $\{s_{\eta_k}\}_{\eta_k \in P_\eta}$. If $\sum_{k=1}^{\infty} \prod_{i=1}^k s_{\eta_i} < \infty$ then*

$$\gamma(\eta) := \lim_{k \rightarrow \infty} f_{\tau_1 \eta} \circ \dots \circ f_{\tau_{k-1} \eta} \circ f_{\tau_k \eta}(x) \quad (2.3)$$

exists for all $x \in C_{P_\eta}$, and is the same element in C_{P_η} .

Proposition 2.4 (Convergence to U_{TM}). *If the conditions of Proposition 2.3 are satisfied for all paths in T with the same invariant domain C , then the following set $U_{TM} \subset C$ is uniquely defined for all $x \in C$:*

$$U_{TM} = \bigcup_{\eta \in B^{[\infty]}} \gamma(\eta) = \bigcup_{\eta \in B^{[\infty]}} \lim_{k \rightarrow \infty} f_{\tau_1 \eta} \circ \dots \circ f_{\tau_{k-1} \eta} \circ f_{\tau_k \eta}(x). \quad (2.4)$$

Definition 2.5 (Convergent TMs). Let the TM be such that for every path in it, the limit (2.3) exists and is the same for all $x \in C \subset X$. Then we call the TM *convergent* and term the set $U_{TM} = \bigcup_{\eta \in B^{[\infty]}} \gamma(\eta)$ the *attractor of TM*.

2.2. Changing the Order of \bigcup and \lim in (2.4).

In order to ensure a practical computation of U_{TM} , we should consider changing the order of the operations \bigcup and \lim in (2.4). Changing the order requires a stronger assumption on the convergence along the paths of TM, namely, a uniform convergence, uniform on all paths in the tree. The objects we deal with are nonvoid compact subsets of X . Hence, the convergence we ask for is in $\mathbb{H}(X)$, the collection of all nonvoid compact subsets of X . Note that $\mathbb{H}(X)$ becomes a complete metric space when endowed with the Hausdorff metric h [2].

Proposition 2.6 (Uniform onvergence to U_{TM}). *Let us ssume that all the maps in T share the same common invariant domain C . We further assume that for all paths in T*

$$\prod_{i=1}^k s_{\tau_i \eta} \leq \delta_k, \quad \forall \eta \in B^{[\infty]}, \quad (2.5)$$

where $\sum_{k=1}^{\infty} \delta_k < \infty$. Then,

$$U_{TM} = \lim_{k \rightarrow \infty} \bigcup_{\eta \in B^{[\infty]}} f_{\tau_1 \eta} \circ \cdots \circ f_{\tau_{k-1} \eta} \circ f_{\tau_k \eta}(x), \quad (2.6)$$

where the convergence is with respect to the Hausdorff metric h .

Proof. The assumption (2.5) implies that

$$\sum_{k=m}^{\infty} \prod_{i=1}^k s_{\tau_i \eta} \leq e_m, \quad \forall \eta \in B^{[\infty]}, \quad (2.7)$$

where $\lim_{m \rightarrow \infty} e_m = 0$. For $x \in C$ and $k \in \mathbb{N}$, let

$$\gamma_k(\eta)(x) := f_{\tau_1 \eta} \circ \cdots \circ f_{\tau_{k-1} \eta} \circ f_{\tau_k \eta}(x). \quad (2.8)$$

Following the proof of convergence of backward trajectories in [14], we note that for every $x \in X$ and $\eta \in B^{[\infty]}$, $\{\gamma_k(\eta)(x)\}$ is a Cauchy sequence in (X, d) and

$$d(\gamma_m(\eta)(x), \gamma_{m+\ell}(\eta)(x)) \leq E e_m, \quad \forall \ell \in \mathbb{N}, \quad (2.9)$$

where $E = \text{diam}(C)$.

Since $\lim_{k \rightarrow \infty} \gamma_k(\eta)(x) = \gamma(\eta)$, it follows that

$$d(\gamma_m(\eta)(x), \gamma(\eta)) \leq E e_m. \quad (2.10)$$

Hence,

$$h \left(\bigcup_{\eta \in B^{[\infty]}} \gamma_m(\eta)(x), \bigcup_{\eta \in B^{[\infty]}} \gamma(\eta) \right) \leq E e_m,$$

which implies the convergence of (2.6) with respect to h . \square

2.2.1. Self-Referential Property.

A convergent TM with maps $\{f_{\eta_k} : \eta_k \in B^{[k]}, k \in \mathbb{N}\}$, defines two subtrees, TM1 with the maps $\{f_{1\eta_k} : \eta_k \in B^{[k]}, k \in \mathbb{N}\}$ and TM2 with the maps $\{f_{2\eta_k} : \eta_k \in B^{[k]}, k \in \mathbb{N}\}$. Assume that TM1 has attractor U_{TM1} and TM2 has attractor U_{TM2} . Then these two attractors are related to the attractor U_{TM} of TM by

$$U_{TM} = f_1(U_{TM1}) \cup f_2(U_{TM2}). \quad (2.11)$$

Remark 2.7 (General TMs). The above results are presented for binary TMs. The case of general TMs involves more complicated indexing but can be treated in exactly the same manner. For general TMs, the union operation in the definition of the limit set U_{TM} in (2.4) is over all the paths in the tree.

2.2.2. Code Dependence and Location Dependence.

Within a TM, we have a collection of maps which are code dependent. We explain below that code dependence can imply location dependence. That is, different paths lead to different locations in the attractor U_{TM} .

Let P_η and P_ν be two paths in a convergent TM with $\eta = (i_1 i_2 i_3 \dots)$ and $\nu = (j_1 j_2 j_3 \dots)$. Let $\gamma(\eta)$, respectively, $\gamma(\nu)$ be the two points in X generated by the limits (2.3). Endowed with the Fréchet metric $d_F : B^{[\infty]} \times B^{[\infty]} \rightarrow \mathbb{R}$,

$$d_F(P_\eta, P_\nu) := \sum_{n=1}^{\infty} \frac{|i_n - j_n|}{3^n},$$

$(B^{[\infty]}, d_F)$ becomes a compact metric space. It is known (cf. for instance, [1, 15]) that there exists a continuous surjection γ from code space $B^{[\infty]}$ to the attractor of an IFS. In the TM setting, the mapping $\gamma(\eta)$ is also a continuous map from $B^{[\infty]}$ to the attractor U_{TM} , as shown below.

Proposition 2.8. *Under the conditions of Proposition 2.4, the mapping $\gamma(\eta)$ is a continuous surjection from $B^{[\infty]}$ onto U_{TM} .*

Proof. To prove continuity at P_η , let us assume that $d_F(P_\eta, P_\nu) < \delta$. It follows that $i_m = j_m$ for $1 \leq m \leq \ell(\delta)$ where $\ell(\delta) = \lceil -\log_3(\delta) \rceil$. Using the expression in (2.3), we examine the distance

$$\epsilon_k := d(f_{\eta_1} \circ \dots \circ f_{\eta_{k-1}} \circ f_{\eta_k}(x), f_{\nu_1} \circ \dots \circ f_{\nu_{k-1}} \circ f_{\nu_k}(x)), \quad x \in C, \quad k > \ell(\delta). \quad (2.12)$$

Since $\eta_m = \tau_m$ for $1 \leq m \leq \ell(\delta)$, it follows by recursive application of

$$d(f_{\eta_m}(a), f_{\eta_m}(b)) \leq s_{\eta_m} d(a, b),$$

that

$$\epsilon_k \leq \prod_{m=1}^{\ell(\delta)} s_{\eta_m} \cdot d(f_{\eta_{\ell(\delta)+1}} \circ \dots \circ f_{\eta_k}(x), f_{\nu_{\ell(\delta)+1}} \circ \dots \circ f_{\nu_k}(x)).$$

Hence,

$$\epsilon_k \leq D \prod_{m=1}^{\ell(\delta)} s_{\eta_m}, \quad \forall k > \ell(\delta),$$

where D is the diameter of C . Consequently,

$$d(\gamma(\eta), \gamma(\nu)) \leq D \prod_{m=1}^{\ell(\delta)} s_{\eta_m}.$$

The proof follows by observing that as $\delta \rightarrow 0$, $\ell(\delta) \rightarrow \infty$ and $\prod_{m=1}^{\ell(\delta)} s_{\eta_m} \rightarrow 0$. □

The continuity of γ implies that the code dependence of the functions in the tree means location dependence. For more details about the relation between codes and points in the attractor of an iterated function system, see e.g. [1, 15].

In the next subsection we apply the above results to trees of function systems consisting of pairs of maps each.

2.3. Trees of Function Systems (TFSs)

Recall that an iterated function system (IFS) is a pair consisting of a complete metric space (X, d) and a finite family of continuous maps $f_i : X \rightarrow X$, $i \in \{1, 2, \dots, n\}$. We denote such an IFS by $\mathcal{F} = \{X; f_i : i = 1, 2, \dots, n\}$. If the f_i are contraction maps, the IFS is called contractive. The contraction constant of \mathcal{F} is $L_{\mathcal{F}} = \max_{i=1,2,\dots,n} \text{Lip}(f_i)$.

Consider the set-valued mapping $\mathcal{F} : \mathbb{H}(X) \rightarrow \mathbb{H}(X)$,

$$\mathcal{F}(B) := \bigcup_{f \in \mathcal{F}} f(B), \quad B \in \mathbb{H}(X), \quad (2.13)$$

where $f(B) := \{f(b) : b \in B\}$. It is well known that for a contractive IFS, \mathcal{F} is a contraction in $(\mathbb{H}(X), h)$ with contraction constant $L_{\mathcal{F}} = \max_{i=1,2,\dots,n} \text{Lip}(f_i)$ [2]. Therefore, by the Banach contraction principle \mathcal{F} has a unique fixed point, denoted here by U_{IFS} , called the attractor of the IFS.

Here, we consider a binary IFS $\mathcal{F} = \{f_1, f_2\}$. If both functions are contractions on X , the attractor of the IFS is given by

$$U_{IFS} = \lim_{k \rightarrow \infty} \bigcup_{(i_1 i_2 \dots i_k) \in B^{[k]}} f_{i_1} \circ f_{i_2} \circ \dots \circ f_{i_k}(A), \quad (2.14)$$

where A is any non-void compact subset of X . By Proposition 2.6, the convergence in (2.14) is ensured.

A binary tree of maps may be considered as a tree of function systems. Each pair of functions $(f_{\eta_{k1}}, f_{\eta_{k2}})$ defines a function system, which we denote by \mathcal{F}_{η_k} . Consequently, the above tree of maps induces an infinite binary tree of function systems (TFS) $\{\mathcal{F}_{\eta_k}\}_{\eta_k \in B^{[k]}, k \in \mathbb{N}}$ as follows. For $k = 0$, set

$$\mathcal{F}_0 = \{f_1, f_2\},$$

and for $\eta_k \in B^{[k]}$, $k \in \mathbb{N}$,

$$\mathcal{F}_{\eta_k} = \{f_{\eta_{k1}}, f_{\eta_{k2}}\}.$$

Following backward paths on the tree of function systems, the natural generalization of the fractal U_{IFS} in (2.14) is

$$U_{TFS} = \lim_{k \rightarrow \infty} \bigcup_{\eta_k \in B^{[k]}} f_{\tau_1 \eta_k} \circ f_{\tau_2 \eta_k} \cdots \circ f_{\eta_k}(A). \quad (2.15)$$

To ensure convergence in (2.15), with respect to h , we assume here that the condition (2.7) in Proposition 2.6 is satisfied, with $s_{\eta_k} = \text{Lip}(\mathcal{F}_{\eta_k})$.

If all the function systems are the same IFS, we have $U_{TFS} = U_{IFS}$. If the function systems $\{\mathcal{F}_{\eta_k}\}$ are all the same for a fixed k , we retrieve the case of sequences of function systems discussed in [14]. In the following example we present a tree of function systems and its attractor.

Example. To each code $\eta_k = (i_1 i_2 \dots i_k)$ we attach a number

$$t(\eta_k) := \sum_{j=1}^k (i_j - 1)2^{-j} \in [0, 1].$$

We define the function systems in the TFS by the following maps on $X = \mathbb{R}^2$:

$$f_{\eta_{k1}} \left(\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \right) = \begin{pmatrix} 0.4 & 0.4 \\ -0.5 & 0.3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} 4 \\ 4 \end{pmatrix},$$

and

$$f_{\eta_{k2}} \left(\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \right) = 0.8 \begin{pmatrix} \cos \theta_k & \sin \theta_k \\ -\sin \theta_k & \cos \theta_k \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} -2 \\ 0 \end{pmatrix},$$

where $\theta_k := 3t(\eta_k) + 1$, $k \in \mathbb{N}$.

All of the above maps are contractive and thus the tree of maps is convergent by Proposition 2.4. In Figure 1 below, we depict its attractor. The noticeable property of this attractor is that it has different structures at different locations. The challenge is to investigate how to design desirable structures at specific locations.

3. Staircase Maps on a Sequence of Metric Spaces and their Trajectories

Let (X, d_X) and (Y, d_Y) be two complete metric spaces and let $T : X \rightarrow Y$ be a continuous map. Consider a non-void compact set $C \subset X$.

Definition 3.1. The Lipschitz constant associated with f on C is defined by

$$\text{Lip}_C(T) := \sup_{a, b \in C, a \neq b} \frac{d_Y(T(a), T(b))}{d_X(a, b)}.$$

A map T is said to be Lipschitz map with respect to C if $\text{Lip}_C(T) < +\infty$ and a contraction with respect to C if $\text{Lip}_C(T) < 1$.




FIGURE 1. An attractor U_{TFS} showing different structures at different locations.

Let us consider an infinite sequence of complete metric spaces $\{(X_i, d_i)\}_{i \in \mathbb{N}_0}$ and an associated sequence of continuous maps $\{T_i\}_{i \in \mathbb{N}}$,

$$T_i : X_i \rightarrow X_{i-1}, \quad i \in \mathbb{N}. \quad (3.1)$$

Correspondingly, we assume the existence of non-void compact subsets $\{C_i\}_{i \in \mathbb{N}_0}$, $C_i \subset X_i$, of bounded diameters, $\text{diam}(C_i) \leq D$, such that

$$T_i : C_i \rightarrow C_{i-1}, \quad i \in \mathbb{N}. \quad (3.2)$$

We further assume that each T_i is Lipschitz with respect to C_i , $i \in \mathbb{N}$. We denote its Lipschitz constant by s_i :

$$s_i := \text{Lip}_{C_i}(T_i), \quad i \in \mathbb{N}. \quad (3.3)$$

In order to consider backward trajectories in the present general setting of maps between different spaces, we need to modify their definition. We relate a trajectory of $\{T_i\}_{i \in \mathbb{N}}$ to a base sequence of points

$$\bar{x} = \{x_i\}_{i \in \mathbb{N}}, \quad x_i \in C_i. \quad (3.4)$$

We refer to such a sequence as a *base sequence for a backward staircase trajectory*.

Definition 3.2 (Backward Staircase Trajectories of $\{T_i\}$ with respect to \bar{x}). Under (3.1) and (3.2), a trajectory of the sequence of maps $\{T_i\}_{i \in \mathbb{N}}$ with

respect to a sequence of base points \bar{x} as in (3.4) is the sequence of points $\{p_k(\bar{x})\}_{k \in \mathbb{N}} \subset X_0$ defined by

$$p_k(\bar{x}) = T_1 \circ T_2 \circ \cdots \circ T_k(x_k), \quad k \in \mathbb{N}. \quad (3.5)$$

Proposition 3.3 (Asymptotic Equivalence of Backward Staircase Trajectories). *Let $\{(X_i, d_i)\}_{i \in \mathbb{N}_0}$, $\{T_i\}_{i \in \mathbb{N}}$, $\{C_i\}_{i \in \mathbb{N}_0}$, and $\{s_i\}_{i \in \mathbb{N}}$ be as above. If $\lim_{k \rightarrow \infty} \prod_{i=1}^k s_i = 0$, then for any two base sequences \bar{x} and \bar{y} ,*

$$\lim_{k \rightarrow \infty} d_0(p_k(\bar{x}), p_k(\bar{y})) = 0. \quad (3.6)$$

The proof of this result is similar to the proof in [14] for backward trajectories. We present it here for the sake of completeness.

Proof. Let \bar{x}, \bar{y} be two base sequences and consider the backward staircase trajectories defined in (3.5). Using the fact that T_i is a Lipschitz map with Lipschitz constant s_i and that $\text{diam}_i(C_i) \leq D$ for $i \in \mathbb{N}$, we have that

$$\begin{aligned} d_0(p_k(\bar{x}), p_k(\bar{y})) &\leq s_1 d_1(T_2 \circ T_3 \circ \cdots \circ T_k(x_k), T_2 \circ T_3 \circ \cdots \circ T_k(y_k)) \\ &\leq s_1 s_2 d_2((T_3 \circ T_4 \circ \cdots \circ T_k(x_k), T_3 \circ T_4 \circ \cdots \circ T_k(y_k)) \leq \dots \\ &\leq \left(\prod_{i=1}^k s_i \right) d_k(x_k, y_k) \leq D \left(\prod_{i=1}^k s_i \right), \end{aligned} \quad (3.7)$$

from which the result follows. \square

Note that the condition $\lim_{k \rightarrow \infty} \prod_{i=0}^k s_i = 0$ certainly holds if $s_i < s < 1$, for all $i \in \mathbb{N}$.

Proposition 3.4 (Convergence of Backward Staircase Trajectories). *Let*

$\{(X_i, d_i)\}_{i \in \mathbb{N}_0}$, $\{T_i\}_{i \in \mathbb{N}}$, $\{C_i\}_{i \in \mathbb{N}_0}$, and $\{s_i\}_{i \in \mathbb{N}}$ be as above. If $\sum_{k=1}^{\infty} \prod_{i=1}^k s_i < \infty$, then for all base sequences \bar{x} the staircase trajectories $\{p_k(\bar{x})\}$ defined by (3.5) converge to a unique limit in C_0 .

Proof. By (3.5), the relations in (3.7), and the assumption $\text{diam}(C_i) \leq D$ for $i \in \mathbb{N}$, we have that

$$\begin{aligned} d_0(p_{k+1}(\bar{x}), p_k(\bar{x})) &= d_0(T_1 \circ T_2 \circ \cdots \circ T_{k+1}(x_{k+1}), T_1 \circ T_2 \circ \cdots \circ T_k(x_k)) \\ &\leq \left(\prod_{i=1}^k s_i \right) d_k(T_{k+1}(x_{k+1}), x_k) \leq D \left(\prod_{i=1}^k s_i \right). \end{aligned} \quad (3.8)$$

For $m, k \in \mathbb{N}$ with $m > k$, we obtain in view of (3.8),

$$\begin{aligned} d_0(p_m(\bar{x}), p_k(\bar{x})) &\leq d_0(p_m(\bar{x}), p_{m-1}(\bar{x})) + \dots \\ &\quad + d_0(p_{k+2}(\bar{x}), p_{k+1}(\bar{x})) + d_0(p_{k+1}(\bar{x}), p_k(\bar{x})) \\ &\leq D \prod_{i=1}^{m-1} s_i + \dots + D \prod_{i=1}^{k+1} s_i + D \prod_{i=1}^k s_i = D \left(\sum_{j=k}^{m-1} \prod_{i=1}^j s_i \right). \end{aligned} \quad (3.9)$$

$$(3.10)$$

Since $\sum_{k=1}^{\infty} \prod_{i=1}^k s_i < \infty$, Eq. (3.9) asserts that $d_0(p_m(\bar{x}), p_k(\bar{x})) \rightarrow 0$ as $k \rightarrow \infty$. That is, $\{p_k(\bar{x})\}_{k \in \mathbb{N}}$ is a Cauchy sequence in C_0 and due to the completeness of (X_0, d_0) and the compactness of C_0 , it is convergent to a limit in C_0 . The uniqueness of the limit is derived by the equivalence of all trajectories as proved in Proposition 3.3. \square

Remark 3.5 (Invariance to Scaling of the Metrics $\{d_i\}$). In view of Definition 3.1, the Lipschitz constant of a mapping from one metric space to another, it is clear that by a proper scaling of the metrics $\{d_i\}$ we can make all the maps $\{T_i\}$ contractive. By scaling the metrics we mean replacing $\{d_i\}$ with $\{\tilde{d}_i\}$, where

$$\tilde{d}_i(x, y) = \alpha_i d_i(x, y) \quad \forall x, y \in X_i, \quad \alpha_i > 0. \quad (3.11)$$

Such a scaling implies new Lipschitz constants

$$\tilde{s}_i = \frac{\alpha_{i-1}}{\alpha_i} s_i. \quad (3.12)$$

Choosing $\alpha_0 = 1$ we obtain

$$\prod_{i=1}^k \tilde{s}_i = \frac{1}{\alpha_k} \prod_{i=1}^k s_i. \quad (3.13)$$

Let us discuss the implications of the scaling of the metrics. To make the discussion more concrete, consider the case $X_i = \mathbb{R}^i$. A natural possibility is to use the same metric for all these spaces. For example, the metric induced by the maximum norm. If the conditions for asymptotic equivalence or for convergence are not satisfied for these metrics, we can scale the metrics up ($\alpha_i > 1$) in the case of backward trajectories. As we explain below, convergence can be obtained by scaling the metrics.

In view of the proof of Proposition 3.4, the critical quantity for convergence is the expression

$$\left(\prod_{i=1}^k s_i \right) d_k(T_{k+1}(x_{k+1}), x_k)$$

in equation (3.8). Because of (3.11) and (3.7), we find that this term is invariant under the scaling of metrics:

$$\left(\prod_{i=1}^k \tilde{s}_i \right) \tilde{d}_k(T_{k+1}(x_{k+1}), x_k) = \left(\prod_{i=1}^k s_i \right) d_k(T_{k+1}(x_{k+1}), x_k). \quad (3.14)$$

We see that by our method of proof nothing can be gained by scaling the metrics.

Remark 3.6 (Grouping). Sometimes the conditions for the convergence of backward trajectories in Proposition 3.4 are not satisfied but they still may converge. One way of relaxing the conditions is to form groups of maps. Consider the following grouping of ℓ maps:

$$\begin{aligned} G_j^{[\ell]} &: X_{j\ell} \rightarrow X_{(j-1)\ell}, \quad j \in \mathbb{N}, \\ G_j^{[\ell]} &:= T_{(j-1)\ell+1} \circ T_{(j-1)\ell+2} \circ \dots \circ T_{j\ell-1} \circ T_{j\ell}. \end{aligned} \quad (3.15)$$

If for some ℓ the conditions for convergence of backward trajectories of $\{G_j^{[\ell]}\}$ are fulfilled, then the backward trajectories of $\{T_i\}$ will converge for some base sequences.

Corollary 3.7 (Partial Backward Staircase Trajectories). *Let us assume that the backward staircase trajectory of $\Sigma_1 := \{T_i\}_{i \in \mathbb{N}}$ converge to a unique limit for all possible base sequences. Let us denote this limit by τ_0 . Consider the partial sequences $\Sigma_m = \{T_i\}_{i=m}^\infty$, $m > 1$ and assume that the staircase trajectories of each Σ_m converge to a unique limit τ_{m-1} , for all possible base sequences. Then,*

$$\tau_{m-1} = T_m(\tau_m), \quad m \in \mathbb{N}. \quad (3.16)$$

Remark 3.8. As we extended the notion of backward trajectories from sequences of maps in a metric space to staircase trajectories, it is also possible to extend the notion of trees of maps on X to a tree of staircase maps.

4. Sequences of Function Systems on a Sequence of Metric Spaces

Consider an infinite sequence of complete metric spaces, $\{(X_i, d_i)\}_{i \in \mathbb{N}_0}$ and a related sequence of function systems $\{\mathcal{F}_i\}_{i \in \mathbb{N}}$ defined by

$$\mathcal{F}_i = \{f_{1,i}, f_{2,i}, \dots, f_{n_i,i}\},$$

where $f_{r,i} : X_i \rightarrow X_{i-1}$, $r = 1, 2, \dots, n_i$, are continuous maps. The associated set-valued maps are given by

$$\mathcal{F}_i : \mathbb{H}(X_i) \rightarrow \mathbb{H}(X_{i-1}); \quad \mathcal{F}_i(A) = \bigcup_{r=1}^{n_i} f_{r,i}(A).$$

We now assume the existence of non-void compact subsets $\{C_i\}_{i \in \mathbb{N}_0}$, $C_i \subset X_i$ of bounded diameters, $\text{diam}(C_i) \leq D$, such that

$$f_{r,i} : C_i \rightarrow C_{i-1}, \quad r = 1, \dots, n_i, \quad i \in \mathbb{N}. \quad (4.1)$$

We further assume that each $f_{r,i}$ is Lipschitz with respect to C_i , $i \in \mathbb{N}$, and we denote its Lipschitz constant by

$$s_{r,i} = \text{Lip}_{C_i}(f_{r,i}). \quad (4.2)$$

The contraction factor of \mathcal{F}_i in $(\mathbb{H}(X_i), h)$ is $L_{\mathcal{F}_i} = \max_{r=1,2,\dots,n_i} s_{r,i} =: s_i$ [14].

Let

$$\bar{A} = \{A_i\}_{i \in \mathbb{N}}, \quad A_i \subset C_i \quad (4.3)$$

be a base sequence.

The following results follow directly from Proposition 3.4 and Corollary 3.7.

Corollary 4.1 (Convergence of Staircase SFS). *Consider the above staircase structure subject to the above assumptions. Then, if*

$$\sum_{k=1}^{\infty} \prod_{i=1}^k s_i < \infty,$$

the staircase trajectory of $\Sigma_1 = \{\mathcal{F}_i\}_{i \in \mathbb{N}}$,

$$p_k(\bar{A}) = \mathcal{F}_1 \circ \mathcal{F}_2 \circ \dots \circ \mathcal{F}_k(A_k), \quad k \in \mathbb{N}, \quad (4.4)$$

converges for any base sequence \bar{A} to a unique set (attractor) $P_0 \subseteq C_0$.

Corollary 4.2 (Staircase Self-referential Property). *Under the conditions of Corollary 4.1, the staircase trajectories of $\Sigma_m = \{\mathcal{F}_i\}_{i=m}^{\infty}$, namely, the sequence of sets*

$$p_k(\bar{A}) = \mathcal{F}_m \circ \mathcal{F}_{m+1} \circ \dots \circ \mathcal{F}_k(A_k), \quad k \geq m, \quad (4.5)$$

converge for any base sequence to a unique set (attractor) $P_{m-1} \subseteq C_{m-1}$. Furthermore,

$$P_{m-1} = \mathcal{F}_m(P_m) = \bigcup_{r=1}^{n_m} f_{r,m}(P_m), \quad m \in \mathbb{N}. \quad (4.6)$$

Remark 4.3. Equation (4.6) extends the self-referential property of fractals generated by IFSs to the setting of SFSs [2].

5. From Subdivision to Staircase SFS

In this section, we first review the relation between non-stationary subdivision with a mask of fixed size and backward trajectories of a SFS as presented in [14]. Then we suggest a framework for dealing with non-stationary subdivision with masks of increasing size.

Starting with a set of control points $p^0 = \{p_j^0 \in \mathbb{R}^m, j \in \mathbb{Z}\}$ at level 0, a non-stationary binary subdivision processes can be written in matrix form as

$$\bar{p}^{k+1} = S^{[k]} \bar{p}^k, \quad k = 1, 2, \dots, \quad (5.1)$$

where \bar{p}^k is the infinite matrix whose rows are the points at level k of the subdivision, and each $S^{[k]} := S_{a^{[k]}}$ is a ‘‘two-slanted’’ infinite matrix. Namely,

$(S_{a^{[k]}})_{i,j} = a_{i-2j}^{[k]}$, where $a^{[k]}$ is a finite sequence of reals termed the mask of the subdivision process. The refinement rule (5.1) can be written as $p_i^{k+1} = \sum_j a_{i-2j}^{[k]} p_j^k$.

As demonstrated in [8], non-stationary subdivision processes can generate interesting limits which cannot be generated by stationary schemes ($S^{[k]} = S$), e.g., exponential splines. Interpolatory non-stationary subdivision schemes can generate new types of compactly supported orthogonal wavelets as shown in [6].

5.1. Non-Stationary Subdivision with Masks of Fixed Size

It is assumed in [14] that the supports of the masks $a^{[k]}$, $|\sigma(a^{[k]})|$, are of the same size which is smaller than the number of initial control points. For a given finite set of control points, $p^0 = \{p_j^0\}_{j=1}^n$, [14] defines for each k the two square $n \times n$ sub-matrices of each $S^{[k]}$, $S_1^{[k]}$ and $S_2^{[k]}$, in the same way as suggested for a stationary scheme in [18]. Out of the points generated at the first level of the subdivision, the first n points are the rows $S_1^{[1]} \bar{p}^0$ and the last n points are the rows of $S_2^{[1]} \bar{p}^0$. At the second level we apply $S_1^{[2]}$ and $S_2^{[2]}$ to the two resulting vectors of points and so on. The set of all the points generated at level k of the subdivision process is given by

$$p^k = \bigcup_{i_1, i_2, \dots, i_k \in \{1, 2\}} \text{rows} (S_{i_k}^{[k]}, \dots, S_{i_2}^{[2]} S_{i_1}^{[1]} \bar{p}^0) . \quad (5.2)$$

Here, $\text{rows}(\bar{p})$ denotes the set of points comprised of the rows of \bar{p} . If the subdivision is convergent [14],

$$p^k \rightarrow p^\infty \quad \text{as } k \rightarrow \infty, \quad (5.3)$$

where p^∞ is the set of points defined by the non-stationary subdivision process starting with p^0 .

As shown in [14], the same limit is obtained by backward trajectories of a related SFS $\{\mathcal{F}_k\}$, where $\mathcal{F}_k = \{X; f_{1,k}, f_{2,k}\}$ with level dependent maps

$$f_{r,k}(A) = AP^{-1} S_r^{[k]} P, \quad r = 1, 2. \quad (5.4)$$

Here P is the $n \times n$ matrix defined as in the stationary case:

1. The first m columns of P constitutes the vector \bar{p}^0 of given n control points p^0 which are points in \mathbb{R}^m .
2. The last column is a column of 1's.
3. The rest of the columns are defined so that P is non-singular. We assume here that the control points p^0 do not all lie on an $m - 1$ hyper plane so that the first m columns of P will be linearly independent and that the column of 1's is independent of the first m columns.

5.1.1. A Simpler SFS Construction for Non-Stationary Subdivision. Before dealing with masks of increasing size, we present here a simpler SFS replacing the above SFS that was presented in [14]. This simplification uses the theory

in Section 4 which was not available in [14]. The functions in this new SFS operate on row vectors in \mathbb{R}^n by right matrix multiplication.

We use the same sequence of $n \times n$ matrices $\{S_r^{[k]}\}$ defined above and for a row vector $A \in \mathbb{R}^n$ we define:

$$\begin{aligned} f_{r,1} &: \mathbb{R}^n \rightarrow \mathbb{R}^m, \\ f_{r,1}(A) &= AS_r^{[1]}\bar{p}^0, \quad r = 1, 2, \end{aligned} \quad (5.5)$$

and for $k > 1$

$$\begin{aligned} f_{r,k} &: \mathbb{R}^n \rightarrow \mathbb{R}^n, \\ f_{r,k}(A) &= AS_r^{[k]}, \quad r = 1, 2. \end{aligned} \quad (5.6)$$

The rows of $\bar{p}^0 \in \mathbb{R}^{n \times m}$ are the n initial control points in \mathbb{R}^m for the subdivision process.

Let us follow a backward trajectory of the SFS $\Sigma := \{\mathcal{F}_k\}$, $\mathcal{F}_k = \{f_{1,k}, f_{2,k}\}$, starting from $A \in \mathbb{R}^n$:

$$\mathcal{F}_k(A) = f_{1,k}(A) \cup f_{2,k}(A) = AS_1^{[k]} \cup AS_2^{[k]}, \quad k > 1,$$

and for $j, k > 1$

$$\mathcal{F}_j(\mathcal{F}_k(A)) = f_{1,j}(AS_1^{[k]} \cup AS_2^{[k]}) \cup f_{2,j}(AS_1^{[k]} \cup AS_2^{[k]}).$$

Noting that

$$f_{r,j}(AS_i^{[k]}) = AS_i^{[k]}S_r^{[j]}, \quad (5.7)$$

it follows that the set generated at the k th step of a backward trajectory of Σ is

$$\mathcal{F}_1 \circ \mathcal{F}_2 \circ \dots \circ \mathcal{F}_{k-1} \circ \mathcal{F}_k(A) = \bigcup_{i_1, i_2, \dots, i_k \in \{1, 2\}} AS_{i_k}^{[k]} \dots S_{i_2}^{[2]} S_{i_1}^{[1]} \bar{p}^0. \quad (5.8)$$

For a C^0 -stationary subdivision scheme, it is shown in [18] that the maps defined by (5.4) are contractive on Q^{n-1} , the $(n-1)$ -dimensional hyperplane (flat) of vectors of the form $(x_1, \dots, x_{n-1}, 1)$. The new maps (5.5) and (5.6) do not have an evident contraction property and yet a fixed-point theorem holds for the backward trajectories of Σ . Instead of the flat Q^{n-1} we consider another flat in \mathbb{R}^n , namely,

$$K^{n-1} = \left\{ (x_1, x_2, \dots, x_n) : \sum_{i=1}^n x_i = 1 \right\}. \quad (5.9)$$

Notice that if the non-stationary scheme satisfies the constant reproduction property at every subdivision level, then all the maps in the SFS map K^{n-1} into itself. That is,

$$\forall A \in K^{n-1}, \quad AS_r^{[k]} \in K^{n-1}, \quad r = 1, 2, \quad k \in \mathbb{N}. \quad (5.10)$$

Definition 5.1 (The Set K_C^{n-1}).

$$K_C^{n-1} := \left\{ (x_1, x_2, \dots, x_n) : \sum_{i=1}^n x_i = 1, \quad |x_i| \leq C \right\}.$$

The following theorem establishes the convergence in the Hausdorff distance h of the backward trajectories of Σ to the limit curve of the non-stationary subdivision.

Theorem 5.2. *Let $\{S_{a^{[k]}}\}$ be a non-stationary C^0 -convergent subdivision scheme and let $\Sigma = \{\mathcal{F}_k\}_{k=1}^\infty$ be the SFS defined in (5.5) and (5.6). Then the backward trajectories of Σ starting with $\mathbf{A} \subset K_C^{n-1}$ converge to a unique attractor which constitutes the limit curve (in \mathbb{R}^m) of the non-stationary scheme.*

Proof. Since $\{S_{a^{[k]}}\}$ converges, it immediately follows from (5.8), in view of (5.2), that the backward trajectory of Σ initialized with any A converges. We would like to show that all the backward trajectories of Σ initialized with an arbitrary set of points $\mathbf{A} \subset K_C^{n-1}$ converge to the same limit. Since the subdivision scheme converges to a continuous function, it follows that an infinite sequence $\eta = \{i_k\}_{k \in \mathbb{N}}$, $i_k \in \{1, 2\}$, defines a vector \bar{q}_η of n identical points in \mathbb{R}^m ,

$$\bar{q}_\eta = \lim_{k \rightarrow \infty} S_{i_k}^{[k]}, \dots, S_{i_2}^{[2]} S_{i_1}^{[1]} \bar{p}^0 = \begin{pmatrix} q_\eta \\ \cdot \\ \cdot \\ \cdot \\ q_\eta \end{pmatrix}, \quad q_\eta = (q_{\eta,1}, \dots, q_{\eta,m}), \quad (5.11)$$

attached to a parametric value $x_\eta = \sum_{k=1}^\infty (i_k - 1)2^{-k}$. (See Remark 6.2 in [14].)

Starting the backward trajectory with any point $A \in \mathbf{A}$, and following the same sequence η , it follows from (5.8) that the limit is $A\bar{q}$. Recalling (5.9), it now follows that $A\bar{q}_\eta = q_\eta, \forall A \in K^{n-1}$.

Hence,

$$\lim_{k \rightarrow \infty} \mathcal{F}_1 \circ \mathcal{F}_2 \circ \dots \circ \mathcal{F}_{k-1} \circ \mathcal{F}_k(A) = \bigcup_{\eta} A \lim_{k \rightarrow \infty} S_{i_k}^{[k]} \dots S_{i_2}^{[2]} S_{i_1}^{[1]} \bar{p}^0 = \bigcup_{\eta} q_\eta, \quad (5.12)$$

which is the set of all the limit points generated by the subdivision process starting with initial data p^0 . For a bounded set \mathbf{A} this implies that

$$\lim_{k \rightarrow \infty} \mathcal{F}_1 \circ \mathcal{F}_2 \circ \dots \circ \mathcal{F}_{k-1} \circ \mathcal{F}_k(\mathbf{A}) = \bigcup_{A \in \mathbf{A}} \bigcup_{\eta} A \lim_{k \rightarrow \infty} S_{i_k}^{[k]} \dots S_{i_2}^{[2]} S_{i_1}^{[1]} \bar{p}^0 = \bigcup_{\eta} q_\eta, \quad (5.13)$$

□

Above, we assumed that the number of control points n is larger than the size of the subdivision masks. Therefore, when dealing with non-stationary subdivision with increasing mask size, the definition of the related SFS should be revised.

5.2. Non-Stationary Subdivision with Masks of Increasing Size

Consider a binary non-stationary subdivision scheme with increasing mask size. Such schemes are suggested in the subdivision literature for generating highly smooth limit functions of small support [9]. Let us denote the support size of the k -th level subdivision mask by σ_k . For example, the up-function

which is a C^∞ -function of compact support is generated by a non-stationary subdivision with $\sigma_k = k+1$. We hereby suggest a “staircase” SFS $\{\mathcal{F}_k\}$ where $\mathcal{F}_k = \{X_k; f_{1,k}, f_{2,k}\}$ with level dependent backward maps defined on row vectors $A \in \mathbb{R}^{n_k}$ at level k ,

$$f_{r,1} : \mathbb{R}^{n_0} \rightarrow \mathbb{R}^m, \quad f_{r,1}(A) = AS_r^{[1]}\bar{p}^0, \quad (5.14)$$

and for $k > 1$

$$f_{r,k} : \mathbb{R}^{n_k} \rightarrow \mathbb{R}^{n_{k-1}}, \quad f_{r,k}(A) = AS_r^{[k]}, \quad r = 1, 2. \quad (5.15)$$

The vector $p^0 \in \mathbb{R}^{n_0 \times m}$ represents the set of n_0 initial control points in \mathbb{R}^m for the subdivision process.

The matrices $S_r^{[k]}$, $r = 1, 2$, $k \in \mathbb{N}$, are $n_k \times n_{k-1}$ matrices representing the subdivision rules at level k . From a vector of n_{k-1} values, the subdivision at level k generates new $m_k > n_{k-1}$ values at level $k+1$. We set $n_k = \lceil \frac{m_k}{2} \rceil$. $S_1^{[k]}$ generates the first n_k values and $S_2^{[k]}$ generates the last n_k values. We observe that by (5.15) for $k > 1$

$$\mathcal{F}_k : \mathbb{H}(\mathbb{R}^{n_k}) \rightarrow \mathbb{H}(\mathbb{R}^{n_{k-1}}). \quad (5.16)$$

Remark 5.3 (Increase of Mask Size). The growth rate of the support sizes $\{\sigma_k\}$ should be limited so that the subdivision process starting with the initial control points p^0 defines a non-void curve in \mathbb{R}^m .

The above construction leads to a sequence of backward maps $\{\mathcal{F}_k\}$ defined on a sequence of metric spaces $\{\mathbb{H}(\mathbb{R}^{n_k})\}$ and to the question of convergence of the related backward staircase trajectories. Let us develop the expression of the backward staircase trajectory.

For $A \in \mathbb{R}^{n_k}$, let

$$\mathcal{F}_k(A) = f_{1,k}(A) \cup f_{2,k}(A) = AS_1^{[k]} \cup AS_2^{[k]}.$$

Let $\bar{A} = \{A_i\}$, $A_i \in K^{n_i-1}$, $i \in \mathbb{N}$, be any base sequence for a backward staircase trajectory of $\Sigma = \{\mathcal{F}_k\}$. It follows that

$$p_k(\bar{A}) = \mathcal{F}_1 \circ \mathcal{F}_2 \circ \dots \circ \mathcal{F}_{k-1} \circ \mathcal{F}_k(A_k) = \bigcup_{i_1, i_2, \dots, i_k \in \{1, 2\}} A_k S_{i_k}^{[k]} \dots S_{i_2}^{[2]} S_{i_1}^{[1]} \bar{p}^0. \quad (5.17)$$

Hence, if the non-stationary subdivision converges, then many staircase trajectory converges. For example, the backward trajectory with the base sequence elements

$$A_i = (1, 0, 0, \dots, 0), \quad i \in \mathbb{N}. \quad (5.18)$$

Using Corollary 4.1 in order to show convergence of all trajectories to the same limit, one should check the related Lipschitz constants $\{s_i\}$. For the test case of a staircase subdivision generating the up-function, the conditions on $\{s_i\}$ stated in Corollary 4.1 are not satisfied. It can be shown that, for ℓ large enough, a map grouping of order ℓ gives contractive maps and hence the backward trajectories converge. We present below another approach for showing convergence of all staircase backward trajectories to the same limit.

This approach extends the idea developed in Theorem 5.2 and it can be used for all schemes whose mask sizes $\{\sigma_k\}$ exhibit polynomial growth.

Theorem 5.4. *Let $\{S_{a^{[k]}}\}$ be a non-stationary C^1 -convergent subdivision scheme and let $\Sigma = \{\mathcal{F}_k\}_{k=1}^\infty$ be the SFS defined in (5.14) and (5.15). Then the backward trajectories of Σ starting with any base sequence $\bar{A} = \{A_k\}$ with $A_k \subset K_C^{n_k-1}$, for some constant C , converge to a unique attractor which constitutes the limit curve (in \mathbb{R}^m) of the non-stationary scheme.*

Proof. The idea of the proof is similar to that of the proof of Theorem 5.2. However, we do not have a relation like (5.11) since the dimension of $S_{i_k}^{[k]} \dots S_{i_2}^{[2]} S_{i_1}^{[1]} \bar{p}^0$ increases with k . Let us assume that $\{\sigma_k\}$ is growing algebraically with k . Since the subdivision scheme converges to a continuous function, it follows that an infinite sequence $\eta = \{i_k\}_{k \in \mathbb{N}}$, $i_k \in \{1, 2\}$, defines a limit point $q_\eta \in \mathbb{R}^m$ attached to a parametric value $x_\eta = \sum_{k=1}^\infty (i_k - 1)2^{-k}$. Let us denote by $\bar{q}_\eta^{n_k}$ the vector of n_k identical points in \mathbb{R}^m ,

$$\bar{q}_\eta^{n_k} = \begin{pmatrix} q_\eta \\ \cdot \\ \cdot \\ \cdot \\ q_\eta \end{pmatrix}, \quad q_\eta = (q_{\eta,1}, \dots, q_{\eta,m}). \quad (5.19)$$

For k large enough and with $\eta_k = \{i_j\}_{j=1}^k$, $i_j \in \{1, 2\}$, the vector of n_k points in \mathbb{R}^m

$$\bar{p}_{\eta_k} = S_{i_k}^{[k]} \dots S_{i_2}^{[2]} S_{i_1}^{[1]} \bar{p}^0,$$

is close to the smooth limit curve generated by the subdivision scheme. To estimate how close we use here some observations from the theory of subdivision [9].

For a C^1 -convergent scheme it can be shown that

$$\|q_{\eta_k} - q_\eta\| = \mathcal{O}(2^{-k}),$$

$$\|(\bar{p}_{\eta_k})_{i_\cdot} - q_{\eta_k}\| = n_k \mathcal{O}(2^{-k}), \quad i = 1, \dots, n_k.$$

Hence, we obtain

$$\bar{p}_{\eta_k} = \bar{q}_\eta^{n_k} + n_k \mathcal{O}(2^{-k}), \quad \text{as } k \rightarrow \infty. \quad (5.20)$$

Since $A_k \in K_C^{n_k-1}$, it follows that $A_k \bar{q}_\eta^{n_k} = q_\eta$ and an additional n_k factor enters when we apply A_k to the remainder term in (5.20):

$$A_k \bar{p}_{\eta_k} = q_\eta + (n_k)^2 \mathcal{O}(2^{-k}), \quad \text{as } k \rightarrow \infty. \quad (5.21)$$

Now, $\{\sigma_k\}$ is growing algebraically with k and $n_k = n_0 + \sigma_k - \sigma_0$ and thus for any base sequence $\bar{A} = \{A_k\}$ with $A_k \subset K_C^{n_k-1}$ we get

$$A_k S_{i_k}^{[k]} \dots S_{i_2}^{[2]} S_{i_1}^{[1]} \bar{p}^0 \rightarrow q_\eta, \quad \text{as } k \rightarrow \infty. \quad (5.22)$$

The proof now follows by using (5.17). \square

6. Non-Uniform Subdivision

Another class of subdivision processes includes non-uniform schemes in which the subdivision rules depend on the location. An example of such schemes is dealt with in [10]. The approach used in [14] cannot be used for non-uniform schemes not even those with a fixed mask size. The introduction of the notion of staircase trajectories opens up new possibilities. Another option is to use the structure of trees of maps.

Let us recall that the challenge is to show that the limit of the subdivision process can be represented as the unique fixed point of some iterative process starting from a certain set of starting points (or base sequences). Below, we present two ways of approaching the non-uniform case.

6.1. First Approach - Using Staircases of Maps

Consider a general **linear** subdivision process, univariate or multivariate, stationary or non-stationary, uniform or non-uniform. Here we restrict the discussion to univariate binary subdivision. Starting with n_0 control points $p^0 \in \mathbb{R}^m$, there is an $n_1 \times n_0$ matrix $S^{[1]}$ that generates all the n_1 points derived at level 1 of the subdivision process

$$\bar{p}^1 = S^{[1]}\bar{p}^0.$$

Inductively, denote by $S^{[k]}$ the $n_k \times n_{k-1}$ matrix which represents the subdivision rules transforming the n_{k-1} points p^{k-1} attained at level $k-1$ to the n_k points p^k (in \mathbb{R}^m) at level k :

$$\bar{p}^k = S^{[k]}\bar{p}^{k-1}.$$

Note that here $n_k = \mathcal{O}(2^k)$. We have a sequence of metric spaces $\{\mathbb{R}^{n_k}\}$ (endowed with say the discrete L^2 -metric) and a sequence of maps between the metric spaces. If the subdivision process is known to be h -convergent, then the forward trajectory starting with p^0 converges to $p^\infty \subset \mathbb{R}^m$. However, different initial vectors yield different limit points. As done in the previous section, let us view the matrices $\{S^{[k]}\}$ as backward maps. For $k=1$,

$$S^{[1]} : \mathbb{R}^{\ell \times n_1} \rightarrow \mathbb{R}^{\ell \times m}, \quad A \in \mathbb{R}^{\ell \times n_1} \rightarrow AS^{[1]}\bar{p}^0 \in \mathbb{R}^{\ell \times m}, \quad (6.1)$$

and for $k > 1$,

$$S^{[k]} : \mathbb{R}^{\ell \times n_k} \rightarrow \mathbb{R}^{\ell \times n_{k-1}}, \quad A \in \mathbb{R}^{\ell \times n_k} \rightarrow AS^{[k]} \in \mathbb{R}^{\ell \times n_{k-1}}. \quad (6.2)$$

Of course, the above maps are only well defined for linear subdivision schemes.

6.1.1. The Choice of a Base Sequence. The question now is: For which class of base sequences $\bar{A} = \{A_k\}$ do all backward trajectories of the above maps converge to p^∞ ? The choice

$$A_k = I_{n_k \times n_k},$$

the identity matrix in $\mathbb{R}^{n_k \times n_k}$, generates the sequence $\{\bar{p}^k\}$ as its trajectory. Clearly, the backward trajectory with respect to this base sequence converges to p^∞ .

Definition 6.1. Let ρ be a positive integer and let C be a positive constant. For $n_k > \rho$, we denote by $K_C^{n_k, \rho-1}$ the set of all row vectors of length n_k of the form

$$(0, 0, \dots, 0, v, 0, 0, \dots, 0), \quad v \in K_C^{\rho-1}. \quad (6.3)$$

Definition 6.2 (Elements of the Base Sequence). Let ρ be a positive integer, C a positive constant, and $n_k > \rho$. We denote by $A_C^{\ell \times n_k, \rho}$ the set of all $\ell \times n_k$ matrices with rows in $K_C^{n_k, \rho-1}$ such that the sum of the rows does not have zero elements.

Let us further assume that the subdivision process converges to a continuous limit. Then, in the spirit of Theorem 5.2, we can prove the following.

Theorem 6.3. Consider a non-uniform C^ν -convergent subdivision scheme, $\nu > 0$, and let $\{S^{[k]}\}_{k=1}^\infty$ be the backward maps defined above. Then the backward trajectories starting with any base sequence $\bar{A} = \{A_k\}$ with $A_k \in A_C^{n_k \times n_k, \rho}$ for some fixed C and ρ , converge to a unique attractor whose rows constitute the limit curve (in \mathbb{R}^m) of the non-uniform scheme.

Proof. Unlike the case in Theorem 5.2, here the vectors of points \bar{p}^k are of fast increasing length, and as such, even if the subdivision is convergent, the vectors are not converging to a vector with identical rows. However, let us consider any partial vector w of \bar{p}^k containing ρ consecutive rows of \bar{p}^k . Such vectors tend to constant vectors as $k \rightarrow \infty$. Left multiplication of \bar{p}^k by a row vector of the form (6.3) corresponds to left multiplication of w by v . The result, as in Theorem 5.2, would be a point in \mathbb{R}^m approaching p^∞ . The rows of the matrices in the base sequences are all of the form (6.3). The condition that the sum of the rows does not have zero elements ensures that $A_k \bar{p}^k$ generates points in \mathbb{R}^m which tend to a dense set of points in p^∞ . \square

6.2. Second Approach - Using Trees of Maps

To present the idea, let us consider a univariate binary non-uniform subdivision scheme of fixed mask size. Being non-uniform means that the subdivision masks depend upon the geometric location of the control points, or, on the parametric correspondence of the control points at all levels. In other words, in order to generate the n_k points $\{p^k\}$ at level k , we use n_k different masks. As in the uniform case, we would like to attach to the subdivision process a related SFS, a binary SFS. We already know how to define a function system to a given linear subdivision rule. The question is, how to introduce a parametric correspondence into the definition of the system function systems and to the SFS iterations?

As in Section 2, we denote by $B^{[k]}$ the set of binary sequences of length k , i.e.,

$$B^{[k]} = \{\eta_k : \eta_k = \{i_j\}_{j=1}^k, i_j \in \{1, 2\}\}. \quad (6.4)$$

A parametric location at level k is determined by a binary sequence $\eta_k \in B^{[k]}$. In the non-uniform case the matrices $S_1^{[k]}$ and $S_2^{[k]}$ are location dependent. Instead of two matrices at each level we have 2^k matrices at level k , recursively defined as follows.

Definition 6.4 (Recursive Definition of the Matrices $S_{\eta_k}^{[k]}$). $S_1^{[1]}$ and $S_2^{[1]}$ are defined as in the uniform case, i.e., given the n_0 points, p^0 , at level zero, $S_1^{[1]}$ generates the first n_0 points at level 1 and $S_2^{[1]}$ generates the last n_0 points. We denote by p_{η_k} the n_0 points at level k attached to a parametric location η_k :

$$\bar{p}_{\eta_k} = S_{i_1 i_2 \dots i_k}^{[k]} \cdots S_{i_1 i_2 i_3}^{[3]} S_{i_1 i_2}^{[2]} S_{i_1}^{[1]} \bar{p}^0. \quad (6.5)$$

For $k > 1$, given the vector of points \bar{p}_{η_k} , we define two $n_0 \times n_0$ matrices $S_{i_1 i_2 \dots i_k 1}^{[k+1]}$ and $S_{i_1 i_2 \dots i_k 2}^{[k+1]}$. The first one generates the first n_0 points resulting from \bar{p}_{η_k} and the second generates the last n_0 points resulting from \bar{p}_{η_k} .

We hereby define a tree of maps $\{f_{\eta_k}\}$: For $k = 1$, let

$$f_r : \mathbb{R}^{n_0} \rightarrow \mathbb{R}^m, \quad f_r(A) := AS_r^{[1]} \bar{p}^0, \quad r = 1, 2, \quad (6.6)$$

where $\bar{p}^0 \in \mathbb{R}^{n_0 \times m}$ represents the set of n_0 initial control points in \mathbb{R}^m for the subdivision process.

For $k > 1$, let

$$f_{\eta_k} : \mathbb{R}^{n_0} \rightarrow \mathbb{R}^{n_0}, \quad f_{\eta_k}(A) := AS_{\eta_k}^{[k]}. \quad (6.7)$$

Consider the space $X := \mathbb{R}^{n_0}$ and the infinite binary tree of maps on X defined as above in (6.6) and (6.7). Note that for $k > 1$, $f_{\eta_k} : X \rightarrow X$, while for $k = 1$, $f_{\eta_k} : X \rightarrow \mathbb{R}^m$. Thus, convergent backward trajectories along paths in the tree converge to points in \mathbb{R}^m . Using the same steps as in the proof of Theorem 5.2 for each path of the tree, the following theorem holds.

Theorem 6.5. *Consider a linear binary non-uniform C^0 -convergent subdivision scheme and let TM be the tree of maps defined above. Then the backward trajectories along all paths in the tree converge for any $A \in K^{n_0-1}$, and the attractor U_{TM} of the tree constitutes the limit curve (in \mathbb{R}^m) of the non-uniform scheme.*

7. Summary and Conclusions

We considered countable sequences of maps on a complete metric space (X, d) and introduced the novel concept of an infinite tree of maps $X \rightarrow X$. Conditions for the convergence of these trees of maps to a unique attractor were derived. More generally, we investigated countable sequences of complete metric spaces $\{(X_i, d_i)\}$ and an associated countable sequence of maps $T_i : X_i \rightarrow X_{i-1}$. We provided conditions for the convergence of the backward trajectories $\Psi_k = T_1 \circ \cdots \circ T_k$, $k \in \mathbb{N}$, to a unique attractor. As an example, we considered trees of maps arising from iterated function systems and constructed a new class of fractals that are both scale dependent and location dependent. Finally, we exhibited the connections to non-stationary and non-uniform subdivision schemes and showed that this new approach is capable of linking all types of linear subdivision schemes to attractors of sequences of function systems.

References

- [1] M. F. Barnsley, *Fractals Everywhere*, Academic Press, Orlando, Florida (1988).
- [2] M. F. Barnsley, *SuperFractals*, Cambridge University Press (2006).
- [3] M. F. Barnsley, J. H. Elton, D. P. Hardin, *Recurrent iterated function systems*, *Constr. Approx.* 5(1) (1989), 3–31.
- [4] A. S. Cavaretta, W. Dahmen and C. A. Micchelli, *Stationary Subdivision*, *Mem. Amer. Math. Soc.* 93 (453), Providence, R.I. (1991).
- [5] C. Conti, N. Dyn, C. Manni, M.-L. Mazure, *Convergence of univariate non-stationary subdivision schemes via asymptotic similarity*, *Computer Aided Geometric Design* 37 (2015), 1–8.
- [6] N. Dyn, O. Kounchev, D. Levin, H. Render, *Regularity of generalized Daubechies wavelets reproducing exponential polynomials with real-valued parameters*, *Appl. Comput. Harm. Anal.* 37(2) (2014), 288–306.
- [7] N. Dyn, D. Levin, J. A. Gregory, *A 4-point interpolatory subdivision scheme for curve design*, *Computer Aided Geometric Design* 4(4), 257–268 (1987).
- [8] N. Dyn, D. Levin, *Analysis of asymptotically equivalent binary subdivision schemes*, *J. Math. Anal. Appl.* 193, (2), 594–621 (1995).
- [9] N. Dyn, D. Levin, *Subdivision schemes in geometric modelling*, *Acta Numer.*, 1–72 (2002).
- [10] N. Dyn, D. Levin, J. Yoon. *A new method for the analysis of univariate nonuniform subdivision schemes*, *Constr. Approx.* 40 (2) (2014), 173–188.
- [11] J. E. Hutchinson, *Fractals and self-similarity*, *Indiana Univ. Math. J.* 30 (1981), 713–747.
- [12] Y. Fisher, *Fractal Image Compression: Theory and Application*, Springer-Verlag, New York (1995).
- [13] D. Levin, *Using Laurent polynomial representation for the analysis of non-uniform binary subdivision schemes*, *Adv. Comput. Math.* 11 (1999) 41–54.
- [14] D. Levin, N. Dyn, P. Viswanathan, *Non-stationary Versions of Fixed-point Theory, with Applications to Fractals and Subdivision*, *J. Fixed Point Theory Appl.* 21 (2019), 1–25.
- [15] P. R. Massopust, *Fractal functions and their applications*, *Chaos, Solitons and Fractals* 8 (2) (1997), 171–190.
- [16] M. A. Navascues, *Fractal approximation*, *Complex Analysis and Operator Theory*, 4(4) (2010), 953–974.
- [17] H. Prautzsch, W. Boehm, M. Palusny, *Bézier and B-Spline Techniques*, Springer, Germany (2002).
- [18] S. Schaefer, D. Levin, R. Goldman, *Subdivision Schemes and Attractors*. In: M. Desbrun, H. Pottmann (eds) *Eurographics Symposium on Geometry Processing* (2005). Eurographics Association 2005, Aire-la-Ville Switzerland. ACM International Conference Proceeding Series 225 (2005), 171–180.
- [19] P. Viswanathan, A. K. B. Chand, *Fractal rational functions and their approximation properties*, *J. Approx. Th.* 185 (2014), 31–50.

Nira Dyn
School of Mathematical Sciences
Tel Aviv University
Israel
e-mail: niradyn@tau.ac.il

David Levin
School of Mathematical Sciences
Tel Aviv University
Israel
e-mail: levindd@gmail.com

Peter Massopust
Centre of Mathematics
Technical University of Munich
Germany
e-mail: massopust@ma.tum.de