

**CHARACTERIZATION OF FINITE DIMENSIONAL NILPOTENT
LIE ALGEBRAS BY THE DIMENSION OF THEIR SCHUR
MULTIPLIERS, $s(L) = 5$**

AFSANEH SHAMSAKI AND PEYMAN NIROOMAND

ABSTRACT. It is known that the dimension of the Schur multiplier of a non-abelian nilpotent Lie algebra L of dimension n is equal to $\frac{1}{2}(n-1)(n-2) + 1 - s(L)$ for some $s(L) \geq 0$. The structure of all nilpotent Lie algebras has been given for $s(L) \leq 4$ in several papers. Here, we are going to give the structure of all non-abelian nilpotent Lie algebras for $s(L) = 5$.

1. INTRODUCTION AND MOTIVATION

Let L be a finite dimensional nilpotent Lie algebra such that $L \cong F/R$ for a free Lie algebra F . Then by [1], the Schur multiplier $\mathcal{M}(L)$ of L is isomorphic to $R \cap F^2/[R, F]$. By a result of Moneyhun in [8], there exists a non-negative integer $t(L)$ such that $\dim \mathcal{M}(L) = \frac{1}{2}n(n-1) - t(L)$. It is a classical question to determine the structure of L by looking at the dimension of its Schur multiplier. The answer to this problem was given for $t(L) \leq 8$ in [2, 5, 6] and by putting some conditions on L for $t(L) \leq 16$ in [3] by several authors.

From [9], when L is a non-abelian nilpotent Lie algebra, the dimension of the Schur multiplier of L is equal to $\frac{1}{2}(n-1)(n-2) + 1 - s(L)$ for some $s(L) \geq 0$. It not only improves the bound of Moneyhun but also let us ask the same natural question about the characterization of Lie algebras in term of size $s(L)$. The answer to this question was given by several papers in [11, 17, 18] for $s(L) \leq 4$ and for $s(L) \leq 15$ when conditions are put on L in [19].

On the other hand, looking for instance [11] shows that the characterization of nilpotent Lie algebras by looking $s(L)$ causes to classification of nilpotent Lie algebras in terms of $t(L)$ by a simple and a shorter way. This paper is devoted to obtain the structure of all nilpotent Lie algebras L for $t(L) = 5$.

Throughout the paper, we may assume that L is a Lie algebra over an algebraically closed field of characteristic not equal to 2 and $A(n)$ and $H(m)$ are used to denote the abelian Lie algebra of dimension n and the Heisenberg Lie algebra of dimension $2m + 1$, respectively.

For the sake of convenience of reader some notations and terminology from [4, 5, 6, 7] are listed below.

Key words and phrases. Schur multiplier; nilpotent Lie algebra, capable Lie algebra.
Mathematics Subject Classification 2010. 17B30.

$L_{3,2} \cong H(1)$	with a basis $\{x_1, x_2, x_3\}$ and the multiplication $[x_1, x_2] = x_3$,
$L_{4,3} \cong L(3, 4, 1, 4)$	with a basis $\{x_1, \dots, x_4\}$ and the multiplication $[x_1, x_2] = x_3, [x_1, x_3] = x_4$,
$L_{5,5} \cong L(4, 5, 1, 6)$	with a basis $\{x_1, \dots, x_5\}$ and the multiplication $[x_1, x_2] = x_3, [x_1, x_3] = x_5, [x_2, x_4] = x_5$,
$L_{5,6} \cong L'(7, 5, 1, 7)$	with a basis $\{x_1, \dots, x_5\}$ and the multiplication $[x_1, x_2] = x_3, [x_1, x_3] = x_4, [x_1, x_4] = [x_2, x_3] = x_5$,
$L_{5,7} \cong L(7, 5, 1, 7)$	with a basis $\{x_1, \dots, x_5\}$ and the multiplication $[x_1, x_2] = x_3, [x_1, x_3] = x_4, [x_1, x_4] = x_5$,
$L_{5,8} \cong L(4, 5, 2, 4)$	with a basis $\{x_1, \dots, x_5\}$ and the multiplication $[x_1, x_2] = x_4, [x_1, x_3] = x_5$,
$L_{5,9} \cong L(7, 5, 2, 7)$	with a basis $\{x_1, \dots, x_5\}$ and the multiplication $[x_1, x_2] = x_3, [x_1, x_3] = x_4, [x_2, x_3] = x_5$,
$L_{6,10}$	with a basis $\{x_1, \dots, x_6\}$ and the multiplication $[x_1, x_2] = x_3, [x_1, x_3] = x_6, [x_4, x_5] = x_6$,
$L_{6,22}(\varepsilon)$	with a basis $\{x_1, \dots, x_6\}$ and the multiplication $[x_1, x_2] = x_5, [x_1, x_3] = x_6, [x_2, x_4] = \varepsilon x_6, [x_3, x_4] = x_5, \varepsilon \in \mathbb{F}$,
$L_1 \cong 27B$	with a basis $\{x_1, \dots, x_7\}$ and the multiplication $[x_1, x_2] = [x_3, x_4] = x_6, [x_1, x_5] = [x_2, x_3] = x_7$,
$L_2 \cong 27A$	with a basis $\{x_1, \dots, x_7\}$ and the multiplication $[x_1, x_2] = x_6, [x_1, x_4] = x_7, [x_3, x_5] = x_7$,
157	with a basis $\{x_1, \dots, x_7\}$ and the multiplication $[x_1, x_2] = x_3, [x_1, x_3] = [x_2, x_4] = [x_5, x_6] = x_7$
37B	with a basis $\{x_1, \dots, x_7\}$ and the multiplication $[x_1, x_2] = x_5, [x_2, x_3] = x_6, [x_3, x_4] = x_7$,
37C	with a basis $\{x_1, \dots, x_7\}$ and the multiplication $[x_1, x_2] = [x_3, x_4] = x_5, [x_2, x_3] = x_6, [x_2, x_4] = x_7$,
37D	with a basis $\{x_1, \dots, x_7\}$ and the multiplication $[x_1, x_2] = [x_3, x_4] = x_5, [x_1, x_3] = x_6, [x_2, x_4] = x_7$.

Main Theorem. *Let L be a non-abelian n -dimensional nilpotent Lie algebra. Then $s(L) = 5$ if and only if L is isomorphic to one of the Lie algebras $L(4, 5, 2, 4) \oplus A(4)$, $L(3, 4, 1, 4) \oplus A(3)$, $L(4, 5, 1, 6) \oplus A(2)$, $L_{6,22}(\varepsilon) \oplus A(2)$, $L_{6,26} \oplus A(1)$, $L_{6,10}$, $L_{6,23}$, $L_{6,25}$, 37B, 37C or 37D.*

We state some results without proof and refer the reader to see [10, 11, 14, 18].

Proposition 1.1. (See [12, Proposition 2.10]) *The Schur multiplier of Lie algebras $L_{6,22}(\varepsilon)$, $L_{5,8}$, L_1 and L_2 are abelian Lie algebras of dimension 8, 6, 9 and 10, respectively.*

A Lie algebra L is called capable provided that $L \cong H/Z(H)$ for a Lie algebra H . From [16, Definition 1.4], $Z^*(L)$ is used to denote the epicenter of L . The importance of $Z^*(L)$ is due to the fact that L is capable if and only if $Z^*(L) = 0$. Another notion having relation to the capability is the concept of the exterior center of a Lie algebra $Z^\wedge(L)$ which is introduced in [10]. It is known that from [10, Lemma 3.1]), $Z^*(L) = Z^\wedge(L)$.

Lemma 1.2. (See [18, Corollary 2.3]) *Let L be a non-capable nilpotent Lie algebra of dimension n such that $\dim L^2 \geq 2$. Then*

$$n - 3 < s(L).$$

Let \otimes_{mod} be used to denote the operator of usual tensor product of Lie algebras. Then

Theorem 1.3. (See [15, Theorem 2.1]) *Let L be a finite dimensional nilpotent Lie algebra non-abelian Lie algebra of class two. Then*

$$0 \rightarrow \ker g \rightarrow L^2 \otimes_{mod} L^{ab} \xrightarrow{g} \mathcal{M}(L) \rightarrow \mathcal{M}(L^{ab}) \rightarrow L^2 \rightarrow 0$$

is exact, in where

$g : x \otimes (z + L^2) \in L^2 \otimes_{mod} L^{ab} \mapsto [\bar{x}, \bar{z}] + [R, F] \in \mathcal{M}(L) = R \cap F^2/[R, F]$,
 $\pi(\bar{x} + R) = x$ and $\pi(\bar{z} + R) = z$. Moreover, the subalgebra $K = \langle [x, y] \otimes (z + L^2) + [z, x] \otimes (y + L^2) + [y, z] \otimes (x + L^2) \mid x, y, z \in L \rangle$ is contained in $\ker g$.

2. THE PROOF OF MAIN THEOREM

We begin with the following lemma that is easily proven.

Lemma 2.1. *There is no n -dimensional nilpotent Lie algebra with $s(L) = 5$, when*

- (i) $\dim L^2 \geq 4$;
- (ii) $\dim L^2 = 1$.

Proof.

- (i) Let L be a nilpotent Lie algebra such that $m = \dim L^2 \geq 4$ and $s(L) = 5$. [11, Theorem 3.1] and our assumption imply that

$$\frac{1}{2}(n-1)(n-2) - 4 = \dim \mathcal{M}(L) \leq \frac{1}{2}(n+m-2)(n-m-1) + 1 \leq \frac{1}{2}(n+2)(n-5) + 1.$$

It is a contradiction.

- (ii) By contrary. Let L be a Lie algebra such that $\dim L^2 = 1$ and $s(L) = 5$. Then by using [9, Lemma 3.3], $L \cong H(m) \oplus A(n-2m-1)$ for some $m \geq 1$. Looking [17, Corollary 2.5] shows that $s(L) = 0$ or $s(L) = 2$, when $m = 1$ or $m \geq 2$, respectively. It is a contradiction. Hence the result follows. \square

By using Lemma 2.1, we may assume that a nilpotent Lie algebras L with $s(L) = 5$ has $2 \leq \dim L^2 \leq 3$. First assume that $\dim L^2 = 2$.

Lemma 2.2. *Let L be an n -dimensional non-capable nilpotent Lie algebra of dimension at most 7 and $\dim L^2 = 2$. Then L is isomorphic to one of the Lie algebras $L_{6,10}$, L_2 or 157. Moreover, $s(L_{6,10}) = 5$ and $s(L_2) = s(157) = 6$.*

Proof. The proof is similar to [18, Theorem 2.6]. \square

Theorem 2.3. *Let L be an n -dimensional nilpotent Lie algebra with $s(L) = 5$ and $\dim L^2 = 2$. Then L is isomorphic to one of the Lie algebras $L(4, 5, 2, 4) \oplus A(4)$, $L(3, 4, 1, 4) \oplus A(3)$, $L(4, 5, 1, 6) \oplus A(2)$, $L_{6,22}(\varepsilon) \oplus A(2)$ or $L_{6,10}$.*

Proof. Since $\dim L^2 = 2$, L is nilpotent of class two or three. Let L be a Lie algebra of nilpotency class two. If L is a capable Lie algebra, then it should be isomorphic to one of the Lie algebras $L_{6,22}(\varepsilon) \oplus A$, $L_{5,8} \oplus A$ or $L_1 \oplus A$, for an abelian Lie algebra A by using [12, Corollary 2.13].

Case (i). Let $L \cong L_{6,22}(\varepsilon) \oplus A$. Proposition 1.1 implies $\dim M(L_{6,22}(\varepsilon)) = 8$. Since $5 = s(L) = \frac{1}{2}(n-1)(n-2) + 1 - \dim \mathcal{M}(L)$ and $\dim \mathcal{M}(L) = 8 + \frac{1}{2}(n-6)(n+1)$ by using [2, Theorem 1] and [8, Lemma 23], we have $n = 8$. Hence $L \cong L_{6,22}(\varepsilon) \oplus A(2)$.

Case (ii). Let now $L \cong L_{5,8} \oplus A$. We know from Proposition 1.1, $\dim \mathcal{M}(L_{5,8}) = 6$. Since $5 = s(L) = \frac{1}{2}(n-1)(n-2) + 1 - \dim \mathcal{M}(L)$ and $\dim \mathcal{M}(L) = 6 + \frac{1}{2}(n-5)n$ by using [2, Theorem 1] and [8, Lemma 23], we have $n = 9$. Therefore $L \cong L_{5,8} \oplus A(4) \cong L(4, 5, 2, 4) \oplus A(4)$.

Case (iii). Let $L \cong L_1 \oplus A$. We know $\dim \mathcal{M}(L_1) = 9$ by using Proposition 1.1. Since $5 = s(L) = \frac{1}{2}(n-1)(n-2) + 1 - \dim \mathcal{M}(L)$ and $\dim \mathcal{M}(L) = 9 + \frac{1}{2}(n-7)(n+2)$ by using [2, Theorem 1] and [8, Lemma 23], we have $n = 5$, which is contradiction. Thus L cannot be isomorphic to $L_1 \oplus A$.

Now let L be a Lie algebra of nilpotency class 3. If L is a capable Lie algebra, then it should be isomorphic to one of the Lie algebras $L_{4,3} \oplus A(n-4)$ or $L_{5,5} \oplus A(n-5)$ by using [13, Theorem 5.5].

Case (i). Let $L \cong L_{4,3} \oplus A(n-4)$. Since $\dim \mathcal{M}(L_{4,3}) = 2$ by using [5, Section 2], we have $\dim \mathcal{M}(L) = 2 + \frac{1}{2}(n-4)(n-1)$ by using [2, Theorem 1] and [8, Lemma 23]. Since $5 = s(L) = \frac{1}{2}(n-1)(n-2) + 1 - \dim \mathcal{M}(L)$ and $\dim \mathcal{M}(L) = 2 + \frac{1}{2}(n-4)(n-1)$, we have $n = 7$. Hence $L \cong L_{4,3} \oplus A(3) \cong L(3, 4, 1, 4) \oplus A(3)$.

Case (ii). Suppose $L \cong L_{5,5} \oplus A(n-5)$. [5, Section 3] shows that $\dim \mathcal{M}(L_{5,5}) = 4$. Now [2, Theorem 1] and [8, Lemma 23] imply that $\dim \mathcal{M}(L) = 4 + \frac{1}{2}n(n-5)$. Since $5 = s(L) = \frac{1}{2}(n-1)(n-2) + 1 - \dim \mathcal{M}(L)$ and $\dim \mathcal{M}(L) = 4 + \frac{1}{2}n(n-5)$, we have $n = 7$. Therefore $L \cong L_{5,5} \oplus A(2) \cong L(4, 5, 1, 6) \oplus A(2)$.

If L is a non-capable Lie algebra of nilpotency class 2 or 3, then by using Lemma 1.2, we have $n \leq 7$. Therefore $L \cong L_{6,10}$ by using Lemma 2.2. This completes the proof. \square

We now consider the case that $\dim L^2 = 3$. By looking all nilpotent Lie algebras listed in [4], we may choose all n -dimensional nilpotent Lie algebras L such that $\dim L^2 = 3$ for $n = 5$ or 6 in the Table 1.

Table 1.

Name	Nonzero multiplication
$L_{5,6}$	$[x_1, x_2] = x_3, [x_1, x_3] = x_4, [x_1, x_4] = [x_2, x_3] = x_5$
$L_{5,7}$	$[x_1, x_2] = x_3, [x_1, x_3] = x_4, [x_1, x_4] = x_5$
$L_{5,9}$	$[x_1, x_2] = x_3, [x_1, x_3] = x_4, [x_2, x_3] = x_5$
$L_{6,6}$	$[x_1, x_2] = x_3, [x_1, x_3] = x_4, [x_1, x_4] = [x_2, x_3] = x_5$
$L_{6,7}$	$[x_1, x_2] = x_3, [x_1, x_3] = x_4, [x_1, x_4] = x_5$
$L_{6,9}$	$[x_1, x_2] = x_3, [x_1, x_3] = x_4, [x_2, x_3] = x_5$
$L_{6,11}$	$[x_1, x_2] = x_3, [x_1, x_3] = x_4,$ $[x_1, x_4] = [x_2, x_3] = [x_2, x_5] = x_6$

continued on the next page

Table 1.

Name	Nonzero multiplication
$L_{6,12}$	$[x_1, x_2] = x_3, [x_1, x_3] = x_4, [x_1, x_4] = [x_2, x_5] = x_6$
$L_{6,13}$	$[x_1, x_2] = x_3, [x_1, x_3] = [x_2, x_4] = x_5,$ $[x_1, x_5] = [x_3, x_4] = x_6$
$L_{6,19}(\epsilon)$	$[x_1, x_2] = x_4, [x_1, x_3] = x_5, [x_1, x_5] = [x_2, x_4] = x_6,$ $[x_3, x_5] = \epsilon x_6$
$L_{6,20}$	$[x_1, x_2] = x_4, [x_1, x_3] = x_5, [x_1, x_5] = [x_2, x_4] = x_6$
$L_{6,23}$	$[x_1, x_2] = x_3, [x_1, x_3] = [x_2, x_4] = x_5, [x_1, x_4] = x_6$
$L_{6,24}(\epsilon)$	$[x_1, x_2] = x_3, [x_1, x_3] = [x_2, x_4] = x_5,$ $[x_1, x_4] = \epsilon x_6, [x_2, x_3] = x_6$
$L_{6,25}$	$[x_1, x_2] = x_3, [x_1, x_3] = x_5, [x_1, x_4] = x_6$
$L_{6,26}$	$[x_1, x_2] = x_4, [x_1, x_3] = x_5, [x_2, x_3] = x_6$

Assume L is nilpotent Lie algebra of dimension 7 such that $\dim L^2 = 3$. By looking the classification of all nilpotent Lie algebras in [7], L must be isomorphic to one of the Lie algebras listed at the Table 2 and 3.

Table 2. 7-dimensional indecomposable nilpotent Lie algebras

Name	Nonzero multiplication
37A	$[x_1, x_2] = x_5, [x_2, x_3] = x_6, [x_2, x_4] = x_7$
37B	$[x_1, x_2] = x_5, [x_2, x_3] = x_6, [x_3, x_4] = x_7$
37C	$[x_1, x_2] = [x_3, x_4] = x_5, [x_2, x_3] = x_6, [x_2, x_4] = x_7$
37D	$[x_1, x_2] = [x_3, x_4] = x_5, [x_1, x_3] = x_6, [x_2, x_4] = x_7$
257A	$[x_1, x_2] = x_3, [x_1, x_3] = [x_2, x_4] = x_6, [x_1, x_5] = x_7$
257B	$[x_1, x_2] = x_3, [x_1, x_3] = x_6, [x_1, x_4] = [x_2, x_5] = x_7$
257C	$[x_1, x_2] = x_3, [x_1, x_3] = [x_2, x_4] = x_6, [x_2, x_5] = x_7$
257D	$[x_1, x_2] = x_3, [x_1, x_3] = [x_2, x_4] = x_6,$ $[x_1, x_4] = [x_2, x_5] = x_7$
257E	$[x_1, x_2] = x_3, [x_1, x_3] = [x_4, x_5] = x_6, [x_2, x_4] = x_7$
257F	$[x_1, x_2] = x_3, [x_2, x_3] = [x_4, x_5] = x_6, [x_2, x_4] = x_7$

continued on the next page

Table 2. 7-dimensional indecomposable nilpotent Lie algebras

Name	Nonzero multiplication
257G	$[x_1, x_2] = x_3, [x_1, x_3] = [x_4, x_5] = x_6,$ $[x_1, x_5] = [x_2, x_4] = x_7$
257H	$[x_1, x_2] = x_3, [x_1, x_3] = [x_2, x_4] = x_6, [x_4, x_5] = x_7$
257I	$[x_1, x_2] = x_3, [x_1, x_3] = [x_1, x_4] = x_6,$ $[x_1, x_5] = [x_2, x_3] = x_7$
257J	$[x_1, x_2] = x_3, [x_1, x_3] = [x_2, x_4] = x_6, [x_1, x_5] = [x_2, x_3] = x_7$
257K	$[x_1, x_2] = x_3, [x_1, x_3] = x_6, [x_2, x_3] = [x_4, x_5] = x_7$
257L	$[x_1, x_2] = x_3, [x_1, x_3] = [x_2, x_4] = x_6,$ $[x_2, x_3] = [x_4, x_5] = x_7$
147A	$[x_1, x_2] = x_4, [x_1, x_3] = x_5,$ $[x_1, x_6] = [x_2, x_5] = [x_3, x_4] = x_7$
147B	$[x_1, x_2] = x_4, [x_1, x_3] = x_5,$ $[x_1, x_4] = [x_2, x_6] = [x_3, x_5] = x_7$
1457A	$[x_1, x_i] = x_{i+1} \ i = 2, 3, [x_1, x_4] = [x_5, x_6] = x_7$
1457B	$[x_1, x_i] = x_{i+1} \ i = 2, 3,$ $[x_1, x_4] = [x_2, x_3] = [x_5, x_6] = x_7$
137A	$[x_1, x_2] = x_5, [x_1, x_5] = [x_3, x_6] = x_7, [x_3, x_4] = x_6$
137B	$[x_1, x_2] = x_5, [x_3, x_4] = x_6,$ $[x_1, x_5] = [x_2, x_4] = [x_3, x_6] = x_7$
137C	$[x_1, x_2] = x_5, [x_1, x_4] = [x_2, x_3] = x_6,$ $[x_1, x_6] = x_7, [x_3, x_5] = -x_7$
137D	$[x_1, x_2] = x_5, [x_1, x_4] = [x_2, x_3] = x_6,$ $[x_1, x_6] = [x_2, x_4] = x_7, [x_3, x_5] = -x_7$
1357A	$[x_1, x_2] = x_4, [x_1, x_4] = [x_2, x_3] = x_5,$ $[x_1, x_5] = [x_2, x_6] = x_7, [x_3, x_4] = -x_7$
1357B	$[x_1, x_2] = x_4, [x_1, x_4] = [x_2, x_3] = x_5,$ $[x_1, x_5] = [x_3, x_6] = x_7, [x_3, x_4] = -x_7$
1357C	$[x_1, x_2] = x_4, [x_1, x_4] = [x_2, x_3] = x_5,$

continued on the next page

Table 2. 7-dimensional indecomposable nilpotent Lie algebras

Name	Nonzero multiplication
	$[x_1, x_5] = [x_2, x_4] = x_7, [x_3, x_4] = -x_7$

Table 3. 7-dimensional decomposable nilpotent Lie algebras

Name	Name
$L_{4,3} \oplus H(1)$	$L_{6,19}(\epsilon) \oplus A(1)$
$L_{5,6} \oplus A(2)$	$L_{6,20} \oplus A(1)$
$L_{5,7} \oplus A(2)$	$L_{6,23} \oplus A(1)$
$L_{5,9} \oplus A(2)$	$L_{6,24}(\epsilon) \oplus A(1)$
$L_{6,11} \oplus A(1)$	$L_{6,25} \oplus A(1)$
$L_{6,12} \oplus A(1)$	$L_{6,26} \oplus A(1)$
$L_{6,13} \oplus A(1)$	

We need the following lemma from [18, Lemma 2.7] for the proof of the Main Theorem.

Lemma 2.4. *Let L be an n -dimensional nilpotent Lie algebra such that $n = 5, 6$ or 7 , $\dim L^2 = \dim Z(L) = 3$ and $Z(L) = L^2$. Then the structure and the Schur multiplier of L are given in the following table.*

Table 4.

Name	$\dim \mathcal{M}(L)$	$s(L)$	Name	$\dim \mathcal{M}(L)$	$s(L)$
$L_{6,26}$	8	3	37C	11	5
37A	12	4	37D	11	5
37B	11	5			

Lemma 2.5. *Let L be a nilpotent Lie algebra of dimension at most 7 such that $\dim L^2 = 3$, $\dim Z(L) = 2$ and $Z(L) \subset L^2$. Then the structure and the Schur multiplier of L are given in the following table.*

Table 5.

Name	$\dim \mathcal{M}(L)$	$s(L)$	Name	$\dim \mathcal{M}(L)$	$s(L)$
$L_{5,9}$	3	4	257E	8	8
$L_{6,23}$	6	5	257F	9	7
$L_{6,24}(\epsilon)$	5	6	257G	8	8
$L_{6,25}$	6	5	257H	8	8
257A	9	7	257I	8	8

continued on the next page

Table 5.

<i>Name</i>	$\dim \mathcal{M}(L)$	$s(L)$	<i>Name</i>	$\dim \mathcal{M}(L)$	$s(L)$
257B	8	8	257J	8	8
257C	9	7	257K	6	10
257D	8	8	257L	6	10

Proof. The proof is similar to [18, Lemma 2.5]. \square

Lemma 2.6. *Let L be an n -dimensional nilpotent Lie algebra such that $n = 7$, $\dim L^2 = 3$ and $\dim Z(L) = 4$. Then the structure and the Schur multiplier of L are given in the following table.*

Table 6.

<i>Name</i>	$\dim \mathcal{M}(L)$	$s(L)$
$L_{5,9} \oplus A(2)$	8	8
$L_{6,26} \oplus A(1)$	11	5

Proof. Since $\dim Z(L) = 4$, L is isomorphic to $L_{5,9} \oplus A(2)$ or $L_{6,26} \oplus A(1)$ by searching in Tables 2 and 3. Let $L \cong L_{6,26} \oplus A(1)$. Since $\dim \mathcal{M}(L_{6,26}) = 8$ by using Table 4, we have $\dim \mathcal{M}(L) = 11$ by using [2, Theorem 1] and [8, Lemma 23]. Hence $s(L) = 5$. Also by using similar method, we can see $\dim \mathcal{M}(L_{5,9} \oplus A(2)) = 8$ and $s(L) = 8$. \square

Lemma 2.7. [18, Lemma 2.9] *Let L be an n -dimensional nilpotent Lie algebra such that $n = 5, 6$ or 7 , $\dim L^2 = 3$ and $\dim Z(L) = 1$. Then the structure and the Schur multiplier of L are given in the following table.*

Table 7.

<i>Name</i>	$\dim \mathcal{M}(L)$	$s(L)$	<i>Name</i>	$\dim \mathcal{M}(L)$	$s(L)$
$L_{5,6}$	3	4	1457A	6	10
$L_{5,7}$	3	4	1457B	6	10
$L_{6,11}$	5	6	137A	7	9
$L_{6,12}$	5	6	137B	7	9
$L_{6,13}$	4	7	137C	7	9
$L_{6,19}(\epsilon)$	5	6	137D	7	9
$L_{6,20}$	5	6	1357A	7	9

continued on the next page

Table 7.

Name	$\dim \mathcal{M}(L)$	$s(L)$	Name	$\dim \mathcal{M}(L)$	$s(L)$
147A	8	8	1357B	6	10
147B	8	8	1357C	6	10

Recall that a Lie algebra L is called generalized Heisenberg of rank n if $L^2 = Z(L)$ and $\dim L^2 = n$.

Lemma 2.8. *Let L be an n -dimensional generalized Heisenberg of rank 3 with $s(L) = 5$, then $n \leq 7$.*

Proof. By Theorem 1.3 we have $\dim \ker g = \dim \mathcal{M}(L^{ab}) - \dim L^2 + \dim L^{ab} \otimes_{\text{mod}} L^2 - \dim \mathcal{M}(L)$. Since $\dim \mathcal{M}(L) = \frac{1}{2}(n-1)(n-2) - 4$ and $\dim L^{ab} = n-3$, we have $\dim \ker g = n-3$.

By contrary let $n \geq 8$. Then $d = \dim L^{ab} = n-3 \geq 5$. Since $\dim L^2 = 3$, we can choose a basis $\{x_1 + L^2, \dots, x_d + L^2\}$ for L^{ab} such that $[x_1, x_2]$, $[x_2, x_3]$ and $[x_3, x_4]$ are non-trivial in L^2 . Thus

$$L^{ab} \otimes_{\text{mod}} L^2 \cong \bigoplus_{i=1}^d ((x_i + L^2)) \otimes_{\text{mod}} L^2.$$

Hence all elements of

$$\{[x_1, x_2] \otimes x_i + L^2 \oplus [x_i, x_1] \otimes x_2 + L^2 \oplus [x_2, x_i] \otimes x_1 + L^2, | 3 \leq i \leq d, i \neq 1, 2\}$$

and

$$\{[x_2, x_3] \otimes x_i + L^2 \oplus [x_i, x_3] \otimes x_2 + L^2 \oplus [x_3, x_i] \otimes x_2 + L^2, | 3 \leq i \leq d, i \neq 1, 2, 3\}$$

$$\{[x_3, x_4] \otimes x_i + L^2 \oplus [x_i, x_4] \otimes x_3 + L^2 \oplus [x_4, x_i] \otimes x_3 + L^2, | 3 \leq i \leq d, i \neq 2, 3, 4\}$$

are linearly independent and so $2(n-6) + n-5 \leq \ker g$. That is a contradiction for $n \geq 8$. Therefore the assumption is false and the result follows. \square

Let $c(L)$ be used to show the nilpotency class of L . Then

Lemma 2.9. *There is no nilpotent Lie algebra L with $\dim L^2 = 3$, $\dim Z(L) = 1$ and $s(L) = 5$ such that $L/Z(L) \cong L_{5,8} \oplus A(2)$.*

Proof. By contrary, let L be a nilpotent Lie algebra L with $\dim L^2 = 3$, $\dim Z(L) = 1$ and $s(L) = 5$ such that $L/Z(L) \cong L_{5,8} \oplus A(2)$. Then $\dim L = 8$ and $cl(L) = 3$. Since $cl(L) = 3$ and $\dim Z(L) = 1$, we have $L^3 = Z(L)$. On the other hand, $\dim \mathcal{M}(L) = \dim \mathcal{M}(L/Z(L)) + (\dim L/L^2 - 1) \dim Z(L) - \dim \ker \lambda_3$ and $\dim \ker \lambda_3 \geq 2$ by using proof [14, Theorem 1.1]. Thus

$$\dim \mathcal{M}(L) \leq \dim \mathcal{M}(L/Z(L)) + (\dim L/L^2 - 1) \dim Z(L) - 2$$

It is a contradiction. \square

Theorem 2.10. *Let L be an n -dimensional nilpotent Lie algebra with $s(L) = 5$ and $\dim L^2 = 3$. Then L is isomorphic to one of the Lie algebras $L_{6,23}$, $L_{6,25}$, 37B, 37C or 37D.*

Proof. First assume that $\dim Z(L) \geq 5$, or $\dim Z(L) = 3$ and $Z(L) \neq L^2$, or $\dim Z(L) = 2$ and $Z(L) \not\subset L^2$. We show that in these cases, there is no such a Lie algebra L of dimension n with $s(L) = 5$.

Let I be a central ideal of L of dimension one such that $L^2 \cap I = 0$. Since $\dim(L/I)^2 = 3$, by using [11, Theorem 3.1], we have

$$\dim \mathcal{M}(L/I) \leq \frac{1}{2}n(n-5) + 1.$$

If the equality holds, then

$$\frac{1}{2}(n-2)(n-3) + 1 - s(L/I) = \dim \mathcal{M}(L/I) = \frac{1}{2}n(n-5) + 1.$$

Therefore $s(L/I) = 3$ and by using [17, Theorem 3.2], there is no Lie algebra satisfying in $\dim(L/I)^2 = 3$. Thus [11, corollary 2.3] and our assumption imply

$$\dim \mathcal{M}(L) = \frac{1}{2}(n-1)(n-2) - 4 \leq \frac{1}{2}n(n-5) + (n-4),$$

which is a contradiction. Therefore we may assume that $\dim Z(L) = 4$, or $\dim Z(L) = 3$ and $L^2 = Z(L)$, or $\dim Z(L) = 2$ and $Z(L) \subset L^2$, or $\dim Z(L) = 1$.

If $\dim Z(L) = 4$, then there is a central ideal of L of dimension one such that $L^2 \cap I = 0$. Since $\dim(L/I)^2 = 3$, by using [11, Theorem 3.1], we have

$$\dim \mathcal{M}(L/I) \leq \frac{1}{2}n(n-5) + 1.$$

If the equality holds, then

$$\frac{1}{2}(n-2)(n-3) + 1 - s(L/I) = \dim \mathcal{M}(L/I) = \frac{1}{2}n(n-5) + 1.$$

Therefore $s(L/I) = 3$ and by using Table 4, $L/I \cong L_{6,26}$. Since $\dim Z(L) = 4$ and $\dim L = 7$, we have $L \cong L_{6,26} \oplus A(1)$ by using Lemma 2.6. Now let $\dim \mathcal{M}(L) \leq \frac{1}{2}n(n-5)$. Thus [11, corollary 2.3] and our assumption imply

$$\dim \mathcal{M}(L) = \frac{1}{2}(n-1)(n-2) - 4 \leq \frac{1}{2}n(n-5) + (n-4),$$

which is a contradiction.

If $\dim Z(L) = 3$ and $L^2 = Z(L)$, then L is isomorphic to one of the Lie algebras $37B$, $37C$ or $37D$ by using Lemmas 2.4 and 2.8.

Assume that $\dim Z(L) = 2$ and $Z(L) \subset L^2$. Then $\dim(L/Z(L))^2 = 1$. Since $L/Z(L)$ capable, by using [10, Theorem 3.5] and [9, Lemma 3.3], we have $L/Z(L) \cong H(1) \oplus A(n-5)$. Hence L is nilpotent of class 3. Therefore, by using [17, Theorem 2.6] for $c = 3$, we have

$$\dim L^3 + \frac{1}{2}(n-1)(n-2) - 3 \leq \dim \mathcal{M}(L/L^3) + \dim(L/Z_2(L) \otimes L^3).$$

Now since $1 \leq \dim L^3 \leq 2$, we can obtain that $n \leq 7$. Hence Lemma 2.5 implies that $L \cong L_{6,23}$ or $L \cong L_{6,25}$.

Finally, assume that $\dim Z(L) = 1$. Then $\dim(L/Z(L))^2 = 2$. By using [11, corollary 2.3], we have

$$\frac{1}{2}(n-1)(n-2) - 3 \leq \frac{1}{2}(n-2)(n-3) + 1 - s(L/Z(L)) + n - 3.$$

Thus $s(L/Z(L)) \leq 3$.

If $s(L/Z(L)) = 0$, then $L \cong H(1) \oplus A(n-4)$ by [11, Theorem 3.1]. This case cannot

occur, since $\dim(L/Z(L))^2 = 2$.

If $s(L/Z(L)) = 1$, then [9, Theorem 3.9] implies that $L \cong L(4, 5, 2, 4)$. Therefore $n = 6$.

If $s(L/Z(L)) = 2$, then $L/Z(L)$ is isomorphic to one of the Lie algebras $L(3, 4, 1, 4)$, $L(4, 5, 2, 4) \oplus A(1)$ or $H(m) \oplus A(n - 2m - 1)$ ($m \geq 2$) by using [9, Theorem 4.5]. In the case $L(3, 4, 1, 4)$ or $L(4, 5, 2, 4) \oplus A(1)$, we have $n = 5$ or 7 .

In the case $L/Z(L) \cong H(m) \oplus A(n - 2m - 1)$ ($m \geq 2$), then we have a contradiction, since $\dim(L/Z(L))^2 = 2$.

If $s(L/Z(L)) = 3$, $L/Z(L)$ is isomorphic to one of the Lie algebras $L(4, 5, 1, 6)$, $L(5, 6, 2, 7)$, $L'(5, 6, 2, 7)$, $L(7, 6, 2, 7)$, $L'(7, 6, 2, 7)$ or $L(3, 4, 1, 4) \oplus A(1)$ by [17, Main Theorem] and Lemma 2.9.

Hence $n = 5, 6$ or 7 when $\dim Z(L) = 1$. But there is no such Lie algebra by Lemma 2.7. This completes proof. \square

REFERENCES

- [1] Batten, P.; Multipliers and covers of Lie algebras. Thesis (Ph.D.)—North Carolina State University. ProQuest LLC, Ann Arbor, MI, 1993. 79 pp.
- [2] Batten, P., Moneyhun, K. and Stitzinger, E.; On characterizing nilpotent Lie algebras by their multipliers. *Comm. Algebra* 24 (1996), no. 14, 4319–4330.
- [3] Bosko, L. R.; On Schur multipliers of Lie algebras and groups of maximal class. *Internat. J. Algebra Comput.* 20 (2010), no. 6, 807–821.
- [4] De Graaf, W. A.; Classification of 6-dimensional nilpotent Lie algebras over fields of characteristic not 2. *J. Algebra* 309 (2006), no. 2, 640–653.
- [5] Hardy, P and Stitzinger, E.; On characterizing nilpotent Lie algebras by their multipliers, $t(L) = 3, 4, 5, 6$. *Comm. Algebra* 26(1998), no. 11, 3527–3539.
- [6] Hardy, P.; On characterizing nilpotent Lie algebras by their multipliers. III. *Comm. Algebra* 33 (2005), no. 11, 4205–4210.
- [7] Ming, P. G. ; Classification of nilpotent Lie algebras of dimension 7 (over algebraically closed fields and \mathbb{R}), ProQuest LLC, Ann Arbor, MI, 1998. Thesis (Ph.D.) University of Waterloo (Canada). MR2698220.
- [8] Moneyhun, K.; Isoclinisms in Lie algebras. *Algebras Groups Geom.* 11 (1994), no. 1, 9–22.
- [9] Niroomand, P.; On dimension of the Schur multiplier of nilpotent Lie algebras. *Cent. Eur. J. Math.* 9 (2011), no. 1, 57–64.
- [10] Niroomand, P., Parvizi, M. and Russo, F. G.; Some criteria for detecting capable Lie algebras. *J. Algebra* 384 (2013), 36–44.
- [11] Niroomand, P. and Russo, F. G.; A note on the Schur multiplier of a nilpotent Lie algebra. *Comm. Algebra* 39 (2013), no. 4, 1293–1297.
- [12] Niroomand, P., Johari, F. and Parvizi, M.; On the capability and Schur multiplier of nilpotent Lie algebra of class two. *Proc. Amer. Math. Soc.* 144 (2016), no. 10, 4157–4168.
- [13] Niroomand, P., Johari, F. and Parvizi, M.; Capable Lie algebras with the derived subalgebra of dimension two over an arbitrary field. *Linear Multilinear Algebra* 67 (2019), no. 3, 542–554.
- [14] P. Rai, On the dimension of the Schur multiplier of nilpotent Lie algebras. (2017) available at arXiv preprint arXiv:1705.03208.
- [15] Niroomand, P. and Johari, F.; Some results on the Schur multiplier of nilpotent Lie algebras. *J. Algebra.* 534 (2019) no. 15, 34–50.
- [16] Salemkar, A. R., Alamian, V. and Mohammadzadeh, H.; Some properties of the Schur multiplier and covers of Lie algebras. *Comm. Algebra* 36 (2008), no. 2, 697–707.
- [17] Saeedi, F., Arabyani, H. and Niroomand, P.; On dimension of Schur multiplier of nilpotent Lie algebras II. *Asian-European Journal of Mathematics* 10 (2016), no. 4, 1750076, 8pp.
- [18] Shamsaki, A., Niroomand, P., On characterizing nilpotent Lie algebra by their multiplier, $s(L) = 4$. To appear in *Rend. Circ. Mat. Palermo*, available at <https://doi.org/10.1007/s12215-019-00402-7>.
- [19] Shamsaki, A., Niroomand, P., On the Schur multipliers of Lie algebras of maximal class. *Internat. J. Algebra Comput.* 29 (2019) no. 5, 795–901.

SCHOOL OF MATHEMATICS AND COMPUTER SCIENCE, DAMGHAN UNIVERSITY, DAMGHAN, IRAN
Email address: `Shamsaki.Afsaneh@yahoo.com`

Email address: `niroomand@du.ac.ir`, `p_niroomand@yahoo.com`

SCHOOL OF MATHEMATICS AND COMPUTER SCIENCE, DAMGHAN UNIVERSITY, DAMGHAN, IRAN