

BOOLEAN FIP RING EXTENSIONS

GABRIEL PICAUVET AND MARTINE PICAUVET-L'HERMITTE

ABSTRACT. We characterize extensions of commutative rings $R \subseteq S$ whose sets of subextensions $[R, S]$ are finite (i.e. $R \subseteq S$ has the FIP property) and are Boolean lattices, that we call Boolean FIP extensions. Some characterizations involve “factorial” properties of the poset $[R, S]$. A non trivial result is that each subextension of a Boolean FIP extension is simple (i.e. $R \subseteq S$ is a simple pair).

1. INTRODUCTION AND NOTATION

We consider the category of commutative and unital rings, whose epimorphisms will be involved. If $R \subseteq S$ is a (ring) extension, we denote by $[R, S]$ the set of all R -subalgebras of S and set $]R, S[:= [R, S] \setminus \{R, S\}$ (with a similar definition for $[R, S[$ or $]R, S]$).

A *lattice* is a poset L such that every pair $a, b \in L$ has a supremum and an infimum. For an extension $R \subseteq S$, the poset $([R, S], \subseteq)$ is a *complete* lattice where the supremum of any non void subset is the compositum which we call *product* from now on and denote by Π when necessary, and the infimum of any non void subset is the intersection. We are aiming to study some lattice properties of the poset $([R, S], \subseteq)$, mainly the Boolean property. As a general rule, an extension $R \subseteq S$ is said to have some property of lattices if $[R, S]$ has this property.

The extension $R \subseteq S$ is said to have FIP (for the “finitely many intermediate algebras property”) or an FIP extension if $[R, S]$ is finite. A *chain* of R -subalgebras of S is a set of elements of $[R, S]$ that are pairwise comparable with respect to inclusion. When $[R, S]$ is a chain, the extension $R \subseteq S$ is called a λ -*extension* by some authors. We will say that $R \subseteq S$ is chained. We also say that the extension $R \subseteq S$ has FCP (or is an FCP extension) if each chain in $[R, S]$ is finite. Clearly, each extension that satisfies FIP must also satisfy FCP. Dobbs and the authors characterized FCP and FIP extensions [11].

2010 *Mathematics Subject Classification*. Primary:13B02,13B21, 13B22, 06E05, 06C10; Secondary: 13B30, 12F10, 06E25.

Key words and phrases. FIP, FCP extension, minimal extension, integral extension, support of a module, lattice, Boolean lattice, algebraic field extension, Galois extension.

Let $R \subseteq S$ be an FCP extension, then $[R, S]$ is a complete Noetherian Artinian lattice, with R as the least element and S as the largest element. We use lattice definitions and properties described in [24].

Our main tool are the minimal (ring) extensions, a concept that was introduced by Ferrand-Olivier [18]. Recall that an extension $R \subseteq S$ is called *minimal* if $[R, S] = \{R, S\}$. An extension $R \subseteq S$ is called a *simple* extension if $S = R[t]$ for some $t \in S$ and a *simple pair* if $R \subseteq T$ is a simple extension for each $T \in [R, S]$. A minimal extension is simple. The key connection between the above ideas is that if $R \subseteq S$ has FCP, then any maximal (necessarily finite) chain of R -subalgebras of S , $R = R_0 \subset R_1 \subset \cdots \subset R_{n-1} \subset R_n = S$, with *length* $n < \infty$, results from juxtaposing n minimal extensions $R_i \subset R_{i+1}$, $0 \leq i \leq n - 1$. An FCP extension is finitely generated, and (module) finite if integral. For any extension $R \subseteq S$, the *length* $\ell[R, S]$ of $[R, S]$ is the supremum of the lengths of chains of R -subalgebras of S . Notice that if $R \subseteq S$ has FCP, then there *does* exist some maximal chain of R -subalgebras of S with length $\ell[R, S]$ [12, Theorem 4.11].

1.1. A summary of the main results. Any undefined material is explained at the end of the section or in the next sections.

Section 2 is devoted to some general properties of lattices $[R, S]$, mainly in the context of FCP and FIP extensions. Since Boolean extensions are distributive, we have evidently to work on distributive extensions, which is done in this section. We discuss the decomposition of elements of $[R, S]$ into irreducible elements. When $[R, S]$ has finitely many atoms and each element of $[R, S]$ is a product of atoms, then Proposition 2.17 shows that $R \subseteq S$ has FIP and is almost-Prüfer, and Theorem 2.11 shows that $R \subseteq S$ is a simple pair.

For extensions $R \subseteq S$ of integral domains, Ayache considered *Boolean lattices* (also called Boolean algebras) $([R, S], \cap, \cdot)$, that are distributive lattices such that each $T \in [R, S]$ has a (unique) complement [1], [2] and [3]. In particular, [2, Problem 45] asked under which conditions $[R, S]$ is a finite Boolean lattice. This question is completely answered in Sections 3 and 4, where we get in Theorem 3.29 a characterization of Boolean extensions. In particular, Theorem 3.1 shows that an FCP Boolean extension $R \subseteq S$ has FIP, each element of $[R, S]$ has a unique representation by a finite product of atoms, and $R \subseteq S$ a simple pair. Section 3 is devoted to the study of arbitrary FIP extensions. The canonical decomposition of a ring extension is crucial. It consists of the tower $R \subseteq {}^+_S R \subseteq {}^t_S R \subseteq \overline{R} \subseteq S$, where ${}^+_S R$ (resp. ${}^t_S R$) is the *seminormalization* (resp. *t-closure*) of R in S (see Section 3). This decomposition allows us to only consider special extensions: subintegral,

seminormal infra-integral, t -closed and integrally closed. The t -closed case is reduced to the context of field extensions and is the subject of Section 4. In particular, for a field extension $k \subset L$ with separable closure T and radical closure U such that $U, T \notin \{k, L\}$, Theorem 4.2 shows that $k \subset L$ is Boolean if and only if $k \subset U$ and $T \subset L$ are minimal, $[k, L] = [k, T] \cup [U, L]$, $k \subset T$ and $k \subset U$ are linearly disjoint and $[k, T]$ is a Boolean lattice. Boolean separable field extensions needs a special study. A striking result is Theorem 4.19: A Galois finite extension (hence FIP) $k \subset L$ is Boolean if and only if $k \subset L$ is a cyclic extension whose dimension is square free.

1.2. Some conventions and notation. A *local* ring is here what is called elsewhere a quasi-local ring. As usual, $\text{Spec}(R)$ and $\text{Max}(R)$ are the set of prime and maximal ideals of a ring R . The support of an R -module E is $\text{Supp}_R(E) := \{P \in \text{Spec}(R) \mid E_P \neq 0\}$, and $\text{MSupp}_R(E) := \text{Supp}_R(E) \cap \text{Max}(R)$. When $R \subseteq S$ is an extension, we will set $\text{Supp}_R(T/R) := \text{Supp}(T/R)$ and $\text{Supp}_R(S/T) := \text{Supp}(S/T)$ for each $T \in [R, S]$, unless otherwise specified.

If $R \subseteq S$ is a ring extension and $P \in \text{Spec}(R)$, then S_P is both the localization $S_{R \setminus P}$ as a ring and the localization at P of the R -module S . We denote by $\kappa_R(P)$ the residual field R_P/PR_P at P . An extension $R \subseteq S$ is called *locally minimal* if $R_P \subset S_P$ is minimal for each $P \in \text{Supp}(S/R)$ or equivalently for each $P \in \text{MSupp}(S/R)$.

We denote by $(R : S)$ the conductor of $R \subseteq S$. The integral closure of R in S is denoted by \overline{R}^S (or by \overline{R} if no confusion can occur).

Recall that an extension $R \subseteq S$ is *Prüfer* (or a normal pair) if $R \subseteq T$ is a flat epimorphism for each $T \in [R, S]$. The *Prüfer hull* of an extension $R \subseteq S$ is the greatest *Prüfer* subextension \widetilde{R} of $[R, S]$ [28]. An extension $R \subseteq S$ is called *almost-Prüfer* if $\widetilde{R} \subseteq S$ is integral, or equivalently, when $R \subseteq S$ is FIP, if $S = \widetilde{R}\overline{R}$ [30, Theorem 4.6].

A poset (X, \leq) is called a *tree* if $x_1, x_2 \leq x_3$ in X implies that x_1 and x_2 are comparable (with respect to \leq). We also say that (X, \leq) is *treed*. A subset Y of X is called an *antichain* if no two distinct elements of Y are comparable.

Finally, $|X|$ is the cardinality of a set X , \subset denotes proper inclusion and, for a positive integer n , we set $\mathbb{N}_n := \{1, \dots, n\}$. The characteristic of an integral domain k is denoted by $c(k)$. For a, b, c in a ring R , if c divides $a - b$, we write $a \equiv b \pmod{c}$.

2. LATTICES PROPERTIES OF THE POSET $[R, S]$

2.1. Some definitions on the lattice $[R, S]$. In the context of a lattice $[R, S]$, some definitions and properties of lattices have the following formulations.

An element $T \in [R, S]$ is called \cap -irreducible (resp.; Π -irreducible) (see [24]) if $T = T_1 \cap T_2$ (resp.; $T = T_1 T_2$) implies $T = T_1$ or $T = T_2$.

An element T of $[R, S]$ is an *atom* (resp.; *co-atom*) if and only if $R \subset T$ (resp.; $T \subset S$) is a minimal extension. Therefore, an atom (resp.; co-atom) is Π -irreducible (resp.; \cap -irreducible). We denote by \mathcal{A} the set of atoms of $[R, S]$ and by \mathcal{CA} the set of co-atoms of $[R, S]$.

Now $R \subset S$ is called:

(a) *atomic* (resp.; *atomistic*) if each $T \in]R, S[$ contains some atom (resp.; is the product of atoms (contained in T)) [32, page 80].

(b) *co-atomic* (resp.; *co-atomistic*) if each $T \in [R, S[$ is contained in some co-atom (resp.; is the intersection of co-atoms (containing T)).

(c) *distributive* if intersection and product are each distributive with respect to the other. Actually, each distributivity implies the other [24, Exercise 5, page 33].

(d) *factorial* (resp.; *co-factorial*) if each element of $[R, S]$ has a unique irredundant representation by a finite product of atoms (resp.; a unique irredundant representation by a finite intersection of co-atoms.)

An FCP extension is both atomic and co-atomic.

We introduce a definition reminiscent of arithmetic rings [29].

Definition 2.1. A ring extension $R \subseteq S$ is called *arithmetic* if $[R_P, S_P]$ is a chain for each $P \in \text{Spec}(R)$.

Example 2.2. The extension $R \subset S$ is arithmetic in the following cases ([29, Example 5.13] for (2), (3) and (4)):

- (1) $R \subset S$ is locally minimal.
- (2) $R \subset S$ has FCP and is integrally closed.
- (3) $R \subset S$ is FIP subintegral and $|R/M| = \infty$ for each $M \in \text{MSupp}(S/R)$.
- (4) $R \subset S$ is FIP t -closed integral such that $R_M/MR_M \subset S_M/MR_M$ is radicial for each $M \in \text{MSupp}(S/R)$.
- (5) We also have examples in [31, Theorem 6.1] of arithmetic extensions of length 2 $R \subset S$ when $|[R, S]| = 3$.

We proved in [29, Proposition 5.18]:

Proposition 2.3. *An arithmetic extension is distributive.*

The following proposition will make easier many proofs.

Proposition 2.4. *Let $R \subseteq S$ be a ring extension. The following statements are equivalent:*

- (1) $R \subseteq S$ is distributive ;
- (2) $R_M \subseteq S_M$ is distributive for each $M \in \text{MSupp}(S/R)$;
- (3) $R_P \subseteq S_P$ is distributive for each $P \in \text{Supp}(S/R)$;
- (4) $R/I \subseteq S/I$ is distributive for each ideal I shared by R and S .
- (5) $R/I \subseteq S/I$ is distributive for some ideal I shared by R and S .

Proof. We have obviously (1) \Rightarrow (3) \Rightarrow (2) and (1) \Leftrightarrow (4) \Rightarrow (5) \Rightarrow (1). Conversely, assume that $R_M \subseteq S_M$ is distributive for each $M \in \text{MSupp}(S/R)$. Then, $R_M \subseteq S_M$ is distributive for each $M \in \text{Max}(R)$. It follows that the distributivity property holds in $[R, S]$ since it holds in any $[R_M, S_M]$. \square

Proposition 2.5. [21, Theorem 1, p. 172] *In a distributive lattice of finite length, all maximal chains between two comparable elements have the same length (the Jordan-Hölder chain condition or condition (JH)).*


In [16], Dobbs and Shapiro defined a field extension $k \subset L$ of finite length to be *catenarian* if all maximal chains of fields between k and L have the same length, and get examples of catenarian field extensions. It follows that a distributive field extension is catenarian.

Proposition 2.6. [8, Remarks, page 9 and Theorem 1.7] *A distributive extension $R \subset S$ satisfies the upper covering condition (UCC), which means that for each $T, U \in [R, S]$ such that $T \cap U \subset T$ is minimal, then $U \subset TU$ is minimal.*


2.2. Some distributive extensions are simple. We are going to show that some special subextensions of an FCP distributive extension are simple. Before, next lemma is needed.

Lemma 2.7. *If $R \subset S$ is a distributive extension, then $R[x, y] = R[x + y]$ whenever $y \in S \setminus R$, $x \in S \setminus R[y]$ and $R \subset R[x]$ is minimal.*

Proof. Consider the diagram



Since $R[x]R[y] = R[x, y]$ and $R \subset R[x]$ is minimal, we get that $R[y] \subset R[x, y]$ is minimal by UCC. There is another diagram



where $R[x] \cap (R[y] \cap R[x + y]) = R$ and $R \subset R[x]$ is minimal. Using again UCC, we get that $R[y] \cap R[x + y] \subset R[x](R[y] \cap R[x + y])$ is minimal. But $R[x](R[y] \cap R[x + y]) = R[x]R[y] \cap R[x]R[x + y] = R[x, y]$ by distributivity, so that $R[y] \cap R[x + y] \subset R[x, y]$ is minimal. From $R[y] \cap R[x + y] \subseteq R[y] \subset R[x, y]$, we deduce $R[y] = R[y] \cap R[x + y]$, which implies $R[y] \subset R[x + y] \subseteq R[x, y]$ because $x \notin R[y]$, and then $R[x + y] = R[x, y]$. \square

Proposition 2.8. *Let $R \subset S$ be a distributive extension. Let $T \in [R, S]$ be a product of finitely many atoms. Then, $R \subset T$ is simple. More precisely, if $T = \prod_{i=1}^n R[x_i]$, where the $R \subset R[x_i]$ are minimal distinct extensions, then, $T = R[\sum_{i=1}^n x_i]$.*

Proof. We prove the two statements by induction on n . There is nothing to prove when $n = 1$. Assume that the induction hypothesis holds for $n - 1$ and set $T' := \prod_{i=1}^{n-1} R[x_i] \neq R$, so that $T' = R[x]$ with $x := \sum_{i=1}^{n-1} x_i$. Then, $T = R[x]R[x_n]$, with $x \notin R$ and $x_n \in S \setminus R[x]$. Deny. Then $R[x_n] \subseteq \prod_{i=1}^{n-1} R[x_i]$ would imply $R[x_n] \subseteq R[x_i]$ for some $i \in \{1, \dots, n-1\}$, a contradiction [32, Theorem 4.30]. Now, use Lemma 2.7 to get the result. \square

2.3. The lattice $[R, S]$ for an FCP or FIP extension.

Proposition 2.9. [24, Proposition 1.4.4] *If $R \subseteq S$ has FCP, then any $T \in [R, S]$ is a finite intersection (resp.; product) of \cap -irreducible (resp.; Π -irreducible) elements of $[R, S]$.*

Lemma 2.10. *If $R \subseteq S$ is a distributive extension and $T \in [R, S]$ has an irredundant representation $T = T_1 \cdots T_m$ by Π -irreducible elements of $[R, S]$, (resp.; $T = U_1 \cap \cdots \cap T_r$ by \cap -irreducible elements of $[R, S]$) (for example by atoms (resp.; co-atoms)), the representation is unique.*

If in addition $R \subseteq S$ has FCP, $[R, S]$ has exactly n Π -irreducible (resp.; \cap -irreducible) elements if and only if $\ell[R, S] = n$. If these conditions hold, $R \subseteq S$ has FIP.

Proof. [32, Theorem 4.30], [17, Theorem 147] and [23, Lemma 4, page 486] combine to yield the result. The FIP property is then an easy consequence. \square

Theorem 2.11. *An extension $R \subset S$ is atomistic, distributive and FIP if and only if $R \subset S$ is factorial (respectively, is co-factorial). If these conditions hold, then $R \subset S$ is a simple pair.*

Proof. If $R \subset S$ is an atomistic distributive FIP extension, then $[R, S]$ has finitely many atoms, and then is factorial by Lemma 2.10.

Conversely, assume that $R \subset S$ is factorial, whence atomistic. Let $T, U, V \in [R, S]$. Obviously, $(T \cap U)(T \cap V) \subseteq T \cap UV$. Since $R \subset S$ is factorial, we can write $T = \prod_{\alpha \in I} A_\alpha$, $U = \prod_{\beta \in J} A_\beta$ and $V = \prod_{\gamma \in K} A_\gamma$, for finite subsets $\{A_\alpha \mid \alpha \in I\}$, $\{A_\beta \mid \beta \in J\}$ and $\{A_\gamma \mid \gamma \in K\}$ of \mathcal{A} . Then, $UV = \prod_{\beta \in J \cup K} A_\beta$, so that $T \cap UV = (\prod_{\alpha \in I} A_\alpha) \cap (\prod_{\beta \in J \cup K} A_\beta)$. Write $T \cap UV =: \prod_{\delta \in L} A_\delta = (\prod_{\alpha \in I} A_\alpha) \cap (\prod_{\beta \in J \cup K} A_\beta)$. Then, for any $\delta \in L$, we have $A_\delta \subseteq T$ and $A_\delta \subseteq \prod_{\beta \in J \cup K} A_\beta$, so that there exist some $\alpha \in I$ such that $A_\delta = A_\alpha$ and some $\beta \in J \cup K$ such that $A_\delta = A_\beta$. If $\beta \in J$, then $A_\delta \subseteq U$, so that $A_\delta \subseteq T \cap U \subseteq (T \cap U)(T \cap V)$. If $\beta \in K$, then $A_\delta \subseteq V$, so that $A_\delta \subseteq T \cap V \subseteq (T \cap U)(T \cap V)$. In both cases, $A_\delta \subseteq (T \cap U)(T \cap V)$, which yields $T \cap UV \subseteq (T \cap U)(T \cap V)$, and then $T \cap UV = (T \cap U)(T \cap V)$. Therefore, $R \subset S$ is distributive. Moreover, $R \subset S$ has FIP since \mathcal{A} is finite (S is the product of all elements of \mathcal{A}).

Now, let $R \subset S$ be factorial, whence distributive. Set $n := |\mathcal{A}|$, and for $A_\alpha \in \mathcal{A}$, set $B_\alpha := \prod_{\beta \neq \alpha} A_\beta$. Obviously, $\mathcal{C}\mathcal{A} = \{B_\alpha \mid \alpha \in \mathbb{N}_n\}$. Let $T \in [R, S]$, with $T = \prod_{\alpha \in I} A_\alpha$. For $J = \mathbb{N}_n \setminus I$, an easy calculation shows that $T = \bigcap_{\beta \in J} B_\beta$ in a unique way. Hence, $R \subset S$ is co-factorial.

To end, assume that $R \subset S$ is co-factorial. We get that $R \subset S$ is factorial and distributive, mimicking the previous proof. It is enough to exchange product and intersection, and atoms and co-atoms. In fact, we use the fact that $R \subset S$ is co-atomistic. If these conditions hold, then $R \subset S$ is a simple pair by Proposition 2.8. \square

The following notions and results are deeply involved in the sequel.

Definition 2.12. [7, Theorem 4.5] An extension $R \subset S$ is called *M-crucial* if there is some unique $M \in \text{Max}(R)$, called the **crucial (maximal) ideal** $\mathcal{C}(R, S)$ of $R \subset S$, such that $R_P = S_P$ for each $P \in \text{Spec}(R) \setminus \{M\}$.

Theorem 2.13. [18, Théorème 2.2] *A minimal extension is crucial and is either integral (finite) or a flat epimorphism.*

Lemma 2.14. [11, Corollary 3.2] *If there exists a maximal chain $R = R_0 \subset \dots \subset R_i \subset \dots \subset R_n = S$ of extensions, where $R_i \subset R_{i+1}$ is minimal, then $\text{Supp}(S/R) = \{\mathcal{C}(R_i, R_{i+1}) \cap R \mid i = 0, \dots, n-1\}$.*

Lemma 2.15. [31, Lemma 1.8] *Let $R \subset S$ be an FCP extension. and $M \in \text{MSupp}(S/R)$. There is some $T \in [R, S]$ such that $R \subset T$ is minimal and $\mathcal{C}(R, T) = M$.*

If $R \subseteq S$ has FCP, set $\mathcal{T} := \{T \in [R, S] \mid |\text{MSupp}_R(T/R)| = 1\}$ and $\mathcal{T}_M := \{T \in \mathcal{T} \mid \text{MSupp}_R(T/R) = \{M\}\}$. We are able to give dual results with $\mathcal{T}^* := \{T \in [R, S] \mid |\text{MSupp}_R(S/T)| = 1\}$, but they do not appear in this paper because they are not used in the sequel.

Proposition 2.16. *A subextension $R \subset T$ of an FCP extension $R \subset S$ is M -crucial when $T \in \mathcal{T}_M$ is such that $\text{Supp}(T/R) \subseteq \text{Max}(R)$. Moreover, \mathcal{A} is the set of all minimal elements of \mathcal{T} .*

In case $R \subset S$ has FIP, \mathcal{T}_M has a greatest element $s(M) := \prod_{T \in \mathcal{T}_M} T$.

Proof. Let $T \in \mathcal{T}_M$ be such that $\text{Supp}(T/R) \subseteq \text{Max}(R)$. It follows that $\text{Supp}_R(T/R) = \text{MSupp}(T/R) = \{M\}$ and $R_P = T_P$ for each $P \in \text{Spec}(R) \setminus \{M\}$. Hence $R \subset T$ is M -crucial. If $U \in [R, S]$ is an atom, $R \subset U$ is minimal and $M := \mathcal{C}(R, U)$ with $\text{MSupp}(U/R) = \{M\}$, whence $U \in \mathcal{T}$. Since $R \subset U$ is minimal, U is a minimal element of \mathcal{T} . Conversely, let U be a minimal element of \mathcal{T} . If U is not an atom, there is $U' \in [R, S]$ with $R \subset U' \subset U$. Then, $\emptyset \neq \text{MSupp}(U'/R) \subseteq \text{MSupp}(U/R) = \{M\}$ implies $\text{MSupp}(U'/R) = \{M\}$ giving $U' \in \mathcal{T}$, contradicting the minimality of U in \mathcal{T} . Therefore, U is an atom.

Since $s(M) := \prod_{T \in \mathcal{T}_M} T$, we get that $T \subseteq s(M)$ for each $T \in \mathcal{T}_M$ and $R \subset s(M)$, whence $\text{MSupp}(s(M)/R) \neq \emptyset$. Since $R_P = T_P$ for each $T \in \mathcal{T}_M$ and $P \in \text{Spec}(R) \setminus \{M\}$, we get $s(M)_P = R_P$, so that $\text{MSupp}(s(M)/R) = \{M\}$ and $s(M)$ is the greatest element of \mathcal{T}_M . \square

Proposition 2.17. *Let $R \subset S$ be an FCP atomistic extension such that $|\mathcal{A}| < \infty$, with $\text{MSupp}(S/R) = \{M_1, \dots, M_n\}$. Set $\mathcal{A}_M = \mathcal{A} \cap \mathcal{T}_M$ and, for each $k \in \mathbb{N}_n$, $V_k := \prod_{i=1}^k s(M_i)$ and $V_0 := R$.*

Then, the following statements hold.

- (1) $R \subset S$ has FIP.
- (2) For each $M \in \text{MSupp}(S/R)$, $s(M) = \prod_{A \in \mathcal{A}_M} A$.
- (3) For each $k \in \mathbb{N}_n$, $\text{Supp}(V_k/V_{k-1}) = \{M_k\}$ and $\{V_k\}_{k=0}^n$ is an increasing chain such that $V_n = S$.
- (4) For each $T \in]R, S]$, there exists $T_i \in \mathcal{T}_{M_i}$, for each $M_i \in \text{MSupp}(T/R)$, such that $T = \prod_{M_i \in \text{MSupp}(T/R)} T_i$.
- (5) $R \subset S$ is almost-Prüfer.

Proof. (1) Since $|\mathcal{A}| < \infty$ and any element of $[R, S]$ is a product of atoms, then $|[R, S]| < \infty$ and $R \subset S$ has FIP.

(2) Let $M \in \text{MSupp}(S/R)$. Any element of \mathcal{T}_M is a product of atoms, which are necessarily in \mathcal{T}_M , and then in \mathcal{A}_M . It follows from Proposition 2.16 that $s(M) = \prod_{A \in \mathcal{A}_M} A$.

(3) Let $M_j \in \text{MSupp}(S/R)$ for some $j \in \mathbb{N}_n$. Assume $1 < k < n$. Since $V_k = \prod_{i=1}^k s(M_i)$ and $V_{k-1} = \prod_{i=1}^{k-1} s(M_i)$, we have $(V_{k-1})_{M_j} = (V_k)_{M_j} = R_{M_j}$ if $j > k$, so that $M_j \notin \text{Supp}(V_k/V_{k-1})$. If $j = k > k-1$, then, $(V_{k-1})_{M_k} = R_{M_k}$ and $(V_k)_{M_k} = s(M_k)_{M_k} \neq R_{M_k}$, so that $M_k \in \text{Supp}(V_k/V_{k-1})$. At last, if $j < k$, then, $(V_{k-1})_{M_j} = (V_k)_{M_j} = s(M_j)_{M_j}$, so that $M_j \notin \text{Supp}(V_k/V_{k-1})$. Hence, $\text{Supp}(V_k/V_{k-1}) = \{M_k\}$. If $k \in \{1, n\}$, the same reasoning holds. The end of (3) is obvious.

(4) Let $T \in [R, S]$. Let $I \subset \mathbb{N}_n$ be such that $\text{Supp}(T/R) = \{M_i \mid i \in I\}$. Since T is a product of atoms A_α , for each $i \in I$, set $T_i = \prod [A_\alpha \mid A_\alpha \in \mathcal{T}_{M_i}]$. Then, $T_i \in \mathcal{T}_{M_i}$ and $T = \prod_{M_i \in \text{MSupp}(T/R)} T_i$.

(5) Since $R \subset A$ is minimal for any $A \in \mathcal{A}$, either $R \subset A$ is integral or $R \subset A$ is integrally closed. Moreover, by [31, Lemma 1.5], for a given $M \in \text{MSupp}(S/R)$, minimal extensions $R \subset A$, for $A \in \mathcal{T}_M$, are either all integral, or all integrally closed. Reorder $\text{MSupp}(S/R)$ such that for some $k \in \mathbb{N}_n$, $R \subset A$ is integrally closed for all $A \in \mathcal{T}_{M_i}$ and for any $i \leq k$ and $R \subset A$ is integral for all $A \in \mathcal{T}_{M_i}$ and for any $i > k$. Then, $R \subset V_k$ is Prüfer and $V_k \subset S$ is integral, so that $R \subset S$ is almost-Prüfer. \square

Proposition 2.18. *Let $R \subset S$ be an FCP extension, such that (\mathcal{T}, \subseteq) is a tree. Then,*

- (1) *The elements of \mathcal{T} are Π -irreducible in $[R, S]$.*
- (2) *For each $M \in \text{MSupp}(S/R)$, \mathcal{T}_M is a chain, whose least element $i(M)$ is the only $T \in [R, S]$ satisfying $R \subset T$ is minimal and $M = \mathcal{C}(R, T)$.*
- (3) *Let $M \in \text{MSupp}(S/R)$. For each $T \in [R, S]$ such that $M \in \text{MSupp}(T/R)$, we have $i(M) \subseteq T$.*

Proof. (1) Let $T \in \mathcal{T}$ and $M \in \text{MSupp}(S/R)$ such that $\text{MSupp}(T/R) = \{M\}$. It follows that $R_M \neq T_M$ and $R_{M'} = T_{M'}$ for each $M' \in \text{Max}(R) \setminus \{M\}$. Let $T_1, T_2 \in [R, S]$ be such that $T = T_1 T_2$. Then we have $R \subseteq T_i \subseteq T$ for $i \in \mathbb{N}_2$, giving $(T_i)_{M'} = R_{M'} = T_{M'}$ for each $M' \in \text{Max}(R) \setminus \{M\}$, and $R_M \subseteq (T_i)_M \subseteq T_M$ for $i \in \mathbb{N}_2$, giving, for each $i \in \mathbb{N}_2$, either $R = T_i$ (a), or $\text{MSupp}(T_i/R) = \{M\}$ (b). Fix i and let $j \in \mathbb{N}_2 \setminus \{i\}$. Case (a) gives $T = T_j$. Case (b) gives that $T_i \in \mathcal{T}$. In this case, either $T_j = R$, giving $T = T_i$, or $T_j \in \mathcal{T}$. Hence T_i and T_j are comparable, because \mathcal{T} is a tree. Therefore, T is equal to the greatest element of $\{T_1, T_2\}$ and T is Π -irreducible.

(2) Let $M \in \text{MSupp}(S/R)$ and $T_1, T_2 \in \mathcal{T}_M$. Set $U := T_1 T_2$, so that $T_i \subseteq U$ for $i \in \mathbb{N}_2$ and then $(T_i)_{M'} = R_{M'} = U_{M'}$ for each $M' \in \text{Max}(R) \setminus \{M\}$, and $R_M \subset (T_i)_M \subseteq U_M$. Then, $\text{MSupp}(U/R) = \{M\}$. Therefore, $U \in \mathcal{T}$, so that T_1 and T_2 are comparable and \mathcal{T}_M is a chain.

By Lemma 2.15, there is $T \in [R, S]$ such that $R \subset T$ is minimal with $M = \mathcal{C}(R, T)$. Obviously, we have $\text{MSupp}(T/R) = \{M\}$, so that $T \in \mathcal{T}_M$. Since \mathcal{T}_M is a chain, T is the least element of \mathcal{T}_M , because any $U \in \mathcal{T}_M$ is comparable to T , and we cannot have $U \subset T$. Moreover, since any $T' \in [R, S]$ such that $R \subset T'$ is minimal with $M = \mathcal{C}(R, T')$ satisfies $T' \in \mathcal{T}_M$, we have just proved that there is only one $T' \in [R, S]$ such that $R \subset T'$ is minimal with $M = \mathcal{C}(R, T')$. We set $i(M) := T'$.

(3) Let $T \in [R, S]$ with $M \in \text{MSupp}(T/R)$. For each $M' \in \text{Max}(R) \setminus \{M\}$, we have $R_{M'} = i(M)_{M'} \subseteq T_{M'}$ and a maximal chain $R = R_0 \subset \cdots \subset R_i \subset \cdots \subset R_n = T$, where $R_{i-1} \subset R_i$ is minimal for each $i \in \mathbb{N}_n$. Since $M \in \text{MSupp}(T/R)$, there is some $i \in \mathbb{N}_n$ such that $M = \mathcal{C}(R_{i-1}, R_i) \cap R$, by Lemma 2.14. Let $k \in \mathbb{N}_n$ be the least i satisfying this property. Assume $k \neq 1$, so that $M \notin \text{MSupp}(R_{k-1}/R)$. Again by Lemma 2.15, there is $R'_1 \in [R, R_k]$ such that $R \subset R'_1$ is minimal with $M = \mathcal{C}(R, R'_1)$. From (2), we deduce that $R'_1 = i(M)$, so that $i(M) \subseteq T$. If $k = 1$, then $R_{k-1} = R$ and $R_k = i(M) \subseteq T$. \square

We will see in Proposition 2.21, that, under some conditions, for an FCP extension $R \subset S$, any element of $[R, S]$ can be written in a unique way, as a product of elements of \mathcal{T} . But Remark 2.24 shows that this property does not always hold.

Now, we look at some properties of arithmetic FCP extensions.

Theorem 2.19. *Let $R \subset S$ be an arithmetic FCP extension. Then,*

- (1) $R \subset S$ has FIP and $|[R, S]| \leq \prod_{M \in \text{MSupp}(S/R)} (1 + \ell[R_M, S_M])$.
- (2) \mathcal{T}_M is a chain for each $M \in \text{MSupp}(S/R)$ and (\mathcal{T}, \subseteq) is treed.

Proof. Clearly, $R_M \subset S_M$ has FCP for each $M \in \text{MSupp}(S/R)$ ([11, Proposition 3.7(a)]).

(1) Therefore, $|[R_M, S_M]| < \infty$ follows, since $[R_M, S_M]$ is a chain and $R_M \subset S_M$ has FIP for each $M \in \text{MSupp}(S/R)$. Hence, $R \subset S$ has FIP [11, Proposition 3.7(b)]. By [11, Theorem 3.6 (a)], we have $|[R, S]| \leq \prod_{M \in \text{MSupp}(S/R)} |[R_M, S_M]|$. But $|[R_M, S_M]| = 1 + \ell[R_M, S_M]$ holds since $[R_M, S_M]$ is a chain, giving the requested inequality.

(2) Let $T_1, T_2 \in \mathcal{T}_M$, so that $\{M\} = \text{MSupp}(T_i/R)$

for $i \in \mathbb{N}_2$. Then, $(T_1)_{M'} = (T_2)_{M'} = R_{M'}$ for each $M' \in \text{MSupp}(S/R) \setminus \{M\}$. Moreover, $(T_i)_M \in [R_M, S_M]$ for $i \in \mathbb{N}_2$, so that $(T_1)_M$ and $(T_2)_M$ are comparable, and so are T_1 and T_2 . Then, \mathcal{T}_M is a chain.

Let $T_1, T_2, T \in \mathcal{T}$ be such that $T_i \subseteq T$ for $i \in \mathbb{N}_2$ and $M \in \text{MSupp}(S/R)$ such that $\text{MSupp}(T/R) = \{M\}$. We get, for $i \in \mathbb{N}_2$, that $\emptyset \neq \text{MSupp}(T_i/R) \subseteq \text{MSupp}(T/R) = \{M\}$, so that $\text{MSupp}(T_i/R) = \{M\}$. It follows that $T_1, T_2 \in \mathcal{T}_M$, which is a chain, and then T_1 and T_2 are comparable. Therefore, \mathcal{T} is a tree. \square

Let $R \subset S$ be an FCP extension and $\text{MSupp}(S/R) := \{M_1, \dots, M_n\}$. We will use the maps $\varphi : [R, S] \rightarrow \prod_{i=1}^n [R_{M_i}, S_{M_i}]$ defined by $\varphi(T) := (T_{M_1}, \dots, T_{M_n})$ and $\varphi_M : [R, S] \rightarrow [R_M, S_M]$ defined by $\varphi_M(T) := T_M$, for each $M \in \text{MSupp}(S/R)$. Then φ is injective [11, Theorem 3.6]. If φ is bijective, $R \subseteq S$ is called a \mathcal{B} -extension (\mathcal{B} stands for bijective).

Proposition 2.20. *An FCP extension $R \subseteq S$ is a \mathcal{B} -extension if and only if R/P is local for each $P \in \text{Supp}(S/R)$.*

The above “local” condition on the factor domains R/P holds in case $\text{Supp}(S/R) \subseteq \text{Max}(R)$, and, in particular, if $R \subset S$ is integral.

Proof. One implication appears in the proof of [11, Theorem 3.6(b)] which uses [11, Lemma 3.5].

Conversely, assume that φ is bijective and that there is some $P \in \text{Supp}(S/R)$ contained in two elements $M_1, M_2 \in \text{Max}(R)$. Consider a maximal chain $\mathcal{C} : R = R_0 \subset \dots \subset R_i \subset \dots \subset R_n = S$, where $R_{i-1} \subset R_i$ is minimal for each $i \in \mathbb{N}_n$. Since $P \in \text{Supp}(S/R)$, there exists some $i \in \mathbb{N}_n$ such that $P = N \cap R$, where $N := \mathcal{C}(R_{i-1}, R_i)$ by Lemma 2.14. In particular, $P \in \text{Supp}(R_i/R)$, which implies that $M_j \in \text{Supp}(R_i/R)$ for $j \in \mathbb{N}_2$. Since φ is surjective, there is $T \in [R, S]$ such that $T_{M_1} = (R_i)_{M_1}$ and $T_M = R_M$ for each $M \in \text{MSupp}(S/R) \setminus \{M_1\}$. In particular, $M_2 \notin \text{Supp}(T/R)$. Localizing \mathcal{C} at M_1 , we get that $P_{M_1} \in \text{Supp}((R_i)_{M_1}/R_{M_1}) = \text{Supp}(T_{M_1}/R_{M_1})$ so that $P \in \text{Supp}(T/R)$, Now $M_2 \in \text{Supp}(T/R)$ is absurd. Then, R/P is a local ring for each $P \in \text{Supp}(S/R)$.

If $\text{Supp}(S/R) \subseteq \text{Max}(R)$, it follows from Lemma 2.14 that each $\mathcal{C}(R_{i-1}, R_i)$ lies over a maximal ideal of R (and so R/P is a field, hence local for each $P \in \text{Supp}(S/R)$). Finally, it is standard that maximal ideals lie over maximal ideals in any integral extension (cf. [22, Theorem 44]). \square

Under an additional assumption, next proposition gives a converse to Theorem 2.19. Set $\text{MSupp}(S/R) := \{M_1, \dots, M_n\}$ for an FCP extension $R \subset S$. It follows that the elements of $\text{MSupp}(T/R)$ are some M_i when $T \in]R, S]$.

Proposition 2.21. *Let $R \subset S$ be an FCP \mathcal{B} -extension.*

- (1) *For each $M \in \text{MSupp}(S/R)$, we have $[R_M, S_M] = \varphi_M(\mathcal{T}) \cup \{R_M\} = \varphi_M(\mathcal{T}_M) \cup \{R_M\}$.*

- (2) Any $T \in]R, S]$ is a product of $|\text{MSupp}(T/R)|$ distinct elements $E_i \in \mathcal{T}$ in a unique way such that $E_i \in \mathcal{T}_{M_i}$, for each $M_i \in \text{MSupp}(T/R)$.
- (3) $R \subset S$ is arithmetic if and only if \mathcal{T} is a tree.
- (4) Assume that $R \subset S$ is arithmetic. Then, \mathcal{T} is the set of Π -irreducible elements of $]R, S]$.

$$\text{Moreover, } |[R, S]| = \prod_{M \in \text{MSupp}(S/R)} (1 + \ell[R_M, S_M]).$$

Proof. (1) The following inclusions are obvious: $\varphi_M(\mathcal{T}_M) \cup \{R_M\} \subseteq \varphi_M(\mathcal{T}) \cup \{R_M\} \subseteq [R_M, S_M]$. Let $E \in]R_M, S_M]$. Since φ is a bijection, there exists $T \in]R, S]$ such that $T_M = E$ and $T_{M'} = R_{M'}$ for each $M' \in \text{MSupp}(S/R) \setminus \{M\}$. The result follows from $E = \varphi_M(T)$ and $T \in \mathcal{T}_M$.

(2) Let $T \in]R, S]$. From $\text{MSupp}(T/R) \neq \emptyset$, we infer that some $M_i \in \text{MSupp}(T/R)$. Since φ is a bijection, there exists a unique $E_i \in [R, S]$ such that $(E_i)_{M_i} = T_{M_i}$ and $(E_i)_M = R_M$ for $M \neq M_i$. It follows that $E_i \in \mathcal{T}_{M_i}$ for each $M_i \in \text{MSupp}(T/R)$. Set $E_j = R$ when $M_j \notin \text{MSupp}(T/R)$ and $E := E_1 \cdots E_n$. For each $j \in \mathbb{N}_n$, we get that $E_{M_j} = T_{M_j}$, so that $E = T$ is the product of $|\text{MSupp}(T/R)|$ distinct elements of \mathcal{T} . The uniqueness of these elements is obvious.

(3) One implication is Theorem 2.19. Assume that \mathcal{T} is a tree. Let $M \in \text{MSupp}(S/R)$. By Proposition 2.18, \mathcal{T}_M is a chain. Since φ_M preserves order, (1) implies that $[R_M, S_M]$ is a chain.

(4) Assume that $R \subset S$ is arithmetic. Then, \mathcal{T} is a tree. It results from Proposition 2.18 that the elements of \mathcal{T} are Π -irreducible. Conversely, let $T \in]R, S]$ be Π -irreducible. In view of (2), T is a product of elements $E_i \in \mathcal{T}_{M_i}$, where $M_i \in \text{MSupp}(T/R)$, and then, of only one element of \mathcal{T} , so that $T \in \mathcal{T}$.

Now $\prod_{M \in \text{MSupp}(S/R)} |[R_M, S_M]| = \prod_{M \in \text{MSupp}(S/R)} (1 + \ell[R_M, S_M]) = |[R, S]|$, since φ is bijective and each $R_M \subset S_M$ a chain. \square

For the definition of the Goldie dimension of a distributive lattice, the reader may look at [24, p. 14 and Exercise 8, p. 33]. We will use the following results.

Proposition 2.22. [24, Theorem 1.5.9] *If $R \subseteq S$ is an FCP arithmetic extension, its Goldie dimension is the integer n such that $R = B_1 \cap \cdots \cap B_n$ is an irredundant representation of R by \cap -irreducible elements.*

Proposition 2.23. *Let $R \subset S$ be an FCP arithmetic \mathcal{B} -extension extension. Then, the Goldie dimension of $[R, S]$ is $n := |\text{MSupp}_R(S/R)|$.*

Proof. For each $M_i \in \text{MSupp}(S/R)$ there exists a unique $E'_i \in [R, S]$ such that $(E'_i)_{M_i} = R_{M_i}$ and $(E'_i)_M = S_M$ for $M \neq M_i$. Set $E' :=$

$E'_1 \cap \dots \cap E'_n$. For each i , we get that $(E'_i)_{M_i} = R_{M_i}$, so that $E' = R$ is the intersection of $|\text{MSupp}(S/T)|$ distinct elements of $[R, S]$. The uniqueness of these elements is obvious.

We claim that the E'_i are \cap -irreducible. Let $T, T' \in [R, S]$ be such that $E'_i = T \cap T'$. Then we have $R \subset E'_i \subseteq T, T'$, giving $(E'_i)_M = S_M = T_M = T'_M$ (*) for each $M \in \text{Max}(R) \setminus \{M_i\}$, and $R_{M_i} = (E'_i)_{M_i} \subseteq T_{M_i}, T'_{M_i}$. Since $R \subset S$ is arithmetic, T_{M_i} and T'_{M_i} are comparable, and so are T and T' by (*). It follows that E'_i is the least element of $\{T, T'\}$, and then is \cap -irreducible so that n is the Goldie dimension of $[R, S]$ by Proposition 2.22. \square

Remark 2.24. Using [11, Remark 3.4(b)], we exhibit a non \mathcal{B} -extension, for which some statements of Proposition 2.21 do not hold.

We now summarize the context of the above quoted remark.

Let R be a two-dimensional Prüfer domain with exactly two height-2 maximal ideals, N and N' , each of which containing the unique height 1 prime ideal P of R . Set $R_0 := R$, $R_1 := R_N$, $R'_1 := R_{N'}$, $R_2 := R_P$ and $R_3 := K$, the quotient field of R . Since each overring of a Prüfer domain is an intersection of localizations [19, Theorem 26.2], it is easy to check that $R_0 \subset R_1$, $R_0 \subset R'_1$, $R_1 \subset R_2$, $R'_1 \subset R_2$ and $R_2 \subset R_3$ are Prüfer minimal extensions.

Moreover, $\mathcal{C}(R_0, R_1) = N'$, $\mathcal{C}(R_1, R_2) = NR_N$ and $\mathcal{C}(R_2, R_3) = P$, which is an ideal of R_2 because $P = (R_0 : R_2)$. It follows that $\text{Supp}(K/R) = \{P, N, N'\}$ and $\text{MSupp}(K/R) = \{N, N'\}$. By Proposition 2.20, the map φ is not bijective, since R/P is not local. The poset $\mathcal{T} = \{R_1, R'_1\}$ is a tree. Moreover, $[R_N, K_N] = \{R_N, R_P, K\} = \{R_1, R_2, R_3\}$ and $[R_{N'}, K_{N'}] = \{R_{N'}, R_P, K\} = \{R'_1, R_2, R_3\}$ are chains. However, $\varphi_N(\mathcal{T}) = \{R_1, R_2\}$, because $(R_{N'})_N = R_P$. Then, Proposition 2.21(1) is not satisfied. In the same way, Proposition 2.21(2) is not satisfied, because K is not a product of elements of \mathcal{T} . In particular, some Π -irreducible element as $R_3 = K$ of $[R, S]$ is not in \mathcal{T} .

The conditions of Proposition 2.21 hold in the following context and provide us a structure of Boolean lattices in Proposition 3.14.

Proposition 2.25. *Let $R \subset S$ be an FCP extension. Assume that $\text{Supp}(T/R) \cap \text{Supp}(S/T) = \emptyset$ for all $T \in [R, S]$. Then:*

- (1) $\text{Supp}(S/R) \subseteq \text{Max}(R)$ and $R \subseteq S$ is a \mathcal{B} -extension.
- (2) \mathcal{T} is an antichain.

Proof. (1) Assume that $\text{Supp}(T/R) \cap \text{Supp}(S/T) = \emptyset$ for all $T \in [R, S]$. We use Proposition 2.20 to show that φ is a bijection. Let $P \in \text{Supp}(S/R)$. Consider a maximal chain $\mathcal{C} : R = R_0 \subset \dots \subset R_i \subset \dots \subset R_n = S$, where $R_{i-1} \subset R_i$ is minimal for each $i \in \mathbb{N}_n$. Since $P \in$

$\text{Supp}(S/R)$, there exists some $k \in \mathbb{N}_n$ such that $P = N \cap R$, where $N := \mathcal{C}(R_{k-1}, R_k)$, in view of Lemma 2.14. In particular, $P \in \text{Supp}(R_k/R)$, and, more precisely, $P \in \text{Supp}(R_k/R_{k-1}) \subseteq \text{Supp}(S/R_{k-1})$. Assume that P is not a maximal ideal, and let $M \in \text{Max}(R)$ be such that $P \subset M$. Then, $M \in \text{Supp}(R_k/R)$. Moreover, there is some $j < k$ such that $M = N' \cap R$, where $N' := \mathcal{C}(R_{j-1}, R_j)$. Indeed, $j \neq k$ because $N \cap R = P \neq M$. Therefore $M \in \text{Supp}(R_j/R) \subseteq \text{Supp}(R_{k-1}/R)$, since $j \leq k-1$ and $M \in \text{Supp}(R_{k-1}/R) \cap \text{Supp}(S/R_{k-1})$ is absurd. Hence, $\text{Supp}(S/R) \subseteq \text{Max}(R)$, and φ is a bijection by Proposition 2.20.

(2) Let $T, T' \in \mathcal{T}$ be such that $T' \subset T$ and $M \in \text{Max}(R)$ such that $\text{Supp}(T/R) = \{M\}$. Since $T' \subset T$, we get that $\text{Supp}(T'/R) \subseteq \text{Supp}(T/R)$, giving $\text{Supp}(T'/R) = \{M\}$. But, $\emptyset \neq \text{Supp}(T/T') \subseteq \text{Supp}(T/R) = \{M\}$ implies that $\{M\} = \text{Supp}(T/T') \subseteq \text{Supp}(S/T')$. Therefore, $M \in \text{Supp}(T'/R) \cap \text{Supp}(S/T') = \emptyset$, a contradiction. Then, two distinct elements of \mathcal{T} are incomparable, and \mathcal{T} is an antichain. \square

Let $R \subset S$ be an FCP extension. In the following, we will meet the condition that $\text{Supp}(T/R) \cap \text{Supp}(S/T) = \emptyset$ for some $T \in [R, S]$. Here is a theorem which gives a stronger result.

Theorem 2.26. *If $R \subset S$ has FCP, then $\text{Supp}(T/R) \cap \text{Supp}(S/T) = \emptyset$ for all $T \in [R, S]$ if and only if $R \subset S$ is locally minimal. In this case, $R \subset S$ is an FIP factorial extension.*

Proof. Assume that $\text{Supp}(T/R) \cap \text{Supp}(S/T) = \emptyset$ for all $T \in [R, S]$. By Propositions 2.25 and 2.21, $R \subset S$ is arithmetic. Let $T \in [R, S]$, there is $U \in [R, S]$ such that $U \cap T = R$ and $UT = S$ [28, Lemma 3.7], which gives $U_M \cap T_M = R_M$ and $U_M T_M = S_M$ for each $M \in \text{MSupp}(S/R)$, so that $\{U_M, T_M\} = \{R_M, S_M\}$, whence, $R_M \subset S_M$ is minimal.

Conversely, if there exists some $N \in \text{MSupp}(T/R) \cap \text{MSupp}(S/T)$ for some $T \in [R, S]$, we get that $T_N \neq R_N, S_N$, so that $|[R_N, S_N]| \geq 2$. But $R_N \subset S_N$ is minimal, a contradiction.

If these conditions hold, then $R \subseteq S$ has FIP by Theorem 2.19, and is factorial by Propositions 2.21 and 2.16. \square

3. BOOLEAN FCP EXTENSIONS

3.1. General properties of Boolean extensions. Let $R \subseteq S$ be an extension and $T \in [R, S]$. Then, $T' \in [R, S]$ is called a *complement* of T if $T \cap T' = R$ and $TT' = S$. If $R \subseteq S$ is distributive, then T has at most one complement [24, Exercise 9, page 33]. We denote this complement by T° when it exists.

We recall that $R \subseteq S$ is Boolean if and only if $R \subseteq S$ is distributive and each $T \in [R, S]$ has a (unique) complement T° [32, Definition

page 129]. In a Boolean FIP extension $R \subseteq S$, any $T \in]R, S[$ is, in a unique way, a product of finitely many atoms and the intersection of finitely many co-atoms. To see this, use Lemma 2.10, [32, Theorems 5.1 and 6.3], and the fact that $[R, S]$ is a complete Boolean lattice. In particular, if A is an atom, then A° is a co-atom [32, Theorem 3.43]. Next Theorem characterizes Boolean FCP extensions amid distributive FCP extensions:

Theorem 3.1. *Let $R \subseteq S$ be an FCP distributive extension and set $n := \ell[R, S]$. Then, the following conditions are equivalent:*

- (1) $R \subseteq S$ is a Boolean extension.
- (2) $n = |\mathcal{A}|$, where \mathcal{A} is the set of atoms of $[R, S]$.
- (3) $|[R, S]| = 2^n$.

If these conditions hold, then $R \subseteq S$ is FIP factorial and a simple pair. Moreover, for each $T, U \in [R, S]$ with $T \subseteq U$, all maximal chains of $[T, U]$ have the same length.

Proof. To begin with, $R \subseteq S$ has FIP by Lemma 2.10. Moreover, $[R, S]$ has exactly n Π -irreducible element by Lemma 2.10. Since any atom is Π -irreducible, the equivalences of (1), (2) and (3) are a translation of the equivalences given in [33, page 292].

If these conditions hold, then $R \subseteq S$ is factorial and a simple pair by Theorem 2.11. Since a Boolean lattice is a distributive lattice, the last results comes from Proposition 2.5. \square

Example 3.2. (1) An extension $R \subset S$ is Boolean and chained if and only if it is minimal. One implication is obvious. Conversely, assume that $[R, S]$ is chained and a Boolean lattice. If there is some $T \in]R, S[$, it has a complement T' . Then, $T \cap T' = R$ and $TT' = S$ implies $\{T, T'\} = \{R, S\}$ since T and T' are comparable, a contradiction.

(2) Let $R \subset S$ be a Boolean FCP extension and $x \in S \setminus R$. We intend to compute $R[x]^\circ$. Let $\mathcal{C}\mathcal{A} := \{B_1, \dots, B_n\}$. Then $R[x] = \cap[B_\alpha \in Y]$ for a unique family $Y := \{B_\alpha\}$ of co-atoms ([32, Theorem 6.3]), so that $x \in B_\alpha$ for each $B_\alpha \in Y$. Let $B_\beta \in \mathcal{C}\mathcal{A}$ which does not contain x . Then $R[x] \not\subseteq B_\beta$ and Y is the set of co-atoms containing x . Now, $(R[x])^\circ = (\cap[B_\alpha \in Y])^\circ = \Pi[(B_\alpha)^\circ \mid B_\alpha \in Y]$, where the $(B_\alpha)^\circ$ are atoms. Then, $(R[x])^\circ$ is the product of atoms which are complements of the co-atoms containing x . But we also have $(R[x])^\circ = \cap[B_\beta \in \mathcal{C}\mathcal{A} \setminus Y]$. Indeed, set $T := \cap[B_\beta \in \mathcal{C}\mathcal{A} \setminus Y]$. Obviously, $R[x] \cap T = R$ and assume that $R[x]T \neq S$, so that $R[x]T = \cap[B_\gamma \in X]$ for some $X \subseteq \mathcal{C}\mathcal{A}$. Let $B_\gamma \in X$. Then, $R[x] = \cap[B_\alpha \in Y] \subseteq B_\gamma$ implies $B_\alpha = B_\gamma$ for some $B_\alpha \in Y$. In the same way, $B_\beta = B_\gamma$ for some $B_\beta \in \mathcal{C}\mathcal{A} \setminus Y$, a contradiction. Then, $R[x]T = S$ and $T = (R[x])^\circ$.

(3) By Theorem 3.1, an FCP Boolean extension $R \subseteq S$ verifies $|\llbracket R, S \rrbracket| = 2^n$, where $n = \ell[R, S]$. But an extension $R \subseteq S$ may be distributive with $|\llbracket R, S \rrbracket| = 2^n$ for some integer n without being Boolean. It is enough to consider a chained extension of length $2^n - 1$.

The following lemma is needed for the next proposition. See the close notion of *patching* due to Dobbs-Shapiro [15].

Lemma 3.3. *Let $R \subseteq S$ be an FCP extension and $M \in \text{MSupp}(S/R)$. For any $T' \in [R_M, S_M]$ such that $R_M \subset T'$ is minimal, there exists $T \in [R, S]$ such that $R \subset T$ is minimal with $T_M = T'$.*

Proof. Let $\varphi : S \rightarrow S_M$ be the canonical ring morphism and set $T'' := \varphi^{-1}(T')$. Then $T'' \in [R, S]$ is such that $T' = T''_M \neq R_M$, so that $M \in \text{MSupp}(T''/R)$. From Lemma 2.15 we deduce the existence of some $T \in [R, T''] \subseteq [R, S]$ such that $R \subset T$ is minimal with $M = \mathcal{C}(R, T)$. Hence, $R_M \subset T_M \subseteq T''_M = T'$ gives $T_M = T'$. \square

Proposition 3.4. *Let $R \subseteq S$ be an FCP extension. The following statements are equivalent:*

- (1) $R \subseteq S$ is Boolean.
- (2) $R_M \subseteq S_M$ is Boolean for each $M \in \text{MSupp}(S/R)$.
- (3) $R_P \subseteq S_P$ is Boolean for each $P \in \text{Supp}(S/R)$.
- (4) $R/I \subseteq S/I$ is Boolean for each ideal I shared by R and S .
- (5) $R/I \subseteq S/I$ is Boolean for some ideal I shared by R and S .

Proof. We have obviously (1) \Rightarrow (3) \Rightarrow (2) and (1) \Leftrightarrow (4) \Rightarrow (5) \Rightarrow (1). Conversely, assume that $R_M \subseteq S_M$ is Boolean for each $M \in \text{MSupp}(S/R)$. Then, $R_M \subseteq S_M$ is Boolean for each $M \in \text{Max}(R)$. It follows that the distributivity property holds in $[R, S]$ since it holds in any $[R_M, S_M]$. It remains to show that any $T \in [R, S]$ has a complement. Let $M \in \text{MSupp}(S/R)$. Then, T_M has a complement $(T_M)^\circ$ in $[R_M, S_M]$ satisfying $T_M \cap (T_M)^\circ = R_M$ (*) and $T_M(T_M)^\circ = S_M$ (**). Since $[R_M, S_M]$ is Boolean, any of its element is a product of its atoms [32, Theorem 5.2]. Then, $(T_M)^\circ = \prod_{i \in I_M} R'_{i,M}$, where the $R'_{i,M}$ are atoms of $[R_M, S_M]$, that is $R_M \subset R'_{i,M}$ is minimal. By Lemma 3.3, for each $R'_{i,M} \in [R_M, S_M]$, there is $R_{i,M} \in [R, S]$ such that $R \subset R_{i,M}$ is minimal, with $R'_{i,M} = (R_{i,M})_M$. In particular, $(R_{i,M})_{M'} = R_{M'}$ for each $M' \in \text{MSupp}(S/R) \setminus \{M\}$. Setting $T' := \prod_{M \in \text{MSupp}(S/R)} (\prod_{i \in I_M} R_{i,M})$, we get, for each $M \in \text{MSupp}(S/R)$ that $T'_M = \prod_{i \in I_M} R'_{i,M} = (T_M)^\circ$, so that (*) and (**) give $T_M \cap T'_M = R_M$ and $T_M T'_M = S_M$, and, to end, $T \cap T' = R$ and $TT' = S$ showing that T' is the complement of T . Therefore, $R \subseteq S$ is Boolean. \square

Corollary 3.5. *Let $R \subset S$ be an FIP extension. The following statements are equivalent:*

- (1) $R \subset S$ is atomistic and arithmetic;
- (2) $R \subset S$ is Boolean and arithmetic;
- (3) $R \subset S$ is locally minimal.

Assume that these conditions hold and let the set of atoms be $\mathcal{A} = \{A_1, \dots, A_a\}$ where a is some integer. Then, the complement of any $T = \prod_{i \in I} A_i \in [R, S]$, where $I \subseteq \mathbb{N}_a$, is $T^\circ := \prod_{j \in J} A_j$ where $J := \mathbb{N}_a \setminus I$. If in addition $R \subset S$ is integral, then $\ell[R, S] = |\text{MSupp}(S/R)|$.

Proof. (1) \Rightarrow (2) by Proposition 2.3 and the equivalences given in [33, page 292], (2) \Rightarrow (1) by [32, Theorem 5.2] and (2) \Leftrightarrow (3) by Proposition 3.4 and Example 3.2(1).

Assume that these conditions hold. We observe that S is the product of all atoms. Let $T = \prod_{i \in I} A_i \in [R, S]$, where $I \subseteq \mathbb{N}_a$. Since T° is a product of atoms, the relations $T \cap T^\circ = R$ and $TT^\circ = S$ give $T^\circ := \prod_{j \in J} A_j$ where $J := \mathbb{N}_a \setminus I$.

In case $R \subseteq S$ is integral, we use $\ell[R, S] = \sum_{M \in \text{MSupp}(S/R)} \ell[R_M, S_M]$ [12, Proposition 4.6]. \square

Next proposition uses the notation of [4, Proposition 10, p.52].

Proposition 3.6. *Let $R \subset S$ be a ring extension, $f : R \rightarrow R'$ a faithfully flat ring morphism and $S' := R' \otimes_R S$. Assume that $R' \subset S'$ is distributive. Then,*

- (1) $R \subset S$ is distributive.
- (2) Let $T \in [R, S]$ be such that $R'T$ is Π -irreducible (resp. an atom) in $[R', S']$. Then, T is Π -irreducible (resp. an atom) in $[R, S]$.
- (3) In case $R'T$ is Π -irreducible for any Π -irreducible $T \in [R, S]$ and $R' \subset S'$ is an FIP Boolean extension, so is $R \subset S$.

Proof. (1) The ring morphism $\varphi : S \rightarrow S'$ defines a map $\theta : [R', S'] \rightarrow [R, S]$ while there is a map $\psi : [R, S] \rightarrow [R', S']$, defined by $\psi(T) = R' \otimes_R T$ and such that $\theta \circ \psi$ is the identity of $[R, S]$ by [4, Proposition 10, p.52] (it is enough to take $F = S$ and to observe that if M is an R -submodule of S , then with the notation of the above reference, $R'M$ identifies to $M \otimes_R R'$). In particular, ψ is injective. The same reference shows that $\psi(T \cap U) = \psi(T) \cap \psi(U)$ for $U, T \in [R, S]$. It is easy to show that $\psi(TU) = \psi(T)\psi(U)$. If $R' \subset S'$ is distributive, it follows that for $T, U, V \in [R, S]$, we get $\psi[T(U \cap V)] = \psi(T)[\psi(U) \cap \psi(V)] = [\psi(T)\psi(U)] \cap [\psi(T)\psi(V)] = \psi[(TU) \cap (TV)]$, giving $T(U \cap V) = (TU) \cap (TV)$. Then, $R \subset S$ is distributive.

(2) Let $T \in [R, S]$ be such that $R'T$ is Π -irreducible in $[R', S']$ and let $U, V \in [R, S]$ be such that $T = UV$. Then, $R'T = (R'U)(R'V)$, so that either $R'T = R'U$, or $R'T = R'V$, which implies either $T = U$, or $T = V$ and T is Π -irreducible in $[R, S]$.

Let $T \in [R, S]$ be such that $R'T$ is an atom in $[R', S']$. Assume that T is not an atom in $[R, S]$. There exists $U \in]R, T[$ which yields $R' \subset R'U \subset R'T$, a contradiction. Then, T is an atom in $[R, S]$.

(3) Assume that $R'T$ is Π -irreducible in $[R', S']$ for any Π -irreducible $T \in [R, S]$. This holds if θ (or ψ) is bijective. If $R' \subset S'$ is an FIP Boolean extension, any Π -irreducible element of $[R', S']$ is an atom in $[R', S']$ in view of [33, page 292]. Moreover, $R \subset S$ is also an FIP extension since θ is surjective. Let $T \in [R, S]$ be Π -irreducible in $[R, S]$. Then, $R'T$ is Π -irreducible in $[R', S']$. Then $R'T$ is an atom in $[R', S']$, so that T is an atom in $[R, S]$ by (2). Since Π -irreducible elements of $[R, S]$ are atoms, the same reference shows that $R \subset S$ is Boolean. \square

Proposition 3.7. *An FCP extension $R \subset S$, whose Nagata extension $R(X) \subset S(X)$ has FIP and is Boolean, has FIP and is Boolean.*

Proof. In view of ([12, Corollary 3.5]), $R \subset S$ is an FCP extension implies that $S(X) = R(X) \otimes_R S$, so that we can use Proposition 3.6 because $R(X) \subset S(X)$ is distributive. Since $R(X) \subset S(X)$ has FIP, the map $\psi : [R, S] \rightarrow [R(X), S(X)]$ defined by $\psi(T) = T(X)$ in Proposition 3.6 is an order-isomorphism by [13, Theorem 32]. Let $T \in [R, S]$ be Π -irreducible. Then, $\psi(T) = T(X) = TR(X)$ is Π -irreducible, so that $R \subset S$ is Boolean by Proposition 3.6. \square

Proposition 3.8. *Let $R \subset S$ be a ring extension, $f : R \rightarrow R'$ a flat ring epimorphism and $S' := R' \otimes_R S$. If $R \subset S$ is a distributive extension (resp. a FIP Boolean extension), then so is $R' \subset S'$.*

Proof. The proof is a consequence of the following facts. Let $f : R \rightarrow R'$ be a flat epimorphism and $Q \in \text{Spec}(R')$, lying over P in R , then $R_P \rightarrow R'_Q$ is an isomorphism. Moreover, we have $(R' \otimes_R S)_Q \cong R'_Q \otimes_{R_P} S_P$, so that $R_P \rightarrow S_P$ identifies to $R'_Q \rightarrow (R' \otimes_R S)_Q = S'_Q$.

Assume that $R \subset S$ is distributive (resp.: FIP Boolean). Then, so is $R_P \rightarrow S_P$ for each $P \in \text{Spec}(R)$ by Proposition 2.4 (resp. Proposition 3.4 and [11, Proposition 3.7]). Let $Q \in \text{Spec}(R')$ and $P := f^{-1}(Q) \in \text{Spec}(R)$. Since $R_P \rightarrow S_P$ identifies to $R'_Q \rightarrow S'_Q$, we get that $R'_Q \subset S'_Q$ is distributive (resp.; FIP Boolean) for each $Q \in \text{Spec}(R')$. It follows that $R' \subset S'$ is distributive (resp.: FIP Boolean) by the same references. Indeed, in the FIP Boolean case, since $R \subset S$ has FIP, so has $R' \subset S'$, because $\text{Spec}(R') \rightarrow \text{Spec}(R)$ is injective. \square

Proposition 3.9. *Let $R \subset S$ be an FCP distributive extension. The following statements are equivalent:*

- (1) $R \subset S$ is Boolean;
- (2) Any Π -irreducible element is an atom;
- (3) Any \cap -irreducible element is a co-atom.

Proof. (1) \Rightarrow (2) and (3). Assume that $R \subset S$ is Boolean. By [32, Theorem 5.2], the Π -irreducible elements of $[R, S]$ are the atoms of $[R, S]$. Using complements, we deduce that the \cap -irreducible elements of $[R, S]$ are the co-atoms of $[R, S]$.

(2) \Rightarrow (1) by the equivalences given in [33, page 292] since $R \subset S$ has FIP by Lemma 2.10.

(3) \Rightarrow (2) It is enough to exchange products and intersections, and atoms and co-atoms. \square

Theorem 3.10. *Let $R \subset S$ be an FCP extension. The following statements are equivalent, each of them implying that $R \subset S$ has FIP:*

- (1) $R \subset S$ is Boolean;
- (2) $R \subset S$ is factorial;
- (3) $R \subset S$ is co-factorial.

Proof. (1) \Rightarrow (2) by Theorem 3.1 and (1) \Rightarrow (3) because, by Proposition 2.9, any $T \in [R, S]$ is a finite intersection of \cap -irreducible elements, and then a finite intersection of co-atoms of $[R, S]$ by Proposition 3.9.

(2) \Rightarrow (1). By Theorem 2.11, $R \subset S$ is an atomistic distributive FIP extension. Then, use the equivalences given in [33, page 292].

(3) \Rightarrow (2) It is enough to exchange products and intersections, and atoms and co-atoms. \square

Proposition 3.11. [17, Theorems 107 and 158] *Let $R \subset S$ be a Boolean extension. Then, $U \subset T$ is Boolean for any $U, T \in [R, S]$ such that $U \subseteq T$ and the complement of $V \in [U, T]$ in $[U, T]$ is $U(T \cap V^\circ)$.*

We can now generalize Ayache's result [3, Theorem 7] in case of an arbitrary ring extension.

Proposition 3.12. *When $R \subset S$ has a maximal chain of length n from R to S such that $|\text{Supp}(S/R)| = |\text{MSupp}(S/R)| = n$, then $R \subset S$ is FIP Boolean, any maximal chain of $[R, S]$ has length n and $|[R, S]| = 2^n$.*

Proof. Let $R_0 := R \subset \dots \subset R_i \subset \dots \subset R_n := S$ be a maximal chain of R -subalgebras of length n . For each $i \in \mathbb{N}_n$, set $M_{i-1} := \mathcal{C}(R_{i-1}, R_i) \cap R$. Then, $\text{Supp}(S/R) = \{M_i\}_{i=0}^{n-1} \subseteq \text{Max}(R)$ in view of Lemma 2.14. It follows that $M_i \neq M_j$ for each $i \neq j$, so that $R_{M_i} = (R_i)_{M_i} \subset (R_{i+1})_{M_i} = S_{M_i}$ is minimal (and then has FIP), so

that $R \subset S$ has FIP by [11, Proposition 3.7]. Now, $R_{M_i} \subseteq S_{M_i}$ is Boolean (see Example 3.2(1)), and so is $R \subseteq S$ by Proposition 3.4. The last results follow from Theorem 3.1. \square

Let $R \subset S$ be an FCP Boolean extension and let $R_0 := R \subset \dots \subset R_i \subset \dots \subset R_n := S$ be a maximal chain of R -subalgebras of S . By Proposition 3.11, $R \subset R_{n-1}$ is an Boolean FCP extension and $R_{n-1} \subset S$ is minimal. Next theorem gives a kind of converse which allows us to check by induction that an FIP extension is Boolean.

Theorem 3.13. *An FIP extension $R \subset S$, which is not minimal, is Boolean if and only if there exist $U, T \in]R, S[$ such that the conditions (1), (2), (3) and (4) hold, if and only if there exist $U, T \in]R, S[$ such that the conditions (1), (4) and (5) hold:*

- (1) $[R, S] = [R, T] \cup [U, S]$.
- (2) $[U, S] = \{UL \mid L \in [R, T]\}$.
- (3) $L \subset UL$ is a minimal extension for each $L \in [R, T]$.
- (4) $[R, T]$ is a Boolean lattice.
- (5) The map $\varphi : [R, T] \rightarrow [U, S]$ defined by $\varphi(L) = UL$, for $L \in [R, T]$, is bijective.

Moreover, if these conditions hold, U is an atom, $T = U^\circ$ is a co-atom. In fact, these conditions hold for any atom U' and its complement T' , and $[R, T'] \cap [U', S] = \emptyset$.

Proof. Assume that $R \subset S$ has FIP and is not minimal. Set $\mathcal{A} := \{A_1, \dots, A_n\}$. We will prove the Theorem in four steps:

- (a) $R \subset S$ is Boolean \Rightarrow (1)+(2)+(3)+(4).
- (b) (1)+(2)+(3)+(4) \Rightarrow (1)+(4)+(5).
- (c) (1)+(4)+(5) \Rightarrow (1)+(2)+(3)+(4).
- (d) (1)+(2)+(3)+(4)+(5) \Rightarrow $R \subset S$ is Boolean.

(a) Assume that $R \subset S$ is Boolean. Set $U := A_1 \in [R, S]$ and $T := U^\circ \in [R, S]$, the complement of U , so that $U \cap T = R$ and $UT = S$. Moreover, $T = \prod_{i=2}^n A_i$ [32, Theorems 3.43 and 5.1].

(1) Let $L \in [R, S]$ and assume that $L \notin [U, S]$. We have $U \subseteq L^\circ$, the complement of L [32, Theorem 5.1] and $U \subseteq L^\circ$ implies $L \subseteq T$ by [32, Theorem 5.1], so that $L \in [R, T]$. Hence (1) is proved. Moreover, $[R, T] \cap [U, S] = \emptyset$ because $U \subseteq L \subseteq T$, for some $L \in [R, T] \cap [U, S]$ leads to the contradiction $U \cap T = U = R$.

(2) Each element of $[R, S]$ is a product of some A_i s by Theorem 3.10. Let $L' := \prod_{i \in I} A_i \in [U, S]$ for some $I \subseteq \mathbb{N}_n$. Then, $1 \in I$ because $U \subseteq L'$ (if not, $U \cap L' = R = U$ is absurd). In particular, $L' = UL$, where $L := \prod_{i \in I \setminus \{1\}} A_i \subseteq T$ and (2) follows since $UL \in [U, S]$.

(3) Let $L := \prod_{i \in I} A_i \in [R, T]$, so that $1 \notin I$ by (1). Then, $UL = \prod_{i \in I \cup \{1\}} A_i$. Let $L' \in [L, UL]$. There exists some $J \subseteq \mathbb{N}_n$ such that $L' := \prod_{i \in J} A_i$. By the uniqueness of this writing, $I \subseteq J \subseteq I \cup \{1\}$, so that we have either $I = J$ or $J = I \cup \{1\}$, giving either $L' = L$ or $L' = UL$. It follows that $L \subset UL$ is minimal and (3) holds.

(4) By Proposition 3.11 $R \subset T$ is Boolean. Remark that (1), (2), (3) and (4) hold for any atom U' with complement T' .

$$\begin{array}{ccccc} R & \rightarrow & L & \rightarrow & T \\ \downarrow & & \downarrow & & \downarrow \\ U & \rightarrow & UL & \rightarrow & S \end{array}$$

(b) Assume that (1)+(2)+(3)+(4) holds. Then φ is surjective by (2). Let $L, L' \in [R, T]$ be such that $\varphi(L) = \varphi(L')$, that is $UL = UL'$. Since $L, L' \subseteq T$, we get that $LL' \in [R, T]$. Moreover, $U(LL') = (UL)(UL') = UL$. Then, $L, L' \subseteq LL' \subseteq UL = UL'$ gives by (3) either $LL' \neq UL = UL'$, so that $LL' = L = L'$ or $LL' = UL \in [U, S]$. But, in this last case, $UL = LL' \subset ULL' = UL$ is minimal, a contradiction, because $LL' \in [R, T]$. Then, φ is bijective and (5) holds.

(c) Assume that (1)+(4)+(5) holds. Then (2) holds by (5). Let $L \in [R, T]$. Since $UL \in [U, S]$, we get $L \neq UL$. Deny, so that $U \subseteq L \subseteq T$ yields $S = UT = T$, a contradiction. Assume that there exists some $L' \in]L, UL[$. It follows that $UL \subset UL' \subset U(UL) = UL$, a contradiction. Then, $L \subset UL$ is minimal, giving (3).

(d) Assume that (1)+(2)+(3)+(4)+(5) holds for some $U, T \in [R, S]$ and that $[R, T] \cap [U, S] \neq \emptyset$. If $L \in [R, T] \cap [U, S]$, then $L = UL$ is a contradiction with (3). So, $[R, T] \cap [U, S] = \emptyset$. We are going to prove that $[R, S]$ is a complemented distributive lattice. We first show that T is the complement of U . We get that $R \subset U$ is minimal by (3), so that $R = U \cap T$, because $U \cap T = U$ would imply $U \subseteq T$, a contradiction. Indeed, $UT = S$ by (2) because the map $[R, T] \rightarrow [U, S]$ defined by $L \mapsto UL$ is increasing. Then, T is a complement of U .

Now, conditions (1), (2) and (5) show that $[R, S]$ is a *decomposable* lattice, that is for any $L \in [R, S]$, there is a unique pair $(L_1, L_2) \in [R, T] \times [R, U]$ such that $L = L_1 L_2$ (see [36, p.57]). Set $L_1 := L$ and $L_2 := R$ when $L \in [R, T]$. Set $L_2 := U$ and take $L_1 \in [R, T]$ such that $L = \varphi(L_1)$ when $L \in [U, S]$. The uniqueness of such (L_1, L_2) is obvious in each case. Then, $[R, S]$ is isomorphic as a lattice to $[R, T] \times [R, U]$. In particular, since $[R, T]$ and $[R, U]$ are each Boolean, and then distributive, so is $[R, S]$ by [36, Proposition 3.5.1].

Let $L \in [R, S] = [R, T] \cup [U, S]$. Assume first that $L \in [R, T]$ with complement L'' in $[R, T]$ because $R \subseteq T$ is Boolean, so that $L \cap L'' = R$ and $LL'' = T$. Setting $L' := L''U$, we get $LL' = LL''U = TU = S$.

Moreover $L \cap L' = L \cap L''U = (L \cap L'')(L \cap U) = R$. Then, L' is a complement of L in $[R, S]$.

Now, assume that $L \in [U, S]$. Set $L'' = L \cap T \in [R, T]$, and let $L' \in [R, T]$ be its complement in $[R, T]$. Then, $L \cap L' = L \cap L' \cap T = L'' \cap L' = R$. Moreover, $LL' = LUL' \supseteq UL''L' = UT = S$ gives that $LL' = S$. Then, L has a complement in $[R, S]$.

So, any element of $[R, S]$ has a complement in $[R, S]$, which is unique by distributivity. To conclude, $R \subset S$ is Boolean. \square

We are now in position to generalize and improve Ayache's result [2, Theorem 38] for an arbitrary FIP extension, using a completely different method. We will need next results.

Proposition 3.14. *An FCP extension $R \subset S$, such that $\text{Supp}(T/R) \cap \text{Supp}(S/T) = \emptyset$ for all $T \in [R, S]$, is a Boolean extension.*

Proof. By Theorem 2.26, $R_M \subset S_M$ is minimal, whence Boolean for $M \in \text{MSupp}(S/R)$, so that $R \subset S$ is Boolean by Proposition 3.4. \square

Proposition 3.15. *Let $R \subset S$ be an FCP extension and let $T \in [R, S]$ such that $\text{MSupp}(S/T) \cap \text{MSupp}(T/R) = \emptyset$. Then $R \subset S$ is Boolean if and only if $R \subset T$ and $T \subset S$ are Boolean.*

Proof. Assume first that $R \subseteq T$ and $T \subseteq S$ are Boolean. Let $M \in \text{MSupp}(S/R)$. Then either $M \in \text{MSupp}(S/T)$ (*) or $M \in \text{MSupp}(T/R)$ (**). In case (*), $M \notin \text{MSupp}(T/R)$, so that $T_M = R_M$. Hence, $[R_M, S_M] = [T_M, S_M]$ is Boolean by Proposition 3.4. In case (**), $M \notin \text{MSupp}(S/T)$, so that $T_M = S_M$. Then, $[R_M, S_M] = [R_M, T_M]$ is Boolean by Proposition 3.4. Therefore, $[R_M, S_M]$ is Boolean for each $M \in \text{MSupp}(S/R)$, and Proposition 3.4 gives that $R \subset S$ is Boolean.

The converse is given by Proposition 3.11. \square

3.2. Characterization of Boolean extensions.

Theorem 3.16. *An FIP extension $R \subset S$ is Boolean if and only if (1) and (2) hold, in which case \tilde{R} is the complement of \overline{R} :*

- (1) $\text{Supp}(\overline{R}/R) \cap \text{Supp}(S/\overline{R}) = \emptyset$.
- (2) $[R, \overline{R}]$ and $[\overline{R}, S]$ are Boolean lattices.

Proof. If $[R, S]$ is Boolean, set $T := (\overline{R})^\circ$. Then $\overline{R} \cap T = R$ and $\overline{R}T = S$, so that $\text{Supp}(\overline{R}/R) \cap \text{Supp}(S/\overline{R}) = \emptyset$ [28, Proposition 3.6], and (1) holds. The same proposition shows that $\tilde{R} = (\overline{R})^\circ$. Now (2) results from Proposition 3.15.

Conversely, assume that (1) and (2) hold. Then, Proposition 3.15 implies that $[R, S]$ is Boolean. \square

We get a generalization of Ayache's result [2, Theorem 38] as a corollary thanks to the following lemma.

Lemma 3.17. *Let $R \subset S$ be an FCP extension. Set $X := \{(T', T'') \in [R, \overline{R}] \times [\overline{R}, S] \mid \text{Supp}_{T'}(\overline{R}/T') \cap \text{Supp}_{T''}(T''/\overline{R}) = \emptyset\}$. There is a bijection $\varphi : [R, S] \rightarrow X$ defined by $\varphi(T) := (T \cap \overline{R}, \overline{R}T)$ for each $T \in [R, S]$. In particular, if $R \subset S$ has FIP, then $|[R, S]| \leq |[R, \overline{R}]| |\overline{R}, S|$.*

Proof. Let $(T', T'') \in [R, \overline{R}] \times [\overline{R}, S]$. Then, \overline{R} is also the integral closure $\overline{T'}$ of T' in T'' (and in S).

Let $T \in [R, S]$. Set $T' := T \cap \overline{R}$ and $T'' := \overline{R}T$. Then $(T', T'') \in [R, \overline{R}] \times [\overline{R}, S]$. If $T' = T''$, then $T' = T'' = \overline{R}$ implies $T = \overline{R}$ and $\text{Supp}_{T'}(\overline{R}/T') = \text{Supp}_{T''}(T''/\overline{R}) = \emptyset$. Now assume that $T' \neq T''$, then $\text{Supp}_{T'}(\overline{R}/T') \cap \text{Supp}_{T''}(T''/\overline{R}) = \emptyset$ [28, Proposition 3.6]. Hence we can define $\varphi : [R, S] \rightarrow X$ by $\varphi(T) := (T \cap \overline{R}, \overline{R}T)$ for each $T \in [R, S]$.

Now, let $T_1, T_2 \in [R, S]$ be such that $\varphi(T_1) = \varphi(T_2) = (T', T'')$. Assume $T' \neq T''$. Another use of [28, Proposition 3.6] gives that $T_1 = T_2$. If $T' = T''$, then, $T' = T'' = \overline{R}$, so that $T_1 = T_2 = \overline{R}$. It follows that φ is injective. The same reference gives that φ is bijective. \square

Corollary 3.18. *The extension $R \subset S$ is FIP Boolean if and only if the two following conditions hold:*

- (1) $R \subseteq \overline{R}$ and $\overline{R} \subseteq S$ are FIP Boolean extensions.
- (2) $|[R, S]| = |[R, \overline{R}]| |\overline{R}, S|$.

Proof. If $R \subset S$ is FIP Boolean, Proposition 3.11 implies that so are $R \subseteq \overline{R}$ and $\overline{R} \subseteq S$, giving (1). Since $\text{Supp}(\overline{R}/R) \cap \text{Supp}(S/\overline{R}) = \emptyset$ by Theorem 3.16, we infer that $|[R, S]| = |[R, \overline{R}]| |\overline{R}, S|$ [28, Lemma 3.7].

Conversely, assume that conditions (1) and (2) (which are equivalent to those of Theorem 3.16 because $R \subset S$ has FIP) hold. The map φ defined in Lemma 3.17 being injective, condition (2) shows that $\varphi([R, S]) = [R, \overline{R}] \times [\overline{R}, S]$. Therefore, there is $T \in [R, S]$ such that $\varphi(T) = (R, S)$, inducing by the properties of φ that $\text{Supp}_R(\overline{R}/R) \cap \text{Supp}_R(S/\overline{R}) = \emptyset$, (actually, the condition of Theorem 3.16(1), asserting that $R \subset S$ is Boolean). \square

Corollary 3.19. *Let $R \subset S$ be an FIP Boolean extension, where R is a local ring. Then, $R \subset S$ is either Prüfer, or integral.*

Proof. Use Theorem 3.16 and [30, Proposition 4.16]. \square

By Corollary 3.19 and Proposition 3.4, the characterization of Boolean FIP extension $R \subseteq S$ can be reduced to those that are either Prüfer or integral, with R local.

For the Prüfer case, we recover and generalize some Ayache's results on extensions of integral domains [2, Proposition 35].

Proposition 3.20. *An integrally closed extension $R \subseteq S$ is Boolean and FIP if and only if $R \subseteq S$ is a Prüfer extension and $|\text{Supp}(S/R)| = |\text{MSupp}(S/R)| < \infty$. If these conditions hold, $|[R, S]| = 2^{|\text{Supp}(S/R)|}$.*

Proof. Assume first that $|\text{Supp}(S/R)| = |\text{MSupp}(S/R)| < \infty$ and that $R \subseteq S$ is Prüfer. From [11, Proposition 6.9], we infer that $R \subseteq S$ has FIP. Moreover, we proved in [11, Proposition 6.12], that, under these conditions, $|[R_M, S_M]| = 2$ for each $M \in \text{Supp}(S/R)$, so that $R_M \subset S_M$ is Boolean for each $M \in \text{Supp}(S/R)$ by Example 3.2(1), and then $[R, S]$ is Boolean by Proposition 3.4. From [11, Proposition 6.12], we deduce that $|[R, S]| = 2^{|\text{Supp}(S/R)|}$.

Conversely, suppose that $R \subseteq S$ is Boolean and has FIP. Then, [11, Proposition 6.9] implies that $R \subseteq S$ is Prüfer and $\text{Supp}(S/R)$ is finite. Since $R \subseteq S$ is Boolean, so is $R_M \subset S_M$ for each $M \in \text{Supp}(S/R)$ by Proposition 3.4. But, $[R_M, S_M]$ is chained [11, Theorem 6.10]. It follows that $R_M \subset S_M$ is minimal for each $M \in \text{Supp}(S/R)$ by Example 3.2 (1), so that $|\text{Supp}_{R_M}(S_M/R_M)| = 1$ for each $M \in \text{Supp}(S/R)$ and then $\text{Supp}(S/R) \subseteq \text{MSupp}(S/R)$ completes the proof. \square

Corollary 3.21. *An integrally closed FIP extension $R \subset S$, with $\text{MSupp}(S/R) := \{M_1, \dots, M_n\}$, is Boolean if and only if $R \subset S$ is locally minimal. In this case, $R \subset S$ is a \mathcal{B} -extension.*

Proof. The two properties are equivalent by the proof of the above proposition. Since $\text{Supp}(S/R) = \text{MSupp}(S/R)$ holds in this case, [11, Theorem 3.6] shows that φ is a bijection. \square

The following definitions are needed for the sequel.

Definition 3.22. An integral extension $R \subseteq S$ is called *infra-integral* [26] (resp.; *subintegral* [34]) if all its residual extensions $\kappa_R(P) \rightarrow \kappa_S(Q)$, (with $Q \in \text{Spec}(S)$ and $P := Q \cap R$) are isomorphisms (resp.; and the natural map $\text{Spec}(S) \rightarrow \text{Spec}(R)$ is bijective). An extension $R \subseteq S$ is called *t-closed* (cf. [26]) if the relations $b \in S$, $r \in R$, $b^2 - rb \in R$, $b^3 - rb^2 \in R$ imply $b \in R$. The *t-closure* ${}^t_S R$ of R in S is the smallest element $B \in [R, S]$ such that $B \subseteq S$ is t-closed and the greatest element $B' \in [R, S]$ such that $R \subseteq B'$ is infra-integral. An extension $R \subseteq S$ is called *seminormal* (cf. [34]) if the relations $b \in S$, $b^2 \in R$, $b^3 \in R$ imply $b \in R$. The *seminormalization* ${}^{\dagger}_S R$ of R in S is the smallest element $B \in [R, S]$ such that $B \subseteq S$ is seminormal and the greatest element $B' \in [R, S]$ such that $R \subseteq B'$ is subintegral. The extension $R \subseteq {}^{\dagger}_S R \subseteq {}^t_S R \subseteq \overline{R} \subseteq S$ is called the *canonical decomposition* of $R \subseteq S$.

Three types of minimal integral extensions exist, characterized in the next theorem, (a consequence of the fundamental lemma of Ferrand-Olivier), so that there are four types of minimal extensions.

Theorem 3.23. [11, Theorems 2.2 and 2.3] *Let $R \subset T$ be an extension and $M := (R : T)$. Then $R \subset T$ is minimal and finite if and only if $M \in \text{Max}(R)$ and one of the following three conditions holds:*

(a) **inert case:** $M \in \text{Max}(T)$ and $R/M \rightarrow T/M$ is a minimal field extension.

(b) **decomposed case:** There exist $M_1, M_2 \in \text{Max}(T)$ such that $M = M_1 \cap M_2$ and the natural maps $R/M \rightarrow T/M_1$ and $R/M \rightarrow T/M_2$ are both isomorphisms; or, equivalently, there exists $q \in T \setminus R$ such that $T = R[q]$, $q^2 - q \in M$, and $Mq \subseteq M$.

(c) **ramified case:** There exists $M' \in \text{Max}(T)$ such that $M'^2 \subseteq M \subset M'$, $[T/M : R/M] = 2$, and the natural map $R/M \rightarrow T/M'$ is an isomorphism; or, equivalently, there exists $q \in T \setminus R$ such that $T = R[q]$, $q^2 \in M$, and $Mq \subseteq M$.

It remains to solve [2, Problem 45]: under which conditions an integral extension $R \subset S$ is Boolean and has FIP? The study is quite complicated. We are going to use the canonical decomposition of an integral ring extension and Proposition 3.4 which allows us to only consider local rings R .

Proposition 3.24. *Let $R \subset S$ be an integral FIP Boolean extension where R is local. Then, $R \subset S$ is either infra-integral or t-closed.*

Proof. Let $T := {}_t^t R$ be the t-closure of the local ring (R, M) in S , and let $T^\circ \in [R, S]$ be its complement. Let T' be the t-closure of R in T° , so that $R \subseteq T'$ is infra-integral, and then $T' \subseteq T$. It follows that $T' \subseteq T^\circ \cap T = R$. Then $T' = R$ and $R \subseteq T^\circ$ is t-closed. In the same way, let T'' be the t-closure of T° in S , so that $T'' \subseteq S$ is t-closed and then $T \subseteq T''$. Hence $S = T^\circ T \subseteq T''$, so that $T'' = S$ and $T^\circ \subseteq S$ is infra-integral.

Assume that $R \subset S$ is neither t-closed, nor infra-integral, so that $T, T^\circ \neq R, S$. Then, there are $R_1 \in [R, T]$ and $R'_1 \in [R, T^\circ]$ such that $R \subset R_1$ is minimal infra-integral, and $R \subset R'_1$ is minimal inert, with both the same crucial maximal ideal M . By [14, Propositions 7.1 and 7.4], there are two maximal chains from R to $R_1 R'_1$ with different lengths, and the same statement holds for $R \subset S$. This contradicts Condition (JH) of Proposition 2.5 since $R \subset S$ is distributive. Then, $R \subset S$ is either infra-integral, or t-closed. \square

Remark 3.25. Proposition 3.24 is no longer true if R is not local. Take a ring R with two distinct maximal ideals M_1 and M_2 , and two minimal

extensions $R \subset T_1$ ramified and $R \subset T_2$ inert with $M_1 := \mathcal{C}(R, T_1)$ and $M_2 := \mathcal{C}(R, T_2)$. Assume that $S := T_1 T_2$ exists, so that $R_{M_1} \subset S_{M_1} = (T_1)_{M_1}$ is minimal ramified, and then Boolean and $R_{M_2} \subset S_{M_2} = (T_2)_{M_2}$ is minimal inert, and then Boolean. It follows from Proposition 3.4 that $R \subset S$ is Boolean although being neither infra-integral nor t-closed. To get such a situation, we may take $S := \mathbb{Z}[i]$, $T_2 := \mathbb{Z} + 2S$, $T_1 := \mathbb{Z} + 3S$ and $R := T_1 \cap T_2 = \mathbb{Z} + 6S$. It is well known that 2 is ramified in S and 3 is inert in S . Then, $T_2 \subset S$ is a minimal ramified extension with conductor $2S$ and $T_1 \subset S$ is a minimal inert extension with conductor $3S$. Moreover, $2S$ and $3S$ are incomparable. Setting $M_1 := 2S \cap R = 2T_1$ and $M_2 := 3S \cap R = 3T_2$, [14, Proposition 6.6(a)] shows that $R \subset T_1$ is minimal ramified and $R \subset T_2$ is minimal inert with $M_1 := \mathcal{C}(R, T_1)$, $M_2 := \mathcal{C}(R, T_2)$ and $S = T_1 T_2$.

We first consider the infra-integral case for which we need the next lemma.

Lemma 3.26. *A subintegral FIP extension (resp. seminormal and infra-integral) $R \subset S$, where (R, M) is local, is Boolean if and only if $R \subset S$ is minimal ramified (resp. decomposed).*

Proof. One implication is Example 3.2 (1).

Conversely, assume that $[R, S]$ is a Boolean lattice. The atoms of $[R, S]$ are of the form $R_i := R + Rx_i$, for $i \in I := \mathbb{N}_n$, $n := |\mathcal{A}|$, where $x_i \in S$ is such that $x_i^2 \in M$ (resp. $x_i^2 - x_i \in M$) and $x_i M \subseteq M$, because $R \subset R_i$ is minimal ramified (resp.; decomposed), with $M = (R : R_i)$. Then, $S = \prod_{i \in I} R_i$ by Theorem 3.1. Let $M_i := M + Rx_i$ be the (resp.; one) maximal ideal of R_i . Assume that $n > 1$. Let $i, j \in I$ be such that $i \neq j$, so that $R = R_i \cap R_j$, with $x_i x_j \in R_i R_j$. But, $R_i \subset R_i R_j$ and $R_j \subset R_i R_j$ are minimal Theorem 3.1, so that $x_i \in M_i = (R_i : R_i R_j)$ and $x_j \in M_j = (R_j : R_i R_j)$. In the decomposed case, we may choose x_i and x_j in order that M_i and M_j are the needed conductors. It follows that M_i and M_j are ideals of $R_i R_j$, and so is $M_i \cap M_j$, which contains $M_i M_j$. But $M_i \cap M_j \subseteq R_i \cap R_j = R$. This implies that $x_i x_j \in M_i \cap M_j = M_i \cap R \cap M_j = M$, the maximal ideal of R . Then, $x_i x_j \in M$, giving $S = R + \sum_{i \in I} Rx_i$. Set $x := x_i + x_j \notin R$ and $R_x := R + Rx \neq R_i, R_j$. We get that $x^2 = x_i^2 + x_j^2 + 2x_i x_j \in M$ (resp. $x^2 = x_i + x_j + m = x + m$, where $m \in M$). Moreover, $xM \subseteq x_i M + x_j M \subseteq M$, so that $R \subset R_x$ is minimal by Theorem 3.23, and R_x is an atom of $[R, S]$. But we have $R_x \subseteq R_i R_j$, so that $R_x = R_x \cap R_i R_j = (R_x \cap R_i)(R_x \cap R_j) = R$, a contradiction. Then, $n = 1$, $S = R_1$ and $R \subset S$ is minimal. \square

It may be asked if extensions of Boolean rings and Boolean extensions are linked. Next result shows that they are quite never linked.

Proposition 3.27. *Let S be a Boolean ring and R a subring of S such that $R \subset S$ is a finite extension. Then $R \subset S$ is seminormal infra-integral and $R \subset S$ is Boolean if and only if $R \subset S$ is locally minimal.*

Proof. Since S is a Boolean ring, $R \subset S$ is seminormal integral. Because a Boolean local ring is isomorphic to $\mathbb{Z}/2\mathbb{Z}$, the residual extensions of $R \subset S$ are isomorphisms, so that $R \subset S$ is infra-integral. Now, $R \subset S$ is finite implies that it has FCP by [11, Theorem 4.2].

Assume that $R \subset S$ is Boolean. Then, $R \subset S$ has FIP by Theorem 3.1 because FCP distributive. Moreover, an appeal to Lemma 3.26 shows that $R \subset S$ is locally minimal.

Conversely, if $R \subset S$ is locally minimal, then $R \subset S$ is Boolean in view of Corollary 3.5. Indeed, $R \subset S$ has FIP [11, Proposition 3.7]. \square

Next proposition gives a characterization of arbitrary infra-integral Boolean FIP extensions.

Proposition 3.28. *An infra-integral FIP extension $R \subset S$, where (R, M) is local, is Boolean if and only if there exist $x, y \in S$ such that $S = R[x, y]$, where $x^2, xy, y^2 - y \in M$ and $xM, yM \subseteq M$.*

If these conditions hold, then $R[x, y] = R[x + y]$.

Proof. If $R \subset S$ is subintegral, we choose $y = 0$, if $R \subseteq S$ is seminormal, we choose $x = 0$. In both cases, we use Lemma 3.26 to get the equivalence. From now on, we assume that $R \subset S$ is neither subintegral, nor seminormal.

Assume that $R \subset S$ is Boolean and set $T := \textstyle\bigoplus_S R \neq R, S$. From Proposition 3.11, we deduce that $[R, T]$ and $[T, S]$ are Boolean. Then, $R \subset T$ is minimal ramified, which implies that T is also local. It follows that $T \subset S$ is minimal decomposed by Lemma 3.26, so that $\ell[R, S] = 2$, because all maximal chains of $[R, S]$ have the same length ([11, Lemma 5.4]). If $U := T^\circ$, then, $U \neq R, S$, so that $R \subset U$ and $U \subset S$ are minimal. Since $U \cap T = R$, we get that $R \subset U$ is decomposed, because it cannot be ramified. For the same reason, $U \subset S$ is minimal ramified. Let $x, y \in S$ be such that $T = R[x]$ and $U = R[y]$, so that $S = R[x, y]$, where $x^2, y^2 - y \in M$ and $xM, yM \subseteq M$. Set $M' := M + Rx$ which is the only maximal ideal of T , so that $M' = (T : S)$. Set $M'' := M + Ry \in \text{Max}(U)$. We can assume that $M'' = (U : S)$. Then, $xy \in M' \cap M'' \subseteq T \cap U \cap M' = M$.

Conversely, assume that there exist $x, y \in S$ such that $S = R[x, y]$, where $x^2, xy, y^2 - y \in M$ and $xM, yM \subseteq M$, so that $M = (R : S)$. Set $T := R[x]$ and $U := R[y]$. Then, $R \subset T$ is minimal ramified and $M + Rx$ is the maximal ideal of T , whereas, $R \subset U$ is minimal

decomposed and $M + Ry \in \text{Max}(U)$. Moreover, $T \cap U = R$ and $TU = S$. Since $(M + Rx)(M + Ry) \subseteq M$, it follows from [14, Proposition 7.6] that $R \subset S$ is an FCP infra-integral extension of length 2, so that $T \subset S$ is minimal decomposed and $U \subset S$ is minimal ramified. Hence, $[R, S] = \{R, T, U, S\}$ by [31, Theorem 6.1] and is Boolean by Theorem 3.13.

Assume that the preceding conditions hold, so that $R \subset S$ is Boolean, whence simple by Theorem 3.1. Since $R[x + y] \neq R, T, U$, we have $S = R[x + y]$. \square

We can now sum up the previous results in order to get a characterization of Boolean FIP extensions.

Theorem 3.29. *An FIP extension $R \subset S$ is Boolean if and only if, for each $M \in \text{MSupp}(S/R)$, one of the following conditions holds:*

- (1) $R_M \subset S_M$ is a minimal extension.
- (2) There exist $U, T \in [R_M, S_M]$ such that $R_M \subset T$ is minimal ramified, $R_M \subset U$ is minimal decomposed and $[R_M, S_M] = \{R_M, T, U, S_M\}$.
- (3) $R_M \subset S_M$ is a Boolean t -closed extension (equivalently, $R_M \subset S_M$ is t -closed and $\kappa_R(M) \subset \kappa_S(N)$ is a Boolean field extension, where N is the only maximal ideal of S lying above M).

Proof. From Proposition 3.4, we deduce that $[R, S]$ is Boolean if and only if, for each $M \in \text{MSupp}(S/R)$, $[R_M, S_M]$ is Boolean. An appeal to Corollary 3.19 shows that this statement is equivalent to the following: for each $M \in \text{MSupp}(S/R)$, either $R_M \subset S_M$ is Boolean and integrally closed (*), or $R_M \subset S_M$ is Boolean integral (**). In case (*) Corollary 3.21 gives that $R_M \subset S_M$ is Boolean integrally closed if and only if $R_M \subset S_M$ is Prüfer minimal. In case (**), when $R_M \subset S_M$ is integral, Proposition 3.24 says that $R_M \subset S_M$ is Boolean if and only if either $R_M \subset S_M$ is infra-integral Boolean (a), or $R_M \subset S_M$ is t -closed Boolean (b). To sum up, if $[R, S]$ is Boolean, for each $M \in \text{MSupp}(S/R)$, either (*) or (**) (a) or (**) (b) holds. Case (*) implies (1). Case (**) (a) implies either (1) or (2). Indeed, by Proposition 3.28, there exist $x, y \in S_M$ such that $S_M = R_M[x, y]$, where $x^2, xy, y^2 - y \in MR_M$ and $xMR_M, yMR_M \subseteq MR_M$. Since $M \in \text{MSupp}(S/R)$, we have $R_M \neq S_M$, so that either $x \notin R_M$ or $y \notin R_M$. If both $x, y \notin R_M$, setting $T := R_M[x]$ and $U := R_M[y]$, we get (2). If only one of $x, y \notin R_M$, we get (1). Case (**) (b) is (3). Conversely, if (1) holds, then $R_M \subset S_M$ is Boolean by Example 3.2(1). If (2) holds, then $R_M \subset S_M$ is Boolean by Proposition 3.28, setting $T := R_M[x]$ and $U := R_M[y]$ for some

$x \in T$, $y \in U$ such that $x^2, xy, y^2 - y \in MR_M$ and $xMR_M, yMR_M \subseteq MR_M$. At last, (3) is (**) (b).

We now show that, for some $M \in \text{MSupp}(S/R)$, $R_M \subseteq S_M$ is Boolean t-closed is equivalent to $R_M \subseteq S_M$ is t-closed and $\kappa(M) \subseteq \kappa(N)$ is a Boolean field extension, where N is the only maximal ideal of S lying above M . Set $R' := R_M$, $M' := MR_M$ and $S' := S_M$. Since $R' \subseteq S'$ is t-closed, $M' = (R' : S')$ and (S', M') is local, [12, Lemma 3.17]. It follows that there is only one maximal ideal N in S lying over M , so that $S_M = S_N$ ([5, Proposition 2, page 40]) and $\kappa_S(N) = S'/M'$. Then $\kappa_R(M) = R'/M' \subseteq S'/M' = \kappa_S(N)$ is an FIP field extension. Now, $R' \subseteq S'$ is Boolean if and only if $R'/M' \subseteq S'/M'$ is Boolean by Proposition 3.4. \square

It follows that the remaining t-closed case can be reduced to the case of FIP field extensions. The case of fields is the subject of the next section, because of its complexity.

Proposition 3.30. *A ring extension $R \subseteq S$ has FIP and is Boolean if and only if $R(X) \subseteq S(X)$ has FIP and is Boolean.*

Proof. One part of the proof is Proposition 3.7. So, assume that $R \subseteq S$ has FIP and is Boolean. Then, $R(X) \subseteq S(X)$ has FIP if and only if $R \subseteq {}^+_S R$ is arithmetic [29, Corollary 4.3]. If this conditions holds, the map $\psi : [R, S] \rightarrow [R(X), S(X)]$ defined by $T \mapsto T(X)$ is an order-isomorphism [13, Theorem 32]. It follows that $R(X) \subseteq S(X)$ is Boolean since Boolean conditions are preserved through ψ . To complete the proof, we need only to show that $R(X) \subseteq S(X)$ has FIP. It follows from Proposition 3.11 that $[R, {}^+_S R]$ is finite and Boolean. But, under these conditions, Theorem 3.29 yields that either $R_M = ({}^+_S R)_M$ or $R_M \subset ({}^+_S R)_M$ is minimal, for each $M \in \text{MSupp}(S/R)$. This means that $R \subseteq {}^+_S R$ is arithmetic, so that $R(X) \subseteq S(X)$ has FIP. \square

4. BOOLEAN FIP FIELD EXTENSIONS

The characterization of a Boolean extension of fields is quite different from those obtained in Theorem 3.13 and needs a special study.

4.1. FIP non separable field extensions. We will call in this paper *radicial* any purely inseparable field extension. We recall that a minimal field extension is either separable or radicial ([27, p. 371]). We will use the separable closure of a FIP algebraic field extension. In this subsection, we only consider FIP field extensions. Indeed, a finite algebraic field extension is not necessarily FIP. For instance a radicial extension $k \subseteq L$ has not FIP, when $p := c(k)$, $L := k[x, y]$, with $x^p, y^p \in k$ and $[L : k] = p^2$.

Lemma 4.1. *An FIP radical field extension $k \subseteq K$ is chained.*

Proof. Since $k \subseteq K$ has FIP, there exists $\alpha \in K$ such that $K = k[\alpha]$ by the Primitive Element Theorem. Moreover, $c(k) = p$ is a prime integer because $k \subseteq K$ is radical. Then, the monic minimal polynomial of α over k is of the form $f(X) := X^{p^n} - a = (X - \alpha)^{p^n}$, where $a := \alpha^{p^n}$ for some positive integer n .

The map $\varphi : \{0, \dots, n\} \rightarrow [k, K]$ defined by $\varphi(m) := k[\alpha^{p^m}]$ is strictly decreasing. Let $L \in [k, K]$ and $g(X)$ be the monic minimal polynomial of α over L . Then, $g(X)$ divides $f(X)$ in $K[X]$ and is of the form $g(X) = (X - \alpha)^{p^m}$ for some $m \in \{0, \dots, n\}$ because $L \subseteq K = L[\alpha]$ is radical and then the degree of $g(X)$ is a power of p . It follows that $g(X) = X^{p^m} - \alpha^{p^m} = X^{p^m} - \beta$, where $\beta := \alpha^{p^m} \in L$. By the proof of the Primitive Element Theorem, L is generated over k by the coefficients of $g(x)$, so that $L = k[\beta] = k[\alpha^{p^m}] = \varphi(m)$. Then, φ is a bijection and $[k, K]$ is chained. \square

We here take the opportunity to correct a miswriting in the proof of [29, Proposition 2.3]. The sentence: “It follows that there is only one maximal chain composing $K \subseteq L$, and it has length n ” has to be replaced with “It follows that any maximal chain composing $K \subseteq L$ has length n ”.

Theorem 4.2. *An FIP field extension $k \subset K$, with separable closure T and radical closure U such that $U, T \notin \{k, K\}$, is Boolean if and only if the following conditions hold:*

- (1) $k \subset U$ and $T \subset K$ are minimal.
- (2) $[k, K] = [k, T] \cup [U, K]$.
- (3) $k \subset T$ and $k \subset U$ are linearly disjoint.
- (4) $[k, T]$ is a Boolean lattice.

If these conditions hold, then $[k, T] \cap [U, K] = \emptyset$ and $U = T^\circ$.

Proof. Since $k \subset U$ is radical, $c(k) = p$ is a prime integer.

Assume that $k \subset K$ is Boolean. If T° is the complement of T and T' the separable closure of k in T° , then $T' \subseteq T^\circ \cap T = k$ entails that $k \subseteq T^\circ$ is radical, so that $T^\circ \subseteq U$. But $K = T^\circ T \subseteq UT$ gives $UT = K$. Moreover, $k = U \cap T$ shows that $T^\circ = U$.

We claim that $k \subset U$ is minimal. Deny, and let $U_1 \in]k, U[$ be such that $k \subset U_1$ is minimal. Since $k \subset U$ is radical, $[k, U]$ is a chain by Lemma 4.1. Let $U_1^\circ \in]T, K[$ be the complement of U_1 in $[k, K]$. Then, $K = U_1 U_1^\circ \subseteq U U_1^\circ$ implies $K = U U_1^\circ$. Moreover, $k \subseteq U \cap U_1^\circ \subseteq U \Rightarrow U \cap U_1^\circ \in [k, U]$. If $k \neq U \cap U_1^\circ$, then $U_1 \subseteq U \cap U_1^\circ \subseteq U_1^\circ \Rightarrow U_1 U_1^\circ = U_1^\circ = K$ is absurd because $U_1 \neq k$. Hence, $U \cap U_1^\circ = k$, so that U_1° is also the complement of U , which is absurd. It follows that $k \subset U$ is

minimal. Then (1), (2) and (4) hold by Theorem 3.13(3),(1),(4). We get that $[U : k] = [K : T] = p$ since $k \subset U$ and $T \subset K$ are minimal radical. But, $K = TU \Rightarrow [TU : T] = [U : k]$, and then $k \subset T$ and $k \subset U$ are linearly disjoint [6, Proposition 5, A V.13].

Conversely, assume that (1), (2), (3) and (4) hold. Then, the above conditions (2) and (4) on T and U coincide with Theorem 3.13(1),(4), so that we can use the diagram appearing in its proof, where $L \in [k, T]$:

$$\begin{array}{ccccc} k & \rightarrow & L & \rightarrow & T \\ \downarrow & & \downarrow & & \downarrow \\ U & \rightarrow & UL & \rightarrow & K \end{array}$$

Then, $L \subseteq LU$ and $L \subseteq T$ are linearly disjoint [6, Proposition 8, A V.14]. In particular, $[LU : L] = [K : T] = p$ shows that $L \subset LU$ is minimal as in Theorem 3.13(3). Let $L \in [U, K]$, and set $L' := L \cap T$, which is the separable closure of $k \subseteq L$. Then, $L' \subseteq L$ is radical. But, $UL' \subseteq L$ is both separable (because so is $U \subseteq K$ by [6, Proposition 16, page 45]) and radical, because $L' \subseteq L$ is radical, so that $L'U = L = U(L \cap T)$. Then, $[U, K] = \{UL' \mid L' \in [k, T]\}$ implies Theorem 3.13(2). Finally, $[k, K]$ is Boolean by Theorem 3.13. The missing statement $[k, T] \cap [U, K] = \emptyset$ hold by Theorem 3.13. \square

Hence, we need only to consider either radical or separable Boolean field extensions to have a complete answer. We recall [20] that a finite field extension $k \subset L$ is said to be *exceptional* if k is the radical closure and L is not the separable closure. Then, a finite exceptional field extension $k \subset L$ is never Boolean. Deny. Using notation of Theorem 4.2, $U = k = T^\circ$ implies $T = L$, a contradiction.

Proposition 4.3. *An FIP radical field extension $k \subset K$ is Boolean if and only if $k \subset K$ is minimal if and only if $c(k) = [K : k]$.*

Proof. Use Example 3.2 (1) since $[k, K]$ is a chain by Lemma 4.1 for the first equivalence, the second comes from [27, Proposition 2.2]. \square

A Galois extension $k \subset K$ is minimal if and only if $[K : k]$ is a prime integer [27, Proposition 2.2]. But this equivalence does not always hold for an arbitrary finite separable extension. Theorem 4.5 characterizes minimal separable extensions, independently of Galois Theory, contrary to Philippe’s methods [25]. She proved that a separable extension $k \subset k(x)$ is minimal if and only if the Galois group of the minimal polynomial of x is primitive [27, Proposition 2.2(3)]. See also Cox [9, page 414] for the characterization and links between primitive groups and “primitive” separable polynomials, a non trivial theory.

4.2. Finite separable field extensions. Let $k \subset L$ be a finite separable field extension, (whence FIP). Then any $T \in [k, L[$ is an intersection of finitely many \cap -irreducible elements of $[k, L[$ by Proposition 2.9. We give an upper bound of $|[k, L[$ and recall a result from Dobbs-Mullins.

Proposition 4.4. *Let $k \subset L$ be a finite separable field extension of degree n .*

- (1) *Then, $|[k, L[\leq B_n$, where B_n is the n th Bell number.*
- (2) *[10, Theorem 2.7] If k is an infinite field, then, $|[k, L[\leq 2^{n-2} + 1$.*

Proof. (1) $k \subset L$ is étale since a finite separable field extension, and then, letting Ω be an algebraic closure of k , [6, Proposition 2, page AV.29] shows that the Ω -algebra $\Omega \otimes_k L$ is diagonalizable, and then isomorphic to Ω^n , for some integer n . But $|\Omega, \Omega^n| = B_n$ by [12, Proposition 4.15], so that $[k, L[\leq B_n$. \square

Moreover, if $k \subset L$ is Boolean, any \cap -irreducible element is a co-atom by Proposition 3.9. In fact, Theorem 3.10 says that $k \subset L$ is Boolean if and only if any $T \in [k, L[$ is an intersection in a unique way of finitely many co-atoms. Thanks to principal subfields introduced in [35] and some of their properties we studied in [31], we are able to characterize co-atoms of a finite separable field extension, and then give a characterization of a finite separable Boolean field extension by using [35], from van Hoeij, Klüners and Novocin, that gives an algorithm to compute subextensions of a finite separable field extension. We recall the notation of [31], ($k_u[X]$ is the set of monic polynomials of $k[X]$).

From now on, our riding hypotheses for the subsection will be: $L := k[x]$ is a finite separable (whence FIP) field extension of k with degree n and $f(X) \in k_u[X]$ is the minimal polynomial of x over k . If $g(X) \in L_u[X]$ divides $f(X)$, we denote by K_g the k -subalgebra of L generated by the coefficients of g . For any $K \in [k, L[$, we denote by $f_K(X) \in K_u[X]$ the minimal polynomial of x over K . The proof of the Primitive Element Theorem shows that $K = K_{f_K} (*)$. Of course, $f_K(X)$ divides $f(X)$ in $K[X]$ (and in $L[X]$). If $f(X) := (X - x)f_1(X) \cdots f_r(X)$ is the decomposition of $f(X)$ into irreducible factors of $L_u[X]$, we set $\mathcal{F} := \{f_1, \dots, f_r\}$ because the f'_α 's are different by separability. For each $\alpha \in \mathbb{N}_r$, we set $L_\alpha := \{g(x) \in L \mid g(X) \in k[X], g(X) \equiv g(x) (f_\alpha(X)) \text{ in } L[X]\}$. The L'_α 's are called the *principal subfields* of $k \subset L$. It may be that $L_\alpha = L_\beta$ for some $\alpha \neq \beta$ (see [31, Example 5.17 (1)]). To get rid of this situation, we defined in [31] $\Phi : \mathcal{F} \rightarrow [k, L[$ by $\Phi(f_\alpha) = L_\alpha$. If $t := |\Phi(\mathcal{F})|$, we set $\Phi(\mathcal{F}) := \{E_1, \dots, E_t\} := \mathcal{E}$. We denote by E_β the common value of these L_α 's and by $\mathcal{E} := \Phi(\mathcal{F})$ the set of all distinct principal subfields of $k \subset L$. We set $m_\beta := f_{E_\beta}$ for $\beta \in \mathbb{N}_t$.

For $K \in [k, L[$, we set $I(K) := \{\alpha \in \mathbb{N}_r \mid f_K(X)/f_\alpha(X) \in L_u[X]\}$. We also set $J(K) := \{\beta \in \mathbb{N}_t \mid \Phi(f_\alpha) = E_\beta \text{ for all } \alpha \in I(K)\}$. For each $\beta \in \mathbb{N}_t$, we set $\mathcal{F}_\beta := \{f_\alpha \in \mathcal{F} \mid f_\alpha \text{ divides } m_\beta \text{ in } L[X]\}$.

Theorem 4.5. [31, Theorem 5.5] *Let $K \in [k, L[$. Then, $f_K(X) = (X - x) \prod_{\alpha \in I(K)} f_\alpha(X)$ and $K = \{g(x) \in L \mid g(X) \in k[X], g(X) \equiv g(x) \pmod{f_K(X)} \text{ in } K[X]\} = \cap_{\beta \in J(K)} E_\beta = \cap_{\alpha \in I(K)} L_\alpha$. In particular, $[[k, L]] \leq 2^t$ and $k \subset L$ is minimal if and only if $t = 1$.*

Proof. The inequality comes from that any $K \in [k, L]$ is an intersection of some principal subfields and gives the equivalence. \square

The inequality in Theorem 4.5 gives a better bound than the one of Proposition 4.4 because $2^t \leq 2^{n-1} \leq B_n$ thanks to the induction formula $B_{n+1} = \sum_{i=0}^n C_n^i B_i$. In case $k \subset L$ is not a Galois extension, we get a better bound than the bound of [10, Theorem 2.7].

Corollary 4.6. *Let $k \subset L$ be a finite separable field extension which is not Galois and of degree n . Then, $[[k, L]] \leq 2^{n-2}$.*

Proof. Let $f(X)$ be the minimal polynomial of the extension $k \subset L$ and set $f(X) := (X - x)f_1(X) \cdots f_r(X)$ the decomposition of $f(X)$ into irreducible factors of $L_u[X]$. Since $k \subset L$ is not Galois, at least one f_i is not of degree 1, so that $n = 1 + \sum_{i=1}^r \deg(f_i) \geq 1 + 2 + (r-1) = r + 2$. It follows that $t \leq r \leq n - 2$ implies $[[k, L]] \leq 2^t \leq 2^{n-2}$. \square

In the following, we write $K_\alpha := K_{g_\alpha}$, where $g_\alpha(X) := (X - x)f_\alpha(X)$.

Lemma 4.7. *For $K, K' \in [k, L]$, $K \subseteq K' \Leftrightarrow f_{K'}(X) \mid f_K(X)$ in $L[X]$.*

Proof. Assume that $K \subseteq K'$. Then, $f_K(X) \in K'[X]$ satisfies $f_K(x) = 0$, so that $f_{K'}(X)$ divides $f_K(X)$ in $K'[X]$, and also in $L[X]$.

Conversely, assume that $f_{K'}$ divides f_K in $L[X]$. Since Theorem 4.5 implies $K = \cap_{\alpha \in I(K)} L_\alpha$, $K' = \cap_{\alpha \in I(K')} L_\alpha$ and $f_{K'}$ divides f_K in $L[X]$, any f_α which divides $f_{K'}$ divides f_K , so that $I(K') \subseteq I(K)$ and then $K = \cap_{\alpha \in I(K)} L_\alpha \subseteq \cap_{\alpha \in I(K')} L_\alpha = K'$. \square

Proposition 4.8. *If $K, K' \in [k, L]$, then $\text{lcm}(f_K, f_{K'})$ divides $f_{K \cap K'}$ and $f_{KK'}$ divides $\text{gcd}(f_K, f_{K'})$ in $L[X]$.*

Proof. Use Lemma 4.7 applied to $K \cap K' \subseteq K, K' \subseteq KK'$. \square

We set $\mathcal{D} := \{f_K \mid K \in [k, L]\}$. Then, (\mathcal{D}, \leq) is a poset for the order \leq defined as follows: if $f_K, f_{K'} \in \mathcal{D}$, then $f_K \leq f_{K'}$ if and only if $f_K \mid f_{K'}$ in $L[X]$, which is equivalent to $K' \subseteq K$ by Lemma 4.7. In particular, sup and inf are respectively lcm and gcd in \mathcal{D} .

Corollary 4.9. *The map $\varphi : [k, L] \rightarrow \mathcal{D}$ defined by $K \mapsto f_K$ is a reversing order bijection such that $\sup(f_K, f_{K'}) = f_{K \cap K'}$ and $f_{KK'} = \inf(f_K, f_{K'})$ for $K, K' \in [k, L]$.*

Proof. φ is obviously surjective and is injective since $K = K_{f_K}$ by (*). It is reversing order by Lemma 4.7. Let $K, K' \in [k, L]$, we deduce from Proposition 4.8 that $f_{KK'} \leq f_K, f_{K'} \leq f_{K \cap K'}$. Let $K_1 \in [k, L]$ be such that $f_{K_1} \leq f_K, f_{K'}$. It follows that $K, K' \subseteq K_1$ so that $KK' \subseteq K_1$, whence $f_{K_1} \leq f_{KK'}$ and then, $f_{KK'} = \inf(f_K, f_{K'})$. A similar proof shows that $\sup(f_K, f_{K'}) = f_{K \cap K'}$. \square

We denote by $\mathcal{CA} := \{K \in [k, L] \mid K \subset L \text{ minimal}\}$ the set of co-atoms of $[k, L]$.

Proposition 4.10. *Assume that $k \subset L$ is not minimal and let $K \in]k, L[$. If $K \in \mathcal{CA}$, there is some $\beta \in \mathbb{N}_t$ such that $K = E_\beta$. Moreover, for any $\beta \in \mathbb{N}_t$, the following conditions are equivalent:*

- (1) $E_\beta \in \mathcal{CA}$.
- (2) \mathcal{F}_β is a minimal element in the set $\{\mathcal{F}_\gamma \mid \gamma \in \mathbb{N}_t\}$.
- (3) $\varphi(E_\beta)$ is a minimal element in $\mathcal{D} \setminus \{X - x\}$.

Proof. By [31, Lemma 5.10], $K = E_\beta$ for some $\beta \in \mathbb{N}_t$ since K is \cap -irreducible. Moreover, $m_\beta(X) = (X - x) \prod_{f_\alpha \in \mathcal{F}_\beta} f_\alpha(X)$ by the definition of \mathcal{F}_β .

(1) \Rightarrow (2) Assume that $E_\beta \subset L$ is minimal. We claim that \mathcal{F}_β is minimal in the poset $\{\mathcal{F}_\gamma \mid \gamma \in \mathbb{N}_t\}$. Deny, then there is some $\beta' \in \mathbb{N}_t$ such that $\mathcal{F}_{\beta'} \subset \mathcal{F}_\beta$, so that $m_{\beta'}$ divides strictly m_β . We get $E_\beta \subset E_{\beta'} \subset L$, contradicting $E_\beta \subset L$ minimal and (2) holds.

(2) \Rightarrow (3) Let E_β for some $\beta \in \mathbb{N}_t$ satisfying (2) and assume that $\varphi(E_\beta)$ is not minimal in $\mathcal{D} \setminus \{X - x\}$. Then, there is some $K \in [k, L]$ such that f_K divides strictly m_β . It follows that $E_\beta \subset K$ by Lemma 4.7. But K is an intersection of some E_γ 's by Theorem 4.5. In particular, we have $E_\beta \subset K \subseteq E_\gamma$ which implies that m_γ divides strictly m_β , so that $\mathcal{F}_\gamma \subset \mathcal{F}_\beta$, a contradiction with (2).

(3) \Rightarrow (1) Let E_β for some $\beta \in \mathbb{N}_t$ satisfying (3). Assume that $E_\beta \subset L$ is not minimal, so that there exists some $K \in [k, L]$ such that $E_\beta \subset K \subset L$. Using again Theorem 4.5, we exhibit some E_γ such that $K \subseteq E_\gamma \subset L$, giving $E_\beta \subset K \subseteq E_\gamma$. Then, m_γ divides strictly m_β , contradicting (3) and then, $E_\beta \in \mathcal{CA}$. \square

In case $k \subset L$ is Galois, we can give a characterization of \mathcal{CA} from the Galois group of the extension.

Proposition 4.11. *Let $k \subset L$ be a finite Galois extension with $n := [L : k] = \prod_{i \in \mathbb{N}_m} p_i^{e_i}$ and Galois group G .*

- (1) Let $K \in [k, L]$. Then $K \in \mathcal{CA} \Leftrightarrow$ there exists some $i \in \mathbb{N}_m$ such that $[L : K] = p_i \Leftrightarrow$ there exists some subgroup H of G of order p_i such that K is the fixed field of H in L .
- (2) $|\mathcal{CA}| \geq m$.

Proof. (1) An appeal to the Fundamental Theorem of Galois Theory shows that $K \in \mathcal{CA} \Leftrightarrow$ the group H of K -automorphisms of L has no proper subgroup $\Leftrightarrow |H| = p_i$ for some $i \in \mathbb{N}_m$ since $|G| = n \Leftrightarrow$ there exists some $i \in \mathbb{N}_m$ such that $[L : K] = |H| = p_i$ because K is the fixed field of H .

(2) For each $i \in \mathbb{N}_m$, there exists a subgroup of G of order p_i and therefore an element of \mathcal{CA} by (1), which yields $|\mathcal{CA}| \geq m$. \square

Since each element of \mathcal{CA} is some E_β , we can reorder them so that $\mathcal{CA} = \{E_1, \dots, E_s\}$ with $s \leq t$.

If $k \subset L$ is a finite separable field extension, Theorem 4.5 tells us that any $K \in [k, L]$ is an intersection of some E_β s, that we can suppose \cap -irreducible. In order to have a Boolean extension, any irreducible E_β must belong to \mathcal{CA} .

Remark 4.12. An \cap -irreducible element is not necessarily a co-atom. It is enough to take a Galois cyclic extension $k \subset L$ such that $[L : k] = p^n$, $n \geq 3$, where p is a prime integer. Then the Galois group of the extension is a chain, and so is $[k, L]$, with $\ell[k, L] = n$. Any $K \in [k, L]$ such that $\ell[k, K] \in \{1, \dots, n - 2\}$ is \cap -irreducible but not a co-atom.

We have seen in Lemma 4.7 that for each $K \in [k, L]$, there exists $g(X) = (X - x)g'(X)$, where $g'(X) \in L[X]$ is a product of some of the $f_\alpha(X)$, and satisfying $g = f_K$. Let $g(X) = (X - x)g'(X)$, where $g'(X) \in L[X]$ is a product of some of the $f_\alpha(X)$. A necessary and sufficient condition in order that there is $K \in [k, L]$ such that $g = f_K$ is gotten for $k = \mathbb{Q}$ in [35, Remark 6], a result without proof that we supply for an arbitrary field k .

Proposition 4.13. *If $g(X) \in L_u[X]$, there exists $K \in [k, L]$ such that $g = f_K \Leftrightarrow g \in \mathcal{D} \Leftrightarrow g(x) = 0$ and $[L : K_g] = \deg(g)$.*

Proof. If $g = f_K$ for some $K \in [k, L]$, then, $K = K_g$ by (*). Obviously, $g(x) = 0$. Moreover, $[L : K] = \deg(f_K) = \deg(g) = [L : K_g]$.

Conversely, if $g(x) = 0$ and $[L : K_g] = \deg(g)$ hold, set $K := K_g$. Then, $f_K(X)$ divides $g(X)$ in $K[X]$ since $g(x) = 0$. Moreover, $[L : K] = \deg(f_K) = [L : K_g] = \deg(g)$, so that $g = f_K$.

If $g(X) = X - x$, we get that $L = K$. \square

This result allows to characterize Boolean and finite separable extensions using only polynomials with the following result.

Theorem 4.14. *Let $k \subset L := k[x]$ be a finite separable field extension. The following conditions are equivalent:*

- (1) $k \subset L$ is a Boolean extension;
- (2) For any $K \in [k, L[$, there is a unique subset T of \mathcal{CA} such that $f_K = \sup\{m_\beta \mid E_\beta \in T\}$;
- (3) For any $g \in \mathcal{D} \setminus \{X - x\}$, there is a unique subset $I \subseteq \mathbb{N}_s$ such that $g = \sup\{m_\beta \mid \beta \in I\}$.

Proof. Theorem 3.10, states that $k \subset L$ is Boolean if and only if each $K \in [k, L[$ is of the form $K = \cap_{E_\beta \in T} E_\beta$ for some unique $T \subseteq \mathcal{CA}$.

(1) \Rightarrow (2) Assume first that $k \subset L$ is Boolean. Let $K \in [k, L[$ and $T := \{E_\gamma \in \mathcal{CA} \mid K = \cap_{E_\gamma \in T} E_\gamma\}$, which is unique. Corollary 4.9 yields that $f_K = \sup\{m_\beta \mid E_\beta \in T\}$. Let $T' \subseteq \mathcal{CA}$ be such that $f_K = \sup\{m_\gamma \mid E_\gamma \in T'\}$. Then, $K = \cap_{E_\gamma \in T'} E_\gamma$ and $T = T'$ follows from the uniqueness property.

(2) \Rightarrow (1) Assume that for any $K \in [k, L[$, there is a unique subset T of \mathcal{CA} such that $f_K = \sup\{m_\gamma \mid E_\gamma \in T\}$. This implies that $K = \cap_{E_\gamma \in T} E_\gamma$ by Corollary 4.9. Assume that there is some $T' \neq T$ such that $K = \cap_{E_\gamma \in T'} E_\gamma$ with $T' \subseteq \mathcal{CA}$. It follows that $f_K = \sup\{m_\gamma \mid E_\gamma \in T'\}$ and $T = T'$ because of the assumption on T . Therefore, $k \subset L$ is Boolean by Theorem 3.10.

(2) \Leftrightarrow (3) Use Corollary 4.9 and the bijection φ . □

Scholium. Here are the different steps in order to check that a finite separable extension $k \subset L$, with minimal polynomial $f(X)$, is Boolean according to Theorem 4.14:

- (1) Decompose $f(X)$ into irreducible elements of $L[X]$.
- (2) Determine the set \mathcal{E} of principal subfields.
- (3) Determine \mathcal{CA} using Proposition 4.10.
- (4) Determine \mathcal{D} using Proposition 4.13.
- (5) Check if condition (3) of Theorem 4.14 holds.

Remark 4.15. (1) Assume that $k \subset L$ is a finite separable Boolean field extension of degree n . Using the previous notation, we have $|\mathcal{CA}| = s$, which is also the value of $|\mathcal{A}|$. Using Theorem 3.1 and Proposition 4.4, we get that $2^s \leq B_n$. It follows that if we want to calculate the elements of \mathcal{CA} , it is enough to calculate the E_1, \dots, E_t , and to stop as soon as we get r distinct elements of \mathcal{CA} such that $B_n < 2^{r+1}$.

(2) Let $k \subset L := k[x]$ be a finite separable Boolean field extension and let $\mathcal{CA} = \{E_1, \dots, E_s\}$ be the set of co-atoms of the extension. Let $K := k[z] \in]k, K[$. Then Example 3.2 implies that, if $K = \cap[E_\alpha \in Y]$, where $Y \subseteq \mathcal{CA}$, we have $Y = \{E_\alpha \in \mathcal{CA} \mid z \in E_\alpha\}$. Moreover, $K^\circ = \cap[E_\beta \in \mathcal{CA} \setminus Y]$. But, since the extension is finite separable, there

exists $y \in L$ such that $K^\circ = k[y]$. Then, $\mathcal{CA} \setminus Y = \{E_\beta \in \mathcal{CA} \mid y \in E_\beta\} = \{E_\beta \in \mathcal{CA} \mid z \notin E_\beta\}$. It follows that $f_K = \sup(m_\alpha \mid E_\alpha \in Y)$ and $f_{k[y]} = \sup(m_\beta \mid E_\beta \in \mathcal{CA} \setminus Y)$. We recall that the sup is considered in \mathcal{D} , the set of the minimal polynomials of the elements of $[R, S]$.

Proposition 4.16. *A finite separable field extension $k \subset L := k[x]$ such that $\mathcal{D} = \{g(X) \in L_u[X] \mid g(x) = 0, g(X) \mid f(X) \text{ in } L[X]\}$ is Boolean. If, in addition $k \subset L$ is Galois and k an infinite field, then $k \subset L$ is minimal of degree 2.*

Proof. For $\alpha \in \mathbb{N}_r$, we have $g_\alpha(X) = (X - x)f_\alpha(X) \in \mathcal{D}$, giving that there exists some $K \in [k, L]$ such that $g_\alpha = f_K$ with $L \neq K$ because $f_L(X) = X - x$. It follows that $K \subset L$ is minimal by [31, Lemma 5.7]. Hence, $K = E_\gamma$ for some $\gamma \in \mathbb{N}_s$. Let $K' \in [k, L]$ and set $f_{K'}(X) := (X - x) \prod_{\alpha \in I} f_\alpha(X)$ for some $I \subseteq \mathbb{N}_r$. Moreover, $f_{K'} = \text{lcm}_{\alpha \in I}(\{m_\alpha\})$, for a unique I , and then a unique subset $T = \{L_\alpha\}_{\alpha \in I}$ of \mathcal{CA} satisfying the hypotheses of Theorem 4.14. (In fact, the L_α are **all** distinct and are the E_β .) Therefore, $k \subset L$ is Boolean and $\ell[k, L] = r$ by Theorem 3.1, because $|\mathcal{CA}| = |\mathcal{A}| = r$.

Now if $k \subset L$ is Galois, any f_α has degree 1. Set $n := \deg(f)$, so that $r = n - 1$, with the previous notation. Then $\ell[k, L] = n - 1$ and $||[k, L]|| = 2^{n-1}$ by Theorem 3.1. But k is infinite, which implies that $2^{n-1} = ||[k, L]|| \leq 2^{n-2} + 1$ by Proposition 4.4, which gives $n = 2$. \square


There exist finite separable Boolean extensions $k \subset L$ such that $\mathcal{D} = \{g(X) \in L_u[X] \mid g(x) = 0, g(X) \mid f(X) \text{ in } L[X]\}$ and $k \subset L$ is not Galois. Take $k := \mathbb{Q}$ and $L := k[x]$, where $x := \sqrt[3]{2}$. Then $k \subset L$ is finite separable and not Galois, because not normal. Indeed, the minimal polynomial of x is $f(X) = X^3 - 2 = (X - x)(X^2 + xX + x^2)$, with $X^2 + xX + x^2$ irreducible in $L[X]$. Then, $k \subset L$ is Boolean by Proposition 4.16.

Here is an example of Boolean extension where we show how the irreducible divisors of the minimal polynomial provides the subextensions of a finite separable extension of fields.

Example 4.17. [29, Remark 5.19] Let $k := \mathbb{Q}$, $x := \sqrt[6]{2}$ and set $L := k[x]$, which is a finite separable extension of k , but not Galois.

The monic minimal polynomial of x over k is $f(x) := X^6 - 2 = (X - x)(X + x)(X^2 + xX + x^2)(X^2 - xX + x^2)$, which is its decomposition into irreducible polynomials over L . Set $f_1(X) := X + x$, $f_2(X) := X^2 + xX + x^2$, $f_3(X) := X^2 - xX + x^2$ and $g_\alpha(X) := (X - x)f_\alpha(X)$, for $\alpha = 1, 2, 3$. Then, $g_1(X) = X^2 - x^2 = X^2 - \sqrt[3]{2}$, $g_2(X) = X^3 - x^3 = X^3 - \sqrt{2}$ and $g_3(X) = X^3 - 2xX^2 + 2x^2X - x^3$. It follows that $K_1 = k[\sqrt[3]{2}]$, $K_2 = k[\sqrt{2}]$ and $K_3 = L$. Then, $L_1 = K_1$ and $L_2 = K_2$ by

[31, Lemma 5.10]. Moreover, no subextension $K \in [k, L[$ is such that $g_3 = f_K$ since $K_3 = L$. Let $K \in [k, L]$ be such that g_3 divides strictly f_K in $L[X]$. Then, $[L : K] = \deg(f_K) > 3$ gives that $[L : K] = 6$, so that $K = k = L_3$ because $f_{L_3} = f$ [31, Proposition 5.8]. To end, $L_1 \cap L_2 = k = L_3$. Hence, $[k, L] = \{k, L_1, L_2, K\}$ is a Boolean lattice by Theorem 3.13, and we get the following diagram:



Remark 4.18. [31, Example 5.17 (2)] Let $k := \mathbb{Q}$ and $L = k[x]$, where $x := \sqrt{2} + \sqrt{3}$. The monic minimal polynomial of x over k is $f(X) = X^4 - 10X^2 + 1 = (X - x)(X + x)(X - x^{-1})(X + x^{-1})$. Set $f_1(X) := X + x$, $f_2(X) = X - x^{-1}$, $f_3(X) = X + x^{-1}$. We get $K_1 = L_1 = k[\sqrt{6}]$, $K_2 = L_2 = k[\sqrt{3}]$ and $K_3 = L_3 = k[\sqrt{2}]$. In particular, $L_\alpha = E_\alpha$ for each α and $L_\alpha \cap L_\beta = k$ for $\alpha \neq \beta$, $\alpha, \beta \in \mathbb{N}_3$, which shows that $[k, L] = \{k, L_1, L_2, L_3, L\}$ is not Boolean because $\ell[k, L] = 2$ and $|[k, L]| = 5 \neq 2^2$, but $k \subset L$ is Galois. Therefore, although $k \subset L_\alpha$ is Boolean for $\alpha \in \mathbb{N}_2$, the product $k \subset L_1 L_2 = L$ is not Boolean. We also observe that despite the fact that $k \subset L_\alpha$ and $L_\alpha \subset L$ are Boolean for $\alpha \in \mathbb{N}_3$, $k \subset L$ is not Boolean.

However, we are able to characterize Boolean Galois extensions.

Theorem 4.19. *Let $k \subset L$ be a finite separable extension with normal closure N . If $k \subset N$ is a cyclic extension with a square free degree, then $k \subset L$ is a Boolean extension.*

In particular, a finite Galois extension $k \subset L$ with Galois group G is Boolean if and only if $k \subset L$ is cyclic whose degree is square free.

Proof. We begin to prove the second part of the Theorem. Let \mathcal{G} be the set of subgroups of G . For $H, H' \in \mathcal{G}$, we denote by $\langle H, H' \rangle$ the subgroup of \mathcal{G} generated by H and H' . Define $\varphi : [k, L] \rightarrow \mathcal{G}$ by $\varphi(K) := \text{Aut}_K(L)$, the group of K -automorphisms of L , for each $K \in [k, L]$ and $\psi : \mathcal{G} \rightarrow [k, L]$ by $\psi(H) := \text{Fix}(H)$, the fixed field of H in L , for each $H \in \mathcal{G}$. The Fundamental Theorem of Galois Theory for finite extensions shows that φ and ψ are reversing order isomorphisms of lattices, with $\varphi = \psi^{-1}$. Therefore, $[k, L]$ is a Boolean lattice if and only if \mathcal{G} is a Boolean lattice. To conclude, use [37, Corollary 2] which says that for a finite group G , the lattice of its subgroups is a Boolean lattice if and only if G is a cyclic group whose order is square free.

Now, if $k \subset L$ is a finite separable extension whose normal closure is N such that $k \subset N$ is a cyclic extension with a square free degree, then $k \subset L$ is Boolean by Proposition 3.11 because $k \subset N$ is Boolean. \square

Remark 4.20. The last part of the Theorem has no converse, unless adding some new assumptions as in Theorem 3.13. Consider Example 4.17. The normal closure N of the extension $k \subset L$ is generated over k by x and its conjugates, which are the zeroes $\{\pm x, \pm jx, \pm j^2x\}$, of $f(X) = X^6 - 2$, where $j = (1 + i\sqrt{3})/2$. Then, $N = k[x, jx, j^2x]$. Moreover, $k \subset N$ is Galois. Assume that $k \subset N$ is Boolean. Then, $k \subset N$ is a cyclic extension by Theorem 4.19. Set $K'_1 := k[j^2x^2] \subset N$. Since $(j^2x^2)^3 = 2$, we get that $[K'_1 : k] = 3 = [K_1 : k]$, and we have two subextensions of $k \subset N$ of degree 3, a contradiction for a cyclic extension (see [6, AV, page 81]). Then, $k \subset N$ is not Boolean.

The two next examples exhibit Galois Boolean field extensions.

Example 4.21. (1) Let n be a positive integer, $n \geq 2$. In view of [6, AV.152, Exercice 3)], there exists a cyclic extension of \mathbb{Q} of degree n . It is enough to take a square free integer n and to use Theorem 4.19 to get a Boolean extension.

(2) Let $k := \mathbb{F}_2 = \mathbb{Z}/2\mathbb{Z}$ be the finite field with two elements, and let K_n be the cyclic extension of k of degree n . Set $L := K_{30}$. The subfields of L are the K_n , where n divides 30. In view of Theorem 4.19, $k \subset L$ is a Boolean extension, because cyclic of degree a square free integer, and $[k, L] = \{K_n \mid n = 1, 2, 3, 5, 6, 10, 15, 30\}$.

Proposition 4.22. *Let $k \subset L$ be a finite Galois Boolean extension and let $T, U \in]R, S[$. Then $U = T^\circ$ if and only if $k \subset T$ and $k \subset U$ are linearly disjoint with $L = TU$.*

Proof. If $k \subset T$ and $k \subset U$ are linearly disjoint, then $T \cap U = k$, so that $U = T^\circ$ since $TU = L$.

Conversely, assume that $U = T^\circ$. We are going to show how U is build from T . Since $k \subset L$ is a finite Galois Boolean extension, Theorem 4.19 shows that $k \subset L$ is cyclic, $n := [L : k]$ is square free, and so is $[T : k]$. Set $n = p_1 \cdots p_k p_{k+1} \cdots p_r$ where the p_i 's are distinct prime integers ordered such that $m := [T : k] = p_1 \cdots p_k$, and set $l := n/m$. Since $k \subset L$ is cyclic, there exists $V \in [k, L]$ such that $[V : k] = l$. It follows that $(m, l) = ([T : k], [V : k]) = 1$, so that $V \cap T = k$ and $TV := L$. Indeed, $T, V \subset TV \subseteq L$ shows that m, l dividing $[TV : k]$, gives that $[L : k] = n = ml$ divides $[TV : k]$, which leads to $TV = L$. Then, $V = T^\circ = U$. Under these conditions, we have $[TU : k] = [L : k] = ml = [T : k][U : k]$, which shows that $k \subset T$ and $k \subset U$ are linearly disjoint. \square

We can say more about distributive Galois extension non necessarily Boolean, involving a result from Dobbs-Mullins.

Proposition 4.23. *Let $k \subset L$ be a finite Galois extension with degree $n (= \prod_{i=1}^m p_i^{e_i}$ the factorization into prime integers).*

- (1) [10, Proposition 2.2] *If $k \subset L$ is Abelian, then $\ell[k, L] = \sum_{i=1}^m e_i$.*
- (2) *If in addition $k \subset L$ is distributive, then $k \subset L$ is cyclic, $\ell[k, L] = \sum_{i=1}^m e_i$ and $|[k, L]| = \tau(n)$, where $\tau(n)$ is the number of divisors of n .*
- (3) *If in addition $k \subset L$ is Boolean, $\ell[k, L] = m$ and $|[k, L]| = 2^m$.*

Proof. (2) Let G be the Galois group of $k \subset L$ and \mathcal{G} be the set of subgroups of G . Since $k \subset L$ is distributive, so is \mathcal{G} . Then, G is cyclic by [32, Page 97], and so is the extension $k \subset L$. The first part of (2) comes from (1) since a cyclic extension is Abelian. Moreover, there is a bijection between the subgroups of a cyclic group of order n and the divisors of n , as there is a bijection between \mathcal{G} and $[k, L]$. This gives the last equality.

(3) Assume that $k \subset L$ is Boolean, then cyclic by Theorem 4.19 and n is square free, so that $e_i = 1$ for each i . Hence, $\ell[k, L] = m$ and $|[k, L]| = 2^m$, since $k \subset L$ is Boolean by Theorem 3.1. \square

In a recent paper [31], we characterized ring extensions $R \subset S$ of length 2 and gave the value of $|[R, S]|$. It is then easy to characterize an extension of length 2 which is Boolean.

Proposition 4.24. *Let $R \subset S$ be an FIP extension of length 2. Then $R \subset S$ is Boolean if and only if $|[R, S]| = 4$, and, if and only if one of the following condition holds:*

- (1) $|\text{Supp}(S/R)| = 2$ and $\text{Supp}(S/R) \subseteq \text{Max}(R)$.
- (2) $R \subset S$ is infra-integral such that $\text{Supp}(S/R) = \{M\}$, ${}^+_S R \neq R, S$ and $(R : S) = M$.
- (3) $R \subset S$ is t -closed integral such that $\text{Supp}(S/R) = \{M\}$, $M = (R : S) \in \text{Max}(S)$, and one of the following conditions holds:
 - (a) $R/M \subset S/M$ is neither radical nor separable, nor exceptional.
 - (b) $R/M \subset S/M$ is a finite separable field extension and $t = 2$, where t is the number of principal subfields of S/M different from R/M .

Proof. Assume that $R \subset S$ is Boolean. Then $|[R, S]| = 4$ by Theorem 3.1. Conversely, if $|[R, S]| = 4$, then $[R, S] = \{R, T, U, S\}$ for some $U, T \in]R, S[$, where T and U are incomparable, so that $R \subset S$ is Boolean by Theorem 3.13. Now the second equivalence comes from [31, Theorem 6.1] which gives the different cases for a length 2 extension $R \subset S$ to satisfy $|[R, S]| = 4$. \square

REFERENCES

- [1] A. Ayache, When is the set of intermediate rings a finite boolean algebra?, *Inter. Elec. J. Algebra*, **8**, (2010), 129–139.
- [2] A. Ayache, A constructive study about the set of intermediate rings, *Comm. Algebra*, **41** (2013), 4637–4661.
- [3] A. Ayache, The set of intermediate rings as a boolean algebra, *Comm. Algebra*, **43** (2015), 1377–1386.
- [4] N. Bourbaki; Algèbre Commutative, Chs. 1–2, Hermann, Paris, 1961.
- [5] N. Bourbaki, Algèbre Commutative, Chs. 5–6, Hermann, Paris, 1964.
- [6] N. Bourbaki, Algèbre, Chs. 4–7, Masson, Paris, 1981.
- [7] P.-J. Cahen, D. E. Dobbs and T.G. Lucas, Valuative domains, *J. Algebra Appl.*, **9** (2010), 43–72.
- [8] G. Calugareanu, Lattice concepts of module theory, Springer, New York, 2000.
- [9] D. A. Cox, Galois theory, Wiley/Interscience, 2004.
- [10] D. E. Dobbs and B. Mullins, On the lengths of maximal chains of intermediate fields in a field extension, *Comm. Algebra*, **29** (2001), 4487–4507.
- [11] D. E. Dobbs, G. Picavet and M. Picavet-L’Hermitte, Characterizing the ring extensions that satisfy FIP or FCP, *J. Algebra*, **371** (2012), 391–429.
- [12] D. E. Dobbs, G. Picavet and M. Picavet-L’Hermitte, Transfer results for the FIP and FCP properties of ring extensions, *Comm. Algebra*, **43** (2015), 1279–1316.
- [13] D. E. Dobbs, G. Picavet and M. Picavet-L’Hermitte, When an extension of Nagata rings has only finitely many intermediate rings, each of those is a Nagata ring?, *Int. J. Math. Math. Sci.*, **Vol 2014** (2012), Article ID 315919, 13 pp..
- [14] D. E. Dobbs, G. Picavet, M. Picavet-L’Hermitte and J. Shapiro, On intersections and composites of minimal ring extensions, *JPJANTA*, **26** (2012), 103–158.
- [15] D. E. Dobbs and J. Shapiro, Patching together a minimal overring, *Houston J. Math.* **36** (2010), 985–995.
- [16] D. E. Dobbs and J. Shapiro, Catenarian Numbers, *Comm. Algebra* **42** (2014), 2603–2623.
- [17] T. Donnellan, Lattice Theory, Pergamon Press, Oxford, 1968.
- [18] D. Ferrand and J.-P. Olivier, Homomorphismes minimaux d’anneaux, *J. Algebra*, **16** (1970), 461–471.
- [19] R. Gilmer, Multiplicative Ideal Theory, Dekker, New York, 1972.
- [20] R. Gilmer and W. Heinzer, On the existence of exceptional field extensions, *Bull. Amer. Math. Soc.*, **74** (1968), 545–547.
- [21] G. Grätzer, General Lattice Theory, Academic Press, New York (1978).
- [22] I. Kaplansky, Commutative Rings, rev. ed., Univ. Chicago Press, Chicago, 1974.
- [23] S. Mac Lane, G. Birkhoff, Algebra, American Mathematical Soc., 1999.
- [24] C. Năstăsescu and F. Van Oystaeyen, Dimensions of Ring Theory, Mathematics and its applications, D. Reidel Publishing Co., Dordrecht, 1987.
- [25] A. Philippe, Extension minimale de corps, Mémoire de D.E.A., Paris (1969), unpublished.

- [26] G. Picavet and M. Picavet-L'Hermitte, T-Closedness, pp. 369–386, in: *Non-Noetherian Commutative Ring Theory, Math. Appl. 520*, Kluwer, Dordrecht, 2000.
- [27] G. Picavet and M. Picavet-L'Hermitte, About minimal morphisms, pp. 369–386, in: *Multiplicative Ideal Theory in Commutative Algebra*, Springer, New York, 2006.
- [28] G. Picavet and M. Picavet-L'Hermitte, Prüfer and Morita hulls of FCP extensions, *Comm. Algebra*, **43** (2015), 102–119.
- [29] G. Picavet and M. Picavet-L'Hermitte, Some more combinatorics results on Nagata extensions, *Palestine J. of Maths*, **5** (2016), 1–15.
- [30] G. Picavet and M. Picavet-L'Hermitte, Quasi-Prüfer extensions of rings, pp. 307–336, in: *Rings, Polynomials and Modules*, Springer, 2017.
- [31] G. Picavet and M. Picavet-L'Hermitte, Ring extensions of length two, JAA, DOI: 10.1142/S0219498819501743.
- [32] S. Roman, Lattices and ordered sets, Springer, New York, 2008.
- [33] R. P. Stanley, Enumerative combinatorics, Vol. 1, Second edition, Version of 15 July 2011.
- [34] R. G. Swan, On seminormality, *J. Algebra*, **67** (1980), 210–229.
- [35] M. van Hoeij, J. Klüners and A. Novocin, Generating subfields, *Journal of Symbolic Computation*, **52** (2013), 17–34.
- [36] N. White, Theory of Matroids, Cambridge University Press, London, New York, 1986.
- [37] G. Whitson, Finite groups whose subgroup, composition subgroup, or normal subgroup lattice is an ortholattice, *Algebra Universalis*, **8** (1978), 123–127.

MATHÉMATIQUES, 8 RUE DU FOREZ, 63670 - LE CENDRE, FRANCE
E-mail address: picavet.mathu (at) orange.fr