

# ON THE FUNDAMENTAL GROUP OF COMPLETE MANIFOLDS WITH ALMOST EUCLIDEAN VOLUME GROWTH

JIANMING WAN

ABSTRACT. It is proved that the fundamental group of a complete Riemannian manifold with nonnegative Ricci curvature and certain volume growth conditions is trivial or finite.

## 1. INTRODUCTION

Throughout the paper  $M$  denotes a complete noncompact Riemannian  $n$ -manifold with nonnegative Ricci curvature. Let  $V_p(r)$  be the volume of the metric ball  $B_p(r)$  origin at  $p$  with radius  $r$  in  $M$ . By Bishop-Gromov volume comparison,  $\frac{V_p(r)}{\omega_n r^n}$  is a decreasing function, where  $\omega_n$  is the volume of unit ball in  $\mathbb{R}^n$ . So the limit is existent as  $r$  goes to infinite. Denote the *volume growth* of  $M$  by

$$\alpha_M := \lim_{r \rightarrow \infty} \frac{V_p(r)}{\omega_n r^n}.$$

The  $\alpha_M$  is independent of  $p$  and so a global geometric invariant. Moreover, the volume comparison also implies that  $0 \leq \alpha_M \leq 1$  and  $\alpha_M = 1$  if and only if  $M$  is isometric to  $\mathbb{R}^n$ . We say that  $M$  has *Euclidean volume growth* (or large volume growth) if  $\alpha_M > 0$ .

The volume comparison theorem implies that

$$0 < \frac{V_p(2r)}{V_p(r)} \leq 2^n$$

for all  $r > 0$  and Euclidean volume growth condition implies that

$$\lim_{r \rightarrow \infty} \frac{V_p(2r)}{V_p(r)} = 2^n.$$

The main result of this note is

**Theorem 1.1.** *Given  $n$ , there is a constant  $C(n) < 2^n$  such that if an open  $n$ - manifold  $M$  satisfies*

1)

$$(1.1) \quad \frac{V_p(2r)}{V_p(r)} > C(n)$$

for some  $p \in M$  and **all**  $r > 0$ , then  $M$  is simple connected.

2)

$$(1.2) \quad \liminf_{r \rightarrow \infty} \frac{V_p(2r)}{V_p(r)} > C(n),$$

for some  $p \in M$ , then the fundamental group  $\pi_1(M)$  is finite.

---

2010 *Mathematics Subject Classification.* 53C20.

*Key words and phrases.* nonnegative Ricci curvature, volume growth, fundamental group.

We should note that even though  $M$  has Euclidean volume growth, one can not deduce that  $M$  is simple connected. So formula (1.1) holding for all  $r > 0$  is important.

Set  $\epsilon(n) = n - \log_2 \frac{C(n)+2^n}{2}$ . We see immediately that 2) of Theorem 1.1 implies the following

**Corollary 1.2.** *Given  $n$ , there is a constant  $\epsilon(n)$  such that if an open  $n$ - manifold  $M$  satisfies*

$$(1.3) \quad \lim_{r \rightarrow \infty} \frac{V_p(r)}{r^{n-\epsilon}} > 0,$$

for some  $p \in M$ ,  $0 \leq \epsilon < \epsilon(n)$ , then  $\pi_1(M)$  is finite.

This shows a gap phenomenon for a well-known result of Peter Li [3] and Anderson [1] states that  $\pi_1(M)$  is finite provided  $M$  has Euclidean volume growth.

On the other hand, Anderson has proved that (see Theorem 1.1 in [1]) under condition (1.3), every finitely generated subgroup of  $\pi_1(M)$  is actually of polynomial growth of order  $\leq \epsilon < 1$ . In [6] Bingye Wu proved that under condition (1.3)  $\pi_1(M)$  is finitely generated. But every infinite group of finitely generated has polynomial growth of order at least 1 (I thank the referee for pointing out this fact. See Section 3). So Corollary 1.2 is also a consequence of Anderson and Wu's results.

**Acknowledgment:** I would like to thank the referee for his (her) invaluable suggestions. The referee's explanation clarify some understanding of mine on Anderson's paper [1].

## 2. A RELATED VOLUME RATIO

In this section we prove an estimate on the volume ratio  $V_p(2r)/V_p(r)$  related to certain generated elements of  $\pi_1(M)$  (Lemma 2.4 below). The main ingredients are Sormani's uniform cut lemma [5] and some ideas due to Shen [4].

**2.1. A uniform estimate.** Let  $g \in \pi_1(M, p)$  and  $\pi : \tilde{M} \rightarrow M$  be the universal cover. Following [5], we say that  $\gamma$  is a *minimal representative geodesic loop* (based at  $p$ ) of  $g$  if  $\gamma = \pi \circ \tilde{\gamma}$ , where  $\tilde{\gamma}$  is a minimal geodesic connecting  $\tilde{p}$  and  $g\tilde{p}$ . So  $L(\gamma) = d_{\tilde{M}}(\tilde{p}, g\tilde{p})$ .

Given a group  $G$ , we say that  $\{g_1, g_2, g_3, \dots\}$  is an *ordered set of independent generators* of  $G$  if every  $g_i$  can not be expressed as the previous generators and their inverses,  $g_1, g_1^{-1}, \dots, g_{i-1}, g_{i-1}^{-1}$ .

In [5] Sormani proved the following two lemmas.

**Lemma 2.1.** (*halfway lemma*) *There exists an ordered set of independent generators  $\{g_1, g_2, \dots\}$  of  $\pi_1(M, p)$  with minimal representative geodesic loops  $\gamma_k$  of length  $d_k$  such that*

$$d_M(\gamma_k(0), \gamma_k(d_k/2)) = d_k/2.$$

*In particular, we have a sequence of such generators if  $\pi_1(M, p)$  is infinitely generated.*

**Lemma 2.2.** (*uniform cut lemma*) *Let  $M^n$  ( $n \geq 3$ ) be a complete manifold with  $\text{Ric} \geq 0$ . Let  $\gamma$  be a noncontractible geodesic loop based at  $p_0$  of length  $L(\gamma) = 2D$  such that*

- (1) *If  $\sigma$  based at  $p_0$  is a loop homotopic to  $\gamma$ , then  $L(\sigma) \geq 2D$ ;*
- (2) *The  $\gamma$  is minimal on  $[0, D]$  and  $[D, 2D]$ .*

*Then there is a constant  $S_n$  depending on  $n$  such that if  $x \in \partial B_{p_0}(rD)$  where  $r \geq 1 + 2S_n$ , then*

$$d(x, \gamma(D)) \geq (r-1)D + 4S_n D.$$

The main idea of proof of uniform cut lemma is to lift geodesic loop to the universal covering space and research carefully the related excess function. It contains a nice application of Abresch-Gromoll's estimate on excess function [2]. The above two lemmas allow her to show that Milnor conjecture holds for the manifold with so called small linear diameter growth.

Let  $\gamma$  be a minimal representative geodesic loops based at  $p$  of  $L(\gamma) = d$  satisfying Lemma 2.1. The below estimate is important for our purpose.

**Lemma 2.3.** *Let  $\sigma$  be a geodesic issuing from  $p$  such that  $\sigma(t)$  is minimal on  $[0, d)$ . Then there is a constant  $S(n)$  such that*

$$h := d_M(\gamma(d/2), \sigma|_{[0, d)}) \geq S(n)d.$$

*Proof.* We set  $h_1 := d_M(\gamma(d/2), \sigma|_{[0, d/2)})$  and  $h_2 := d_M(\gamma(d/2), \sigma|_{[d/2, d)})$ . By Lemma 2.2, we have

$$H = d_M(\gamma(d/2), \sigma(d/2)) \geq 2S(n)d,$$

where  $S(n)$  is a universal constant,  $S(n) = \frac{n}{n-1} \frac{1}{4} \frac{1}{3^n} (\frac{n-2}{n-1})^{n-1}$ .

Suppose that  $h_1 = d_M(\gamma(d/2), \sigma(r_0))$ . By the triangle inequality one has

$$h_1 \geq \frac{d}{2} - r_0$$

and

$$h_1 \geq H - (\frac{d}{2} - r_0).$$

Then

$$h_1 \geq H/2 \geq S(n)d.$$

We also note that  $d_M(\gamma(d/2), \sigma(d)) \geq d/2$ . So similarly one has

$$h_2 \geq H/2 \geq S(n)d.$$

It follows that

$$h = \min(h_1, h_2) \geq S(n)d.$$

□

**2.2. A volume's ratio.** Continuing with notations  $p, d$  in Lemma 2.3, we shall prove

**Lemma 2.4.** *We have the following ratio of volume*

$$\frac{V_p(2d)}{V_p(d)} \leq (1 - (\frac{2S(n)}{3})^n)(2^n - 1) + 1.$$

Before giving the proof of Lemma 2.4, (following [4]) we introduce some necessary notations. Let  $\Sigma_p$  be a close subset of unit tangent sphere  $S_p M \subset T_p M$ . Let  $B_{\Sigma_p}(r)$  be the set of points  $x \in B_p(r)$  such that there exists a minimal geodesic  $\gamma$  from  $p$  to  $x$  with  $\gamma'(0) \in \Sigma_p$ . We write  $V_{\Sigma_p}(r)$  for the volume of  $B_{\Sigma_p}(r)$ .

We denote by  $\Sigma_p(r)$  the set of unit vectors  $v \in S_p M$  such that  $\gamma(t) = \exp_p(tv)$  is minimal on  $[0, r)$ .

*Proof. of Lemma 2.4.* We write  $y = \gamma(d/2)$ . Since  $h \leq d/2$ , we have  $B_p(d) \supset B_y(h) \cup B_{\Sigma_p(d)}(d)$ . By the definition of  $h$ , this gives  $V_p(d) \geq V_y(h) + V_{\Sigma_p(d)}(d)$ , i.e.

$$(2.1) \quad 1 \geq \frac{V_y(h)}{V_p(d)} + \frac{V_{\Sigma_p(d)}(d)}{V_p(d)}.$$

**Claim 1:**

$$(2.2) \quad \frac{V_y(h)}{V_p(d)} \geq \left(\frac{2S(n)}{3}\right)^n.$$

By the Bishop-Gromov comparison theorem,  $\mu^n V_p(r) \geq V_p(\mu r)$  for  $\mu \geq 1$ . By Lemma 2.3 we have  $h \geq S(n)d$ . So

$$V_y(h) \geq V_y(S(n)d) \geq \left(\frac{2S(n)}{3}\right)^n V_y(3d/2).$$

Since  $B_y(3d/2) \supset B_p(d)$ , we have  $V_y(3d/2) \geq V_p(d)$ . Thus we obtain (2.2).

**Claim 2:**

$$(2.3) \quad \frac{V_{\Sigma_p(d)}(d)}{V_p(d)} \geq \frac{1}{2^n - 1} \left( \frac{V_p(2d)}{V_p(d)} - 1 \right).$$

Following the observation of Shen (c.f. [4] Lemma 2.4), we see that

$$B_p(2r) \setminus B_p(r) \subset B_{\Sigma_p(r)}(2r) \setminus B_{\Sigma_p(r)}(r).$$

Then we have

$$\begin{aligned} V_p(2r) - V_p(r) &\leq V_{\Sigma_p(r)}(2r) - V_{\Sigma_p(r)}(r) \\ &\leq (2^n - 1) V_{\Sigma_p(r)}(r). \end{aligned}$$

The second in equality follows from the generalized volume comparison (Lemma 2.2 of [4]). Thus

$$\frac{V_{\Sigma_p(r)}(r)}{V_p(r)} \geq \frac{1}{2^n - 1} \left( \frac{V_p(2r)}{V_p(r)} - 1 \right).$$

Jointing formulas (2.1), (2.2) and (2.3), we establish the lemma.  $\square$

### 3. A PROOF OF THEOREM 1.1

We set

$$C(n) = \left(1 - \left(\frac{2S(n)}{3}\right)^n\right)(2^n - 1) + 1.$$

If  $\frac{V_p(2r)}{V_p(r)} > C(n)$  for all  $r > 0$ , then there is no nontrivial generator satisfying Lemma 2.4. So  $M$  is simple connected. Thus the first part of Theorem 1.1 is proved.

The proof of second part of Theorem 1.1 is divided into two steps.

Firstly,  $\pi_1(M, p)$  is finitely generated. We argue by contradiction. Assume  $\pi_1(M, p)$  is infinitely generated, then by Lemma 2.2, there is a sequence  $\{d_k\}$ ,  $d_k \rightarrow \infty$  as  $k \rightarrow \infty$  satisfying Lemma 2.4, i.e.

$$\frac{V_p(2d_k)}{V_p(d_k)} \leq C(n),$$

for all  $k \geq 1$ . This contradicts to condition (1.2).

Secondly, condition (1.2) implies that  $V_p(r) \geq C \cdot r^{n-\epsilon}$  for some  $\epsilon < 1$  and sufficiently large  $r$ . So by Anderson's result [1],  $\pi_1(M)$  has polynomial growth of order  $\leq \epsilon < 1$ .

The form of  $C(n)$  allows us to write  $C(n) = 2^{n-\epsilon}$ ,  $\epsilon < 1$ . By condition (1.2), there exists  $r_0 > 0$ , for all  $r \geq r_0$ , one has

$$V_p(2r) \geq 2^{n-\epsilon} V_p(r).$$

So

$$V_p(r_0) \leq (2^{n-\epsilon})^{-1} V_p(2r_0) \leq \cdots \leq (2^{n-\epsilon})^{-k} V_p(2^k r_0),$$

$k \in N$ . For any  $r \geq r_0$ , we can assume  $r \in [2^k r_0, 2^{k+1} r_0]$  for some  $k \in N$ . Then

$$\begin{aligned} V_p(r) \geq V_p(2^k r_0) &\geq \frac{V_p(r_0)}{r_0^{n-\epsilon}} (2^k r_0)^{n-\epsilon} \\ &\geq \frac{V_p(r_0)}{r_0^{n-\epsilon}} (2^k r_0 \cdot \frac{r}{2^{k+1} r_0})^{n-\epsilon} \\ &= \frac{V_p(r_0)}{(2r_0)^{n-\epsilon}} r^{n-\epsilon}. \end{aligned}$$

It follows that  $V_p(r) \geq C \cdot r^{n-\epsilon}$  for all  $r > r_0$ .

An algebraic fact: If  $\Gamma$  is an infinite group with generators  $S = \{g_1, \dots, g_k\}$ , then  $\#U(r) \geq r$  for all  $r \in N$ , where  $U(r)$  is the set of elements with word length  $\leq r$  with respect to  $S$ . In particular,  $\Gamma$  has polynomial growth of order at least 1. (This proof is provided by referee) To see this we argue by contradiction. Let  $r$  be the smallest integer so that  $\#U(r) < r$ , then  $r-1 \leq \#U(r-1) \leq \#U(r) < r$ . This shows  $U(r) = U(r-1)$ . In other words, any word of length  $r$  can be expressed as a word of length  $\leq r-1$ . It follows that  $\Gamma = U(r-1)$ , which is finite, a contradiction.

The second part of Theorem 1.1 follows from above immediately.

*Remark 3.1.* Our proof of finite generation of  $\pi_1(M)$  is much different to Wu's arguments under condition (1.3). Wu's proof was based on the estimate of ordered set of independent generators with minimal representative geodesic loops.

#### REFERENCES

1. Anderson, M.T. *On the topology of complete manifolds of nonnegative Ricci curvature*. Topology 29 (1990), no. 1, 41-55.
2. Abresch, U. and Gromoll, D. *On complete manifolds with nonnegative Ricci curvature*. J. Amer. Math. Soc. 3 (1990), no. 2, 355-374.
3. Li, Peter *Large time behavior of the heat equation on complete manifolds with nonnegative Ricci curvature*. Ann. of Math. (2) 124 (1986), no. 1, 1-21.
4. Shen, Zhongmin *Complete manifolds with nonnegative Ricci curvature and large volume growth*. Invent. Math. 125 (1996), no. 3, 393-404.
5. Sormani, Christina *Nonnegative Ricci curvature, small linear diameter growth and finite generation of fundamental groups*. J. Differential Geom. 54 (2000), no. 3, 547-559.
6. Wu, Bing Ye *On the fundamental group of Riemannian manifolds with nonnegative Ricci curvature*. Geom. Dedicata 162 (2013), 337-344.

SCHOOL OF MATHEMATICS, NORTHWEST UNIVERSITY, XI'AN 710127, CHINA  
E-mail address: wanj\_m@aliyun.com