QUANTIFIER ALTERNATION IN A CLASS OF RECURSIVELY DEFINED TREE PROPERTIES

MOUMANTI PODDER

ABSTRACT. Alternating quantifier depth is a natural measure of difficulty required to express first order logical sentences. We define a sequence of first order properties on rooted, locally finite trees in a recursive manner, and provide rigorous arguments for finding the alternating quantifier depth of each property in the sequence, using Ehrenfeucht-Fraïssé games.

1. INTRODUCTION

We study the alternating quantifier depths of a class of recursively defined first order properties on rooted, locally finite trees. A tree T is a connected graph on either a finite or an infinite vertex set that does not contain any cycle. We in particular shall consider trees as directed graphs, with clear distinction of parent and child between adjacent vertices. A rooted tree has a special vertex called the root, which we denote by R. A tree is called locally finite when each vertex of the tree has finite degree. We denote by V = V(T) the vertex set of the tree T. For a given rooted tree T and a vertex $v \in V(T)$, we let T(v) denote the subtree of T that consists of v and all its descendants.

We now define the first order (FO) language on rooted trees. This consists of the root R of the tree as a constant symbol, the equality of vertices (denoted x = y which means that the vertices x and y coincide), and the parent-child relation (denoted $\pi(y) = x$ which means that the vertex x is the parent of the vertex y). We denote vertices in the tree other than the root by letters such as x, y, z, \ldots usually in lower case. Sentences are finite and comprise Boolean connectives such as $\neg, \land, \lor, \Longrightarrow, \Leftrightarrow$ etc., and existential (denoted \exists) and universal (denoted \forall) quantification over vertices. We refer the reader to any of [2], [3] and [4] for detailed references on first order logic. The quantifier depth, abbreviated as q.d., of an FO sentence A is the minimum number of nested quantifiers required to express A, whereas the alternating quantifier depth of A, which we abbreviate henceforth as the a.q.d. of A, is the minimum number, the minimum being taken over all formulas that express A, of the maximum number of alternating nested quantifiers required to write each such formula. That is, a.q.d. gives us the minimum number of times we have to switch from existential to universal or universal to existential quantification in a nested sequence of quantifiers in order to express the sentence. Trivially, the a.q.d. of any FO sentence is always bounded above by its q.d. Purely existential or purely universal FO sentences are defined to have a.q.d. 0.

²⁰¹⁰ Mathematics Subject Classification. 05C57, 05C05, 05C75, 03C07, 03C13, 68Q19.

Key words and phrases. First order logic, alternating quantifier depth, recursive properties of rooted trees, Ehrenfeucht games.

The author was partially supported by the grant NSF DMS-1444084.

Example 1.1. Consider the FO sentence: there exists a vertex with precisely one child. This can be written as

$$\exists x \left[\exists y \left[\left[\pi(y) = x \right] \land \forall z \left[\pi(z) = x \implies z = y \right] \right] \right]. \tag{1.1}$$

The q.d. of this sentence is 3 whereas the a.q.d. is 1, because we have two existential followed by one universal quantifier in the nested sequence.

One of the classically studied family of questions by mathematical logicians comprise the model-theoretic results about the expressive power of logical languages. In general, given two languages A and B such that A is a subset of B, the task is to show that the two are not tautologically equal, i.e. there exists a sentence which is expressible in B but not in A. Quantifier alternation hierarchy is a natural tool for proving such results. We now discuss some literature on the a.g.d. of FO, and also touch briefly on a.g.d. of monadic second order (MSO) logic. [5] is a survey paper that discusses the usefulness of quantifier alternation hierarchy in membership algorithms – algorithms that are used to decide whether a given regular language of finite words is definable by a sentence from FO logic or not. They beautifully express the necessity to understand quantifier alternation hierarchy as a measure of the difficulty of defining a language – a language is considered complicated if many switches need to be made between blocks of existential quantifiers and blocks of universal quantifiers. [6] considers quantifier alternation hierarchy within FO language comprising two variables over finite words with linear order and binary successor predicate. They show that for a given regular language and any non-negative integer m, it is decidable whether the language is definable by an FO sentence of a.q.d. at most m. [7] also considers quantifier alternation hierarchy in FO logic on finite words – they prove that one can decide membership of a regular language to the levels of FO sentences that have a.q.d. 1 or FO sentences with a.q.d. 2 and beginning with an existential quantifier. [8] provides an effective characterization of tree languages that are definable by FO sentences with a.q.d. 1. [15] proves the strictness of FO quantifier alternation hierarchy over the class of finite labeled graphs – for each positive integer k, they exhibit a property of finite labeled directed graphs that is expressible as an FO sentence of a.q.d. k+1 but not as any FO sentence of a.q.d. k. [16] shows that FO quantifier alternation hierarchy is equivalent to dot-depth alternation hierarchy for FO formulas over word models with a total ordering on the alphabet rather than the successor relation on word positions. [13], [14] and [17] discuss the strictness of MSO quantifier alternation, i.e. sentences in prenex normal form having a prefix of k+1 many alternations of set quantifiers can describe strictly more graph properties than those having a prefix of only k many alternations of set quantifiers.

All the results we have been able to find in the literature pertaining to quantifier alternation hierarchy as an important tool to understand the expressive power of languages, are in the premise of graphs, trees, grids or pictures where each vertex or position is labeled by some element from a finite alphabet. In particular, in case of trees, some form of ordering is considered on the vertices, and each vertex is assigned a label from the pre-fixed, finite alphabet, and the unary predicate that specifies the label of a vertex is considered. We hope that our attempt to investigate a.q.d.'s of FO properties of rooted trees without the involvement of any alphabet will serve to begin a new direction of study with new definitions of FO and MSO languages on rooted, locally finite trees. Studies in this direction may reveal strictness, or lack thereof, of quantifier alternation hierarchy of the FO language, as we define it, on rooted trees. Continuing with this hope, we cite here [12], where the authors define the FO language on graphs using the binary relations of adjacency and equality of vertices (without the involvement of any alphabet). They show that the minimum number of quantifier alternations that an FO sentence A must have in order to fail to have a 0-1law on $G(n, n^{-\alpha})$ for infinitely many values of α is 3. We also hope that future work in this area would bring to light many more classes of recursively defined properties on rooted trees whose analysis may reveal more information on quantifier alternation hierarchy.

1.1. Organization of the paper: Our paper is organized in the following manner. In Section 2 we describe the class of recursively defined properties on rooted, locally finite trees that we examine. In Section 3 we describe the main tool we use to understand the a.q.d. of these properties – the Ehrenfeucht games, also known as the Ehrenfeucht-Fraüssé games. In Section 4 we describe the detailed inductive construction of the trees where the Ehrenfeucht games are played, and finally, in Section 5, we describe, also along an inductive argument, the winning strategy for Duplicator, that gives us the final conclusion.

2. Description of the problem

We define, for any rooted, locally finite tree T with root R, and any $x \in V(T)$, the property $P_0(x)$ which states that x has no child, which can be expressed as

$$P_0(x) = \forall y \neg [\pi(y) = x].$$
(2.1)

We now define the class of properties $P_i(x)$, for $i \in \mathbb{N}$ and any $x \in V(T)$, recursively as follows:

$$P_i(x) := \forall \ y \left[\pi(y) = x \implies \neg P_{i-1}(y) \right].$$

$$(2.2)$$

For example, $P_1(x)$ denotes the property that x has no child with no child. In particular, we define the property $\text{KEIN}_i = P_i(R)$ for every $i \in \mathbb{N}$. The aim of this paper is to show that the a.q.d. of KEIN_i is i for every $i \in \mathbb{N} \cup \{0\}$. It is immediate to see that KEIN_0 has a.q.d. 0, since it is a purely universal sentence. From our recursive definition (2.2), we can also see that the a.q.d. of KEIN_i is at least i. It remains to be shown that i is indeed the minimum number of alternations of nested quantifiers required to express the sentence KEIN_i for every $i \in \mathbb{N}$.

We mention here an analogous family of properties defined for graphs. On graphs, the corresponding FO language will consist of vertices denoted by x, y, z, \ldots etc. the equality of vertices (again, denoted x = y) and the adjacency of vertices (denoted $x \sim y$), the Boolean connectives and the existential and universal quantifiers over vertices. One can define, for any vertex x, the property $Q_0(x)$ that x has no neighbour, or, in other words, x being an isolated vertex, and set $N_0 = \exists x [Q_0(x)]$. Then one can define

$$Q_1(x) = \forall y [y \sim x \implies [\exists z [\neg [z = x] \land [z \sim y]]]], \qquad (2.3)$$

or, in other words, for every neighbour y of x, there exists some neighbour z of y which is distinct from x, or, in other words, no neighbour of x is a degree 1 vertex. We then define $N_1 = \exists x [Q_1(x)]$. Finally, we recursively define

$$Q_i(x) = \forall y [y \sim x \implies \neg [Q_{i-1}(y)]], \qquad (2.4)$$

and set $N_i = \exists x[Q_i(x)]$. Notice that although defined in a rather similar fashion to the tree properties KEIN_s, the properties N_s are harder to analyze. This is because of the following reason. As we shall see in the subsequent sections, our argument hinges upon the use of Ehrenfeucht games on two trees that we construct recursively on s, such that one of them satisfies KEIN_s and the other satisfies $\neg \text{KEIN}_s$. The attempt to do a similar recursive construction in the graph case fails, because there, the property $\neg N_s$ claims something much stronger than $\neg \text{KEIN}_s$ – it claims that a certain property fails to hold for *every* vertex in the graph. The study of the properties N_s for $s \in \mathbb{N}$ and their a.q.d. remains of keen interest to us in future research in this area. The comparison of such classes of properties on graphs and rooted trees may reveal significant difference between FO on graphs and FO on rooted trees as we have defined them.

3. The Ehrenfeucht game for alternating quantifiers

As mentioned above, the main objective of this paper is to show that KEIN_s has a.q.d. precisely equal to s for every $s \in \mathbb{N}$. The rigorous proof of this statement relies on a special version of the well-known Ehrenfeucht-Fraïssé games, which we henceforth refer to as simply the Ehrenfeucht games. For standard definition of these combinatorial games and their connection to mathematical logic, we refer the reader to any one of the references [1] and [3]. For the special version of the game we are about to make use of in this paper, we refer the reader to Definition 2.7, Definition 2.8 and Theorem 2.9 of [10], and a formal proof of the connection of this special version with a.q.d. of FO sentences in [11].

We first state here, in the premise of rooted trees, the version of the Ehrenfeucht games described in [10] and [11], that determines the maximum among the a.q.d.'s of all FO sentences that hold true in both the structures on which the game is being played. We first state here a few general rules and terminology which apply to all three of Definitions 3.1, 3.3 and 3.5. The game is played, for a given number of rounds, on two given rooted trees T_1 and T_2 , by two players known as Spoiler and Duplicator. Each round of the game consists of two parts: a *move* by Spoiler followed by a move by Duplicator. By a move, we mean the action by any player of choosing a vertex from one of the two trees. In each round, once Spoiler has made his selection of a vertex from one of the trees, Duplicator must make her selection of a vertex from the *other* tree. Thus, in every round, there is precisely one vertex chosen from T_1 and one from T_2 .

If in the *i*-th round, for some $1 \le i \le r-1$ where *r* is the total number of rounds, Spoiler makes his move on T_1 and in the (i + 1)-st round makes his move on T_2 , or vice versa, then we say that a *switch* has happened.

Definition 3.1. [Ehrenfeucht game with given maximum number of alternations] Given two rooted trees T_1 with root R_1 and T_2 with root R_2 , and two positive integers r and s with $r \ge s$, this game, denoted EHR $[T_1, T_2, s, r]$, consists of r many rounds. Spoiler is allowed to make his move in the first round on *any* of the two trees T_1 and T_2 , but throughout the game, he is allowed to make at most s many switches.

Let x_i be the vertex selected from T_1 and y_i that from T_2 in round i, for $1 \le i \le r$. We set $x_0 = R_1$ and $y_0 = R_2$. Duplicator wins the game if all of the following conditions hold: for all $i, j \in [r]$,

(Main 1) $\pi(x_j) = x_i \Leftrightarrow \pi(y_j) = y_i;$ (Main 2) $x_i = x_j \Leftrightarrow y_i = y_j,$

where for any positive integer n, we denote by [n] the set $\{0, 1, \ldots, n\}$.

The importance of this version of the Ehrenfeucht games is illustrated in the following theorem, which is Theorem 2.9 of [10] stated for rooted, locally finite trees.

Theorem 3.2. For any two rooted trees T_1 and T_2 , for positive integers r and s, Duplicator wins EHR $[T_1, T_2, s, r]$ if and only if for every FO sentence A of q.d. at most r and a.q.d. at most s, we have $T_1 \models A \Leftrightarrow T_2 \models A$.

Here, for any rooted tree T and any FO sentence A, the notation $T \models A$ implies that A holds in T. The conclusion of Theorem 3.2 is that, if Duplicator wins EHR $[T_1, T_2, s, r]$, then for every FO sentence A of q.d. at most r and a.q.d. at most s, either A holds for both T_1 and T_2 , or it holds for neither.

The version of the Ehrenfeucht games that we use is slightly different from that given in Definition 3.1. We show in Lemma 3.4 that if, on two given trees T_1 and T_2 , Duplicator wins the Ehrenfeucht game described in Definition 3.3 with sufficiently large values of the parameters concerned, then she also wins the Ehrenfeucht game described in Definition 3.1.

Definition 3.3. [Ehrenfeucht game with alternation after every k rounds for fixed k] Given two rooted trees T_1 with root R_1 and T_2 with root R_2 , and two positive integers k and s, this game, denoted $\text{EHR}_{\text{alt}}[T_1, T_2, s; k]$, consists of sk rounds. The sk many rounds are divided into s many batches of k rounds each. Spoiler, before the very first round, selects any one of T_1 and T_2 and in the first k rounds makes his moves on that tree, in the next k rounds he makes his moves on the other tree, and so on, i.e. he makes a switch after every jk-th round for every $1 \le j \le s - 1$ (in particular, there can be no switch possible if s = 1).

If x_i is the vertex selected from T_1 and y_i that from T_2 in round *i*, for $i \in [sk]$, setting $x_0 = R_1$ and $y_0 = R_2$, the winning conditions for Duplicator are (Main 1) and (Main 2), as described in Definition 3.1.

Lemma 3.4. For any two positive integers r and s, and two given rooted trees T_1 and T_2 , if Duplicator wins $\text{EHR}_{\text{alt}}[T_1, T_2, s + 1; r]$, then she also wins $\text{EHR}[T_1, T_2, s, r]$.

Proof. Suppose Spoiler makes t many switches during the game EHR $[T_1, T_2, s, r]$, where $t \leq s$. Without loss of generality, let us assume that Spoiler starts playing on T_1 . Let the first switch happen after the i_1 -th round, the second switch after the $(i_1 + i_2)$ -th round, and so on, where t, i_1, \ldots, i_t are entirely dependent on Spoiler's decision and are unknown to Duplicator a priori. Clearly, for each $1 \leq j \leq t$, we have $i_j \leq r$.

Having a winning strategy W for Duplicator for the game $\text{EHR}_{\text{alt}}[T_1, T_2, s + 1; r]$ means that, whatever sequence of (s + 1)r moves Spoiler plays according to the rules of the game, Duplicator has a sequence of (s + 1)r responses such that she can maintain all the winning conditions. Without loss of generality, suppose Spoiler starts playing $\text{EHR}_{\text{alt}}[T_1, T_2, s + 1; r]$ on T_1 , and let x_1, \ldots, x_r denote his moves in T_1 in the first r rounds, and let y_1, \ldots, y_r denote the corresponding responses of Duplicator in T_2 according to W; let y_{r+1}, \ldots, y_{2r} denote the moves made by Spoiler in T_2 in rounds $r + 1, \ldots, 2r$, and let x_{r+1}, \ldots, x_{2r} denote the corresponding responses of Duplicator in T_1 according to W, and so on.

Now, we construct the winning strategy for Duplicator for the game EHR $[T_1, T_2, s, r]$. As mentioned above, let Spoiler start playing on tree T_1 . Let a_1, \ldots, a_{i_1} denote the moves made by Spoiler in rounds $1, \ldots, i_1$ in tree T_1 . Setting $x_j = a_j$ for all $1 \leq j \leq i_1$, Duplicator replies with moves $b_j = y_j$ for all $1 \leq j \leq i_1$ where the y_j 's, as mentioned above, are chosen according to strategy W. Note that Duplicator can do this because $i_1 \leq r$. Next, for $i_1+1 \leq j \leq i_1+i_2$, Spoiler plays on T_2 and selects vertices $b_j, i_1+1 \leq j \leq i_1+i_2$. Duplicator, setting $y_{r+j} = b_{i_1+j}$ for all $1 \leq j \leq i_2$, selects vertices $a_{i_1+j} = x_{r+j}$, for $1 \leq j \leq i_2$, in T_1 according to winning strategy W. Once again, this is possible since $i_2 \leq r$. She continues to play like this until all r rounds are done.

Clearly, the winning conditions listed in Definition 3.1 now only need $\{(x_j, y_j), 1 \le j \le i_1\} \quad \cup \quad \{(x_j, y_j), r+1 \le j \le r+i_2\} \quad \cup$ hold for subset to a $\cdots \{ (x_j, y_j), (t-1)r + 1 \le j \le (t-1)r + i_t \} \cup \{ (x_j, y_j), tr + 1 \le j \le tr + (r - \sum_{\ell=1}^t i_\ell) \}$ of the set $\{(x_i, y_i), 1 \leq j \leq rs\}$, and we know that the winning conditions already hold for the bigger set because Duplicator chose her responses according to the winning strategy W. This shows that she wins $\text{EHR}[T_1, T_2, s, r]$.

We give here yet another version of the Ehrenfeucht games, which comes in handy in the description of winning strategies for Duplicator in Subsection 5.1.

Definition 3.5. Given rooted trees T_1 with root R_1 and T_2 with root R_2 , and positive integers $s, i_1, i_2, \ldots i_s$, this version of the Ehrenfeucht game, denoted $\text{EHR}_{\text{weak}}[T_1, T_2; i_1, \ldots i_s]$, consists of $i_1 + i_2 \ldots + i_s$ rounds. First, Spoiler chooses any of T_1 and T_2 , and makes his first i_1 moves on that tree, while Duplicator makes the corresponding i_1 moves on the other tree. Spoiler makes the first switch after the i_1 -th round, the second switch after the $(i_1 + i_2)$ -th round, \ldots , and finally, the (s-1)-st switch after the $(i_1 + \cdots + i_{s-1})$ -th round.

As before, if x_i is the vertex selected from T_1 and y_i that from from T_2 in round *i*, for $1 \leq i \leq \sum_{j=1}^{s} i_j$, then, setting $x_0 = R_1$ and $y_0 = R_2$, Duplicator wins the game if (Main 1) and (Main 2) hold.

Notice that this is weaker than the game described in Definition 3.1 because here the values i_1, \ldots, i_s are known to both Spoiler and Duplicator before the start of the game.

Lemma 3.6. For any two positive integers s and k and rooted trees T_1 and T_2 , if Duplicator wins $\text{EHR}_{\text{alt}}[T_1, T_2, s; k]$, then she also wins $\text{EHR}_{\text{weak}}[T_1, T_2; i_1, \ldots i_s]$ for any $1 \leq i_1, \ldots i_s \leq k$.

Proof. The proof of this lemma is very similar to that of Lemma 3.4, and is in fact simpler, and the details are therefore omitted. \Box

We now describe the way we put the Ehrenfeucht game described in Definition 3.3 to use in proving that KEIN_s has a.q.d. s for each $s \in \mathbb{N}$. For any positive integer k, we construct two rooted trees T_1 and T_2 (these trees will obviously depend on k and s), such that $T_1 \models \text{KEIN}_s$ and $T_2 \models \neg \text{KEIN}_s$, and Duplicator wins $\text{EHR}_{\text{alt}}[T_1, T_2, s; k]$. From this, we can draw the following conclusion: using Lemma 3.4, we know that Duplicator wins $\text{EHR}[T_1, T_2, s - 1, k]$, and hence, from Theorem 3.2, we can tell that for every FO sentence A that has q.d. at most k and a.q.d. at most s - 1, either A holds for both T_1 and T_2 , or $\neg A$ holds for both

 $\mathbf{6}$



Each such structure represents m childless children.

FIGURE 1. $T_1^{(1,k,m)}$ and $T_2^{(1,k,m)}$

 T_1 and T_2 . But notice that KEIN_s holds for T_1 whereas \neg KEIN_s holds for T_2 . This clearly means that KEIN_s either fails to have q.d. at most k, or it fails to have a.q.d. at most s - 1. Since we are able to construct T_1 and T_2 and provide a winning strategy for Duplicator for arbitrary k, clearly it is the upper bound on the a.q.d. of KEIN_s that fails. This shows that KEIN_s must have a.q.d. at least s. Since we have noted before that its a.q.d. is at most s, it must be precisely equal to s. This completes the proof of our main result.

4. Construction of the trees

In this section, we describe the construction of the trees T_1 and T_2 depending on the given parameters s and k. We take into account an additional parameter m, whose role becomes clear from the construction. The constructions are described inductively on s, starting with the base case of s = 1, and arbitrary k, which we now describe.

For any positive integer $m \geq k$, we construct the rooted trees $T_1^{(1,k,m)}$ and $T_2^{(1,k,m)}$ as follows:

- (i) In $T_1^{(1,k,m)}$, the root R_1 has m + 1 children $u_1, \ldots u_{m+1}$, and each of them has m childless children of its own.
- (ii) In $T_2^{(1,k,m)}$, the root R_2 has m + 1 children $v_1, \ldots v_{m+1}$. Each of $v_1, \ldots v_m$ has m childless children of its own; v_{m+1} has no child.

An illustration is given in Figure 1.

Suppose we know how to construct the trees $T_1^{(s',k,m)}$ and $T_2^{(s',k,m)}$ for all $s' \leq s$ where s is some positive integer, for arbitrary k, and for all $m \geq s'k$, such that $T_1^{(s',k,m)} \models$ KEIN_{s'} and $T_2^{(s',k,m)} \models \neg$ KEIN_{s'}, and Duplicator wins EHR_{alt} $\left[T_1^{(s',k,m)}, T_2^{(s',k,m)}, s'; k\right]$. Now, fixing an arbitrary k, we construct $T_1^{(s+1,k,m)}$ and $T_2^{(s+1,k,m)}$ for all $m \geq (s+1)k$, such that $T_1^{(s+1,k,m)} \models$ KEIN_{s+1} whereas $T_2^{(s+1,k,m)} \models \neg$ KEIN_{s+1}, and Duplicator wins EHR_{alt} $\left[T_1^{(s+1,k,m)}, T_2^{(s+1,k,m)}, s+1; k\right]$.

Note that we make use of $T_1^{(s,k,m)}$ and $T_2^{(s,k,m)}$ in constructing $T_1^{(s+1,k,m)}$ and $T_2^{(s+1,k,m)}$ for $m \ge (s+1)k$, and since this last condition ensures that m > sk, hence it makes sense to talk about $T_1^{(s,k,m)}$ and $T_2^{(s,k,m)}$. The inductive construction is as follows:

- (i) In $T_1^{(s+1,k,m)}$, the root R_1 has m+1 children $u_1, \ldots u_{m+1}$, and from each of them hangs a copy of $T_2^{(s,k,m)}$.
- (ii) In $T_2^{(s+1,k,m)}$, the root R_2 has m+1 children, $v_1, \ldots v_{m+1}$. From each of $v_1, \ldots v_m$, hangs a copy of $T_2^{(s,k,m)}$, and from v_{m+1} hangs a copy of $T_1^{(s,k,m)}$.

An illustration is given in Figure 2.



FIGURE 2. Trees $T_1^{(s+1,k,m)}$ and $T_2^{(s+1,k,m)}$

We make sure here that indeed we have $T_1^{(s+1,k,m)} \models \text{KEIN}_{s+1}$ and $T_2^{(s+1,k,m)} \models \neg \text{KEIN}_{s+1}$. Recall from (2.2) that a rooted tree T satisfies KEIN_{s+1} if for every child x of the root R, the property $\neg P_s(x)$ holds. By induction hypothesis, $T_2^{(s,k,m)} \models \neg \text{KEIN}_s$. Since for each child u_i of the root in $T_1^{(s+1,k,m)}$, we have

$$T_1^{(s+1,k,m)}(u_i) \cong T_2^{(s,k,m)}$$

(where the notation \cong indicates a tree ismorphism map which maps the root of one tree to the root of the other), hence $\neg P_s(u_i)$ holds for every $1 \le i \le m+1$. On the other hand, also by induction hypothesis, $T_1^{(s,k,m)} \models \text{KEIN}_s$. In $T_2^{(s+1,k,m)}$, the root has one child v_{m+1} such that

$$T_2^{(s+1,k,m)}(v_{m+1}) \cong T_1^{(s,k,m)}$$

and hence $P_s(v_{m+1})$ holds. This tells us that $\neg \text{KEIN}_{s+1}$ must hold for $T_2^{(s+1,k,m)}$.

5. Winning strategy for Duplicator for arbitrary s

We first describe Duplicator's winning strategy for s = 1. For s = 1, no alternation is allowed in EHR_{alt} $[T_1, T_2, s; k]$, i.e. either Spoiler plays the entire game on T_1 while Duplicator answers on T_2 , or Spoiler plays the entire game on T_2 and Duplicator answers on T_1 . Also, there are a total of k many rounds now. We now describe the winning strategy for Duplicator in either scenario.

We introduce here a terminology that we shall use in our exposition of the winning strategy for Duplicator henceforth.

Definition 5.1. Suppose *i* rounds of the game $\text{EHR}_{\text{alt}}[T_1, T_2, s; k]$ have been played. For any vertex $u \in T_1$, we call *u* free up to round *i* if no x_j , $1 \leq j \leq i$, has been selected from $T_1(u)$. Similarly, for any vertex $v \in T_2$, we call *v* free up to round *i* if no y_j , $1 \leq j \leq i$, has been selected from $T_2(v)$.

QUANTIFIER ALTERNATION IN A CLASS OF RECURSIVELY DEFINED TREE PROPERTIES 9

First, consider the case where Spoiler plays on tree $T_2^{(1,k,m)}$. Throughout the game, Duplicator maintains the following conditions, and we show that she is able to maintain them using an inductive argument. Suppose *i* rounds of the game have been played. Then, for $1 \le j \le i$,

- (i) if Spoiler selects $y_j = v_\ell$ for some $1 \le \ell \le m+1$, then Duplicator selects $x_j = u_\ell$;
- (ii) if Spoiler selects y_j to be a child of the vertex v_ℓ , for some $1 \le \ell \le m$, then Duplicator selects x_j to be a child of u_ℓ , making sure that (Main 2) is maintained (note that these two conditions together imply that $x_j = R_1$ if and only if $y_j = R_2$).

Suppose Duplicator has been able to maintain these conditions up to and including round *i*. Now suppose Spoiler selects y_{i+1} to be equal to some v_{ℓ} for $1 \leq \ell \leq m+1$. Then Duplicator sets $x_{i+1} = u_{\ell}$. If Spoiler selects y_{i+1} to be a child of v_{ℓ} for some $1 \leq \ell \leq m$, and this child was not chosen before, then Duplicator selects x_{i+1} to be a child of u_{ℓ} that has not been chosen before, and notice that such a child she can always find because each u_{ℓ} for $1 \leq \ell \leq m$ has *m* children and $m \geq k$, and there are only *k* many rounds. If Spoiler selects y_{i+1} to be a child of v_{ℓ} that was already chosen before, say in the *j*-th round for some $j \leq i$, then Duplicator, to maintain (Main 2), simply sets $x_{i+1} = x_j$. It is straightforward to see that these choices do guarantee the satisfaction of both (Main 1) and (Main 2) at the end of the game.

Now suppose Spoiler plays on tree $T_1^{(1,k,m)}$. Once again, Duplicator maintains the following conditions throughout the game, which we prove via inductive arguments that she is able to. Suppose *i* rounds of the game have been played. Then for all $j, j' \in [i]$,

- (i) for $j \neq j'$, $x_j = x_{j'}$ if and only if $y_j = y_{j'}$ (this implies that $x_j = R_1$ if and only if $y_j = R_2$);
- (ii) $x_j \in \{u_1, \dots, u_{m+1}\}$ if and only if $y_j \in \{v_1, \dots, v_m\}$;
- (iii) x_j is a child of u_ℓ for some $1 \le \ell \le m+1$ if and only if y_j is a child of $v_{\ell'}$ for some $1 \le \ell' \le m$;
- (iv) for $j \neq j'$, $\pi(x_{j'}) = x_j$ if and only if $\pi(y_{j'}) = y_j$;
- (v) $x_j = u_\ell$ for some $1 \le \ell \le m + 1$ such that u_ℓ has been free up to round j 1 if and only if $y_j = v_{\ell'}$ for some $1 \le \ell' \le m$ such that $v_{\ell'}$ has been free up to round j - 1;
- (vi) for $j \neq j'$, x_j and $x_{j'}$ are siblings, i.e. they share a common parent, if and only if y_j and $y_{j'}$ are siblings.

Suppose Duplicator has been able to maintain all of these conditions up to and including round *i*, for some $i \leq k - 1$. Of course, if Spoiler selects $x_{i+1} = x_j$ for some $j \leq i$, then Duplicator sets $y_{i+1} = y_j$. So, for the rest of the inductive argument, assume that x_{i+1} is distinct from all previously chosen vertices.

Suppose Spoiler selects x_{i+1} to be some u_{ℓ} for $1 \leq \ell \leq m+1$ such that u_{ℓ} has been free up to round *i*. Note that $m \geq k$ and there are *k* rounds in total, hence there has to be at least one vertex $v_{\ell'}$ for some $1 \leq \ell' \leq m$ free up to round *i*. Hence Duplicator finds such a $v_{\ell'}$ and sets $y_{i+1} = v_{\ell'}$. Suppose Spoiler selects x_{i+1} to be u_{ℓ} such that there exists $j \leq i$ with x_j a child of u_{ℓ} ; by induction hypothesis, we know that y_j is a child of $v_{\ell'}$ for some $1 \leq \ell' \leq m$, and we set $y_{i+1} = v_{\ell'}$.

If Spoiler selects x_{i+1} to be a child of some vertex u_{ℓ} for $1 \leq \ell \leq m+1$ such that u_{ℓ} has been free up to round *i*, then Duplicator once again finds $v_{\ell'}$ for some $1 \leq \ell' \leq m$ that has

been free up to round *i* (possible for the same reason as argued above), and then selects y_{i+1} to be any child of $v_{\ell'}$. Suppose Spoiler selects x_{i+1} to be a child of u_{ℓ} , for some $1 \leq \ell \leq m+1$, such that for some $j \leq i$, we have $x_j = u_{\ell}$. Then by induction hypothesis, we know that y_j must equal $v_{\ell'}$ for some $1 \leq \ell' \leq m$. Duplicator selects a child of $v_{\ell'}$, distinct from all previously chosen vertices, as y_{i+1} . As there are *m* children of $v_{\ell'}$ and a total of *k* rounds with $m \geq k$, hence she is able to make such a selection. Suppose Spoiler selects x_{i+1} to be a child of u_{ℓ} for some $1 \leq \ell \leq m+1$ such that for some $j \leq i$, the vertex x_j is also a child of u_{ℓ} . Again by induction hypothesis we know that y_j is a child of $v_{\ell'}$ for some $1 \leq \ell \leq m+1$ such that y_j is a child of $v_{\ell'}$ for some $1 \leq \ell' \leq m$, and Duplicator sets y_{i+1} to be a child of $v_{\ell'}$ that is different from y_j as well as any other previously chosen vertex. Note that, since each u_{ℓ} and each $v_{\ell'}$ has *m* children and $m \geq k$, where *k* is the total number of rounds, there is always a choice of y_{i+1} as a child of $v_{\ell'}$ which does not coincide with any previously chosen vertex.

This exhausts all possible moves by Spoiler and we have shown that Duplicator is able to respond in each case such that all the conditions mentioned above are maintained.

We now come to the winning strategy for Duplicator on $\operatorname{EHR}_{\operatorname{alt}}\left[T_1^{(s,k,m)}, T_2^{(s,k,m)}, s; k\right]$ for arbitrary k and $m \geq sk$, and this is described as an inductive strategy where the induction happens on s. Thus, we assume that we already have a winning strategy for Duplicator for the game $\operatorname{EHR}_{\operatorname{alt}}\left[T_1^{(s,k,m)}, T_2^{(s,k,m)}, s; k\right]$, and then devise a strategy for her for the game $\operatorname{EHR}_{\operatorname{alt}}\left[T_1^{(s+1,k,m)}, T_2^{(s+1,k,m)}, s+1; k\right]$ for the same k and $m \geq (s+1)k$. The analysis is split into two parts, according to which tree Spoiler starts playing on, and these are described separately, in detail, in Subsections 5.1 and 5.2.

5.1. Spoiler starts playing on $T_2^{(s+1,k,m)}$. Suppose Spoiler starts playing the game $\operatorname{EHR}_{\operatorname{alt}}\left[T_1^{(s+1,k,m)}, T_2^{(s+1,k,m)}, s+1; k\right]$ on $T_2^{(s+1,k,m)}$. Recall, from the inductive construction of the trees in Section 4, that $T_1^{(s,k,m)}$ and $T_2^{(s,k,m)}$ are as illustrated in Figure 3.



FIGURE 3. Trees $T_1^{(s,k,m)}$ and $T_2^{(s,k,m)}$

Induction hypothesis 5.2. For every $1 \le t \le m+1$ and $1 \le t' \le m$, fix a tree isomorphism $\varphi_{t,t'}^{(s)}: T_1^{(s,k,m)}(u_t) \to T_2^{(s,k,m)}(v_{t'})$ (this is possible as both are copies of $T_2^{(s-1,k,m)}$), such that $\varphi_{t,t'}^{(s)}(u_t) = v_{t'}$. Fix ℓ pairs of vertices $(x_1, y_1), \ldots, (x_\ell, y_\ell)$ in $T_1^{(s,k,m)} \times T_2^{(s,k,m)}$ such that $\ell \le k$ and they satisfy the following conditions:

(IH1) for each $1 \le i \le \ell$, we have $x_i \in \bigcup_{t=1}^{m+1} T_1^{(s,k,m)}(u_t)$ and $y_i \in \bigcup_{t'=1}^m T_2^{(s,k,m)}(v_{t'})$.

- (IH2) For $1 \le i \ne j \le \ell$, if $x_i, x_j \in T_1^{(s,k,m)}(u_t)$ for some $1 \le t \le m+1$, then we can find some $1 \le t' \le m$ such that $y_i, y_j \in T_2^{(s,k,m)}(v_{t'})$. The converse also holds, i.e. if for $1 \leq i \neq j \leq \ell$, if $y_i, y_j \in T_2^{(s,k,m)}(v_{t'})$ for some $1 \leq t' \leq m$, then we can find some $1 \le t \le m+1 \text{ such that } x_i, x_j \in T_1^{(s,k,m)}(u_t).$ (IH3) For every $1 \le i \le \ell$, if $x_i \in T_1^{(s,k,m)}(u_t)$ and $y_i \in T_2^{(s,k,m)}(v_{t'})$ for some $1 \le t \le m+1$
- and $1 \le t' \le m$, then $y_i = \varphi_{t,t'}^{(s)}(x_i)$.

As in Definitions 3.1 and 3.3, we set $x_0 = R_1$, the root of $T_1^{(s,k,m)}$, and $y_0 = R_2$, the root of $T_2^{(s,k,m)}$. Then Duplicator will be able to win EHR_{alt} $\left[T_1^{(s,k,m)}, T_2^{(s,k,m)}, s; k\right]$, where Spoiler starts playing on $T_2^{(s,k,m)}$, with the pairs $(x_0, y_0), \ldots (x_{\ell}, y_{\ell})$ considered as designated pairs.

Remark 5.3. It is helpful to imagine designated pairs of vertices as pairs chosen before the game has even begun. For example, in Induction hypothesis 5.2, one could visualize the designated pairs $(x_0, y_0), \ldots, (x_\ell, y_\ell)$ as pairs already selected / provided to the players from $\ell + 1$ many rounds that occurred before the actual game, and then the game $\operatorname{EHR}_{\operatorname{alt}}\left[T_{1}^{(s,k,m)}, T_{2}^{(s,k,m)}, s; k\right]$ of sk many rounds begins, which Duplicator is able to win, taking into account these $(\ell+1)$ many pairs as well, if the conditions given in Induction hypothesis 5.2 hold. In other words, suppose in EHR_{alt} $\left[T_1^{(s,k,m)}, T_2^{(s,k,m)}, s; k\right]$, the pair selected in round *i* from $T_1^{(s,k,m)} \times T_2^{(s,k,m)}$ is denoted by $(x_{i+\ell}, y_{i+\ell})$, for all $1 \leq i \leq sk$. Then Duplicator has to maintain *all* of the following conditions to win EHR_{alt} $\left[T_1^{(s,k,m)}, T_2^{(s,k,m)}, s; k\right]$ with $(x_0, y_0), \ldots, (x_\ell, y_\ell)$ as designated pairs:

(i) $x_i = \pi(x_j) \Leftrightarrow y_i = \pi(y_j)$ for all $0 \le i \ne j \le sk + \ell$; (ii) $x_i = x_j \Leftrightarrow y_i = y_j$ for all $0 \le i \ne j \le sk + \ell$.

Armed with Induction hypothesis 5.2, we now prove that the corresponding claim holds on $T_1^{(s+1,k,m)}$ and $T_2^{(s+1,k,m)}$ where $m \ge (s+1)k$. First, we need to fix $\ell \le k$ designated pairs that satisfy conditions analogous to ((IH1)) through ((IH3)). So, referring to Figure 2, we fix, for every $1 \le t \le m+1$ and every $1 \le t' \le m$, an isomorphism $\varphi_{t,t'}^{(s+1)}: T_1^{(s+1,k,m)}(u_t) \to T_2^{(s+1,k,m)}(v_{t'})$ (again possible because both $T_1^{(s+1,k,m)}$ and $T_2^{(s+1,k,m)}(v_{t'})$ are both copies of $T_2^{(s,k,m)}$; next, we fix an arbitrary collection of pairs $(x_1, y_1), \ldots, (x_\ell, y_\ell)$ for any non-negative integer $\ell \leq k$ (if $\ell = 0$, then we choose only the roots R_1 and R_2 as x_0 and y_0 respectively), such that the following hold:

- (C1) for each $1 \le i \le \ell$, we have $x_i \in \bigcup_{t=1}^{m+1} T_1^{(s+1,k,m)}(u_t)$ and $y_i \in \bigcup_{t'=1}^m T_2^{(s+1,k,m)}(v_{t'})$. (C2) For $1 \le i \ne j \le \ell$, if $x_i, x_j \in T_1^{(s+1,k,m)}(u_t)$ for some $1 \le t \le m+1$, then we can find some $1 \le t' \le m$ such that $y_i, y_j \in T_2^{(s+1,k,m)}(v_{t'})$. The converse also holds, i.e. if for $1 \le i \ne j \le \ell$, if $y_i, y_j \in T_2^{(s+1,k,m)}(v_{t'})$ for some $1 \le t' \le m$, then we can find some $1 \le t \le m+1$ such that $x_i, x_j \in T_1^{(s+1,k,m)}(u_t)$. (C3) For every $1 \le i \le \ell$, if $x_i \in T_1^{(s+1,k,m)}(u_t)$ and $y_i \in T_2^{(s+1,k,m)}(v_{t'})$, then $y_i =$
- $\varphi_{t.t'}^{(s+1)}(x_i).$

We now provide a strategy for Duplicator to win $\text{EHR}_{\text{alt}}\left[T_1^{(s+1,k,m)}, T_2^{(s+1,k,m)}, s+1; k\right]$ with $(x_0, y_0), (x_1, y_1) \dots (x_\ell, y_\ell)$ as designated vertices. Recall again that $x_0 = R_1$, the root of $T_1^{(s+1,k,m)}$, and $y_0 = R_2$, the root of $T_2^{(s+1,k,m)}$.

Assumption 5.4. Since $\ell \leq k$ and $m \geq (s+1)k$ with $s \geq 2$, there exists at least one $1 \leq i_0 \leq m+1$ such that no $x_j, 1 \leq j \leq \ell$, has been chosen from $T_1^{(s+1,k,m)}(u_{i_0})$. Because for each $1 \leq i \leq m+1$, the subtree $T_1^{(s+1,k,m)}(u_i)$ is a copy of $T_2^{(s,k,m)}$, we can therefore assume without loss of generality that $i_0 = m+1$.

Assumption 5.4 is made throughout Subsection 5.1.

Let us now take a closer look at the trees $T_1^{(s+1,k,m)}$ and $T_2^{(s+1,k,m)}$, especially in more detail the subtrees $T_1^{(s+1,k,m)}(u_{m+1})$ and $T_2^{(s+1,k,m)}(v_{m+1})$. These are illustrated in Figures (4) and (5).



FIGURE 4. $T_1^{(s+1,k,m)}$ with detailed view of $T_1^{(s+1,k,m)}(u_{m+1})$



FIGURE 5. $T_2^{(s+1,k,m)}$ with detailed view of $T_2^{(s+1,k,m)}(v_{m+1})$

Let $(x_{i+\ell}, y_{i+\ell})$ be the pair selected from $T_1^{(s+1,k,m)} \times T_2^{(s+1,k,m)}$ in round *i* for $1 \le i \le (s+1)k$. As is our convention for Subsection 5.1, Spoiler plays the first *k* rounds on $T_2^{(s+1,k,m)}$. In the following paragraph, we state some conditions Duplicator maintains on the configuration $\{(x_0, y_0), \ldots, (x_\ell, y_\ell)\} \cup \{(x_{i+\ell}, y_{i+\ell}) : 1 \le i \le k\}$ (i.e. on the pairs of vertices resulting from the first *k* rounds, along with the designated pairs). And we prove that she can indeed maintain these conditions by using an inductive argument (within the first *k* rounds).

QUANTIFIER ALTERNATION IN A CLASS OF RECURSIVELY DEFINED TREE PROPERTIES 13

First, Duplicator fixes any tree isomorphism $\varphi_{t,t'}^{(s)}: T_1^{(s+1,k,m)}(u_{m+1,t}) \to T_2^{(s+1,k,m)}(v_{m+1,t'})$, with $\varphi_{t,t'}^{(s)}(u_{m+1,t}) = v_{m+1,t'}$, for all $1 \leq t \leq m$ and $1 \leq t' \leq m+1$. This is possible as each $T_1^{(s+1,k,m)}(u_{m+1,t}), 1 \leq t \leq m$, as well as each $T_2^{(s+1,k,m)}(v_{m+1,t'}), 1 \leq t' \leq m+1$, is a copy of $T_2^{(s-1,k,m)}$. Suppose $p \leq k-1$ rounds of the game have been played. The conditions on the configuration $\{(x_0, y_0), \ldots, (x_\ell, y_\ell)\} \cup \{(x_{\ell+1}, y_{\ell+1}), \ldots, (x_{\ell+p}, y_{\ell+p})\}$ are as follows:

(A1) $x_{i+\ell} = R_1 \Leftrightarrow y_{i+\ell} = R_2$. If $x_{i+\ell} \in T_1^{(s+1,k,m)}(u_t)$ for some $1 \leq t \leq m$, then $y_{i+\ell} \in T_2^{(s+1,k,m)}(v_{t'})$ for some $1 \leq t' \leq m$, and vice versa; moreover, in this case, $y_{i+\ell} = \varphi_{t,t'}^{s+1}(x_{i+\ell})$.

Note that if $x_{i+\ell} \in T_1^{(s+1,k,m)}(u_{m+1})$, then actually $x_{i+\ell} \in \bigcup_{t=1}^m \in T_1^{(s+1,k,m)}(u_{m+1,t}) \bigcup \{u_{m+1}\}$. If $x_{i+\ell} = u_{m+1}$ then $y_{i+\ell} = v_{m+1}$, and vice versa. We have $x_{i+\ell} \in T_1^{(s+1,k,m)}(u_{m+1,t})$ for some $1 \leq t \leq m$ if and only if $y_{i+\ell} \in T_2^{(s+1,k,m)}(u_{m+1,t'})$ for some $1 \leq t' \leq m+1$, and moreover, in this case, $y_{i+\ell} = \varphi_{t,t'}^{(s)}(x_{i+\ell})$.

(A2) Suppose we have $x_{i+\ell}, x_j \in T_1^{(s+1,k,m)}(u_t)$ for some $1 \le i \le p$, some $1 \le j \le p+\ell$, and some $1 \le t \le m$. Then there exists some $1 \le t' \le m$ such that $y_{i+\ell}, y_j \in T_2^{(s+1,k,m)}(v_{t'})$.

The converse of this statement is true as well. That is, if for some $1 \leq i \leq p$, some $1 \leq j \leq p + \ell$, and some $1 \leq t' \leq m$, we have $y_{i+\ell}, y_j \in T_2^{(s+1,k,m)}(v_{t'})$, then there exists some $1 \leq t' \leq m$ such that $x_{i+\ell}, x_j \in T_1^{(s+1,k,m)}(u_t)$.

(A3) Suppose for $1 \leq i \neq j \leq p$ we have $x_{i+\ell}, x_{j+\ell} \in T_1^{(s+1,k,m)}(u_{m+1,t})$ for some $1 \leq t \leq m$. Then there exists $1 \leq t' \leq m+1$ such that $y_{i+\ell}, y_{j+\ell} \in T_2^{(s+1,k,m)}(v_{m+1,t'})$. The converse of this statement is also true, i.e. if for $1 \leq i \neq j \leq p$, we have $y_{i+\ell}, y_{j+\ell} \in T_2^{(s+1,k,m)}(v_{m+1,t'})$ for some $1 \leq t' \leq m+1$, then there exists $1 \leq t \leq m$ such that $x_{i+\ell}, x_{j+\ell} \in T_1^{(s+1,k,m)}(u_{m+1,t})$.

Suppose Duplicator has been able to maintain all these conditions up to and including round p. Now, Spoiler chooses $y_{\ell+p+1}$ from $T_2^{(s+1,k,m)}$. Here is how Duplicator replies:

- (i) If $y_{p+\ell+1} = R_2$, the root of $T_2^{(s+1,k,m)}$, then Duplicator chooses $x_{p+\ell+1} = R_1$, the root of $T_1^{(s+1,k,m)}$.
- (ii) Suppose $y_{p+\ell+1} \in \bigcup_{t'=1}^{m} T_2^{(s+1,k,m)}(v_{t'})$. Then there are a few possible cases, as follows: (a) Suppose there exists $1 \leq j \leq \ell + p$ such that y_j and $y_{p+\ell+1}$ both belong to the same $T_2^{(s+1,k,m)}(v_{t'})$ for some $1 \leq t' \leq m$. Then Duplicator finds t such that $x_j \in T_1^{(s+1,k,m)}(u_t)$. Note that if $1 \leq j \leq \ell$, then $1 \leq t \leq m$ because of Assumption 5.4), and if $\ell + 1 \leq j \leq p + \ell$, then $1 \leq t \leq m$ because of induction hypothesis (A1). Then Duplicator sets $x_{p+\ell+1} = \left\{\varphi_{t,t'}^{(s+1)}\right\}^{-1}(y_{p+\ell+1})$.
 - (b) Suppose $y_{p+\ell+1} \in T_2^{(s+1,k,m)}(v_{t'})$ for some $1 \le t' \le m$, such that $v_{t'}$ is free up to round p, i.e. $y_j \notin T_2^{(s+1,k,m)}(v_{t'})$ for all $1 \le j \le \ell + p$ (recall Definition 5.1). Then Duplicator finds a $1 \le t \le m$ such that u_t is free up to round p. She

can always find such a t as $m \ge (s+1)k > 2k > p+\ell$. She then sets $x_{p+\ell+1} = \left\{\varphi_{t,t'}^{(s+1)}\right\}^{-1} (y_{p+\ell+1}).$

- (iii) Now suppose $y_{p+\ell+1} \in T_2^{(s+1,k,m)}(v_{m+1})$. Again, there are a few possible cases:
 - (a) If $y_{p+\ell+1} = v_{m+1}$, then Duplicator sets $x_{p+\ell+1} = u_{m+1}$.
 - (b) If $y_{p+\ell+1} \in T_2^{(s+1,k,m)}(v_{m+1,t'})$ for some $1 \le t' \le m+1$ such that there exists some $1 \le i \le p$ with $y_{i+\ell} \in T_2^{(s+1,k,m)}(v_{m+1,t'})$, then Duplicator finds the t (where $1 \le t \le m$ by induction hypothesis (A1)) such that $x_{i+\ell} \in T_1^{(s+1,k,m)}(u_{m+1,t})$. Then she sets $x_{p+\ell+1} = \left\{\varphi_{t,t'}^{(s)}\right\}^{-1}(y_{p+\ell+1})$.
 - Then she sets $x_{p+\ell+1} = \left\{\varphi_{t,t'}^{(s)}\right\}^{-1} (y_{p+\ell+1}).$ (c) If $y_{p+\ell+1} \in T_2^{(s+1,k,m)}(v_{m+1,t'})$ for some $1 \leq t' \leq m+1$ such that $y_{i+\ell} \notin T_2^{(s+1,k,m)}(v_{m+1,t'})$ for all $1 \leq i \leq p$, then Duplicator finds a $1 \leq t \leq m$ such that $u_{m+1,t}$ is free up to round p (again, possible since $m \geq (s+1)k > 2k > p+\ell$) and sets $x_{p+\ell+1} = \left\{\varphi_{t,t'}^{(s)}\right\}^{-1} (y_{p+\ell+1}).$

Let us focus now on $T_2^{(s+1,k,m)}(v_{m+1})$ and $T_1^{(s+1,k,m)}(u_{m+1})$ only. Suppose the only pairs selected up to round k that are in $T_1^{(s+1,k,m)}(u_{m+1}) \times T_2^{(s+1,k,m)}(v_{m+1})$ are $(x_{i_1+\ell}, y_{i_1+\ell}), \dots, (x_{i_r+\ell}, y_{i_r+\ell})$. Now, $T_1^{(s+1,k,m)}(u_{m+1})$ is a copy of $T_2^{(s,k,m)}$ and $T_2^{(s+1,k,m)}(v_{m+1})$ is a copy of $T_1^{(s,k,m)}$. From (A1) and (A3), we can see that the pairs $(y_{i_1+\ell}, x_{i_1+\ell}), \dots, (y_{i_r+\ell}, x_{i_r+\ell})$ satisfy Conditions (IH1) through (IH3).

Conclusion 5.5. So Duplicator, by Induction hypothesis 5.2, will win $\operatorname{EHR}_{\operatorname{alt}}[T_2^{(s+1,k,m)}(v_{m+1}), T_1^{(s+1,k,m)}(u_{m+1}), s; k]$ with designated pairs $(y_{i_1+\ell}, x_{i_1+\ell}), \ldots, (y_{i_r+\ell}, x_{i_r+\ell})$ (notice the deliberate writing of $T_2^{(s+1,k,m)}(v_{m+1})$ before $T_1^{(s+1,k,m)}(u_{m+1})$).

Now consider the subsequent game of remaining sk rounds. We call this the second part of the game. For this part of the game, we really can split the tree $T_1^{(s+1,k,m)}$ into

$$\{R_1\} \cup S_1 \cup T_1^{(s+1,k,m)}(u_{m+1}), \text{ where } S_1 = \left\{\bigcup_{t=1}^m T_1^{(s+1,k,m)}(u_t)\right\},$$
(5.1)

and the tree $T_2^{(s+1,k,m)}$ into

$$\{R_2\} \cup S_2 \cup T_2^{(s+1,k,m)}(v_{m+1}), \text{ where } S_2 = \left\{\bigcup_{t'=1}^m T_1^{(s+1,k,m)}(v_{t'})\right\}.$$
(5.2)

Suppose p rounds of the game have been played, with $k . Duplicator maintains the following conditions on the configuration <math>\{(x_i, y_i) : 0 \le i \le p + \ell\}$:

- (B1) For $0 \le i \le p$, we have $x_i = R_1 \Leftrightarrow y_i = R_2$.
- (B2) For $0 \le i \le p$, we have $x_i \in S_1 \Leftrightarrow y_i = S_2$. In this case, if $x_i \in T_1^{(s+1,k,m)}(u_t)$ for some $1 \le t \le m$ and $y_i \in T_2^{(s+1,k,m)}(v_{t'})$ for some $1 \le t' \le m$, then $y_i = \varphi_{t,t'}^{(s+1)}(x_i)$.
- (B3) For $0 \le i, i' \le p$, if $x_i, x_{i'} \in T_1^{(s+1,k,m)}(u_t)$ for some $1 \le t \le m$, then there exists $1 \le t' \le m$ with $y_i, y_{i'} \in T_2^{(s+1,k,m)}(v_{t'})$.

QUANTIFIER ALTERNATION IN A CLASS OF RECURSIVELY DEFINED TREE PROPERTIES 15(B4) For $0 \le j \le p$, we have $x_j \in T_1^{(s+1,k,m)}(u_{m+1}) \Leftrightarrow y_j \in T_2^{(s+1,k,m)}(v_{m+1})$. Suppose $(i-1)k+1 \le p \le ik$, where $2 \le i \le s+1$. Suppose the number of pairs of vertices selected from $T_2^{(s+1,k,m)}(v_{m+1}) \times T_1^{(s+1,k,m)}(u_{m+1})$ between rounds k+1 and 2k is j_1 , between rounds 2k+1 and 3k is j_2, \ldots , between rounds (i-2)k+1 and (i-1)k is j_{i-2} , between rounds (i-1)k+1 and p is j'. Then Duplicator wins EHR_{weak} $T_2^{(s+1,k,m)}(v_{m+1}), T_1^{(s+1,k,m)}(u_{m+1}); j_1, j_2, \dots, j_{i-2}, j'+ik-p, \underbrace{k, \dots, k}_{(s+1-i) \text{ many}}$ with designated pairs $(y_{i_1+\ell}, x_{i_1+\ell}), \dots, (y_{i_r+\ell}, x_{i_r+\ell})$, where the first $j_1+j_2+\dots, j_{i-2}+j_{i-2}$ rounds have been played out, with Spoiler starting on $T_1^{(s+1,k,m)}(u_{m+1})$.

It is straightforward to see that Duplicator is able to maintain all the conditions stated above, via a similar inductive argument as has been detailed in this paper before. She is able to maintain Condition (B4) because of Conclusion 5.5 and Lemma 3.6. This brings us to the end of the inductive proof.

5.2. When Spoiler starts playing on $T_1^{(s,k,m)}$. The construction remains the same as before. The induction hypothesis in this case is simpler to state.

Induction hypothesis 5.6. Duplicator wins $\text{EHR}_{\text{alt}}\left[T_1^{(s,k,m)}, T_2^{(s,k,m)}, s; k\right]$ when Spoiler starts on $T_1^{(s,k,m)}$.

We show here the proof by induction. Refer to Figures 4 and 5. Again, we chalk out the detailed strategy of Duplicator for the first k rounds, where we do not use Induction hypothesis 5.6. For every $1 \leq t \leq m+1$ and $1 \leq t' \leq m$, she fixes any tree isomorphism $\varphi_{t,t'}^{(s+1)}: T_1^{(s+1,k,m)}(u_t) \to T_2^{(s+1,k,m)}(v_{t'})$ with $\varphi_{t,t'}^{(s+1)}(u_t) = v_{t'}$. This is possible since each of $T_1^{(s+1,k,m)}(u_t)$ and $T_2^{(s+1,k,m)}(v_{t'})$ is a copy of $T_2^{(s,k,m)}$ for all $1 \leq t \leq m+1$ and $1 \leq t' \leq m$. For the first p rounds, for all $1 \le p \le k$, she maintains the following conditions on the configuration $\{(x_i, y_i) : 1 \le i \le p\}$:

- (A'1) For all $1 \le i \le p$, we have $x_i = R_1 \Leftrightarrow y_i = R_2$. (A'2) For all $1 \le i \le p$, we have $x_i \in T_1^{(s+1,k,m)}(u_t)$ for some $1 \le t \le m+1$ if and only if $y_i \in T_2^{(s+1,k,m)}(v_{t'})$ for some $1 \le t' \le m$. Furthermore $y_i = \varphi_{t,t'}^{(s+1)}(x_i)$.
- (A'3) If $x_i, x_j \in T_1^{(s+1,k,m)}(u_t)$ for some $1 \le i \ne j \le p$ and some $1 \le t \le m+1$, then there exists some $1 \le t' \le m$ such that $y_i, y_j \in T_2^{(s+1,k,m)}(v_{t'})$. The converse is also true, i.e. if $y_i, y_j \in \overline{T_2^{(s+1,k,m)}}(v_{t'})$ for some $1 \le i \ne j \le p$ and some $1 \le t' \le m$, then there exists some $1 \le t \le m+1$ such that $x_i, x_j \in T_1^{(s+1,k,m)}(u_t)$.

That Duplicator is able to maintain these conditions can be shown by induction on pwhere $1 \le p \le k-1$. So, suppose the first p rounds have been played. In the (p+1)-st round Spoiler chooses x_{p+1} from $T_1^{(s+1,k,m)}$. Duplicator replies as follows:

(i) If $x_{p+1} = R_1$, then Duplicator sets $y_{p+1} = R_2$.

- (ii) If $x_{p+1} \in T_1^{(s+1,k,m)}(u_t)$ for some $1 \le t \le m+1$ such that there exists some $1 \le j \le p$ with $x_j \in T_1^{(s+1,k,m)}(u_t)$ as well, then Duplicator finds the $1 \le t' \le m$ such that $y_j \in T_2^{(s+1,k,m)}(v_{t'})$ and sets $y_{p+1} = \varphi_{t,t'}^{(s+1)}(x_{p+1})$.
- (iii) If $x_{p+1} \in T_1^{(s+1,k,m)}(u_t)$ for some $1 \leq t \leq m+1$ such that $x_j \notin T_1^{(s+1,k,m)}(u_t)$ for all $1 \leq j \leq p$, then Duplicator finds $1 \leq t' \leq m$ such that $v_{t'}$ has been free up to round p. She can always find such a $v_{t'}$ since $m \ge (s+1)k > k > p$. Then she sets $y_{p+1} = \varphi_{t,t'}^{(s+1)}(x_{p+1}).$

Assumption 5.7. Since $m \ge (s+1)k > k$, hence there will be at least one $1 \le i_0 \le m+1$ such that u_{i_0} is free up to round k. Since for all $1 \leq t \leq m+1$, $T_1^{(s+1,k,m)}(u_t)$ is a copy of $T_2^{(s,k,m)}$, hence we can assume without loss of generality that $i_0 = m + 1$.

We make Assumption 5.7 throughout the rest of Subsection 5.2.

Now we come to the second part of the game, where there are sk many rounds, and Spoiler plays the first k rounds of this part on $T_2^{(s+1,k,m)}$. Note that $T_1^{(s+1,k,m)}(u_{m+1})$ is a copy of $T_2^{(s,k,m)}$ whereas $T_2^{(s+1,k,m)}(v_{m+1})$ is a copy of $T_1^{(s,k,m)}$.

 $\begin{array}{c|c} \textbf{Conclusion} & \textbf{5.8.} & By \quad induction & hypothesis, & Duplicator \\ \text{EHR}_{\text{alt}} \left[T_2^{(s+1,k,m)}(v_{m+1}), T_1^{(s+1,k,m)}(u_{m+1}), s; k \right] & when \quad Spoiler \quad starts \quad pla \end{array}$ wins playing on $T_2^{(s+1,k,\bar{m})}(v_{m+1}).$

We can again split up the two trees as follows:

$$T_1^{(s+1,k,m)} = \{R_1\} \cup S_1 \cup T_1^{(s+1,k,m)}(u_{m+1}), \text{ where } S_1 = \bigcup_{t=1}^m T_1^{(s+1,k,m)}(u_t), \qquad (5.3)$$

and

$$T_2^{(s+1,k,m)} = \{R_2\} \cup S_2 \cup T_2^{(s+1,k,m)}(v_{m+1}), \text{ where } S_2 = \bigcup_{t'=1}^m T_2^{(s+1,k,m)}(v_{t'}).$$
(5.4)

In the second part of the game, for every $k \leq p \leq (s+1)k$, Duplicator maintains the following conditions on the configuration $\{(x_i, y_i) : 0 \le i \le p\}$:

- (B'1) For $0 \le i \le p$, we have $x_i = R_1 \Leftrightarrow y_i = R_2$.
- (B'2) For $0 \le i \le p$, we have $x_i \in S_1 \Leftrightarrow y_i \in S_2$. If $x_i \in T_1^{(s+1,k,m)}(u_t)$ for some $1 \le t \le m$ and $y_i \in T_2^{(s+1,k,m)}(v_{t'})$ for some $1 \le t' \le m$, then $y_i = \varphi_{t,t'}^{(s+1)}(x_i)$.
- (B'3) For $1 \le i \ne j \le p$, we have $x_i, x_j \in T_1^{(s+1,k,m)}(u_t)$ for some $1 \le t \le m$ if and only if
- (B'4) For $0 \le j \le p$, we have $x_j \in T_1^{(s+1,k,m)}(v_{t'})$. (B'4) For $0 \le j \le p$, we have $x_j \in T_1^{(s+1,k,m)}(u_{m+1}) \Leftrightarrow y_j \in T_2^{(s+1,k,m)}(v_{m+1})$. Suppose $(i-1)k+1 \le p \le ik$, where $2 \le i \le s+1$. Suppose the number of pairs of nodes selected from $T_2^{(s+1,k,m)}(v_{m+1}) \times T_1^{(s+1,k,m)}(u_{m+1})$ between rounds k+1 and 2k is j_1 ; between rounds 2k + 1 and 3k is j_2 ; ... between rounds (i - 2)k + 1 and (i-1)k is j_{i-2} ; between rounds (i-1)k+1 and p is j'. Then Duplicator wins

Г

EHR_{weak}
$$\left| T_{2}^{(s+1,k,m)}(v_{m+1}), T_{1}^{(s+1,k,m)}(u_{m+1}); j_{1}, j_{2}, \dots, j_{i-2}, j'+ik-p, \underbrace{k, \dots, k}_{(s+1-i) \text{ many}} \right|,$$

٦

where the first $j_1 + j_2 + ... j_{i-2} + j'$ rounds have been played out, with Spoiler starting on $T_2^{(s+1,k,m)}(u_{m+1})$.

Again, it is straightforward to see that Duplicator is able to maintain all the conditions stated above. She is able to maintain the Condition (B'4) because of Conclusion 5.8 and Lemma 3.6. This brings us to the end of the inductive proof.

6. Acknowledgements

The author expresses her sincere gratitude to her doctoral advisor Joel Spencer for suggesting this beautiful combinatorial problem to her. She also extends her thanks to Maksim Zhukovskii for carefully perusing through the proofs in this paper for correctness of content, and for suggesting further scopes of research in this area. This research has been supported partially by the grant NSF DMS-1444084.

References

- J. Spencer, The Strange Logic of Random Graphs, Springer Publishing Company, Inc. 2010; Series : Algorithms and Combinatorics, Vol. 22; ISBN:3642074995 9783642074998.
- [2] N. Immerman, Descriptive complexity, Springer Science & Business Media, 2012; ISBN-13: 978-0387986005.
- [3] D. Marker, Model theory: an introduction, Springer Science & Business Media, 2006; ISBN: 978-1-4419-3157-3.
- [4] G. Erich, K. G. Phokion, L. Libkin, M. Marx, J. Spencer, M. Y. Vardi, Y. Venema and S. Weinstein, Finite Model Theory and its applications, Springer Science & Business Media, 2007; eBook ISBN: 978-3-540-68804-4.
- [5] T. Place and M. Zeitoun, The tale of the quantifier alternation hierarchy of first-order logic over words, ACM SIGLOG News, Volume 2, Issue 3, Pages 4–17, 2015, ACM.
- [6] M. Kufleitner and A. Lauser, Quantifier alternation in two-variable first-order logic with successor is decidable, arXiv preprint arXiv:1212.6500, 2012.
- [7] T. Place and M. Zeitoun, Going higher in the first-order quantifier alternation hierarchy on words, International Colloquium on Automata, Languages, and Programming, Pages 342–353, 2014, Springer.
- [8] M. Bojańczyk and L. Segoufin, Tree languages defined in first-order logic with one quantifier alternation, International Colloquium on Automata, Languages, and Programming, Pages 233–245, 2008, Springer.
- [9] H. J. Keisler and W. B. Lotfallah, First order quantifiers in monadic second order logic, The Journal of Symbolic Logic, Volume 69, Issue 1, Pages 118–136, 2004, Cambridge University Press.
- [10] E. Pezzoli, Computational Complexity of Ehrenfeucht-Frass Games on Finite Structures, International Workshop on Computer Science Logic, Pages 159–170, 1998, Springer.
- [11] E. Pezzoli, On the computational complexity of type-two functionals and logical games on finite structures, Ph. D. thesis, Stanford University, 1998.
- [12] A. D. Matushkin, and M. E. Zhukovskii, ME First order sentences about random graphs: small number of alternations, Discrete Applied Mathematics, Volume 236, Pages 329–346, 2018, Elsevier.
- [13] O. Matz, N. Schweikardt and W. Thomas, The monadic quantifier alternation hierarchy over grids and graphs, Information and Computation, Volume 179, Issue 2, Pages 356–383, 2002, Academic Press.
- [14] O. Matz, Dot-depth, monadic quantifier alternation, and first-order closure over grids and pictures, Theoretical Computer Science, Volume 270, Issue 1-2, Pages 1–70, 2002, Elsevier.

- [15] A. Chandra and D. Harel, Structure and complexity of relational queries, Journal of Computer and System Sciences, Volume 25, Issue 1, Pages 99–128, 1982, Elsevier.
- [16] W. Thomas, Classifying regular events in symbolic logic, Journal of Computer and System Sciences, Volume 25, Issue 3, Pages 360–376, 1982, Elsevier.
- [17] O. Matz and N. Schweikardt, Expressive power of monadic logics on words, trees, pictures, and graphs, Logic and Automata, Volume 2, Pages 531–552, 2008.

MOUMANTI PODDER, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WASHINGTON, C-524

PADELFORD HALL, WEST STEVENS WAY NORTHEAST, SEATTLE, WA 98105, UNITED STATES. E-mail address: mpodder3@uw.edu.