

PERTURBATION THEORY OF KMS STATES

S. EJIMA AND Y. OGATA

ABSTRACT. We extend the new perturbation formula of equilibrium states by Hastings to KMS states of general W^* -dynamical systems.

1. INTRODUCTION

Perturbation theory of equilibrium states is a basic problem in quantum statistical physics which has been studied from old time. For a finite system with Hamiltonian H , equilibrium state with inverse temperature β is defined by the Gibbs state:

$$\frac{\mathrm{Tr}(e^{-\beta H} A)}{\mathrm{Tr}(e^{-\beta H})}. \quad (1)$$

If we perturb H by V , $e^{-\beta H}$ is replaced by $e^{-\beta(H+V)}$. Using the Duhamel formula, $e^{-\beta(H+V)}$ can be obtained from $e^{-\beta H}$ by multiplying

$$E_V^\tau \left(\frac{i\beta}{2} \right) = \sum_{n \geq 0} \left(-\frac{\beta}{2} \right)^n \int_{0 \leq s_n \leq \dots \leq s_1 \leq 1} ds_1 \cdots ds_n \tau_{\frac{i\beta}{2} s_n}(V) \cdots \tau_{\frac{i\beta}{2} s_1}(V) \quad (2)$$

and its adjoint from left and right respectively. Here τ is the dynamics given by H , i.e., $\tau_t(A) := e^{itH} A e^{-itH}$. Namely, we have

$$e^{-\beta(H+V)} = E_V^\tau \left(\frac{i\beta}{2} \right) e^{-\beta H} \left(E_V^\tau \left(\frac{i\beta}{2} \right) \right)^*. \quad (3)$$

In the thermodynamic limit, this $E_V^\tau \left(\frac{i\beta}{2} \right)$ can diverge. In spite of that, an analogous representation exists in infinite systems as well, thanks to Araki [A]. (See Theorem 9.) Corresponding to the possible divergence of $E_V^\tau \left(\frac{i\beta}{2} \right)$, the representation by Araki is written in terms of an unbounded operator.

In [H], Hastings introduced a new representation of $e^{-\beta(H+V)}$ with $E_V^\tau \left(\frac{i\beta}{2} \right)$ replaced by some bounded operator $\mathcal{O}(V)$:

$$e^{-\beta(H+V)} = \mathcal{O}(V) e^{-\beta H} (\mathcal{O}(V))^*. \quad (4)$$

The main result of this paper is the extension of Hastings's result to W^* -dynamical systems. For the notations and known facts about W^* -dynamical systems used in this paper, see Appendix A and Appendix B. Let \mathcal{H} be a Hilbert space, $\mathfrak{M} \subset B(\mathcal{H})$ a von Neumann algebra. We denote the set of self-adjoint elements in \mathfrak{M} by $\mathfrak{M}_{\mathrm{sa}}$. The predual of \mathfrak{M} is denoted by \mathfrak{M}_* . Let $\Omega \in \mathcal{H}$ be a cyclic and separating unit vector for \mathfrak{M} . Define the functional $\omega \in \mathfrak{M}_*$ by $\omega(A) = (\Omega, A\Omega)$ for all $A \in \mathfrak{M}$. Let Δ be the modular

operator and J the modular conjugation associated to (\mathfrak{M}, Ω) . Let σ be the modular automorphism group, and σ^Q its perturbation by $Q \in \mathfrak{M}_{sa}$. Define Liouvillean L of σ by $L = \log \Delta$. (See Appendix B.) For $V \in \mathfrak{M}_{sa}$, define $\Omega_V = e^{\frac{L+V}{2}} \Omega$ and the functional $\omega_V \in \mathfrak{M}_*$ by $\omega_V(A) = (\Omega_V, A\Omega_V)$ for all $A \in \mathfrak{M}$. In [A], it was proven that $\frac{\omega_V}{\omega_V(1)}$ is a $(\sigma^V, -1)$ -KMS state. The explicit expansion of ω_V is given, in terms of unbounded operator, the Liouvillean L . This corresponds to the expansion (3). (See Appendix Theorem 9 (e).)

Our main Theorem is an alternative representation of ω_V , in terms of bounded operators, which corresponds to the expression by Hastings (4).

Define the real function f on \mathbb{R} by

$$f(t) = \begin{cases} \sum_{n=0}^{\infty} \frac{2}{n + \frac{1}{2}} e^{-2\pi(n+\frac{1}{2})|t|} = -2 \log \left(\tanh \frac{\pi|t|}{2} \right), & (t \neq 0) \\ 0 & (t = 0) \end{cases} \quad (5)$$

Then, $f \in L^1(\mathbb{R})$. Throughout this paper we fix this f . Properties of f are collected in Appendix C.

Define the maps $\Phi : \mathfrak{M}_{sa} \times \mathbb{R} \rightarrow \mathfrak{M}_{sa}$ by

$$\Phi(V; u) = \int_{\mathbb{R}} f(t) \sigma_t^{uV}(V) dt, \quad (6)$$

for all $V \in \mathfrak{M}_{sa}$ and $u \in \mathbb{R}$. The right hand side integral is defined in the σ -weak topology [BR1]. Furthermore, $\Phi(V; u)$ is continuous in the norm topology with respect to $u \in \mathbb{R}$. (See Lemma 7.) Properties of $\Phi(V; u)$ we use in this paper are shown in Section 4. Define $\theta : \mathfrak{M}_{sa} \rightarrow \mathfrak{M}$ by

$$\theta(V) = \sum_{n=0}^{\infty} \int_0^1 du_1 \int_0^{u_1} du_2 \cdots \int_0^{u_{n-1}} du_n \Phi(V; u_1) \cdots \Phi(V; u_n), \quad (7)$$

for all $V \in \mathfrak{M}_{sa}$. The integral and summation is in the norm topology. Here is our main Theorem.

Theorem 1. *For any $V \in \mathfrak{M}_{sa}$, we have*

$$\omega_V(A) = (\Omega, \theta(V)^* A \theta(V) \Omega), \quad A \in \mathfrak{M}. \quad (8)$$

2. PROOF OF THEOREM 1 FOR ANALYTIC V

In this section, we show Theorem 1 for analytic V . We will use facts in Appendix B. For $s \in \mathbb{R}$, define the vector $\Omega_s := \Omega_{sV} = e^{\frac{L+sV}{2}} \Omega$ and the functional $\omega_s \in \mathfrak{M}_*$ by $\omega_s(A) := \omega_{sV}(A) = (\Omega_s, A\Omega_s)$ for all $A \in \mathfrak{M}$. Especially, we have $\Omega_0 = \Omega$ and $\omega_0 = \omega$. For $s \in \mathbb{R}$, the W^* -dynamics σ^{sV} satisfies $\sigma_t^{sV}(A) := e^{it(L+sV)} A e^{-it(L+sV)}$ for all $A \in \mathfrak{M}$ and $t \in \mathbb{R}$. Define $\Delta_{\Omega_s} = e^{(L+sV-sJVJ)}$ which is the modular operator of the pair (\mathfrak{M}, Ω_s) . By Theorem 9(d), Ω_s is continuous in the norm topology with respect to $s \in \mathbb{R}$.

Lemma 2. *Let $V \in \mathfrak{M}$ be a σ -entire analytic and self-adjoint element. Then Ω_s is differentiable with respect to $s \in \mathbb{R}$ in the norm topology, and we have*

$$\frac{d}{ds}\Omega_s = \frac{1}{2} \int_0^1 du \sigma_{-\frac{iu}{2}}^{sV}(V)\Omega_s. \quad (9)$$

Proof. First we recall several equalities which can be shown using identity theorem: We have $\Omega \in D(e^{z(L+sV)})$ and

$$e^{z(L+sV)}\Omega = E_{sV}(-iz)\Omega, \quad (10)$$

for all $z \in \mathbb{C}$ and $s \in \mathbb{R}$. We also have $\Omega \in D\left(e^{z\frac{L+sV}{2}}e^{(1-z)\frac{L+\tilde{s}V}{2}}\right)$ for all $z \in \mathbb{C}$ and $s, \tilde{s} \in \mathbb{R}$,

$$e^{z\left(\frac{L+sV}{2}\right)}e^{(1-z)\left(\frac{L+\tilde{s}V}{2}\right)}\Omega = E_{sV}\left(-\frac{iz}{2}\right)E_{\tilde{s}V}\left(\frac{i\tilde{z}}{2}\right)^*e^{\frac{L+\tilde{s}V}{2}}\Omega. \quad (11)$$

The proofs are analogous to that of (15), (16) below that we omit them.

Since the right-hand side of (11) is analytic in norm, the left-hand side is analytic too. Therefore, it is differentiable with respect to z in norm. We claim that $\Omega \in D\left(e^{z\frac{L+sV}{2}}Ve^{(1-z)\frac{L+\tilde{s}V}{2}}\right)$ and the derivative is given by

$$\frac{d}{dz}\left(e^{z\left(\frac{L+sV}{2}\right)}e^{(1-z)\left(\frac{L+\tilde{s}V}{2}\right)}\Omega\right) = \frac{s-\tilde{s}}{2}e^{z\left(\frac{L+sV}{2}\right)}Ve^{(1-z)\left(\frac{L+\tilde{s}V}{2}\right)}\Omega \quad (12)$$

for all $z \in \mathbb{C}$ and $s, \tilde{s} \in \mathbb{R}$.

To prove this, define the subspace $D_s \subset \mathcal{H}$ for $s \in \mathbb{R}$, by

$$D_s := \bigcup_{N \in \mathbb{N}} E_{[-N, N]}\mathcal{H}, \quad (13)$$

where $\{E_\lambda\}_{\lambda \in \mathbb{R}}$ is the projection-valued measure such that $\int_{\mathbb{R}} dE_\lambda \lambda = L + sV$. Then, D_s is a core for $e^{z(L+sV)}$ for all $z \in \mathbb{C}$.

For $\xi \in D_s$, we have

$$\begin{aligned} & \left(\xi, \frac{d}{dz}e^{z\left(\frac{L+sV}{2}\right)}e^{(1-z)\left(\frac{L+\tilde{s}V}{2}\right)}\Omega\right) = \frac{d}{dz}\left(e^{\tilde{z}\left(\frac{L+sV}{2}\right)}\xi, e^{(1-z)\left(\frac{L+\tilde{s}V}{2}\right)}\Omega\right) \\ & = \left(\left(\frac{L+sV}{2}\right)e^{\tilde{z}\left(\frac{L+sV}{2}\right)}\xi, e^{(1-z)\left(\frac{L+\tilde{s}V}{2}\right)}\Omega\right) + \left(e^{\tilde{z}\left(\frac{L+sV}{2}\right)}\xi, -\left(\frac{L+\tilde{s}V}{2}\right)e^{(1-z)\left(\frac{L+\tilde{s}V}{2}\right)}\Omega\right) \\ & = \frac{s-\tilde{s}}{2}\left(e^{\tilde{z}\left(\frac{L+sV}{2}\right)}\xi, Ve^{(1-z)\left(\frac{L+\tilde{s}V}{2}\right)}\Omega\right). \end{aligned} \quad (14)$$

We claim

$$Ve^{(1-z)\left(\frac{L+\tilde{s}V}{2}\right)}\Omega \in D\left(e^{z\left(\frac{L+sV}{2}\right)}\right) \quad (15)$$

and

$$\begin{aligned} & e^{z(\frac{L+sV}{2})} V e^{(1-z)(\frac{L+\tilde{s}V}{2})} \Omega \\ &= E_{sV}^\sigma \left(-\frac{iz}{2} \right) \sigma_{-\frac{iz}{2}}(V) \left(E_{sV}^\sigma \left(-\frac{\overline{iz}}{2} \right) \right)^* E_{sV}^\sigma \left(-\frac{iz}{2} \right) \left(E_{\tilde{s}V}^\sigma \left(-\frac{\overline{iz}}{2} \right) \right)^* e^{(\frac{L+\tilde{s}V}{2})} \Omega. \end{aligned} \quad (16)$$

From (14) and (15), we obtain (12).

To prove (15) and (16), let ξ be an arbitrary element of D_s and consider the following two complex functions on \mathbb{C} :

$$z \mapsto \left(e^{\bar{z}(\frac{L+sV}{2})} \xi, V e^{(1-z)(\frac{L+\tilde{s}V}{2})} \Omega \right) \quad (17)$$

and

$$z \mapsto \left(\xi, E_{sV}^\sigma \left(-\frac{iz}{2} \right) \sigma_{-\frac{iz}{2}}(V) \left(E_{sV}^\sigma \left(-\frac{\overline{iz}}{2} \right) \right)^* E_{sV}^\sigma \left(-\frac{iz}{2} \right) \left(E_{\tilde{s}V}^\sigma \left(-\frac{\overline{iz}}{2} \right) \right)^* e^{(\frac{L+\tilde{s}V}{2})} \Omega \right). \quad (18)$$

These two functions coincide on $i\mathbb{R}$ since for $t \in \mathbb{R}$ we have

$$\begin{aligned} & \left(e^{it(\frac{L+sV}{2})} \xi, V e^{(1-it)(\frac{L+\tilde{s}V}{2})} \Omega \right) \\ &= \left(\xi, e^{it(\frac{L+sV}{2})} V e^{-it(\frac{L+\tilde{s}V}{2})} e^{\frac{L+\tilde{s}V}{2}} \Omega \right) \\ &= \left(\xi, \sigma_{\frac{t}{2}}^{sV}(V) e^{it(\frac{L+sV}{2})} e^{-it\frac{L}{2}} \left(e^{it(\frac{L+\tilde{s}V}{2})} e^{-it\frac{L}{2}} \right)^* e^{\frac{L+\tilde{s}V}{2}} \Omega \right) \\ &= \left(\xi, E_{sV}^\sigma \left(\frac{t}{2} \right) \sigma_{\frac{t}{2}}(V) \left(E_{sV}^\sigma \left(\frac{t}{2} \right) \right)^* E_{sV}^\sigma \left(\frac{t}{2} \right) \left(E_{\tilde{s}V}^\sigma \left(\frac{t}{2} \right) \right)^* e^{\frac{L+\tilde{s}V}{2}} \Omega \right). \end{aligned} \quad (19)$$

The second equality follows by (76) and the third equality follows due to (75), (77). Since these two functions are analytic (recall (74) and (10)), they coincide on \mathbb{C} . Since D_s is a core for $e^{\bar{z}(\frac{L+sV}{2})}$, we obtain the claim (15) and (16). Hence we have completed the proof of (12).

Finally, we show (9). To prove this, first note that from (16) and the expansion of $E_{sV}^\sigma(z)$ (79), we obtain the following continuity in the norm:

$$\lim_{s \rightarrow \tilde{s}} e^{z(\frac{L+sV}{2})} V e^{(1-z)(\frac{L+\tilde{s}V}{2})} \Omega = e^{z(\frac{L+\tilde{s}V}{2})} V e^{(1-z)(\frac{L+\tilde{s}V}{2})} \Omega. \quad (20)$$

For $s, \tilde{s} \in \mathbb{R}$, since the map

$$\mathbb{R} \ni u \mapsto e^{u(\frac{L+sV}{2})} e^{(1-u)(\frac{L+\tilde{s}V}{2})} \Omega \in \mathcal{H} \quad (21)$$

is differentiable in the norm topology and satisfies

$$\frac{d}{du} e^{u(\frac{L+sV}{2})} e^{(1-u)(\frac{L+\tilde{s}V}{2})} \Omega = \left(\frac{s - \tilde{s}}{2} \right) e^{u(\frac{L+sV}{2})} V e^{(1-u)(\frac{L+\tilde{s}V}{2})} \Omega, \quad (22)$$

(from (12)), we have

$$\begin{aligned}
 \Omega_s - \Omega_{\tilde{s}} &= e^{\frac{L+sV}{2}} \Omega - e^{\frac{L+\tilde{s}V}{2}} \Omega \\
 &= \int_0^1 du \frac{d}{du} e^{u(\frac{L+sV}{2})} e^{(1-u)(\frac{L+\tilde{s}V}{2})} \Omega \\
 &= \frac{s - \tilde{s}}{2} \int_0^1 du e^{u(\frac{L+sV}{2})} V e^{(1-u)(\frac{L+\tilde{s}V}{2})} \Omega
 \end{aligned} \tag{23}$$

Therefore from the continuity (20) and (16), using the Lebesgue's dominated convergence theorem, Ω_s is differentiable with respect to s in the norm topology and we have

$$\begin{aligned}
 \frac{d}{ds} \Omega_s &= \frac{1}{2} \int_0^1 du e^{u(\frac{L+sV}{2})} V e^{(1-u)(\frac{L+sV}{2})} \Omega \\
 &= \frac{1}{2} \int_0^1 du \sigma_{-\frac{iu}{2}}^{sV}(V) \Omega_s.
 \end{aligned} \tag{24}$$

□

Lemma 3. *Let $V \in \mathfrak{M}$ be a self-adjoint σ -entire analytic element, and $A \in \mathfrak{M}$. Then $\omega_s(A)$ is differentiable with respect to s and*

$$\frac{d}{ds} \omega_s(A) = \int_0^1 du (\Omega_s, \sigma_{iu}^{sV}(V) A \Omega_s) \tag{25}$$

for all $s \in \mathbb{R}$.

Proof. By Lemma 2, $\omega_s(A)$ is differentiable and we have

$$\begin{aligned}
 \frac{d}{ds} \omega_s(A) &= \frac{d}{ds} (\Omega_s, A \Omega_s) \\
 &= \left(\frac{1}{2} \int_0^1 du \sigma_{-\frac{iu}{2}}^{sV}(V) \Omega_s, A \Omega_s \right) + \left(\Omega_s, A \left(\frac{1}{2} \int_0^1 du \sigma_{-\frac{iu}{2}}^{sV}(V) \Omega_s \right) \right) \\
 &= \frac{1}{2} \int_0^1 du \left(\left(\Omega_s, \sigma_{\frac{iu}{2}}^{sV}(V) A \Omega_s \right) + \left(\Omega_s, A \sigma_{-\frac{iu}{2}}^{sV}(V) \Omega_s \right) \right).
 \end{aligned} \tag{26}$$

We obtain

$$\left(\Omega_s, A \sigma_{-\frac{iu}{2}}^{sV}(V) \Omega_s \right) = \left(\Omega_s, \sigma_{i(1-\frac{u}{2})}^{sV}(V) A \Omega_s \right) \tag{27}$$

since we have

$$\begin{aligned}
& \left(\Omega_s, A\sigma_{-\frac{iu}{2}}^{sV}(V)\Omega_s \right) = \left(A^*\Omega_s, \sigma_{-\frac{iu}{2}}^{sV}(V)\Omega_s \right) \\
& = \left(S_{\Omega_s}A\Omega_s, S_{\Omega_s}\sigma_{\frac{iu}{2}}^{sV}(V)\Omega_s \right) = \left(\Delta_{\Omega_s}^{\frac{1}{2}}\sigma_{\frac{iu}{2}}^{sV}(V)\Omega_s, \Delta_{\Omega_s}^{\frac{1}{2}}A\Omega_s \right) \\
& = \left(\Delta_{\Omega_s}^{1-\frac{u}{2}}V\Omega_s, A\Omega_s \right) = \left(\sigma_{-i(1-\frac{u}{2})}^{sV}(V)\Omega_s, A\Omega_s \right) \\
& = \left(\Omega_s, \sigma_{i(1-\frac{u}{2})}^{sV}(V)A\Omega_s \right). \tag{28}
\end{aligned}$$

Therefore we have

$$\begin{aligned}
\frac{d}{ds}\omega_s(A) &= \frac{1}{2} \int_0^1 du \left(\Omega_s, \sigma_{\frac{iu}{2}}^{sV}(V)A\Omega_s \right) + \frac{1}{2} \int_0^1 du \left(\Omega_s, \sigma_{i(1-\frac{u}{2})}^{sV}(V)A\Omega_s \right) \\
&= \int_0^{\frac{1}{2}} du \left(\Omega_s, \sigma_{iu}^{sV}(V)A\Omega_s \right) + \int_{\frac{1}{2}}^1 du \left(\Omega_s, \sigma_{iu}^{sV}(V)A\Omega_s \right) \\
&= \int_0^1 du \left(\Omega_s, \sigma_{iu}^{sV}(V)A\Omega_s \right). \tag{29}
\end{aligned}$$

□

Lemma 4. *Let $V \in \mathfrak{M}$ be a σ -entire analytic and self-adjoint element, and $A \in \mathfrak{M}$. Let $f \in L^1(\mathbb{R})$ be the function defined in (5). Then for all $s \in \mathbb{R}$, we have*

$$\int_0^1 du \left(\Omega_s, \sigma_{iu}^{sV}(V)A\Omega_s \right) = (\Phi(V; s)\Omega_s, A\Omega_s) + (\Omega_s, A\Phi(V; s)\Omega_s), \tag{30}$$

where

$$\Phi(V; s) = \int_{\mathbb{R}} f(t) \sigma_t^{sV}(V) dt. \tag{31}$$

Proof. Let G be a function defined on $\mathbb{R}_{>0}$ by

$$G(\lambda) = (1 + \lambda)^{-1} \int_0^1 du \lambda^u, \quad \lambda > 0. \tag{32}$$

Define a function F on \mathbb{R} by

$$F(x) = \begin{cases} \frac{e^x - 1}{e^x + 1} \frac{1}{x} & (x \neq 0) \\ \frac{1}{2} & (x = 0) \end{cases}. \tag{33}$$

It follows that $F(x) = G(e^x)$ for all $x \in \mathbb{R}$. Therefore, from (87) in Appendix C, we have

$$G(e^x) = \int_{\mathbb{R}} f(t) e^{itx} dt, \tag{34}$$

for all $x \in \mathbb{R}$. We now show (30) for any σ -entire analytic element A . We have

$$\begin{aligned}
 \int_0^1 du (\Omega_s, \sigma_{iu}^{sV}(V)A\Omega_s) &= \int_0^1 du (\sigma_{-iu}^{sV}(V)\Omega_s, A\Omega_s) \\
 &= \left(\int_0^1 du \Delta_{\Omega_s}^u V\Omega_s, A\Omega_s \right) \\
 &= \left(\int_0^1 du \Delta_{\Omega_s}^u V\Omega_s, (1 + \Delta_{\Omega_s})^{-1}(1 + \Delta_{\Omega_s})A\Omega_s \right) \\
 &= \left((1 + \Delta_{\Omega_s})^{-1} \int_0^1 du \Delta_{\Omega_s}^u V\Omega_s, (1 + \Delta_{\Omega_s})A\Omega_s \right). \quad (35)
 \end{aligned}$$

We have $A\Omega_s \in D(\Delta_{\Omega_s})$ due to the σ -entire analyticity of A .

We have

$$(1 + \Delta_{\Omega_s})^{-1} \int_0^1 du \Delta_{\Omega_s}^u V\Omega_s = G(\Delta_{\Omega_s})V\Omega_s = \int_{\mathbb{R}} dt f(t) \Delta_{\Omega_s}^{it} V \Delta_{\Omega_s}^{-it} \Omega_s = \Phi(V; s)\Omega_s. \quad (36)$$

Therefore we have

$$\begin{aligned}
 \int_0^1 du (\Omega_s, \sigma_{iu}^{sV}(V)A\Omega_s) &= \left((1 + \Delta_{\Omega_s})^{-1} \int_0^1 du \Delta_{\Omega_s}^u V\Omega_s, (1 + \Delta_{\Omega_s})A\Omega_s \right) \\
 &= (\Phi(V; s)\Omega_s, (1 + \Delta_{\Omega_s})A\Omega_s) \\
 &= (\Phi(V; s)\Omega_s, A\Omega_s) + (\Phi(V; s)\Omega_s, \Delta_{\Omega_s}A\Omega_s) \\
 &= (\Omega_s, \Phi(V; s)A\Omega_s) + (\Delta_{\Omega_s}^{\frac{1}{2}}\Phi(V; s)\Omega_s, \Delta_{\Omega_s}^{\frac{1}{2}}A\Omega_s) \\
 &= (\Omega_s, \Phi(V; s)A\Omega_s) + (J\Delta_{\Omega_s}^{\frac{1}{2}}A\Omega_s, J\Delta_{\Omega_s}^{\frac{1}{2}}\Phi(V; s)\Omega_s) \\
 &= (\Omega_s, \Phi(V; s)A\Omega_s) + (\Omega_s, A\Phi(V; s)\Omega_s). \quad (37)
 \end{aligned}$$

We repeatedly used the self-adjointness of $\Phi(V; s)$.

To extend (30) to general $A \in \mathfrak{M}$, we just need to notice that both sides of (30) are continuous in A with respect to the strong-topology and recall that \mathfrak{M}_σ is strong-dense in \mathfrak{M} . \square

Due to Lemma 7, we can define the map $\theta : \mathfrak{M}_{sa} \times \mathbb{R} \rightarrow \mathfrak{M}_{sa}$ by

$$\theta(V; s) = \sum_{n=0}^{\infty} \int_0^s du_1 \int_0^{u_1} du_2 \cdots \int_0^{u_{n-1}} du_n \Phi(V; u_1) \cdots \Phi(V; u_n). \quad (38)$$

for all $V \in \mathfrak{M}_{sa}$ and $s \in \mathbb{R}$.

We now prove the statement of Theorem 1 for σ -entire analytic V .

Lemma 5. *Let $V \in \mathfrak{M}$ be a σ -entire analytic and self-adjoint element. Then we have*

$$\omega_s(A) = \omega(\theta(V; s)^* A \theta(V; s)), \quad s \in \mathbb{R}, \quad A \in \mathfrak{M}. \quad (39)$$

Proof. For each $s \in \mathbb{R}$, we define the map $\mathcal{L}_s : \mathfrak{M} \rightarrow \mathfrak{M}$ by

$$\mathcal{L}_s(X) = \Phi(V; s)X + X\Phi(V; s), \quad X \in \mathfrak{M}. \quad (40)$$

By Lemma 3 and Lemma 4, for any $A \in \mathfrak{M}$, we have

$$\begin{aligned} \frac{d}{ds}\omega_s(A) &= \int_0^1 du (\Omega_s, \sigma_{iu}^{sV}(V)A\Omega_s) \\ &= (\Phi(V; s)\Omega_s, A\Omega_s) + (\Omega_s, A\Phi(V; s)\Omega_s). \\ &= \omega_s \circ \mathcal{L}_s(A). \end{aligned} \quad (41)$$

Next, define the state $\phi_s \in \mathfrak{M}_*$ by

$$\phi_s(A) = \omega(\theta(V; s)^*A\theta(V; s)), \quad A \in \mathfrak{M}. \quad (42)$$

We now show that ϕ_s satisfies the same differential equation as ω_s :

$$\frac{d}{ds}\phi_s(A) = \phi_s \circ \mathcal{L}_s(A), \quad A \in \mathfrak{M}. \quad (43)$$

By Lemma 7, the map $\mathbb{R} \ni s \mapsto \Phi(V; s) \in \mathfrak{M}$ is continuous in the norm topology. Therefore $\theta(V; s)$ is differentiable in the norm topology and we have

$$\frac{d}{ds}\theta(V; s) = \Phi(V; s)\theta(V; s). \quad (44)$$

Since $\Phi(V; s)$ is self-adjoint, we obtain (43):

$$\begin{aligned} \frac{d}{ds}\phi_s(A) &= \omega(\theta(V; s)^*\Phi(V; s)^*A\theta(V; s)) + \omega(\theta(V; s)^*A\Phi(V; s)\theta(V; s)) \\ &= \omega(\theta(V; s)^*\mathcal{L}_s(A)\theta(V; s)) \\ &= \phi_s \circ \mathcal{L}_s(A). \end{aligned} \quad (45)$$

As ϕ_s and ω_s satisfies the same differential equation and $\omega_0 = \phi_0$, we obtain $\omega_s = \phi_s$ for all $s \in \mathbb{R}$. \square

3. PROOF OF THEOREM 3

In order to extend the result to general V , we need the following continuity of $\theta(V; s)\Omega$.

Lemma 6. *Let $\{V_m\}_{m \in \mathbb{N}} \subset \mathfrak{M}_{sa}$ be a sequence such that $V_m \rightarrow V \in \mathfrak{M}_{sa}$ strongly as $m \rightarrow \infty$. Then we have $\|\theta(V_m; s)\Omega - \theta(V; s)\Omega\| \rightarrow 0$.*

Proof. From the uniform boundedness principle, we have $c := \sup_m \|V_m\| < \infty$. By Lemma 7, $\Phi(V_m; s)$ converges to $\Phi(V; s)$ strongly. From this convergence and the uniform boundedness

$$\sup_m \|\Phi(V_m; s)\| \leq \|f\|_{L^1} \sup_m \|V_m\| \leq c\|f\|_{L^1}, \quad (46)$$

we have

$$\lim_{m \rightarrow \infty} \|\Phi(V_m; u_1) \dots \Phi(V_m; u_n)\Omega - \Phi(V; u_1) \dots \Phi(V; u_n)\Omega\| = 0. \quad (47)$$

We also have

$$\|\Phi(V_m; u_1) \dots \Phi(V_m; u_n) \Omega - \Phi(V; u_1) \dots \Phi(V; u_n) \Omega\| \leq (c^n + \|V\|^n) \|f\|_{L^1}^n, \quad (48)$$

for any $u_1, \dots, u_n \in \mathbb{R}$ and $n \in \mathbb{N}$. From the definition of $\theta(V; s)$, we have

$$\|\theta(V_m; s)\Omega - \theta(V; s)\Omega\| \quad (49)$$

$$\leq \sum_{n=0}^{\infty} \int_0^s du_1 \int_0^{u_1} du_2 \dots \int_0^{u_{n-1}} du_n \|(\Phi(V_m; u_1) \dots \Phi(V_m; u_n) - \Phi(V; u_1) \dots \Phi(V; u_n)) \Omega\| \quad (50)$$

From (47) and (48), applying Lebesgue dominated convergence Theorem, we obtain

$$\lim_{m \rightarrow \infty} \|\theta(V_m; s)\Omega - \theta(V; s)\Omega\| = 0. \quad (51)$$

□

Proof of Theorem 1. Note that $\theta(V) = \theta(V; 1)$. There exists a sequence $\{V_m\} \subset \mathfrak{M}_{sa}$ such that V_m is σ -entirely analytic for all $m \in \mathbb{N}$ and that $V_m \rightarrow V$ strongly. Let $A \in \mathfrak{M}$. By Lemma 6, we have

$$(\Omega, \theta(V_m)^* A \theta(V_m) \Omega) \rightarrow (\Omega, \theta(V)^* A \theta(V) \Omega) \quad (52)$$

as $m \rightarrow \infty$.

By Theorem 9(d), ω_{V_m} converges to ω_V in the norm topology. By Lemma 5 we have

$$\omega_{V_m}(A) = (\Omega, \theta(V_m)^* A \theta(V_m) \Omega). \quad (53)$$

for all $m \in \mathbb{N}$. Therefore we conclude that

$$\omega_V(A) = (\Omega, \theta(V)^* A \theta(V) \Omega). \quad (54)$$

□

4. PROPERTIES OF $\Phi(V; s)$

In this section, we collect some properties of Φ used in Section 2 and Section 3.

Lemma 7. (1) For all $V \in \mathfrak{M}_{sa}$ and $s \in \mathbb{R}$, we have

$$\|\Phi(V; s)\| \leq \|f\|_{L^1} \|V\|. \quad (55)$$

(2) For all $V \in \mathfrak{M}_{sa}$ the map defined by $\mathbb{R} \ni s \mapsto \Phi(V; s) \in \mathfrak{M}$ is norm-continuous.

(3) For any $s \in \mathbb{R}$ and sequence V_N in \mathfrak{M}_{sa} such that $V_N \rightarrow V \in \mathfrak{M}_{sa}$ in the strong operator topology, we have $\Phi(V_N; s) \rightarrow \Phi(V; s)$ in the strong operator topology.

Proof. For each $V \in \mathfrak{M}_{sa}$, $t \in \mathbb{R}$ and $n = 0, 1, \dots$, set

$$\mathcal{T}_n(V; t) := i^n \int_{0 \leq s_n \leq \dots \leq s_1 \leq t} ds_1 \dots ds_n (\sigma_{s_n}(V) \dots \sigma_{s_1}(V)). \quad (56)$$

Here the integral is defined in σ -weak topology. Note that the map $\mathbb{R} \ni t \mapsto \mathcal{T}_n(V; t) \in \mathfrak{M}$ is norm-continuous. Furthermore, its norm is bounded

$$\|\mathcal{T}_n(V; t)\| \leq \frac{|t|^n \|V\|^n}{n!}. \quad (57)$$

We have

$$E_{sV}^\sigma(t) = \sum_{n \geq 0} s^n \mathcal{T}_n(V; t), \quad (58)$$

where the summation converges in norm. From (75), we have

$$\sigma_t^{sV}(A) = E_{sV}^\sigma(t) \sigma_t(A) E_{sV}^\sigma(t)^* = \sum_{n \geq 0} \sum_{m \geq 0} s^{n+m} \mathcal{T}_n(V; t) \sigma_t(A) (\mathcal{T}_m(V; t))^*, \quad (59)$$

where the summation in the last term converges in the norm topology.

For each $\varepsilon > 0$, we fix $M_\varepsilon > 0$ such that

$$\int_{|t| \geq M_\varepsilon} dt |f(t)| < \varepsilon. \quad (60)$$

Now let us prove the claim of the Lemma. The first statement of the Lemma is trivial. Let us prove the second statement of the Lemma. For all $V \in \mathfrak{M}_{sa}$, $\xi \in \mathcal{H}$, $s, \tilde{s} \in \mathbb{R}$ and $\varepsilon > 0$, using (57), (60), we have

$$\begin{aligned} & \|(\Phi(V; \tilde{s}) - \Phi(V; s)) \xi\| \\ & \leq \int_{|t| \leq M_\varepsilon} dt f(t) \|(\sigma_t^{\tilde{s}V}(V) - \sigma_t^{sV}(V)) \xi\| + \int_{|t| \geq M_\varepsilon} dt f(t) \|(\sigma_t^{\tilde{s}V}(V) - \sigma_t^{sV}(V)) \xi\| \\ & \leq \int_{|t| \leq M_\varepsilon} dt f(t) \sum_{n \geq 0} \sum_{m \geq 0} |\tilde{s}^{n+m} - s^{n+m}| \|(\mathcal{T}_n(V; t) \sigma_t(V) (\mathcal{T}_m(V; t))^*) \xi\| + 2\varepsilon \|\xi\| \|V\| \\ & \leq \left(\sum_{n \geq 0} \sum_{m \geq 0} |\tilde{s}^{n+m} - s^{n+m}| \|f\|_{L^1} \frac{|M_\varepsilon|^{n+m}}{n!m!} \|V\|^{n+m+1} + 2\varepsilon \|V\| \right) \|\xi\| \end{aligned} \quad (61)$$

Therefore we have

$$\|\Phi(V; \tilde{s}) - \Phi(V; s)\| \leq \sum_{n \geq 0} \sum_{m \geq 0} |\tilde{s}^{n+m} - s^{n+m}| \|f\|_{L^1} \frac{|M_\varepsilon|^{n+m} \|V\|^{n+m+1}}{n!m!} + 2\varepsilon \|V\|, \quad (62)$$

for any $\varepsilon > 0$, $s, \tilde{s} \in \mathbb{R}$ and $V \in \mathfrak{M}_{sa}$. As $\tilde{s} \rightarrow s$, the first term converges to 0 by Lebesgue dominated convergence theorem and we obtain

$$\limsup_{\tilde{s} \rightarrow s} \|\Phi(V; \tilde{s}) - \Phi(V; s)\| \leq 2\varepsilon \|V\|, \quad (63)$$

for any $\varepsilon > 0$, $s \in \mathbb{R}$ and $V \in \mathfrak{M}_{sa}$. Therefore, we have

$$\lim_{\tilde{s} \rightarrow s} \|\Phi(V; \tilde{s}) - \Phi(V; s)\| = 0, \quad (64)$$

for any $s \in \mathbb{R}$ and $V \in \mathfrak{M}_{sa}$. This proves the second statement of Lemma.

To prove the third statement, let V_N in \mathfrak{M}_{sa} be a sequence such that $V_N \rightarrow V \in \mathfrak{M}_{sa}$ in the strong operator topology. By the uniform boundedness principle, we have $c := \sup_N \|V_N\| < \infty$. Fix arbitrary $\varepsilon > 0$, $\xi \in \mathcal{H}$, and $s \in \mathbb{R}$. We have

$$\begin{aligned}
& \|(\Phi(V_N; s) - \Phi(V; s)) \xi\| \\
& \leq \int_{|t| \leq M_\varepsilon} dt f(t) \|(\sigma_t^{sV_N}(V_N) - \sigma_t^{sV}(V)) \xi\| + \int_{|t| \geq M_\varepsilon} dt f(t) \|(\sigma_t^{sV_N}(V_N) - \sigma_t^{sV}(V)) \xi\| \\
& \leq \int_{|t| \leq M_\varepsilon} dt f(t) \sum_{n \geq 0} \sum_{m \geq 0} |s^{n+m}| \|(\mathcal{T}_n(V_N; t) \sigma_t(V_N) (\mathcal{T}_m(V_N; t))^* - \mathcal{T}_n(V; t) \sigma_t(V) (\mathcal{T}_m(V; t))^*) \xi\| \\
& + \varepsilon \|\xi\| (\|V\| + c), \tag{65}
\end{aligned}$$

for all $N \in \mathbb{N}$. Note that

$$\begin{aligned}
& \|(\mathcal{T}_n(V_N; t) \sigma_t(V_N) (\mathcal{T}_m(V_N; t))^* - \mathcal{T}_n(V; t) \sigma_t(V) (\mathcal{T}_m(V; t))^*) \xi\| \\
& \leq \int_{0 \leq s_n \leq \dots \leq s_1 \leq t} ds_1 \cdots ds_n \int_{0 \leq u_n \leq \dots \leq u_1 \leq t} du_1 \cdots du_m \\
& \|(\sigma_{s_n}(V_N) \cdots \sigma_{s_1}(V_N) \sigma_t(V_N) \sigma_{u_1}(V_N) \cdots \sigma_{u_m}(V_N) - \sigma_{s_n}(V) \cdots \sigma_{s_1}(V) \sigma_t(V) \sigma_{u_1}(V) \cdots \sigma_{u_m}(V)) \xi\|. \tag{66}
\end{aligned}$$

From Lemma 8, the integrand on the last line converges to 0 as $N \rightarrow \infty$, point wise. On the other hand, they are bounded by $\|V\|^{n+m+1} + c^{n+m+1}$. Therefore, from Lebesgue dominated convergence Theorem, we see that

$$\lim_{N \rightarrow \infty} \|(\mathcal{T}_n(V_N; t) \sigma_t(V_N) (\mathcal{T}_m(V_N; t))^* - \mathcal{T}_n(V; t) \sigma_t(V) (\mathcal{T}_m(V; t))^*) \xi\| = 0. \tag{67}$$

From (66), we also have

$$\|(\mathcal{T}_n(V_N; t) \sigma_t(V_N) (\mathcal{T}_m(V_N; t))^* - \mathcal{T}_n(V; t) \sigma_t(V) (\mathcal{T}_m(V; t))^*) \xi\| \leq \frac{M_\varepsilon^{n+m}}{n!m!} (\|V\|^{n+m+1} + c^{n+m+1}) \|\xi\|, \tag{68}$$

for any $|t| \leq M_\varepsilon$. From (67) and (68), applying the Lebesgue dominated convergence Theorem, we have

$$\lim_{N \rightarrow \infty} \int_{|t| \leq M_\varepsilon} dt f(t) \sum_{n \geq 0} \sum_{m \geq 0} |s^{n+m}| \|(\mathcal{T}_n(V_N; t) \sigma_t(V_N) (\mathcal{T}_m(V_N; t))^* - \mathcal{T}_n(V; t) \sigma_t(V) (\mathcal{T}_m(V; t))^*) \xi\| = 0. \tag{69}$$

Substituting this to (65), we obtain

$$\limsup_{N \rightarrow \infty} \|(\Phi(V_N; s) - \Phi(V; s)) \xi\| \leq \varepsilon \|\xi\| (\|V\| + c), \tag{70}$$

for any $\varepsilon > 0$, $\xi \in \mathcal{H}$, and $s \in \mathbb{R}$. As ε is arbitrary, we have

$$\lim_{N \rightarrow \infty} \|(\Phi(V_N; s) - \Phi(V; s)) \xi\| = 0, \tag{71}$$

for any $\xi \in \mathcal{H}$, and $s \in \mathbb{R}$. This proves the third statement. \square

APPENDIX A. W^* -DYNAMICAL SYSTEMS

We collect known facts on W^* -dynamical systems. See [BR2] and [DJP] and references therein for more information. Let \mathcal{H} be a Hilbert space and $\mathfrak{N} \subset B(\mathcal{H})$ a von Neumann algebra. We denote by \mathfrak{N}_* the predual of \mathfrak{N} . We also use \mathfrak{N}_{sa} the set of all selfadjoint elements in \mathfrak{N} . Let $\text{Aut}(\mathfrak{N})$ denote the group of all $*$ -automorphisms of the von Neumann algebra \mathfrak{N} . A family $\{\tau_t\}_{t \in \mathbb{R}} \subset \text{Aut}(\mathfrak{N})$ is called W^* -dynamics if it is satisfied that for all $t, s \in \mathbb{R}$, $\tau_s \circ \tau_t = \tau_{s+t}$, $\tau_0 = 1$ and $\tau_t(A)$ converges to A in the σ -weak topology as $t \rightarrow 0$. As τ_t is an automorphism, for any $A \in \mathfrak{N}$, $\mathbb{R} \ni t \mapsto \tau_t(A) \in \mathfrak{N}$ is continuous with respect to the σ -strong topology. The pair (\mathfrak{N}, τ) is called a W^* -dynamical system. For an entire τ -analytic $A \in \mathfrak{N}$, we denote by $\tau_z(A)$, for $z \in \mathbb{C}$ the analytic continuation of $\mathbb{R} \ni t \mapsto \tau_t(A) \in \mathfrak{N}$. We denote the set of all τ -entire analytic elements in \mathfrak{N} by \mathfrak{N}_τ . Then \mathfrak{N}_τ is a $*$ -subalgebra in \mathfrak{N} , which is dense in \mathfrak{N} with respect to the strong topology. Let $\beta \in \mathbb{R}$. A normal state ω on \mathfrak{N} is called a (τ, β) -KMS state if for all $A \in \mathfrak{N}$ and $B \in \mathfrak{N}_\tau$,

$$\omega(A\tau_{i\beta}(B)) = \omega(BA). \quad (72)$$

Any (τ, β) -KMS state is τ -invariant.

Finally, we consider the perturbation theory of the dynamics. Let (\mathfrak{N}, τ) be a W^* -dynamical system and $Q \in \mathfrak{N}_{sa}$. For $t \in \mathbb{R}$, we define τ_t^Q by

$$\tau_t^Q(A) = \sum_{n \geq 0} (it)^n \int_{0 \leq s_n \leq \dots \leq s_1 \leq 1} ds_1 \cdots ds_n [\tau_{s_n t}(Q), [\dots, [\tau_{s_1 t}(Q), \tau_t(A)] \cdots]]. \quad (73)$$

for all $A \in \mathfrak{N}$. The integral is defined in the σ -weak topology and the summation is in the norm topology.

Then τ^Q is a W^* -dynamics. For all $t \in \mathbb{R}$, define $E_Q^\tau(t) \in \mathfrak{N}$ by

$$\begin{aligned} E_Q^\tau(t) &= \sum_{n \geq 0} (it)^n \int_{0 \leq s_n \leq \dots \leq s_1 \leq 1} ds_1 \cdots ds_n \tau_{s_n t}(Q) \cdots \tau_{s_1 t}(Q) \\ &= \sum_{n \geq 0} i^n \int_{0 \leq s_n \leq \dots \leq s_1 \leq t} ds_1 \cdots ds_n \tau_{s_n}(Q) \cdots \tau_{s_1}(Q). \end{aligned} \quad (74)$$

The integral is defined in the σ -weak topology and the summation is in the norm topology. This $E_Q^\tau(t)$ is a unitary element of \mathfrak{N} and

$$\tau_t^Q(A) = E_Q^\tau(t) \tau_t(A) E_Q^\tau(t)^* \quad (75)$$

for all $t \in \mathbb{R}$ and $A \in \mathfrak{N}$. Suppose that there exists a self-adjoint operator L on \mathcal{H} satisfying $\tau_t(A) = e^{itL} A e^{-itL}$ for all $t \in \mathbb{R}$ and $A \in \mathfrak{M}$. Then, for all $t \in \mathbb{R}$, $A \in \mathfrak{M}$ and $Q \in \mathfrak{M}_{sa}$,

$$\tau_t^Q(A) = e^{it(L+Q)} A e^{-it(L+Q)} \quad (76)$$

and

$$E_Q^\tau(t) = e^{it(L+Q)} e^{-itL}. \quad (77)$$

If A, Q are entire-analytic for τ , then A is τ^Q -entire analytic and $\mathbb{R} \ni t \mapsto E_Q^\tau(t) \in \mathfrak{M}$ has an entire analytic continuation. We have the expansion

$$\tau_z^Q(A) = \sum_{n \geq 0} (iz)^n \int_{0 \leq s_n \leq \dots \leq s_1 \leq 1} ds_1 \cdots ds_n [\tau_{s_n z}(Q), [\dots, [\tau_{s_1 z}(Q), \tau_z(A)] \cdots]], \quad (78)$$

and

$$E_Q^\tau(z) = \sum_{n \geq 0} (iz)^n \int_{0 \leq s_n \leq \dots \leq s_1 \leq 1} ds_1 \cdots ds_n \tau_{s_n z}(Q) \cdots \tau_{s_1 z}(Q) \quad (79)$$

for all $z \in \mathbb{C}$ respectively. The integration and the summation converges in the norm topology. ([DJP])

APPENDIX B. TOMITA-TAKESAKI THEORY

We collect known facts Tomita-Takesaki Theory. See [BR2] and [DJP] and references therein for more information. Let \mathcal{H} be a Hilbert space, $\mathfrak{M} \subset B(\mathcal{H})$ a von Neumann algebra, and $\Omega \in \mathcal{H}$ a cyclic and separating vector for \mathfrak{M} . Define the operator S_Ω^0 on \mathcal{H} with the domain $\mathfrak{M}\Omega$ by

$$S_\Omega^0 A \Omega = A^* \Omega, \quad A \in \mathfrak{M}. \quad (80)$$

Then S_Ω^0 is anti-linear and closable. Let S_Ω be its closure. Let $S_\Omega = J_\Omega \Delta_\Omega^{\frac{1}{2}}$ be the polar decomposition of S_Ω . The operators Δ_Ω and J_Ω are called respectively the modular operator and the modular conjugation for (\mathfrak{M}, Ω) . The modular operator Δ_Ω is non-singular. The modular operator and the modular conjugation satisfy $J_\Omega \Delta_\Omega J_\Omega = \Delta_\Omega^{-1}$, $\Delta_\Omega^z \Omega = \Omega$ for all $z \in \mathbb{C}$ and $\log \Delta_\Omega \Omega = 0$. Furthermore, J_Ω interchange \mathfrak{M} and \mathfrak{M}' , i.e., $J_\Omega \mathfrak{M} J_\Omega = \mathfrak{M}'$. From the fact that $\Delta_\Omega^{it} \mathfrak{M} \Delta_\Omega^{-it} = \mathfrak{M}$ all $t \in \mathbb{R}$, we can define a *-automorphism σ_t on \mathfrak{M} by $\sigma_t(A) = \Delta_\Omega^{it} A \Delta_\Omega^{-it}$, $A \in \mathfrak{M}$. This W^* -dynamics σ is called the modular automorphism associated to Ω . We set $L_\Omega = \log \Delta_\Omega$ and call it the Liouvillean of σ . One can check that the normalized state of ω is a $(\sigma, -1)$ -KMS state.

The set $\mathfrak{M}_\sigma \Omega$ is a core for S_Ω and $\Delta_\Omega^{\frac{1}{2}}$. (Proposition 2.5.22 of [BR1].) We use the following lemma.

Lemma 8. *Let $Q \in \mathfrak{M}$, $n \in \mathbb{N}$ and $t_1, \dots, t_n \in \mathbb{R}$. Let $\{Q_m\}_{m \in \mathbb{N}} \subset \mathfrak{M}$ satisfy that $Q_m \rightarrow Q$ and $Q_m^* \rightarrow Q^*$ strongly as $m \rightarrow \infty$. Then*

$$\Delta_\Omega^{it_n} Q_m \cdots \Delta_\Omega^{it_1} Q_m \xi \rightarrow \Delta_\Omega^{it_n} Q \cdots \Delta_\Omega^{it_1} Q \xi \quad (81)$$

as $m \rightarrow \infty$, for any $\xi \in \mathcal{H}$.

The following theorem is shown in [A].

Theorem 9. *Let $Q \in \mathfrak{M}_{sa}$. Then $\Omega \in D\left(e^{\frac{L_\Omega + Q}{2}}\right)$.*

Define the vectors $\Omega_Q := e^{\frac{L_\Omega + Q}{2}} \Omega$, and the functional ω_Q on \mathfrak{M} by

$$\omega_Q(A) = (\Omega_Q, A \Omega_Q) \quad (82)$$

for all $A \in \mathfrak{M}$. Then the following conditions are satisfied;

- (a) Ω_Q is cyclic and separating for \mathfrak{M} ;
- (b) The normalized state of ω_Q is a $(\sigma^Q, -1)$ -KMS state;
- (c) By (a), we can define the modular operator Δ_{Ω_Q} and the modular conjugation J_{Ω_Q} for the pair (\mathfrak{M}, Ω_Q) . Then $J_\Omega = J_{\Omega_Q}$,

$$\log \Delta_{\Omega_Q} = L_\Omega + Q - J_\Omega Q J_\Omega \quad (83)$$

and

$$\sigma_t^Q(A) = \Delta_{\Omega_Q}^{it} A \Delta_{\Omega_Q}^{-it} = e^{it(L_\Omega + Q)} A e^{-it(L_\Omega + Q)}; \quad (84)$$

- (d) Assume that a sequence $\{Q_n\}_{n \in \mathbb{N}} \subset \mathfrak{M}_{sa}$ converges to $Q \in \mathfrak{M}_{sa}$ strongly. Then $\Omega_{Q_n} \rightarrow \Omega_Q$ and $\omega_{Q_n} \rightarrow \omega_Q$ in the norm topology;
- (e) For any $0 \leq s_n \leq \dots \leq s_1 \leq 1$, Ω is in the domain of $e^{\frac{1}{2}s_n L} Q \cdot e^{\frac{1}{2}(-s_n + s_{n-1})L} Q \dots e^{\frac{1}{2}(-s_2 + s_1)L} Q$ and Ω_Q has an expansion

$$\sum_{n \geq 0} \left(\frac{1}{2}\right)^n \int_{0 \leq s_n \leq \dots \leq s_1 \leq 1} ds_1 \dots ds_n e^{\frac{1}{2}s_n L} Q \cdot e^{\frac{1}{2}(-s_n + s_{n-1})L} Q \dots e^{\frac{1}{2}(-s_2 + s_1)L} Q \Omega. \quad (85)$$

Due to (c) of this theorem, let us denote J the modular conjugation if there is no danger of confusion.

APPENDIX C. PROPERTIES OF f

In this section we consider some properties of f . Clearly f is in $L^1(\mathbb{R})$ by Fubini's Theorem. This f and F in (33) are Fourier transform of each other.

Lemma 10. *For f defined in (5) and F defined in (33), we have*

$$\lim_{R \rightarrow \infty} \int_{-R}^R F(w) e^{-iwt} dw = f(t) \quad (86)$$

for all $t \in \mathbb{R} \setminus \{0\}$ and


$$F(x) = \int_{-\infty}^{\infty} f(t) e^{ixt} dt \quad (87)$$

for all $x \in \mathbb{R}$.

Proof. In order to prove this we consider the analytic continuation of F (defined in (33)) to $\mathbb{C} \setminus (2\pi i (\mathbb{Z} + \frac{1}{2}))$:

$$F(z) = \begin{cases} \frac{e^z - 1}{e^z + 1} \frac{1}{z} & (z \in \mathbb{C} \setminus (2\pi i (\mathbb{Z} + \frac{1}{2}) \cup \{0\})) \\ \frac{1}{2} & (z = 0). \end{cases} \quad (88)$$

The claim (86) for $t < 0$ follows applying the residue theorem to $F(z)e^{-izt}$ along the path in the following picture.



Each pole $z = 2\pi i \left(n + \frac{1}{2}\right)$ of $F(z)e^{-izt}$ is a simple pole of $F(z)e^{-izt}$, and

$$\begin{aligned} \operatorname{Res}\left(F(z)e^{-izt}; 2\pi i \left(n + \frac{1}{2}\right)\right) &= \lim_{z \rightarrow 2\pi i \left(n + \frac{1}{2}\right)} \left(z - 2\pi i \left(n + \frac{1}{2}\right)\right) F(z)e^{-izt} \\ &= -\frac{i}{\pi \left(n + \frac{1}{2}\right)} e^{2\pi \left(n + \frac{1}{2}\right)t}. \end{aligned} \quad (89)$$

From the residue Theorem, we obtain

$$\begin{aligned} \int_{-R}^R dw F(w)e^{-iwt} &= \int_{\gamma_1} F(z)e^{-izt} dz \\ &= 2\pi i \sum_{k=0}^{n-1} \operatorname{Res}\left(F(z)e^{-izt}; 2\pi i \left(k + \frac{1}{2}\right)\right) \\ &\quad - \int_{\gamma_2} F(z)e^{-izt} dz - \int_{\gamma_3} F(z)e^{-izt} dz - \int_{\gamma_4} F(z)e^{-izt} dz. \end{aligned} \quad (90)$$

The integral along γ_3 goes to 0 as $n \rightarrow \infty$. The integral along γ_2, γ_4 goes to 0 as $R \rightarrow \infty$. Hence we obtain (86) for all $t < 0$.

To show that (86) for $t > 0$, we note

$$\int_{-R}^R F(w)e^{-iwt} dw = \overline{\int_{-R}^R \overline{F(w)} e^{-iw(-t)} dw} = \overline{\int_{-R}^R F(w)e^{-iw(-t)} dw}. \quad (91)$$

The final term converges to $f(-t) = f(t)$ as $R \rightarrow \infty$ due to $-t < 0$. Hence we obtain (86) for all $t \neq 0$.

Finally we show (87). Note that $F|_{\mathbb{R}} \in L^2(\mathbb{R})$. Let $\hat{F} \in L^2(\mathbb{R})$ be the Fourier transform of $F|_{\mathbb{R}}$. For $R > 0$, we define the function ϕ_R on \mathbb{R} by

$$\phi_R(t) = \int_{-R}^R F(x)e^{-ixt} dx. \quad (92)$$

for all $t \in \mathbb{R}$. Then we have $\|\phi_R - \hat{F}\|_2 \rightarrow 0$ as $R \rightarrow \infty$ by Theorem 9.13 of [R]. Therefore there exists a sequence $\{n_k\}_{k \in \mathbb{N}}$ such that $\phi_{n_k}(t) \rightarrow \hat{F}(t)$ as $k \rightarrow \infty$ a.e. t . Since we have

shown

$$\lim_{R \rightarrow \infty} \phi_R(t) = f(t) \quad (93)$$

for all $t \in \mathbb{R} \setminus \{0\}$, it follows that $f(t) = \hat{F}(t)$ a.e. t . Hence $\hat{F} = f$ belongs to $L^1(\mathbb{R})$. By Theorem 9.14 of [R], we have

$$F(x) = \int_{-\infty}^{\infty} \hat{F}(t)e^{ixt} dt = \int_{-\infty}^{\infty} f(t)e^{ixt} dt \quad (94)$$

a.e. x . The left-hand side is continuous. The right-hand side is continuous too due to $f \in L^1(\mathbb{R})$. Therefore the equality holds for all $x \in \mathbb{R}$. □

REFERENCES

- [A] H. Araki: Expansionals in Banach algebras, *Ann. Sci. Ecole Norm. Sup.* 6 , **6**, 67, 1973.
- [BR1] O. Bratteli and D. W. Robinson: *Operator Algebras and Quantum Statistical Mechanics 1* *Springer-Verlag* 1987.
- [BR2] O. Bratteli and D. W. Robinson: *Operator Algebras and Quantum Statistical Mechanics 2* *Springer-Verlag* 1996.
- [DJP] J. Dereziński, V. Jaksic and C.-A. Pillet: Perturbation theory of W^* -dynamics, Liouvilleans and KMS-states. *Reviews in Mathematical Physics*,15-05, 447–489, 2003.
- [H] M.B. Hastings: Quantum belief propagation: An algorithm for thermal quantum systems. *Phys. Rev. B* **76**, 201102 2007.
- [R] W. Rudin: *Real and Complex Analysis*. *McGraw-Hill Publishing Company*, 1966.