

ALEXANDER AND JONES POLYNOMIALS OF SURGERIZED TST LINKS

WILSON WONG, FRANKY MOK

Dedicated to our team members Sam Lau, Aaron Chan and Kenith Chan, and our teacher Mr. S. H. Chan, with whom we worked together on a HLMA project in 2012.

ABSTRACT. This paper is a continuation on the 2012 paper on "Cutting Twisted Solid Tori (TSTs)", in which we considered twisted solid torus links (tst links). We generalize the notion of tst links to "surgerized tst links": recall that when performing $\Phi^\mu(n(\tau), d(\tau), M)$ on a tst $\langle \tau \rangle$ where M is odd, we obtain the tst link, $[\Phi^\mu(n(\tau), d(\tau), M)]$ that contains a trivial knot as one of its components. We then perform another operation $\Phi^{\mu'}(n(\tau'), d(\tau'), M')$ on that trivial knot to create a new link, which we call a "surgerized tst link" (stst link). If M' is odd, we can repeat the process to give more complicated stst links. We compute braid words, Alexander and Jones polynomials of such links.

INTRODUCTION

Recall that a twisted solid torus link (tst link) $[\Phi^\mu(n(\tau), d(\tau), M)]$ is obtained by cutting a twisted solid torus (tst) $\langle \tau \rangle$ into M pieces with chosen denominator $d(\tau)$ by inserting $\mu(M-1)$ blades. For details please see [5]. The resultant link has $(\mu P + I)$ components, where P and I are defined in the paper [5] using the number of partial tsts attached to different faces in the fundamental graph $\Lambda(n(\tau), d(\tau), M)$. When $N(\tau) \neq \tau$, P and I are respectively the same as the multiplicity of knotted tst $\langle N(\tau) \rangle_K$ and $\langle \tau \rangle_K$ in the knotted tst sum $\Phi(n(\tau), d(\tau), M)$. When $N(\tau) = \tau$, the multiplicity of $\langle \tau \rangle_K$ is $P + I$. In particular I has a simple formula:

$$I = \frac{1 - (-1)^M}{2} = M \pmod{2}$$

This means that I is 1 when M is odd and 0 when M is even. The tst link $[\Phi^\mu(n(\tau), d(\tau), M)]$ is a $(-\mu PN(\tau), \mu PD(\tau))$ -torus link when M is even, where each component is a $\langle N(\tau) \rangle_K$, and a $(-\mu PN(\tau), \mu PD(\tau))$ -torus link with a trivial knot, the components being $\langle N(\tau) \rangle_K$ and $\langle \tau \rangle_K$ respectively when M is odd.

Thus, if M is odd, we can consider the following surgery: first, we drill $\langle \tau \rangle_K$ from S^3 with the $(-\mu PN(\tau), \mu PD(\tau))$ -torus link left untouched. Then we replace the part that is drilled away with a new twisted solid torus $\langle \tau' \rangle$. Afterwards we perform the operation $\Phi^{\mu'}(n(\tau'), d(\tau'), M')$ on this new tst.

The resultant knotted tst sum is then denoted $\Phi(n(\tau), d(\tau), M, \mu; n(\tau'), d(\tau'), M', \mu')$, and the resultant link is denoted $[\Phi(n(\tau), d(\tau), M, \mu; n(\tau'), d(\tau'), M', \mu')]$ and called

Date: January 31, 2019.

2010 Mathematics Subject Classification. 57N99, 57R56.

Key words and phrases. Knot Theory, Algebraic Topology, Topological Quantum Field Theory.

a surgerized tst link (stst link). If we have

$$\Phi^{\mu'}(n(\tau'), d(\tau'), M') = \mu' P' \langle N(\tau') \rangle_K + I' \langle \tau' \rangle_K$$

then $[\Phi(n(\tau), d(\tau), M, \mu; n(\tau'), d(\tau'), M', \mu')]$ has $(\mu P + \mu' P' + I')$ components, and

$$\Phi(n(\tau), d(\tau), M, \mu; n(\tau'), d(\tau'), M', \mu') = \mu P \langle N(\tau) \rangle_K + \mu' P' \langle N(\tau') \rangle_K + I' \langle \tau' \rangle_K$$

If M' is odd, we can repeat the process and obtain the stst link

$$[\Phi(n(\tau), d(\tau), M, \mu; n(\tau'), d(\tau'), M', \mu'; n(\tau''), d(\tau''), M'', \mu'')]$$

and so forth. Performing the surgery $(k-1)$ times and performing the cut $\Phi^{\mu_j}(n(\tau_j), d(\tau_j), M_j)$ the $(j-1)^{\text{st}}$ time, where M_1, M_2, \dots, M_{k-1} are odd, we obtain the stst link

$$[\Phi(n(\tau_1), d(\tau_1), M_1, \mu_1; n(\tau_2), d(\tau_2), M_2, \mu_2; \dots; n(\tau_k), d(\tau_k), M_k, \mu_k)]$$

or more compactly denoted by $[\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j)]$. We also have the knotted tst sum given by

$$\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j) = \sum_{j=1}^k \mu_j P_j \langle N(\tau_j) \rangle_K + I_k \langle \tau_k \rangle_K$$

where $\Phi^{\mu_j}(n(\tau_j), d(\tau_j), M_j) = \mu_j P_j \langle N(\tau_j) \rangle_K + \langle \tau_j \rangle_K$ for $j = 1, 2, \dots, k-1$, and $\Phi^{\mu_k}(n(\tau_k), d(\tau_k), M_k) = \mu_k P_k \langle N(\tau_k) \rangle_K + I_k \langle \tau_k \rangle_K$. The link $[\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j)]$ has $(\sum_{j=1}^k \mu_j P_j + I_k)$ components, and consists of $\mu_j P_j$ torus knots of the $(-N(\tau_j), D(\tau_j))$ type, and I_k trivial knot(s).

In this paper, we will compute a braid word for each of the stst links, then a Seifert matrix using the braid words. We will then compute the Alexander polynomials using techniques of Heegaard splitting and Fox calculus. Then, we will compute the Jones polynomials using techniques of topological quantum field theory and Heegaard splitting in a different manner.

1. BRAID WORDS

Recall that a braid word of $[\Phi^\mu(n(\tau), d(\tau), M)]$ is $(\sigma_1^{-1} \sigma_2^{-1} \dots \sigma_{\mu P D(\tau) - 1}^{-1})^{\mu P N(\tau)}$ when M is even, and $(\sigma_1^{-2} \sigma_2^{-1} \dots \sigma_{\mu P D(\tau)}^{-1})^{\mu P N(\tau)}$ when M is odd as we derived in [5]. We now write $N(\tau_j) = N_j, D(\tau_j) = D_j$ and $\mu_j P_j = S_j$ and consider a braid word for the stst link $[\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j)]$.

For $k = 2$ and M_2 even, the braid word is given by

$$\left(\prod_{i=1}^{i=S_2 D_2} \sigma_i^{-1} \prod_{i=1}^{S_1 D_1 + S_2 D_2 - 1} \sigma_i^{-1} \right)^{S_1 N_1} \left(\prod_{i=1}^{S_2 D_2 - 1} \sigma_i^{-1} \right)^{S_2 N_2} \in B_{S_1 D_1 + S_2 D_2}$$

where $\prod_{i=a}^b \beta_i = \beta_a \beta_{a+1} \dots \beta_b$, $\prod_a^{i=b} \beta_i = \beta_b \beta_{b-1} \dots \beta_a$, β_i is a braid-valued function on i and B_n is the braid group of n strings.

For M_2 odd, the braid word is given by

$$\left(\prod_{i=1}^{i=S_2 D_2 + 1} \sigma_i^{-1} \prod_{i=1}^{S_1 D_1 + S_2 D_2} \sigma_i^{-1} \right)^{S_1 N_1} \left(\sigma_1^{-1} \prod_{i=1}^{S_2 D_2} \sigma_i^{-1} \right)^{S_2 N_2} \in B_{S_1 D_1 + S_2 D_2 + 1}$$

Generalizing, a braid word of the stst link for even M_k is given by

$$\begin{aligned} & \left(\prod_{i=1}^{i=\sum_{j=2}^k S_j D_j} \sigma_i^{-1} \prod_{i=1}^{\sum_{j=1}^k S_j D_j - 1} \sigma_i^{-1} \right)^{S_1 N_1} \left(\prod_{i=1}^{i=\sum_{j=3}^k S_j D_j} \sigma_i^{-1} \prod_{i=1}^{\sum_{j=2}^k S_j D_j - 1} \sigma_i^{-1} \right)^{S_2 N_2} \cdots \left(\prod_{i=1}^{S_k D_{k-1}} \sigma_i^{-1} \right)^{S_k N_k} \\ & = \prod_{l=1}^k \left(\prod_{i=1}^{i=\sum_{j=l+1}^k S_j D_j} \sigma_i^{-1} \prod_{i=1}^{\sum_{j=1}^k S_j D_j - 1} \sigma_i^{-1} \right)^{S_l N_l} \in B_{\sum_{j=1}^k S_j D_j} \end{aligned}$$

where $\sum_{j=k+1}^k$ is interpreted as 0 and $\prod_{i=1}^{i=0}$ is interpreted as the identity braid, whereas for odd M_k , it is given by

$$\prod_{l=1}^k \left(\prod_{i=1}^{i=\sum_{j=l+1}^k S_j D_j + 1} \sigma_i^{-1} \prod_{i=1}^{\sum_{j=1}^k S_j D_j} \sigma_i^{-1} \right)^{S_l N_l} \in B_{\sum_{j=1}^k S_j D_j + 1}$$

2. ALEXANDER POLYNOMIALS

To find the Alexander polynomials of the stst link $[\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j)]$, we first review how the Alexander polynomials of the tst link $[\Phi^\mu(n(\tau), d(\tau), M)]$ is computed.

2.1. Tst links $[\Phi^\mu(n(\tau), d(\tau), M)]$ with even M . For even integers M , the link is the same as the torus link $K_{-\mu PN(\tau), \mu PD(\tau)}$. Following [3], the Alexander polynomial of the link is given by

$$\Delta_{[\Phi^\mu(n(\tau), d(\tau), M)]}(t) = (-1)^{\mu P - 1} t^{-\frac{(\mu PN(\tau) - 1)(\mu PD(\tau) - 1)}{2}} \frac{(1-t)(1-t^{\mu PN(\tau)D(\tau)})^{\mu P}}{(1-t^{\mu PN(\tau)})(1-t^{\mu PD(\tau)})}$$

Murasugi claimed that this can be computed from the Seifert matrix of the link [3], but it is in fact very difficult to do so, and which is probably why the book skipped the derivation.

We now derive this result using Fox calculus and Seifert-van Kampen Theorem. For the sake of simplicity we write $\mu P = S$, $N(\tau) = N$ and $D(\tau) = D$ and consider the Alexander polynomial of the link $L = K_{SN, SD}$. Let $n(C)$ denote the tubular neighborhood of a link C in S^3 . Firstly we note that if we put all the S knots onto the same torus ∂U that bounds a solid torus U and separate the complement of L along the torus into the manifolds $M = S^3 \setminus (U \cup n(L))$ and $N = \overline{U} \setminus n(L)$, their intersection $M \cap N$ is not connected, therefore Seifert-van Kampen Theorem cannot be applied (at least not in its common form). Instead, we consider S solid tori $U_S \subset \cdots \subset U_2 \subset U_1$ that share the same longitudinal circle. On each of their boundaries ∂U_i , we place one torus knot $K_{N, D}$ and let K_i denote that knot. Consider the 3-manifolds $X_i = \overline{U_i \setminus (U_i \cap n(K_i)) \setminus (U_{i+1} \cup n(K_{i+1}))}$ for $1 \leq i \leq S-1$, $X_0 = S^3 \setminus (\overline{U_1 \cup n(K_1)})$ and $X_S = \overline{U_S \setminus (U_S \cap K_S)}$. Then $\bigcup_{i=0}^S X_i = S^3 \setminus n(L) = S^3 \setminus L$. Now $X_{i-1} \cap X_i = \overline{\partial U_i \setminus (\partial U_i \cap n(K_i))}$ is connected, so we can apply Seifert-van Kampen Theorem. Let a_i and b_i be the meridian and longitudinal circles respectively inside the solid torus U_i for $1 \leq i \leq S$. Note that a_S is contractible inside X_S . We also introduce its meridian circle of the solid torus $V = S^3 \setminus U_1$ to be a_0 . The longitudinal circle of V is contractible in X_0 . The fundamental group $\pi_1(X_0)$ of X_0 is generated by a_0 , and that of X_i is

$$\pi_1(X_i) \cong \langle a_i, b_i \mid a_i b_i = b_i a_i \rangle$$

where $1 \leq i \leq S-1$. Now we consider the fundamental group of $X_{i-1} \cap X_i$: it is generated by g_i , which is the knot K_i slightly translated, so that it sits on the intersection. Now note that g_i winds inside U_{i-1} around the longitudinal circle D times, and N times around the meridian circle, so we have $g_i = a_{i-1}^N b_{i-1}^D$. Similarly we have $g = a_i^N b_i^D$ inside U_i . Repeatedly applying Seifert-van Kampen Theorem gives the fundamental group of the link complement:

$$\pi_1(S^3 \setminus L) \cong \left\langle \begin{array}{l} a_0, a_1, a_2, \dots, a_{S-1}, \left| \begin{array}{l} a_0^N = a_1^N b_1^D, a_i^N b_i^D = a_{i+1}^N b_{i+1}^D = b_S^D, \quad 1 \leq i \leq S-2 \\ b_1, b_2, \dots, b_S \end{array} \right. \\ a_j b_j = b_j a_j, 1 \leq j \leq S-1 \end{array} \right\rangle$$

Applying a Tietze type II transformation gives

$$\pi_1(S^3 \setminus L) \cong \left\langle \begin{array}{l} a_0, a_1, a_2, \dots, a_{S-1}, \left| \begin{array}{l} a_0^N = a_i^N b_i^D = b_S^D, \quad 1 \leq i \leq S-1 \\ b_1, b_2, \dots, b_S \end{array} \right. \\ a_j b_j = b_j a_j, \quad 1 \leq j \leq S-1 \end{array} \right\rangle$$

Next we use Fox calculus to find the Alexander matrix. Recall that under the Fox differential operator we have

$$\frac{\partial(fg)}{\partial x} = \frac{\partial f}{\partial x} + f \frac{\partial g}{\partial x}$$

Thus we have

$$\begin{aligned} \frac{\partial x^n}{\partial x} &= \frac{\partial x^{n-1}}{\partial x} + x^{n-1} \frac{\partial x}{\partial x} \\ &= \frac{\partial x^{n-1}}{\partial x} + x^{n-1} \\ &= \frac{\partial x^{n-2}}{\partial x} + x^{n-2} + x^{n-1} \\ &\vdots \\ &= \frac{\partial x}{\partial x} + x + \dots + x^{n-2} + x^{n-1} \\ &= 1 + x + \dots + x^{n-1} \\ &= \frac{1 - x^n}{1 - x} \end{aligned}$$

We then find the Fox derivative of an inverse note that $ff^{-1} = 1$ and $\frac{\partial(1)}{\partial x} = 0$:

$$\begin{aligned} \frac{\partial(ff^{-1})}{\partial x} &= \frac{\partial f}{\partial x} + f \frac{\partial f^{-1}}{\partial x} \\ 0 &= \frac{\partial f}{\partial x} + f \frac{\partial f^{-1}}{\partial x} \\ 0 &= f^{-1} \frac{\partial f}{\partial x} + \frac{\partial f^{-1}}{\partial x} \\ \frac{\partial f^{-1}}{\partial x} &= -f^{-1} \frac{\partial f}{\partial x} \end{aligned}$$

$$\begin{aligned}
& \begin{array}{c} \mathbf{C} \\ \mathbf{D} \end{array} \begin{array}{c} \left. \begin{array}{ccccccc} -\frac{1-b_1^D}{1-b_1} & & & & & & \\ & -\frac{1-b_2^D}{1-b_2} & & & & & \\ & & \ddots & & & & \\ & & & \frac{1-a_0^N}{1-a_0} & & & \\ & & & \frac{1-a_0^N}{1-a_0} & & & \\ & & & \vdots & & & \\ & & & \frac{1-a_0^N}{1-a_0} & & & \\ & & & \vdots & & & \\ & & & \frac{1-a_0^N}{1-a_0} & & -\frac{1-b_{S-1}^D}{1-b_{S-1}} & \\ & & & & & & -\frac{1-b_S^D}{1-b_S} \end{array} \right| \\ \begin{array}{ccccccc} a_1 - 1 & & & & & & \\ & a_2 - 1 & & & & & \\ & & \ddots & & & & \\ & & & 0 & & & \\ & & & & \ddots & & \\ & & & & & a_{S-1} - 1 & \end{array} \end{array} \\
& = (-1)^j \frac{1-b_S^D}{1-b_S} \begin{array}{c} \left. \begin{array}{ccccccc} 1 - a_1^N b_1^D & & & & & & \\ & 1 - a_2^N b_2^D & & & & & \\ & & \ddots & & & & \\ & & & -\frac{(1-b_j)(1-a_0^N)}{1-a_0} & & & \\ & & & -\frac{(1-b_j)(1-a_0^N)}{1-a_0} & & & \\ & & & \vdots & & & \\ & & & -\frac{(1-b_j)(1-a_0^N)}{1-a_0} & & & \\ & & & \vdots & & & \\ & & & -\frac{(1-b_j)(1-a_0^N)}{1-a_0} & & \ddots & \\ & & & & & & 1 - a_{S-1}^N b_{S-1}^D \end{array} \right| \\ \prod_{i=1,2,\dots,\hat{j},\dots,S-1} (1 - a_i^N b_i^D) \end{array} \\
& = (-1)^{j+1} \frac{1-b_S^D}{1-b_S} \frac{(1-b_j)(1-a_0^N)}{1-a_0} \\
& = (-1)^j \frac{(1-b_j)(1-a_0^N)^S}{(1-b_S)(1-a_0)}
\end{aligned}$$

where $\mathbf{C} = \text{diag}\left(\frac{1-a_1^N}{1-a_1}, \frac{1-a_2^N}{1-a_2}, \dots, \frac{1-a_{S-1}^N}{1-a_{S-1}}\right)$ and $\mathbf{D} = \text{diag}(1-b_1, 1-b_2-1, \dots, 1-b_{S-1})$ are diagonal matrix blocks.

Type IV: Here the $2S^{\text{th}}$ column is deleted.

$$\begin{array}{c}
\left| \begin{array}{cccccccc}
\frac{1-a_0^N}{1-a_0^N} & -\frac{1-a_1^N}{1-a_1} & & & -a_1^N \frac{1-b_1^D}{1-b_1} & & & \\
\frac{1-a_0^N}{1-a_0} & & -\frac{1-a_2^N}{1-a_2} & & & -a_2^N \frac{1-b_2^D}{1-b_2} & & \\
\vdots & & & \ddots & & & \ddots & \\
\frac{1-a_0^N}{1-a_0} & & & & \frac{1-a_{S-1}^N}{1-a_{S-1}} & & & -a_{S-1}^N \frac{1-b_{S-1}^D}{1-b_{S-1}} \\
\frac{1-a_0^N}{1-a_0} & & & & & & & \\
1-b_1 & & & & a_1-1 & & & \\
& 1-b_2 & & & & a_2-1 & & \\
& & \ddots & & & & \ddots & \\
& & & 1-b_{S-1} & & & & a_{S-1}-1
\end{array} \right| \\
= (-1)^{S+1} \frac{1-a_0^N}{1-a_0} \prod_{i=1}^{S-1} (1-a_i^N b_i^D) \\
= (-1)^{S+1} \frac{(1-a_0^N)^S}{1-a_0}
\end{array}$$

The Alexander ideal is thus given by

$$\left(\frac{(1-a_0^N)^S}{1-b_S}, \left\{ \frac{(1-a_j)(1-a_0^N)^S}{(1-b_S)(1-a_0)} \right\}_{j=1}^{S-1}, \left\{ \frac{(1-b_j)(1-a_0^N)^S}{(1-b_S)(1-a_0)} \right\}_{j=1}^{S-1}, \frac{(1-a_0^N)^S}{1-a_0} \right)$$

The abelianization map ϕ is given by the linking numbers, namely,

$$\begin{aligned}
\text{lk}(a_0, L) &= SD, \\
\text{lk}(a_j, L) &= (S-j)D, \\
\text{lk}(b_j, L) &= jN
\end{aligned}$$

and $\phi(x) = t^{\text{lk}(x, L)}$.

The Alexander ideal under the abelianization map is given by

$$\begin{aligned}
& \left(\frac{(1-t^{SND})^S}{1-t^{SN}}, \left\{ \frac{(1-t^{(S-j)D})(1-t^{SND})^S}{(1-t^{SN})(1-t^{SD})} \right\}_{j=1}^{S-1}, \left\{ \frac{(1-t^{jN})(1-t^{SND})^S}{(1-t^{SN})(1-t^{SD})} \right\}_{j=1}^{S-1}, \frac{(1-t^{SND})^S}{1-t^{SD}} \right) \\
&= \frac{(1-t^{SND})^S}{(1-t^{SN})(1-t^{SD})} \left(1-t^{SD}, \left\{ 1-t^{(S-j)D} \right\}_{j=1}^{S-1}, \left\{ 1-t^{jN} \right\}_{j=1}^{S-1}, 1-t^{SN} \right) \\
&= \frac{(1-t^{SND})^S}{(1-t^{SN})(1-t^{SD})} \left(1-t^{SD}, 1-t^D, \left\{ 1-t^{(S-j)D} \right\}_{j=2}^{S-1}, 1-t^N, \left\{ 1-t^{jN} \right\}_{j=2}^{S-1}, 1-t^{SN} \right) \\
&= \frac{(1-t^{SND})^S}{(1-t^{SN})(1-t^{SD})} (1-t)
\end{aligned}$$

since N and D are coprime integers.

Then, to obtain the Alexander polynomial, we have to multiply by a power of t so that the function gives the same Laurent polynomial in \sqrt{t} when evaluated in t and t^{-1} : the power is $-\frac{1}{2}(1+S^2ND-SN-SD) = -\frac{1}{2}(SN-1)(SD-1)$. Besides it is customary to add the factor $(-1)^{S-1}$ so that the leading term has positive coefficient.

Hence we have

$$\Delta_L(t) = (-1)^{S-1} t^{-\frac{(SN-1)(SD-1)}{2}} \frac{(1-t)(1-t^{SND})^S}{(1-t^{SN})(1-t^{SD})}$$

Now we note that $K_{-SN,SD}$ is the mirror image of $K_{SN,SD}$ [3], and mirror images have the same Alexander polynomial [4], so we have the desired result.

2.2. Tst links $[\Phi^\mu(n(\tau), d(\tau), M)]$ with odd M . Next we consider the Alexander polynomials of the tst link $[\Phi^\mu(n(\tau), d(\tau), M)]$ when M is odd, which we conjectured in [5] to be

$$\Delta_{[\Phi^\mu(n(\tau), d(\tau), M)]}(t) = (-1)^{\mu P} t^{-\frac{(\mu P N(\tau)-1)(\mu P D(\tau)+1)}{2}} \frac{(1-t)(1-t^{(\mu P D(\tau)+1)N(\tau)})^{\mu P}}{1-t^{\mu P D(\tau)+1}}$$

Here we add a factor of $(-1)^{\mu P}$. We shall now prove this result.

When M is odd, the link is given by a $K_{-\mu P N(\tau), \mu P D(\tau)}$ on the surface of the torus and inside the solid torus bounded by it there is a trivial knot along the longitudinal direction. As above we shall write $N = N(\tau)$, $D = D(\tau)$, $\mu P = S$ and consider the mirror image L where $K_{SN,SD}$ is placed on the surface of the torus instead of $K_{-SN,SD}$.

Like the case where M is even, we introduce S solid tori U_1, U_2, \dots, U_S with the same longitudinal circle where $U_S \subset \dots \subset U_2 \subset U_1$ and place a component $K_{N,D}$ of the torus link on each of the boundaries ∂U_i . By Seifert-van Kampen Theorem the fundamental group of $S^3 \setminus L$ is given by

$$\pi_1(S^3 \setminus L) \cong \left\langle a_0, a_1, a_2, \dots, a_S, \left. \begin{array}{l} a_0^N = a_i^N b_i^D, 1 \leq i \leq S \\ b_1, b_2, \dots, b_S \end{array} \right| \begin{array}{l} a_j b_j = b_j a_j, 1 \leq j \leq S \end{array} \right\rangle$$

Now we set the relators $r_i = a_0^N (a_i^N b_i^D)^{-1}$ and $s_j = (a_j b_j)(b_j a_j)^{-1}$. The Fox derivatives are given by

$$\begin{aligned} \frac{\partial r_i}{\partial a_0} &= \frac{1 - a_0^N}{1 - a_0} \\ \frac{\partial r_i}{\partial a_i} &= -\frac{1 - a_i^N}{1 - a_i} \\ \frac{\partial r_i}{\partial b_i} &= -a_i^N \frac{1 - b_i^D}{1 - b_i} \\ \frac{\partial s_j}{\partial a_j} &= 1 - b_j \\ \frac{\partial s_j}{\partial b_j} &= a_j - 1 \end{aligned}$$

where $1 \leq i, j \leq S$ and other Fox derivatives are zero.

Under this map the Alexander ideal is

$$\begin{aligned}
& \left((1 - t^{(SD+1)N})^S, \left\{ \frac{(1 - t^{(S-j)D+1})(1 - t^{(SD+1)N})^S}{1 - t^{SD+1}} \right\}_{j=1}^S, \left\{ \frac{(1 - t^{jN})(1 - t^{(SD+1)N})^S}{1 - t^{SD+1}} \right\}_{j=1}^S \right) \\
&= \frac{(1 - t^{(SD+1)N})^S}{1 - t^{SD+1}} \left(1 - t^{SD+1}, \left\{ 1 - t^{(S-j)D+1} \right\}_{j=1}^S, \left\{ 1 - t^{jN} \right\}_{j=1}^S \right) \\
&= \frac{(1 - t^{(SD+1)N})^S}{1 - t^{SD+1}} \left(1 - t^{SD+1}, \left\{ 1 - t^{(S-j)D+1} \right\}_{j=1}^{S-1}, 1 - t, \left\{ 1 - t^{jN} \right\}_{j=1}^S \right) \\
&= \frac{(1 - t^{(SD+1)N})^S}{1 - t^{SD+1}} (1 - t)
\end{aligned}$$

The generator of the ideal is thus given by

$$\frac{(1 - t)(1 - t^{(SD+1)N})^S}{1 - t^{SD+1}}$$

The power of t to multiply is $t^{-\frac{1}{2}(1+(SD+1)NS-(SD+1))} = t^{-\frac{(SD+1)(SN-1)+1}{2}}$. We also multiply a power of -1 so that the leading term has positive coefficient, which is given by $(-1)^S$. Hence

$$\Delta_L(t) = (-1)^S t^{-\frac{(SN-1)(SD+1)+1}{2}} \frac{(1 - t)(1 - t^{(\mu PD(\tau)+1)N(\tau)})^{\mu P}}{1 - t^{\mu PD(\tau)+1}}$$

For odd M , the link $[\Phi^\mu(n(\tau), d(\tau), M)]$ is the mirror image of L , and since mirror images have the same Alexander polynomial [4], the claimed result follows.

2.3. Stst links $[\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j)]$ **with odd** M_k . We now turn to more general surgerized twisted solid tori links (stst links) $[\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j)]$. We introduce the notations $N(\tau_j) = N_j, D(\tau_j) = D_j, \mu_j P_j = S_j$. We first consider the case where M_k is odd. We use the fact that mirror images have the same Alexander polynomial [4], and consider $\sum_{j=1}^k S_j$ solid tori with the same longitudinal circle, $U_{k,S_k} \subset \cdots \subset U_{k,1} \subset \cdots \subset U_{1,S_1} \subset \cdots \subset U_{1,1}$, inside S^3 . Then we place the S_1 components of the torus link $K_{S_1 N_1, S_1 D_1}$ on the surfaces $\partial U_{S_1,1}, \dots, \partial U_{S_1,S_1}$, and so on, up to the S_k components of the torus link $K_{S_k N_k, S_k D_k}$ on the surfaces $\partial U_{S_k,1}, \dots, \partial U_{S_k,S_k}$, and place a trivial knot inside the solid torus U_{S_k,S_k} . We denote the collection of all these torus knots and the trivial knot by L . The fundamental group of the complement of L is given by

$$\pi_1(S^3 \setminus L) \cong \left\langle \begin{array}{l} a_0, a_{1,1}, \dots, a_{1,S_1}, \dots, a_{k,1}, \dots, a_{k,S_k} \\ b_{1,1}, \dots, b_{1,S_1}, \dots, b_{k,1}, \dots, b_{k,S_k} \end{array} \left| \begin{array}{l} a_0^{N_1} = a_{1,i_1}^{N_1} b_{1,i_1}^{D_1}, a_{l-1,S_{l-1}}^{N_l} b_{l-1,S_{l-1}}^{D_l} = a_{l,i_l}^{N_l} b_{l,i_l}^{D_l} \\ a_{m,n} b_{m,n} = b_{m,n} a_{m,n} \end{array} \right. \right\rangle$$

where $1 \leq i_j \leq S_j, 2 \leq l \leq k, 1 \leq m \leq k$, and $1 \leq n \leq S_m$.

The Alexander matrix is given by

$$\begin{bmatrix} \mathbf{v}_0 & A_1 & & & C_1 & & & & \\ & B_1 & A_2 & & D_1 & C_2 & & & \\ & & \ddots & \ddots & & \ddots & \ddots & & \\ & & & B_{k-1} & A_k & & & & \\ \mathbf{0} & & E & & & F & D_{k-1} & C_k & \end{bmatrix}$$

where $\mathbf{v}_0 = \frac{1-a_0^N}{1-a_0} [1 \ 1 \ \cdots \ 1]^T$ has S_1 rows,

$$\begin{aligned} A_j &= -\text{diag} \left(\frac{1-a_{j,1}^{N_j}}{1-a_{j,1}}, \frac{1-a_{j,2}^{N_j}}{1-a_{j,2}}, \dots, \frac{1-a_{j,S_j}^{N_j}}{1-a_{j,S_j}} \right) \\ B_j &= \frac{1-a_{j,S_j}^{N_{j+1}}}{1-a_{j,S_j}} \begin{bmatrix} 0 & \cdots & 0 & 1 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 0 & 1 \end{bmatrix} \\ C_j &= -\text{diag} \left(a_{j,1}^{N_j} \frac{1-b_{j,1}^{D_j}}{1-b_{j,1}}, a_{j,2}^{N_j} \frac{1-b_{j,2}^{D_j}}{1-b_{j,2}}, \dots, a_{j,S_j}^{N_j} \frac{1-b_{j,S_j}^{D_j}}{1-b_{j,S_j}} \right) \\ D_j &= a_{j,S_j}^{N_{j+1}} \frac{1-b_{j,S_j}^{D_{j+1}}}{1-b_{j,S_j}} \begin{bmatrix} 0 & \cdots & 0 & 1 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 0 & 1 \end{bmatrix} \\ E &= \text{diag}(1-b_{1,1}, \dots, 1-b_{1,S_1}, \dots, 1-b_{k,1}, \dots, 1-b_{k,S_k}) \\ F &= \text{diag}(a_{1,1}-1, \dots, a_{1,S_1}-1, \dots, a_{k,1}-1, \dots, a_{k,S_k}-1) \end{aligned}$$

and B_j and D_j both have S_{j+1} rows and S_j columns. The Alexander ideal is generated by the following types of determinants:

Type I: The first column of the Alexander matrix is deleted. The determinant is given by

$$\prod_{j=1}^k \prod_{i=1}^{S_j} (1 - a_{j,i}^{N_j} b_{j,i}^{D_j}) = \prod_{j=1}^k (1 - a_{j,1}^{N_j} b_{j,1}^{D_j})^{S_j}$$

Type II.i: The $(i+1)^{\text{st}}$ column of the Alexander matrix is deleted, where $1 \leq i \leq S_1$. The determinant is given by

$$\begin{aligned} & (-1)^{i+1} \left(\prod_{j=2}^k \prod_{m=1}^{S_j} (1 - a_{j,m}^{N_j} b_{j,m}^{D_j}) \right) \left(\prod_{l=1,2,\dots,\hat{i},\dots,S_1} (1 - a_{1,l}^{N_1} b_{1,l}^{D_1}) \right) \frac{(1-a_{1,i})(1-a_0^{N_1})}{1-a_0} \\ &= (-1)^{i+1} \frac{(1-a_{1,i}) \prod_{j=1}^k (1 - a_{j,1}^{N_j} b_{j,1}^{D_j})^{S_j}}{1-a_0} \end{aligned}$$

The determinant of the matrix where the $(i+1)^{\text{st}}$ column of the Alexander matrix is deleted is zero for $S_1 + 1 \leq i \leq 1 + \sum_{j=1}^k S_k$.

Type III.i: The $(i+1 + \sum_{j=1}^k S_k)^{\text{th}}$ column of the Alexander matrix is deleted, where $1 \leq i \leq S_1$. The determinant is given by

$$\begin{aligned} & (-1)^{i+1+\sum_{j=1}^k S_k} \left(\prod_{j=2}^k \prod_{m=1}^{S_j} (1 - a_{j,m}^{N_j} b_{j,m}^{D_j}) \right) \left(\prod_{l=1,2,\dots,\hat{i},\dots,S_1} (1 - a_{1,l}^{N_1} b_{1,l}^{D_1}) \right) \frac{(1-b_{1,i})(1-a_0^{N_1})}{1-a_0} \\ &= (-1)^{i+1+\sum_{j=1}^k S_k} \frac{(1-b_{1,i}) \prod_{j=1}^k (1 - a_{j,1}^{N_j} b_{j,1}^{D_j})^{S_j}}{1-a_0} \end{aligned}$$

The determinant of the matrix where the $(i + 1 + \sum_{j=1}^k S_j)$ th column of the Alexander matrix is deleted is zero for $S_1 + 1 \leq i \leq \sum_{j=1}^k S_j$.

The abelianization map ϕ is given by $\phi(a_0) = t^{1+\sum_{j=1}^k S_j D_j}$, $\phi(a_{j,i}) = t^{1+(S_j-i)D_j + \sum_{l=j+1}^k S_l D_l}$ and $\phi(b_{j,i}) = t^{iN_j + \sum_{l=1}^{j-1} S_l N_l}$, where the sum $\sum_{l=k+1}^k$ and $\sum_{l=1}^0$ are interpreted as 0.

Writing

$$\begin{aligned} \Pi &= \prod_{j=1}^k \left(1 - t^{(1+(S_j-1)D_j + \sum_{l=j+1}^k S_l D_l)N_j + (N_j + \sum_{l=1}^{j-1} S_l N_l)D_j} \right)^{S_j} \\ &= \prod_{j=1}^k \left(1 - t^{N_j + S_j N_j D_j + N_j \sum_{l=j+1}^k S_l D_l + D_j \sum_{l=1}^{j-1} S_l N_l} \right)^{S_j} \end{aligned}$$

the Alexander ideal under the abelianization map is given by

$$\begin{aligned} & \left(\Pi, \left\{ \frac{(1 - t^{1-iD_1 + \sum_{l=1}^k S_l D_l}) \Pi}{1 - t^{1+\sum_{j=1}^k S_j D_j}} \right\}_{i=1}^{S_1}, \left\{ \frac{(1 - t^{iN_1}) \Pi}{1 - t^{1+\sum_{j=1}^k S_j D_j}} \right\}_{i=1}^{S_1} \right) \\ &= \frac{\Pi}{1 - t^{1+\sum_{j=1}^k S_j D_j}} \left(1 - t^{1+\sum_{j=1}^k S_j D_j}, \left\{ 1 - t^{1+(S_1-i)D_1 + \sum_{l=2}^k S_l D_l} \right\}_{i=1}^{S_1}, \left\{ 1 - t^{iN_1} \right\}_{i=1}^{S_1} \right) \\ &= \frac{\Pi}{[1 + \sum_{j=1}^k S_j D_j]} \left(\left[1 + \sum_{j=1}^k S_j D_j \right], \left\{ \left[1 + (S_1 - i)D_1 + \sum_{l=1}^k S_l D_l \right] \right\}_{i=1}^{S_1}, \{[iN_1]\}_{i=1}^{S_1} \right) \end{aligned}$$

where to write more compactly, we introduce the notation $[\lambda] = 1 - t^\lambda$ for integers λ .

We note that the following diagram is commutative:

$$\begin{array}{ccc} \mathbb{Z}^n & \xrightarrow{[\cdot]^n} & \mathbb{Z}[t]^n \\ \text{gcd} \downarrow & & \downarrow \text{gcd} \\ \mathbb{Z} & \xrightarrow{[\cdot]} & \mathbb{Z}[t] \end{array}$$

We thus have

$$\begin{aligned}
& \gcd \left(\left[1 + \sum_{j=1}^k S_j D_j \right], \left\{ \left[1 + (S_1 - i) D_1 + \sum_{l=1}^k S_l D_l \right] \right\}_{i=1}^{S_1}, \{[iN_1]\}_{i=1}^{S_1} \right) \\
&= \left[\gcd \left(1 + \sum_{j=1}^k S_j D_j, \left\{ 1 + (S_1 - i) D_1 + \sum_{l=1}^k S_l D_l \right\}_{i=1}^{S_1}, \{iN_1\}_{i=1}^{S_1} \right) \right] \\
&= \left[\gcd \left(\gcd \left(1 + (S_1 - 1) D_1 + \sum_{l=1}^k S_l D_l, 1 + (S_1 - 2) D_1 + \sum_{l=1}^k S_l D_l \right), N_1, \dots \right) \right] \\
&= \left[\gcd \left(\gcd \left(D_1, 1 + (S_1 - 2) D_1 + \sum_{l=1}^k S_l D_l \right), N_1, \dots \right) \right] \\
&= [\gcd(D_1, N_1, \dots)] \\
&= [1] \\
&= 1 - t
\end{aligned}$$

where we assumed $S_1 \geq 2$.

The generator of the Alexander ideal is thus

$$\frac{(1-t) \prod_{j=1}^k \left(1 - t^{N_j + S_j N_j D_j + N_j \sum_{l=j+1}^k S_l D_l + D_j \sum_{l=1}^{j-1} S_l N_l} \right)^{S_j}}{1 - t^{1 + \sum_{j=1}^k S_j D_j}}$$

Multiplying by powers of t and (-1) , the Alexander polynomial of L is

$$\Delta_L(t) = (-1)^{\sum_{j=1}^k S_j} t^{\frac{\sigma}{2}} \frac{(1-t) \prod_{j=1}^k \left(1 - t^{N_j + S_j N_j D_j + N_j \sum_{l=j+1}^k S_l D_l + D_j \sum_{l=1}^{j-1} S_l N_l} \right)^{S_j}}{1 - t^{1 + \sum_{j=1}^k S_j D_j}}$$

where $\sigma = \sum_{j=1}^k \left(S_j D_j - (N_j + S_j N_j D_j + N_j \sum_{l=j+1}^k S_l D_l + D_j \sum_{l=1}^{j-1} S_l N_l) \right)$.

2.4. Stst links $[\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j)]$ with even M_k . We now consider stst links $[\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j)]$ with even M_k . Writing $N(\tau_j) = N_j, D(\tau_j) = D_j, \mu_j P_j = S_j$, we also use the fact that mirror images have the same Alexander polynomial [4]. The link $[\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j)]$ contains $\sum_{j=1}^k S_j$ torus knots. We denote the collection of the mirror images of these torus knots by L . Using the technique as above, the fundamental group of the complement of L is given by

$$\pi_1(S^3 \setminus L) \cong \left\langle \begin{array}{l} a_0, a_{1,1}, \dots, a_{k-1, S_{k-1}}, a_{k,1}, \dots, a_{k, S_{k-1}}, \\ b_{1,1}, \dots, b_{1, S_1}, \dots, b_{k,1}, \dots, b_{k, S_k} \end{array} \middle| \begin{array}{l} a_0^{N_1} = a_{1, i_1}^{N_1} b_{1, i_1}^{D_1}, a_{l-1, S_{l-1}}^{N_l} b_{l-1, S_{l-1}}^{D_l} = a_{l, i_l}^{N_l} b_{l, i_l}^{D_l} \\ a_{k-1, S_{k-1}}^{N_{k-1}} b_{k-1, S_{k-1}}^{D_{k-1}} = b_{k, S_k}^{D_k}, a_{m,n} b_{m,n} = b_{m,n} a_{m,n} \end{array} \right\rangle$$

where $1 \leq i_j \leq S_j$ for $1 \leq j \leq k-1$ and $1 \leq i_k \leq S_k-1, 2 \leq l \leq k, 1 \leq n \leq S_m$ for $1 \leq m \leq k-1$, and $1 \leq n \leq S_k-1$ for $m = k$.

The Alexander matrix is given by

$$\begin{bmatrix} \mathbf{v}_0 & A_1 & & & & C_1 & & & & \\ & B_1 & A_2 & & & D_1 & C_2 & & & \\ & & \ddots & \ddots & & & \ddots & \ddots & & \\ & & & B_{k-1} & A_k & & & D_{k-1} & C_k & \\ \mathbf{0} & E & & & & F & & & & \end{bmatrix}$$

where $\mathbf{v}_0 = \frac{1-a_0^N}{1-a_0} [1 \ 1 \ \cdots \ 1]^T$ has S_1 rows,

$$A_j = -\text{diag} \left(\frac{1-a_{j,1}^{N_j}}{1-a_{j,1}}, \frac{1-a_{j,2}^{N_j}}{1-a_{j,2}}, \dots, \frac{1-a_{j,S_j}^{N_j}}{1-a_{j,S_j}} \right)$$

$$A_k = \begin{bmatrix} -\text{diag} \left(\frac{1-a_{k,1}^{N_k}}{1-a_{k,1}}, \frac{1-a_{k,2}^{N_k}}{1-a_{k,2}}, \dots, \frac{1-a_{k,S_{k-1}}^{N_k}}{1-a_{k,S_{k-1}}} \right) \\ \mathbf{0} \end{bmatrix}$$

$$B_l = \frac{1-a_{j,S_j}^{N_{j+1}}}{1-a_{j,S_j}} \begin{bmatrix} 0 & \cdots & 0 & 1 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 0 & 1 \end{bmatrix}$$

$$C_j = -\text{diag} \left(a_{j,1}^{N_j} \frac{1-b_{j,1}^{D_j}}{1-b_{j,1}}, a_{j,2}^{N_j} \frac{1-b_{j,2}^{D_j}}{1-b_{j,2}}, \dots, a_{j,S_j}^{N_j} \frac{1-b_{j,S_j}^{D_j}}{1-b_{j,S_j}} \right)$$

$$C_k = -\text{diag} \left(a_{k,1}^{N_k} \frac{1-b_{k,1}^{D_k}}{1-b_{k,1}}, a_{k,2}^{N_k} \frac{1-b_{k,2}^{D_k}}{1-b_{k,2}}, \dots, a_{k,S_{k-1}}^{N_{k-1}} \frac{1-b_{k,S_{k-1}}^{D_k}}{1-b_{k,S_{k-1}}}, \frac{1-b_{k,S_k}^{D_k}}{1-b_{k,S_k}} \right)$$

$$D_l = a_{j,S_j}^{N_{j+1}} \frac{1-b_{j,S_j}^{D_{j+1}}}{1-b_{j,S_j}} \begin{bmatrix} 0 & \cdots & 0 & 1 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 0 & 1 \end{bmatrix}$$

$$E = \text{diag}(1-b_{1,1}, \dots, 1-b_{1,S_1}, \dots, 1-b_{k,1}, \dots, 1-b_{k,S_k})$$

$$F = \text{diag}(a_{1,1}-1, \dots, a_{1,S_1}-1, \dots, a_{k,1}-1, \dots, a_{k,S_k}-1)$$

where B_l and D_l both have S_{l+1} rows and S_l columns, $1 \leq l \leq k-1$ and $1 \leq j \leq k-1$

The Alexander ideal is generated by the following types of determinants:

Type I: The first column of the Alexander matrix is deleted. The determinant, up to sign, is given by

$$\left(\prod_{j=1}^{k-1} \prod_{i=1}^{S_j} (1-a_{j,i}^{N_j} b_{j,i}^{D_j}) \right) \frac{1-b_{k,S_k}^{D_k}}{1-b_{k,S_k}} \prod_{i=1}^{S_{k-1}} (1-a_{k,i}^{N_k} b_{k,i}^{D_k}) = \frac{\prod_{j=1}^k (1-a_{j,1}^{N_j} b_{j,1}^{D_j})^{S_j}}{1-b_{k,S_k}}$$

Type II.i: The $(i+1)^{\text{st}}$ column of the Alexander matrix is deleted, where $1 \leq i \leq S_1$. The determinant, up to sign, is given by

$$\begin{aligned} & \left(\prod_{j=2}^k \prod_{m=1}^{S_j} (1 - a_{j,m}^{N_j} b_{j,m}^{D_j}) \right) \left(\prod_{l=1,2,\dots,\hat{i},\dots,S_1} (1 - a_{1,i}^{N_1} b_{1,i}^{D_1}) \right) \frac{(1 - a_{1,i})(1 - a_0^{N_1})}{(1 - a_0)(1 - b_{k,S_k})} \\ &= \frac{(1 - a_{1,i}) \prod_{j=1}^k (1 - a_{j,1}^{N_j} b_{j,1}^{D_j})^{S_j}}{(1 - a_0)(1 - b_{k,S_k})} \end{aligned}$$

The determinant of the matrix where the $(i+1)^{\text{st}}$ column of the Alexander matrix is deleted is zero for $S_1 + 1 \leq i \leq -1 + \sum_{j=1}^k S_k$.

Type III.i: The $(i+1 + \sum_{j=1}^k S_k)^{\text{th}}$ column of the Alexander matrix is deleted, where $1 \leq i \leq S_1$. Up to sign, the determinant is given by

$$\begin{aligned} & \left(\prod_{j=2}^k \prod_{m=1}^{S_j} (1 - a_{j,m}^{N_j} b_{j,m}^{D_j}) \right) \left(\prod_{l=1,2,\dots,\hat{i},\dots,S_1} (1 - a_{1,i}^{N_1} b_{1,i}^{D_1}) \right) \frac{(1 - b_{1,i})(1 - a_0^{N_1})}{(1 - a_0)(1 - b_{k,S_k})} \\ &= \frac{(1 - b_{1,i}) \prod_{j=1}^k (1 - a_{j,1}^{N_j} b_{j,1}^{D_j})^{S_j}}{(1 - a_0)(1 - b_{k,S_k})} \end{aligned}$$

The determinant of the matrix where the $(i + \sum_{j=1}^k S_k)^{\text{th}}$ column of the Alexander matrix is deleted is zero for $S_1 + 1 \leq i \leq \sum_{j=1}^k S_k$.

Type IV: The last column of the Alexander matrix is deleted. The determinant of the matrix is zero.

The abelianization map ϕ is given by $\phi(a_0) = t^{\sum_{j=1}^k S_j D_j}$, $\phi(a_{j,i}) = t^{(S_j - i)D_j + \sum_{l=j+1}^k S_l D_l}$ and $\phi(b_{j,i}) = t^{iN_j + \sum_{l=1}^{j-1} S_l N_l}$, where the sum $\sum_{l=k+1}^k$ and $\sum_{l=1}^0$ are interpreted as 0.

Writing

$$\begin{aligned} \Pi &= \prod_{j=1}^k \left(1 - t^{(S_j - 1)N_j D_j + N_j \sum_{l=j+1}^k S_l D_l + D_j N_j + D_j \sum_{k=1}^{j-1} S_l N_l} \right)^{S_j} \\ &= \prod_{j=1}^k \left(1 - t^{S_j N_j D_j + N_j \sum_{l=j+1}^k S_l D_l + D_j \sum_{k=1}^{j-1} S_l N_l} \right)^{S_j} \end{aligned}$$

the Alexander ideal under the abelianization map is given by

$$\begin{aligned}
& \left(\frac{\Pi}{1 - t^{\sum_{j=1}^k S_j N_j}}, \left\{ \frac{(1 - t^{(S_1-i)D_1 + \sum_{l=2}^k S_l D_l}) \Pi}{(1 - t^{\sum_{j=1}^k S_j D_j}) (1 - t^{\sum_{j=1}^k S_j N_j})} \right\}_{i=1}^{S_1}, \left\{ \frac{(1 - t^{iN_1}) \Pi}{(1 - t^{\sum_{j=1}^k S_j D_j}) (1 - t^{\sum_{j=1}^k S_j N_j})} \right\}_{i=1}^{S_1} \right) \\
&= \frac{\Pi}{(1 - t^{\sum_{j=1}^k S_j D_j}) (1 - t^{\sum_{j=1}^k S_j N_j})} \left(1 - t^{\sum_{j=1}^k S_j D_j}, \{1 - t^{(S_1-i)D_1 + \sum_{l=2}^k S_l D_l}\}_{i=1}^{S_1}, \{1 - t^{iN_1}\}_{i=1}^{S_1} \right) \\
&= \frac{\Pi}{(1 - t^{\sum_{j=1}^k S_j D_j}) (1 - t^{\sum_{j=1}^k S_j N_j})} \left(\left[\gcd \left(\sum_{j=1}^k S_j D_j, \left\{ (S_1 - i)D_1 + \sum_{l=2}^k S_l D_l \right\}_{i=1}^{S_1}, \{iN_1\}_{i=1}^{S_1} \right) \right] \right) \\
&= \frac{\Pi}{(1 - t^{\sum_{j=1}^k S_j D_j}) (1 - t^{\sum_{j=1}^k S_j N_j})} (1 - t)
\end{aligned}$$

where again we have assumed $S_1 \geq 2$. The Alexander polynomial is thus given by

$$\Delta_L(t) = (-1)^{-1 + \sum_{j=1}^k S_j} t^{\frac{\sigma}{2}} \frac{(1-t) \prod_{j=1}^k \left(1 - t^{S_j N_j D_j + N_j \sum_{l=j+1}^k S_l D_l + D_j \sum_{k=1}^{j-1} S_l N_l} \right)^{S_j}}{(1 - t^{\sum_{j=1}^k S_j D_j}) (1 - t^{\sum_{j=1}^k S_j N_j})}$$

where

$$\sigma = -1 + \sum_{j=1}^k S_j (N_j + D_j) - \sum_{j=1}^k S_j \left(S_j N_j D_j + N_j \sum_{l=j+1}^k S_l D_l + D_j \sum_{k=1}^{j-1} S_l N_l \right)$$

3. JONES POLYNOMIALS

Jones, using representation theory and properties of Temperley-Lieb algebra, computed the Jones polynomial of the torus knot $K_{p,q}$ to be

$$V_{K_{p,q}}(t) = t^{\frac{(p-1)(q-1)}{2}} \frac{1 - t^{p+1} - t^{q+1} + t^{p+q+1}}{1 - t^2}$$

for coprime integers p and q [6]. We now extend this to tst links. First we note that if K^* denotes the mirror image of the knot K , $V_{K^*}(t) = V_K(t^{-1})$, hence

$$V_{K_{-p,q}}(t) = V_{K_{p,q}}(t^{-1}) = t^{-\frac{(p-1)(q-1)}{2}} \frac{1 - t^{-p-1} - t^{-q-1} + t^{-p-q-1}}{1 - t^{-2}}$$

since $K_{-p,q}$ is the mirror image of $K_{p,q}$ [3].

3.1. Tst links $[\Phi^\mu(n(\tau), d(\tau), M)]$ with even M . We consider the case where M is even. Note that the link is the same as the torus link $K_{-\mu PN(\tau), \mu PD(\tau)}$. We can thus use the technique of topological quantum field theory. First recall that the unnormalized Jones polynomial $\langle L \rangle$ of a given link L is related to the vacuum expectation value of the Wilson lines in the $(2+1)$ -dimensional Chern-Simons theory as follows [2]. First the partition function over the 3-manifold M is given by

$$Z(M) = \int [DA] \exp \left(\frac{ik}{8\pi\hbar} \int_M \text{Tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right) \right)$$

where $S_{CS} = \frac{k}{8\pi} \int_M \text{Tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right)$ is called the level- k Chern-Simons action. The $\mu(L)$ components of L are taken as Wilson lines and the partition function of M with L inside it is given by

$$Z(M; L) = \int [DA] \exp \left(\frac{i}{\hbar} S_{CS} \right) \prod_{i=1}^{\mu(L)} \text{Tr}_R \mathcal{P} \exp \int_{L_i} A$$

where $L = \{L_1, L_2, \dots, L_{\mu(L)}\}$ contains the components L_i , $W_R(C) = \text{Tr}_R \mathcal{P} \exp \int_C A$ is called the Wilson operator, R is the 2-dimensional representation of $SU(2)$ in the case of the usual Jones polynomial, and \mathcal{P} is the path-ordering operator.

Then we can split the manifold M containing L into two parts M_1 and M_2 along a Riemann surface Σ such that $M = M_1 \cup_{\Sigma} M_2$ and no parts of L appear in Σ . Suppose L_1, L_2, \dots, L_n lie in M_1 and the remaining components lie in M_2 . Then we can define coordinates z and \bar{z} on the surface Σ , and the gauge connections A_z and $A_{\bar{z}}$. A_z and $A_{\bar{z}}$ satisfy a conjugate commutator relation. Choosing a metric γ on Σ , we also define the wave functional associated with M_1 to be

$$\zeta_1[A_{\bar{z}}] = \int [DA] \prod_{i=1}^n W_R(L_i) \exp \left(\frac{ik}{\hbar} S_{CS} - \frac{k}{2\pi\hbar} \int_{\Sigma} \frac{i}{2} dz d\bar{z} \sqrt{\gamma} \gamma^{z\bar{z}} \text{Tr}(A_z A_{\bar{z}}) \right)$$

The functional corresponding to M_2 is given by

$$\zeta_2[A_{\bar{z}}] = \int [DA] \prod_{i=n+1}^{\mu(L)} W_R(L_i) \exp \left(\frac{ik}{\hbar} S_{CS} - \frac{k}{2\pi\hbar} \int_{\Sigma} \frac{i}{2} dz d\bar{z} \sqrt{\gamma} \gamma^{z\bar{z}} \text{Tr}(A_z A_{\bar{z}}) \right)$$

Then the inner product defined by

$$\langle \zeta_1 | \zeta_2 \rangle = \int [DA_z DA_{\bar{z}}] \exp \left(\frac{k}{\pi\hbar} \int_{\Sigma} \frac{i}{2} dz d\bar{z} \sqrt{\gamma} \gamma^{z\bar{z}} \text{Tr}(A_z A_{\bar{z}}) \right) \overline{\zeta_1[A_{\bar{z}}]} \zeta_2[A_{\bar{z}}]$$

gives both the correct commutator relation and the partition function, i.e.

$$Z(M; L) = \langle \zeta_1 | \zeta_2 \rangle$$

Following [1], we consider the manifolds M_1, M_2 without the link components. Let the corresponding wave functionals be ζ'_1 and ζ'_2 respectively, then

$$\begin{aligned} Z(M) &= \langle \zeta'_1 | \zeta'_2 \rangle \\ Z(M_1; L_1, \dots, L_n) &= \langle \zeta'_1 | \zeta'_2 \rangle \\ Z(M_2; L_{n+1}, \dots, L_{\mu(L)}) &= \langle \zeta'_1 | \zeta'_2 \rangle \end{aligned}$$

We further note that

$$\begin{aligned}
Z(M; L)Z(M) &= \langle \zeta_1 | \zeta_2 \rangle \langle \zeta'_1 | \zeta'_2 \rangle \\
&= \int [DA_z DA_{\bar{z}}] \exp\left(\frac{k}{\pi\hbar} \int_{\Sigma} \frac{i}{2} dz d\bar{z} \sqrt{\gamma} \gamma^{z\bar{z}} \text{Tr}(A_z A_{\bar{z}})\right) \overline{\zeta_1[A_{\bar{z}}]} \zeta_2[A_{\bar{z}}] \\
&\quad \times \int [DA_z DA_{\bar{z}}] \exp\left(\frac{k}{\pi\hbar} \int_{\Sigma} \frac{i}{2} dz d\bar{z} \sqrt{\gamma} \gamma^{z\bar{z}} \text{Tr}(A_z A_{\bar{z}})\right) \overline{\zeta'_1[A_{\bar{z}}]} \zeta'_2[A_{\bar{z}}] \\
&= \int \int [DA_z DA_{\bar{z}}] \exp\left(\frac{k}{\pi\hbar} \int_{\Sigma} idz d\bar{z} \sqrt{\gamma} \gamma^{z\bar{z}} \text{Tr}(A_z A_{\bar{z}})\right) \overline{\zeta_1[A_{\bar{z}}]} \zeta_2[A_{\bar{z}}] \overline{\zeta'_1[A_{\bar{z}}]} \zeta'_2[A_{\bar{z}}] \\
&= \int \int [DA_z DA_{\bar{z}}] G^2 \overline{\zeta'_1[A_{\bar{z}}]} \zeta_2[A_{\bar{z}}] \overline{\zeta_1[A_{\bar{z}}]} \zeta'_2[A_{\bar{z}}] \\
&= \int [DA_z DA_{\bar{z}}] G \overline{\zeta'_1[A_{\bar{z}}]} \zeta_2[A_{\bar{z}}] \int [DA_z DA_{\bar{z}}] G \overline{\zeta_1[A_{\bar{z}}]} \zeta'_2[A_{\bar{z}}] \\
&= \langle \zeta'_1 | \zeta_2 \rangle \langle \zeta_1 | \zeta'_2 \rangle \\
&= Z(M_1; L_1, \dots, L_n) Z(M_2; L_{n+1}, \dots, L_{\mu(L)})
\end{aligned}$$

where $G = \exp\left(\frac{k}{\pi\hbar} \int_{\Sigma} \frac{i}{2} dz d\bar{z} \sqrt{\gamma} \gamma^{z\bar{z}} \text{Tr}(A_z A_{\bar{z}})\right)$. Rearranging gives

$$\frac{Z(M; L)}{Z(M)} = \frac{Z(M_1; L_1, \dots, L_n)}{Z(M)} \frac{Z(M_2; L_{n+1}, \dots, L_{\mu(L)})}{Z(M)}$$

Hence we are motivated to define the unnormalized Jones polynomial $\langle L \rangle$ of the link L to be

$$\langle L \rangle = \frac{Z(M; L)}{Z(M)}$$

so that

$$(3.1) \quad \langle L \rangle = \langle L_1, \dots, L_n \rangle \langle L_{n+1}, \dots, L_{\mu(L)} \rangle$$

Now the normalized Jones polynomial is given by

$$V_L(t) = \frac{\langle L \rangle}{\langle O \rangle}$$

where O denotes the unknot. From [1], we also have

$$\langle O \rangle = \frac{t - t^{-1}}{t^{\frac{1}{2}} - t^{-\frac{1}{2}}} = t^{-\frac{1}{2}} \frac{1 - t^2}{1 - t} = t^{\frac{1}{2}} \frac{1 - t^{-2}}{1 - t^{-1}}$$

To consider the case where $L = [\Phi^\mu(n(\tau), d(\tau), M)]$ with even M , we first compute $\langle K_{-N, D} \rangle$:

$$\begin{aligned}
\langle K_{-N, D} \rangle &= V_{K_{-N, D}}(t) t^{-\frac{1}{2}} \frac{1 - t^2}{1 - t} \\
&= t^{-\frac{(N-1)(D-1)}{2}} \frac{1 - t^{-N-1} - t^{-D-1} + t^{-N-D-1}}{1 - t^{-2}} t^{\frac{1}{2}} \frac{1 - t^{-2}}{1 - t^{-1}} \\
&= t^{\frac{1-(N-1)(D-1)}{2}} \frac{1 - t^{-N-1} - t^{-D-1} + t^{-N-D-1}}{1 - t^{-1}} \\
&= t^{\frac{N+D-ND}{2}} \frac{1 - t^{-N-1} - t^{-D-1} + t^{-N-D-1}}{1 - t^{-1}}
\end{aligned}$$

We then consider $(S-1)$ solid tori U_1, U_2, \dots, U_{S-1} that share a common longitudinal circle such that $U_{S-1} \subset \dots \subset U_2 \subset U_1$. We then consider another set of S solid

tori V_1, V_2, \dots, V_S that share the same longitudinal circle such that $\partial V_i \subset U_{i-1} \setminus U_i$ for $i \geq 2$ and $\partial V_1 \subset S^3 \setminus U_1$ and place each component $K_{-N,D}$ on ∂V_i . Then we can split $M = S^3$ into the spaces $M_1 = \overline{S^3 \setminus U_1}$ and $M_2 = \overline{U_1}$ so that $\Sigma = \partial U_1$ and $M = M_1 \cup_\Sigma M_2$, and apply equation (3.1) to give

$$\langle L \rangle = t^{\frac{N+D-ND}{2}} \frac{1 - t^{-N-1} - t^{-D-1} + t^{-N-D-1}}{1 - t^{-1}} \langle L' \rangle$$

where L' contains the remaining $(S-1)$ components. We then further split $\overline{U_1}$ into $\overline{U_2}$ and $\overline{U_1 \setminus U_2}$ and apply (3.1) again, until we are left with the last component. Thus we have

$$\begin{aligned} \langle K_{-SN,SD} \rangle &= \left(t^{\frac{N+D-ND}{2}} \frac{1 - t^{-N-1} - t^{-D-1} + t^{-N-D-1}}{1 - t^{-1}} \right)^S \\ &= t^{\frac{S(N+D-ND)}{2}} \left(\frac{1 - t^{-N-1} - t^{-D-1} + t^{-N-D-1}}{1 - t^{-1}} \right)^S \end{aligned}$$

The normalized Jones polynomial is given by

$$\begin{aligned} V_{[\Phi^\mu(n(\tau), d(\tau), M)]}(t) &= \frac{\langle K_{-SN,SD} \rangle}{\langle O \rangle} \\ &= \frac{t^{\frac{S(N+D-ND)}{2}} \left(\frac{1 - t^{-N-1} - t^{-D-1} + t^{-N-D-1}}{1 - t^{-1}} \right)^S}{t^{\frac{1}{2}} \frac{1 - t^{-2}}{1 - t^{-1}}} \\ &= t^{\frac{S(N+D-ND)-1}{2}} \frac{(1 - t^{-N-1} - t^{-D-1} + t^{-N-D-1})^S}{(1 - t^{-1})^S (1 + t^{-1})} \end{aligned}$$

3.2. Tst links $[\Phi^\mu(n(\tau), d(\tau), M)]$ with odd M . We now turn to the case where M is odd. We can simply set $M_2 = \overline{U}$ to be a solid torus containing the trivial knot and $M_1 = \overline{S^3 \setminus U}$ to contain the torus link. Then $\Sigma = \partial U$ and we can apply equation (3.1) to give

$$\langle [\Phi^\mu(n(\tau), d(\tau), M)] \rangle = \langle K_{-SN,SD} \rangle \langle O \rangle$$

The normalized Jones polynomial is given by

$$V_{[\Phi^\mu(n(\tau), d(\tau), M)]}(t) = \frac{\langle K_{-SN,SD} \rangle \langle O \rangle}{\langle O \rangle} = t^{\frac{S(N+D-ND)}{2}} \left(\frac{1 - t^{-N-1} - t^{-D-1} + t^{-N-D-1}}{1 - t^{-1}} \right)^S$$

3.3. Stst links $[\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j)]$ with even M_k . We now turn to the stst link $[\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j)]$. We write $N(\tau_j) = N_j$, $D(\tau_j) = D_j$ and $\mu_j P_j = S_j$. For even M_k , we consider k solid tori with the same longitudinal circle, $V_k \subset \dots \subset V_2 \subset V_1$ and another $(k-1)$ solid tori with also the same longitudinal circle, $U_{k-1} \subset \dots \subset U_2 \subset U_1$ and place ∂V_1 inside $\overline{S^3 \setminus U_1}$ and ∂V_i inside $\overline{U_{i-1} \setminus U_i}$ for $i \geq 2$. On each ∂V_j , we place the torus link $[\Phi^{\mu_j}(n(\tau_j), d(\tau_j), M_j)]$. We now

repeatedly apply equation(3.1) so that

$$\begin{aligned}
 V_{[\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j)]}(t) &= \frac{\prod_{j=1}^k t^{\frac{S_j(N_j+D_j-N_j D_j)}{2}} \left(\frac{1-t^{-N_j-1}-t^{-D_j-1}+t^{-N_j-D_j-1}}{1-t^{-1}} \right)^{S_j}}{t^{\frac{1}{2}} \frac{1-t^{-2}}{1-t^{-1}}} \\
 &= t^{\frac{-1+\sum_{j=1}^k S_j(N_j+D_j-N_j D_j)}{2}} \frac{\prod_{j=1}^k (1-t^{-N_j-1}-t^{-D_j-1}+t^{-N_j-D_j-1})^{S_j}}{(1-t^{-1})^{\sum_{j=1}^k S_j} (1+t^{-1})}
 \end{aligned}$$

3.4. Stst links $[\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j)]$ with odd M_k . We now turn to the case where M_k is odd. This is similar to the case where M_k is even, but we introduce one more solid torus U_k that shares the same longitudinal circle with other U 's such that the trivial knot is placed inside U_k . We can now apply equation (3.1) so that

$$V_{[\Phi_{j=1}^k(n(\tau_j), d(\tau_j), M_j, \mu_j)]}(t) = t^{\frac{\sum_{j=1}^k S_j(N_j+D_j-N_j D_j)}{2}} \frac{\prod_{j=1}^k (1-t^{-N_j-1}-t^{-D_j-1}+t^{-N_j-D_j-1})^{S_j}}{(1-t^{-1})^{\sum_{j=1}^k S_j}}$$

CONCLUSIONS

In the paper we introduced stst links, computed their braid words, Alexander and Jones polynomials. Further research directions include finding other knot or link invariants such as unknotting numbers, hyperbolic volumes, HOMFLY polynomials, Kontsevich integrals and other Vassiliev invariants. Further generalizations also include satellite links built from stst links, and virtual knots. These could be investigated in the future.

REFERENCES

- [1] Edward Witten. "Quantum field theory and the Jones polynomial" In: *Communications in Mathematical Physics*. 121.3 (1989), pp.351-399.
- [2] J.M.F. Labastida, P.M. Llatas and A.V. Ramallo. "Knot operators in Chern-Simons Gauge Theory". In: *Nuclear Physics B*. 348.3 (1991), pp. 651-692.
- [3] Kunia Murasugi. "Torus Knots". In: *Knot Theory and Its Applications*. Trans. by Bohdan Kurpita. Boston: Birkäuser, 1996. Chap. 7, pp. 132-151.
- [4] Julia Collins. *The Alexander Polynomial: The woefully overlooked granddaddy of knot polynomials*. 2007. URL: <http://infohost.nmt.edu/~starrett/Fall12014/TopologyOfChaos/KnotTheory/GeomClub.pdf>.
- [5] Kai Lai Chan, Tsz Nam Chan, Ho Yin Lau, Kai Shing Mok and Yiu Shing Wong. "Cutting Twisted Solid Tori (TSTs)". In: *Hong Kong Mathematics Awards Collection of Winning Papers 2012*. Ed. by Thomas Au and Kwok Wai Chan. CUHK: Institute of Mathematical Sciences, 2012, pp. 25-174.
- [6] Vaughan Jones. *The Jones polynomial for dummies*. 2014. URL: <https://math.berkeley.edu/~vfr/jonesakl.pdf>.

EMAIL: WILSONWYS@YAHOO.COM.HK