

# The invariance and formulas for solutions of some fifth-order difference equations

D. Nyirenda and M. Folly-Gbetoula\*

School of Mathematics, University of the Witwatersrand, 2050,  
Johannesburg, South Africa.

## Abstract

Lie group analysis of the difference equations of the form

$$x_{n+1} = \frac{x_{n-4}x_{n-3}}{x_n(a_n + b_n x_{n-4}x_{n-3}x_{n-2}x_{n-1})},$$

where  $a_n$  and  $b_n$  are real sequences, is performed and non-trivial symmetries are derived. Furthermore, we find formulas for exact solutions of the equations. This work generalizes a recent result by Elsayed [Elsayed, E.M.: Expression and behavior of the solutions of some rational recursive sequences. *Math. Meth. Appl. Sci.* **2016:39**,5682–5694 (2016)].

**Keywords** Difference equation; symmetry; reduction; group invariant solutions

**Mathematics Subject Classification** 39A10

## 1 Introduction

In recent years, following the work of Sophus Lie [8] on differential equations, various researchers showed interest in symmetries. Lie investigated the group of transformations which leaves the differential equations invariant. The idea of symmetry is also connected to conservation laws and this connection between the two areas has led to greater motivation in researchers, after the work of Noether [15]. It is known that so long as the symmetries and first integrals are related via the invariance condition, one can implement the double reduction of the differential equations [14, 17]. The notion of

---

\*Corresponding author: Mensah.Folly-Gbetoula@wits.ac.za

using symmetries has had its extension to difference equations thanks to Maeda [9, 10]. On symmetries in difference equations, refer to [2–5, 7, 11, 16]. Hydon [4] established a symmetry based algorithm that makes solution finding possible. Despite the fact that Hydon [5] emphasized on lower-order difference equations, his procedure works for any order. However, for higher-order equations, computations are cumbersome as such certain assumptions are put in order to lessen the burden of computation.

In this paper, we are inspired by the work of Elsayed [1], who studied the following recursive sequences:

$$x_{n+1} = \frac{x_{n-3}x_{n-4}}{x_n(\pm 1 \pm x_{n-1}x_{n-2}x_{n-3}x_{n-4})}, \quad n = 0, 1, \dots, \quad (1)$$

where the initial conditions are arbitrary real numbers. Clearly, (1) are special cases of a more general form

$$x_{n+1} = \frac{x_{n-3}x_{n-4}}{x_n(a_n + b_nx_{n-1}x_{n-2}x_{n-3}x_{n-4})}, \quad n = 0, 1, \dots, \quad (2)$$

where  $(a_n)$  and  $(b_n)$  are real sequences. Our aim is to utilize symmetry methods to solve this more general difference equation (2). Equivalently, we study the forward difference equation

$$u_{n+5} = \frac{u_n u_{n+1}}{u_{n+4}(A_n + B_n u_n u_{n+1} u_{n+2} u_{n+3})}, \quad (3)$$

since we follow the notation of [4].

## 2 Preliminaries

This section provides background to difference equations in the context of Lie symmetry analysis.

**Definition 2.1** *Let  $G$  be a local group of transformations acting on a manifold  $M$ . A subset  $\mathcal{S} \subset M$  is called  $G$ -invariant, and  $G$  is called symmetry group of  $\mathcal{S}$ , if whenever  $x \in \mathcal{S}$ , and  $g \in G$  is such that  $g \cdot x$  is defined, then  $g \cdot x \in \mathcal{S}$ .*

**Definition 2.2** *Let  $G$  be a connected group of transformations acting on a manifold  $M$ . A smooth real-valued function  $\mathcal{V} : M \rightarrow \mathbb{R}$  is an invariant function for  $G$  if and only if*

$$X(\mathcal{V}) = 0 \quad \text{for all} \quad x \in M,$$

and every infinitesimal generator  $X$  of  $G$ .

**Definition 2.3** A parameterized set of point transformations,

$$\Gamma_\varepsilon : x \mapsto \hat{x}(x; \varepsilon), \quad (4)$$

where  $x = x_i$ ,  $i = 1, \dots, p$  are continuous variables, is a one-parameter local Lie group of transformations if the following conditions are satisfied:

1.  $\Gamma_0$  is the identity map if  $\hat{x} = x$  when  $\varepsilon = 0$
2.  $\Gamma_a \Gamma_b = \Gamma_{a+b}$  for every  $a$  and  $b$  sufficiently close to 0
3. Each  $\hat{x}_i$  can be represented as a Taylor series (in a neighborhood of  $\varepsilon = 0$  that is determined by  $x$ ), and therefore

$$\hat{x}_i(x; \varepsilon) = x_i + \varepsilon \xi_i(x) + O(\varepsilon^2), \quad i = 1, \dots, p. \quad (5)$$

Assume that the forward  $r$ th-order difference equation takes the form

$$u_{n+r} = F(n, u_n, u_{n+1}, \dots, u_{n+r-1}), \quad n \in D \quad (6)$$

for some smooth function  $F$  and a regular domain  $D \subset \mathbb{Z}$ . So as to compute a symmetry group of (6), we take into consideration the group of point transformations given as

$$\hat{n} = n, \quad (7a)$$

$$\hat{u}_n = u_n + \epsilon Q(n, u_n) + O(\epsilon^2), \quad (7b)$$

$$\hat{u}_{n+j} = u_{n+j} + \epsilon S^j Q(n, u_n) + O(\epsilon^2), \quad (7c)$$

where  $\epsilon$  ( $\epsilon$  is sufficiently small) is the parameter,  $Q = Q(n, u_n)$  is a continuous function, referred to as characteristic and  $S$  is the shift operator defined as  $S : n \mapsto n + 1$ . The criterion of invariance is then

$$\hat{u}_{n+r} = F(\hat{n}, \hat{u}_n, \hat{u}_{n+1}, \dots, \hat{u}_{n+r-1}), \quad (8)$$

which yields the linearized symmetry condition

$$S^r Q - XF = 0 \quad (9)$$

by substituting (7) in (8). Observe that

$$X = Q(n, u_n) \frac{\partial}{\partial u_n} + SQ(n, u_n) \frac{\partial}{\partial u_{n+1}} + \cdots + S^{r-1}Q(n, u_n) \frac{\partial}{\partial u_{n+r-1}} \quad (10)$$

is the corresponding ‘prolonged’ infinitesimal of the group of transformations (7). Upon knowledge of the function(s)  $Q$ , one is able to obtain the invariant  $\mathcal{V}$  by using the canonical coordinate [6]

$$S_n = \int \frac{du_n}{Q(n, u_n)}. \quad (11)$$

Generally, the steps involves are lengthy even though very exact and do not give room to guess work on the perfect choice of invariants.

For more understanding on Lie analysis of differential and difference equations, see [4, 12].

### 3 Main results

We are studying the equation

$$u_{n+5} = F = \frac{u_n u_{n+1}}{u_{n+4}(A_n + B_n u_n u_{n+1} u_{n+2} u_{n+3})}. \quad (12)$$

Applying condition (9) to (12), we get

$$\begin{aligned} Q(n+5, F) + \frac{A_n u_n u_{n+1} Q(n+4, u_{n+4})}{u_{n+4}(A_n + B_n u_n u_{n+1} u_{n+2} u_{n+3})^2} + \\ \frac{B_n u_n^2 u_{n+1}^2 u_{n+2} u_{n+3} Q(n+4, u_{n+4})}{u_{n+4}^2 (A_n + B_n u_n u_{n+1} u_{n+2} u_{n+3})^2} + \frac{B_n u_n^2 u_{n+1}^2 u_{n+2} Q(n+3, u_{n+3})}{u_{n+4}(A_n + B_n u_n u_{n+1} u_{n+2} u_{n+3})^2} \\ + \frac{B_n u_n^2 u_{n+1}^2 u_{n+3} Q(n+2, u_{n+2})}{u_{n+4}(A_n + B_n u_n u_{n+1} u_{n+2} u_{n+3})^2} - \frac{A_n u_{n+1} Q(n, u_n)}{u_{n+4}(A_n + B_n u_n u_{n+1} u_{n+2} u_{n+3})^2} = 0. \end{aligned} \quad (13)$$

Eliminating  $F$  is achieved by applying implicit differentiation with respect to  $u_n$  (regarding  $u_{n+4}$  as a function of  $u_n, u_{n+1}, u_{n+2}, u_{n+3}$  and  $u_{n+5}$ ) via the differential operator

$$L = \frac{\partial}{\partial u_n} + \frac{\partial u_{n+4}}{\partial u_n} \frac{\partial}{\partial u_{n+4}} = \frac{\partial}{\partial u_n} - \left[ \left( \frac{\partial F}{\partial u_n} \right) / \left( \frac{\partial F}{\partial u_{n+4}} \right) \right] \frac{\partial}{\partial u_{n+4}}.$$

With some simplification, we get

$$\begin{aligned}
& (A_n + B_n u_n u_{n+1} u_{n+2} u_{n+3}) Q'(n+4, u_{n+4}) + B_n u_n u_{n+1} u_{n+2} Q(n+3, u_{n+3}) + \\
& B_n u_n u_{n+1} u_{n+3} Q(n+2, u_{n+2}) + B_n u_n u_{n+2} u_{n+3} Q(n+2, u_{n+2}) + \\
& B_n u_n u_{n+2} u_{n+3} Q(n+1, u_{n+1}) - (A_n + B_n u_n u_{n+1} u_{n+2} u_{n+3}) Q'(n, u_n) + \\
& 2B_n u_{n+1} u_{n+2} u_{n+3} Q(n, u_n) + \frac{A_n}{u_n} Q(n, u_n) = 0. \tag{14}
\end{aligned}$$

The symbol ' denotes the derivative with respect to the continuous variable. Differentiating (14) with respect to  $u_n$  twice, keeping  $u_{n+4}$  constant, yields

$$\begin{aligned}
& -B_n u_n u_{n+1} u_{n+2} u_{n+3} Q'''(n, u_n) - A_n Q'''(n, u_n) + \frac{A_n}{u_n} Q''(n, u_n) \\
& - \frac{2A_n}{u_n^2} Q'(n, u_n) + \frac{2A_n}{u_n^3} Q(n, u_n) = 0. \tag{15}
\end{aligned}$$

The characteristic in (15) is a function of  $u_n$  only and thus we split (15) to get the system

$$1 : Q'''(n, u_n) - \frac{1}{u_n} Q''(n, u_n) + \frac{2}{u_n^2} Q'(n, u_n) - \frac{2}{u_n^3} Q(n, u_n) = 0 \tag{16a}$$

$$u_{n+1} u_{n+2} u_{n+3} : Q'''(n, u_n) = 0. \tag{16b}$$

One obtains the solution to (16) as

$$Q(n, u_n) = \alpha_n u_n^2 + \beta_n u_n \tag{17}$$

for some arbitrary functions  $\alpha_n$  and  $\beta_n$  of  $n$ . Substituting (17) and its shifts in (13), and making a replacement of the expression of  $u_{n+5}$  given in (12) in the resulting equation leads to

$$\begin{aligned}
& B_n u_n u_{n+1} u_{n+2} u_{n+3}^2 u_{n+4} \alpha_{n+3} + B_n u_n u_{n+1} u_{n+2} u_{n+3} u_{n+4}^2 \alpha_{n+4} + \\
& B_n u_n u_{n+1} u_{n+2}^2 u_{n+3} u_{n+4} \alpha_{n+2} + B_n u_n u_{n+1} u_{n+2} u_{n+3} u_{n+4} (\beta_{n+2} + \beta_{n+3} + \\
& \beta_{n+4} + \beta_{n+5}) + A_n u_{n+4}^2 \alpha_{n+4} - A_n u_n u_{n+4} \alpha_n - A_n u_{n+1} u_{n+4} \alpha_{n+1} - \\
& A_n u_{n+4} (\beta_n + \beta_{n+1} - \beta_{n+4} - \beta_{n+5}) + u_n u_{n+1} \alpha_{n+5} = 0. \tag{18}
\end{aligned}$$

Now equate coefficients of all powers of shifts of  $u_n$  to zero, i.e.,

$$u_n u_{n+1} u_{n+2} u_{n+3}^2 u_{n+4} : \alpha_{n+3} = 0 \quad (19)$$

$$u_n u_{n+1} u_{n+2} u_{n+3} u_{n+4}^2 : \alpha_{n+4} = 0 \quad (20)$$

$$u_n u_{n+1} u_{n+2}^2 u_{n+3} u_{n+4} : \alpha_{n+2} = 0 \quad (21)$$

$$u_n u_{n+1} u_{n+2} u_{n+3} u_{n+4} : (\beta_{n+2} + \beta_{n+3} + \beta_{n+4} + \beta_{n+5}) = 0 \quad (22)$$

$$u_{n+4}^2 : \alpha_{n+4} = 0 \quad (23)$$

$$u_n u_{n+4} : \alpha_n = 0 \quad (24)$$

$$u_{n+1} u_{n+4} : \alpha_{n+1} = 0 \quad (25)$$

$$u_{n+4} : (\beta_n + \beta_{n+1} - \beta_{n+4} - \beta_{n+5}) = 0 \quad (26)$$

$$u_n u_{n+1} : \alpha_{n+5} = 0. \quad (27)$$

So the system above is reduced to

$$\alpha_n = 0, \quad (28)$$

$$\beta_n + \beta_{n+1} + \beta_{n+2} + \beta_{n+3} = 0. \quad (29)$$

The three independent solutions of the linear third-order difference equation (29) are given by

$$(-1)^n, \quad \beta^n \quad \text{and} \quad \bar{\beta}^n \quad (30)$$

where  $\beta = \exp\{i\pi/2\}$  and  $\bar{\beta}$  denotes its complex conjugate. The characteristics are then given by

$$Q_1(n, u_n) = (-1)^n u_n, \quad Q_2(n, u_n) = \beta^n u_n \quad \text{and} \quad Q_3(n, u_n) = \bar{\beta}^n u_n, \quad (31)$$

and therefore, the symmetry operators admitted by (12) are given by

$$\begin{aligned} X_1 &= \sum_{j=0}^4 (-1)^{n+j} u_{n+j} \frac{\partial}{\partial u_{n+j}}, \quad X_2 = \sum_{j=0}^4 \beta^{n+j} u_{n+j} \frac{\partial}{\partial u_{n+j}}, \\ X_3 &= \sum_{j=0}^4 \bar{\beta}^{n+j} u_{n+j} \frac{\partial}{\partial u_{n+j}}. \end{aligned} \quad (32)$$

One can choose any one of the characteristics to write the canonical coordinate. We select  $Q_2$ . Thus

$$S_n = \int \frac{du_n}{Q_2(n, u_n)} = \int \frac{du_n}{\beta^n u_n} = \frac{1}{\beta^n} \ln |u_n| \quad (33)$$

and we use relation (29) to derive the invariant function  $\tilde{V}_n$  as follows:

$$\tilde{V}_n = S_n \beta^n + S_{n+1} \beta^{n+1} + S_{n+2} \beta^{n+2} + S_{n+3} \beta^{n+3}. \quad (34)$$

Actually,

$$X_1(\tilde{V}_n) = (-1)^n + (-1)^{n+1} + (-1)^{n+2} + (-1)^{n+3} = 0, \quad (35)$$

$$X_2(\tilde{V}_n) = \beta^n + \beta^{n+1} + \beta^{n+2} + \beta^{n+3} = 0 \quad (36)$$

and

$$X_3(\tilde{V}_n) = \bar{\beta}^n + \bar{\beta}^{n+1} + \bar{\beta}^{n+2} + \bar{\beta}^{n+3} = 0. \quad (37)$$

For the sake of simplicity, we utilize

$$|V_n| = \exp\{-\tilde{V}_n\} \quad (38)$$

instead. In other words,  $V_n = \pm 1/(u_n u_{n+1} u_{n+2} u_{n+3})$ . One can show via (12) and (38) that

$$V_{n+2} = A_n V_n \pm B_n. \quad (39)$$

By utilizing the plus sign (one is allowed to choose), the solution of (39) can be presented in closed form as follows:

$$V_{2n+j} = V_j \left( \prod_{k_1=0}^{n-1} A_{2k_1+j} \right) + \sum_{l=0}^{n-1} \left( B_{2l+j} \prod_{k_2=l+1}^{n-1} A_{2k_2} \right), \quad j = 0, 1. \quad (40)$$

From the above equation, obtaining the solution of (12) is easier. We first use (33) to get

$$|u_n| = \exp(\beta_n S_n). \quad (41)$$

Secondly, we use (34) to get

$$\begin{aligned} |u_n| = \exp & \left[ \beta^n c_1 + \bar{\beta}^n c_2 + (-1)^n c_3 - \left( \frac{1}{4} - \frac{i}{4} \right) \sum_{k_1=0}^{n-1} \beta^n \bar{\beta}^{k_1} |\tilde{V}_{k_1}| \right. \\ & \left. - \left( \frac{1}{4} + \frac{i}{4} \right) \sum_{k_2=0}^{n-1} \bar{\beta}^n \beta^{k_2} |\tilde{V}_{k_2}| - \frac{1}{2} \sum_{k_3=0}^{n-1} (-1)^{n-k_3} |\tilde{V}_{k_3}| \right]. \quad (42) \end{aligned}$$

Finally, invoking (38) yields

$$\begin{aligned}
|u_n| &= \exp \left[ \beta^n c_1 + \bar{\beta}^n c_2 + (-1)^n c_3 + \left( \frac{1}{4} - \frac{i}{4} \right) \sum_{k_1=0}^{n-1} \beta^n \bar{\beta}^{k_1} \ln |V_{k_1}| \right. \\
&\quad \left. + \left( \frac{1}{4} + \frac{i}{4} \right) \sum_{k_2=0}^{n-1} \bar{\beta}^n \beta^{k_2} \ln |V_{k_2}| + \frac{1}{2} \sum_{k_3=0}^{n-1} (-1)^{n-k_3} \ln |V_{k_3}| \right] \\
&= \exp \left( H_n + \frac{1}{2} \sum_{k=0}^{n-1} \left[ \sqrt{2} \cos \left( \frac{\pi(2k-2n+1)}{4} \right) + (-1)^{k-n} \right] \ln |V_k| \right),
\end{aligned} \tag{43}$$

where  $H_n = \beta^n c_1 + \bar{\beta}^n c_2 + (-1)^n c_3$ . Replacing  $n$  with  $4n+j$  for  $j = 0, 1, 2, 3$  yields

$$|u_{4n+j}| = \exp \left[ H_j + \frac{1}{2} \sum_{k=0}^{n-1} \left( \sqrt{2} \cos \left( \frac{\pi(2k-2j+1)}{4} \right) + (-1)^{k-j} \right) \ln |V_k| \right]. \tag{44}$$

Set  $j = 0$  in (44) to get

$$|u_{4n}| = \exp(H_0) \prod_{s=0}^{n-1} \left| \frac{V_{4s}}{V_{4s+1}} \right|.$$

But substituting  $n = 0$  in (43) leads to  $|u_0| = \exp(H_0)$ . Furthermore, using (12) and (38), it can be shown that there is no need of absolute values. Hence

$$\begin{aligned}
u_{4n} &= u_0 \prod_{s=0}^{n-1} \frac{V_{4s}}{V_{4s+1}} \\
&= u_0 \prod_{s=0}^{n-1} \frac{V_0 \left( \prod_{k_1=0}^{2s-1} A_{2k_1} \right) + \sum_{l=0}^{2s-1} \left( B_{2l} \prod_{k_2=l+1}^{2s-1} A_{2k_2} \right)}{V_1 \left( \prod_{k_1=0}^{2s-1} A_{2k_1+1} \right) + \sum_{l=0}^{2s-1} \left( B_{2l+1} \prod_{k_2=l+1}^{2s-1} A_{2k_2} \right)} \\
&= \frac{u_4}{u_0^{n-1}} \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{2s-1} A_{2k_1} \right) + u_0 u_1 u_2 u_3 \sum_{l=0}^{2s-1} \left( B_{2l} \prod_{k_2=l+1}^{2s-1} A_{2k_2} \right)}{\left( \prod_{k_1=0}^{2s-1} A_{2k_1+1} \right) + u_1 u_2 u_3 u_4 \sum_{l=0}^{2s-1} \left( B_{2l+1} \prod_{k_2=l+1}^{2s-1} A_{2k_2+1} \right)}.
\end{aligned}$$



For  $j = 1$ , we find that

$$u_{4n+1} = u_1 \prod_{s=0}^{n-1} \frac{V_{4s+1}}{V_{4s+2}}$$

so that

$$\begin{aligned} u_{4n+1} &= u_1 \prod_{s=0}^{n-1} \frac{V_1 \left( \prod_{k_1=0}^{2s-1} A_{2k_1+1} \right) + \sum_{l=0}^{n-1} \left( B_{2l+1} \prod_{k_2=l+1}^{2s-1} A_{2k_2+1} \right)}{V_0 \left( \prod_{k_1=0}^{2s} A_{2k_1} \right) + \sum_{l=0}^{2s} \left( B_{2l} \prod_{k_2=l+1}^{2s} A_{2k_2} \right)} \\ &= \frac{u_0^n u_1}{u_4^n} \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{2s-1} A_{2k_1+1} \right) + u_1 u_2 u_3 u_4 \sum_{l=0}^{2s-1} \left( B_{2l+1} \prod_{k_2=l+1}^{2s-1} A_{2k_2+1} \right)}{\left( \prod_{k_1=0}^{2s} A_{2k_1} \right) + u_0 u_1 u_2 u_3 \sum_{l=0}^{2s} \left( B_{2l} \prod_{k_2=l+1}^{2s} A_{2k_2} \right)}. \end{aligned}$$

For  $j = 2$ , we have

$$u_{4n+2} = u_2 \prod_{s=0}^{n-1} \frac{V_{4s+2}}{V_{4s+3}}$$

which evaluates to

$$\begin{aligned} u_{4n+2} &= u_2 \prod_{s=0}^{n-1} \frac{V_0 \left( \prod_{k_1=0}^{2s} A_{2k_1} \right) + \sum_{l=0}^{2s} \left( B_{2l} \prod_{k_2=l+1}^{2s} A_{2k_2} \right)}{V_1 \left( \prod_{k_1=0}^{2s} A_{2k_1+1} \right) + \sum_{l=0}^{2s} \left( B_{2l+1} \prod_{k_2=l+1}^{2s} A_{2k_2+1} \right)} \\ &= \frac{u_4^n u_2}{u_0^n} \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{2s} A_{2k_1} \right) + u_0 u_1 u_2 u_3 \sum_{l=0}^{2s} \left( B_{2l} \prod_{k_2=l+1}^{2s} A_{2k_2} \right)}{\left( \prod_{k_1=0}^{2s} A_{2k_1+1} \right) + u_1 u_2 u_3 u_4 \sum_{l=0}^{2s} \left( B_{2l+1} \prod_{k_2=l+1}^{2s} A_{2k_2+1} \right)}. \end{aligned}$$

Finally, for  $j = 3$ , we obtain

$$u_{4n+3} = u_3 \prod_{s=0}^{n-1} \frac{V_{4s+3}}{V_{4s+4}}$$

so that

$$\begin{aligned}
u_{4n+3} &= u_3 \prod_{s=0}^{n-1} \frac{V_1 \left( \prod_{k_1=0}^{2s} A_{2k_1+1} \right) + \sum_{l=0}^{2s} \left( B_{2l+1} \prod_{k_2=l+1}^{2s} A_{2k_2+1} \right)}{V_0 \left( \prod_{k_1=0}^{2s+1} A_{2k_1} \right) + \sum_{l=0}^{2s+1} \left( B_{2l} \prod_{k_2=l+1}^{2s+1} A_{2k_2} \right)} \\
&= \frac{u_0^n u_3}{u_4^n} \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{2s} A_{2k_1+1} \right) + u_1 u_2 u_3 u_4 \sum_{l=0}^{2s} \left( B_{2l+1} \prod_{k_2=l+1}^{2s} A_{2k_2+1} \right)}{\left( \prod_{k_1=0}^{2s+1} A_{2k_1} \right) + u_0 u_1 u_2 u_3 \sum_{l=0}^{2s+1} \left( B_{2l} \prod_{k_2=l+1}^{2s+1} A_{2k_2} \right)}.
\end{aligned}$$

Hence the solution to (2) is given by

$$x_{4n-4} = \frac{x_0^n}{x_{-4}^{n-1}} \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{2s-1} a_{2k_1} \right) + x_{-4} x_{-3} x_{-2} x_{-1} \sum_{l=0}^{2s-1} \left( b_{2l} \prod_{k_2=l+1}^{2s-1} a_{2k_2} \right)}{\left( \prod_{k_1=0}^{2s-1} a_{2k_1+1} \right) + x_{-3} x_{-2} x_{-1} x_0 \sum_{l=0}^{2s-1} \left( b_{2l+1} \prod_{k_2=l+1}^{2s-1} a_{2k_2+1} \right)},$$

which can be rearranged as

$$x_{4n} = \frac{x_0^{n+1}}{x_{-4}^n} \prod_{s=0}^n \frac{\left( \prod_{k_1=0}^{2s-1} a_{2k_1} \right) + x_{-4} x_{-3} x_{-2} x_{-1} \sum_{l=0}^{2s-1} \left( b_{2l} \prod_{k_2=l+1}^{2s-1} a_{2k_2} \right)}{\left( \prod_{k_1=0}^{2s-1} a_{2k_1+1} \right) + x_{-3} x_{-2} x_{-1} x_0 \sum_{l=0}^{2s-1} \left( b_{2l+1} \prod_{k_2=l+1}^{2s-1} a_{2k_2+1} \right)}.$$

The term  $s = 0$  in the product (indexed by  $s$ ) is equal to 1 using the facts that  $\sum_{i=0}^{-1} a_i = 0$  and  $\prod_{j=0}^{-1} a_j = 1$ . As a result, we can still rewrite the solution as

$$x_{4n} = \frac{x_0^{n+1}}{x_{-4}^n} \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{2s+1} a_{2k_1} \right) + x_{-4} x_{-3} x_{-2} x_{-1} \sum_{l=0}^{2s+1} \left( b_{2l} \prod_{k_2=l+1}^{2s+1} a_{2k_2} \right)}{\left( \prod_{k_1=0}^{2s+1} a_{2k_1+1} \right) + x_{-3} x_{-2} x_{-1} x_0 \sum_{l=0}^{2s+1} \left( b_{2l+1} \prod_{k_2=l+1}^{2s+1} a_{2k_2+1} \right)}. \tag{45a}$$

Furthermore, observe that

$$x_{4n-3} = \frac{x_{-4}^n x_{-3}}{x_0^n} \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{2s-1} a_{2k_1+1} \right) + x_{-3} x_{-2} x_{-1} x_0 \sum_{l=0}^{2s-1} \left( b_{2l+1} \prod_{k_2=l+1}^{2s-1} a_{2k_2+1} \right)}{\left( \prod_{k_1=0}^{2s} a_{2k_1} \right) + x_{-4} x_{-3} x_{-2} x_{-1} \sum_{l=0}^{2s} \left( b_{2l} \prod_{k_2=l+1}^{2s} a_{2k_2} \right)}, \quad (45b)$$

$$x_{4n-2} = \frac{x_0^n x_{-2}}{x_{-4}^n} \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{2s} a_{2k_1} \right) + x_{-4} x_{-3} x_{-2} x_{-1} \sum_{l=0}^{2s} \left( b_{2l} \prod_{k_2=l+1}^{2s} a_{2k_2} \right)}{\left( \prod_{k_1=0}^{2s} a_{2k_1+1} \right) + x_{-3} x_{-2} x_{-1} x_0 \sum_{l=0}^{2s} \left( b_{2l+1} \prod_{k_2=l+1}^{2s} a_{2k_2+1} \right)} \quad (45c)$$

and

$$x_{4n-1} = \frac{x_{-4}^n x_{-1}}{x_0^n} \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{2s} a_{2k_1+1} \right) + x_{-3} x_{-2} x_{-1} x_0 \sum_{l=0}^{2s} \left( b_{2l+1} \prod_{k_2=l+1}^{2s} a_{2k_2+1} \right)}{\left( \prod_{k_1=0}^{2s+1} a_{2k_1} \right) + x_{-4} x_{-3} x_{-2} x_{-1} \sum_{l=0}^{2s+1} \left( b_{2l} \prod_{k_2=l+1}^{2s+1} a_{2k_2} \right)} \quad (45d)$$

as long as any of the denominators does not vanish.

In the following sections, we look at some special cases.

## 4 The case $a_n, b_n$ are 1-periodic

In this case  $a_n = a$  and  $b_n = b$  where  $a, b \in \mathbb{R}$ .

### 4.1 The case $a \neq 1$

From (45), the solution is given by

$$x_{4n} = \frac{x_0^{n+1}}{x_{-4}^n} \prod_{s=0}^{n-1} \frac{a^{2s+2} + b x_{-4} x_{-3} x_{-2} x_{-1} \frac{1-a^{2s+2}}{1-a}}{a^{2s+2} + b x_{-3} x_{-2} x_{-1} x_0 \frac{1-a^{2s+2}}{1-a}}, \quad (46a)$$

$$x_{4n-3} = \frac{x_{-4}^n x_{-3}}{x_0^n} \prod_{s=0}^{n-1} \frac{a^{2s} + b x_{-3} x_{-2} x_{-1} x_0 \frac{1-a^{2s}}{1-a}}{a^{2s+1} + b x_{-4} x_{-3} x_{-2} x_{-1} \frac{1-a^{2s+1}}{1-a}}, \quad (46b)$$

$$x_{4n-2} = \frac{x_0^n x_{-2}}{x_{-4}^n} \prod_{s=0}^{n-1} \frac{a^{2s+1} + bx_{-4}x_{-3}x_{-2}x_{-1} \frac{1-a^{2s+1}}{1-a}}{a^{2s+1} + bx_{-3}x_{-2}x_{-1}x_0 \frac{1-a^{2s+1}}{1-a}} \quad (46c)$$

and

$$x_{4n-1} = \frac{x_{-4}^n x_{-1}}{x_0^n} \prod_{s=0}^{n-1} \frac{a^{2s+1} + bx_{-3}x_{-2}x_{-1}x_0 \frac{1-a^{2s+1}}{1-a}}{a^{2s+2} + bx_{-4}x_{-3}x_{-2}x_{-1} \frac{1-a^{2s+2}}{1-a}} \quad (46d)$$

where  $x_{-4}, x_0 \neq 0$  and for all  $(i, s) \in \{0, 1\} \times \{0, 1, 2, 3, \dots, n-1\}$ ,

$$(1-a)a^{2s+i} + (1-a^{2s+i})bx_{-3}x_{-2}x_{-1}x_0 \neq 0$$

and

$$(1-a)a^{2s+1+i} + (1-a^{2s+1+i})bx_{-4}x_{-3}x_{-2}x_{-1} \neq 0.$$

#### 4.1.1 The case $a = -1$

In this case, the solution which for  $b = \pm 1$  appears in [1] (see Theorems 3 and 8), is given by

$$x_{4n} = \frac{x_0^{n+1}}{x_{-4}^n}, \quad (47a)$$

$$x_{4n-3} = \frac{x_{-4}^n x_{-3}}{x_0^n} (-1 + bx_{-4}x_{-3}x_{-2}x_{-1})^{-n}, \quad (47b)$$

$$x_{4n-2} = \frac{x_0^n x_{-2}}{x_{-4}^n} \left( \frac{-1 + bx_{-4}x_{-3}x_{-2}x_{-1}}{-1 + bx_{-3}x_{-2}x_{-1}x_0} \right)^n \quad (47c)$$

and

$$x_{4n-1} = \frac{x_{-4}^n x_{-1}}{x_0^n} (-1 + bx_{-3}x_{-2}x_{-1}x_0)^n \quad (47d)$$

where  $x_{-4}, x_0 \neq 0$ ,  $bx_{-4}x_{-3}x_{-2}x_{-1} \neq 1$  and  $bx_{-3}x_{-2}x_{-1}x_0 \neq 1$ .

## 4.2 The case $a = 1$

From (45), the solution, which for  $b = \pm 1$  appears in [1] (see Theorems 1 and 6), is given by

$$x_{4n} = \frac{x_0^{n+1}}{x_{-4}^n} \prod_{s=0}^{n-1} \frac{1 + (2s+2)bx_{-4}x_{-3}x_{-2}x_{-1}}{1 + (2s+2)bx_{-3}x_{-2}x_{-1}x_0}, \quad (48a)$$

$$x_{4n-3} = \frac{x_{-4}^n x_{-3}}{x_0^n} \prod_{s=0}^{n-1} \frac{1 + 2sbx_{-3}x_{-2}x_{-1}x_0}{1 + (2s+1)bx_{-4}x_{-3}x_{-2}x_{-1}}, \quad (48b)$$

$$x_{4n-2} = \frac{x_0^n x_{-2}}{x_{-4}^n} \prod_{s=0}^{n-1} \frac{1 + (2s+1)bx_{-4}x_{-3}x_{-2}x_{-1}}{1 + (2s+1)bx_{-3}x_{-2}x_{-1}x_0} \quad (48c)$$

and

$$x_{4n-1} = \frac{x_{-4}^n x_{-1}}{x_0^n} \prod_{s=0}^{n-1} \frac{1 + (2s+1)bx_{-3}x_{-2}x_{-1}x_0}{1 + (2s+2)bx_{-4}x_{-3}x_{-2}x_{-1}} \quad (48d)$$

where  $x_{-4}, x_0 \neq 0$ ,  $2jbx_{-4}x_{-3}x_{-2}x_{-1} \neq -1$ ,  $(2j-1)bx_{-4}x_{-3}x_{-2}x_{-1} \neq -1$ ,  $2jbx_{-3}x_{-2}x_{-1}x_0 \neq -1$  and  $(2j-1)bx_{-3}x_{-2}x_{-1}x_0 \neq -1$  for all  $j = 1, 2, 3, \dots, n$ .

## 5 The case $a_n, b_n$ are 2-periodic

We assume that  $\{a_n\}_{n=0}^\infty = a_0, a_1, a_0, a_1, \dots$  and  $\{b_n\}_{n=0}^\infty = b_0, b_1, b_0, b_1, \dots$ . Then, from (45), we have

$$x_{4n} = \frac{x_0^{n+1}}{x_{-4}^n} \prod_{s=0}^{n-1} \frac{a_0^{2s+2} + b_0x_{-4}x_{-3}x_{-2}x_{-1} \sum_{l=0}^{2s+1} a_0^l}{a_1^{2s+2} + b_1x_{-3}x_{-2}x_{-1}x_0 \sum_{l=0}^{2s+1} a_1^l}, \quad (49a)$$

$$x_{4n-3} = \frac{x_{-4}^n x_{-3}}{x_0^n} \prod_{s=0}^{n-1} \frac{a_1^{2s} + b_1x_{-3}x_{-2}x_{-1}x_0 \sum_{l=0}^{2s-1} a_1^l}{a_0^{2s+1} + b_0x_{-4}x_{-3}x_{-2}x_{-1} \sum_{l=0}^{2s} a_0^l}, \quad (49b)$$

$$x_{4n-2} = \frac{x_0^n x_{-2}}{x_{-4}^n} \prod_{s=0}^{n-1} \frac{a_0^{2s+1} + b_0x_{-4}x_{-3}x_{-2}x_{-1} \sum_{l=0}^{2s} a_0^l}{a_1^{2s+1} + b_1x_{-3}x_{-2}x_{-1}x_0 \sum_{l=0}^{2s} a_1^l} \quad (49c)$$

and

$$x_{4n-1} = \frac{x_{-4}^n x_{-1}}{x_0^n} \prod_{s=0}^{n-1} \frac{a_1^{2s+1} + b_1x_{-3}x_{-2}x_{-1}x_0 \sum_{l=0}^{2s} a_1^l}{a_0^{2s+2} + b_0x_{-4}x_{-3}x_{-2}x_{-1} \sum_{l=0}^{2s+1} a_0^l} \quad (49d)$$

as long as  $x_{-4}, x_0 \neq 0$  and for all  $(i, s) \in \{0, 1\} \times \{0, 1, 2, \dots, n-1\}$ ,  $a_0^{2s+1+i} + b_0 x_{-4} x_{-3} x_{-2} x_{-1} \sum_{l=0}^{2s+i} a_0^l \neq 0$  and  $a_1^{2s+1+i} + b_1 x_{-3} x_{-2} x_{-1} x_0 \sum_{l=0}^{2s+i} a_1^l \neq 0$ .

## 5.1 The case $a_0 = 1$ and $a_1 = -1$

The solution is given by

$$x_{4n} = \frac{x_0^{n+1}}{x_{-4}^n} \prod_{s=0}^{n-1} (1 + (2s+2)b_0 x_{-4} x_{-3} x_{-2} x_{-1}), \quad (50a)$$

$$x_{4n-3} = \frac{x_{-4}^n x_{-3}}{x_0^n} \prod_{s=0}^{n-1} \frac{1}{1 + (2s+1)b_0 x_{-4} x_{-3} x_{-2} x_{-1}}, \quad (50b)$$

$$x_{4n-2} = x_{-2} \left( \frac{x_0}{x_{-4}(-1 + b_1 x_{-3} x_{-2} x_{-1} x_0)} \right)^n \prod_{s=0}^{n-1} (1 + (2s+1)b_0 x_{-4} x_{-3} x_{-2} x_{-1}), \quad (50c)$$

and

$$x_{4n-1} = x_{-1} \left( \frac{x_{-4}(-1 + b_1 x_{-3} x_{-2} x_{-1} x_0)}{x_0} \right)^n \prod_{s=0}^{n-1} \frac{1}{1 + (2s+2)b_0 x_{-4} x_{-3} x_{-2} x_{-1}} \quad (50d)$$

where  $x_{-4}, x_0 \neq 0$ ,  $b_1 x_{-3} x_{-2} x_{-1} x_0 \neq 1$  and  $j b_0 x_{-4} x_{-3} x_{-2} x_{-1} \neq -1$  for all  $j = 1, 2, 3, \dots, 2n$ .

## 5.2 The case $a_0 = -1$ and $a_1 = 1$

In this case, we obtain

$$x_{4n} = \frac{x_0^{n+1}}{x_{-4}^n} \prod_{s=0}^{n-1} \frac{1}{1 + (2s+2)b_1 x_{-3} x_{-2} x_{-1} x_0}, \quad (51a)$$

$$x_{4n-3} = x_{-3} \left( \frac{x_{-4}}{x_0(-1 + b_0 x_{-4} x_{-3} x_{-2} x_{-1})} \right)^n \prod_{s=0}^{n-1} (1 + 2s b_1 x_{-3} x_{-2} x_{-1} x_0), \quad (51b)$$

$$x_{4n-2} = x_{-2} \left( \frac{x_0(-1 + b_0 x_{-4} x_{-3} x_{-2} x_{-1})}{x_{-4}} \right)^n \prod_{s=0}^{n-1} \frac{1}{1 + (2s+1)b_1 x_{-3} x_{-2} x_{-1} x_0} \quad (51c)$$

and

$$x_{4n-1} = \frac{x_{-4}^n x_{-1}}{x_0^n} \prod_{s=0}^{n-1} (1 + (2s+1)b_1 x_{-3} x_{-2} x_{-1} x_0) \quad (51d)$$

where  $x_{-4}, x_0 \neq 0$ ,  $b_0 x_{-4} x_{-3} x_{-2} x_{-1} \neq 1$  and  $j b_1 x_{-3} x_{-2} x_{-1} x_0 \neq -1$  for all  $j = 1, 2, 3, \dots, 2n$ .

## 6 Conclusion

Our work in this paper was twofold. First, we found non-trivial Lie symmetry generators of the difference equations (2). Second, we derived explicit formulas for solutions of difference equations in question. Consequently, this generalised what Elsayed found in [1] where the values of  $a_n$  and  $b_n$  were only confined to  $\pm 1$ . We showed that in those particular cases, our results yielded Elsayed's results.

## References

- [1] E.M. Elsayed, *Expression and behavior of the solutions of some rational recursive sequences*, Math. Meth. Appl. Sci., **2016:39** (2016) 5682–5694.
- [2] M. Folly-Gbetoula and A.H. Kara, Symmetries, conservation laws, and integrability of difference equations, *Advances in Difference Equations*, **2014**, 2014.
- [3] M. Folly-Gbetoula (2017), Symmetry, reductions and exact solutions of the difference equation  $u_{n+2} = (au_n)/(1+bu_n u_{n+1})$ , *Journal of Difference Equations and Applications*, **23:6** (2017).
- [4] P. E. Hydon, *Difference Equations by Differential Equation Methods*, Cambridge University Press (2014).
- [5] P. E. Hydon, *Symmetries and first integrals of ordinary difference equations*, Proc. Royal Soc. A, **456:2004** (2000) 2835–2855.
- [6] N. Joshi and P. Vassiliou, The existence of Lie Symmetries for First-Order Analytic Discrete Dynamical Systems, *Journal of Mathematical Analysis and Applications* **195** (1995) 872-887.

- [7] D. Levi, L. Vinet and P. Winternitz, Lie group formalism for difference equations, *J. Phys. A: Math. Gen.* **30:2** (1997) 633-649.
- [8] S. Lie, *Classification und Integration von gewöhnlichen Differentialgleichungen zwischen  $xy$ , die eine Gruppe von Transformationen gestatten I*, *Math. Ann.*, **22** (1888) 213–253.
- [9] S. Maeda, Canonical structure and symmetries for discrete systems, *Math. Japonica* **25** (1980) 405–420.
- [10] S. Maeda, The similarity method for difference equations, *IMA J. Appl. Math.* **38** (1987) 129-134.
- [11] D. Nyirenda and M. Folly-Gbetoula, Invariance analysis and exact solutions of some sixth-order difference equations, *J. Nonlinear Sci. Appl.*, **10** (2017) 6262-6273.
- [12] P. J. Olver, *Applications of Lie Groups to Differential Equations*, Second Edition, Springer, New York (1993).
- [13] G. R. W. Quispel and R. Sahadevan, Lie symmetries and the integration of difference equations, *Physics Letters A*, **184** (1993) 64-70.
- [14] R. Morris, A. H. Kara, A. Chowdhury, A. Biswas, *Soliton solutions, conservation laws, and reductions of certain classes of nonlinear wave equations*, *Zeitschrift für Naturforschung Section A*, **67:10-11** (2012) 613–620.
- [15] E. Noether, *Invariante variations probleme*, *Mathematisch-Physikalische Klasse* **2** (1918), 235–257.
- [16] G. R. W. Quispel, R. Sahadevan, *Lie symmetries and the integration of difference equations*, *Phys. Lett. A* **184:1** (1993), 64–70.
- [17] W. Sarlet, F. Cantrijn, *Generalizations of Noether's theorem in classical mechanics*, *SIAM Review* **23:4** (1981), 467–494.