

# A group theory approach towards some rational difference equations

Mensah Folly-Gbetoula\*, Nkosingiphile Mnguni and A. H. Kara

School of Mathematics, University of the Witwatersrand, Private Bag 3,  
Johannesburg 2050, South Africa.

## Abstract

A full Lie point symmetry analysis of rational difference equations is performed. Non-trivial symmetries are derived and exact solutions using these symmetries are obtained.

**Keywords:** Difference equation; symmetry; canonical coordinates; group invariant solutions

MSC: 39A10; 39A13; 39A99

## 1 Introduction

Over a century ago, symmetries became a centre of interest of several authors after the work of Sophus Lie [6] on differential equations. He studied the continuous group of transformations that leaves the differential equations invariant. This concept of symmetries is strongly related to the existence of conservation laws and the relationship between them has attracted great interest among researchers following the work of Noether [7]. The extension of this idea to difference equations is now well-documented (see [4] and references herein). In [4], Hydon developed a symmetry based algorithm enabling one to derive solutions of difference equations without making any special lucky guesses. Hydon emphasized on second-order difference equations, although his algorithm is valid for any order. When it comes to higher-order equations, computations can be cumbersome and extra ansatz may be needed to ease the calculations.

---

\*Mensah.Folly-Gbetoula@wits.ac.za

We aim to extend the work by Elsayed [1] where the author investigated the dynamics and solutions of

$$x_{n+1} = \frac{x_{n-5}}{\pm 1 \pm x_{n-1}x_{n-3}x_{n-5}}, \quad (1)$$

where the initial conditions  $x_{-5}, x_{-4}, x_{-3}, x_{-2}, x_{-1}$  and  $x_0$  are arbitrary non-zero real numbers. For related work, see [2, 3]. One can notice that equations (1) are just special cases of a more general form

$$x_{n+1} = \frac{x_{n-5}}{a_n + b_n x_{n-1}x_{n-3}x_{n-5}}, \quad (2)$$

where  $(a_n)_{n \in \mathbb{N}_0}$  and  $(b_n)_{n \in \mathbb{N}_0}$  are arbitrary sequences. We will use a symmetry based method to solve (2). Equivalently, we study

$$u_{n+6} = \frac{u_n}{A_n + B_n u_n u_{n+2} u_{n+4}} \quad (3)$$

instead, where  $(A_n)_{n \in \mathbb{N}_0}$  and  $(B_n)_{n \in \mathbb{N}_0}$  are arbitrary sequences. This means we can only compare  $x_i$  with  $u_{i+5}$ . Furthermore, we use the same technique to obtain exact solutions of

$$x_{n+1} = \frac{x_{n-4}x_{n-3}x_{n-2}}{x_{n-1}x_n(\lambda + \mu x_{n-4}x_{n-3}x_{n-2})}, \quad (4)$$

where  $\lambda, \mu \in \mathbb{R}$ , and again we study

$$u_{n+5} = \frac{u_n u_{n+1} u_{n+2}}{u_{n+3} u_{n+4} (\lambda + \mu u_n u_{n+1} u_{n+2})}, \quad (5)$$

instead. Note that solutions of (4) were found in [8]; however, their method is completely different from ours.

## 2 Definitions and notation

The definitions are taken from Hydon [4] and most of the notation follows from the same book.

**Definition 1** *A parameterized set of point transformations,*

$$\Gamma_\varepsilon : x \mapsto \hat{x}(x; \varepsilon), \quad (6)$$

where  $x = x_i, i = 1, \dots, p$  are continuous variables, is a one-parameter local Lie group of transformations if the following conditions are satisfied:

1.  $\Gamma_0$  is the identity map if  $\hat{x} = x$  when  $\varepsilon = 0$
2.  $\Gamma_a \Gamma_b = \Gamma_{a+b}$  for every  $a$  and  $b$  sufficiently close to 0
3. Each  $\hat{x}_i$  can be represented as a Taylor series (in a neighborhood of  $\varepsilon = 0$  that is determined by  $x$ ), and therefore

$$\hat{x}_i(x; \varepsilon) = x_i + \varepsilon \xi_i(x) + O(\varepsilon^2), i = 1, \dots, p. \quad (7)$$

Consider the  $p$ th-order difference equation

$$u_{n+p} = \Omega(n, u_n, \dots, u_{n+p-1}), \quad (8)$$

for some function  $\Omega$ . Assume the point transformations are of the form

$$\hat{n} = n; \quad \hat{u}_n = u_n + \varepsilon Q(n, u_n) + O(\varepsilon^2) \quad (9)$$

with the corresponding infinitesimal symmetry generator

$$X = Q(n, u_n) \frac{\partial}{\partial u_n} + S Q(n, u_n) \frac{\partial}{\partial u_{n+1}} + \dots + S^{(p-1)} Q(n, u_n) \frac{\partial}{\partial u_{n+p-1}}, \quad (10)$$

where  $S$  is the shift operator, i.e.,  $S : n \mapsto n + 1$ . The symmetry condition is defined as

$$\hat{u}_{n+p} = \Omega(n, \hat{u}_n, \hat{u}_{n+1}, \dots, \hat{u}_{n+p-1}), \quad (11)$$

whenever (8) is true. Substituting the Lie point symmetries (9) into the symmetry condition (11) leads to the linearized symmetry condition

$$S^{(p)} Q - X \Omega = 0 \quad (12)$$

whenever (8) holds.

**Definition 2**  $V_n$  is invariant under the Lie group of transformations (9) if and only if  $X V_n = 0$ .

We define the functions

$$\Theta^n(\theta_j, s, l) = \prod_{j=s}^l \theta_{2j+n}, \quad n = 0, 1, \quad (13)$$

and we adopt the standard conventions

$$\prod_{j=s}^l \theta_j = 1 \text{ when } s > l \text{ and } \sum_{j=n}^{l_0} \theta_j = 0 \text{ when } n > l_0. \quad (14)$$

We refer the reader to [4] for a deeper understanding of the concept of symmetry analysis of difference equations.

### 3 Main results

#### 3.1 On the difference equations (3)

Consider the sixth-order difference equations of the form (3), that is,

$$u_{n+6} = \frac{u_n}{A_n + B_n u_n u_{n+2} u_{n+4}}. \quad (15)$$

We impose the symmetry condition (12) and simplify the resulting equation to get

$$S^6 Q + \frac{B_n u_n^2 u_{n+2} S^4 Q + B_n u_n^2 u_{n+4} S^2 Q - A_n Q}{(B_n u_n u_{n+2} u_{n+4} + A_n)^2} = 0. \quad (16)$$

To solve for  $Q$ , we first differentiate (16) with respect to  $u_n$  (keeping  $\Omega$  fixed and viewing  $u_{n+2}$  as a function of  $u_n, u_{n+4}$  and  $\Omega$ ). This leads, after simplification, to

$$S^4 Q + u_{n+4} (S^2 Q)' - u_{n+4} Q' + \frac{2u_{n+4}}{u_n} Q = 0, \quad (17)$$

where  $'$  denotes the derivative with respect to the independent variable. We then differentiate (17) with respect to  $u_n$  to get

$$Q''(n, u_n) - \frac{2}{u_n} Q'(n, u_n) + \frac{2}{u_n^2} Q(n, u_n) = 0. \quad (18)$$

The solutions of (18) are given by

$$Q(n, u_n) = \alpha_n u_n^2 + \beta_n u_n \quad (19)$$

for some functions  $\alpha_n$  and  $\beta_n$  of  $n$ . To obtain more information on  $\alpha_n$  and  $\beta_n$ , we substitute (19) in (16) and we split the resulting equation to get the following:

$$u_n u_{n+2} u_{n+4}^2 : B_n \alpha_{n+4} = 0 \quad (20a)$$

$$u_n u_{n+2}^2 u_{n+4} : B_n \alpha_{n+2} = 0 \quad (20b)$$

$$u_n u_{n+2} u_{n+4} : B_n (\beta_{n+2} + \beta_{n+4} + \beta_{n+6}) = 0 \quad (20c)$$

$$u_n : -A_n \alpha_n + \alpha_{n+6} = 0 \quad (20d)$$

$$1 : A_n (\beta_n - \beta_{n+6}) = 0. \quad (20e)$$

These equations (20) reduce to

$$\alpha_n = 0, \quad \beta_{n+4} + \beta_{n+2} + \beta_n = 0. \quad (21)$$

The expression of  $\beta_n$  in (21) is merely obtained by solving the corresponding characteristic equation  $r^4 + r^2 + 1 = 0$  for  $r$ . Assuming that  $r_i$ ,  $i = 1, 2, 3, 4$ , are the solutions of this characteristic equation, then  $\beta_n$  is a linear combination of the  $r_i^n$ 's. In other words, the solutions of (21) are

$$\alpha_n = 0 \quad \text{and} \quad \beta_n = (-1)^n \beta^n c_1 + (-1)^n \bar{\beta}^n c_2 + \bar{\beta}^n c_3 + \beta^n c_4 \quad (22)$$

for some arbitrary constants  $c_i$ ,  $i = 1, \dots, 4$ , and where  $\beta = \exp(\pi i/3)$ . Thus, we obtain four characteristics given by

$$Q_1 = (-1)^n \beta^n u_n, \quad Q_2 = (-1)^n \bar{\beta}^n u_n, \quad Q_3 = \bar{\beta}^n u_n, \quad Q_4 = \beta^n u_n. \quad (23)$$

The four corresponding symmetry generators  $X_1$ ,  $X_2$ ,  $X_3$  and  $X_4$  are given by

$$X_1 = (-1)^n \beta^n u_n \partial_{u_n} - (-1)^n \beta^{n+1} u_{n+1} \partial_{u_{n+1}} + (-1)^n \beta^{n+2} u_{n+2} \partial_{u_{n+2}} - (-1)^n \beta^{n+3} u_{n+3} \partial_{u_{n+3}} + (-1)^n \beta^{n+4} u_{n+4} \partial_{u_{n+4}} - (-1)^n \beta^{n+5} u_{n+5} \partial_{u_{n+5}}, \quad (24a)$$

$$X_2 = (-1)^n \bar{\beta}^n u_n \partial_{u_n} - (-1)^n \bar{\beta}^{n+1} u_{n+1} \partial_{u_{n+1}} + (-1)^n \bar{\beta}^{n+2} u_{n+2} \partial_{u_{n+2}} - (-1)^n \bar{\beta}^{n+3} u_{n+3} \partial_{u_{n+3}} + (-1)^n \bar{\beta}^{n+4} u_{n+4} \partial_{u_{n+4}} - (-1)^n \bar{\beta}^{n+5} u_{n+5} \partial_{u_{n+5}}, \quad (24b)$$

$$X_3 = \bar{\beta}^n u_n \partial_{u_n} + \bar{\beta}^{n+1} u_{n+1} \partial_{u_{n+1}} + \bar{\beta}^{n+2} u_{n+2} \partial_{u_{n+2}} + \bar{\beta}^{n+3} u_{n+3} \partial_{u_{n+3}} + \bar{\beta}^{n+4} u_{n+4} \partial_{u_{n+4}} + \bar{\beta}^{n+5} u_{n+5} \partial_{u_{n+5}}, \quad (24c)$$

$$X_4 = \beta^n u_n \partial_{u_n} + \beta^{n+1} u_{n+1} \partial_{u_{n+1}} + \beta^{n+2} u_{n+2} \partial_{u_{n+2}} + \beta^{n+3} u_{n+3} \partial_{u_{n+3}} + \beta^{n+4} u_{n+4} \partial_{u_{n+4}} + \beta^{n+5} u_{n+5} \partial_{u_{n+5}}. \quad (24d)$$

Here, using  $Q_4$ , we introduce the canonical coordinate [5]

$$s_n = \int \frac{du_n}{\beta^n u_n} = \frac{1}{\beta^n} \ln |u_n|. \quad (25)$$

In view of (21), we introduce the variable

$$r_n = \beta^{n+4} s_{n+4} + \beta^{n+2} s_{n+2} + \beta^n s_n. \quad (26)$$

It is easy to check that

$$X_i r_n = \beta^{n+4} + \beta^{n+2} + \beta^n = 0, \quad i = 1, \dots, 4. \quad (27)$$

Therefore  $r_n$  is invariant under  $X_4$ . It is advantageous to use

$$|\tilde{r}_n| = \exp(-r_n), \quad (28)$$

that is,  $\tilde{r}_n = \pm 1/(u_n u_{n+2} u_{n+4})$ . Here, we choose to use the plus sign and we have shown, using (3), that

$$\tilde{r}_{n+2} = A_n \tilde{r}_n + B_n. \quad (29)$$

Therefore,

$$\tilde{r}_{2n+k} = \tilde{r}_k \left( \prod_{k_1=0}^{n-1} A_{2k_1+k} \right) + \sum_{l=0}^{n-1} \left( B_{2l+k} \prod_{k_2=l+1}^{n-1} A_{2k_2+k} \right), \quad k = 0, 1. \quad (30)$$

It is worthwhile to mention that equations in (30) give the solution of (29) for all  $n$ . By reversing all the change of variables, we have

$$\begin{aligned} |u_n| = \exp & \left[ (-\beta)^n c_5 + (\beta)^n c_6 + (\bar{\beta})^n c_7 + (-\bar{\beta})^n c_8 + \right. \\ & (-\beta)^{n+1} \sum_{k_1=0}^{n-1} \frac{-i\sqrt{3}}{6} (-\bar{\beta})^{k_1} r_{k_1} + (\beta)^{n+1} \sum_{k_2=0}^{n-1} \frac{i\sqrt{3}}{6} (\bar{\beta})^{k_2} r_{k_2} + \\ & \left. (\bar{\beta})^{n+1} \sum_{k_3=0}^{n-1} \frac{-i\sqrt{3}}{6} \beta^{k_3} r_{k_3} + (-\bar{\beta})^{n+1} \sum_{k_4=0}^{n-1} \frac{i\sqrt{3}}{6} (-\beta)^{k_4} r_{k_4} \right] \\ = & \Gamma_n \exp \left( \sum_{k=0}^{n-1} \frac{\sqrt{3}}{3} [(-1)^{n+k} + 1] \operatorname{Im} [\gamma(n+1, k)] \ln |\tilde{r}_k| \right), \quad (31) \end{aligned}$$

where  $\tilde{r}_k$  is given in (30),  $\Gamma_n = \exp\{(-\beta)^n c_5 + \beta^n c_6 + \bar{\beta}^n c_7 + (-\bar{\beta})^n c_8\}$  and  $\gamma(l, k) = \beta^l \bar{\beta}^k$ .

**Note.** Equation (31) gives the solution of (3) in a unified manner.

We can simplify (31) further by splitting it into six categories. We have

$$|u_{6n}| = \Gamma_{6n} \exp \left( \sum_{k=0}^{6n-1} \frac{\sqrt{3}}{3} [(-1)^{6n+k} + 1] \operatorname{Im} [\gamma(6n+1, k)] \ln |\tilde{r}_k| \right) \quad (32)$$

$$u_{6n} = u_0 \prod_{s=0}^{n-1} \frac{\tilde{r}_{6s}}{\tilde{r}_{6s+2}}. \quad (33)$$

Similarly, after a straightforward but lengthy computation, we get

$$u_{6n+i} = u_i \prod_{s=0}^{n-1} \frac{\tilde{r}_{6s+i}}{\tilde{r}_{6s+2+i}}, i = 0, \dots, 5. \quad (34)$$

**Note.** We can obtain (34) using (28) which need not the use of absolute values.

Now, using (30) and (34), we obtain the solutions of (3) as follows:

$$u_{6n} = u_0 \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{3s-1} A_{2k_1} \right) + u_0 u_2 u_4 \sum_{l=0}^{3s-1} \left( B_{2l} \prod_{k_2=l+1}^{3s-1} A_{2k_2} \right)}{\left( \prod_{k_1=0}^{3s} A_{2k_1} \right) + u_0 u_2 u_4 \sum_{l=0}^{3s} \left( B_{2l} \prod_{k_2=l+1}^{3s} A_{2k_2} \right)}, \quad (35a)$$

$$u_{6n+1} = u_1 \frac{\left( \prod_{k_1=0}^{3s-1} A_{2k_1+1} \right) + u_1 u_3 u_5 \sum_{l=0}^{3s-1} \left( B_{2l+1} \prod_{k_2=l+1}^{3s-1} A_{2k_2+1} \right)}{\left( \prod_{k_1=0}^{3s} A_{2k_1+1} \right) + u_1 u_3 u_5 \sum_{l=0}^{3s} \left( B_{2l+1} \prod_{k_2=l+1}^{3s} A_{2k_2+1} \right)}, \quad (35b)$$

$$u_{6n+2} = u_2 \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{3s} A_{2k_1} \right) + u_0 u_2 u_4 \sum_{l=0}^{3s} \left( B_{2l} \prod_{k_2=l+1}^{3s} A_{2k_2} \right)}{\left( \prod_{k_1=0}^{3s+1} A_{2k_1} \right) + u_0 u_2 u_4 \sum_{l=0}^{3s+1} \left( B_{2l} \prod_{k_2=l+1}^{3s+1} A_{2k_2} \right)}, \quad (35c)$$

$$u_{6n+3} = u_3 \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{3s} A_{2k_1+1} \right) + u_1 u_3 u_5 \sum_{l=0}^{3s} \left( B_{2l+1} \prod_{k_2=l+1}^{3s} A_{2k_2+1} \right)}{\left( \prod_{k_1=0}^{3s+1} A_{2k_1+1} \right) + u_1 u_3 u_5 \sum_{l=0}^{3s+1} \left( B_{2l+1} \prod_{k_2=l+1}^{3s+1} A_{2k_2+1} \right)}, \quad (35d)$$

$$u_{6n+4} = u_4 \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{3s+1} A_{2k_1} \right) + u_0 u_2 u_4 \sum_{l=0}^{3s+1} \left( B_{2l} \prod_{k_2=l+1}^{3s+1} A_{2k_2} \right)}{\left( \prod_{k_1=0}^{3s+2} A_{2k_1} \right) + u_0 u_2 u_4 \sum_{l=0}^{3s+2} \left( B_{2l} \prod_{k_2=l+1}^{3s+2} A_{2k_2} \right)}, \quad (35e)$$

$$u_{6n+5} = u_5 \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{3s+1} A_{2k_1+1} \right) + u_1 u_3 u_5 \sum_{l=0}^{3s+1} \left( B_{2l+1} \prod_{k_2=l+1}^{3s+1} A_{2k_2+1} \right)}{\left( \prod_{k_1=0}^{3s+2} A_{2k_1+1} \right) + u_1 u_3 u_5 \sum_{l=0}^{3s+2} \left( B_{2l+1} \prod_{k_2=l+1}^{3s+2} A_{2k_2+1} \right)}, \quad (35f)$$

whenever the denominators do not vanish, i.e.,  $u_0 u_2 u_4 \sum_{l=0}^s B_{2l} \Theta^0(A_{k_2}, l+1, s) \neq -\Theta^0(A_{k_1}, 0, s)$  and  $u_1 u_3 u_5 \sum_{l=0}^s B_{2l+1} \Theta^1(A_{k_2}, l+1, s) \neq -\Theta^1(A_{k_1}, 0, s)$ . The implication is that solutions of (2) are given by

$$x_{6n-5} = x_{-5} \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{3s-1} a_{2k_1} \right) + x_{-5} x_{-3} x_{-1} \sum_{l=0}^{3s-1} \left( b_{2l} \prod_{k_2=l+1}^{3s-1} a_{2k_2} \right)}{\left( \prod_{k_1=0}^{3s} a_{2k_1} \right) + x_{-5} x_{-3} x_{-1} \sum_{l=0}^{3s} \left( b_{2l} \prod_{k_2=l+1}^{3s} a_{2k_2} \right)}, \quad (36a)$$

$$x_{6n-4} = x_{-4} \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{3s-1} a_{2k_1+1} \right) + x_{-4} x_{-2} x_0 \sum_{l=0}^{3s-1} \left( b_{2l+1} \prod_{k_2=l+1}^{3s-1} a_{2k_2+1} \right)}{\left( \prod_{k_1=0}^{3s} a_{2k_1+1} \right) + x_{-4} x_{-2} x_0 \sum_{l=0}^{3s} \left( b_{2l+1} \prod_{k_2=l+1}^{3s} a_{2k_2+1} \right)}, \quad (36b)$$

$$x_{6n-3} = x_{-3} \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{3s} a_{2k_1} \right) + x_{-5} x_{-3} x_{-1} \sum_{l=0}^{3s} \left( b_{2l} \prod_{k_2=l+1}^{3s} a_{2k_2} \right)}{\left( \prod_{k_1=0}^{3s+1} a_{2k_1} \right) + x_{-5} x_{-3} x_{-1} \sum_{l=0}^{3s+1} \left( b_{2l} \prod_{k_2=l+1}^{3s+1} a_{2k_2} \right)}, \quad (36c)$$

$$x_{6n-2} = x_{-2} \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{3s} a_{2k_1+1} \right) + x_{-4} x_{-2} x_0 \sum_{l=0}^{3s} \left( b_{2l+1} \prod_{k_2=l+1}^{3s} a_{2k_2+1} \right)}{\left( \prod_{k_1=0}^{3s+1} a_{2k_1+1} \right) + x_{-4} x_{-2} x_0 \sum_{l=0}^{3s+1} \left( b_{2l+1} \prod_{k_2=l+1}^{3s+1} a_{2k_2+1} \right)}, \quad (36d)$$

$$x_{6n-1} = x_{-1} \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{3s+1} a_{2k_1} \right) + x_{-5} x_{-3} x_{-1} \sum_{l=0}^{3s+1} \left( b_{2l} \prod_{k_2=l+1}^{3s+1} a_{2k_2} \right)}{\left( \prod_{k_1=0}^{3s+2} a_{2k_1} \right) + x_{-5} x_{-3} x_{-1} \sum_{l=0}^{3s+2} \left( b_{2l} \prod_{k_2=l+1}^{3s+2} a_{2k_2} \right)}, \quad (36e)$$



$$x_{6n} = x_0 \prod_{s=0}^{n-1} \frac{\left( \prod_{k_1=0}^{3s+1} a_{2k_1+1} \right) + x_{-4}x_{-2}x_0 \sum_{l=0}^{3s+1} \left( b_{2l+1} \prod_{k_2=l+1}^{3s+1} a_{2k_2+1} \right)}{\left( \prod_{k_1=0}^{3s+2} a_{2k_1+1} \right) + x_{-4}x_{-2}x_0 \sum_{l=0}^{3s+2} \left( b_{2l+1} \prod_{k_2=l+1}^{3s+2} a_{2k_2+1} \right)}, \quad (36f)$$

whenever the denominators do not vanish, i.e.,  $x_{-5}x_{-3}x_{-1} \sum_{l=0}^s b_{2l} \Theta^0(a_{k_2}, l+1, s) \neq -\Theta^0(a_{k_1}, 0, s)$  and  $x_{-4}x_{-2}x_0 \sum_{l=0}^s b_{2l+1} \Theta^1(a_{k_2}, l+1, s) \neq -\Theta^1(a_{k_1}, 0, s)$ .

### 3.2 The case where $(a_n)$ and $(b_n)$ are two-periodic sequences

Let  $a_n = (\lambda, \mu, \lambda, \mu, \lambda, \dots)$ ,  $b_n = (\eta, \zeta, \eta, \zeta, \dots)$ ,  $\Phi = x_{-5}x_{-3}x_{-1}$  and  $\Psi = x_{-4}x_{-2}x_0$ . The solution in this case is given by the equations

$$x_{6n-5} = x_{-5} \prod_{s=0}^{n-1} \frac{\lambda^{3s} + \eta\Phi \sum_{l=0}^{3s-1} \lambda^l}{\lambda^{3s+1} + \eta\Phi \sum_{l=0}^{3s} \lambda^l}, \quad x_{6n-4} = x_{-4} \prod_{s=0}^{n-1} \frac{\mu^{3s} + \zeta\Psi \sum_{l=0}^{3s-1} \mu^l}{\mu^{3s+1} + \zeta\Psi \sum_{l=0}^{3s} \mu^l}, \quad (37a)$$

$$x_{6n-3} = x_{-3} \prod_{s=0}^{n-1} \frac{\lambda^{3s+1} + \eta\Phi \sum_{l=0}^{3s} \lambda^l}{\lambda^{3s+2} + \eta\Phi \sum_{l=0}^{3s+1} \lambda^l}, \quad x_{6n-2} = x_{-2} \prod_{s=0}^{n-1} \frac{\mu^{3s+1} + \zeta\Psi \sum_{l=0}^{3s} \mu^l}{\mu^{3s+2} + \zeta\Psi \sum_{l=0}^{3s+1} \mu^l}, \quad (37b)$$

$$x_{6n-1} = x_{-1} \prod_{s=0}^{n-1} \frac{\lambda^{3s+2} + \eta\Phi \sum_{l=0}^{3s+1} \lambda^l}{\lambda^{3s+3} + \eta\Phi \sum_{l=0}^{3s+2} \lambda^l}, \quad x_{6n} = x_0 \prod_{s=0}^{n-1} \frac{\mu^{3s+2} + \zeta\Psi \sum_{l=0}^{3s+1} \mu^l}{\mu^{3s+3} + \zeta\Psi \sum_{l=0}^{3s+2} \mu^l}, \quad (37c)$$

where  $\left( \sum_{l=0}^{s-1} \lambda^l \right) \eta\Phi \neq -\lambda^s$  and  $\left( \sum_{l=0}^{s-1} \mu^l \right) \zeta\Psi \neq -\mu^s$ ,  $s \leq 3n$ .

### 3.3 The case where $(a_n)$ and $(b_n)$ are constants

Here,  $a_n = \mu = \lambda$ ,  $b_n = \eta = \zeta$ ,  $\Phi = x_{-5}x_{-3}x_{-1}$  and  $\Psi = x_{-4}x_{-2}x_0$ . Equation (2) becomes  $x_{n+1} = x_{n-5}/(\lambda + \eta x_{n-1}x_{n-3}x_{n-5})$ .

### 3.3.1 $\lambda = 1$

The solution given in (36) simplifies to

$$x_{6n-5} = x_{-5} \prod_{s=0}^{n-1} \frac{1 + (3s)\eta\Phi}{1 + (3s+1)\eta\Phi}, x_{6n-4} = x_{-4} \prod_{s=0}^{n-1} \frac{1 + (3s)\eta\Psi}{1 + (3s+1)\eta\Psi}, \quad (38a)$$

$$x_{6n-3} = x_{-3} \prod_{s=0}^{n-1} \frac{1 + (3s+1)\eta\Phi}{1 + (3s+2)\eta\Phi}, x_{6n-2} = x_{-2} \prod_{s=0}^{n-1} \frac{1 + (3s+1)\eta\Psi}{1 + (3s+2)\eta\Psi}, \quad (38b)$$

$$x_{6n-1} = x_{-1} \prod_{s=0}^{n-1} \frac{1 + (3s+2)\eta\Phi}{1 + (3s+3)\eta\Phi}, x_{6n} = x_0 \prod_{s=0}^{n-1} \frac{1 + (3s+2)\eta\Psi}{1 + (3s+3)\eta\Psi}, \quad (38c)$$

where  $s\eta\Phi \neq -1$  and  $s\eta\Psi \neq -1$ ,  $s \leq 3n$ .

- If  $\eta = 1$ , then equations in (38) are exactly the ones obtained in Theorem 2.1 in [1] for

$$x_{n+1} = \frac{x_{n-5}}{1 + x_{n-1}x_{n-3}x_{n-5}} \quad (39)$$

and their restriction (the initial conditions are arbitrary nonzero positive real numbers) is a special case of our restriction (the initial conditions are arbitrary nonzero real numbers and  $s\Phi \neq -1$ ,  $s\Psi \neq -1$ ,  $0 \leq s \leq 3n$ ).

- If  $\eta = -1$ , then equations in (38) are exactly the ones obtained in Theorem 3.1 in [1] for

$$x_{n+1} = \frac{x_{n-5}}{1 - x_{n-1}x_{n-3}x_{n-5}} \quad (40)$$

and their restriction (the initial conditions are arbitrary nonzero real numbers and  $j b d f \neq 1$ ,  $j a c e \neq 1$ ) coincides with our restriction (the initial conditions are arbitrary nonzero real numbers and  $s\Phi \neq 1$ ,  $s\Psi \neq 1$ ,  $0 \leq s \leq 3n$ ).

### 3.3.2 The case where $\lambda \neq 1$

In this case, the solution given in (36) simplifies to

$$\begin{aligned}
x_{6n-5} &= x_{-5} \prod_{s=0}^{n-1} \frac{\lambda^{3s} + \eta\Phi \left( \frac{1-\lambda^{3s}}{1-\lambda} \right)}{\lambda^{3s+1} + \eta\Phi \left( \frac{1-\lambda^{3s+1}}{1-\lambda} \right)}, & x_{6n-4} &= x_{-4} \prod_{s=0}^{n-1} \frac{\lambda^{3s} + \zeta\Psi \left( \frac{1-\lambda^{3s}}{1-\lambda} \right)}{\lambda^{3s+1} + \zeta\Psi \left( \frac{1-\lambda^{3s+1}}{1-\lambda} \right)}, \\
x_{6n-3} &= x_{-3} \prod_{s=0}^{n-1} \frac{\lambda^{3s+1} + \eta\Phi \left( \frac{1-\lambda^{3s+1}}{1-\lambda} \right)}{\lambda^{3s+2} + \eta\Phi \left( \frac{1-\lambda^{3s+2}}{1-\lambda} \right)}, & x_{6n-2} &= x_{-2} \prod_{s=0}^{n-1} \frac{\lambda^{3s+1} + \zeta\Psi \left( \frac{1-\lambda^{3s+1}}{1-\lambda} \right)}{\lambda^{3s+2} + \zeta\Psi \left( \frac{1-\lambda^{3s+2}}{1-\lambda} \right)}, \\
x_{6n-1} &= x_{-1} \prod_{s=0}^{n-1} \frac{\lambda^{3s+2} + \eta\Phi \left( \frac{1-\lambda^{3s+2}}{1-\lambda} \right)}{\lambda^{3s+3} + \eta\Phi \left( \frac{1-\lambda^{3s+3}}{1-\lambda} \right)}, & x_{6n} &= x_0 \prod_{s=0}^{n-1} \frac{\lambda^{3s+2} + \zeta\Psi \left( \frac{1-\lambda^{3s+2}}{1-\lambda} \right)}{\lambda^{3s+3} + \zeta\Psi \left( \frac{1-\lambda^{3s+3}}{1-\lambda} \right)},
\end{aligned} \tag{41}$$

where  $(1 - \lambda^s)\eta\Phi \neq -\lambda^s(1 - \lambda)$  and  $(1 - \lambda^s)\zeta\Psi \neq -\lambda^s(1 - \lambda)$ ,  $s \leq 3n$ .

**Note.** If  $\lambda = -1$  then the solution given in (41) simplifies to

$$\begin{aligned}
x_{12n-5} &= x_{6(2n)-5} = x_{-5}, & x_{12n-4} &= x_{-4}, & x_{12n-3} &= x_{-3}, & x_{12n-2} &= x_{-2}, \\
x_{12n-1} &= x_{-1}, & x_{12n} &= x_0, & x_{12n+1} &= \frac{x_{-5}}{-1 + \eta\Phi}, & x_{12n+2} &= \frac{x_{-4}}{-1 + \zeta\Psi}, \\
x_{12n+3} &= x_{-3}(-1 + \eta\Phi), & x_{12n+4} &= x_{-2}(-1 + \zeta\Psi), & x_{12n+5} &= \frac{x_{-1}}{(-1 + \eta\Phi)}, \\
x_{12n+6} &= \frac{x_0}{-1 + \zeta\Psi}, & x_{12n+7} &= x_{6(2n+2)-5} = x_{-5}.
\end{aligned} \tag{42}$$

- When setting  $\eta = 1$  in (42), we get the result obtained in Theorem 4.1 in [1] for

$$x_{n+1} = \frac{x_{n-5}}{-1 + x_{n-1}x_{n-3}x_{n-5}} \tag{43}$$

and their restriction coincides with our restriction (the initial conditions are arbitrary nonzero real numbers,  $\Phi \neq 1$  and  $\Psi \neq 1$ ).

- When setting  $\eta = -1$  in (42), we get the result obtained in Theorem 5.1 in [1] for

$$x_{n+1} = \frac{x_{n-5}}{-1 - x_{n-1}x_{n-3}x_{n-5}} \tag{44}$$

and their restriction coincides with our restriction (the initial conditions are arbitrary nonzero real numbers,  $\Phi \neq -1$  and  $\Psi \neq -1$ ).

## 4 On the difference equations (5)

Consider the fifth-order difference equations of the form (5), i.e.,

$$u_{n+5} = \Omega = \frac{u_n u_{n+1} u_{n+2}}{u_{n+3} u_{n+4} (\lambda + \mu u_n u_{n+1} u_{n+2})}.$$

Here, the procedure for finding the characteristics of (5) is similar as above and is as follows:

- Impose the symmetry condition (12) to (5).
- Differentiate with respect to  $u_n$  (keeping  $\Omega$  fixed) and viewing  $u_{n+3}$  as a function of  $u_n, u_{n+1}, u_{n+2}$  and  $\Omega$ .
- Differentiate with respect to  $u_n$  twice (keeping  $u_{n+1}$  fixed).
- Use the method of separation.

After performing this series of operations, we obtain the characteristics

$$Q_5(n, u_n) = \beta^n u_n \quad \text{and} \quad Q_6 = \bar{\beta}^n u_n, \quad (45)$$

where  $\beta = \exp\left(\frac{-2\pi i}{3}\right)$ . Thus, we obtain two characteristics with corresponding generators given by

$$X_5 = \bar{\beta}^n u_n \partial_{u_n} + \bar{\beta}^{n+1} u_{n+1} \partial_{u_{n+1}} + \beta \bar{\beta}^n u_{n+2} \partial_{u_{n+2}} + \bar{\beta}^n u_{n+3} \partial_{u_{n+3}} + \bar{\beta}^{n+1} u_{n+4} \partial_{u_{n+4}}, \quad (46a)$$

$$X_6 = \beta^n u_n \partial_{u_n} + \beta^{n+1} u_{n+1} \partial_{u_{n+1}} + \bar{\beta} \beta^n u_{n+2} \partial_{u_{n+2}} + \beta^n u_{n+3} \partial_{u_{n+3}} + \beta^{n+1} u_{n+4} \partial_{u_{n+4}}. \quad (46b)$$

From the characteristic equations

$$\frac{du_n}{\bar{\beta}^n u_n} = \frac{du_{n+1}}{\bar{\beta}^{n+1} u_{n+1}} = \frac{du_{n+2}}{\bar{\beta}^{n+2} u_{n+2}} = \frac{du_{n+3}}{\bar{\beta}^{n+3} u_{n+3}} = \frac{du_{n+4}}{\bar{\beta}^{n+4} u_{n+4}} \left( = \frac{V_n}{0} \right), \quad (47)$$

we obtain the invariants  $c_1 = u_{n+1}^\beta / u_n$ ,  $c_2 = u_{n+2}^{\bar{\beta}} / u_n$ ,  $c_3 = u_n u_{n+3}$ ,  $c_4 = u_{n+4}^\beta / u_n$  and  $c_5 = V_n$ . We readily notice that

$$S^3 \left( \frac{du_n}{\bar{\beta}^n u_n} = \frac{du_{n+1}}{\bar{\beta}^{n+1} u_{n+1}} \right) = \frac{du_{n+3}}{\bar{\beta}^{n+3} u_{n+3}} = \frac{du_{n+4}}{\bar{\beta}^{n+4} u_{n+4}} \quad (48)$$

and we choose  $V_n = c_1^{\bar{\beta}} c_2^{\beta}$ , i.e.,

$$V_n = u_n u_{n+1} u_{n+2}. \quad (49)$$

By shifting (49) thrice, we get

$$V_{n+3} = \frac{V_n}{\lambda + \mu V_n} \quad (50)$$

whose solution is given by

$$V_n = \begin{cases} \left( \frac{\mu [((-1)^{2/3} - 1)n - (-1)^{1/3} + 1 + \beta]}{3((-1)^{2/3} - 1)} + c_6 + \bar{\beta}^n c_7 + \beta^n c_8 \right)^{-1} & \text{if } \lambda = 1, \\ (c_{10} \lambda^{n/3} + [(-1)^{2/3} \lambda^{1/3}]^n c_8 + [ -(-1)^{1/3} \lambda^{1/3} ]^n c_9 + \frac{\mu}{1-\lambda})^{-1} & \text{if } \lambda \neq 1. \end{cases} \quad (51)$$

The constants  $c_i$ ,  $i = 6, \dots, 10$ , can be obtained from the following equations:

$$c_6 + c_7 + c_8 = \frac{1}{u_0 u_1 u_2} - \frac{\mu [ -(-1)^{1/3} + 1 + \beta ]}{3((-1)^{2/3} - 1)} \quad (52a)$$

$$c_6 + \bar{\beta} c_7 + \beta c_8 = \frac{1}{u_1 u_2 u_3} - \frac{\mu [\beta - (-1)^{1/3} + (-1)^{2/3}]}{3((-1)^{2/3} - 1)} \quad (52b)$$

$$c_6 + \beta c_7 + \bar{\beta} c_8 = \frac{1}{u_2 u_3 u_4} - \frac{\mu [ -1 + 2(-1)^{2/3} - (-1)^{1/3} + \beta ]}{3((-1)^{2/3} - 1)} \quad (52c)$$

$$c_{10} + c_8 + c_9 = \frac{1}{u_0 u_1 u_2} - \frac{\mu}{1 - \lambda} \quad (52d)$$

$$\lambda^{1/3} c_{10} + [(-1)^{2/3} \lambda^{1/3}] c_8 + [ -(-1)^{1/3} \lambda^{1/3} ] c_9 = \frac{1}{u_1 u_2 u_3} - \frac{\mu}{1 - \lambda} \quad (52e)$$

$$\lambda^{2/3} c_{10} + [(-1)^{2/3} \lambda^{1/3}]^2 c_8 + [ -(-1)^{1/3} \lambda^{1/3} ]^2 c_9 = \frac{1}{u_2 u_3 u_4} - \frac{\mu}{1 - \lambda}. \quad (52f)$$

Thanks to (49), we can express  $u_n$  in terms of  $V_n$  as follows:

$$u_n = \exp \left( \beta^n c_{11} + \bar{\beta}^n c_{12} - \frac{2}{\sqrt{3}} \left[ \sum_{k=0}^{n-1} \text{Im}(\gamma(n, k)) \ln V_k \right] \right), \quad (53a)$$

where  $V_k$  is given in (51) with  $\gamma(n, k) = \beta^n \bar{\beta}^{k+1}$ . The constants  $c_{11}$  and  $c_{12}$  must satisfy

$$c_{11} + c_{12} = \ln u_0, \quad \beta c_{11} + \bar{\beta} c_{12} = \ln u_1. \quad (53b)$$

Equations in (53) give the solutions of (5) in a unified manner. For the sake of clarification, we split solutions (53a) to realise the solutions in the existing literature. Using (53a) and (53b) , we have

$$u_{6n} = u_0 \prod_{s=1}^{2n} \frac{V_{3s-2}}{V_{3s-3}}. \quad (54)$$

Using the same approach, we have shown that

$$u_{6n+i} = u_i \prod_{s=1}^{2n} \frac{V_{3(s-1)+i+1}}{V_{3(s-1)+i}}, \quad i = 0, \dots, 5. \quad (55)$$

#### 4.1 The case of $\lambda = 1$

Equation (5) becomes

$$u_{n+5} = \frac{u_n u_{n+1} u_{n+2}}{u_{n+3} u_{n+4} (1 + \mu u_n u_{n+1} u_{n+2})}. \quad (56)$$

and we said earlier that the solution of (50), in this case, is (51), i.e.,

$$V_n = \left( \frac{\mu [((-1)^{2/3} - 1)n - (-1)^{1/3} + 1 + \beta]}{3((-1)^{2/3} - 1)} + c_6 + \bar{\beta}^n c_7 + \beta^n c_8 \right)^{-1}. \quad (57)$$

We have

$$V_{3s} = \left( \frac{\mu [((-1)^{2/3} - 1)(3s) - (-1)^{1/3} + 1 + \beta]}{3((-1)^{2/3} - 1)} + c_6 + c_7 + c_8 \right)^{-1} \quad (58)$$

and using (52a) in (58), we get

$$V_{3s} = \frac{u_0 u_1 u_2}{1 + \mu s u_0 u_1 u_2}. \quad (59)$$

Using the same approach, we have shown that

$$V_{3s} = \frac{u_0 u_1 u_2}{1 + \mu s u_0 u_1 u_2}; \quad V_{3s+1} = \frac{u_1 u_2 u_3}{1 + \mu s u_1 u_2 u_3}; \quad V_{3s+2} = \frac{u_2 u_3 u_4}{1 + \mu s u_2 u_3 u_4}. \quad (60)$$

Let  $a = x_{-4}$ ,  $b = x_{-3}$ ,  $c = x_{-2}$ ,  $d = x_{-1}$  and  $A = u_0$ ,  $B = u_1$ ,  $C = u_2$ ,  $D = u_3$  and  $E = u_4$ . Using (60) in (55), we obtain the solution of (56) as follows:

$$\begin{aligned}
u_{6n} &= \frac{D^{2n}}{A^{2n-1}} \prod_{s=1}^{2n-1} \frac{1 + \mu s ABC}{1 + \mu s BCD}, u_{6n+1} = \frac{E^{2n}}{B^{2n-1}} \prod_{s=1}^{2n-1} \frac{1 + \mu s BCD}{1 + \mu s CDE}, \\
u_{6n+2} &= \frac{CA^{2n} B^{2n}}{D^{2n} E^{2n}} \prod_{s=0}^{2n-1} \frac{1 + \mu s CDE}{1 + \mu(s+1) ABC}, u_{6n+3} = \frac{D^{2n+1}}{A^{2n}} \prod_{s=1}^{2n} \frac{1 + \mu s ABC}{1 + \mu s BCD}, \\
u_{6n+4} &= \frac{E^{2n+1}}{B^{2n}} \prod_{s=0}^{2n} \frac{1 + \mu s BCD}{1 + \mu s CDE}, u_{6n+5} = \frac{C(AB)^{2n+1}}{(DE)^{2n+1}} \prod_{s=0}^{2n} \frac{1 + \mu s CDE}{1 + \mu(s+1) ABC}
\end{aligned} \tag{61}$$

whenever the denominators do not vanish.

## 4.2 The case of $\lambda \neq 1$

In this case, as we found earlier, the solution of (50) is given by (51), i.e.,

$$V_n = \left( c_{10} \lambda^{n/3} + [(-1)^{2/3} \lambda^{1/3}]^n c_8 + [-(-1)^{1/3} \lambda^{1/3}]^n c_9 + \frac{\mu}{1-\lambda} \right)^{-1}. \tag{62}$$

Using this, we get

$$V_{3s} = \left( c_{10} \lambda^s + c_8 \lambda^s + c_9 \lambda^s + \frac{\mu}{1-\lambda} \right)^{-1} \tag{63}$$

and using (52d) in (63), we find

$$V_{3s} = \frac{ABC}{\lambda^s + \mu ABC \left( \frac{1-\lambda^s}{1-\lambda} \right)}. \tag{64}$$

Using the same approach, we have shown that

$$V_{3s+i} = \frac{u_i u_{i+1} u_{i+2}}{\lambda^s + \mu u_i u_{i+1} u_{i+2} \left( \frac{1-\lambda^s}{1-\lambda} \right)}, \quad i = 0, 1, 2. \tag{65}$$

Using (65) in (55), we obtain the solution of (5) as follows:

$$u_{6n} = \frac{D^{2n}}{A^{2n-1}} \prod_{s=2}^{2n} \frac{\lambda^{s-1} + \mu \Delta_0 \sum_{j=0}^{s-2} \lambda^j}{\lambda^{s-1} + \mu \Delta_1 \sum_{j=0}^{s-2} \lambda^j}, u_{6n+1} = \frac{E^{2n}}{B^{2n-1}} \prod_{s=2}^{2n} \frac{\lambda^{s-1} + \mu \Delta_1 \sum_{j=0}^{s-2} \lambda^j}{\lambda^{s-1} + \mu \Delta_2 \sum_{j=0}^{s-2} \lambda^j},$$

$$\begin{aligned}
u_{6n+2} &= C \frac{\Delta_0^{2n}}{\Delta_1^{2n}} \frac{\prod_{s=2}^{2n} \left( \lambda^{s-1} + \mu \Delta_2 \sum_{j=0}^{s-2} \lambda^j \right)}{\prod_{s=1}^{2n} \left( \lambda^s + \mu \Delta_0 \sum_{j=0}^{s-1} \lambda^j \right)}, u_{6n+3} = \frac{D^{2n+1}}{A^{2n}} \prod_{s=2}^{2n+1} \frac{\lambda^{s-1} + \mu \Delta_0 \sum_{j=0}^{s-2} \lambda^j}{\lambda^{s-1} + \mu \Delta_1 \sum_{j=0}^{s-2} \lambda^j}, \\
u_{6n+4} &= E \frac{\Delta_2^{2n}}{\Delta_1^{2n}} \prod_{s=2}^{2n+1} \frac{\lambda^{s-1} + \mu \Delta_1 \sum_{j=0}^{s-2} \lambda^j}{\lambda^{s-1} + \mu \Delta_2 \sum_{j=0}^{s-2} \lambda^j}, u_{6n+5} = C \frac{\Delta_0^{2n+1}}{\Delta_1^{2n+1}} \frac{\prod_{s=2}^{2n+1} \left( \lambda^{s-1} + \mu \Delta_2 \sum_{j=0}^{s-2} \lambda^j \right)}{\prod_{s=1}^{2n+1} \left( \lambda^s + \mu \Delta_0 \sum_{j=0}^{s-1} \lambda^j \right)},
\end{aligned} \tag{66}$$

where  $\Delta_i = u_i u_{i+1} u_{i+2}$ . Equations in (66) give the exact solution of (5) for any real values of  $\lambda$  and  $\mu$  provided that the denominators do not vanish. Recall that we acted the shift operator on (4) to get (5). Hence, the solutions of (4) are obtained, using (66), as follows:

$$\begin{aligned}
x_{6n-4} &= \frac{d^{2n}}{a^{2n-1}} \prod_{s=2}^{2n} \frac{\lambda^{s-1} + \mu abc \sum_{j=0}^{s-2} \lambda^j}{\lambda^{s-1} + \mu bcd \sum_{j=0}^{s-2} \lambda^j}, x_{6n-3} = \frac{e^{2n}}{b^{2n-1}} \prod_{s=2}^{2n} \frac{\lambda^{s-1} + \mu bcd \sum_{j=0}^{s-2} \lambda^j}{\lambda^{s-1} + \mu cde \sum_{j=0}^{s-2} \lambda^j}, \\
x_{6n-2} &= \frac{c(ab)^{2n}}{(de)^{2n}} \frac{\prod_{s=2}^{2n} \left( \lambda^{s-1} + \mu cde \sum_{j=0}^{s-2} \lambda^j \right)}{\prod_{s=1}^{2n} \left( \lambda^s + \mu abc \sum_{j=0}^{s-1} \lambda^j \right)}, x_{6n-1} = \frac{d^{2n+1}}{a^{2n}} \prod_{s=2}^{2n+1} \frac{\lambda^{s-1} + \mu abc \sum_{j=0}^{s-2} \lambda^j}{\lambda^{s-1} + \mu bcd \sum_{j=0}^{s-2} \lambda^j}, \\
x_{6n} &= \frac{e^{2n+1}}{b^{2n}} \prod_{s=2}^{2n+1} \frac{\lambda^{s-1} + \mu bcd \sum_{j=0}^{s-2} \lambda^j}{\lambda^{s-1} + \mu cde \sum_{j=0}^{s-2} \lambda^j}, x_{6n+1} = \frac{c(ab)^{2n+1}}{(de)^{2n+1}} \frac{\prod_{s=2}^{2n+1} \left[ \lambda^{s-1} + \mu cde \sum_{j=0}^{s-2} \lambda^j \right]}{\prod_{s=1}^{2n+1} \left( \lambda^s + \mu abc \sum_{j=0}^{s-1} \lambda^j \right)},
\end{aligned} \tag{67}$$

for any real values of  $\lambda$  and  $\mu$  as long as the denominators do not vanish.



- When  $\lambda = 1$  and  $\mu = 1$ , equations in (67) yield the results obtained by Yasin Yazlik in Theorem 5 in [9] for

$$x_{n+1} = \frac{x_{n-2}x_{n-3}x_{n-4}}{x_n x_{n-1}(1 + x_{n-2}x_{n-3}x_{n-4})}, \quad n = 0, 1, 2, \dots \quad (68)$$

where  $a, b, c, c, d$  and  $e$  are positive real numbers.

- When  $\lambda = 1$  and  $\mu = -1$ , equations in (67) yield the results obtained by Yasin Yazlik in Theorem 9 in [9] for

$$x_{n+1} = \frac{x_{n-2}x_{n-3}x_{n-4}}{x_n x_{n-1}(1 - x_{n-2}x_{n-3}x_{n-4})}, \quad n = 0, 1, 2, \dots \quad (69)$$

where  $a, b, c, c, d$  and  $e$  are positive real numbers with  $abc \neq 1$  and  $cde \neq 1$ .

**Note.** There should not be a minus sign right after the expression of  $x_{3n-2}$  in Theorem 9 in [9].

- When  $\lambda = -1$  and  $\mu = 1$ , equations in (67) yield the results obtained by Yasin Yazlik in Theorem 7 in [9] for

$$x_{n+1} = \frac{x_{n-2}x_{n-3}x_{n-4}}{x_n x_{n-1}(-1 + x_{n-2}x_{n-3}x_{n-4})}, \quad n = 0, 1, 2, \dots \quad (70)$$

where  $a, b, c, d$  and  $e$  are non zero real numbers with  $abc \neq 1$ ,  $bcd \neq 1$  and  $cde \neq 1$ .

- When  $\lambda = -1$  and  $\mu = -1$ , equations in (67) yield the results obtained by Yasin Yazlik in Theorem 11 in [9] for

$$x_{n+1} = \frac{x_{n-2}x_{n-3}x_{n-4}}{x_n x_{n-1}(-1 - x_{n-2}x_{n-3}x_{n-4})}, \quad n = 0, 1, 2, \dots \quad (71)$$

where  $a, b, c, c, d$  and  $e$  are non zero real numbers with  $abc \neq -1$ ,  $bcd \neq -1$  and  $cde \neq -1$ .

**Note.** There should be a minus sign right after the expression of  $x_{6n+1}$  in Theorem 11 in [9].

## 5 Conclusion

In this paper, we have obtained nontrivial symmetries of some rational ordinary difference equations and their exact solutions were obtained. Most importantly, we note that (21) gives a clear idea, without making any lucky guesses, for the most convenient choice of the invariant of (3).

## Conflict of interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

## Data availability statement

No data were used to support this study.

## Funding statement

The authors received no specific funding for this work.

## References

- [1] E. M. Elsayed, Dynamics of a rational recursive Sequence, *Int. J. Difference Equ.*, 4:2 (2009), 185-200.
- [2] M. Folly-Gbetoula, Symmetry, reductions and exact solutions of difference equations  $u_{n+2} = au_n/(1 + bu_nu_{n+1})$ , *J. Differ. Equations Appl.*, 23:6 (2017), 1017–1024.
- [3] M. Folly-Gbetoula and A.H. Kara, Symmetries, conservation laws, and integrability of difference equations, *Adv. Difference Equ.*, 2014, 2014.
- [4] P. E. Hydon, *Difference Equations by Differential Equation Methods*, Cambridge University Press, Cambridge, 2014.
- [5] N. Joshi and P. Vassiliou, The existence of Lie Symmetries for First-Order Analytic Discrete Dynamical Systems, *J. Math. Anal. Appl.*, 195 (1995), 872-887.
- [6] S. Lie, Klassifikation und Integration von gewöhnlichen Differentialgleichungen zwischen x, y, die eine Gruppe von Transformationen gestatten. I, *Math. Ann.*, 22 (1888), 213- 253.
- [7] E. Noether, Invariante variationsprobleme, *Mathematisch- Physikalische Klasse*, 2 (1918), 235–257.

- [8] S. Stevic, J. Diblik, B. Iricanin and Z. Smarda, On a fifth-order difference equation, *J. Computational Analysis and Applications*, 20:7 (2016), 1214–1227.
- [9] Yasin Yazlik, On the solutions and behavior of rational difference equations, *J. Computational Analysis and Applications*, 17:3 (2014), 584–594.