

MULTIFRACTAL FORMALISM FOR GENERALISED LOCAL DIMENSION SPECTRA OF GIBBS MEASURES ON THE REAL LINE

JOHANNES JAERISCH AND HIROKI SUMI

ABSTRACT. We extend the multifractal formalism for the local dimension spectrum of a Gibbs measure μ supported on the attractor Λ of a conformal iterated functions system on the real line. Namely, for $\alpha \in \mathbb{R}$, we establish the multifractal formalism for the Hausdorff dimension of the set of $x \in \Lambda$ for which the μ -measure of a ball of radius r_n centred at x obeys a power law r_n^α , for a sequence $r_n \rightarrow 0$. This allows us to investigate the Hölder regularity of various fractal functions, such as distribution functions and conjugacy maps associated with conformal iterated function systems.

1. INTRODUCTION AND STATEMENT OF RESULTS

Multifractal analysis has its origin in statistical physics (see [Man74, FP85, HJK⁺86]). For the mathematical theory of multifractal formalism and its relation to thermodynamic formalism, we recommend [Pes97]. In this article we extend the multifractal formalism for local dimension spectra of Gibbs measures supported on the attractor $\Lambda \subset \mathbb{R}$ of a conformal iterated function system on the real line. For a Borel measure μ on \mathbb{R} and with $B(x, r) := \{y \in \mathbb{R} \mid |y - x| < r\}$, the local dimension spectrum of μ is given by the level sets

$$\mathcal{F}(\alpha) := \left\{ x \in \Lambda \mid \lim_{r \rightarrow 0} \frac{\log \mu(B(x, r))}{\log r} = \alpha \right\}, \quad \alpha \in \mathbb{R}.$$

It is well known ([Pat97, PW97a, PW97b, Pes97]) for finitely generated contracting conformal iterated function systems satisfying the open set condition that, if μ is the Gibbs measure of a Hölder continuous potential, then the dimension spectrum f given by

$$f(\alpha) := \dim_H \mathcal{F}(\alpha)$$

is equal to the Legendre transformation of a certain function t defined implicitly by solving a topological pressure equation. To denote this state of affairs, we say that the multifractal formalism holds. In our setting, the function t will be given by equation (1.1) below. For related results on the multifractal formalism for non-uniformly hyperbolic systems and graph-directed constructions, we refer to [JR11, RU09].

For self-conformal measures of finitely generated contracting conformal iterated function systems it is also known ([BS00]) that the set of divergence points

$$\left\{ x \in \Lambda \mid \liminf_{r \rightarrow \infty} \frac{\log \mu(B(x, r))}{\log r} < \limsup_{r \rightarrow \infty} \frac{\log \mu(B(x, r))}{\log r} \right\}$$

has full Hausdorff dimension, unless the spectrum f is degenerate. In [LWX12] it is shown that, for $x \in \Lambda$, the set of accumulation points $A(x)$ of $(\log(r)^{-1} \log \mu(B(x, r)))_{r \geq 0}$ is either a singleton or a closed interval. Moreover, it is shown that

$$\dim_H \{A(x) = [a, b]\} = \inf_{\alpha \in [a, b]} \dim_H \mathcal{F}(\alpha).$$

Recently, the following modified level sets have attracted a lot of attention in studying the regularity of various fractal functions ([JS15], [JS18], [Ota18], [BKK18], [All18]). For $\alpha \in \mathbb{R}$, we define

$$R_*(\alpha) := \left\{ x \in \Lambda \mid \liminf_{r \rightarrow 0} \frac{\log \mu(B(x, r))}{\log r} = \alpha \right\}, \quad R^*(\alpha) := \left\{ x \in \Lambda \mid \limsup_{r \rightarrow 0} \frac{\log \mu(B(x, r))}{\log r} = \alpha \right\}.$$

In [JS15] it is shown that the level sets $R_*(\alpha)$ and $R^*(\alpha)$ satisfy the multifractal formalism in the context of semigroups of rational maps on the Riemann sphere satisfying the separating condition. Allaart ([All18]) proved the multifractal formalism for the level sets $R_*(\alpha)$ for self-similar measures satisfying the open set condition. In [JS18] we established the multifractal formalism for $R_*(\alpha)$ for self-conformal measures supported on attractors of conformal iterated function systems satisfying the open set condition. Allaart ([All18]) raised the question whether the multifractal formalism holds for $R^*(\alpha)$. Moreover, it is of interest whether the formalism for R_* and R^* extends to arbitrary Gibbs measures of Hölder continuous potentials.

In this article we establish the multifractal formalism for the generalised level sets

$$\mathcal{F}^*(\alpha) := \left\{ x \in \Lambda \mid \exists (r_k) \rightarrow \infty \lim_{k \rightarrow \infty} \frac{\log \mu(B(x, r_k))}{\log r_k} = \alpha \right\}, \quad \alpha \in \mathbb{R},$$

for Gibbs measures of Hölder continuous potentials supported on attractors of conformal iterated function systems satisfying the open set condition. Combining with the previously known results for $\mathcal{F}(\alpha)$, we thus obtain that the multifractal formalism holds in particular for the level sets $R_*(\alpha)$ and $R^*(\alpha)$.

Let us now introduce the necessary terminology to state our main theorem precisely. For an index set I with $\#I < \infty$ let $\Phi = (\phi_i)_{i \in I}$, $\phi_i : X \rightarrow X$, be a contracting $\mathcal{C}^{1+\varepsilon}$ conformal iterated function system on a compact subset $X \subset \mathbb{R}$. We refer to [MU96] for further details and basic properties. For simplicity, we will also assume that $\phi'_i > 0$ on X for each $i \in I$. Let $\pi : I^{\mathbb{N}} \rightarrow \mathbb{R}$ denote the coding map of Φ which is for $\omega = (\omega_1, \omega_2, \dots) \in I^{\mathbb{N}}$ given by

$$\bigcap_{n \geq 1} \phi_{\omega_1} \circ \dots \circ \phi_{\omega_n}(X) = \{\pi(\omega)\}.$$

Let $\Lambda := \pi(I^{\mathbb{N}})$. We say that Φ satisfies the open set condition if there exists a non-empty open interval $U \subset \mathbb{R}$ such that $\phi_i(U) \subset U$, for all $i \in I$, and $\phi_i(U) \cap \phi_j(U) = \emptyset$ for all $i, j \in I$ with $i \neq j$. To utilize the symbolic thermodynamic formalism (see [Bow75]) we will also need the following definitions. We denote by $\sigma : I^{\mathbb{N}} \rightarrow I^{\mathbb{N}}$ the left-shift on $I^{\mathbb{N}}$ which becomes a compact metric space endowed with the shift metric. Let $\varphi : I^{\mathbb{N}} \rightarrow \mathbb{R}$ be the geometric potential of Φ given by

$$\varphi(\omega) := \log \phi'_{\omega_1}(\pi(\sigma(\omega))), \quad \omega = (\omega_1, \omega_2, \dots) \in I^{\mathbb{N}}.$$

Note that φ is Hölder continuous. Let $\psi : I^{\mathbb{N}} \rightarrow \mathbb{R}$ be Hölder continuous and let μ_ψ denote the unique Gibbs measure for ψ in the sense of Bowen ([Bow75]). For $n \geq 1$ we denote by $S_n \psi := \sum_{k=0}^{n-1} \psi \circ \sigma^k$ the ergodic sum. The range of the multifractal spectrum is defined by

$$\alpha_- := \inf_{\omega \in I^{\mathbb{N}}} \liminf_{n \rightarrow \infty} \frac{S_n \psi(\omega)}{S_n \varphi(\omega)}, \quad \alpha_+ := \sup_{\omega \in I^{\mathbb{N}}} \limsup_{n \rightarrow \infty} \frac{S_n \psi(\omega)}{S_n \varphi(\omega)}.$$

Recall that $\alpha_- \leq \dim_H \Lambda$ with equality if and only if $\alpha_- = \alpha_+$ (see [Pes97]). Since $\varphi < 0$ we have that for each $\beta \in \mathbb{R}$ there exists a unique $t(\beta)$ such that

$$(1.1) \quad \mathcal{P}(t(\beta)\varphi + \beta\psi) = 0,$$

where $\mathcal{P}(f)$ refers to the topological pressure of a continuous function $f : I^{\mathbb{N}} \rightarrow \mathbb{R}$ with respect to σ . We denote by

$$t^*(\alpha) := \sup_{x \in \mathbb{R}} (\alpha x - t(x))$$

the Legendre transform of t .

Theorem 1.1. *Let $\Phi = (\phi_i : X \rightarrow X)_{i \in I}$ be a finitely generated $\mathcal{C}^{1+\varepsilon}$ conformal iterated function system on $X \subset \mathbb{R}$ satisfying the open set condition. Let $\psi : I^{\mathbb{N}} \rightarrow \mathbb{R}$ be Hölder continuous and let $\mu := \mu_\psi \circ \pi^{-1}$. Then we have for all $\alpha \in [\alpha_-, \alpha_+]$,*

$$\dim_H R_*(\alpha) = \dim_H R^*(\alpha) = \dim_H \mathcal{F}(\alpha) = \dim_H \mathcal{F}^*(\alpha) = -t^*(-\alpha),$$

and for $\alpha \notin [\alpha_-, \alpha_+]$ we have $R_*(\alpha) = R^*(\alpha) = \mathcal{F}(\alpha) = \mathcal{F}^*(\alpha) = \emptyset$.

We proceed with two applications of our main result to the regularity of fractal functions.

1.1. Distribution functions of Gibbs measures. For a function $F : \mathbb{R} \rightarrow \mathbb{R}$ we define the pointwise Hölder exponent of F at $x \in \mathbb{R}$ by

$$\text{Höl}(F, x) := \sup \left\{ \alpha > 0 \mid \limsup_{y \rightarrow x} \frac{|F(y) - F(x)|}{|y - x|^\alpha} < \infty \right\}.$$

Under the assumptions of Theorem 1.1, we consider the distribution function of $\mu = \mu_\psi \circ \pi^{-1}$ given by

$$F_\mu : \mathbb{R} \rightarrow [0, 1], \quad F_\mu(x) := \mu((-\infty, x]).$$

It is shown in [JS15, Lemma 5.1] that

$$(1.2) \quad \text{Höl}(F_\mu, x) = \liminf_{r \rightarrow 0} \frac{\log \mu(B(x, r))}{\log r}.$$

Corollary 1.2. *Under the assumptions of Theorem 1.1, the distribution function $F_\mu : \mathbb{R} \rightarrow [0, 1]$ of $\mu = \mu_\psi \circ \pi^{-1}$ satisfies for every $\alpha \in [\alpha_-, \alpha_+]$*

$$\dim_H \{x \in \Lambda \mid \text{Höl}(F_\mu, x) = \alpha\} = -t^*(-\alpha),$$

where $t : \mathbb{R} \rightarrow \mathbb{R}$ is defined implicitly by $\mathcal{P}(t(\beta)\phi + \beta\psi) = 0$.

For results on points of non-differentiability and related properties of F_μ we refer to [Fal04, KS09, Tro14].

1.2. Conjugacy maps between expanding piecewise $\mathcal{C}^{1+\varepsilon}$ interval maps. In this section we apply the multifractal formalism to conjugacy maps between expanding piecewise $\mathcal{C}^{1+\varepsilon}$ interval maps as considered in [JKPS09]. In fact, we slightly extend the framework by allowing one of the repellers of the expanding interval maps to be a proper subset of $[0, 1]$, whereas in [JKPS09] both repellers are equal to $[0, 1]$.

Let us now briefly introduce the setting. Let f be an expanding piecewise $\mathcal{C}^{1+\varepsilon}$ interval map with $s \geq 2$ full branches, i.e., there exist closed intervals $J_1, \dots, J_s \subset [0, 1]$ with non-empty, pairwise disjoint interiors such that $f|_{J_i}$ has a $\mathcal{C}^{1+\varepsilon}$ extension to a neighbourhood of J_i satisfying $f'_{|J_i} > 1$ and $f(J_i) = [0, 1]$, for $1 \leq i \leq s$. We will always assume that the intervals J_1, \dots, J_s are given in increasing order (i.e., $\sup J_i \leq \inf J_j$ if $i < j$). The repeller of f is denoted by Λ and the restriction $f|_\Lambda : \Lambda \rightarrow \Lambda$ is conjugate to the shift $\sigma : I^{\mathbb{N}} \rightarrow I^{\mathbb{N}}$ with $I := \{1, \dots, s\}$. The conjugacy is given by the coding map $\pi_f : I^{\mathbb{N}} \rightarrow \Lambda$ of the conformal iterated function systems Φ_f , which is given by the contracting inverse branches $(f|_{J_i})^{-1} : [0, 1] \rightarrow [0, 1]$, $1 \leq i \leq s$. Similarly, let $g : [0, 1] \rightarrow [0, 1]$ be an expanding $\mathcal{C}^{1+\varepsilon}$ interval map with s full branches. We assume that the repeller of g is the interval $[0, 1]$. Again, g is conjugate to $(I^{\mathbb{N}}, \sigma)$ with conjugacy map given by the coding map $\pi_g : I^{\mathbb{N}} \rightarrow [0, 1]$ of the associated conformal iterated function system Φ_g . Thus, there is a natural conjugacy map $\Theta : \Lambda \rightarrow [0, 1]$ given by $\Theta := \pi_g \circ \pi_f^{-1}$ satisfying $\Theta \circ f|_\Lambda = g \circ \Theta$ (see [JKPS09] for the case $\Lambda = [0, 1]$). Note that $\#(\pi_f^{-1}(x)) \leq 2$ and that $\pi_g \circ \pi_f^{-1}$ is a well-defined non-decreasing function on the real line, which is strictly increasing function on Λ . In particular, we have for $x \in \Lambda$ that

$$(1.3) \quad \pi_g^{-1} \left((-\infty, \pi_g \circ \pi_f^{-1}(x)] \right) = \pi_f^{-1}((-\infty, x]).$$

Denote by $\varphi_g : I^{\mathbb{N}} \rightarrow \mathbb{R}$ (resp. $\varphi_f : I^{\mathbb{N}} \rightarrow \mathbb{R}$) the geometric potential of Φ_g (resp. Φ_f) given by

$$\varphi_g := -\log g' \circ \pi_g, \quad \varphi_f := -\log f' \circ \pi_f.$$

Let λ denote the Lebesgue measure on $[0, 1]$ and recall that $\lambda = \mu_{\varphi_g} \circ \pi_g^{-1}$ where μ_{φ_g} is the unique Gibbs measure for φ_g on $I^{\mathbb{N}}$. By (1.3) we then have for $x \in \Lambda$,

$$\Theta(x) = \mu_{\varphi_g} \circ \pi_g^{-1}((-\infty, \Theta(x)]) = \mu_{\varphi_g} \circ \pi_g^{-1}((-\infty, \pi_g \circ \pi_f^{-1}(x)]) = \mu_{\varphi_g} \circ \pi_f^{-1}((-\infty, x]).$$

So, the conjugacy $\Theta : \Lambda \rightarrow [0, 1]$ coincides with the distribution function of $\mu_{\varphi_g} \circ \pi_f^{-1}$ (cf. [JKPS09] for the case $\Lambda = [0, 1]$). Therefore, we have

$$\{x \in \Lambda \mid \text{Höl}(\Theta, x) = \alpha\} = \{x \in \Lambda \mid \text{Höl}(F_{\mu}, x) = \alpha\},$$

where $\mu = \mu_{\varphi_g} \circ \pi_f^{-1}$. Hence, with

$$\alpha_- = \inf_{\omega \in I^{\mathbb{N}}} \liminf_{n \rightarrow \infty} \frac{S_n \varphi_g(\omega)}{S_n \varphi_f(\omega)}, \quad \alpha_+ := \sup_{\omega \in I^{\mathbb{N}}} \limsup_{n \rightarrow \infty} \frac{S_n \varphi_g(\omega)}{S_n \varphi_f(\omega)},$$

we obtain the following corollary as a consequence of Corollary 1.2.

Corollary 1.3. *Let f and g be two expanding piecewise $\mathcal{C}^{1+\varepsilon}$ interval maps with $s \geq 2$ full branches and coding maps $\pi_f, \pi_g : I^{\mathbb{N}} \rightarrow [0, 1]$. Let $\Lambda := \pi_f(I^{\mathbb{N}})$ and suppose that $\pi_g(I^{\mathbb{N}}) = [0, 1]$. Then the conjugacy map $\Theta : \Lambda \rightarrow [0, 1]$, given by $\Theta := \pi_g \circ \pi_f^{-1}$, satisfies for every $\alpha \in [\alpha_-, \alpha_+]$*

$$\dim_H \{x \in \Lambda \mid \text{Höl}(\Theta, x) = \alpha\} = -t^*(-\alpha),$$

where $t : \mathbb{R} \rightarrow \mathbb{R}$ is defined implicitly by $\mathcal{P}(t(\beta)\varphi_f + \beta\varphi_g) = 0$.

2. PROOF OF THEOREM 1.1

Proof. First observe that $\mathcal{F}(\alpha)$ is a subset of each of the level sets considered in the theorem. Therefore, the lower bound for the Hausdorff dimension follows from the well-known multifractal formalism for $\mathcal{F}(\alpha)$. Therefore, to complete the proof of the theorem, it suffices to show the upper bound for the Hausdorff dimension of $\mathcal{F}^*(\alpha)$. Throughout, we may assume $I = \{1, \dots, s\}$, for $s \geq 2$, and $\phi_i(x) \leq \phi_j(y)$ for all $i, j \in I$ with $i < j$ and for all $x, y \in U$, where U is the open set in the open set condition.

Denote by $I^* := \bigcup_{k \geq 1} I^k$ the set of finite words in the alphabet I . We first observe that there exists $C > 1$ such that for all $\gamma \in I^*$ and $i \in I$,

$$C \cdot \text{diam}(\pi[\gamma i]) \geq \text{diam}(\pi[\gamma]),$$

where $\text{diam}(A) := \sup\{|x - y| \mid x, y \in A\}$ refers to the diameter of a set $A \subset \mathbb{R}$.

Let $x \in \mathcal{F}^*(\alpha)$, $x = \pi(\omega)$ for some $\omega \in I^{\mathbb{N}}$. There exists a sequence $r_k \rightarrow 0$ such that

$$\lim_{k \rightarrow \infty} \frac{\log \mu(B(x, r_k))}{\log r_k} = \alpha.$$

We define for $k \geq 1$

$$n_k := \min\{n \geq 1 \mid \text{diam}(\pi[\omega_1, \dots, \omega_n]) < Cr_k\}.$$

For $a \in I$ and $m \in \mathbb{N}$ we denote $a^m := (a, \dots, a) \in I^m$. We define two sequences of words $(v_k), (v'_k) \in I^*$, $k \geq 1$, as follows. If $(\omega_1, \dots, \omega_{n_k})$ takes the form

$$(\omega_1, \dots, \omega_{n_k}) = (\tau j s^{\ell_k}),$$

for some $\tau \in I^*$, $j \leq s-1$ and $\ell_k \geq 1$ then let

$$\ell'_k := \max \left\{ l \geq 1 \mid \text{diam} \left(\pi \left[\tau(j+1)1^l \right] \right) \geq r_k \right\}.$$

We remark that, if the constant C above is large enough, then ℓ'_k is well defined. This follows because our definition of n_k implies $\text{diam}(\pi[\tau j]) \geq Cr_k$. We then define

$$\mathbf{v}_k := (\tau j s^{\ell_k - 1}) = (\omega_1, \dots, \omega_{n_k - 1}), \quad \mathbf{v}'_k := (\tau(j+1)1^{\ell'_k}).$$

If $(\omega_1, \dots, \omega_{n_k})$ takes the form $(\omega_1, \dots, \omega_{n_k}) = (\tau(j+1)1^{\ell_k})$, then we define $\mathbf{v}_k := (\omega_1, \dots, \omega_{n_k - 1})$ and $\mathbf{v}'_k := (\tau j s^{\ell_k})$ in a similar fashion. Finally, if $(\omega_1, \dots, \omega_{n_k})$ satisfies $\omega_{n_k} \notin \{1, s\}$ then we define $\mathbf{v}_k := \mathbf{v}'_k := (\omega_1, \dots, \omega_{n_k - 1})$. Let $n'_k := |\mathbf{v}'_k|$.

It is important to note that, by the definition of \mathbf{v}_k and \mathbf{v}'_k , we have as $k \rightarrow \infty$,

$$(2.1) \quad \text{diam}(\pi[\mathbf{v}_k]) \asymp \text{diam}(\pi[\mathbf{v}'_k]) \asymp r_k,$$

where, for sequences of positive numbers (a_k) and (b_k) the notation $a_k \asymp b_k$ means that a_k/b_k is bounded away from zero and infinity. We will show that this implies the existence of a uniform constant D such that

$$(2.2) \quad \ell'_k \cdot \varphi(\bar{1}) - D \leq \ell_k \cdot \varphi(\bar{s}) \leq \ell'_k \cdot \varphi(\bar{1}) + D,$$

where we have set $\bar{\gamma} := (\gamma\gamma\dots) \in I^{\mathbb{N}}$, for $\gamma \in I^*$. To prove (2.2) suppose that $(\omega_1, \dots, \omega_{n_k}) = (\tau j s^{\ell_k})$. The other case $(\omega_1, \dots, \omega_{n_k}) = (\tau(j+1)1^{\ell_k})$, can be handled analogously. By the bounded distortion property of the geometric potential we have, as $k \rightarrow \infty$,

$$\begin{aligned} \text{diam}(\pi[\mathbf{v}_k]) &= \text{diam} \left(\pi \left[(\tau j s^{\ell_k - 1}) \right] \right) \asymp \text{diam}(\pi[\tau]) e^{S_{\ell_k} \varphi(\bar{s})} \\ \text{diam}(\pi[\mathbf{v}'_k]) &= \text{diam} \left(\pi \left[\tau(j+1)1^{\ell'_k} \right] \right) \asymp \text{diam}(\pi[\tau]) e^{S_{\ell'_k} \varphi(\bar{1})}, \end{aligned}$$

which proves (2.2).

We will show that for $x \in \Lambda = \pi(I^{\mathbb{N}})$ and $k \geq 1$ we have

$$B(x, r_k) \cap \Lambda \subset \pi([\mathbf{v}_k]) \cup \pi([\mathbf{v}'_k]).$$

First suppose that $(\omega_1, \dots, \omega_{n_k}) = (\tau j s^{\ell_k})$. Then $\pi[\mathbf{v}_k] = \pi[(\tau j s^{\ell_k - 1})] \supset \pi[(\tau j s^{\ell_k - 1} 1)]$. Since $x \in \pi[(\tau j s^{\ell_k})]$ we have $x \geq \max \pi[(\tau j s^{\ell_k - 1} 1)]$. By the definition of C and n_k we have

$$\text{diam} \left(\pi \left[(\tau j s^{\ell_k - 1} 1) \right] \right) \geq C^{-1} \text{diam} \left(\pi \left[(\tau j s^{\ell_k - 1}) \right] \right) = C^{-1} \text{diam}(\pi[(\omega_1, \dots, \omega_{n_k - 1})]) \geq r_k.$$

Hence,

$$\pi[\mathbf{v}_k] \supset [x, x - r_k] \cap \Lambda.$$

Further, by the definition of ℓ'_k we have

$$\text{diam}(\pi[\mathbf{v}'_k]) = \text{diam} \left(\pi \left[\tau(j+1)1^{\ell'_k} \right] \right) \geq r_k,$$

so $[x, x + r_k] \cap \Lambda \subset \pi[\mathbf{v}_k] \cup \pi[\mathbf{v}'_k]$. This proves that $B(x, r_k) \cap \Lambda \subset \pi([\mathbf{v}_k]) \cup \pi([\mathbf{v}'_k])$.

Let $\varepsilon > 0$. We will derive from our assumption $x \in \mathcal{F}^*(\alpha)$ that there exists $N \geq 1$ such that for all $k \geq N$,

$$(2.3) \quad \frac{S_{|\mathbf{v}_k|} \Psi(\overline{\mathbf{v}_k})}{S_{|\mathbf{v}_k|} \varphi(\overline{\mathbf{v}_k})} \leq \alpha + \varepsilon \quad \text{or} \quad \frac{S_{|\mathbf{v}'_k|} \Psi(\overline{\mathbf{v}'_k})}{S_{|\mathbf{v}'_k|} \varphi(\overline{\mathbf{v}'_k})} \leq \alpha + \varepsilon.$$

To prove (2.3), we first note that by the Gibbs property of μ_ψ we have for every $\gamma \in I^*$,

$$\mu_\psi([\gamma]) \leq C_\psi \exp(S_{|\gamma|} \Psi(\bar{\gamma})),$$

where $C_\psi \geq 0$ is a uniform constant depending on ψ . Suppose for a contradiction that (2.3) does not hold. Then, by passing to a subsequence of (n_k) we may assume that for all k and for all $v \in \{v_k, v'_k\}$ we have $S_{|v|}\psi(\bar{v})/S_{|v|}\varphi(\bar{v}) > \alpha + \varepsilon$, and so, taking C' large enough, we have

$$\mu_\psi([v]) \leq C_\psi \exp(S_{|v|}\psi(\bar{v})) \leq C' r_k^{\alpha+\varepsilon}.$$

Since $B(x, r_k) \subset \pi([v_k]) \cup \pi([v'_k])$, we conclude that

$$\lim_{k \rightarrow \infty} \frac{\log \mu(B(x, r_k))}{\log r_k} \geq \alpha + \varepsilon.$$

This contradiction proves (2.3).

We will prove the proposition only in the case when $\alpha \leq \alpha_0 := \int \psi d\mu_{t(0)\varphi} / \int \varphi d\mu_{t(0)\varphi}$. The case $\alpha \geq \alpha_0$ can be considered in a similar fashion (see also Remark 2.1 below). Let $\beta > 0$, $\eta > 0$ and let $b = t(\beta) + \beta(\alpha + \varepsilon) + \eta$. We define

$$\mathcal{C}_{\alpha+\varepsilon} := \left\{ \tau \in I^* \mid \frac{S_{|\tau|}\psi(\bar{\tau})}{S_{|\tau|}\varphi(\bar{\tau})} \leq \alpha + \varepsilon \right\}.$$

We obtain a covering \mathcal{C} of $\mathcal{F}^*(\alpha)$ by cylinders of sufficiently small diameters as follows. For each $x \in \mathcal{F}^*(\alpha)$ we define the sequence n_k as above and define

$$\mathcal{C} := \{v_k \mid k \in \mathbb{N}\},$$

where $v_k \in I^*$ is defined as above satisfying $x \in \pi([v_k])$ and (2.3). To verify that the corresponding sum of diameters $\sum_{v \in \mathcal{C}} \text{diam}(\pi([v]))^b$ converges, we proceed as follows. If $v \notin \mathcal{C}_{\alpha+\varepsilon}$ then, by (2.3), we can replace v by $v' \in \mathcal{C}_{\alpha+\varepsilon}$, because $\text{diam}(\pi([v])) \asymp \text{diam}(\pi([v']))$ by (2.1). This defines a map $v \mapsto v'$ from $\mathcal{C} \setminus \mathcal{C}_{\alpha+\varepsilon}$ to $\mathcal{C}_{\alpha+\varepsilon}$. Since the involved numbers ℓ_k and ℓ'_k in the definition of v and v' satisfy (2.2), we have that the map $v \mapsto v'$ is at most M -to-1 for some uniform constant $M \in \mathbb{N}$. Since

$$(2.4) \quad \sum_{\omega \in \mathcal{C}_{\alpha+\varepsilon}} \text{diam}(\pi([\omega]))^b \asymp \sum_{\omega \in \mathcal{C}_{\alpha+\varepsilon}} e^{(t(\beta)+\beta(\alpha+\varepsilon)+\eta)S_{|\omega|}\varphi(\bar{\omega})} \leq \sum_{\omega \in \mathcal{C}_{\alpha+\varepsilon}} e^{(t(\beta)+\eta)S_{|\omega|}\varphi(\bar{\omega})+\beta S_{|\omega|}\psi(\bar{\omega})} < \infty,$$

we therefore conclude that the b -dimensional Hausdorff measure of $\mathcal{F}^*(\alpha)$ is finite. Now, first assume that $\alpha \in [\alpha_-, \alpha_0]$. Since ε and η are arbitrary, it follows that

$$\dim_H \mathcal{F}^*(\alpha) \leq \inf_{\beta > 0} \{t(\beta) + \beta\alpha\} = -t^*(-\alpha),$$

where we have used $\alpha \leq \alpha_0$ for the last equality. Finally, if $\alpha < \alpha_-$ then $\mathcal{C}_{\alpha+\varepsilon} = \emptyset$ if $\alpha + \varepsilon < \alpha_-$. By the above construction of the covering of $\mathcal{F}^*(\alpha)$ it thus follows that $\mathcal{F}^*(\alpha) = \emptyset$. The proof is complete. \square

Remark 2.1. For the dimension spectrum of the level sets $R_*(\alpha)$ with $\alpha \geq \alpha_0$, the upper bound of the Hausdorff dimension can also be deduced from [LWX12, Theorem 1.1 (2)] (see also [All18, Proof of Proposition 7.6 ii]).

REFERENCES

- [All18] P. C. Allaart, *Differentiability and Hölder spectra of a class of self-affine functions*, Adv. Math. **328** (2018), 1–39. MR 3771122
- [BKK18] B. Bárány, G. Kiss, and I. Kolossváry, *Pointwise regularity of parameterized affine zipper fractal curves*, Nonlinearity **31** (2018), no. 5, 1705–1733. MR 3816651
- [Bow75] R. Bowen, *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, Springer-Verlag, Berlin, 1975, Lecture Notes in Mathematics, Vol. 470. MR MR0442989 (56 #1364)
- [BS00] L. Barreira and J. Schmeling, *Sets of “non-typical” points have full topological entropy and full Hausdorff dimension*, Israel J. Math. **116** (2000), 29–70. MR MR1759398 (2002d:37040)
- [Fal04] K. Falconer, *One-sided multifractal analysis and points of non-differentiability of devil’s staircases*, Math. Proc. Cambridge Philos. Soc. **136** (2004), no. 1, 167–174. MR 2034020 (2005b:28013)

- [FP85] U. Frisch and G. Parisi, *On the singularity structure of fully developed turbulence*, Turbulence and predictability in geophysical fluid dynamics and climate dynamics (North Holland Amsterdam), 1985, pp. 84–88.
- [HJK⁺86] T. C. Halsey, M. H. Jensen, L. P. Kadanoff, I. Procaccia, and B. J. Shraiman, *Fractal measures and their singularities: The characterization of strange sets*, Phys. Rev. A **85** (1986), no. 33, 1141–1151.
- [JKPS09] T. Jordan, M. Kesseböhmer, M. Pollicott, and B. O. Stratmann, *Sets of nondifferentiability for conjugacies between expanding interval maps*, Fund. Math. **206** (2009), 161–183. MR 2576266
- [JR11] T. Jordan and M. Rams, *Multifractal analysis of weak Gibbs measures for non-uniformly expanding C^1 maps*, Ergodic Theory Dynam. Systems **31** (2011), no. 1, 143–164. MR 2755925
- [JS15] J. Jaerisch and H. Sumi, *Multifractal formalism for expanding rational semigroups and random complex dynamical systems*, Nonlinearity **28** (2015), 2913–2938.
- [JS18] ———, *Spectral gap property for random dynamics on the real line and multifractal analysis of generalised takagi functions*, Preprint available on arXiv:1808.08421 (2018).
- [KS09] M. Kesseböhmer and B. O. Stratmann, *Hölder -differentiability of Gibbs distribution functions*, Math. Proc. Cambridge Philos. Soc. **147** (2009), no. 2, 489–503. MR MR2525939
- [LWX12] J. Li, M. Wu, and Y. Xiong, *Hausdorff dimensions of the divergence points of self-similar measures with the open set condition*, Nonlinearity **25** (2012), no. 1, 93–105. MR 2864378
- [Man74] B. B. Mandelbrot, *Intermittent turbulence in self-similar cascades: divergence of high moments and dimension of the carrier*, Journal of Fluid Mechanics Digital Archive **62** (1974), no. 02, 331–358.
- [MU96] R. D. Mauldin and M. Urbański, *Dimensions and measures in infinite iterated function systems*, Proc. London Math. Soc. (3) **73** (1996), no. 1, 105–154. MR 1387085 (97c:28020)
- [Ota18] A. Otani, *Fractal dimensions of graph of Weierstrass-type function and local Hölder exponent spectra*, Nonlinearity **31** (2018), no. 1, 263–292. MR 3746637
- [Pat97] N. Patzschke, *Self-conformal multifractal measures*, Adv. in Appl. Math. **19** (1997), no. 4, 486–513. MR 1479016
- [Pes97] Y. B. Pesin, *Dimension theory in dynamical systems*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1997, Contemporary views and applications. MR MR1489237 (99b:58003)
- [PW97a] Y. Pesin and H. Weiss, *A multifractal analysis of equilibrium measures for conformal expanding maps and Moran-like geometric constructions*, J. Statist. Phys. **86** (1997), no. 1-2, 233–275. MR 1435198 (97m:58118)
- [PW97b] ———, *The multifractal analysis of Gibbs measures: motivation, mathematical foundation, and examples*, Chaos **7** (1997), no. 1, 89–106. MR 1439809 (98e:58130)
- [RU09] M. Roy and M. Urbański, *Multifractal analysis for conformal graph directed Markov systems*, Discrete Contin. Dyn. Syst. **25** (2009), no. 2, 627–650. MR 2525196
- [Tro14] S. Troscheit, *Hölder differentiability of self-conformal devil's staircases*, Math. Proc. Cambridge Philos. Soc. **156** (2014), no. 2, 295–311. MR 3177871

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ENGINEERING, SHIMANE UNIVERSITY, NISHIKAWATSU 1060, MATSUE, SHIMANE, 690-8504 JAPAN

E-mail address: jaerisch@riko.shimane-u.ac.jp

URL: <http://www.math.shimane-u.ac.jp/~jaerisch>

COURSE OF MATHEMATICAL SCIENCE, DEPARTMENT OF HUMAN COEXISTENCE, GRADUATE SCHOOL OF HUMAN AND ENVIRONMENTAL STUDIES, KYOTO UNIVERSITY YOSHIDA-NIHONMATSU-CHO, SAKYO-KU, KYOTO 606-8501, JAPAN

E-mail address: sumi@math.h.kyoto-u.ac.jp

URL: <http://www.math.h.kyoto-u.ac.jp/~sumi/index.html>