

THE SOLUTIONS OF THE 3RD AND 4TH CLAY MILLENNIUM PROBLEMS

The first about the P vs NP in computational complexity and the second about the Navier-Stokes equations

Konstantinos E. Kyritsis

PROLOGUE

In this treatise we present the solutions of the 3rd Clay Millennium problem P vs NP in the Computational Complexity and the 4th Clay Millennium problem in classical fluid dynamics about the Navier-Stokes equations.

Some initial but incorrect solutions of the 3rd Clay Millennium problem about P vs NP had already been published by me in *International Journal of Pure and Applied Mathematics Volume 120 No. 3 2018, pp 497-510* ISSN: 1311-8080 (printed version); ISSN: 1314-3395 (on-line version) url: <http://www.ijpam.eu> doi: 10.12732/ijpam.v120i3.1

But also, in:

Kyritsis C. *On the solution of the 3rd Clay Millennium problem. A short and elegant proof that $P \neq NP$ in the context of deterministic Turing machines and Zermelo-Frankel set theory.* Proceedings of the first ICQSBEI 2017 conference, Athens, Greece, <https://books.google.gr/books?id=BSUsDwAAQBAJ&pg> pp 170-181

And also, in:

Kyritsis K. Review of the Solutions of the Clay Millennium Problem about $P \neq NP = EXPTIME$ World Journal of Research and Review (WJRR) ISSN:2455-3956, Volume-13, Issue-3, September 2021 Pages 21-26

But also in Chapter 3 a 3rd and drastically shorter solution which was presented in the in the 6th International conference on quantitative, social, biomedical and economic issues, ICQSBE 2022 1st July 2022,

<https://icqsbei2022.blogspot.com/2022/06/blog-post.html>

<http://books.google.com/books/about?id=xZnCEAAAQBAJ>

Eventually after discussions and lectures about P vs NP in the School of electrical and Computer engineering in the National Technical University of Athens in 2023, I corrected the solutions to two new solutions presented here

https://www.researchgate.net/publication/378592494_TWO_NEW_SOLUTIONS_OF_THE_P_VERSUS_NP_PROBLEM One theoretical another by counter-example by the Pell's Diophantine equation This is an extract from a lecture

and here

https://www.researchgate.net/publication/376170410_The_millennium_problem_Polynomial_complexity_versus_non-deterministic_polynomial_complexity_What_is_the_state_of_the_art_today_Ill_pose_d_aspects_of_the_problem_Example_of_a_reasonable_solution_Perspec

The solution of the 4th Clay Millennium problem about the Navier-Stokes equations is in at least two papers (Chapter 4,5,6) both of which have already been published

See KYRITSIS, K. August 2021. A solution of the 4th Clay Millennium problem about the Navier-Stokes equations. *World Journal of Research and Review WJRR.org* ISSN 2455-3956 Volume 13 Issue 20 August 21 pp 25-40

And:

Kyritsis, K. (2022) A Short and Simple Solution of the Millennium Problem about the Navier-Stokes Equations and Similarly for the Euler Equations. *Journal of Applied Mathematics and Physics*, **10**, 2538-2560. doi: 10.4236/jamp.2022.108172. <https://www.scirp.org/journal/paperinformation.aspx?paperid=119569>

Before the final solution of this millennium problem there was also a partial solution based on the hypotheses of conservation of particles here

Kyritsis, K. November 2017 “On the 4th Clay Millennium problem: Proof of the regularity of the solutions of the Euler and Navier-Stokes equations, based on the conservation of particles” *Journal of Scientific Research and Studies* Vol 4 (11) , pp304-317, November 2017.

It seems that at the beginning of each century has become a tradition to state a list of significant and usually difficult problems in the mathematics, that it is considered that their solution will advance significantly the mathematical sciences. At the beginning of the 20th century (1900) it was D. Hilbert who formulated and listed 23 problems that most of them have been solved till today (see e.g. https://en.wikipedia.org/wiki/Hilbert%27s_problems) . Those problems from the 23 that have been solved proved to be indeed critical for the overall evolution of mathematics and their applications. Continuing this tradition, the Clay Mathematical Institute formulated in 2000, 7 critical problems and this time there is a monetary award for their solution (see e.g. <http://www.claymath.org/millennium-problems>) . From them, the 6th problem (Poincare Hypothesis) it has been accepted that it has been solved by Grigoriy Perelman in 2003. It is not presented here a common or joint method of solution of the 3rd and 4th Clay millenniums problems. It is only because I am an interdisciplinary researcher that I have worked, on both of them. And of course, I had both the advantages and disadvantages of an interdisciplinary researcher. The disadvantage was that I had to sharpen by specialized knowledge in two different areas of Computer science and Mathematical physics , that specialist would not need not do it, while the advantage , that turned out to be more important, were that “I was not blinded by the trees so as to see the forest”; In other words I used new heuristic methods from other disciplines to discover the correct direction of solutions and afterwards I worked out a standard classical proof for each one of them. This is well known in the history of mathematics. E.g. Archimedes found at first the correct formulae of volumes of the sphere, cylinder etc with water, sand and balanced moments of forces experiments before he worked out logically complete proofs of them in the context of Euclidean geometry. Similarly, Newton discovered the laws of gravitation for earth, sun, moon etc. with his, at that time unpublished calculus of fluxes or infinitesimals, and then worked strict proofs within Euclidean geometry in his famous Principia Mathematica.

Similarly, I used myself a heuristic methodology based on statistical mechanics and the particle structure of fluids. Unfortunately, the mathematical models of the fluid dynamic within which this Millennium problem has been formulated are based on the concept of **infinite divisible matter** (before the discovery in the science of physics that matter consists from finite many atoms) and this is a main source of difficulty involving this problem.

Both problems had at least two different directions of solution. For the 3rd Clay Millennium problem, it is:

- 1) that the non-deterministic polynomial complexity symbolized by NP is equal to a polynomial complexity symbolized by P (in which case the usual setting of passwords and messages might be unsafe) or
- 2) to a higher e.g. EXPTIME (in which case the usual setting of passwords and messages is as expected to be safe). The heuristic analysis gave that it should hold $NP=EXPTIME$, which was eventually proved in two different ways.

And for the 4th Clay Millennium problem two different directions of solution would be that:

- 1) There exist a Blow-up of velocities in finite time.
- 2) No blow-up exist in finite time and the solutions of the Navier-Stokes equations are regular.

The heuristic analysis gave that because of finite initial energy and energy conservation there cannot be a Blow-up which was eventually proved within the context of classical fluid dynamics that allows for infinite limits etc. More on the logic and strategy of proof for each problem in the next two parts of this treatise.

TABLE OF CONTENTS

PART A.

CHAPTER 1

THE SOLUTION OF THE MILLENNIUM PROBLEM ABOUT
THE P vs NP IN COMPUTATIONAL COMPLEXITY. INFORMAL DISCUSSION
.....

CHAPTER 2-3

THE SOLUTION OF THE P VS NP MILLENNIUM PROBLEM WITH GRAPHS:
P $\neq$ NP. RUSSELL'S STANDARDS OF LOGICAL REASONING AGAINST
GOEDEL'S ARGUMENTS OF INCOMPLETENESS AND THE P VS NP
MILLENNIUM PROBLEM OF MATHEMATICAL COMPUTATION.
.....

.....

PART B

CHAPTER 4

THE SOLUTION OF THE MILLENNIUM PROBLEM
ABOUT THE NAVIER STOKES EQUATIONS

CHAPTER 5

ON THE SOLUTION OF THE 4TH MILLENNIUM PROBLEM.
PROOF OF THE REGULARITY OF THE SOLUTIONS OF THE EULER
AND NAVIER-STOKES EQUATIONS,
BASED ON THE CONSERVATION OF PARTICLES

CHAPTER 6

A SHORT AND SIMPLE SOLUTION OF THE 4TH CLAY MILLENNIUM
PROBLEM ABOUT THE NAVIER-STOKES EQUATIONS
AND SIMILARLY FOR THE EULER EQUATIONS.....

PART A.

CHAPTER 1

THE SOLUTION OF THE MILLENNIUM PROBLEM ABOUT THE P vs NP IN COMPUTATIONAL COMPLEXITY. INFORMAL DISCUSSION

Prologue

The standard formulation of the 3rd Clay Millennium problem can be found in **(Cook, Stephen April 2000 *The P versus NP Problem (PDF)*, Clay Mathematics Institute site.** <http://www.claymath.org/millennium-problems/p-vs-np-problem>

<http://www.claymath.org/sites/default/files/pvsnp.pdf>)

1) The P versus NP is a difficult problem, that has troubled the scientific community for some decades

2) It may have simple proofs of a few paragraphs, hopefully not longer than the proof of the Time Hierarchy theorem, which seems to be a deeper result.

3) But it can also have very lengthily and complex proofs, that may take dozens of pages.

What the final proof in the next published is or is not:

1) It does not introduce new theoretical concepts in computational complexity theory so as to solve the P versus NP.

2) It does not use relativization and oracles

3) It does not use diagonalization arguments, although the main proof, utilizes results from the time hierarchy theorem

4) It is not based on improvements of previous bounds of complexity on circuits

5) It is proved with the method of counter-example. Thus, it is transparent short and “simple”. It takes any Exptime-complete DTM decision problem, and from it, it derives in the context of deterministic Turing machines a decision problem language

which it is apparent that it belongs in the NP class decision problems while it does not belong the class P of decision problems.

6) It seems a “simple” proof because it chooses the right context to make the arguments and constructions and the key-abstraction mentioned above. So, it helps that the scientific community will accept that this 3rd Clay Millennium problem has already been solved.

RS0) The disruptive role for the computational complexity of the solution of the millennium problem P vs NP. The Russell’s disallowed impredicative predicates and contradictions.

My odyssey when I tried to re-solve the mystery of P vs NP in the context of computational complexity.

(In the symbolism of paragraphs we utilize RS meaning resolution)

MY INITIAL APPROACH IN MY ODYSSEY

When I finished solving the Millennium problem about the Navier-Stokes equations in fluid dynamics, in 2017 (See [7]) I started trying to solve also the P vs NP millennium problem. I was afraid that it would be more difficult compared to that in fluid dynamics, because fluid dynamics is a centuries old specialization of mathematics and it has proved practically all the necessary tools about it. On the other hand Computational Complexity was not really more than 50-years old, and it was expected that it has not proved yet all its necessary tools and results that are essential for this specialization. But I never expected my Odyssey that followed and the disruptive role that the solution of the millennium problem P vs NP would have for Computational Complexity.

Initially in 2017 I thought that I proved that $NP=EXPTIME$ thus P is not equal to NP (see e.g. [4], [5] , [6] .[8])

But when I lectured the solution in National Technical University of Athens in 2023, I discovered that it had an error.

So I spent all my sabbatical of 2023-2024 in trying to re-solve it.

I was much attracted by the solution of the 10th Hilbert problem as presented by Martin Davis in [47] and that the recursive enumerable sets of m-tuples of natural numbers is identical with the Diophantine sets. Not all of the valuable findings in this problem have been absorbed and applied in Computational Complexity.

I founded relevant to the P vs NP problem, because of the existence of polynomials that can define a corresponding decidable language even though the complexity of the decidability may be exponential.

I did not feel initially very comfortable in working with classes like P, EXP, NP etc instead of sets, because it is well known in set theory that only few of the valid operations and definitions of sets, are also valid for classes.

The distinction between sets and classes was initially done so as to avoid the famous initial contradiction of Cantor set theory discovered by B Russell e.g. the set of sets, which after all is a class and not a set. And there are definable and non-definable classes.

(see e.g. Thomas Jech [11] page 3. A class is defined by a formula $\varphi(x, p_1, p_2, \dots, p_n)$, only for sets, where $x, p_1, \dots, p_n$ are sets, as a class all sets x that satisfy $C = \{x / \varphi(x, p_1, p_2, \dots, p_n)\}$. **We say that the class C is definable** from the sets $p_1, p_2, \dots, p_n$ if φ has one only free variable x . BUT the $x, p_1, \dots, p_n$ must be sets not classes, and the formula φ must be strictly for sets.)

Thus I used to prefer to fix an alphabet Σ , or a Cartesian powers $V(N, m) = N^m$ of the natural numbers and work with sets of languages EXP_Σ , NP_Σ , EXP_V , NP_V etc rather than classes.

From my experience in solving the Clay millennium problem in fluid dynamics about the Navier-Stokes equations (See references [7]), I knew that a probable failure in solving it is that the mainstream of research has forgotten, or disregarded as too old research, some old but significant researchers who had accumulated significant results and theorem relevant to the subject. In the case of the Navier-Stokes equations it was the invariants discovered by Helmholtz, Kelvin and Stokes. Here it seemed as if it was the work on recursive functions, by K. Goedel, S. Kleene, and the solution of the 10th Hilbert problem by many significant researchers. At their time the concept of complexity was not the main concern, but it could be traced and assessed in a posterior reading in their theorems. **The context of the natural numbers instead of formal languages of words, is older and much more promising in existing tools, theorems and conclusions.**

Among the 3 schools of equivalent definitions of computability (and consequent Church thesis) 1) Recursive functions of natural numbers, 2) Normal Markov algorithms, 3) Turing machines computability, The 1) is the oldest going back to the 19th century and with the greater number of proved theorems and tools. Therefore I preferred this school to find tools for a proof of the P vs NP millennium problem.

As Gauss used to say, there is no better University studies that to study the work of great mathematical masters. The ideas and tools to solve the P vs NP problem came

to me after studying the solution of the 10th Hilbert problem (see e.g. David Martin references [47]) but also the **metamathematical work** of K. Goedel with the K. Goedel numbering who solved the 1st of the 1900- Hilbert problems about the consistency and completeness of Peano arithmetic, after inventing a model of logical provability inside the natural numbers and on the numbers computability by recursive functions. (see e.g. references [44] Goedel Kurt “Über formal unentscheidbare sätze der Principia Mathematica und verwandter Systeme I, Monatshefte für mathematik und Physik, vol. 38,(1931) pp.173-198.) A result of this work is the **metamathematical Kleene’s normal form** of the recursive functions as described e. g. in references [51] Kleene S.C. 1936 “General Recursive functions of natural numbers “ Mathematics Annalen vol. 112 (1936) pp 727-742. Also the metamathematical study by Kleene of the Arithmetic Hierarchy of predicates and sets, as in references [52] Kleene S.C. 1943 “Recursive predicates and quantifiers “

Transactions of the American mathematical Society vol. 53 (1943) pp 41-73

I must stress here that this normal form of the recursive functions depends is essentially based on 2nd order predicates according to **Russell’s ramified logical orders** , thus it is a **3rd logical order formula** , that does not exist in 1st order countable logic, thus strictly speaking is not a recursive function (which is of logical order=1) but it says something very interesting about the recursive functions. We will discuss about it below.

As I am an interdisciplinary researcher, is a pleasure to me to search all possible research even distantly relevant to this millennium problem. I wanted to be sure, that I did not miss any results that could be used.

When in the beginning I tried to solve the P vs NP millennium problem, I could not imagine, that I had to go back 125 years to read the axiom of reducibility in Principia Mathematica of B Russell and discover that the famous arguments of incompleteness by K Goedel are refuted by the axiom of reducibility. Happily the Richard contradiction is also refuted this way (and probably this is the only way to refute it). *And of course I could not expect that I would find proofs that $P \neq NP$ is not provable at least within the 1st order **countable logic** of set theory* (See RS3, and RS4 below). But the happy end came with the meta-mathematical approach of **descriptive complexity**, which allows for a proof that $P \neq NP$. This meta-mathematical approach of descriptive complexity is equivalent to a symbolic logic of **transfinite logical order $\omega+n$** .

1) K. Goedel’s claims of incompleteness of the logical system of Principia Mathematica from the point of view of B. Russell. How can we vanquish (refute) the Richard contradiction in mathematical computation? What will happen to Goedel’s arguments if we vanquish the Richard contradiction?

- 2) Russell's disallowed impredicative predicates and consequences for the millennium problem P vs NP of mathematical computational complexity.
- 3) The way to vanquish the Richard contradiction in mathematical computation, after Russell's standards unfortunately assesses as incomplete the current proof of the time hierarchy theorem and the proof by of the non-decidability of the acceptance problem of Turing machines. Both proofs though could be repaired to fit the B Russel standards.
- 4) Valid after the Russell's standards proof that P is not equal to NP, within the meta-mathematical approach of descriptive complexity (equivalent to a transfinite logical order $\omega+n$, symbolic logic)

CHAPTER 2-3

- 1) The solution of the P vs NP millennium problem in the context of Descriptive complexity.
- 2) Russell's standards of logical reasoning against Goedel's arguments of incompleteness (Although 1st incompleteness does hold !).
- 3) The non-valid Richard antinomy style diagonal arguments of Computational Complexity according to set theory and Russell's standards of logical Reasoning.

By Kyritsis Konstantinos University of Ioannina

(This was originated with a long lecture here

[The millennium problem "Polynomial complexity versus non-deterministic polynomial complexity". What is the state of the art today? Ill posed aspects of the problem? Example of a reasonable solution. Perspectives in the theory of computational complexity](#)

https://www.researchgate.net/publication/376170410_The_millennium_problem_Polynomial_complexity_versus_non-deterministic_polynomial_complexity_What_is_the_state_of_the_art_today_Ill_posed_aspects_of_the_problem_Example_of_a_reasonable_solution_Perspec)

Abstract

In this analysis we present the method that B Russell vanquishes both the syntactical and semantical antinomies in the logical system of Principia Mathematica, through his ramified logical orders, and disallowed predicative predicates (axiom of reducibility). We apply it to the Richard antinomy diagonal argument (in the validity of propositions) which is the twin brother of the Goedel's diagonal arguments (in the provability of propositions). We discuss the consequences of this for the work of Goedel. We also present the consequences of Russell's logical standards for the diagonal method proofs of the time hierarchy theorem and the acceptance problem of the Turing machines. Also the impact of the logical standards of Russell for the solution of the P vs NP millennium problem. We present FINALLY a proof acceptable by the standards of logic by B Russell, that P is not equal to NP within the meta-mathematical approach of descriptive complexity which is equivalent to a $\omega+n$ transfinite logical order, symbolic logic. We discuss both the modification of the Church thesis so as to allow meta-machines but also restriction of the Church thesis to practical algorithms that do not involve at all the infinite in the set of inputs or run time and space (bounded run time and space).

RS1. THE DISRUPTIVE ROLE FOR THE COMPUTATIONAL COMPLEXITY OF THE SOLUTION OF THE MILLENNIUM PROBLEM P vs NP. THE RUSSEL DISALLOWED IMPREDICATIVE PREDICATES , THE RICHARD DIAGONAL ANTINOMY AND CONTRADICTIONS. MY ODYSSEY WHEN I TRIED TO RE-SOLVE ITS MYSTERY.

(In the symbolism of paragraphs RS=resolution)

- 1) K. Goedel's claims of incompleteness of the logical system of Principia Mathematica from the point of view of B. Russell. How can we vanquish (refute) the Richard contradiction in mathematical computation? What will happen to Goedel's arguments if we vanquish the Richard contradiction?
- 2) Russell's disallowed impredicative predicates and consequences for the millennium problem P vs NP of mathematical computational complexity.
- 3) The way to vanquish the Richard contradiction in mathematical computation, after Russell's standards unfortunately assesses as incomplete the current proof of the time hierarchy theorem and the proof by of the non-decidability of the acceptance problem of Turing machines. Both proofs though could be repaired to fit the B Russell standards.
- 4) Proof of the non-provability of $P \neq NP$ within countable symbolic logic of set theory. Valid after the Russell's standards proof that $P \neq NP$, within the meta-mathematical approach of descriptive complexity (equivalent to a transfinite logical order $\omega+n$, symbolic logic)

MY INITIAL APPROACH IN MY ODYSSEY

When I finished solving the Millennium problem about the Navier-Stokes equations in fluid dynamics, in 2017 (See [7]) I started trying to solve also the P vs NP millennium problem. I was afraid that it would be more difficult compared to that in fluid dynamics, because fluid dynamics is a centuries old specialization of mathematics and it has proved practically all the necessary tools about it. On the other hand Computational Complexity was not really more than 50-years old, and it was expected that it has not proved yet all its necessary tools and results that are essential for this specialization. But I never expected my Odyssey that followed and the disruptive role that the solution of the millennium problem P vs NP would have for Computational Complexity.

Initially in 2017 I thought that I proved that $NP=EXPTIME$ thus P is not equal to NP (see e.g. [4], [5] , [6] .[8])

But when I lectured the solution in National Technical University of Athens in 2023, I discovered that it had an error.

So I spent all my sabbatical of 2023-2024 in trying to re-solve it.

I was much attracted by the solution of the 10th Hilbert problem as presented by Martin Davis in [47] and that the recursive enumerable sets of m-tuples of natural numbers is identical with the Diophantine sets. Not all of the valuable findings in this problem have been absorbed and applied in Computational Complexity.

I founded relevant to the P vs NP problem, because of the existence of polynomials that can define a corresponding decidable language even though the complexity of the decidability may be exponential.

I did not feel initially very comfortable in working with classes like P, EXP, NP etc instead of sets, because it is well known in set theory that only few of the valid operations and definitions of sets, are also valid for classes.

The distinction between sets and classes was initially done so as to avoid the famous initial contradiction of Cantor set theory discovered by B Russell e.g. the set of sets, which after all is a class and not a set. And there are definable and non-definable classes.

(see e.g. Thomas Jech [11] page 3. A class is defined by a formula $\varphi(x, p_1, p_2, \dots, p_n)$, only for sets, where $x, p_1 \dots p_n$ are sets, as a class all sets x that satisfy $C = \{x / \varphi(x, p_1, p_2, \dots, p_n)\}$. **We say that the class C is definable** from the sets $p_1, p_2, \dots, p_n$ if φ has one only free variable x . BUT the $x, p_1 \dots p_n$ must be sets not classes, and the formula φ must be strictly for sets.)

Thus I used to prefer to fix an alphabet Σ , or a Cartesian powers $V(N, m) = N^m$ of the natural numbers and work with sets of languages $EXP_\Sigma, NP_\Sigma, EXP_V, NP_V$ etc rather than classes.

From my experience in solving the Clay millennium problem in fluid dynamics about the Navier-Stokes equations (See references [7]), I knew that a probable failure in solving it is that the mainstream of research has forgotten, or disregarded as too old research, some old but significant researchers who had accumulated significant results and theorem relevant to the subject. In the case of the Navier-Stokes equations it was the invariants discovered by Helmholtz, Kelvin and Stokes. Here it seemed as if it was the work on recursive functions, by K. Goedel, S. Kleene, and the solution of the 10th Hilbert problem by many significant researchers. At their time the concept of complexity was not the main concern, but it could be traced and assessed in a posterior reading in their theorems. **The context of the natural numbers instead of**

formal languages of words, is older and much more promising in existing tools, theorems and conclusions.

Among the 3 schools of equivalent definitions of computability (and consequent Church thesis) 1) Recursive functions of natural numbers, 2) Normal Markov algorithms, 3) Turing machines computability, The 1) is the oldest going back to the 19th century and with the greater number of proved theorems and tools. Therefore I preferred this school to find tools for a proof of the P vs NP millennium problem.

As Gauss used to say, there is no better University studies that to study the work of great mathematical masters. The ideas and tools to solve the P vs NP problem came to me after studying the solution of the 10th Hilbert problem (see e.g. David Martin references [47]) but also the **metamathematical work** of K.Goedel with the K. Goedel numbering who solved the 1st of the 1900- Hilbert problems about the consistency and completeness of Peano arithmetic, after inventing a model of logical provability inside the natural numbers and on the numbers computability by recursive functions. (see e.g. references [44] Goedel Kurt “Uber formal unentscheidbare satze der Principia Mathematica und verwandter Systeme I, Monatshefte fur mathematik und Physik, vol. 38,(1931) pp.173-198.) A result of this work is the **metamathematical Kleene’s normal form** of the recursive functions as described e. g. in references [51] Kleene S.C. 1936 “General Recursive functions of natural numbers “ Mathematics Annalen vol. 112 (1936) pp 727-742. Also the metamathematical study by Kleene of the Arithmetic Hierarchy of predicates and sets, as in references [52] Kleene S.C. 1943 “Recursive predicates and quantifiers “

Transactions of the American mathematical Society vol. 53 (1943) pp 41-73

I must stress here that this normal form of the recursive functions depends is essentially based on 2nd order predicates according to **Russell’s ramified logical orders** , thus it is a **3rd logical order formula** , that does not exist in 1st order countable logic, thus strictly speaking is not a recursive function (which is of logical order=1) but it says something very interesting about the recursive functions. We will discuss about it below.

As I am an interdisciplinary researcher, is a pleasure to me to search all possible research even distantly relevant to this millennium problem. I wanted to be sure, that I did not miss any results that could be used.

When in the beginning I tried to solve the P vs NP millennium problem, I could not imagine, that I had to go back 125 years to read the axiom of reducibility in Principia Mathematica of B Russel and discover that the famous arguments of incompleteness by K Goedel are refuted by the axiom of reducibility. Happily the Richard

contradiction is also refuted this way (and probably this is the only way to refute it). *And of course I could not expect that I would find proofs that $P \neq NP$ is not provable at least within the 1st order **countable logic** of set theory* (See RS3, and RS4 below). But the happy end came with the meta-mathematical approach of **descriptive complexity**, which allows for a proof that $P \neq NP$. This meta-mathematical approach of descriptive complexity is equivalent to a symbolic logic of **transfinite logical order $\omega+n$** .

RS1.1) THE RICHARD CONTRADICTION AND THE WORK OF GOEDEL AS DISALLOWED BY RUSSELL'S IMPREDICATIVE PREDICATES AND CORRECT ASSIGNMENT OF LOGICAL ORDERS.

There were many who rejected and criticized the arguments of incompleteness of Goedel, mentioning that it is essentially the arguments of the Richard antinomy. Among them **Finsler**, **Zermelo** and **Wittgenstein**. (see Wikipedia [75]). The answer of Goedel was that he was by far more detailed with 43 lemmas than the original argument of the antinomy of Richard, and applying them to provability rather than truth, thus original compared to Richards arguments. **Nevertheless, the point that seemed missed was not the originality of Goedel which indeed was so, but that refutability of the Richard antinomy would mean refutability of the Goedel's arguments as well! And refutability of Richard's arguments was certain!**

And here is a high time to point out the general problem of contradictions (antinomies) with the mathematical sciences, axiomatic systems that B. Russell in Principia Mathematica very carefully ruled-out (vanquished), which applies also to Computational Complexity.

We have a non-negotiable requirements in the mathematical sciences that the theory must be **free from contradictions**. In other words that we are in an axiomatic system which is consistent. **If not, if our axiomatic system is not consistent then we practically can prove anything by Deductive explosion**, (see [82] Wikipedia Deductive explosion)

The contradictory approaches in a subject (e.g Euclidean and non-Euclidean geometry) are accommodated with different contradictory between them axiomatic systems that each one though, is consistent (The democracy of axiomatic formal systems of D. Hilbert).

Now if a mathematical discipline is working indiscreetly and simultaneously within two different and contradictory between them axiomatic systems A, B, then it is expected that it will meet with contradictions and practically that it can prove anything.

What is a relevant example of contradictory between two them axiomatic systems A , B?

Well B may be **set theory** as formulated by the Zermelo-Frankel axiomatic system and A can be the context of **meta-mathematics or symbolic logic** which is small computer-size that is **1st order and countable (Goedel's logic)** .

In general , the axiomatic theory that we use in the meta-mathematics of an axiomatic system, must be non-contradictory with the objective-ontology axiomatic system of mathematics, because we often make arguments that jump from mathematics to meta-mathematics and vice-versa (e.g. Goedel's incompleteness arguments or discriptice computational complexity).

Is there an example of a contradiction when we use both theories simultaneously? Yes! It is the **Löwenheim–Skolem** “paradox” or contradiction (see [63] [64] below).

In the next by $|A|$ we will symbolize the cardinality of a set A.

Usually it was stated as a theorem (the **downward** part of the modern **Löwenheim–Skolem theorem**) where **any axiomatic system, that has an infinite model M , has also a countable model N .** In the proof of it (see [63]) we must accept as model N a set N which may be of less cardinality the full set of non-logical constants! (known also as signature σ) In other words that $|N| < |\sigma|$. E.g. A model N of the axiomatic system of the real numbers R , that does not contain all the real numbers $|R| = |\sigma|$ as non-logical constants.

Of course the theorem assumes that any such axiomatic system is within a countable 1st order logic.

Skolem was a person who did not believe in the existence of any infinite higher than the countable. (See e.g. [63]) and he could accept as model sets M, with cardinality $|M|$ less than the signature its self $|M| < |\sigma|$, when in σ we include all constants. That is why was severely criticized, in the “proof” of his **downward part** of the **Löwenheim–Skolem theorem**.

For example such a downward side, statement would imply that the axiomatic system of the real numbers has a model which is only countably infinite. On the other hand in most undergraduate programs of mathematics departments it is proved within informal logic (**thus 2nd or higher order and of predicates of cardinalities at least as those of the subsets and power set**) that all the models of the axiomatic system of the real numbers are isomorphic and each one of them is uncountable infinite!

These two results are obviously contradictory!

So the correct **downward side** statement of the **Löwenheim–Skolem theorem**, is derived from a more decent statement which occurs e.g. in the book by C. Papadimitriou in [18] part II, Logic Chapter 5 Corrolary 6 page 111, and it should be

In any axiomatic system, within 1st order countable logic , the part of its informal logic which is countable and 1st order, has a countable model (but in general not a model of the axiomatic system itself).

This Löwenheim–Skolem Theorem was eventually corrected in 1936 with the **upward part** by the famous A. I. Malcev ,in to that **any axiomatic system, if it has an infinite model M , with $|M| \geq |\sigma|$, then it has also a model N , for any larger cardinality $\kappa = |N|$, and $\kappa > |M| \geq |\sigma|$.** (see [63])

So the downward part, also must be corrected with the inequality **$|M| \geq |\sigma|$.**

The upward side is not contradictory with what we know about the axiomatic system of the real numbers being categorical (all models are isomorphic and have cardinality of the continuum) . **The reason is that, the cause for this inability to capture that the axiomatic system is categorical, is the inadequacy of the countable 1st order logic.** It is not adequate because

- 1) it is countable and it cannot contain all the uncountable many non-logical constants (signature σ) of the real numbers
- 2) It is 1st order, and the axiom of continuity (or axiom of supremum) requires a 2nd order proposition-axiom.

The inadequacy of the 1st order countable logic, leads also to that the Peano Arithmetic has models for any cardinality larger than the countable infinite.

(strictly speaking the axiom of induction requires a 2nd order logic).

We conclude that the informal adequate logic, of Peano arithmetic, of Real numbers, of Euclidean geometry, of set theory etc, is neither 1st order, neither countable!

(It may very well turn out that the 1st order countable logic is inadequate as well to prove the famous millennium problem that P is not equal to NP).

Of course a countable formal logic for the axiomatic system of the real numbers cannot even separate the different subsets of the real numbers, with its predicates.

Now here is the extraordinary situation with Computational Complexity that it is working in the context of both of these contradictory axiomatic system! Goedel numbers, Goedel words of Turing machines, Kleenes normal form of recursive functions , numerical hierarchies of predicates are all in the context of the axiomatic system A of symbolic logic and meta-mathematics of the small computer size that is 1st order and countable (denoted e.g. by $L_{1,\omega}$) . While definitions of Turing machines through sets and functions, and statements like that all decidable languages are only countable many compared to uncountable many in total, are in the context of the

contradictory to A axiomatic system B of set theory and in an informal logic thus possible 2^{nd} or higher order and of as large cardinality α as we want denoted by $L_{n,\alpha}$ $n > 1$, $\alpha > \omega$ and according to the standards of **Principia Mathematica of Russell-Whitehead or Hilbert-Ackermann** (See [72],[73]).

There is no doubt therefore that it is destined to meet with contradictions if we assume that everything happens in $L_{1,\omega}$.

I was very happy to think of this. because now the strangeness of the arguments in Computational Complexity that I had discovered started making sense.

Here is an example of a contradiction arising with the hypothesis that the informal logic e.g. $L_{n,\alpha}$ $n > 1$, $\alpha > \omega$ that we utilize in axiomatic set theory (e.g. Zermelo-Frankel) can also exist in $L_{1,\omega}$ the (computers mainly) 1^{st} order countable logic.

In the A axiomatic system (Goedel meta-mathematics, 1st order countable symbolic logic) there is the implicit fallacious axiom that whatever we do in mathematics it has its counterpart in this small size symbolic logic $L_{1,\omega}$ which is at most countable infinite.

Let any set S of the axiomatic system of (Zermelo-Frankel) Z-F set theory which is definable by a predicate P of the symbolic logic based on the axioms of Zermelo-Frankel Z-F. Since therefore in symbolic logic $L_{1,\omega}$ all such predicates are at most countable. Let $P_1, P_2, \dots, P_n, \dots$ an enumeration of them defining the sets $S_1, S_2, \dots, S_n, \dots$

Then we use the diagonal method

We define as set D, as follows for each S_i , we create a new different set

$S_i^* = S_i \cup \{a_i\}$ where a_i is a set not contained in S_i . How we now that it exist such an a_i ? Because otherwise S_i would be a set containing all sets, and the set of all sets does not exist. Thus we result with a set $S^* = (S_1^*, \dots, S_n^*, \dots)$

Which is different from all S_i for all $i=1, 2, \dots, n, \dots$


Since S^* is a set, there exist in full formalization in $L_{1,\omega}$ a predicate P^* which defines it, based on the formal axioms of Zermelo-Frankel axiomatic system, therefore P^* is already one of the P_i for some $i=k$. (Here is the mistake, because scanning all the predicates $P_1, P_2, \dots, P_n, \dots$ essentially it requires a **2nd order predicate** according to B. Russell and D Hilbert logic with all possible higher order predicates and all possible cardinalities see [72], [73]. Therefore the P^* is outside the $L_{1,\omega}$ and all the rest of the arguments is empty. Besides not all of the predicates of sets of set theory can exist in a countable logic of it.)

But then $S^* = S_k$ and also S^* is not equal to S_k **Contradiction!** QED.

This contradiction is actually a proof by reduction to contradiction and it proves that the informal logic, predicates, theorems etc that we use in an axiomatic theory even when symbolized, but the axiomatic theory has uncountable many objects, they cannot all of them exist in a small size logic, like a countable 1^{st} order logic $L_{1,\omega}$.

We shall call it **Anti-Skolem Logical antivirus**.

THINKING (by logic as meta-mathematics) cannot be of lower size (infinite) than (creative) **ACTING** and ontology inside mathematics. (THINKING=LOGIC, ACTING=MATHMETICS)



We can state the elements of the Anti-Skolem logical antivirus :

1. In an axiomatic system, where uncountable many or more , totalities exist (defined by predicates), the required formal logic as defined in Principia Mathematica, must contain **not less cardinality** of predicates.

In general , the axiomatic theory that we use in the meta-mathematics of an axiomatic system (external aspect) , must be non-contradictory with the objective-ontology axiomatic system of mathematics (internal aspect) , because we often make arguments that jump from mathematics to meta-mathematics and vice-versa

2. Let an axiomatic system, where uncountable many or more , totalities exist (defined by properties and thus defining predicates), for which a countable only large formal logic $L_{n, \omega}$ is considered. When we consider an arbitrary totality defined by a property, and symbolized by a predicate P , **we cannot claim** that P belongs always to the countable logic $L_{n, \omega}$ that we consider.

We can present a similar diagonal arguments with contradiction even for the axiomatic system of the natural numbers (within set theory) by considering the uncountable many subsets of the natural numbers.

The size of the utilized formal Logic for an axiomatic system depends on

1) The cardinality of the predicates etc of the logic. If not mentioned (as in Principia Mathematica in [73] or its outline by Hilbert-Ackermann in [72]) then it can be as large as we want.

2) The Russell orders of the predicates (1^{st} order, 2^{nd} order etc). If not put a maximum finite order then it is the full logic or as called by Hilbert-Ackermann the ω - predicate calculus (See [72]) Although it is called ω - predicate calculus, all logical orders in it, are less than ω and are all finite natural numbers denoted by n .

As we shall discuss later the problem $P=NP$ or $P \neq NP$ **does** also depend on the **assumed formal logic** of computational complexity. As we shall see in $L_{n,\omega}$ (Goedel's logic) both are **formally unprovable**, but in higher order or larger cardinalities logics $L_{n,\alpha}$ $n > 1$, $\alpha > \omega$ **they could be proved**.

B. Russell devoted a very large part of his life to design the mathematical Logic so that it rules out antinomies like the Russell antinomy in set theory and the semantic antinomies as well like the Richard antinomy (see [72] page 151) . The method was of course the logical orders of the predicates, and his ramified theory of order types with the axiom of reducibility (See [73]) . We shall call it the **Russell's logical antivirus**. In [72] Hilbert and Ackermann for University teaching goals present the simple (not the ramified) theory of types). His axioms of Logic are simple and few, and like the abstract axioms of group theory, they can mean finite groups, countable infinite groups or any large cardinal group. Russell did not put any axiom of the logic being finite or infinite. But as philosopher he was well aware that Thinking (=Logic) must be larger in cardinality from acting (=doings in mathematics). We first aim then we shoot. We do not shoot 5 times in different directions and aim only one in one. **Thus even for the natural numbers, its informal logic may very well involve all high cardinalities formal logic of Principia Mathematica.**

The elimination of the contradictions in mathematical theories is of the outmost importance. That is why his 3 volumes work on Logic and sets, by B. Russell-A. Whitehead in [73] Principia Mathematica, the same title that I. Newton gave to his monumental work on the infinitesimal calculus.

At his time the discovery of the Russell contradiction in the early Cantorian set theory (which allowed sets that contain , themselves as elements) did change all of the set theory. This resulted in the axiomatic system of **Zermelo-Frankel** and the equivalent of **Bernays**.

In this history of mathematics this had happened also with the infinitesimals of Newton-Leibniz , that were never really defined, and were leading to incorrect proofs of theorems and even contradictions. This lasted for almost 2 centuries till the intervention of a mathematician much less great compared to Newton, namely **K. Weierstrass** , Weierstrass with his ε - δ definitions of topological convergence, eliminated the need for infinitesimals and gave secure and correct proofs of theorems free from contradictions.

Similarly in the early history of the Cantorian set theory, it was allowed to define and use sets that contained as elements themselves. After the discovery of the Russell

contradiction, they disallowed such sets, and the Zermelo-Frankel axiomatic system was created.

That the elimination of contradictions is of the 1st priority is shown by B. Russell, in Principia Mathematica where he described his “Logical antivirus” for the logical viruses or contradictions, in the introduction (See [73] Vol I, Chapter II, The theory of Logical types, pp 37-65. And also in section B, *12 pp 161-172).

He also identifies the contradictions or Logical viruses, with the **principle of vicious circle**, which he states as follows (See [73] Vol I, Chapter II, The theory of Logical types, pp 37)

“The vicious circle arises from supposing that a collection of objects may have members which can only be defined by means of the collection as a whole”

And he continues

“Thus for example, the collection of propositions will be supposed to contain a proposition stating that “all propositions are so and so”

He continues.

“The principle which enables us to avoid illegitimate totalities may be stated as follows “Whatever involves *all* of a collection must not be one of the collection” or conversely “Of provided a certain collection had a total, it would have members only definable in terms of the total, then the said collection has no total”

The method of Russell to vanquish contradictions is based on that any predicate or sentence of the formal logic has a unique logical order which is a natural number.

We symbolize the function of the logical order of the proposition p by $Lo(p)$

For example, a predicate $P(x)$, over natural number x , will have logical order 1, $Lo(P(x))=1$. (sometimes x is assumed to have logical order 0, and $P(x)$, logical order 1). While a predicate $Q(P)$, which is over P and P is a variable over 1st order predicates, will have logical order 2, $Lo(Q(P))=2$.

This somehow is the **simple theory of types**, where the logical order is the visible one.

But in the **ramified theory of types (=logical orders)** a predicate may have hidden higher logical order than the visible.

E.g. The predicate $P(x)$ over the natural numbers, seems as if of 1st order, but it may have hidden quantifications over 1st order predicate variables that are bounded variables as e.g.

$$P(x) = \exists R Q(R) \vee S(x)$$

The predicate $Q(R)$ is 2nd order $Lo(Q(R))=2$, since that variable R is 1st order.

But finally it does not appear in $P(x)$ because the variable R is bounded by the quantifier $\exists$.

So B. Russell demands that in order to define the logical order of the $R(x)$ $Lo(R(x))$, all the hidden quantifiers must be eliminated, and so obtain the predicate $Q(R) \vee S(x)$

Which calls **matrix-predicate** or **predicative predicate**.

So the predicate $T(R,x) = Q(R) \vee S(x)$ is a matrix or predicative predicate and he symbolizes it with an exclamation mark ! to denote this property.

Thus $T!(R, x)$ means that $T!(R, x)$ is a matrix or predicative predicate.

Now this matrix or predicative predicate has logical order 2 $Lo(T!(x))=2$, thus so is the logical order of $R(x)$ $Lo((x))=2$, and not the visible one.

From this point of view the predicate

$$P(x) = \exists R Q(R) \vee S(x)$$

Is an **impredicative predicate**.

Furthermore B. Russel imposes an axiom, in the for the formal logic of any axiomatic system , the axiom of reducibility (see [73] Vol I, Chapter II, The theory of Logical types, pp 55-56. And also in section B, *12 pp 167)

Axiom of Reducibility (strong version)

For any impredicative predicate in the formal logic of an axiomatic theory it is provable within the axiomatic system that it is logically equivalent to a predicative predicate (matrix).

In symbols ($\vdash$ means it is provable) e.g

$$\vdash (\exists T): R(x) \Leftrightarrow T!(x).$$

And for two variables it would be

$$\vdash (\exists T): R(x, y) \Leftrightarrow T!(x, y).$$

Therefore an impredicative predicate which cannot be proved to be logical equivalent in the axiomatic system to a predicative predicate (matrix) is disallowed in the logic of the axiomatic system , in the definitions, or in the proofs and theorems.

Neverthhelss it seems that there is a **weak version** of the axiom of reducibility as it seems , to interpret the ramified logical orders D. Hilbert in [72] Principles of mathematical Logic , in Chapter IV paragraph 5 in page 153, where *one could allow in the logic of the axiomatic system impredicative predicates as long as they are assigned the correct (ramified) logical order.*

We state the elements of the Russell logical antivirus.

1. Every logical formula f has a unique logical order $Lo(f)=n$, n in N
2. In a predicate of order n we can substitute in his variables logical formulae only of lower logical order. This prevents **syntactical antinomies (contradictions)**.

3) (Weak version of axiom of reducibility) In the logic we assign logical orders with the ramified way (**ramified logical orders**) . Given an impredicative predicate, we eliminate all hidden quantifications (of bounded variables) and we obtain a predicative predicate or matrix. The logical order of this matrix predicate is the

logical order of the original impredicative predicate. This prevents semantical antinomies (contradictions).

3. (*Strong version of the axiom of reducibility*) In the logic of the axiomatic system, we require that the axiom of reducibility holds. Thus we accept only impredicative predicates that are proven to be logically equivalent to predicative predicates and we **disallow impredicative predicates that cannot be proved to be logical equivalent to predicative predicates** (matrix) This prevents semantical antinomies (contradictions).

We may remark that it is easy to prove that any impredicative predicate $R(x)$ over a n th logical order variable x with hidden bounded variables S of logical order m that makes it m -logical order, $m > n$, can be converted to an equivalent m -logical order (impredicative) predicate $Q(P)$ over a free variable of m -order. Thus the visible non-hidden logical order of the predicate $Q(P)$ (here the m -order variable P) is also the correct (non-hidden now) logical order. We achieve this by utilizing e.g. $n+1$ logical order predicates P_x that have as truth set only one value of the n -logical order variable x , thus substituting the n -order variable x with the $n+1$ order variables P_x and raising the visible logical order till it becomes equal with the hidden logical order.

We may notice also that the strong version of the axiom of reducibility disallows impredicative predicates, that cannot be proved logically equivalent in the axiomatic system with predicative predicates. What would this mean for set theory, where in the axiom of replacement (see e.g. [11] Thomas, Jech . (1978) Set Theory. Academic Press, Cambridge.)

with a predicate F for a function we define a new set $S = \{F(x) / x \text{ in } A\}$ from an already existing set A ? And similarly for the axiom of Separation, where given a set A and a predicate $R(x)$ with x values in the elements of A , we define a new set $B = \{x / x \text{ belongs to } A \text{ and } R(x) = \text{true}\}$? If we violate the axiom of reducibility strange things will happen. E.g. if we assume that the logical order of the elements x of A as above is n , $Lo(x) = n$ and the logical order of A is $n+1$, $Lo(A) = n+1$, then if R is an impredicative predicate with hidden $n+k$ quantification and R is not provable to a predicate over the x 's of order n (violates the axiom of Reducibility of Russell) then it will hold that $Lo(R) = n+k$, and thus $Lo(B) = n+k > n+1 = Lo(A)$. Should we define less sets after the above restriction by Russell?

Russell (but also Hilbert, Frege, Ackermann etc)) were hoping that we could have a foundation of all of the mathematics, in Logic, after reinterpreting sets as predicates. E.g. the set α belongs to the set S , would be interpreted that the predicate α , satisfies the predicate S , or $S(\alpha) = \text{true}$. But then the axioms of set theory, as in the Zermelo-Frankel set theory surprised them as in such an interpretation (**of a possible Zermelo-Frankel Logic and Zermelo-Frankel predicates**), **were introduced extra axioms in logic, not known before.** (e.g. axiom of choice, axiom of replacement, axiom of well foundation etc). In addition, the complement of a set-predicate S , existed only relative to an other set-predicate Ω , that contains S , (as in

intuitionistic logic) and not in an absolute way as in classical Boolean logic of the non-Zermelo-Frankel set-predicates,(unless we accept proper classes-predicates as well as in the equivalent axiomatic system of Bernays-von-Neumann.).

BITCOIN BLOCKCHAINS AND IMPREDICATIVITY.

From some point of view impredicativity , resembles that blockchains as in Bitcoin. The reason is that an impredicative predicates with hidden higher order quantification that are not reducible to predicative predicates, resort via the higher order quantification to non-local and visible properties of the totality. This exactly the case of with blockchains at the bitcoin files.

Two different files of 5 bitcoins, as a file, would not differ without the blockchains that describes the non-local and non-visible history of these 5 bitcoins from the moment that they were found by mining till the present moment. Another example is when a collector of banknotes has two banknotes of 100 euros from the central bank ECB of the euro. But the first has a certificate (impredicativity) signed by the president of the ECB, that this bank note was printed historically by the first machine for euros that ECB used. While the 2nd banknote of 100 euros has a similar certificate that it was printed by the latest such machine of 2020. Unless there is a detectable difference (even at the chemistry of the inks) on the two bank notes, the difference as property will remain in the realm of irreducible impredicativity, as postulated by B. Russel.

THE RICHARD LOGICAL VIRUS WITHIN THE GOEDEL ARITHMETIZATION MODEL OF ANY ORDER, COUNTABLE LOGIC OF PEANO ARITHMETIC .

Goedel in his very famous work on incompleteness in [49] he applies his method of arithmetization not only to countable 1st order logic $L_{1,\omega}$ but to all of the higher orders logic too, but always countable. Thus to the union

$U_n(L_{n,\omega})=PM_\omega$ Where U_n is the Union for all n .

That he denotes by PM (for Principia Mathematica).

Obviously what Goedel considers is not all of the logic of Principia Mathematica PM but only its smallest countable part PM_ω .

PM contains PM_α for higher cardinal numbers α .

The countable logic PM_ω considered by Goedel is the perfect logic for a computer, but nor for a human mind which will utilize higher cardinalities in the totalities of predicates.

The Richard antinomy was the main inspiration of Goedel to create his proof according to its logic as he mentions himself. (See e.g. Goedel [49] introduction 4th page or [75])

Richard antinomy is one of the **semantical antinomies** which is also a **disguised form of the Russell antinomy**, and B Russell in the logic of Principia mathematical in [73] (outlined by Hilbert-Ackermann in [72]) managed to rule out from the good standards of logic

The Richard diagonal argument is a twin brother of the Goedel diagonal argument, so we shall present them one after the other. The first (Richards) is using validity (truth of propositions) while the second (Goedel's) provability of propositions.

For the Richard's diagonal arguments see e.g. [88] , while for Goedel's diagonal argument see e.g. Gödel [49], Newman [74] , Kleene [83]

Here is an outline within the arithmetization model inside the natural numbers N of any logical order countable logic $\mathbf{PM}_\omega$ by Goedel.

Assume that we enumerate all possible countable many 1st order predicates of $\mathbf{PM}_\omega$.

$P_1(x) P_2(x) \dots, P_n(x) \dots$

Now for each of them it exist a Goedel number $gn(P_i)$ in N . The Goedel Arithmetization **gn**: $\mathbf{PM}_\omega \rightarrow N$ seem computable (we will discuss more about it), it is 1-1 , and **gn**($\mathbf{PM}_\omega$) as subset of N is decidable. Furthermore some of the decidabilities are proven to be by primitive recursive functions. The natural numbers are included in the logic $\mathbf{PM}_\omega$, as the number 0 and symbol s which means "sequent of". If we have a formula $F=s_1 s_2 \dots s_n$ of n -symbols then the Goedel number **gn**(F) of F is defined by considering the first n prime numbers and raising them correspondingly to the Goedel numbers $gn(s_1), \dots, gn(s_n)$. The same if we have a finite sequence of Formulae $F_1 F_2 \dots F_n$. In this way of course The Goedel number **gn**(F) is much greater than the Goedel number **gn**(s_i) of each constituent symbol **gn**(F)> **gn**(s_i) since not all of the **gn**(s_i) are zero. . We may call it **part-whole inequality** of the Goedel arithmetization.

This modeling is very interesting because it **models the provability of $L_{1,\omega}$ in to the computability of N** . I believe this **is the most important result after the 45 lemmas of Goedel** that we did not know in such detail earlier.

Not the incompleteness "results" that somehow were known to Russell and Hilbert, in a bit different form.

So for the sequence $P_1(x) P_2(x) \dots, P_n(x) \dots$

We get a sequence of Goedel numbers

$gn(P_1(x))=gn_1, gn(P_2(x))=gn_2, \dots, gn(P_n(x))=gn_n, \dots$

We may assume that the sequence is ordered by the well order of the Goedel numbers of the predicates.

$P_{gn_1}(x) P_{gn_2}(x) \dots, P_{gn_n}(x) \dots$

Predicates describe properties, of numbers and so they define sets of them as their truth-sets (the sets of numbers that make them true or valid).

It is interesting to ask if e.g. the Goedel number gn_i of any of them satisfies or not the predicate $P_{gni}(x)$, in other words if $P_{gni}(gn_i)=\text{true}$ or false.

As with the Russell paradox of the set of all sets that do not contain themselves as elements, let us define by a new predicate P^* , all the Goedel numbers that do not satisfy their predicate in the above sequence . That is gn as Goedel number belongs to P^* if and only if $P_{gni}(gn(P_{gni}(x)))=\text{false}$ for some P_{gni} .

Now continues Richard since this a predicate, it belongs to $\mathbf{PM}_\omega$ and is one of the sequence $P_1(x) P_2(x) \dots, P_n(x) \dots$ E.g. the $P_{i^*}(x)$ or $P_{gni^*}(x)$. Next it is easy prove that $R=(P_{gni^*}(gni^*))=\text{false}$ is equivalent to $\sim R=(P_{gni^*}(gni^*))=\text{true}$, R equivalents to its own negation $\sim R$, so ***the $\mathbf{PM}_\omega$ as a logic of any axiomatic system which includes the Peano Arithmetic is proven by Richard and with the help of Goedel arithmetization as a contradictory! Not incomplete or unable to prove formally its consistency but plainly...inconsistent.***

We may notice that the Goedel number of $P_{gni^*}(gni^*)$ is not the gn^* , as some authors claim. gn^* is the Godel number of the predicate (not proposition) $P_{gni^*}(x)$.

The situation is terrible, because it is not a contradiction say of a particular axiomatic system, but practically of any, which includes the Peano arithmetic, like the axiomatic system of the **real numbers**, the axiomatic system of **Euclidean geometry**, the differential and integral **calculus**.

And of course a contradiction existing in the theory of mathematical computation, mathematical computational complexity etc.

As it is know in a contradictory axiomatic system, the proof by reduction contradiction id not valid, and practically it can be proved any possible proposition together with its negation. A phenomenon called **deductive explosion** (see [82]). In [82] is presented also the general argument , The **Lewis argument**, with which we can prove anything. E.g. in Computational Complexity we can prove both $P=NP$, and $P!=NP$.

The Lewis argument goes as follows. Let us denote by Q the proposition $P=NP$. And R the formula of Richard.

Step	Proposition	Derivation
1	$R \text{ And } \sim R$	Premise ^[8]
2	R	Conjunction elimination
3	$\sim R$	Conjunction elimination
4	$R \text{ or } Q$	Disjunction introduction

Thus $Q \Leftrightarrow (P=NP)$ is true!

Similarly if we wanted to prove true the $P \neq NP$

In other lectures, and uploaded papers we have presented

a) A proof that in the countable logic of Peano Arithmetic it is not possible to prove that $P \neq NP$.

Nevertheless, this proof is in the standards of Goedel and stanrd computational Complexity, but not acceptable by the stanrds of B Russell's logic.

b) A proof that $P \neq NP$, which is acceptable by the standards of B Russel, but where we have utilized the reducibility axiom of Russel for the symbolic logic of graphs (in general finite structures) where the Fagin's theorem has reduced the P vs NP problem.

Here we have a small list of very important questions

1) Is Goedel aware of this Richard contradiction in his formal system?

The answer is yes, as he mentions that hiw arguments are inspired by the above arguments of Richards. He also mentions that hiw arguments are inspired by the antinomy of the liar.

2) Can Goedel refute this contradiction?

The answer is No! Not only he cannot refute this contradiction by Richard, but he even copies his arguments as he sais and we shall see below.

3) If Goedel cannot refute this Richard contradiction, is it meaningful to prove that a contradictory system such as his (PM_ω) , cannot prove its consistency formally?

Obviously not. Something worse is holding not only that the forma system cannot prove formally its consistency, but that it is already inconsistent.

4) If Goedel cannot refute this Richard contradiction, is it meaningful to prove that a contradictory system such as his (PM_ω) , is incomplete as far as provability is concerned?

The answer again is No! Actually an inconsistent system is complete as it can prove all possible propositions together with the negations, which contradicts, the

incompleteness of Goedel. But also the proof of completeness as well as incompleteness is possible in a contradictory system.

4) If Goedel was aware of all the previous answers to the questions 1)-4), could he ever think that he the only one to knows it and the rest of the mathematicians are not so clever so as to be able to realize it? (especially when he targets the solution of the 2nd Hilbert problem, which is very famous)?

Indeed I do not want to answer this question. And I could only hope that maybe Goedel was only subconsciously aware of the 1)-4) but by repressing it not consciously.

The situation is like the famous **mysterious tale** of the **naked emperor**. The Emperor (=Goedel) walks naked in front of all (=makes arguments in a contradictory context by the Richard contradiction that he cannot refute) but every body applaud him for his new emperor cloths (=solution of the 2nd Hilbert problem)except of a child that shouted that he is just naked (=He makes arguments in a contradictory by Richard logical context) .

We summarize in a qualitative way the Richard antinomy arguments properties

1) It is a diagonal argument

2) But unlike the Cantor Diagonal argument, it assumes that the entity created by the diagonal process, is not different from all listed other entities in the diagonal, but it one of them already pre-existing there

3) It leads to a contradiction.

4) It is refutable by consideration of formal logic and logical orders of predicates.

THE ARGUMENTS OF GOEDEL

In the next we shall proceed with **the arguments of Goedel** in [49], or as it is outlined by S. Kleene in [83] , which differ from these of Richard in that Goedel is interested not in the satisfiability , true or false of a predicate, but in the provability of the resulting proposition.

So we assume the same sequence of countable 1st order predicates that Richard considered

Assume that we enumerate all possible countable many 1st order predicates of **PM_ω**.

$P_1(x) P_2(x) \dots, P_n(x) \dots$

Or ordered by the Godel numbers

$P_{gn1}(x) P_{gn2}(x) \dots, P_{gnn}(x) \dots$

Contrary to Richard, Goedel is not interested if substituting the Goedel number gni in $P_{gni}(x)$, so after getting the $P_{gni}(gni)$, if it is true or false as proposition, but if it is provable or not.

The informal meta-mathematical property of provability or not in $\mathbf{PM}_\omega$ Goedel formulates also symbolically as a predicate of his own, which he symbolizes by $\text{Bew}(x)$ in his Lemma 46 in [49] from the German word Bewisbar=provable. We shall come back to discuss the $\text{Bew}(x)$ symbolism of Goedel.

So again as Richard did, Goedel because Predicates describe properties, of numbers and so they define sets of them as their truth-sets (the sets of numbers that make them true or valid) he defines as a new predicate P^{**} , (double star not to confuse it with Richards which is simple star) all the Goedel numbers gni that when substituted to their predicate $P_{\text{gni}}(x)$ in the above sequence make the resulting proposition $P_{\text{gni}}(\text{gni})$, non-provable.

That is gni as Goedel number belongs to P^{**} if and only if

$P_i(\text{gn}(P_i(x))) = \text{non-provable}$ for some i .

Now continues Goedel, (exactly as Richard continued) since this a predicate, it belongs to $\mathbf{PM}_\omega$ and is one of the sequence $P_1(x) P_2(x) \dots, P_n(x) \dots$ E.g. the $P_{i^{**}}(x)$ or $P_{\text{gni}^{**}}(x)$

We may notice that the Goedel number of $P_{\text{gni}^{**}}(\text{gni}^{**})$ is not the gni^{**} , as some authors claim. gni^{**} is the Godel number of the predicate (not proposition) $P_{\text{gni}^{**}}(x)$.

What happens now if we substitute the gni for x ? It will become a proposition! So in this way he is able to derive (as Richard did) his famous Goedel formula-proposition $G = P_{\text{gni}^{**}}(\text{gni}^{**})$

(Compare it with Richards which it was $R = P_{\text{gni}^*}(\text{gni}^*)$).

In the next Goedel will not derive a contradiction (as Richard did) but a non-provability.

He reasons as follows. Let us assume that G is provable! Then

$P_{\text{gni}^{**}}(\text{gni}^{**})$ is provable, thus it is true. But what for $G = P_{\text{gni}^{**}}(\text{gni}^{**})$ means that it is true? It means that gni^{**} is a Goedel number belonging in the truth set of the 1st order predicate $P_{\text{gni}^{**}}(x)$. And how was $P_{\text{gni}^{**}}(x)$ defined, by what property? By the property of collecting all Goedel numbers that do not make their predicates provable (when substituted in their variable) ! Therefore gni^{**} will make the $P_{\text{gni}^{**}}(x)$ non-provable if substituted for x . Therefore $P_{\text{gni}^{**}}(\text{gni}^{**})$ is non-provable, which is a contradiction compared to the hypothesis with which we started, that G is provable.

Goedel concludes that G is non-provable, and has proved the incompleteness.

(Notice already that Goedel is using a proof by contradiction, which assumes that the logical system is...consistent when...he and ...we know that due to Richard contradiction it is inconsistent. But about refutation of the Goedel arguments we will consult an expert later, in other words B Russell)

Goedel reasons then about the provability or not of the negation $\sim G$ of his proposition G , and after involving the concept of ω -consistency he concludes, that the formal system cannot prove formally its consistency.

We do not need to follow this last part of his arguments as we have in our hands the proposition G , and $\sim G$, (to consult B. Russell about it), with which he draws his conclusions.

Proof of the consistency of Peano arithmetic by G. Grentzen.

K. Goedel claimed that within the symbolic logic there cannot be any proof that Peano Arithmetic is consistent . But a few years later G. Grentzen with correspondence with D. Hilbert, proved (See [71]) that inside set theory and after using sufficient large infinite ordinal numbers the consistence of Peano Arithmetic can indeed be proved! A seemingly contradictory solution to the solution by K. Goedel.

After the above presentation of the Goedel arguments and formula it becomes more clear, that Goedel is not in a position to refute the Richard contradiction. In the contrary, Richards contradiction endangers seriously his arguments.

Another interesting question is.

6) Is there something in the standards of reasoning in mathematical Computation that would refute the Richard contradiction?

The answer I am afraid is no!. Most of the reasoning in mathematical computation does follow the method of reasoning of Goedel from Church (alternative proof of incompleteness based on the existence of semidecidable sets that are not decidable), to S. Kleene (Kleenes “normal form” after Goedel arithmetization of recursive functions) to , M. Davis (Goedel arithmetization of Turing machines, and a general form like Kleene’s “normal form”), etc. And we will understand it better later why the answer is no after we allow the Russel Logical antivirus to refute the Richard contradiction. In mathematical computation and complexity, the method with which B. Russel Logical anti-virus works is constantly violated.

How the Russel logical standards (Russell Logical antivirus) and also the Anti-Skolem Logical antivirus refute the Richard contradiction, but unfortunately also the (incompleteness etc) arguments of Goedel!

So here I believe that all mathematicians will agree (including hopefully Goedel) that **we need to refute the Richard contradiction**. Such a contradiction is obviously a **logical virus**, under the principle of **vicious circle** as stated by B. Russel, a **cancer tumor in the body of Logic**, that needs to be eliminated before it creates by **metastasis** more such cancer-tumors of logic. In fact we need to eliminate it in a way that no metastasis of similar cancer-tumor of logic will occur. And here I am not taking about the Goedel function of arithmetization gn , but about how it is used by the diagonal arguments of Richard to create a contradiction. Let us see how we can do it.

Richard as we mentioned define his predicate P^* by all the Goedel numbers that do not satisfy their predicate in the sequence

$P_{gn1}(x) P_{gn2}(x) \dots, P_{gnn}(x) \dots$

That is gn as Goedel number belongs to P^* if and only if $P_i(gn(P_i(x)))=false$ for some P_{gni} . Or in symbols

x satisfies $P^*(x) \Leftrightarrow \exists P (P(\text{gn}(P(x))) = \text{false})$

$\Leftrightarrow \exists P (\sim P(\text{gn}(P(x))))$ or equivalently

$\exists P (\sim P(\text{gn}(P)) \text{ And } x = \text{gn}(P))$

We see immediately that this predicate has hidden existential $\exists$ quantification over the 1st order variable P , therefore it is **impredicative predicate** and its matrix or corresponding predicative predicate is the

$(\sim P(\text{gn}(P(x))))$ or $(\sim P(\text{gn}(P)) \text{ And } x = \text{gn}(P))$

Which has logical order $\text{Lo}(P^*) = \max \{ \text{Lo}(\text{gn}), \text{Lo}(P) \} + 1$

The logical order of P is $\text{Lo}(P) = 1$. Thus $\text{Lo}(P^*) > 2$

What is the logical order of the Goedel arithmetization function gn ?

Goedel insists to arithmetize **ALL** the predicates and propositions of PM_ω

That although countable contains all finite logical orders! Therefore the logical order of gn is greater than any finite number, or **we may say that**

$\text{Lo}(\text{gn}) = \omega$ where ω is the first infinite (countable) ordinal number!

We see immediately that the gn is not even belonging to the logical formulae of Principia Mathematica because it has greater logical order than any finite logical order of any formula of the logic of Principia Mathematica!

So we conclude from $\text{Lo}(P^*) = \max \{ \text{Lo}(\text{gn}), \text{Lo}(P) \} + 1$

That **$\text{Lo}(P^*) = \omega + 1$**

This was the application of the **B Russel Logical antivirus**, and even in his weak form.

Since now P^* cannot be any of the predicates of

$P_{\text{gn}1}(x) P_{\text{gn}2}(x) \dots, P_{\text{gnn}}(x) \dots$

The diagonal argument of Richard cannot be performed and no contradiction arises!

Could Richard, utilize a restriction of the Goedel function gn , say φgn , on only the 1st order predicates, so that $\text{Lo}(\varphi\text{gn}) = 2$? He could, and in that case

$\text{Lo}(P^*) = \max \{ \text{Lo}(\varphi\text{gn}), \text{Lo}(P) \} + 1 = \max \{ 2, 1 \} + 1 = 3 > 1$

Again the P^* would not be among the

$P_{\text{gn}1}(x) P_{\text{gn}2}(x) \dots, P_{\text{gnn}}(x) \dots$

and thus no Goedel number gni^* can be assigned to P^* so as to complete the argument.

This is essentially the method that B. Russel utilizes with his ramified Logical orders to refute the Richard contradiction which he does in

[73] Vol I, Chapter II, The theory of Logical types, page 64, where he writes “and in Richard’s paradox,....the number.. which causes the paradox, is found to have a definition which belongs to a higher logical type (=logical order), and thus not to come within the scope of ...the previous definition...”

And what about the Anti-Skolem logical antivirus (see above in the current paragraph)? What has it to say about P^* .

Again the Anti-Skolem logical antivirus, will not allow the assignment of a Goedel number gni^* to P^* , because, the Goedel arithmetization is **only on the countable part of the formal logic of the Peano Arithmetic**. There are uncountably many

different subsets of the natural numbers with their logically non-equivalent predicates, and thus most of them will not belong in PM_{ω} . So there is no guarantee, that P^* will belong in PM_{ω} . And we cannot adjust the definition of the arithmetized, countable part of PM , to include P^* , because the existence of P^* depends on the already arithmetized countable 1st order predicates $P_{gn1}(x)$ $P_{gn2}(x)$..., $P_{gnn}(x)$...

Now unfortunately the same objections and refutations can be made for the Goedel predicate P^{} , that would lead to the Goedel formula G .**

The P^{**} is defined a bit differently based on the provability and not satisfiability. Here we shall utilize a 2-variables predicate to express the provability as it is done by S Kleene in [83] Kleene S. "The work of Goedel" The Journal of Symbolic Logic, Dec 1976, Vol 41, No 4 (Dev 1976) pp 761-778

And in particular in page 765, where he symbolizes it by $A(u,y)$ which means that. y is the Goedel number of a proof of the proposition $P(u)$. derived by the 1st order predicate $P(x)$, which has Goedel number u , $gn(P(x))=u$, when we substitute u for x and obtain the proposition $P(u)$.

This predicate $A(u,y)$ already has hidden quantifications of higher order (impredicative predicate) that we shall reveal here.

(It corresponds to the Goedel predicate $y \vdash u$ in Lemma 45 in [49] meaning y is a proof of the formula u).

Now with the above symbolism in hand we may define formally and not only informally the predicate P^{**} .

$P^{**}(x)$ has a truth set of x defined by $\Leftrightarrow$

$\exists P (x = gn(P) \text{ And } (P(x) = \text{non-provable}))$

(see the similarity with Richards case except here we have "...=provable" instead of "...=true") Or if we utilize the A predicate

$\exists P \forall y (\sim A(x, y) \text{ And } x = gn(P))$

(or equivalently in more compact symbolism

$\exists P \forall y (\sim A(gn(P(x)), y))$)

But if we want to avoid the predicate A ,

$\exists P (x = gn(P) \text{ And } (P(x) = \text{non-provable}))$

Again as with Richard, this is a **impredicative predicate**, with **matrix** derived after we eliminate all quantifiers and make the bounded variables free, that is the

$(\sim A(gn(P(x)), y)) \text{ or } (\sim A(gn(P), y) \text{ And } x = gn(P))$

And thus the logical order $Lo(P^{**})$ of P^{**} is that of its matrix, or

$Lo(P^{**}) = \max \{Lo(gn), Lo(A), Lo(P)\} + 1 = \max \{\omega, 1, 1\} + 1 = \omega + 1$

Where as before $Lo(gn) = \omega$.

Could Goedel, utilize a restriction of the Goedel function gn , say ϕn , on only the 1st order predicates, so that $Lo(\phi n)=2$? He could, and in that case $Lo(P^{**})=\max \{Lo(\phi n), Lo(A), Lo(P)\}+1=\max\{2,1\}+1=3>1$

Again the P^{**} would not be among the

$P_{gn1}(x) P_{gn2}(x) \dots, P_{gnn}(x) \dots$ That have logical order 1.

and thus no Goedel number gni^{**} can be assigned to P^{**} so as to complete the argument.

Of course even if in the sequence

$P_{gn1}(x) P_{gn2}(x) \dots, P_{gnn}(x) \dots$

Goedel would include any possible finite logical order, then again P^{**} would have logical order $\omega+1$, thus outside the PM_{ω}

This was the application of the **B Russel Logical antivirus**, and even in his weak form.

Since now P^{**} cannot be any of the predicates of

$P_{gn1}(x) P_{gn2}(x) \dots, P_{gnn}(x) \dots$

no Goedel number gni^{**} can be assigned to P^{**} so as to complete the diagonal argument.

The diagonal argument of Goedel cannot be performed and no implications about incompleteness etc is possible!

And what about the Anti-Skolem logical antivirus (see above in the current paragraph)? What has it to say about P^{**} .

Again the Anti-Skolem logical antivirus, will not allow the assignment of a Goedel number gni^{**} to P^{**} , because, the Goedel arithmetization **is only on the countable part of the formal logic of the Peano Arithmetic**. There are uncountable many different subsets of the natural numbers with their logically non-equivalent predicates, and thus most of them will not belong in PM_{ω} . So there is no guarantee, that P^{**} will belong in PM_{ω} . And we cannot adjust the definition of the arithmetized, countable part of PM , to include P^{**} , because the existence of P^{**} depends on the already arithmetized countable predicates $P_{gn1}(x) P_{gn2}(x) \dots, P_{gnn}(x) \dots$

Goedel meticulously avoids to symbolize the arithmetization function gn , by any symbol whatever, in spite the heavy use of symbols for everything. Even for the act of substituting a variable. He knows that from the moment it symbolizes it, questions will arise about where such a function will belong. Of course nowadays after S Kleene, M. Davis, A. Church etc we do symbolize the arithmetization function. By symbol gn here is due to M. Davis in [62] in his arithmetization of the Turing machines, to their Goedel numbers. **Similarly since he did not symbolize the arithmetization function gn , he did not give the proofs of his 45 lemmas, that would heavily depend symbolically on gn .**

The diagram below, shows, the logic, the Peano Arithmetic, the arithmetization by gn which is where Goedel stands, outside of both theories. So gn is outside both Peano Arithmetic and also countable part of the formal Logic of Principia Mathematica, and so is his predicate P^{**} .

Outside of both, and also his final propositions if it could be ever defined would be outside PM_{ω} , thus, even if it was not provable in PM_{ω} it is irrelevant as it is an outsider proposition and this would not imply incompleteness.

Maybe there was another way out to save the arguments of Goedel? E.g. what if instead of Logic PM_{ω} we treat them as pure symbols without meaning, and the symbols defined in set theory? So both PM_{ω} and Peano arithmetic be absorbed entirely into set theory? Then the place that Goedel would stand would be the set theory. (Actually Goedel wants to avoid this as he intends to apply his conclusions of incompleteness for the axiomatic system of set theory too. But let us assume for the moment that he would agree in to this).

Then gn would be definable as an ordinary set function of course outside Peano arithmetic and also outside PM_{ω} . But then P^{**} as it depends on gn would be an entity of set theory but not of Peano arithmetic or countable formal Logic PM_{ω} and again his arguments would not be valid.


So we summarize the conclusions

1) Goedel is not in a position to refute the Richard contradiction defined by his Arithmetization. So this contradiction exists in his formal systems

2) Goedel considers as logic of Principia Mathematica, symbolized by him by PM , as only the countable part of it PM_{ω} . Thus strictly speaking he has no right to claim his supposed conclusions for the whole of Principia Mathematica PM .

3) The ramified system of logical orders (impredicative predicates) of the propositions by B. Russel (Russell's weak logical antivirus), prohibits of assigning a Goedel number to Goedel's P^{} predicate, therefore his famous non-provable formula G , actually does not exist in the logic of Principia Mathematica. Similarly it prohibits Richard from assigning a Goedel number to his predicate P^* , thus the final contradiction R , does not exist in Principia Mathematica.**

4) Similarly the Anti-Skolem logical antivirus, prohibits Goedel of assigning a Goedel number to Goedels P^{} predicate, therefore his famous non-provable formula G , actually does not exist in the logic of Principia Mathematica. . Similarly it prohibits Richard from assigning a Goedel number to his predicate P^* , thus the final contradiction R , does not exist in Principia Mathematica.**



More or less facts of consistency and incompleteness were somehow known both to Russell and Hilbert. As far as consistency is concerned **Russell. Hilbert and Ackermann provided proofs of the consistency of the logics $L_{n,\alpha}$ $n \geq 1$, $\alpha \geq \omega$** with finite models. And the modeling would require predicates higher than n , thus outside the logic which prove its consistency (See [72] pages 38,87, 158).

I must remark to the reader, that **I am not against conclusions of incompleteness in logic or axiomatic systems.** E.g. in set theory we know that it is incomplete, as the axiom of choice cannot be proved and also the simple continuum hypothesis is not provable after Cohen's forcing method. **What I am against is the method that Goedel chose to prove such claims.**

There is a moral value of cleaning up mathematical theories like computation and complexity from Russell disallowed impredicative predicates (=not possible to be proved equivalent topredicative predicates) and possible contradictions (cancer tumors of logic) and avoiding the chaos of deductive explosion.

My personal attitude is of course the standards of reasoning according to Russell-Whitehead (and Hilbert-Ackermann) with higher order logical orders (not only 1st order) and with cardinalities of the propositions much higher than countable compared to the computer logic of Goedel. Therefore I do not accept actually the

reasoning of Goedel and its proof, And his conclusion more or less I can draw with other methods and easier by respecting the logical orders of the predicates, and not forgetting the cardinality of the logic compared to other logics and compared to the ontology of the axiomatic system that we reason.

RS2.1) THE UNDECIDABILITY OF THE ACCEPTANCE PROBLEM OF TURING MACHINE AND ITS DIAGONAL RICHARD-LIKE PROOF AS NON-VALID ACCORDING TO THE STANDARDS OF SET THEORY (AND RUSSEL'S STANDARDS OF LOGICAL REASONING).

It was seemingly only Zermelo who noticed that the famous Richard-antinomy-like argument by Goedel, was no-valid, flawed and refutable, exactly as the Richard-antinomy argument. And this objection never became public. Instead Goedel became very famous and many other computer scientists (famous or not) started mimicking him.

In this way the discipline of Computational Complexity, became full of instances of diagonal (pseudo-)definitions in the style of Richard-antinomy, that were used as proof of various theorems.

We concentrate here to the early proof the acceptance problem (or non-halting problem) of Turing machines is undecidable. In this (pseudo-) proof it is used a Richard-antinomy like definition and argument which leads as expected to a contradiction, but then instead of blaming the diagonal definition we blame the hypothesis of decidability of the acceptance.

What we will do here is that we will re-procedure the above Ricjard-antinomy style argument on Turing machines but on arbitrary functions of set theory, so as to derive the contradiction which will be irrelevant to Turing machines and decidability

We recall from the previous paragraphs the qualitative properties of the Richard antinomy.

We summarize in a qualitative way the Richard antinomy arguments properties

- 1) It is a diagonal argument
- 2) But unlike the Cantor Diagonal argument, it assumes that the entity created by the diagonal process, is not different from all listed other entities in the diagonal, but it is one of them, already pre-existing there
- 3) It leads to a contradiction.
- 4) It is refutable by consideration of formal logic and logical orders of predicates.

We describe the Richard antinomy in the context of pure functions (not recursive function or Turing machines) in set theory.

1) We assume that we are given a countable set of functions M_i , $i=1,2,\dots,n,\dots$ over 0-1 words w of Σ^* , where Σ is the binary alphabet $\Sigma=\{0,1\}$ and the values of $M_i(w)$ are either 0, 1 or 2. (such functions could be of course Turing machines, and 1 would mean accept, 0, reject, and 2, is looping or non-halting).

2) We also assume a coding function, denote by $\text{cod}(M_i)$ that sends the functions $\text{cod}: M_i \rightarrow \text{cod}(M_i)$, to words of Σ^* , and it is 1-1 (E.g. if M_i are Turing machines, the $\text{cod}(M_i)$, denoted also by $\langle M_i \rangle$, **could** be the code of the Turing machine as a word of Σ^* . But no need to assume that M_i , is a Turing machine). **We may denote also the function cod as gn** (from the words Goedel number)

3) Then we define a **decider function** called H , on pairs of words, by

$H((\text{cod}(M_i), w)) = M_i(w)$ if $M_i(w) = 0$ or 1, and $H((\text{cod}(M_i), w)) = 0$, if $M_i(w) = 2$.

There is no doubt that according to the axioms of set theory such a decider function is definable and it does exist in set theory.

4) Then we define a **diagonal negation function** denoted by D , by

$D((\text{cod}(M_i))) = 0$ if $H((\text{cod}(M_i), \text{cod}(M_i))) = 1$ and (Eq 00)

$D((\text{cod}(M_i))) = 1$ if $H((\text{cod}(M_i), \text{cod}(M_i))) = 0$ and (Eq 01)

(that is by **negating H at the diagonal**).

Again it is obvious that such a function D does exist, in set theory and it is definable on some words of Σ^* , in particular to all words that are codes of the functions M_i

Next we may speculate, if it would be possible at all that the function D , was already one of the functions M_i !!!

One argument against it is that, when defining D diagonally over the $\text{cod}(M_i)$ if there is i^* , such that $D = M_{i^*}$, then at the definition of D , at $\text{cod}(M_{i^*}) = \text{cod}(D)$, we would have

$D((\text{cod}(D))) = 0$ if $H((\text{cod}(D), \text{cod}(D))) = 1$

BUT, HOW IS IT POSSIBLE TO DEFINE D OVER $\text{cod}(D)$, THAT IS ASSUMING THAT D ALREADY EXISTS , WHEN THE DEFINITION OF D IS ALREADY IN THE PROCESS, AND NOT FULLY COMPLETE?

We may try to go around this difficulty, by formulating the situation a bit differently: All right, I will not define D as, we did, I will not claim that it does exist in set theory , but I will speculate , IF OR NOT A D EXISTS SATISFYING

a) THE diagonal; equations (Eq 00), (Eq, 01),

b) There is an i^* such that $D=M_{i^*}$.

Now the answer to such a speculation is that it is impossible.

One reason is that from the equations (Eq 00) (Eq, 01) we get the next contradictions that $D((\text{cod}(D)))=0$ if $H((\text{cod}(D), \text{cod}(D)))=D(\text{cod}(D))=1$

and

$D((\text{cod}(D)))=1$ if $H((\text{cod}(D), \text{cod}(D)))=D(\text{cod}(D))=0$

In other words, that , if D is definable diagonally, then it cannot exist already as one of the M_i or that if D already exists as one of the M_i , then it cannot be definable diagonally as above.

A second reason, that it is impossible, comes from Logic, and the logical order of the predicates that defines the corresponding sets or functions and it is the standard way that such Richard-antinomy style arguments are refuted.

We notice that the Logical order of the words is zero $Lo(w)=0$, The logical order of the functions of words M_i , is 1, $Lo(M_i)=1$. The logical order of D if defined diagonally over the M_i is necessarily larger than that of cod and M_i , thus $Lo(D)=3$. Now this prohibits D to be one of the M_i , that would mean that it has logical order 1.

Conclusions

1) We cannot accept in set theory definitions of functions over words with a diagonal argument over a countable list of other such functions, and at the same time, assume that the function defined so pre-exist already in the list. The obvious objection is that in order to define such a function we must use itself to define itself.

In most of the books of computational complexity it is stated the halting problem and the acceptance problem of Turing machines and it is proved the undesirability of them , with a diagonal method over Goedel words $\langle M \rangle$ of Turing machines.

See for example C. Papadimitriou [18] Chapter 3, paragraph 3.2 page 59 and by M Sipser [39] Chapter 4 Theorem 4.11 page 202.

We denote by M a Turing machine and by $\langle M \rangle$ its Goedel word (coded conversion of the Turing machine M by a Goedel coding function gn to a Goedel word $gn(M)=\langle M \rangle$. The symbolism gn was used by Martin Davis in his book [62] Computability and Unsolvability in chapter 6 page 56)

The Turing machine acceptance problem is defined as

$A_{TM} = \{(\langle M \rangle, x) / \text{The Turing machine } M \text{ accepts the word } x\}$

Let us denote by H a Turing machine that would recognize or decide it.
The argument in the books, proceeds as the one that we described above
With simple functions irrelevant of Turing machines.

But the books, use the contradiction not to refute the possibility of defining the D as above, while D is already a Turing machine among the M_i , but that there cannot be a decider Turing Machine H !!!

Obviously this is “Cheating” !

Because as we saw that the contradiction already arises, even if M_i , H , D are not Turing machines! The contradiction is irrelevant to the concepts and ontology of Computational Complexity! It is a disguised form of the Richard Antinomy and is relevant to Logic and set theory.

Now the cheating is that once this Ricard-antinomy contradiction is introduced in the web of arguments of Computational Complexity (which is also in the context of set theory) it becomes an inconsistent logical system and deductive explosion applies! That is WE CAN PROVE ANY PROPOSITION P , by reduction to contradiction, blaming for the Richard-antinomy contradiction the proposition P .

WE CONCLUDE THEREFORE THAT THE STANDARD ARGUMENT IN THE BOOKS IN THE PROOF OF THE UNDECIDABILITY OF THE ACCEPTANCE PROBLEM OR HALTING PROBLEM OF TURING MACHINES IS INVALID, AND THAT IN REALITY IT HAS NOT BEING PROVED BY THE STANDARD PSEUDO-PROOF IN THE BOOKS!

If it can be proved in another valid way, or if it can be proved at all, is another matter.

Computational complexity utilizes elsewhere also such Richard-antinomy style diagonal definitions, e.g. in the proofs of **the time and space hierarchy theorems** and also in the **Rice Theorem**.

Similar objections can be raised for these arguments and definitions too.

An other important aspect is that we mentioned in the previous paragraph RS1), that sets of Turing machines **as maximal countable sets are defined only in an artificial way after some restrictions** (e.g. all finite states from the same countable set and all finite alphabets from the same countable set). We should prove in addition that D , has respected these restrictions. And if it was also of logical order 2, then claim that it is one among the M of the $\langle M \rangle$.

Could we change the definition of A_{TM} and include Turing machines of any high logical order? Unfortunately not! Because the decidability of the language of all Goedel words $\langle M \rangle$ of Turing machines as 1st order predicates, depends on the normal way of definition where the structure of the code is standard, and this the (hyper-language) of their Godel words decidable.

This means that the rest of the argument in the proof of the non-decidability of the acceptance (or halting) problem of a Turing machine cannot be performed, because the $\langle D \rangle$ is not in the Goedel words $\langle M \rangle$ of 1st order machines that define the 4th logical order machine D . It is exactly the same as with the way that we vanquished the Richard contradiction in mathematical computation, and we cannot do it in any other way (besides the Skolem logical antivirus too).

The consequences of this are significant. It is not only that the **Rice theorem** and some other theorems cannot be proved. But also **A. Church's proof** of the non-decidability of arithmetic N in 1st order countable logic $L_{1, \omega}$ (see e.g. M Sipser [39] theorem 6.13 page 257) which is an alternative proof to that of Goedel's (for 1st order countable logic though).

In this history of mathematics this had happened also with the infinitesimals of Newton-Leibniz, that were never really defined, and were leading to incorrect proofs of theorems and even contradictions. This lasted for almost 2 centuries till the intervention of a mathematician much less great compared to Newton, namely **K. Weierstrass**, Weierstrass with his ε - δ definitions of topological convergence, eliminated the need for infinitesimals and gave secure and correct proofs of theorems free from contradictions.

Similarly in the early history of the Cantorian set theory, it was allowed to define and use sets that contained as elements themselves. After the discovery of the Russell contradiction, they disallowed such sets, and the Zermelo-Frankel axiomatic system was created.

That the elimination of contradictions is of the 1st priority is shown by B. Russell, in Principia Mathematica where he described his “Logical antivirus” for the logical viruses or contradictions, in the introduction (See [73] Vol I, Chapter II, The theory of Logical types, pp 37-65. And also in section B, *12 pp 161-172).

He also identifies the contradictions or Logical viruses, with the **principle of vicious circle**, which he states as follows (See [73] Vol I, Chapter II, The theory of Logical types, pp 37)

RS2.3) THE META-CHURCH THESIS AND META-MACHINES.

Many computer scientists think of Computational complexity as a ...physical science that of the computer machine! No need for axioms. But it is not so! Computational Complexity introduces at least the countable **infinite** (languages of infinite many words) and nothing in material physical reality of a computer machine is infinite. **Therefore, it is crucial to ask and be aware of what are the axioms of computational Complexity or what are the axioms that we use in an argument or proof in Computational Complexity! Is it set theory axiomatic system? is it Peano arithmetic axiomatic system? And in how large formal logic? Countable? Uncountable? 1st order? Higher order?**

DEFINITION C1 . Conventional Goedel-like reasoning with Russell disallowed impredicative predicates (with the wrong logical ramified order) in mathematical computation.

The **Goedel-like conventional reasoning** in computational complexity. is defined as the reasoning where we accept the Goedel-words (or numbers) as ordinary word-inputs of 0-logical order and where by a Universal Turing machine U , we read all (or infinite many different) Goedel words $gn(M_i)=\langle M_i \rangle$ of Turing machines and define accordingly algorithms that are as if calling infinite different subroutines but still we consider it as a single Turing machine definable with a 2nd logical order predicate. **The above attitude violates the logical standards of ramified logical orders for impredicative predicates of B. Russell.**

Common sense suggests that we are cheating here when we assume that in all situations and possible definitions, when we feed a Universal Turing Machine U though a countable infinite set of all possible Gödel words $\langle M \rangle$, of all possible Turing Machines and with the word $(\langle M \rangle, x)$ where x runs over all possible words x and U simulates ALL the Turing machines (including itself...?) that this **is always equivalent** with a single Turing Machine computing on x with a finite only code. As we shall see later such practices define also meta-machines (or self-programmable machines) that are uncountable many (as the irrational numbers) compared to usual Turing machines that are countable many (as the rational numbers)

How is this situation resolved? We shall see that what a Universal Turing machine does after reading the Goedel-words $\langle M \rangle$ of infinite many other Turing machines (like querying an infinite number of sub-routines) is not a Turing machine but a META-MACHINE. **There are uncountable many such recognizers** Meta-machines, and they can compute and recognize any subset of the full vocabulary Σ^* (Or natural numbers $\mathbb{N}$). meta-machines are also called **self-programmable Turing machines** in the full text of the current lecture.

DEFINITION C2 (META-MACHINES) . When restoring the logical standards of B Russell of ramified logical orders, input languages and machines now appear with higher logical orders **that we call in general Hyper-machines or meta-machines.** The meta-machines are not just Turing machines with higher logical

order. But the converse holds . Turing machines with higher logical order are included in the meta-machines. In this way we are allowed to create new ontology the **meta-machines** (like the irrational numbers) which can call **an infinite number of subroutines** and have variable and unbounded size during computation, so they are uncountable many and can compute any subset of the natural numbers. Of course meta-machines include also as special case the standard Turing machines (analogue of the rational numbers). They include also the non-deterministic Turing machines. Meta-machines is a new theoretical definition but in practice they correspond to coding of programs where we decrease the time run by increasing the storage space during computation when keeping records of the computed (memory of computed) that saves searching at later steps.

But if the reader gets the impression that I just like high cardinals and ordinals and want to introduce them in Computational Complexity, then he has the wrong impression. Actually the last work of my carrier is the re-writing of the Newton-Leibniz calculus (of course in a logically non equivalent way) where the infinite is absolutely exiled and it does not exist. It is the perfect and faster differential and integral calculus for engineers and applied scientists and I call it Democritus or digital differential and integral calculus. (Democritus was the ancient Greek philosopher of the atomic structure of matter.

The infinite is behind all our logical difficulties with the paradoxes or antinomies , but in this particular planet that we live it seems that the concept of the infinite was indispensable in creating mathematics and sciences. Only the infinite was giving to the thinkers winds to free their thinking from negative energies of the matter.

And here is an idea to make Computational Complexity more efficient and realistic by abolishing the countable infinite from the languages from which we give input words or natural numbers to algorithms.

Why should we consider infinite languages at all as inputs to Turing machines or recursive functions? It is never in the real life the case! We only give as input finite many data (words). If we include in the definition of a Turing machine this restriction then all Computational Complexity will become better and more realistic. Here is what will happen.

1) Since it is known that any finite language is decidable, (e.g. by adding its words in the code of the Turing machine) we define the decidability of a finite language **only relative** to the next 3 types of complexity

a) Time complexity (runtime of the software program)

b) Space complexity (size of the RAM memory of the computer)

c) Source-code space complexity (size of the source code of the software program)

The latter type of complexity (source-code size of the Turing machine) is a new but important type of complexity.

Then a finite language L may or may be not decidable by any Turing machine M , if there are strict or loose such 3-types of complexity limits .

Now the time complexity is not a function (exponential or polynomial) but simple a natural number time units.

And the famous P vs NP problem loses its meaning since now that all languages are finite it will hold that $P=NP=EXPTIME$. (Collapsing optimistic case).

THE META-CHURCH THESIS.

All the above lead to the **META-CHURCH THESIS**.

The meta church thesis claims

*MC1) that the algorithms as defined by the Church thesis are not realistic as far as the hypothesis, that an infinite number of input data (infinite languages) are given as inputs to the algorithm (recursive function, Turing machine, or normal Markov algorithm) , The reality is that only finite many input data in total can be considered as given to any algorithm. There are not infinite sets for the computer. So we define the concept of **practical algorithm (or machine)** as **differing** from that of Turing machine by allowing **only finite many input data**. Furthermore , the **running time** and the **space size** (in the Turing machine tape) **must always be finite**, by a timer and space restriction counter. **Practical algorithms** are essentially a **restriction of the symbolic logic** of classical computability to a realm of natural numbers where only finite many constants, terms and variables exist.*


MC2) We may accept in the definition of general recursive functions uncountable many different predicates of finite system of equations, which will make the general recursive functions uncountable. This is done by not requiring that the functional predicates are defined by the system of equations but by allowing one more of them be predefined as one among the uncountable many possible functions of the natural


numbers. Thus there will not necessarily exist a unique maximum countable set of such general recursive functions (The maximum principle of subsets ordered by inclusion applies only to totally ordered by inclusion sets of sets and not without the **axiom of choice**). Similarly for the Turing machines when allowing uncountable many different finite alphabets or finite states of them. There will not necessarily exist a unique maximal countable set of such Turing machines (The maximum principle of subsets ordered by inclusion applies only to totally ordered by inclusion sets of sets and not without the **axiom of choice**).

MC3) Furthermore we may enhance the concept of recursive functions and deterministic Turing machines in a different way by allowing unbounded large and changing during computation code of them (as a non-deterministic machine does). We may call them Hyper-machine or meta-machines. Their behavior suggests in practice to strategies to reduce the running time by increasing the available space size and keeping memory and records in the algorithm, of the already computed data or decisions.

From this point of view the meta-machines, may call an infinite number of different sub-routines, is as good abstract and infinite theoretical ontology as the standard Turing machine is for the **practical algorithms** in 1) situation where there is nothing infinite. And as proof of it is that exactly as when only finite languages are considered we get $P=NP=EXP$, so the same we get when we accept meta-machines for infinite languages.

The next scheme shows the case of modified definition of general recursive functions, that allows them to be uncountable many.





Turing machines are usually defined as countable many by restricting the finite alphabets to have letters from the same countable set A , and finite states from the same countable set Q .

But in arguments involving such countable totalities of Turing machines, sometimes the need of more and different alphabets or states is required. Thus different countable totalities of Turing machines from the initial countable totality are created. From the maximum principle derived from the axiom of choice of set theory, for any set A of Turing machines totally ordered by strict inclusion inside a larger set C of Turing machines, there is a unique set B of Turing machines totally ordered in the strict inclusion which includes A and is maximal such, in C . Thus different maximal sets of Turing machines can be created.

Here I discuss that there is a way of evolution in the world of software and solutions of problems, that can turn in a systematic way exponential time problems to polynomial time problems without somehow violating the church thesis, and without resorting to randomness and non-deterministic Turing machines. And at the same time this other method reflects what has already happened in the evolution of the complexity of software and problem solving.

A characteristic example is the software that drives automatically cars. When humans were writing the many hundreds of thousands lines code, for such a software to drive a car, it was failing dramatically. But after utilizing methods like machine deep-learning, and allowing for the software to write its own code, then many millions of code lines were written, and eventually driving the car became possible and sufficient safe.

The meta-machines or self-programmable machines are inspired by the deep learning, where while we had failed to program the computer with a few 100 thousand lines to drive a car, when we asked the computer to program itself it produced some millions of code lines and finally it managed to drive a car.

Meta-machines can define an uncountable number of recognizers ! Not countable!

And can compute ANY set whatever of the natural numbers. These would appear as contradictions if they would be considered as simple Turing machines. They require infinite information over the Goedel words of all other Turing machines as the Universal Turing machines reads them (corresponding to calling infinite different subroutines) . They are like the irrational numbers with infinite many digits compared to standard Turing machines that are like the rational numbers with only finite many different digits.

Another aspect of meta-machines compared to simple Turing machines is that, since they have variable during computation programming code, they admit a 2nd tape compared to Turing machines where they can keep record of the compute or decided so far. This greatly decreased computation time for later steps at the cost of course of computations pace. It corresponds to real life practice where, during the computation the software program keeps records in a small data base, of the so far computed or decided and so decreasing run time at the cost of increasing ram memory or hard disc storage memory.

And this is the suggested concept of **deterministic self-programmable Turing machine**.

DEFINITION 0

The next sequence of statements specify axiomatically, without too tedious details what is a **deterministic self-programmable Turing machine**.

1. A deterministic self-programmable Turing machine is like an ordinary deterministic Turing machine as far as inputs finite alphabets, tape , head , left-right moves , accept-state , reject -state is concerned.
2. It is deterministic in the sense, that like an ordinary deterministic Turing machine , at each time with input a word it produces a finite sequence C_i of configurations, such that C_i yields C_{i+1} according to the rules of the deterministic non-programmable Turing machine. This sequence either will halt at an acceptance state or a rejection state or loop for ever.
3. There are though some very important differences compared to an ordinary deterministic Turing machine. The self-programmable deterministic Turing machine, **can increase its code (finite states and transition functions) while it is calculating on the tape**. In order to do so it has the next abilities.
4. It has an **infinite long tape alphabet**. Also **infinite many states , and transition functions**, divided each time to the **written (or active) ones that are always finite** and the unwritten yet (or non-active) that are infinite. **The transition functions are always defined as function on finite many states**, only from all the infinite states it has.

5. It has the ability to search along his infinite non-written yet transition functions on a **2-dimension tape called code-editor**, which is divided like a 2-dimensional tape and after finding what it needs to increase the written code. While doing so the head is not moving in the 1-dimensional tape **and the complexity of code increase is not counted in the run-time complexity on the 1-dimensional of the self-programmable Turing machine.**
6. **The increase of its code is triggered by what it reads in its 1-dimensional tape, and its already written finite states and finite transition functions code, in the code-editor and after search in the infinite non-written code.**
7. A self-programmable deterministic Turing machine can take infinite many input data before running, although this ability can be covered and as secondary effect produced by its infinite non-active code in the 2-dimensiona code-editor.
8. **Ordinary deterministic Turing machines are special cases of self-programming deterministic Turing machines.**
9. It holds that if a deterministic self-programmable Turing machine decides a finite language L , then there is an ordinary deterministic Turing machine that decides it also. And conversely. **Therefore on finite languages both types of machines have the same power and are equivalent.** Thus the church thesis is not violated , since it refers to what in reality humans and machines can to.
10. *Nevertheless the two types of machine are not equivalent on infinite languages , which after all do not exist in real life, and thus this difference is important only to the fictional logical game of theoretical computational complexity.*
11. For a self-programmable deterministic Turing machine (besides the concept of **run-time complexity** and **space complexity**) there is also a 3rd type of complexity the concept of **code-complexity** which is measured in the usual way by the increase of the length of the written (active) code while running as a single string, compared to the increase of the length of the input word.
12. **It can be proved that if L is an infinite language over an alphabet Σ , decided by an ordinary non-deterministic Turing machine M , with complexity some function $T(|x|)$ (time constructible function) , then there is a self-programmable deterministic Turing machine, SPM which can decide it in complexity $T(|x|)$. Nevertheless we do not put a bound to the code-increase complexity of SPM. Thus there is a self-programmable deterministic Turing machine which can decide any language of NP in polynomial time. That is $P_{SPM}=NP_{SPM}$.**
13. **The even stronger result holds , that any set is decidable by a meta-machine in polynomial time. Thus $P_{SPM}=NP_{SPM}=EXP_{SPM}$. This results corresponds to the same effect of accepting only finite languages as inputs of algorithms , which is the real life situation, in which case of course it also holds that $P=NP=EXP$.**

RS4) PROOF IN THE METAMATHEMATICAL APPROACH OF DESCRIPTIVE COMPLEXITY (IN TRANSFINITE LOGICAL ORDER $\omega+n$ LOGIC) THAT P is not equal to NP.

With the formulation of the problem P vs NP, any one who tries to solve it would like to have a clear perception that it is well defined as far as proper classes and sets is concerned.

For example would an equation $P=NP$ or $P \neq NP$ would change when we formulate the same problem with sets rather than proper classes?

0) Can we define the P vs NP in a most general and flexible way but with sets instead of proper classes?

While we can define any totality after a logical formula over sets, we cannot do so over classes (see e.g. [12] class existence axioms) So sets are by far more convenient than classes in reasoning. Proper classes appear here because the totality of all possible finite alphabets is a proper class.

But this also means that the totality of Turing Machines is a proper class.

Still we use a **standard normalization**, by considering finite alphabets only as subsets of a countable set, and in this normalization we consider the Turing machines as a countable set and not proper class. We use the same standard normalization and consider the P and NP in the P vs NP problem, as sets rather than classes. **So in the next after this standard normalization we will deal with P and NP as sets and not proper classes.**

So would an equation like $P=NP$ or P is not equal to NP would change when

- 1) Instead of proper classes of languages we restrict to the natural numbers $\mathbb{N}$ and the set of all subsets of them with reference to all recursive functions.**
- 2) Instead of proper classes of languages, we restrict to the full vocabulary Σ^* of a finite alphabet and the set of all subsets of it with reference to all Turing machines with states from a single countable set, and finite alphabets from a single countable set? Is it equivalent with the case in 1)?**
- 3) What if we allow extensions of the set of Turing machines in 2) over different finite alphabets not inside the original countable set? Would the problem still be well defined?**
- 4) What if we shift to Cartesian powers of the natural numbers $\mathbb{N}$ in 1)? Would such equations of complexity change? Do we still have a well defined problem?**

5) What if we shift to Cartesian powers of Σ^* in 2) ? Would such equations of complexity change? Do we still have a well defined problem?

6) What will happen if we extend the formal logic of number theory from countable to uncountable and from 1st order to higher order? Would such equations of complexity change? Do we still have a well defined problem?

Unfortunately it does not seem to exist an explicit published text which would clear out in detail all these issues of well posedness of the P vs NP problem.

The countable 1st and 2nd order predicate logic of the natural numbers, is radar that is not capable of discriminating P from NP, and prove $P \neq NP$.

But there is a happy evolution in computational complexity called Descriptive complexity mainly after the Fagin theorem and the work of Immerman (See [101], [102]. [103]. [104]). In this approach instead of being in the symbolic logic of an infinite number of possible input data structures to an algorithm, we focus to a single symbolic logic $L(k, \tau_n)$ of order k with finite initial non-logical signature τ_n (in other words finite any only initial non-logical constants, functions and relations) for each input which is considered a finite structure (e.g. a graph, a number a word etc).

We must notice though, that although, the initial non-logical symbols (non-logical signature) is finite, the total possible formulae, with higher arity or higher logical order, relations will be countably infinite.

The descriptive complexity is in the context of meta-mathematics as in its objective ontology ,is included the n -order Logic. Therefore the n -th order logic of descriptive complexity being the n th order logic (meta-ontology) of meta-mathematics which includes $n \leq \omega$ order logic , thus the logic of logic, **will be an $\omega + n$ order logic (transfinite) order logic.**

In addition the informal meta-logic of descriptive complexity is not countable logic it is uncountable. E.g. we may consider all possible graphs and their formal logics, and then take subsets of them. All possible such subsets are uncountable many.

On the other hand a logic of the mathematics (and not the informal meta-logic of descriptive complexity) if it is 1st order or 2nd order respectively is also symbolized by $L(1, \tau_n) = FO(\tau_n)$ and $L(2, \tau_n) = SO(\tau_n)$ (see [101])The particular properties of the finite structure make such a logic an axiomatic system too. (See [101], [102]. [103]. [104], but also C. Papadimitriou [18] part II, chapters 4,5).Each such logic e.g. $L(1,$

τ_n)=FO(τ_n), over a specific finite structure $G(n)$ of size n , as an axiomatic system will be the quotient or range of a functor F as in the theory of categories, of the corresponding “free” logic

$$F(FL(1, \tau_n))= L(1, \tau_n).$$

If we restrict the logic to contain only the finite initial relations of the finite structure and no other, then the functor F is a morphism of a finite **free boolean algebra** $FB(m)$ with **0-1 valuation** v (in other words a morphism v in the $\{0,1\}$ boolean algebra) that is generated , by all the atomic relations of the initial relations of the finite structure $G(n)$, to the particular boolean algebra over the particular finite structure $B(G(n))$ with 0-1 valuation,

$m \rightarrow F(FB(m))=B(G(n))$. The information contained in the ideal of this morphism of boolean algebra with valuation , is equivalent to the **assignment of 0,1** on the Cayley matrix (or adjacency matrix of it is graph). In the theory of categories this is usually written as a triangle commutative relation. And it is of course equivalent to the information of the particular **axioms** in $L(1, \tau_n)$ =FO(τ_n), that correspond to the particular finite structure $G(n)$ (e.g. graph after an **adjacency relation**, G). We must remark that any two finite boolean algebras with the same number of elements are isomorphic and they are free boolean algebras. But they may not be isomorphic when they are boolean algebras with 0-1 valuations.

This approach of descriptive complexity is a **meta-mathematical approach** too, as we study simultaneously the finite structures of the inputs and their symbolic logic $L(2, \tau_n)$ =SO(τ_n) or $L(k, \tau_n)$ too. Thus the symbolic logic of such a meta-mathematical approach is a equivalent to a logic that must transcend the studied symbolic logics , $L(k, \tau_n)$ thus of logical order $\omega+m$, for $\omega > k$, and ω being the first infinite countable ordinal number.

In this way, the ”ratio” of utilized “logic” per “data structure” , “logic/[data structure]” increases , and the “radar” becomes of lower “height” so as to detect the small differences of the complexity measures in computability.

Since, metaphorically speaking the ratio **logic/data** is also the **consciousness/matter**, this means that there is **higher intelligence** in this approach.

In the next we state that two basic theorems of interest to us of descriptive complexity, that is **Fagin’s** and **Immerman-Vardi theorems**.

RS4.1 DEFINITION. FOR THE DEFINITION OF LEAST FIX POINT EXTENSION OF 1ST ORDER LOGIC OF FINITE STRUCTURES. see [101] Martin Otto, DEFINITION 7.2.1 PAGE 77.

The idea that a 1st order sentence-scheme $\psi(P)$ can define an operator over relations P , which by iteration from the empty set, can find the relation P , as a least fix point, that satisfies the $\psi(P)$ in polynomial time, was initially observed by J. Moschovakis (see [102]). This is contrasted with the existential 2nd order sentence $\exists P \psi(P)$, that in order to find the appropriate P would search for all possible P , and this would take exponential time. But in order to have that a least fixed point exists, the operator must be monotone (as Knaster-Tarski had proved in an old theorem) and this is guaranteed when the sentence scheme ψ is positive over P . When finding the relation P over the vertices, the sentence-scheme $\psi(P)$ becomes a 1st order sentence. The least fixed point extension of the first order logic over finite structures, denoted usually by $LFP(FO(\tau_n))$, is simply the extension of $FO(\tau_n)$, by taking all possible 1st order sentence-schemes $\psi(P)$, that are positive over relations variables P and extending the sentences of $FO(\tau_n)$, with new sentences $\psi(P)$, where now P is the least fixed point, solved and known relation. The extension is done with a **single relation per formula** each time (see [101] Lemma 7.2.5 page 78, and Corollary 7.2.9)

RS4.2 FAGIN'S THEOREM (1973) (see e.g. [18] chapter 8 Theorem 8.3 or [101] theorem 6.2.3 page 73)

Any language L of the class NP can be considered as a language of finite structures, defined as the finite structures satisfied by an existential 2nd order sentence $\exists P_1 \exists P_2 \dots \exists P_k \phi$ of the 2nd order logic $SO(\tau_n)$ over finite structures. The converse is true also.

Remark. We must remark that from the proof of the theorem we realize that the number of 2nd order variables $\exists P_1 \exists P_2 \dots \exists P_k$ depends on the degree of the polynomial time, of the non-deterministic polynomial decision. Thus although it is fixed and finite for each language L , it will increase and become unbounded when considering all possible languages of NP.

RS4.3 IMMERMANN-VARDI THEOREM (1982)(see e.g. [101] theorem 7.2.8 page 79)

Any language L of the class P can be considered as a language of finite structures, defined as the finite structures satisfied by a 1st order sentence ϕ of the least fixed

point extension $LFP(FO(\tau_n))$ of the 1st order logic over finite structures, where in the initial given relations of the finite structures we also include a default order relation S_0 of the vertices. The converse is true also.

LEMMA RS4.4.0. Finite Boolean algebras Representation theorem

Let any finite Boolean algebra B of n elements. Then B is isomorphic to the Boolean algebra of the power set $p(S)$ of a set S of m elements, and $n=2^m$. The set S can be considered as the set of atoms of B .

Proof. See Theorem 4.6.1 page 210 in references [105]. This theorem can be generalized for infinite Boolean algebras provided that they are atomic and complete (complete $\Leftrightarrow$ any subset has a least upper bound in the standard order of the Boolean algebra). See also the general Stone representation theorem of Boolean algebras in [108]. QED.

In the theory of models of finite structures (e.g. graphs), each particular graph Gr with its adjacent matrix and adjacent binary relation $G(x,y)$, defines an axiomatic system for the logic of the particular graph. To be in conformance with the Immerman-Vardi theorem above, we also assume the default order relation S_0 of the vertices, thus axioms based on the atomic default order relation of the vertices. As an axiomatic system it is a finite boolean algebra that we denote by $B(S_0, G)$, with a 0-1 Boolean valuation v , or equivalently a Boolean homomorphism v , from $B(S_0, G)$ to the 2-elements Boolean algebra $\{0,1\}$. The valuation gives the information which propositions are true and which false in the axiomatic system. This we call the **non-free 1st order Logic of the particular graph Gr** .

The quotient Boolean algebra of $B(S_0, G)$, by the equivalence relation $p \sim q$ if and only if $v(p \leftrightarrow q) = 1$ is as it is known the **non-free Lindenbaum-Tarski algebra** (see references [107])

But when we want to talk about all such possible 1st order logics, independently from the particular graph Gr , then we keep the (finite and free) Boolean algebra $B(S_0, G)$ and we eliminate any 0-1 Boolean valuation. In that case we can still define **the (free) Lindenbaum-Tarski algebra**, by $p \sim q$ if and only if $p \leftrightarrow q$ is tautology in $B(S_0, G)$. Then we talk about the **free 1st order Logic** of any possible graph (or finite structure in general).

LEMMA RS4.4

Any sentence of the 1st order logic $L(1, \tau_n) = FO(\tau_n)$ of finite structures, with quantifications on 1st order variables (thus in total a 2nd order logic formula), that makes them all bounded (since it is a sentence) is equivalent to a sentence or

Boolean polynomial of the 1st order logic, without quantifications and over the atomic initial relations only.

Proof. Because the number of vertices $v_1, v_2, \dots, v_n$ of the finite structure is only finite, then also the possible values of the 1st order variables x, y, z , etc are only finite. Thus the sentence when being in **prenex normal form**, the 1st order quantification, e.g. $\forall x$ can be replaced with a finite number of conjunctions $\bigwedge_x$ and the $\exists x$ can be replaced with a finite number of disjunctions $\bigvee_x$. QED

EXAMPLE RS4.5 The simplest case of a graph with 2 only points.

In particular we shall take as sentence the 2nd order existential sentence that defines a Graph with a Hamiltonian cycle. The sentence is the next (See e.g. [18] part II, Chapter 5 example 5.11)

$$\exists P \phi(P) = \exists P (\phi_1(P) \wedge \phi_2(P) \wedge \phi_3(P) \wedge \phi_4(P))$$

The Hamiltonian cycle is defined by the linear (total) , strict order relation P on the vertices. $P(x, y)$ means $x < y$

Where

$$\phi_1(P) = \forall x \forall y (P(x, y) \vee P(y, x) \vee (x = y))$$

This is simply the statement that the order is total (linear).

$$\phi_2(P) = \forall x (\neg P(x, x))$$

This is simply the statement that the order is strict (no reflexive property).

$$\phi_3(P) = \forall x \forall y \forall z ((P(x, y) \wedge P(y, z)) \rightarrow P(x, z))$$

This is simply the statement of the transitive property of orders.

$$\phi_4(P) = \forall x \forall y ((P(x, y) \wedge \forall z (\neg P(y, z) \vee \neg P(z, y))) \rightarrow G(x, y))$$

This is the main statement for the Hamiltonian cycle. It says that

If $x < y$ and there is no intermediate z , with $x < z < y$, thus x, y consecutive, then, x and y are connected: $G(x, y) = \text{true}$, where G is the initial edge relation from the adjacency matrix of the graph.

Now we may transform the sentences $\phi_1, \phi_2, \phi_3, \phi_4$ according to the Lemma RS4. 4. where now there are not 1st order variables quantifications. They will become

$$x = v_1, v_2, \dots, v_n$$

$$y = v_1, v_2, \dots, v_n$$

$$z=v1, v2, \dots vn$$

$$\phi_1(P) = \bigwedge_x \bigwedge_y (P(x, y) \vee P(y, x) \vee (x = y))$$

$$\phi_2(P) = \bigwedge_x (\neg P(x, x))$$

$$\phi_3(P) = \bigwedge_x \bigwedge_y \bigwedge_z ((P(x, y) \wedge P(y, z)) \rightarrow P(x, z))$$

$$\phi_4(P) = \bigwedge_x \bigwedge_y ((P(x, y) \wedge (\bigwedge_z (\neg P(y, z) \vee \neg P(z, y)))) \rightarrow G(x, y))$$

For example we may take the simplest case of a graph with 2 only points v1, v2.

Then all the possible such graphs will be the next

16 cases

1) $v1. \quad v2.$

2) $v1. \rightarrow v2.$

3) $v1. \leftarrow v2.$

4) $v1. \circ v2.$

5) $v1. \quad v2. \circ$

6) $v1. \circ \quad v2. \circ$

7) $v1. \overset{\circ}{\rightarrow} v2.$

8) $v1. \rightarrow v2. \circ$

9) $v1. \overset{\circ}{\rightarrow} v2. \circ$

10) $v1. \overset{\circ}{\leftarrow} v2.$

11) $v1. \leftarrow v2. \circ$

12) $v1. \overset{\circ}{\leftarrow} v2. \circ$

13) $v1 \begin{array}{c} \rightarrow \\ \leftarrow \end{array} v2$

14) $v1 \begin{array}{c} \circ \rightarrow \\ \leftarrow \end{array} v2$

15) $v1 \begin{array}{c} \rightarrow \\ \leftarrow \end{array} v2 \circ$

16) $v1 \begin{array}{c} \circ \rightarrow \\ \leftarrow \end{array} v2 \circ$

Then for example the first ϕ_1 sentence will become

$$\phi_1(P) = \bigwedge_x \bigwedge_y (P(x, y) \vee P(y, x) \vee (x = y))$$

$$\begin{aligned} \phi_1(P) = & \\ & (P(v1, v1) \vee P(v1, v1) \vee (v1 = v1)) \wedge \\ & (P(v1, v2) \vee P(v2, v1) \vee (v1 = v2)) \wedge \\ & (P(v2, v1) \vee P(v1, v2) \vee (v2 = v1)) \wedge \\ & (P(v2, v2) \vee P(v2, v2) \vee (v2 = v2)) \end{aligned}$$

Which is of course finally equivalent to

$$(P(v1, v2) \vee P(v2, v1))$$

We may as well want the sentences in **Conjunctive Normal form**. In which case we must substitute the $a \rightarrow b$ with the $\sim a \vee b$.

The ϕ_1 and ϕ_2 are already in CNF.

The other will become:

$$\phi_3(P) = \bigwedge_x \bigwedge_y \bigwedge_z (\neg P(x, y) \vee \neg P(y, z) \vee P(x, z))$$

$$\phi_4(P) = \bigwedge_x \bigwedge_y ((\neg P(x, y) \vee (\bigvee_z (P(y, z) \wedge P(z, y)))) \vee G(x, y))$$

LEMMA RS4.6

The free logic $L(k, \tau_n)$ of order $k=1$, over finite structures with finite initial signature τ_n , when it is free, in other words independent of the axioms of the particular properties of any specific finite structure of size n , contains a free finite Boolean algebra $B(S_0, G)$ with generators the finite initial atomic relations of the standard order S_0 of the finite many vertices, the adjacency matrix entries of the adjacency relation G , and any finite set of additional relations on the finite vertices. This will be the logical context of our arguments.

Proof. If e.g. the finite structure $G(n)$ is a graph, with vertices $v1, v2, \dots, v_n$ and the initial finite relations, are 1) An order relation $S(x, y)$ 2) an edges relation $G(x, y)$, then the atomic relations are all the $S(v_i, v_j)$ and the

$G(v_i, v_j)$. Thus in total $m=2n^2$, generators, of a free finite Boolean algebra. QED.

LEMMA RS4.7

The logical equivalence of two elements or polynomials p_1, p_2 of a free finite Boolean algebra $FB(n)$, of n generators can be traced as syntactical equality of p_1 , and p_2 , when converted to conjunctive normal form of minimal length of at the most 2^n , occurrences of the n generators (variables of the polynomial) .

Proof. In a free finite Boolean algebra $FB(n)$, two polynomials or elements p_1, p_2 are equivalent if and only if $p_1 \leftrightarrow p_2 = (\text{tautology})$ or $(\sim p_1 \vee p_2) \wedge (p_1 \vee \sim p_2) = 1$. There is no need for a non-trivial 0-1 valuation. We remind also the reader that any finite Boolean algebra has isomorphism representation to the power set $P(A)$ of the set A of its atoms. Thus isomorphic to the power set $P(A)$ Boolean algebra of a set A . (See [105] Chapter 4, theorem 4.6.1 or lemma 4.4.0 above) From this is derived directly the claimed. The equivalence relation $p_1 \leftrightarrow p_2 = 1 (= \text{tautology})$ defines a quotient free Boolean algebra called free **Lindenbaum-Tarski algebra**, see references [107] QED.

LEMMA RS4.8

When transforming a sentence-scheme $\varphi(P)$ of the 1st order logic of finite structures $L(1, \tau_n) = FO(\tau_n)$ which is positive relative to a relation P , to a logically equivalent but in minimal length conjunctive normal form CNF, the property of being positive on P is preserved.

Proof. We remind the reader, that a relation P as variable in a 1st order formulae $\varphi(P)$ is called of positive occurrence, iff every occurrence of P is under the scope of an even number of negations. When transforming a positive formulae $\varphi(P)$, on P , to CNF, we can apply any of the axioms of the boolean algebra. Which are the next tautologies.

Commutativity

$$a \wedge b = b \wedge a$$

$$a \vee b = b \vee a$$

Associativity

$$(a \wedge b) \wedge c = a \wedge (b \wedge c)$$

$$(a \vee b) \vee c = a \vee (b \vee c)$$

Distribution

$$a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$$

$$a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$$

Absorption

$$a \vee (a \wedge b) = a$$

$$a \wedge (a \vee b) = a$$

Identity

$$a \vee 0 = a$$

$$a \wedge 1 = a$$

Complements

$$a \vee (\neg a) = 1$$

$$a \wedge (\neg a) = 0$$

We may see that all the above equations, have left and right sides that are positive relative to the variables, (zero times negations) except of the Completeness equations. But the complements equations do not have the chance to apply when transforming, equivalently, a positive formula $\varphi(P)$, because they involve only an odd number of negations. Instead we may apply the derived formulae

$$\neg(\neg\alpha \wedge \neg\beta) = \alpha \vee \beta$$

$$\neg(\neg\alpha \vee \neg\beta) = \alpha \wedge \beta$$

Therefore in all such equivalent transformations of a positive formulae $\varphi(P)$, on P , the positivity will be conserved. QED.

LEMMA RS4.9. *There is at least one existential 2^{nd} order sentence $\exists \varphi$ in the 2^{nd} order logic $L(2, \tau_n) = SO(\tau_n)$ of finite structures which defines a relation P of arity 2 and it is impossible to be logically equivalent to any Least Fixed Point positive formula ψ of the Least Fixed Point extension $LFP(FO(\tau_n))$ of the 1^{st} order logic $L(1, \tau_n) = FO(\tau_n)$ on finite structures, that would also define the same relation P .*

Thus the existential 2^{nd} order logic $L^{\exists}(2, \tau_n) = \exists SO(\tau_n)$ of finite structures cannot be reduced to the Least Fixed Point extension $LFP(FO(\tau_n))$ of the 1^{st} order logic $L(1, \tau_n) = FO(\tau_n)$ on finite structures

Proof.

The arguments in the proof are in the context of the free finite Boolean algebra that **Lemma RS 4.6** guarantees. In other words in the Boolean algebra $B(S_0, G, P)$.

We shall take as counter example the 2nd order existential sentence that defines a Graph with a Hamiltonian cycle of the **example rs4.5**. We convert it to a Conjunctive Normal Form CNF of minimal length , after the Lemma RS4.4. as it is done in the example RS4.5 where it is directly apparent that it is not positive on P. Then utilizing Lemma RS4.7 that reduces logical equivalence to syntactical equality, and RS4.8 that guarantees that the normalization transformation preserves the property of positivity, we prove that there is no positive Least Fixed Point extension formulae $\psi(Q)$ of the 1st order logic $FO(\tau_n)$ of finite graphs, that can be logically equivalent to it, when Q is not identical with S_0 , as is the case for general Hamiltonian graphs. What would happen if Q is not a binary relation but of higher arity? Then since that Q is checkable in polynomial time and $\phi(P) \leftrightarrow \psi(Q)$, then also P which is binary is checkable in polynomial time, thus we may assume that Q may be binary. So if there is a positive over a binary relation Q, 1st order sentence scheme $\psi(Q)$, logically equivalent to $\exists \phi(P)$, (we are in the free finite Boolean algebra $B(S_0, G, P)$) then we convert it after Lemmas RS4.7 and RS4.8 to conjunctive normal form $[\psi(Q)]$, (symbolizing by $[\psi(Q)]$ the $\psi(Q)$ in CNF) and it will preserve the positivity over Q. Then it must be syntactically identical with the $[\phi(P)]$ in conjunctive normal form after Lemma RS 4.7. We must notice here that logical equivalence in the non-free 1st order Logic of all particular Hamiltonian graphs will imply logical equivalence in the free 1st order logic (free Lindenbaum-Tarski algebra) of any graph. But as we observe from the $[\phi(P)]$ in CNF in the above example RS4.5 ,it is not positive relative to P, thus it can never be the case that $Q=P$, and $[\phi(P)]=[\psi(Q)]$. QED.

COROLLARY RS4.10

There is no single polynomial time algorithm for every Graph G, which can find a Hamiltonian cycle H of G, if it exists.

Proof: Hint. We look at the Immerman-Vardi theorem RS4.3 above. If the initial given order relation $S(x,y)$ among the vertices of the graph coincides with the Hamiltonian cycle, then the, Hamiltonian cycle is known in polynomial time. But this is not the general case for any graph. If it does not coincide, and the Hamiltonian cycle is defined by another order relation $P(x,y)$, then , using proof by contradiction, from the Immerman-Vardi theorem, this relation $P(x,y)$ since it is found in polynomial time , it must be the Least Fixed Point relation P, of a sentence-scheme $\psi(P)$ which is positive over P. But then from the Lemma RS4.9 and the particular

choice in the above lemma, the P to be the definition of a Hamiltonian cycle, this is not possible. QED

COROLLARY RS4.11

The class P of polynomial time deterministically decidable languages cannot be equal to the class NP of polynomial time non-deterministically decidable languages. Or $P \neq NP$, or P not equal to NP ,

Remark. As we mentioned already above this proof does not contradict the non-provability of $P \neq NP$ in a countable n th- order logic of set theory. This is so because, the above proof of $P \neq NP$ is **within the meta-mathematical approach of Descriptive complexity**. This approach of descriptive complexity is a **meta-mathematical approach**, as we study simultaneously the finite structures of the inputs and their symbolic logic $L(2, \tau_n) = SO(\tau_n)$ or $L(k, \tau_n)$ too. Thus the symbolic logic of such a meta-mathematical approach is a equivalent to a logic that must transcend the studied symbolic logics, $L(k, \tau_n)$ thus of logical order $\omega + m$, for $\omega > k$, and ω being the first infinite countable ordinal number.

RS5) THE EQUIVALENCE OF THE SCHEME OF BOUNDED μ -OPERATOR AND SIMPLE INDUCTION IN THE DEFINITIONS OF THE CLASSICAL RECURSIVE FUNCTIONS.

To create any general recursive functions, we utilize (see references [45], [51], [59], [62]).


0) We start with the basic an elementary recursive functions (The identity map, the constants, the function of the next integer etc. They are all poly-time in complexity).

1) Finite many compositions. (The compositions preserve the poly-time complexity of the components)

2) The simple definition by induction (This scheme in general introduces exponential complexity relative to the digits of the arguments.)

**3) The general unbounded μ -operator over a decidable recursive predicate.
(This scheme in general introduces exponential and higher than exponential complexity relative to the digits of the arguments)**

The recursive functions defined with only the 0, 1, 2, are the old Goedel recursive functions and are called primitive recursive functions. It was the 1926 problem of Hilbert, solved by Sudan- Ackermann (see references [54],[55],[56],[57]) , which proved that the general recursive functions are not identical with the primitive recursive functions and contain the primitive recursive functions.



Definition 6.0. *Generic form of the schemes of bounded μ - operator and simple induction means that we do not impose any restriction on the form, which would exclude any of the possible instances of applications of it. Thus a generic form of the scheme , includes all possible instances of applications of it.*

Lemma 6.0 *The generic scheme 3) of unbounded μ -operator is not reducible (in other words, cannot be substituted), by the generic scheme 2) of simple induction and the generic scheme 1) of composition.*

Proof. Well known in the theory of recursive functions. That the 3) general unbounded μ -operator cannot be reduced to the 2) simple induction scheme, was the result of the solution of the 1926 Hilbert problem by Sudan and Ackermann. That the simple scheme of induction is not reducible to the composition scheme is deduced as following. The exponential function of natural numbers $c=a^b$ is known that it is definable with the simple scheme of induction. And it is also known (from $10^{(\text{Log}b)*\text{Log}a}=\text{Log}c$) that the number of decimal digits of the output is exponential function of the number of decimal digits of the input. Thus it cannot be computed by any algorithm in polynomial time. Nevertheless, it is easy to notice that the composition scheme starting from the basic recursive functions defines only polynomial time functions by itself. Thus the simple induction scheme cannot be reduced to the composition scheme.

QED.

Definitions 6.1 We assume familiarity of the reader with concepts like **predicates** $R(x)$ in formal logic of an axiomatic system, and their **representing function** $C_R(x)=0$, (negation of the characteristic function of the set of x that satisfy R) Following David Martin in the references [48], we define as **(general) recursive predicate** $R(x)$ a single free variable predicate that its representing function $C_R(x)=0$ is recursive thus, the set of natural numbers that satisfy R , is a recursive (computable decidable) set. If the representing function is a primitive recursive function the predicate is called **primitive recursive predicate**. Any single free variable recursive predicate with a finite number of universal and existential quantifications of its bounded variables is called **elementary recursive predicate**. Predicates with an existential quantification in the front (left side) are called **recursive enumerable predicates**. A predicate of the form $P=0$, where P is a polynomial with non-negative integer coefficients is called **polynomial predicate**. A polynomial predicate with an existential quantification in the front (left side) is called **Diophantine predicate**. Finally any predicate formed by any finite number sequence of existential and universal quantifiers placed in front of a polynomial predicate is called **arithmetic predicate**.

Although by the scheme of composition of functions is obvious what we mean, let us be more explicit about the scheme of definition by simple inductions

Definition 6.2. Simple inductive scheme

By scheme of definition by simple induction we mean the

Definition of a function $z = f(\vec{x}, y) : N_0^{n+1} \rightarrow N_0$

Through the functions g and h and the next equations

$$g : N_0^n \rightarrow N_0$$

$$h : N_0^{n+2} \rightarrow N_0$$

$$n \geq 0$$

$$\vec{x} \in N_0^n$$

$$y \in N_0$$

$$f(\vec{x}, 0) = g(\vec{x})$$

$$f(\vec{x}, y+1) = h(\vec{x}, y, f(\vec{x}, y))$$

It is also instructive to define explicitly what are the **unbounded and bounded minimization operators (μ -operators)**, which make an existential quantified implicit relation-predicate to a recursive function.

Definition 6.3. Unbounded minimization μ -operator.

*Let $P(\vec{x}, y)$ be a $n+1$ -variables **recursive predicate**, y in N_0*

$$\vec{x} = (x_1, \dots, x_n)$$

$$\vec{x} \in N_0^n$$

And Y the set of natural numbers that satisfies it.

$$Y = \{y / P(\vec{x}, y)\}$$

We define as the unbounded minimization μ -operator as follows, and symbolize the function f derived from P by μ finally as below.

$$\mu : \vec{x} \rightarrow y = 0, \text{ if } Y = \text{empty}$$

$$\mu : \vec{x} \rightarrow y = \min Y, \text{ if } Y \text{ is not empty}$$

$$f(\vec{x}) = \mu y P(\vec{x}, y)$$

We assumed that the predicate P is recursive (decidable). Let us examine if the function f is also recursive.

1) If there exist y such that $P(\vec{x}, y)$

is satisfied, then we scan in sequence the values $y=0, 1, \dots$ till the first time that we find a y which satisfies P . According to the definition of f , this is exactly the value of the function f , by the μ -operator thus f is recursive.

2) If there is no y that satisfies P then the previous process will not halt. Nevertheless if we make the **normal assumption** (as it is called) that for each

$$\vec{x} = (x_1, \dots, x_n)$$

$$\vec{x} \in N_0$$

there exist y which satisfies P , then the above process of the μ -operator also defines **a recursive function f** .

Definition 6.4. Bounded μ -operator.

Let $P(\vec{x}, y)$ be a $n+1$ -variables **recursive predicate**, y in N_0

$$\vec{x} = (x_1, \dots, x_n)$$

$$\vec{x} \in N_0$$

And Y_0 the set of natural numbers defined by .

$$Y_0 = \{z / P(\vec{x}, z) \wedge 0 \leq z \leq y\}$$

We define as the bounded minimization μ -operator as follows, and symbolize the function f derived from P by μ finally as below.

$$\mu_{z=0}^y : \vec{x} \mapsto z = \{0, \text{if } Y_0 = \text{the empty set}\}$$

$$\mu_{z=0}^y : \vec{x} \mapsto z = \{\min Y_0, \text{if } Y_0 \text{ is not the empty set}\}$$

We Symbolise it by

$$\mu_{z=0}^y P(\vec{x}, z) = z$$

and symbolize the function f derived from P by μ finally as below.

$$f(\vec{x}, y) = \mu_{z=0}^y (z), P(\vec{x}, z)$$

Obviously the essential difference of the bounded μ -operator from the unbounded is when the set Y is empty. We notice that when P is recursive (decidable) even if we do not make the “**normal assumption**” the application of the **bounded μ -operator** will yield a recursive function. We notice also that although the application of the unbounded μ -operator on an $n+1$ predicate P yields a n -variable function, the application of the bounded μ -operator, on the same predicate yields a $n+1$ variables function f , where y is the upper-bound of the μ -operator. .

Lemma 6.1.

Let g_i and P_i , $i=1,\dots,m$ be primitive recursive functions and primitive recursive predicates correspondingly. Then the function f , defined by the next equations is also primitive recursive.

$$f(\vec{x}) = g_i(\vec{x}), \text{ if } P_i(\vec{x}), \text{ holds, } i = 1, \dots, m$$

Proof:

Let, $C_{P_i}(\vec{x})$, be the corresponding characteristic functions of the primitive recursive predicate P_i . We assume that for every $\vec{x}$, there is, exactly, one, P_i , satisfied, by, $\vec{x}$. Then the f can also be written as follows

$$f(\vec{x}) = g_1(\vec{x})C_{P_1}(\vec{x}) + \dots + g_m(\vec{x})C_{P_m}(\vec{x})$$

And it is also primitive recursive. QED.

Corollary 6.0. (Finite sequences are recursive functions)

Any finite sequence a_i , $i=1,\dots,n$ is a recursive function $S(i)=a_i$, $i=1,\dots,n$

Proof: We apply the Lemma 6.0 for $g_i=a_i$, and P_i simply being that the value of the variable is i . QED.

In the next Lemma we prove the **equivalence** of the simple induction and bounded μ -operator as creation schemes of recursive and primitive recursive functions.

Lemma 6.2 The generic bounded μ -operator can be reduced to the simple induction scheme, and conversely the generic simple induction scheme can be reduced to the bounded μ -operator.

Proof.

A part.

The first part, is already known in the theory of primitive recursive functions, but we shall repeat the proof, for reasons of completeness.

So let Let $P(\vec{x}, y)$ be a $n+1$ -variables **recursive predicate**, y in N_0

$$\vec{x} = (x_1, \dots, x_n)$$

$$\vec{x} \in N_0$$

and

$$f(\vec{x}, y) = \mu_{z=0}^y (z), P(\vec{x}, z)$$

We shall prove that f is a primitive recursive function, by reducing it to the simple induction scheme.

We define

$$h(\vec{x}, y, t) = t, \text{ if } \exists z \in \{0, 1, \dots, y\} : P(\vec{x}, z)$$

$$h(\vec{x}, y, t) = y + 1, \text{ if } \neg \exists z \in \{0, 1, \dots, y\} : P(\vec{x}, z) \text{ But, it, holds, } P(\vec{x}, y + 1)$$

$$h(\vec{x}, y, t) = 0, \text{ in, all, other, cases}$$

Thus it holds that

$$\begin{aligned} f(\vec{x}, y+1) &= f(\vec{x}, y), \text{ if } \exists z \in \{0, 1, \dots, y\} : P(\vec{x}, z) \\ f(\vec{x}, y+1) &= y+1, \text{ if } \neg \exists z \in \{0, 1, \dots, y\} : P(\vec{x}, z), \text{ but } P(\vec{x}, y+1) \\ f(\vec{x}, y+1) &= 0, \text{ in, all, other, cases.} \end{aligned}$$

Therefore f satisfies the simple induction scheme

$$\begin{aligned} f(\vec{x}, 0) &= 0 \\ f(\vec{x}, y+1) &= h(\vec{x}, y, f(\vec{x}, y)). \end{aligned}$$

And is therefore primitive recursive.

B part.

We will prove that the generic simple induction scheme as in the Definition 6.0 is reducible equivalently to a bounded μ -operator.

$$g : N_0^n \rightarrow N_0$$

$$h : N_0^{n+2} \rightarrow N_0$$

$$n \geq 0$$

$$\vec{x} \in N_0^n$$

$$y \in N_0$$

$$f(\vec{x}, 0) = g(\vec{x})$$

$$f(\vec{x}, y+1) = h(\vec{x}, y, f(\vec{x}, y))$$

For this we shall use the tools used to prove the **10th Hilbert problem**, presented by David Martin in references [47] end of the page 259, where he proves that Diophantine functions are closed to the scheme of (primitive) simple induction, by reducing it to existential quantification (Diophantine predicates). We simply copy his formula of predicates of David Martin, after simplifying it, in the number of variables, and using our **Corollary 6.0. (Finite sequences are recursive functions)**, with $S(i)$, in place of his number theoretic $S(i, u)$ which has an additional variable u , after his theorem 1.3 in [47], because our version of the finite sequence theorem (corollary 6.0) is simpler without the need of a u . Then the equivalence takes the form

$$f(\vec{x}, z) = S(\mu_{t=1}^z t) \{ [\forall t (s(1) = y(\vec{x}) \wedge [(t \geq z) \vee (S(t+1) = h(t, S(t), \vec{x}))]) \wedge (y = S(z))]\}$$

Since with this formula a bounded μ -operator defines, what a simple induction scheme can define we have completed the proof. QED.

RS6.7) Further perspectives in Computational Complexity A. Surreal (ordinal) natural numbers and the measurement of the computational complexity . The phenomenology of the changes of the finite is the ontology of the infinite. Irreducible minimum complexity of languages?

This paragraph introduces a model of the infinite order of ordinal numbers up to ω^{ω^ω} on to the speeds of convergence of computable functions to the infinite. Such models are not strange after the famous metamathematical Lowenheim-Skolem theorem that any axiomatic theory has a countable model. The Sudan computable function was conceived initially by modelling the order of the ordinal ω^{ω^n} on to appropriate orders of computability defined by the computation scheme. (see [57] Sudan G. 1927)

Great other mathematicians like Grentzen, Hilber, Bernays, Ackermann, have also made models of the order of ordinal numbers in appropriate orders of countable mathematical systems. (See e.g. references [58] Maria Hameen Attila. Nominalistic Ordinals)

Computational complexity, as **an axiomatic system** is essentially the axiomatic system of **set theory** of mathematics. Set theory is an axiomatic theory traditionally used as a foundation of all the mathematics. For example the natural umbers can be defined as von Neumann indicated from the empty set , and sets of sets.

First few von Neumann ordinals

$$0 = \{ \} = \emptyset$$

$$1 = \{0\} = \{\emptyset\}$$

$$2 = \{0, 1\} = \{\emptyset, \{\emptyset\}\}$$

$$3 = \{0, 1, 2\} = \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\}$$

$$4 = \{0, 1, 2, 3\} = \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}, \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\}\}$$

And the set theory in mathematics after the turn of the 20th century and the contribution of G Cantor has officially introduced a phenomenological abstraction in mathematics called **the infinite**. Below is the axiom in the axiomatic system of Zermelo-Frankel of the set theory (see references Wikipedia, “Zermelo Frankel Set Theory”).)

7. Axiom of infinity [\[edit\]](#)

Main article: [Axiom of infinity](#)

Let $S(w)$ abbreviate $w \cup \{w\}$, where w is some set. (We can see that $\{w\}$ is a valid set by applying the axiom of pairing with $x = y = w$ so that the set z is $\{w\}$.) Then there exists a set X such that the empty set $\emptyset$, defined axiomatically, is a member of X and, whenever a set y is a member of X then $S(y)$ is also a member of X .

$$\exists X [\exists e (\forall z \neg (z \in e) \wedge e \in X) \wedge \forall y (y \in X \Rightarrow S(y) \in X)].$$

More colloquially, there exists a set X having infinitely many members. (It must be established, however, that these members are all different because if two elements are the same, the sequence will loop around in a finite cycle of sets. The axiom of regularity prevents this from happening.) The minimal set X satisfying the axiom of infinity is the [von Neumann ordinal](#) ω which can also be thought of as the set of [natural numbers](#) $\mathbb{N}$.

First few von Neumann ordinals	
$0 = \{ \}$	$= \emptyset$
$1 = \{0\}$	$= \{\emptyset\}$
$2 = \{0, 1\}$	$= \{\emptyset, \{\emptyset\}\}$
$3 = \{0, 1, 2\}$	$= \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\}$
$4 = \{0, 1, 2, 3\}$	$= \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}, \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\}\}$

Before Cantor of course, the concept of infinite many was implicit in **Euclid's elements**, in the axiom that between every two different points in a line segment there is always a 3rd different point between them.

And of course **Newton and Leibniz** in their famous infinitesimal calculus utilized concepts of the infinite. Newton term for the instantaneous velocity of a “moving quantity” was “**Fluxion**” and it was his physical intuition for the derivative of a function. Nevertheless if that fluxion is of a moving quantity towards the infinite then such a fluxion is nothing else than **a measure of computational complexity** of a Turing machine.

This *phenomenology of the changes of finite* quantities, turned eventually to become *an ontology of the infinite*.

It is known that in computational Complexity theory is used the big-O

(see references [22] Wikipedia “Big O Notation”). In other words for two sequences a_n, b_n and that converge to infinite we write

$a_n = O(b_n)$ iff, there is real constant $M > 0$ such that $a_n \leq M * b_n$, finally in n .

But a more exact measurement of the speed of the convergence to the infinite is of course with the Newtonian fluxions , in other words we say that two sequences

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 1$$

converging to the infinite, have the same speed iff finally in n . And the speed of

a_n is larger than that of b_n iff $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = M, M > 1 \text{ or } M = \infty$ which of course includes

the **little-o** notation in the case of $M = \infty$ where we write $a_n \gg b_n$, as in the case, where a_n has exponential speed while b_n only polynomial speed. This kind of ordering in classical mathematics is called **non-Archimedean**, or in computer science **pre-emptive** orders.

An order is Archimedean if it is like of two real numbers $a < b$, where there is always an integer N , such that $Na > b$. But in the case of polynomial speed a_n and exponential speed b_n there is no integer N such that the speed of Na_n is larger than that of b_n .

We could conceive the phenomenology of the polynomial and exponential speed of convergence to infinite as **an ontology of infinite numbers**.

Continuing the creation of the natural numbers in set theory as by v. Neumann, the axiom of infinite gives that the set of natural numbers $N = \{1, 2, 3, \dots, n, \dots\}$ does exist and we may consider it an new infinite (ordinal) number denoted by ω . Compared to the linear (total) order of the rational numbers and real numbers, the natural numbers have the additional good property **that every subset of the natural numbers has a smallest element**. Such linear orders are called **well orders**.

This may motivate the standard definition, suggested by John von Neumann at the age of 19, now called definition of **von Neumann ordinals** also for large infinite sets "each ordinal is the well-ordered set of all smaller ordinals".


Definition 7.1 *A set S is an ordinal if and only if S is strictly well-ordered with respect to set membership and every element of S is also a subset of S .*

The natural numbers are thus ordinals by this definition. For instance, 2 is an element of $4 = \{0, 1, 2, 3\}$, and 2 is equal to $\{0, 1\}$ and so it is a subset of $\{0, 1, 2, 3\}$.

It can be shown by transfinite induction that every well-ordered set is order-isomorphic to exactly one of these ordinals, that is, there is an order preserving bijective function between them.

(see references [34] Wikipedia Ordinal Number) Before von-Neumann an ordinal number was simply a well ordered set.

The next diagrams show how the ω the first infinite (limit) ordinal number, then we may define $\omega+1$, $\omega+2$ etc



Or in spiral way

(For a reference to standard symbolisms and definitions for ordinal numbers, see [Cohn P.M. 1965] p.1-36 also [Kuratowski K.-Mostowski A. 1968]).

Definition 7.2. The properties of such an algebra (well ordered commutative semiring with cancellation law) N are the next.

The following hold $(x, y, z, c, x', y', \in N = W(\alpha))$.

$$0) \quad s(x) = x+1 \text{ belongs in } N \quad \text{for every } x \in N$$

$$1) \quad x+y = y+x, \quad x \cdot y = y \cdot x$$

$$2) \quad x+(y+z) = (x+y)+z, \quad x \cdot (y \cdot z) = (x \cdot y) \cdot z$$

$$3) \quad x+0 = 0+x = x,$$

$$x \cdot 0 = 0$$

$$x \cdot 1 = 1 \cdot x = x$$

$$4) \quad x(y+z) = xy+xz$$

$$5) \quad \left. \begin{array}{l} x+c = y+c \Rightarrow x=y \\ c \in w(a) \end{array} \right\} \Rightarrow x=y$$

$$6) \quad \text{If } x > y, x' > y' \text{ then } x+x' > y+y', \text{ and } xx'+yy' > xy+yx'$$

Such well ordered commutative semirings with cancellation law, are not only consisting by natural numbers but may be larger (non-Archimedean) consisting from ordinal numbers.

Let us see now how such **semi-rings of ordinal numbers** (sometimes also called **surreal natural numbers**, after the introduction of the surreal numbers by J.H. Conway (see references [31] [Conway J.H. 1976.] and [35] [Kyritsis C.E. 1991]) can be used to define e.g. the polynomial and exponential complexity measures as well ordered semiring.

Instead of resorting to the definition of the ordinal numbers by v. Neumann and then defining commutative addition and multiplication, we shall have an equivalent but **elementary definition of the computational complexity speeds with polynomials and lexicographic order of them.**

For a Turing machine M as we mentioned it has **polynomial computational complexity** if the duration of the computation in steps, is of the order $O(p(n))$. Where by $O(p(n))$ is meant (see Wikipedia [22] [16]) that if $T(n)$ is the time that the Turing machines terminate as a function of the size n of the initial data, then $T(n) \leq M \cdot p(n)$ for some positive constant M not depending on n , for all n , thus as $n \rightarrow +\infty$.

As we mentioned also above this also means that the fluxion speed of convergence defined by the limit of the ratio of $T(n)$ and $p(n)$ is comparable with real positive constant.

So we start measuring the speeds of convergence to infinite by ordinals.

- 1) The standard polynomial speed of the sequence n we define as the ordinal ω
- 2) The polynomial speed of n^k we define as the ordinal ω^k
- 3) The exponential speed n^{n^k} we define as the ordinal ω^{ω^k}
- 4) In general the polynomial speed $p(n)$ where p is a polynomial of n with positive integer coefficients with the ordinal $p(\omega)$
- 5) In general the exponential speed $n^{p(n)}$ as the ordinal $\omega^{p(\omega)}$
- 6) Furthermore the order of the speeds e.g. of the polynomials $p(n)$ with positive integer coefficients is the lexicographic order of the coefficients of the polynomial (the highest order of n is in priority in the lexicographic order of the coefficients). An this well orderings is isomorphic with the well ordering of the corresponding ordinals.
- 7) The sum and product of such speeds corresponds to the sum and formal product of the corresponding ordinals. These commutative operations of sum and product of the ordinals are known as **the natural sum and product of the ordinals defined by the ω -normal forms of the ordinals.**
- 8) Therefore we get that all such ordinals of the form $q(\omega^{p(\omega)})$ for more than two polynomials q, p , with natural numbers as coefficients, (which are actually all the ordinals less than the ordinal $\omega^{\omega^{\omega}}$) form a well ordered, commutative semiring with cancellation law as the **Definition 7.2** which analogous to the natural numbers and is also a particular size such semiring of **surreal natural numbers**(see references [31] [Conway J.H 1976.] and [35] [Kyritsis C.E. 1991]). **We may denote this semiring by $N(\omega^{\omega^{\omega}})$**

9) *In more detail the elements of $N(\omega^{\omega^{\omega}})$ are expressions of ordinals of the form $a_n \omega^{p_n(\omega)} + \dots + \alpha_1 \omega^{p_1(\omega)} + \alpha_0$ for positive integers a_n, α_1 , and polynomials p_n, p_1 of positive integers, in descending order in their lexicographic order. Such forms are special cases of **ω -normal forms of ordinals***

It is interesting to inquire what happens to higher speeds of convergence or higher measures of computational complexity. The axiom of choice of set theory, in its form as the maximum principle for ordered sets guarantees that there is a maximum linearly ordered set of speeds of convergence to infinite or computational complexity measures, that contain the polynomial and exponential measures. .

Another interesting point relevant to the Newton-Leibniz infinitesimal calculus is the next. If we utilize the semiring $N(\omega^{\omega^{\omega}})$ as a kind of infinite natural numbers, and from this we repeat the construction of the real numbers, by 1) defining quotients for the rational numbers 2) taking the completion of the field of rational numbers by Cauchy fundamental sequences (nevertheless here with transfinite such sequences with indexes by ordinals in $N(\omega^{\omega^{\omega}})$) then we end with a linearly ordered field which we may denote by $R(\omega^{\omega^{\omega}})$, which actually is a sub-field of the field of the surreal numbers of J.H. Conway (see references [31] [Conway J.H. 1976.] and [35] [Kyritsis C.E. 1991]), This also could be called a Newton-Leibniz field of finite, infinite and infinitesimal numbers, and it is exactly what Newton-Leibniz could not define in detail, for their infinitesimal calculus. That is why after 2-3 centuries this approach was abandoned, in favour of the Weierstrass calculus with limits restricted only within the real numbers. **It is surprising that by inverting measures of computational complexity, the so long undefinable infinitesimals of Newton-Leibniz are eventually definable.**

8. More perspectives in Computational Complexity. Seemingly infinite languages and the measurement of the computational complexity.

A completely opposite direction of development of Computational Complexity from towards the infinite is, towards a more realistic perception of the computational complexity where there is no infinite at all, not even the countable infinite. We could call it completely digital computational complexity without the infinite. For example in such a perception, there are no infinite languages. Only finite languages. Theoretically if a language L is finite, we can put all its words, as input data, to Turing machine M , which can decide in polynomial time (relative to the length of an arbitrary word w_n , of length n), if w_n belongs to L or not. Thus much of the computational complexity would become obsolete! Still if the finite language is of a

very large number of words (according to appropriate definitions could be called “seemingly infinite” language), and if we have an **upper bound K on the space complexity (size of the code of M)**, and this K prohibits putting all of L in the code of M, then we might need e.g. exponential time algorithms to decide or enumerate the (“seemingly infinite”) language L. Therefore with more refined and realistic definitions in computational complexity theory that avoid the infinite , we may have a considerable part of the theory of computational complexity appropriately modified, for such large finite languages.

9.Epilogue . The mind creates miracles.

References.

[1] Woeginger, G.J. (2016) The P versus NP Page.
<https://www.win.tue.nl/~gwoegi/P-versus-NP.htm>

- [2] Wikipedia, “P versus NP Problem”. https://en.wikipedia.org/wiki/P_versus_NP_problem
- [3] Yannakakis, M. (1998) Expressing Combinatorial Optimization Problems by Linear Programs. *Proceedings of STOC*, Chicago, 2-4 May 1988, 223-228. <https://doi.org/10.1145/62212.62232>
- [4] Kyritsis, C. (2017) On the Solution of the 3rd Clay Millennium Problem. A Short and Elegant Proof That $P \neq NP$ in the Context of Deterministic Turing Machines and Zermelo-Frankel Set Theory. *Proceedings of the 1st ICQSBEI 2017 Conference*, Athens, Greece, 170-181.
- [5] Kyritsis, K. (2018) The Solution of the 3rd Clay Millennium Problem. A Short Proof That $P \neq NP = \text{Exptime}$ in the Context of Zermelo Frankel Set Theory. *International Journal of Pure and Applied Mathematics*, **120**, 497-510. <http://www.ijpam.eu>
- [6] Kyritsis, K. (2021) Study on the Solution of the Clay Millennium Problem about the P vs. NP: A Short Proof that $P \neq NP = \text{EXPTIME}$ in the Context of Deterministic Turing Machines. In: *New Visions in Science and Technology*, Vol. 6, BP International, India, 60-69. <https://doi.org/10.9734/bpi/nvst/v6/5176F> <https://stm.bookpi.org/NVST-V6/article/view/4135>
- [7] Kyritsis, K. (2022) A Short and Simple Solution of the Millennium Problem about the Navier-Stokes Equations and Similarly for the Euler Equations. *Journal of Applied Mathematics and Physics*, **10**, 2538-2560. <https://www.lap-publishing.com/catalog/details//store/gb/book/978-620-4-72562-8/the-solutions-of-the-3rd-and-4th-millennium-mathematical-problems>
- [8] Kyritsis, K. (2023) A 3rd Shorter Solution of the Clay Millennium Problem about $P \neq NP = \text{EXPTIME}$. *Conference: 6th International Conference on Quantitative, Social, Biomedical and Economic Issues*, Athens, 1 July 2022, 81-89. <https://icqsbei2022.blogspot.com/2022/06/blog-post.html> <http://books.google.com/books/about?id=xZnCEAAQBAJ>
- [9] Frankel, A.A. (1976) *Abstract Set Theory*. North Holland Publishing Company, Amsterdam.
- [10] Wikipedia, “Zermelo Frankel Set Theory”. https://en.wikipedia.org/wiki/Zermelo%E2%80%93Fraenkel_set_theory
- [11] Thomas, Jech . (1978) *Set Theory*. Academic Press, Cambridge.
- [12] Wikipedia “Von Neumann-Bernays-Gödel set theory” https://en.wikipedia.org/wiki/Von_Neumann%E2%80%93Bernays%E2%80%93G%C3%B6del_set_theory
- [13] Manin, Y.I. (2010) *A Course in Mathematical Logic for Mathematicians*. Springer, Berlin. <https://doi.org/10.1007/978-1-4419-0615-1>
- [14] Wikipedia, “Second-order Logic”. https://en.wikipedia.org/wiki/Second-order_logic
- [15] Cook, S. (2000) *The P versus NP Problem (PDF)*, Clay Mathematics Institute Site.

- [16] Wikipedia, “Turing Machine”. https://en.wikipedia.org/wiki/Turing_machine
- [17] Martin, J.C. (1997) Introduction to Languages and the Theory of Computation. 2nd Edition, McGraw-Hill, London.
- [18] Papadimitriou, C. (1994) Computational Complexity. Addison-Wesley, Boston.
- [19] Wikipedia, “Nondeterministic Turing Machine”. https://en.wikipedia.org/wiki/Nondeterministic_Turing_machine
- [20] Wikipedia, “DTIME”. <https://en.wikipedia.org/wiki/DTIME>
- [21] Wikipedia, “Time Complexity”. https://en.wikipedia.org/wiki/Time_complexity#Polynomial_time
- [22] Wikipedia “Big O Notation”. https://en.wikipedia.org/wiki/Big_O_notation
- [23] Wikipedia, “P (Complexity)”. [https://en.wikipedia.org/wiki/P_\(complexity\)](https://en.wikipedia.org/wiki/P_(complexity))
- [24] Lewis, H.R. and Papadimitriou, C.H. (1981) Elements of the Theory of Computation. Prentice-Hall, Englewood Cliffs.
- [25] Wikipedia, “EXPTIME”. <https://en.wikipedia.org/wiki/EXPTIME>
- [26] Trevisan, L. (2009) Notes on Hierarchy Theorems. University of California, Berkeley.
- [27] Hartmanis, J. and Stearns, R.E. (1965) On the Computational Complexity of Algorithms. *Transactions of the American Mathematical Society*, **117**, 285-306. <https://doi.org/10.1090/S0002-9947-1965-0170805-7>
- [28] Stanislav, Ž. (1983) A Turing Machine Time Hierarchy. *Theoretical Computer Science*, **26**, 327-333. [https://doi.org/10.1016/0304-3975\(83\)90015-4](https://doi.org/10.1016/0304-3975(83)90015-4)
- [29] Kyritsis, K. (2021) Review of the Solutions of the Clay Millennium Problem about $P \neq NP = EXPTIME$. *World Journal of Research and Review*, **13**, 21-26. <https://www.wjrr.org/vol-13issue-3>
<https://doi.org/10.31871/WJRR.13.3.8>
- [30] Mathoverflow. If $NP = EXPTIME$, Does Every DTM Have a Succinct “Execution Proof”? <https://mathoverflow.net/questions/114887/if-np-exptime-does-every-dtm-have-a-succinct-execution-proof>
- [31] [Conway J.H 1976.] On numbers and games Academic press 1976 .
- [32][Kuratowski K. -Mostowski A.1968] Set Theory North - Holland 1968.
- [33] [Cohn P.M.1965] Universal Algebra Harper - Row 1965 .
- [34] Wikipedia Ordinal Number
https://en.wikipedia.org/wiki/Ordinal_number

[35] [Kyritsis C.E.1991] Algebraic characterisation of the Hessenberg operations in the ordinal numbers. .

https://www.researchgate.net/publication/341734963_ORDINAL_NATURAL_NUMBERS_1_Alternative_algebraic_definitions_of_the_Hessenberg_natural_operations_in_the_ordinal_numbers

[36] Malinina Natalia “ Necessary and sufficient conditions for dividing the structure of algorithms into non-intersecting sets: polynomial and enumeration algorithms”
<https://journals.rudn.ru/engineering-researches/article/view/31716/20976>

[37] Cook S. 1971. "The complexity of theorem-proving procedures" *Proceedings of the third annual ACM symposium on Theory of computing - STOC '71*. pp. 151–158

[38] Wikipedia P-complete, <https://en.wikipedia.org/wiki/P-complete>

[39] Sipser M. *Introduction to the Theory of Computation*, 3rd edition Gengage Learning 2013

[40] Wikipedia P-complete <https://en.wikipedia.org/wiki/P-complete>

[41] Wikipedia Von Neumann–Bernays–Gödel set theory

https://en.wikipedia.org/wiki/Von_Neumann%E2%80%93Bernays%E2%80%93G%C3%B6del_set_theory

[42] Salil Vadhan, Harvard University Lectures in Computational complexity.
<https://people.seas.harvard.edu/~salil/cs221/spring14/spr10-lec8.pdf>

[42b] Stanford University Rice Theorem.
<http://kilby.stanford.edu/~rvrg/154/handouts/Rice.html>

- [43] Wikipedia Probability axioms
https://en.wikipedia.org/wiki/Probability_axioms
- [44] Timothy Moon-Yew Chan- Ruta Mehta, Lecture 22 P and NP
 University of Illinois [Grainger College of Engineering](#)
- [45] Wikipedia General recursive function
https://en.wikipedia.org/wiki/General_recursive_function
- [45b] Wikipedia Universal Turing machine
https://en.wikipedia.org/wiki/Universal_Turing_machine
- [46] Minsky, Marvin (1962). "[Size and Structure of Universal Turing Machines using Tag Systems, Recursive Function Theory](#)". *Proc. Symp. Pure Mathematics*. Providence RI: American Mathematical Society. 5: 229–238.
- [47] Davis Martin “Hilbert’s 10th problem is unsolvable” *The American Mathematical Monthly* , Vol. 80, 1973, pp 233-269.
- [48] Davis Martin “Arithmetical Problems and Recursive Enumerable Predicates” *The Journal of Symbolic Logic*. Mar. 1953, Vol. 18, No 1 pp 33-41.
- [49] Goedel Kurt “Über formal unentscheidbare satze der Principia Mathematica und verwandter Systeme I, Monatshefte für mathematik und Physik, vol. 38,(1931) pp.173-198.
 Translated by B. Meltzer , Introduction by R.B. Braithwaite .
https://monoskop.org/images/9/93/Kurt_G%C3%B6del_On_Formally_Undecidable_Propositions_of_Principia_Mathematica_and_Related_Systems_1992.pdf
- [50] Wikipedia “Pell’s equation (Pell-Fermat equation)
https://en.wikipedia.org/wiki/Pell%27s_equation
- [51] Kleene S.C. 1936 “General Recursive functions of natural numbers “
Mathematicshe Annallen vol. 112 (1936) pp 727-742
- [52] Kleene S.C. 1943 “Recursive predicates and quantifiers “
Transactions of the American mathematical Society vol. 53 (1943) pp 41-73

- [53] Rosser J. B. Extensions of some theorems of Goedel and Church.
Journal of Symbolic Logic vol 1 (1936) pp 87-91
- [54] Wikipedia Ackermann Function
https://en.wikipedia.org/wiki/Ackermann_function.
- [55] Ackermann W. 1928, "Zum Hilbertschen Aufbau der reellen Zahlen"
Mathematische Annalen 99. pp 118-133

- [56] Wikipedia Sudan function
https://en.wikipedia.org/wiki/Sudan_function
- [57] Sudan G. 1927 Sur le nombre transfini ω^{ω^n} Bulletin Mathematique
de la societe Roumaine de Sciences 30, pp 11-30

- [58] Maria Hameen Attila "Nominalistic Ordinals. Recursion on Higher
types, and Finitism" Bulletin of Symbolic
Logic 25 (1) 2019, pp 101-124.

- [59] Hinman P. 2016 Recursion theoretic Hierarchies , Cambridge
University press.
- [60] Wikipedia Arithmetical Hierarchies
https://en.wikipedia.org/wiki/Arithmetical_hierarchy#:~:text=In%20mathematical%20logic%2C%20the%20arithmetical,a%20classification%20is%20called%20arithmetical.

- [61] McMaster University Colloquium. April 2005 "The complexity of
Diophantine equations <https://www.math.mcgill.ca/~darmon/slides/mcmaster-05/slides.pdf>

- [62] Martin Davis 1958 "Computability and unsolvability" Mc Graw-Hill
http://ndl.ethernet.edu.et/bitstream/123456789/48378/1/Martin%20Davis_1995.pdf

- [63] Wikipedia Löwenheim–Skolem theorem
https://en.wikipedia.org/wiki/L%C3%B6wenheim%E2%80%93Skolem_theorem
- [64] Wikipedia Skolem paradox https://en.wikipedia.org/wiki/Skolem%27s_paradox

- [65] Fibonacci sequence https://en.wikipedia.org/wiki/Fibonacci_sequence
- [66] Linear recursive sequences with constant coefficients
https://en.wikipedia.org/wiki/Linear_recurrence_with_constant_coefficients
- [67] Constant linear recursive sequence https://en.wikipedia.org/wiki/Constant-recursive_sequence
- [68] Lucas numbers https://en.wikipedia.org/wiki/Lucas_number
- [69] Gessel, Ira (October 1972), "[Fibonacci is a Square](#)" (PDF), [The Fibonacci Quarterly](#), **10** (4): 417–19, retrieved April 11, 2012
- [70] Strong version of the Rice theorem .
<https://www.youtube.com/watch?v=oHX2hZInXc0>
- [71] G. Grentzen Consistency of Peano Arithmetic
https://en.wikipedia.org/wiki/Gerhard_Gentzen
- [72] Hilbert D,-Ackermann W. “Principles of Mathematical Logic” Chelsea publishing company 1938.
- [73] Whitehead A. - Russell B. Principia Mathematica Vol 1,2,3 Merchant Books 2009.
- [74] Nagel E.-Newman J. “Goedel’s proof” New York University Press 2001.
- [75] Goedel’s incompleteness theorems. Wikipedia
https://en.wikipedia.org/wiki/G%C3%B6del%27s_incompleteness_theorems
- [76] Matiyasevitch Y. V. Hilbert’s Tenth Problem MIT press 1993.
- [77] M. Davis, Y. Matiyasevitch J. Robinson . 1976. Hilbert’s tenth problem Proceedings of Symposia in Pure mathematics pp 323-378 AMS.
- [78] Matiyasevitch Y. 1976 (in Russian) A new proof of the theorem on exponential Diophantine representation of enumerable sets. Journal of Soviet mathematics 14 (5) pp 1475-1486 .1980.
- [79] Kyritsis K. 2024_Deterministic self-programmable Turing machines and the P vs NP problem (presentation)

https://www.researchgate.net/publication/377111073_DETERMISITIC_SELF-PROGRAMMABLE_TURING_MACHINES_AND_THE_P_vs_NP_PROBLEM

[80] Davis M. Putnam H. Robinson J. 1961 “The decision Problem for Exponential Diophantine Phancements” *Annals of mathematics* Vol 74, No 3 pp 425-436.

[81] Matiyasevitch Y.-Robinson J. 1975 “Reduction of an arbitrary Diophantine equation to one in 13 unknowns.” *Acta Arithmetica* XXVII 1975 pp 521-553.

[82] Deductive explosion (in contradictory systems) Wikipedia

https://en.wikipedia.org/wiki/Principle_of_explosion

[83] Kleene S. “The work of Goedel” *The Journal of Symbolic Logic* , Dec 1976 , Vol 41 , No 4 (Dev 1976) pp 761-778

<https://www.jstor.org/stable/2272394>

[84] Kleene S. “General Recursive functions of natural numbers” *A.M.S.* September 1935

[85] Kleene S. Recursive predicates and Quantifiers *Trans. A.M.S.* September 11, 1940 . vol 53 pp-41-73.

[87] Kleene S. Hierarchies of Number-Theoretic Predicates. Presented in Chicago Math. Society April 30, 1954 *Bulletin of the American Mathematical Society*, 1955 Bd. 61 , S. 193–213

[88] Wikipedia . Richard’s paradox ,

https://en.wikipedia.org/wiki/Richard%27s_paradox

[89] Eilenberg, Samuel; Mac Lane, Saunders (1945). "[General theory of natural equivalences](#)" (PDF). *Transactions of the American Mathematical Society*. **58**: 247. doi:[10.1090/S0002-9947-1945-0013131-6](https://doi.org/10.1090/S0002-9947-1945-0013131-6). ISSN 0002-9947. Archived (PDF) from the original on 2022-10-10.

[90] Schalk, A.; Simmons, H. (2005). [An introduction to Category Theory in four easy movements](#) (PDF). Archived from [the original](#) (PDF) on 2017-03-21. Retrieved 2007-12-03. Notes for a course offered as part of the MSc. in [Mathematical Logic](#), [Manchester University](#).

- [91] Turi, Daniele (1996–2001). ["Category Theory Lecture Notes"](#) (PDF). Retrieved 11 December 2009. Based on [Mac Lane 1998](#).
- [100] [Mac Lane, Saunders](#) (1998). [Categories for the Working Mathematician](#). Graduate Texts in Mathematics. Vol. 5 (2nd ed.). Springer-Verlag. [ISBN 978-0-387-98403-2](#). [MR 1712872](#).
- [101] Otto Martin 2015. Finite model theory.
<https://www2.mathematik.tu-darmstadt.de/~otto/LEHRE/FMT16.pdf>
- [102] Fix-point logic Wikipedia
https://en.wikipedia.org/wiki/Fixed-point_logic
- [103] Fagin's theorem Wikipedia
https://en.wikipedia.org/wiki/Fagin%27s_theorem
- [104] Descriptive complexity , Wikipedia
https://en.wikipedia.org/wiki/Descriptive_complexity_theory#First-order_least_fixed-point_logic
- [105] Robert Stoll. Sets, Logic, and Axiomatic Theories. Editor W.H. Freeman and Company. 1961.
- [106] Free Boolean algebra , Wikipedia
https://en.wikipedia.org/wiki/Free_Boolean_algebra
- [107] Lindenbaum-Tarski algebra, Wikipedia
https://en.wikipedia.org/wiki/Lindenbaum%E2%80%93Tarski_algebra
- [108] Stones representation theorem for Boolean algebras, Wikipedia
https://en.wikipedia.org/wiki/Stone%27s_representation_theorem_for_Boolean_algebras

PART B

CHAPTER 4

THE SOLUTION OF THE MILLENNIUM PROBLEM ABOUT THE NAVIER STOKES EQUATIONS

Prologue.

The standard formulation of the 4th Clay Millennium problem can be found in the site of the Clay Mathematical Institute here:

<http://www.claymath.org/millenniumproblems/navier%E2%80%93stokes-equation> and here <http://www.claymath.org/sites/default/files/navierstokes.pdf>

Roughly speaking it asks if in classical 3 dimensional incompressible fluids , (governed by the Navier-Stokes equations) with finite initial energy and smooth initial conditions (with pressures and velocities falling to zero faster than all polynomial powers as we go to infinite distances away or in short smooth Schwartz initial conditions) the flow will continuous forever smooth or would there be a finite time, where velocities and pressures will blow-up to infinite and smoothness will break? The standard formulation is both with periodic initial conditions or not periodic.

Most of the mathematicians were expecting that, since it has been proved that there is no blow-up in 2-dimensions, this should hold in 3 dimensions too. But as more than half a century has passed with not being able to prove it many researchers started believing that because of the vortex stretching which is possible only in 3-dimensions and not in 2-dimensions a blow-up might exist.

Because it was easier to do at the beginning, I spent about half a year discovering more than a dozen of explicitly formulated cases of axial symmetric flows that lead to blow-up in finite time. Nevertheless, for all of them, it was necessary that they start with infinite initial energy and the initial vorticities were unbounded.

So I went back to the more probable case that no Blow-up can occur in finite time.

My heuristic analysis which took 1-2 years, with statistical mechanics and classical fluid dynamics in digital differential and integral calculus suggested to me that there should not exist in finite time a blow-up. The naïve and simple argument was that a blow up would give that at least one particle of the fluid (and in statistical mechanics or classical fluid dynamics in digital differential and integral calculus, finite many finite particles do exist) would exhibit infinite kinetic energy. Nevertheless, what is easy to prove in heuristic context is not at all easy to prove in the classical context of fluid dynamics where there are not finite many particles of finite and lower bounded size, but infinite many points with zero size.

In this strategy my interdisciplinary approach was an advantage. I did not consider as consistent for sciences that e.g. statistical mechanics would give that there is no-blow up in finite time, while classical fluid dynamics would prove that there is a blow-up in finite time.

The next table makes the comparisons in statistical mechanics and classical fluid dynamics

Table 0

COMPARISON AND MUTUAL SIGNIFICANCE OF DIFFERENT TYPES OF MATHEMATICAL MODELS FOR THE 4TH CLAY PROBLEM (NO EXTERNAL FORCE)	CONTINUOUS FLUID MECHANICS MODEL	STATISTICAL MECHANICS MODEL
SMOOTH SCHWARTZ INITIAL CONDITIONS	YES	POSSIBLE TO IMPOSE
FINITE INITIAL ENERGY	YES	YES
CONSERVATION OF THE PARTICLES	YES(NON-OBVIOUS FORMULATION)	YES (OBVIOUS FORMULATION)
LOCAL SMOOTH EVOLUTION IN A INITIAL FINITE TIME INTERVAL	YES	POSSIBLE TO DERIVE
EMERGENCE OF A BLOW-UP IN FINITE TIME	IMPOSSIBLE TO OCCUR	IMPOSSIBLE TO OCCUR

So as it was easy to prove in statistical mechanics that there is no blow-up in finite time, I thought , so as to increase our confidence for the correct side of the solution of the problem , to add hypotheses to the standard formulation of the 4th Clay Millennium problem that correspond to the conservation of particles during the flow, and which would lead to an accessible solution of this problem (that there is no Blow-up in finite) dew to finite initial energy and energy and particle conservation.

This of course was not the solution of the 4th Clay Millennium problem, and the solution finally is presented in the 2nd and last paper in this part B of this treatise in chapter 6.

So once my confidence was in strength that the correct solution is that there is no Blow-up in finite time, I started attacking the problem for a proof in the classical fluid dynamics only with the hypotheses of the standard formulation of the 4th Clay Millennium problem. In Chapter 6 we give a full solution of the 4th Clay Millennium problem without any extra hypothesis.

The next 1st paper in chapter 5 is an initial version (uploaded in) of the published paper in the Journal of Scientific Research and Studies Vol. 4(11), pp. 304-317, November, 2017 ISSN 2375-8791 Copyright © 2017

It solves the millennium problem after an additional hypotheses of particles conservation.

Then in the chapter 6 the millennium problem is solved. without any additional hypotheses. This solution has also been published at least here

Kyritsis, K. (2022) A Short and Simple Solution of the Millennium Problem about the Navier-Stokes Equations and Similarly for the Euler Equations. *Journal of Applied Mathematics and Physics*, **10**, 2538-2560. doi: 10.4236/jamp.2022.108172. <https://www.scirp.org/journal/paperinformation.aspx?paperid=119569>

What is that we do not understand with the Navier-Stokes equations? The need for more consciousness for interpretations. Why we chose the geometric calculus approach for the solution?

It has been written in the initial formulation of the problem, that our difficulty of solving this millennium problem shows that there are several things that we do not understand very well in the Navier-Stokes equations. In this paragraph we will investigate this issue. We will explain also why the rather elementary geometric calculus approach is better so as to solve the millennium problem, compared to more advanced functional analysis.

1) One primary point, known but often forgotten is the next. The Euler and the Navier-Stokes equations are the equations that are considered to govern the flow of fluids, and had been formulated long ago in mathematical physics before it was known that matter consists from atoms. So actually, they formulated the old **infinite divisible material fluids**. After L. Boltzmann and the discovery of material atoms, the truer model is that of statistical mechanics. We may consider that the two different types of matter, a) infinite divisible b) made from finite atoms, behave the same as far as flows in fluid dynamics, and certainly there are many common

properties but ultimately are mathematically and logically different. One example of the difference is that in the atomic structured material fluid model, the angular velocity of the spin e.g. of electrons, protons, neutrons which is about 1 terahertz (infrared range) can vary increase or decrease, independently from the vorticity, which only the part of the angular velocity which is “geared to the environmental” rotation of the fluid. In the classical Weierstrass calculus of infinite divisible material fluids (Euler and Navier-Stokes equations) this distinction does not exist and all the angular velocity of a point is due to the vorticity. In chapter 6 [21] Muriel, A 2000 a corresponding to the millennium problem in statistical mechanics has been solved in the direction of regularity. Similarly, in chapter 6 [6] Kyritsis, K. November 2017 a solution of the current millennium problem has been proved in the direction of regularity, but only if adding an additional hypothesis to the initial formulation, that of existence of finite atomic particles that are conserved during the flow. Strictly speaking a mathematical model of the material fluids and their flow which will have a high degree of exactness should take in to account that matter consists of atoms, (the electron range of magnitudes is of the order 10^{-15} meters) and this it should avoid utilizing concepts of continuity and smoothness that use $\varepsilon > 0$ $\delta > 0$ in their definition smaller than 10^{-15} meters. To address this difficulty of our current (Weierstrass) calculus the author developed the Democritus digital and finite decimal differential calculus (see chapter 6, [16] Kyritsis K. 2019b , [15] Kyritsis K. 2017 B , [14] Kyritsis K. 2022) In this finite calculus, we define concepts, of seemingly infinitesimal numbers (they are finite), seemingly infinite numbers (they are finite) and feasible finite numbers, so as to develop a differential and integral calculus up to decimal numbers with only a fixed finite number decimal (decimal density of level of precision). Different levels of precision give different definitions of continuity and smoothness. These multi-precision levels Democritus calculi is what an applied mathematician is doing when applying the Newton-Leibniz and Weierstrass calculus with the infinite (and infinitesimals). The Democritus calculus strictly speaking is not logically equivalent to the Newton-Leibniz calculus or to the Weierstrass calculus. E.g. classical Weisstrass calculus continuity corresponds in the Democritus calculus of being continuous not only to a single precision level but to all possible precision levels. Because in the Democritus calculus continuity and smoothness is only up to a precision level, the turbulence can be defined in a way that in Weierstrass calculus cannot be defined. In a turbulent flow, the flow in the Democritus calculus may be smooth relative to a precision level but non-smooth relative to a coarser precision level (or the opposite) in the Weierstrass calculus this is impossible. Furthermore, now when a computer scientist is experimenting with computers to discover if in a flow there will be a blow up or not in finite time, within the Democritus calculus and its Navier-Stokes equations he will have an absolute proof and criterion. If the

vorticity will become seemingly infinite (still finite) in a feasible finite time interval there is a blow up. If it becomes only feasible finite in any feasible finite time interval, there is no blow up. Of course blow-up in the Democritus calculus is not equivalent with a blow up in the Weisstrass calculus. Finally, with the Democritus calculus the applied mathematician acquires the subjective quality of congruence. In other words, what he thinks, says and writes is what he acts and applies. With the infinite in the ontology of calculus this is not possible and it is unavoidable the incongruence, because infinite cannot be acted in the applications in a material reality where all are finite.

2) It is known that when the calculus (which is used in modeling the fluids) was discovered by Newton and Leibniz, the original mathematical ontology was utilizing infinitesimals, smaller than any positive real numbers but not zero. Then later with Weierstrass calculus this ontology was abandoned, we restricted ourselves to the real numbers only, and we utilized limits and convergence. So when we take the law of force (momentum conservation) of Newton $F=m*\gamma$ on a solid finite particle and then take the limit by shrinking it to a point to derive the Euler and Navier-Stokes equations, we must not forget, that originally the limit was not to a point but to an infinitesimal solid body particle. **This is not the same!** In chapter 6, [21] Kyritsis K. 2022, I have restored with strict mathematics the original ontology of infinitesimals of Newton-Leibniz , utilizing algebra of intervals (or inverses of ordinal numbers as J. H Conway has also done with the surreal numbers see chapter 6 [1] J H. Conway and [5] K Kyritsis ordinal real numbers 1,2,3). Then we have a two-density calculus with two different linearly ordered fields, a) the real numbers b) a larger such field of Newton-Leibniz fluxions, with infinitesimal, finite and infinite numbers. The topologies of convergence of a solid finite particle by shrinking it to a point or to an infinitesimal solid particle are different! And this affects the issue of vorticity and angular velocity of infinitesimal particle. When I was a University student, and I was learning about the equations of Navier-Stokes, I was satisfied to see that the simple law of force (momentum conservation) of Newton $F=m*\gamma$ was converted to the Navier-Stokes equations, but I was shocked to realize, that the rest of the independent information about the motion of the solid finite particle, namely its rotational momentum, was not shanked to an angular velocity ω of the infinitesimal solid particle. So we see now that this is not reasonable in the Weisstrass calculus, which shrinks to a point, while it is possible in the older Newton-Leibniz calculus which shrinks to an infinitesimal solid body, and would lead to a different model of flows of fluids, with independent initial data of angular velocities, besides linear velocities and besides the derived from them vorticity.

3) In the current solution of chapter 6 of the millennium problem, we may observe a 20%-80% Pareto rule. In other words, more than 80% of the equations utilized as governing equations of the flow, are those derived from fundamental theorem of the calculus, (in the form of Stokes theorem, divergence theorem, green theorem, Helmholtz-kelvin theorem, fundamental theorem of calculus etc.) and less that 30% the PDE of the Navier-Stokes equations. So I might say that the main equations governing the phenomenon of flow is the machinery of exterior differential algebra (wedge product) differentiation (differential forms) etc. rather than simply PDE equations. For reasons of simplicity and because we are restricted here to only 3 spatial dimensions, we do not utilize the symbolism of the wedge products and differential forms, but only the Stokes theorem, divergence theorem etc.

4) These versions of the fundamental theorem of the calculus (Stokes theorem etc) lead to an extension of the law of momentum conservation of 3D fluid parts to a law of 1D line density (rotatory) momentum conservation (Theorem 4.1 of chapter 6) and law of 2D surface density (rotatory) momentum conservation (Theorem 4.2 of chapter 6). These laws are very valuable for infinite divisible fluids so valuable as the existence of finite atoms in the atomics structured fluids. Without these extra laws of momentum density conservation, we would have a hope to solve the millennium problem. As T. Tao had remarked, only an integral of 3D energy conservation and an integral of 3D momentum conservation is not adequate to derive that momentum point densities $\rho \cdot u$, or energy point densities $(1/2)\rho \cdot u^2$ will not blow up.

5) Besides the forgotten conservation law of finite particles, which unfortunately we cannot utilize in the case of infinite divisible fluids to solve the millennium problem, there are **two more forgotten laws of conservation or invariants**. The first of them is the obvious that during the flow, the physical measuring units dimensions (dimensional analysis) of the involved physical quantities (mass density, velocity, vorticity, momentum, energy, force point density, pressure, etc.) are conserved. It is not very wise to eliminate the physical magnitudes interpretation and their dimensional analysis when trying to solve the millennium problem, because the dimensional analysis is a very simple and powerful interlink of the involved quantities and leads with the physical interpretation, to a transcendental shortcut to symbolic calculations. **By eliminating the dimensional analysis we lose part of the map to reach our goal.**

6) The 2nd forgotten conservation law or invariant, is related to the viscosity (friction). Because we do know that at each point (pointwise), the viscosity is only subtracting kinetic energy, with an irreversible way, and converting it to thermal energy, (negative energy point density), and this is preserved in the flow, (it can

never convert thermal energy to macroscopic kinetic energy), we know that its sign does not change too it is a flow invariant , so the integrated 1D or 2D work density is always of the same sign (negative) and as sign an invariant of the flow. The **conservation or invariance of the sign of work density by the viscosity (friction)** is summarized in the lemma 3.1 below.

7) Finally we must not understate the elementary fact that the force densities F_p due to the pressures p , $F_p = -\nabla p$ are conservative , irrotational vector field, and they do not contribute to the increase or decrease of the rotational momentum and vorticity of the fluid during the flow. Because of this we get that the conserved 1D and 2D densities of momentum in chapter 6 Theorems 4.1 and 4.2 are only of the rotatory type.

8) Anyone who has spent time to try to prove existence of Blow up or regularity in the various physical quantities of the fluid like velocity, vorticity, acceleration, force density, momentum, angular momentum, energy etc. he will observe that in the arguments the regularity and uniform in time boundedness propagates easily from derivatives to lower order of differentiation, while the blowup propagates easily from the magnitudes to their derivatives. The converses are hard in proving. This is due to the usual properties of the calculus derivatives and integrals. The hard part of the proofs, must utilize forms of the fundamental theorem of the calculus like stokes theorem, divergence theorem etc.

9) Based on the above 8 remarks about what is not very well understood with Navier-Stokes equations I decided that **elementary geometric calculus should be the appropriate to solve the millennium problem**, and this I did indeed.

CHAPTER 5

ON THE SOLUTION OF THE 4TH MILLENNIUM PROBLEM. PROOF OF THE REGULARITY OF THE SOLUTIONS OF THE EULER AND NAVIER-STOKES EQUATIONS, BASED ON THE CONSERVATION OF PARTICLES.

Abstract

As more and more researchers tend to believe that with the hypotheses of the official formulation of the 4th Clay Millennium problem a blowup may occur, a new goal is set: to find the simplest and most physically natural enhancement of the hypotheses in the official formulation so that the regularity can be proved in the case of 3 dimensions too. The position of this paper is that the standard assumptions of the official formulation of the 4th Clay millennium problem, although they reflect, the finiteness and the conservation of momentum and energy and the smoothness of the incompressible physical flows, they do not reflect the conservation of particles as local structure. By formulating the later conservation and adding it to the hypotheses, we prove the regularity (global in time existence and smoothness) both for the Euler and the Navier-Stokes equations.

Key words: *Incompressible flows, regularity, Navier-Stokes equations, 4th Clay millennium problem*

Mathematical Subject Classification: 76A02

1. Introduction

This 1st paper is an initial version of the published paper in the Journal of Scientific Research and Studies Vol. 4(11), pp. 304-317, November, 2017 ISSN 2375-8791

The famous problem of the 4th Clay mathematical Institute as formulated in FEFERMAN C. L. 2006 , is considered a significant challenge to the science of mathematical physics of fluids, not only because it has withstand the efforts of the scientific community for decades to prove it (or types of converses to it) but also

because it is supposed to hide a significant missing perception about the nature of our mathematical formulations of the physical flows through the Euler and the Navier-Stokes equations.

When the 4th Clay Millennium problem was officially formulated the majority was hoping that the regularity was holding also in 3 dimensions as it had been proved to hold also in 2 dimensions. But as time passed more and more mathematicians started believing that a Blowup can occur with the hypotheses of the official formulation. Therefore, a new goal is set to find the simplest and most physically natural enhancement of the hypotheses in the official formulation so that the regularity can be proved in the case of 3 dimensions too. This is done by the current paper.

After 3 years of research, in the 4th Clay Millennium problem, the author came to believe that, what most of the mathematicians would want, (and seemingly including the official formulators of the problem too), in other words a proof of the regularity in 3 dimensions as well, cannot be given merely by the assumptions of the official formulation of the problem. In other words, a Blow-up may occur even with compact support smooth initial data with finite energy. But solving the 4th Clay Millennium problem, by designing such a case of Blow-up is I think not interesting from the physical point of view, as it is quite away from physical applications and a mathematical pathological curiosity. On the other hand, discovering what physical aspect of the flows is not captured by the mathematical hypotheses, is I believe a more significant contribution to the science of mathematical physics in this area. Although the mathematical assumptions of the official formulation reflect, the finiteness and the conservation of momentum and energy and the smoothness of the incompressible physical flows, they do not reflect the conservation of particles as local structure. By adding this physical aspect formulated simply in the context of continuous fluid mechanics, the expected result of regularity can be proved.

In statistical mechanical models of incompressible flow, we have the realistic advantage of finite many particles, e.g. like balls $B(x,r)$ with finite diameter r . These particles as they flow in time, **remain particles of the same nature and size** and the velocities and inside them remain approximately constant.

Because space and time dimensions in classical fluid dynamics goes in orders of smallness, smaller and at least as small as the real physical molecules, atoms and particles of the fluids, this might suggest imposing too, such conditions resembling uniform continuity conditions. In the case of continuous fluid dynamics models such natural conditions, emerging from the particle nature of material fluids, together with the energy conservation, the incompressibility and the momentum conservation, as

laws conserved in time, may derive the regularity of the local smooth solutions of the Euler and Navier-Stokes equations. **For every atom or material particle of a material fluid, we may assume around it a ball of fixed radius, called particle range depending on the size of the atom or particle, that covers the particle and a little bit of the electromagnetic, gravitational or quantum vacuum field around it, that their velocities and space-time accelerations are affected by the motion of the molecule or particle.** E.g. for the case water, we are speaking here for molecules of H_2O , that are estimated to have a diameter of 2.75 angstroms or $2r = 2.75 \cdot 10^{-10}$ meters, we may define as water molecule **particle range** the balls $B(r_0)$ of radius $r_0 = 3 \cdot 10^{-10}$ meters around the water molecule. As the fluid flows, especially in our case here of incompressible fluids, the shape and size of the molecules do not change much, neither there are significant differences of the velocities and space-time accelerations of parts of the molecule. Bounds δ_u δ_ω of such differences remain constant as the fluid flows. **We may call this effect as the principle of conservation of particles as a local structure.** This principle must be posed in equal setting as the energy conservation and incompressibility together with the Navier-Stokes or Euler equations. Of course, if the fluid is say of solar plasma matter, such a description would not apply. Nevertheless, then incompressibility is hardly a property of it. But if we are talking about incompressible fluids that the molecule is conserved as well as the atoms and do not change atomic number (as e.g. in fusion or fission) then this principle is physically valid. The principle of conservation of particles as a local structure, blocks the self-similarity effects of concentrating the energy and turbulence in very small areas and creating thus a Blow-up. It is the missing invariant in the discussion of many researchers about supercritical, critical and subcritical invariants in scale transformations of the solutions.

The exact definition of the conservation of particles as local structure is in DEFINITION 5.1 and it is as follows:

(Conservation of particles as local structure in a fluid)

*Let a smooth solution of the Euler or Navier-Stokes equations for incompressible fluids, that exists in the time interval $[0, T)$. We may assume initial data on all of R^3 or only on a connected compact support V_0 . For simplicity let us concentrate only on the latter simpler case. Let us denote by F the displacement transformation of the flow. Let us also denote by g the partial derivatives of 1st order in space and time, that is $|\partial_x^\alpha \partial_t^b u(x)|$, $|\alpha| = 1$, $|b| \leq 1$, and call then space-time accelerations. We say that there is **conservation of the particles in the interval $[0, T)$ in** a derivatives homogenous setting, as a local structure of the solution if and only if:*

There is a small radius r , and small constants $\delta_x, \delta_u, \delta_\omega, >0$ so that for all t in $[0, T)$ there is a finite cover C_t (in the case of initial data on R^3 , it is infinite cover, but finite on any compact subset) of V_t , from balls $B(r)$ of radius r , called **ranges of the particles**, such that:

1) For an x_1 and x_2 in a ball $B(r)$ of V_s , s in $[0, T)$, $\|F(x_1) - F(x_2)\| \leq r + \delta_x$ for all $t \geq s$ in $[0, T)$.

2) For an x_1 and x_2 in a ball $B(r)$ of V_s , s in $[0, T)$, $\|u(F(x_1)) - u(F(x_2))\| \leq \delta_u$ for all $t \geq s$ in $[0, T)$.

3) For an x_1 and x_2 in a ball $B(r)$ of V_s , s in $[0, T)$, $\|g(F(x_1)) - g(F(x_2))\| \leq \delta_\omega$ for all $t \geq s$ in $[0, T)$.

If we state the same conditions 1) 2) 3) for all times t in $[0, +\infty)$, then we say that we have the **strong version** of the conservation of particles as local structure.

We prove in paragraph 5 in PROPOSITION 5.2 that indeed adding the above conservation of particles as local structure in the hypotheses of the official formulation of the 4th Clay Millennium problem, we solve it, in the sense of proving the regularity (global in time smoothness) of the locally in time smooth solutions that are known to exist.

A short outline of the logical structure of the paper is the next.

1) The paragraph 3, contains the official formulation of the 4th Clay millennium problem as in FEFFERMAN C. L. 2006. The official formulation is any one of 4 different conjectures, that two of them, assert the existence of blow-up in the periodic and non-periodic case, and two of them the non-existence of blow-up, that is the global in time regularity in the periodic and non-periodic case. We concentrate on to prove the regularity in the non-periodic case or conjecture (A) with is described by equations 1-6 after adding the conservation of particles as a local structure. The paragraph 3 contains definitions, and more modern symbolism introduced by T, Tao in TAO T. 2013. The current paper follows the formal and mathematical austerity standards that the official formulation has set, together with the suggested by the official formulation relevant results in the literature like in the book MAJDA A.J-BERTOZZI A. L. 2002.

But we try also not to lose the intuition of the physical interpretation, as we are in the area of mathematical physics rather than pure mathematics.

The goal is that reader after reading a dozen of mathematical propositions and their proofs, he must be able at the end to have simple physical intuition, why the conjecture (A) of the 4th Clay millennium together with the conservation of particles in the hypotheses problem holds.

2) The paragraph 4 contains some known theorems and results, that are to be used in this paper, so that the reader is not searching them in the literature and can have a direct, at a glance, image of what holds and what is proved. The most important are a list of necessary and sufficient conditions of regularity (PROPOSITIONS 4.5-4.10) The same paragraph contains also some well-known and very relevant results that are not used directly but are there for a better understanding of the physics.

3) The paragraph 5 contains the main idea that the conservation of particles during the flow can be approximately formulated in the context of continuous fluid mechanics and that is the key missing concept of conservation that acts as subcritical invariant in other words blocks the self-similar concentration of energy and turbulence that would create a Blowup. With this new invariant we prove the regularity in the case of 3 dimensions: PROPOSITIONS 5.2.

4) The paragraph 6 contains the idea of defining **a measure of turbulence** in the context of deterministic mechanics based on the **total variation** of the component functions or norms (DEFINITION 6.1) It is also made the significant observation that the smoothness of the solutions of the Euler and Navier-Stokes equations is not a general type of smoothness but one that would deserve the name **homogeneous smoothness** (Remark 6.2) .

According to CONSTANTIN P. 2007 “...The blowup problem for the Euler equations is a major open problem of PDE, theory of far greater physical importance than the blow-up problem of the Navier-Stokes equation, which is of course known to non-specialists because of the Clay Millennium problem...”

Almost all of our proved propositions and in particular the regularity in paragraphs 4 , 5 and 6 (in particular PROPOSITION 4.11 and PROPOSITION 5.2) are stated not only for the Navier-Stokes but also for the Euler equations.

2. The ontology of the continuous fluid mechanics models versus the ontology of statistical mechanics models. The main physical idea of the proof of the regularity in 3 spatial dimensions.

All researchers discriminate between the physical reality with its natural physical ontology (e.g. atoms, fluids etc) from the mathematical ontology (e.g. sets, numbers, vector fields etc). If we do not do that much confusion will arise. The main difference of the physical reality ontology, from the mathematical reality ontology, is what the mathematician D. Hilbert had remarked in his writings about the infinite. He remarked that nowhere in the physical reality there is anything infinite, while the mathematical infinite, as formulated in a special axiom of the infinite in G. Cantor's

theory of sets, is simply a convenient phenomenological abstraction, at a time that the atomic theory of matter was not well established yet in the mathematical community. In the physical reality ontology, as best captured by statistical mechanics models, the problem of the global 3-dimensional regularity seems easier to solve. For example it is known (See PROPOSITION 4.9 and PROPOSITION 4.12 maximum Cauchy development, and it is referred also in the official formulation of the Clay millennium problem in C. L. FEFFERMAN 2006) that if the global 3D regularity does not hold then the velocities become unbounded or tend in absolute value to infinite as time gets close to the finite Blow-up time. Now we know that a fluid consists from a finite number of atoms and molecules, which also have finite mass and with a lower bound in their size. **If such a phenomenon (Blowup) would occur, it would mean that for at least one particle the kinetic energy, is increasing in an unbounded way.** But from the assumptions (see paragraph 3) the initial energy is finite, so this could never happen. We conclude that the fluid is 3D globally in time regular. Unfortunately, such an argument although valid in statistical mechanics models (see also MURIEL A 2000), is not valid in continuous fluid mechanics models, where there are not atoms or particles with lower bound of finite mass, but only points with zero dimension, and only mass density. We must notice also here that this argument is not likely to be successful if the fluid is compressible. In fact, it has been proved that a blow-up may occur even with smooth compact support initial data, in the case of compressible fluids. One of the reasons is that if there is not lower bound in the density of the fluid, then even without violating the momentum and energy conservation, a density converging to zero may lead to velocities of some points converging to infinite. Nevertheless, if we formulate in the context of continuous fluid mechanics the conservation of particles as a local structure (DEFINITION 5.1) then we can derive a similar argument (see proof of PROPOSITION 5.1) where **if a Blowup occurs in finite time, then the kinetic energy of a finite small ball (called in DEFINITION 5.1 particle-range) will become unbounded**, which is again impossible, due to the hypotheses of finite initial energy and energy conservation.

The next table compares the hypotheses and conclusions both in continuous fluid mechanics models and statistical mechanics models of the 4th Clay millennium problem in its official formulation together with the hypothesis of conservation of particles. It would be paradoxical that we would be able to prove the regularity in statistical mechanics and we would not be able to prove it in continuous fluid mechanics.

Table 1

COMPARISON AND MUTUAL SIGNIFICANCE OF DIFFERENT TYPES OF MATHEMATICAL MODELS FOR THE 4TH CLAY PROBLEM (NO EXTERNAL FORCE)	CONTINUOUS FLUID MECHANICS MODEL	STATISTICAL MECHANICS MODEL
SMOOTH SCHWARTZ INITIAL CONDITIONS	YES	POSSIBLE TO IMPOSE
FINITE INITIAL ENERGY	YES	YES
<u>CONSERVATION OF THE PARTICES</u>	<u>YES(NON-OBVIOUS FORMULATION)</u>	<u>YES (OBVIOUS FORMULATION)</u>
LOCAL SMOOTH EVOLUTION IN A INITIAL FINITE TIME INTERVAL	YES	POSSIBLE TO DERIVE
EMERGENCE OF A BLOW-UP IN FINITE TIME	IMPOSSIBLE TO OCCUR	IMPOSSIBLE TO OCCUR

3. The official formulation of the Clay Mathematical Institute 4th Clay millennium conjecture of 3D regularity and some definitions.

In this paragraph we highlight the basic parts of the official formulation of the 4th Clay millennium problem, together with some more modern, since 2006, symbolism, by relevant researchers, like T. Tao.

In this paper I consider the conjecture (A) of C. L. FEFFERMAN 2006 official formulation of the 4th Clay millennium problem, which I identify throughout the paper as the 4th Clay millennium problem.

The Navier-Stokes equations are given by (by $\mathbb{R}$ we denote the field of the real numbers, $\nu > 0$ is the viscosity coefficient)

$$\frac{\partial}{\partial t} u_i + \sum_{j=1}^n u_j \frac{\partial u_i}{\partial x_j} = - \frac{\partial p}{\partial x_i} + \nu \Delta u_i \quad (x \in \mathbb{R}^3, t \geq 0, n=3) \quad (\text{eq.1})$$

$$\text{div} u = \sum_{i=1}^n \frac{\partial u_i}{\partial x_i} = 0 \quad (x \in \mathbb{R}^3, t \geq 0, n=3) \quad (\text{eq.2})$$

with initial conditions $u(x,0)=u^0(x)$ $x \in \mathbf{R}^3$ and $u_0(x)$ C^∞ divergence-free vector field on $\mathbf{R}^3$ (eq.3)

$\Delta = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2}$ is the Laplacian operator. The Euler equations are when $v=0$

For physically meaningful solutions we want to make sure that $u^0(x)$ does not grow large as $|x| \rightarrow \infty$. This is set by defining $u^0(x)$ and called in this paper **Schwartz initial conditions**, in other words

$$|\partial_x^\alpha u^0(x)| \leq C_{\alpha,K} (1+|x|)^{-K} \text{ on } \mathbf{R}^3 \text{ for any } \alpha \text{ and } K \quad (\text{eq.4})$$

(Schwartz used such functions to define the space of Schwartz distributions)

We accept as physical meaningful solutions only if it satisfies

$$p, u \in C^\infty(\mathbf{R}^3 \times [0, \infty)) \quad (\text{eq.5})$$

and

$$\int_{\mathbf{R}^3} |u(x,t)|^2 dx < C \text{ for all } t \geq 0 \text{ (Bounded or finite energy)} \quad (\text{eq.6})$$

The conjecture (A) of the Clay Millennium problem (case of no external force, but homogeneous and regular velocities) claims that for the Navier-Stokes equations, $v>0$, $n=3$, with divergence free, Schwartz initial velocities, there are for all times $t>0$, smooth velocity field and pressure, that are solutions of the Navier-Stokes equations with bounded energy, **in other words satisfying the equations eq.1, eq.2, eq. 3, eq.4, eq.5 eq.6**. It is stated in the same formal formulation of the Clay millennium problem by C. L. Fefferman see C. L. FEFFERMAN 2006 (see page 2nd line 5 from below) that the conjecture (A) has been proved to hold locally. “..if the time interval $[0, \infty)$, is replaced by a small time interval $[0, T)$, with T depending on the initial data...”. In other words there is $\infty > T > 0$, such that there is continuous and smooth solution $u(x,t) \in C^\infty(\mathbf{R}^3 \times [0, T))$. In this paper, as it is standard almost everywhere, the term smooth refers to the space C^∞

Following TAO, T 2013, we define some specific terminology, about the hypotheses of the Clay millennium problem, that will be used in the next.

*We must notice that the definitions below can apply also to the case of inviscid flows, satisfying the **Euler** equations.*

DEFINITION 3.1 (Smooth solutions to the Navier-Stokes system). *A smooth set of data for the Navier-Stokes system up to time T is a triplet (u_0, f, T) , where $0 < T < \infty$ is a time, the initial velocity vector field $u_0 : \mathbf{R}^3 \rightarrow \mathbf{R}^3$ and the forcing term $f : [0,$*

$T] \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ are assumed to be smooth on $\mathbb{R}^3$ and $[0, T] \times \mathbb{R}^3$ respectively (thus, u_0 is infinitely differentiable in space, and f is infinitely differentiable in space time), and u_0 is furthermore required to be divergence-free:

$$\nabla \cdot u_0 = 0.$$

If $f = 0$, we say that the data is *homogeneous*.

In the proofs of the main conjecture, we will not consider any external force, thus the data will always be homogeneous. But we will state intermediate propositions with external forcing. Next, we are defining simple differentiability of the data by Sobolev spaces.

DEFINITION 3.2 We define the H^1 norm (or enstrophy norm) $H^1(u_0, f, T)$ of the data to be the quantity

$$H^1(u_0, f, T) := \|u_0\|_{H_X^1(\mathbb{R}^3)} + \|f\|_{L_t^\infty H_X^1(\mathbb{R}^3)} < \infty \text{ and say that } (u_0, f, T) \text{ is } H^1 \text{ if}$$

$$H^1(u_0, f, T) < \infty.$$

DEFINITION 3.3 We say that a *smooth set of data* (u_0, f, T) is *Schwartz* if, for all integers $\alpha, m, k \geq 0$, one has

$$\sup_{x \in \mathbb{R}^3} (1 + |x|)^k |\nabla_x^\alpha u_0(x)| < \infty \quad \text{and} \quad \sup_{(t,x) \in [0,T] \times \mathbb{R}^3} (1 + |x|)^k |\nabla_x^\alpha \partial_t^m f(x)| < \infty$$

Thus, for instance, the solution or initial data having Schwartz property implies having the H^1 property.

DEFINITION 3.4 A *smooth solution* to the Navier-Stokes system, or a *smooth solution* for short, is a quintuplet (u, p, u_0, f, T) , where (u_0, f, T) is a *smooth set of data*, and the velocity vector field $u : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ and pressure field $p : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}$ are smooth functions on $[0, T] \times \mathbb{R}^3$ that obey the Navier-Stokes equation (eq. 1) but with external forcing term f ,

$$\frac{\partial}{\partial t} u_i + \sum_{j=1}^n u_j \frac{\partial u_i}{\partial x_j} = -\frac{\partial p}{\partial x_i} + \nu \Delta u_i + f_i \quad (x \in \mathbb{R}^3, t \geq 0, n=3)$$

and also the incompressibility property (eq.2) on all of $[0, T] \times \mathbb{R}^3$, but also the initial condition $u(0, x) = u_0(x)$ for all $x \in \mathbb{R}^3$

DEFINITION 3.5 Similarly, we say that (u, p, u_0, f, T) is H^1 if the associated data (u_0, f, T) is H^1 , and in addition one has

$$\|u\|_{L_t^\infty H_X^1([0,T] \times \mathbb{R}^3)} + \|u\|_{L_t^2 H_X^2([0,T] \times \mathbb{R}^3)} < \infty$$

We say that the solution is *incomplete in $[0,T)$* , if it is defined only in $[0,t]$ for every $t < T$.

We use here the notation of *mixed norms* (as e.g. in TAO, T 2013). That is if $\|u\|_{H_x^k(\Omega)}$ is the classical Sobolev norm ,of smooth function of a spatial domain Ω , $u : \Omega \rightarrow R$, I is a time interval and $\|u\|_{L_t^p(I)}$ is the classical L^p -norm, then the mixed norm is defined by

$$\|u\|_{L_t^p H_x^k(I \times \Omega)} := \left(\int_I \|u(t)\|_{H_x^k(\Omega)}^p dt \right)^{1/p} \text{ and } \|u\|_{L_t^\infty H_x^k(I \times \Omega)} := \operatorname{ess\,sup}_{t \in I} \|u(t)\|_{H_x^k(\Omega)}$$

Similar instead of the Sobolev norm for other norms of function spaces.

We also denote by $C_x^k(\Omega)$, for any natural number $k \geq 0$, the space of all k times continuously differentiable functions $u : \Omega \rightarrow R$, with finite the next norm

$$\|u\|_{C_x^k(\Omega)} := \sum_{j=0}^k \|\nabla^j u\|_{L_x^\infty(\Omega)}$$

We use also the next notation for *hybrid norms*. Given two normed spaces X, Y on the same domain (in either space or time), we endow their intersection $X \cap Y$ with the norm

$$\|u\|_{X \cap Y} := \|u\|_X + \|u\|_Y .$$

In particular in the we will use the next notation for intersection functions spaces, and their hybrid norms.

$$X^k(I \times \Omega) := L_t^\infty H_x^k(I \times \Omega) \cap L_x^2 H_x^{k+1}(I \times \Omega) .$$

We also use the *big O notation*, in the standard way, that is $X = O(Y)$ means

$X \leq CY$ for some constant C . If the constant C depends on a parameter s , we denote it by C_s and we write $X = O_s(Y)$.

We denote the difference of two sets A, B by $A \setminus B$. And we denote Euclidean balls by $B(a, r) := \{x \in R^3 : |x - a| \leq r\}$, where $|x|$ is the Euclidean norm.

With the above terminology the target Clay millennium conjecture in this paper can be restated as the next proposition

The 4th Clay millennium problem (Conjecture A)

(Global regularity for homogeneous Schwartz data). *Let $(u_0, 0, T)$ be a homogeneous Schwartz set of data. Then there exists a smooth finite energy solution $(u, p, u_0, 0, T)$ with the indicated data (notice it is for any $T>0$, thus global in time)*

.

4. Some known or directly derivable, useful results that will be used.

In this paragraph I state, some known theorems and results, that are to be used in this paper, so that the reader is not searching them in the literature and can have a direct, at a glance, image of what holds and what is proved.

A review of this paragraph is as follows:

Propositions 4.1, 4.2 are mainly about the uniqueness and existence locally of smooth solutions of the Navier-Stokes and Euler equations with smooth Schwartz initial data. Proposition 4.3 are necessary or sufficient or necessary and sufficient conditions of regularity (global in time smoothness) for the Euler equations without viscosity. Equations 8-15 are forms of the energy conservation and finiteness of the energy loss in viscosity or energy dissipation. Equations 16-18 relate quantities for the conditions of regularity. Proposition 4.4 is the equivalence of smooth Schwartz initial data with smooth compact support initial data for the formulation of the 4th Clay millennium problem. Propositions 4.5-4.9 are necessary and sufficient conditions for regularity, either for the Euler or Navier-Stokes equations, while Propositions 4.10 is a necessary and sufficient condition of regularity for only the Navier-Stokes with non-zero viscosity.

In the next I want to use, the basic local existence and uniqueness of smooth solutions to the Navier-Stokes (and Euler) equations, that is usually referred also as the well posedness, as it corresponds to the existence and uniqueness of the physical reality causality of the flow. The theory of well-posedness for smooth solutions is summarized in an adequate form for this paper by the Theorem 5.4 in TAO, T. 2013.

I give first the definition of **mild solution** as in TAO, T. 2013 page 9. Mild solutions must satisfy a condition on the pressure given by the velocities. Solutions of smooth initial Schwartz data are always mild, but the concept of mild solutions is a generalization to apply for non-fast decaying in space initial data, as the Schwartz data, but for which data we may want also to have local existence and uniqueness of solutions.

DEFINITION 4.1

We define a H^1 mild solution (u, p, u_0, f, T) to be fields $u, f : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$,

$p : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}$, $u_0 : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, with $0 < T < \infty$, obeying the regularity hypotheses

$$u_0 \in H_x^1(\mathbb{R}^3)$$

$$f \in L_t^\infty H_x^1([0, T] \times \mathbb{R}^3)$$

$$u \in L_t^\infty H_x^1 \cap L_t^2 H_x^2([0, T] \times \mathbb{R}^3)$$

with the pressure p being given by (Poisson)

$$p = -\Delta^{-1} \partial_i \partial_j (u_i u_j) + \Delta^{-1} \nabla \cdot f \quad (\text{eq. 7})$$

(Here the summation conventions is used, to not write the Greek big Sigma).

which obey the incompressibility conditions (eq. 2), (eq. 3) and satisfy the integral form of the Navier-Stokes equations

$$u(t) = e^{t\Delta} u_0 + \int_0^t e^{(t-t')\Delta} (-(u \cdot \nabla) u - \nabla p + f)(t') dt'$$

with initial conditions $u(x, 0) = u^0(x)$.

We notice that the definition holds also for the in viscid flows, satisfying the Euler equations. The viscosity coefficient here has been normalized to $\nu=1$.

In reviewing the local well-posedness theory of H^1 mild solutions, the next can be said. The content of the theorem 5.4 in TAO, T. 2013 (that I also state here for the convenience of the reader and from which derive our PROPOSITION 4.2) is largely standard (and in many cases it has been improved by more powerful current well-posedness theory). I mention here for example the relevant research by PRODI G 1959 and SERRIN, J 1963, The local existence theory follows from the work of KATO, T. PONCE, G. 1988, the regularity of mild solutions follows from the work of LADYZHENSKAYA, O. A. 1967. There are now a number of advanced local well-posedness results at regularity, especially that of KOCH, H., TATARU, D. 2001.

There are many other papers and authors that have proved the local existence and uniqueness of smooth solutions with different methods. As it is referred in C. L. FEFFERMAN 2006 I refer to the reader to the MAJDA A.J-BERTOZZI A. L. 2002 page 104 Theorem 3.4,

I state here for the convenience of the reader the summarizing theorem 5.4 as in TAO T. 2013. I omit the part (v) of Lipchitz stability of the solutions from the statement of the theorem. I use the standard $O()$ notation here, $x=O(y)$ meaning $x \leq cy$ for some absolute constant c . If the constant c depends on a parameter k , we set it as index of $O_k()$.

It is important to remark here that the existence and uniqueness results locally in time (well-posedness), hold also not only for the case of viscous flows following the Navier-Stokes equations, but also for the case of inviscid flows under the Euler equations. There are many other papers and authors that have proved the local existence and uniqueness of smooth solutions both for the Navier-Stokes and the Euler equation with the same methodology, where the value of the viscosity coefficient $\nu=0$, can as well be included. I refer e.g. the reader to the MAJDA A.J-BERTOZZI A. L. 2002-page 104 Theorem 3.4, paragraph 3.2.3, and paragraph 4.1 page 138.

PROPOSITION 4.1 *(Local well-posedness in H^1). Let (u_0, f, T) be H^1 data.*

(i) (Strong solution) If (u, p, u_0, f, T) is an H^1 mild solution, then

$$u \in C_t^0 H_x^1([0, T] \times \mathbb{R}^3)$$

(ii) (Local existence and regularity) If

$$(\|u_0\|_{H_x^1(\mathbb{R}^3)} + \|f\|_{L_t^1 H_x^1(\mathbb{R}^3)})^4 T < c$$

for a sufficiently small absolute constant $c > 0$, then there exists

a H^1 mild solution (u, p, u_0, f, T) with the indicated data, with

$$\|u\|_{X^k([0, T] \times \mathbb{R}^3)} = O(\|u_0\|_{H_x^1(\mathbb{R}^3)} + \|f\|_{L_t^1 H_x^1(\mathbb{R}^3)})$$

and more generally

$$\|u\|_{X^k([0, T] \times \mathbb{R}^3)} = O_k(\|u_0\|_{H_x^k(\mathbb{R}^3)}, \|f\|_{L_t^1 H_x^1(\mathbb{R}^3)}, 1)$$

for each $k \geq 1$. In particular, one has local existence whenever

T is sufficiently small, depending on the norm $H^1(u_0, f, T)$.

(iii) (Uniqueness) There is at most one H^1 mild solution (u, p, u_0, f, T)

with the indicated data.

(iv) (Regularity) If (u, p, u_0, f, T) is a H^1 mild solution, and (u_0, f, T)

is (smooth) Schwartz data, then u and p is smooth solution; in fact, one has

$$\partial_t^j u, \partial_t^j p \in L_t^\infty H^k([0, T] \times \mathbb{R}^3) \text{ for all } j, K \geq 0.$$

For the proof of the above theorem, the reader is referred to the TAO, T. 2013 theorem 5.4, but also to the papers and books, of the above mentioned other authors.

Next I state the local existence and uniqueness of smooth solutions of the Navier-Stokes (and Euler) equations with smooth Schwartz initial conditions, that I will use in this paper, explicitly as a PROPOSITION 4.2 here.

PROPOSITION 4.2 Local existence and uniqueness of smooth solutions or smooth well posedness. *Let $u_0(x)$, $p_0(x)$ be smooth and Schwartz initial data at $t=0$ of the Navier-Stokes (or Euler) equations, then there is a finite time interval $[0, T]$ (in general depending on the above initial conditions) so that there is a unique smooth local in time solution of the Navier-Stokes (or Euler) equations*

$$u(x) , p(x) \in C^\infty(\mathbb{R}^3 \times [0, T])$$

Proof: We simply apply the PROPOSITION 4.1 above and in particular, from the part (ii) and the assumption in the PROPOSITION 4.2, that the initial data are smooth Schwartz , we get the local existence of H^1 mild solution $(u, p, u_0, 0, T)$. From the part (iv) we get that it is also a smooth solution. From the part (iii), we get that it is unique.

As an alternative we may apply the theorems in MAJDA A.J-BERTOZZI A. L. 2002-page 104 Theorem 3.4, paragraph 3.2.3, and paragraph 4.1 page 138, and get the local in time solution, then derive from the part (iv) of the PROPOSITION 4.1 above, that they are also in the classical sense smooth. QED.

Remark 4.1 We remark here that the property of smooth Schwartz initial data, is not in general conserved in later times than $t=0$, of the smooth solution in the Navier-Stokes equations, because it is a very strong fast decaying property at spatially infinity. But for lower rank derivatives of the velocities (and vorticity) we have the **(global and) local energy estimate**, and **(global and) local enstrophy estimate** theorems that reduce the decaying of the solutions at later times than $t=0$, at spatially infinite to the decaying of the initial data at spatially infinite. See e.g. TAO, T. 2013, Theorem 8.2 (Remark 8.7) and Theorem 10.1 (Remark 10.6).

Furthermore, in the same paper of formal formulation of the Clay millennium conjecture, L. FEFFERMAN 2006 (see page 3rd line 6 from above), it is stated that the 3D global regularity of such smooth solutions is controlled by the **bounded accumulation in finite time intervals** of the vorticity (Beale-Kato-Majda). I state this also explicitly for the convenience of the reader, for smooth solutions of the Navier-Stokes equations with smooth Schwartz initial conditions, as the

PROPOSITION 4.6 When we say here **bounded accumulation** e.g. of the deformations D , **on finite internals**, we mean in the sense e.g. of the proposition 5.1 page 171 in the book MAJDA A.J-BERTOZZI A. L. 2002, which is a definition designed to control the existence or not of finite blowup times. In other words, for any finite time interval

$[0, T]$, there is a constant M such that

$$\int_0^t |D|_{L^\infty}(s) ds \leq M$$

I state here for the convenience of the reader, a well-known proposition of equivalent necessary and sufficient conditions of existence globally in time of solutions of the Euler equations, as inviscid smooth flows. It is the proposition 5.1 in MAJDA A.J-BERTOZZI A. L. 2002 page 171.

The *stretching* is defined by

$S(x, t) =: D\xi \cdot \xi$ if $\xi \neq 0$ and $S(x, t) =: 0$ if $\xi = 0$ where $\xi =: \frac{\omega}{|\omega|}$, ω being the vorticity.

PROPOSITION 4.3 *Equivalent Physical Conditions for Potential Singular Solutions of the Euler equations. The following conditions are equivalent for smooth Schwartz initial data:*

(1) *The time interval, $[0, T^*)$ with $T^* < \infty$ is a maximal interval of smooth H^s existence of solutions for the 3D Euler equations.*

(2) *The vorticity ω accumulates so rapidly in time that*

$$\int_0^t |\omega|_{L^\infty}(s) ds \rightarrow +\infty \text{ as } t \text{ tends to } T^*$$

(3) *The deformation matrix D accumulates so rapidly in time that*

$$\int_0^t |D|_{L^\infty}(s) ds \rightarrow +\infty \text{ as } t \text{ tends to } T^*$$

(4) *The stretching factor $S(x, t)$ accumulates so rapidly in time that*

$$\int_0^t [\max_{x \in \mathbb{R}^3} S(x, s)] ds \rightarrow +\infty \text{ as } t \text{ tends to } T^*$$

The next theorem establishes the equivalence of smooth connected compact support initial data with the smooth Schwartz initial data, for the homogeneous version of the 4th Clay Millennium problem. It can be stated either for local in time smooth solutions or global in time smooth solutions. The advantage assuming connected compact support smooth initial data, is obvious, as this is preserved in time by smooth functions and also integrations are easier when done on compact connected sets.

Remark 4.2 Finite initial energy and energy conservation equations:

When we want to prove that the smoothness in the local in time solutions of the Euler or Navier-Stokes equations is conserved, and that they can be extended indefinitely in time, we usually apply a “reduction ad absurdum” argument: Let the maximum finite time T^* and interval $[0, T^*)$ so that the local solution can be extended smooth in it.. Then the time T^* will be a blow-up time, and if we manage to extend smoothly the solutions on $[0, T^*]$. Then there is no finite Blow-up time T^* and the solutions holds in $[0, +\infty)$. Below are listed necessary and sufficient conditions for this extension to be possible. Obviously not smoothness assumption can be made for the time T^* , as this is what must be proved. But we still can assume that at T^* the energy conservation and momentum conservation will hold even for a singularity at T^* , as these are universal laws of nature, and the integrals that calculate them, do not require smooth functions but only integrable functions, that may have points of discontinuity.

A very well known form of the energy conservation equation and accumulative energy dissipation is the next:

$$\frac{1}{2} \int_{R^3} \|u(x, T)\|^2 dx + \int_0^T \int_{R^3} \|\nabla u(x, t)\|^2 dx dt = \frac{1}{2} \int_{R^3} \|u(x, 0)\|^2 dx \quad (\text{eq. 8})$$

Where:

$$E(0) = \frac{1}{2} \int_{R^3} \|u(x, 0)\|^2 dx \quad (\text{eq. 9})$$

is the initial finite energy

$$E(T) = \frac{1}{2} \int_{R^3} \|u(x, T)\|^2 dx \quad (\text{eq. 10})$$

is the final finite energy

$$\text{and } \Delta E = \int_0^T \int_{R^3} \|\nabla u(x, t)\|^2 dx dt \quad (\text{eq. 11})$$

is the accumulative finite energy dissipation from time 0 to time T , because of viscosity in to internal heat of the fluid. For the Euler equations it is zero. Obviously

$$\Delta E \leq E(0) = E(T) \quad (\text{eq. 12})$$

The rate of energy dissipation is given by

$$\frac{dE}{dt}(t) = -\nu \int_{\mathbb{R}^3} \|\nabla u\|^2 dx < 0 \quad (\text{eq. 13})$$

(ν , is the viscosity coefficient. See e.g. MAJDA, A.J-BERTOZZI, A. L. 2002 Proposition 1.13, equation (1.80) pp. 28)

At this point we may discuss, that for the smooth local in time solutions of the Euler equations, in other words for flows without viscosity, it is paradoxical from the physical point of view to assume, that the total accumulative in time energy dissipation is zero while the time or space-point density of the energy dissipation (the former is the $\|\nabla u(x,t)\|_{L_\infty}^2$), is not zero! It is indeed from the physical meaningful point of view unnatural, as we cannot assume that there is a loss of energy from to viscosity at a point and a gain from “anti-viscosity” at another point making the total zero. Neither to assume that the time and point density of energy dissipation is non-zero or even infinite at a space point, at a time, or in general at a set of time and space points of measure zero and zero at all other points, which would still make the total accumulative energy dissipation zero. **The reason is of course that the absence of viscosity, occurs at every point and every time, and not only in an accumulative energy level.** If a physical researcher does not accept such inviscid solutions of the Euler equation as having physical meaning, then for all other solutions that have physical meaning and the $\|\nabla u(x,t)\|_{L_\infty}^2$ is zero (and come so from appropriate initial data), we may apply the PROPOSITION 4.7 below and **deduce directly, that the local in time smooth solutions of the Euler equations, with smooth Schwartz initial data, and finite initial energy, and zero time and space point energy dissipation density due to viscosity, are also regular (global in time smooth).** For such regular inviscid solutions, we may see from the inequality in (eq. 15) below, that the total L2-norm of the vorticity is not increasing by time. We capture this remark in PROPOSITION 4.11 below.

Remark 4.3 The next are 3 very useful inequalities for the unique local in time $[0,T]$, smooth solutions u of the Euler and Navier-Stokes equations with smooth Schwartz initial data and finite initial energy (they hold for more general conditions on initial data, but we will not use that):

By $\|\cdot\|_m$ we denote the Sobolev norm of order m . So, if $m=0$ it is essentially the L_2 -norm. By $\|\cdot\|_{L^\infty}$ we denote the supremum norm, u is the velocity, ω is the vorticity, and c_m, c are constants.

$$1) \|u(x, T)\|_m \leq \|u(x, 0)\|_m \exp\left(\int_0^T c_m \|\nabla(u(x, t))\|_{L^\infty} dt\right) \quad (\text{eq. 14})$$

(see e.g. MAJDA, A.J-BERTOZZI, A. L. 2002, proof of Theorem 3.6 pp117, equation (3.79))

$$2) \|\omega(x, t)\|_0 \leq \|\omega(x, 0)\|_0 \exp\left(c \int_0^t \|\nabla u(x, s)\|_{L^\infty} ds\right) \quad (\text{eq. 15})$$

(see e.g. MAJDA, A.J-BERTOZZI, A. L. 2002, proof of Theorem 3.6 pp117, equation (3.80))

$$3) \|\nabla u(x, t)\|_{L^\infty} \leq \|\nabla u(x, 0)\|_0 \exp\left(\int_0^t \|\omega(x, s)\|_{L^\infty} ds\right) \quad (\text{eq. 16})$$

(see e.g. MAJDA, A.J-BERTOZZI, A. L. 2002, proof of Theorem 3.6 pp118, last equation of the proof)

The next are a list of well known necessary and sufficient conditions, for regularity (global in time existence and smoothness) of the solutions of Euler and Navier-Stokes equations, under the standard assumption in the 4th Clay Millennium problem of smooth Schwartz initial data, that after theorem Proposition 4.4 above can be formulated equivalently with smooth compact connected support data. We denote by T^* be the maximum Blow-up time (if it exists) that the local solution $u(x, t)$ is smooth in $[0, T^*)$.

PROPOSITION 4.5 (Condition for regularity)

The local solution $u(x, t)$, t in $[0, T^)$ of the Euler or Navier-Stokes equations, with smooth Schwartz initial data, can be extended to $[0, T^*]$, where T^* is the maximal time that the local solution $u(x, t)$ is smooth in $[0, T^*)$, if and only if the **Sobolev norm** $\|u(x, t)\|_m$, $m \geq 3/2 + 2$, remains bounded, by the same bound in all of $[0, T^*)$, then, there is no maximal Blow-up time T^* , and the solution exists smooth in $[0, +\infty)$*

Remark 4.4 See e.g. . MAJDA, A.J-BERTOZZI, A. L. 2002, pp 115, line 10 from below)

PROPOSITION 4.6 (Condition for regularity. Beale-Kato-Majda)

The local solution $u(x,t)$, t in $[0, T^*)$ of the Euler or Navier-Stokes equations, with smooth compact connected support initial data, can be extended to $[0, T^*]$, where T^* is the maximal time that the local solution $u(x,t)$ is smooth in $[0, T^*)$, if and only if for the finite time interval $[0, T^*]$, there exist a bound $M > 0$, so that the **vorticity has bounded by M , accumulation** in $[0, T^*]$:

$$\int_0^{T^*} \|\omega(x,t)\|_{L^\infty} dt \leq M \quad (\text{eq17})$$

Then there is no maximal Blow-up time T^* , and the solution exists smooth in $[0, +\infty)$

Remark 4.5 See e.g. . MAJDA, A.J-BERTOZZI, A. L. 2002, pp 115, Theorem 3.6. Also page 171 theorem 5.1 for the case of inviscid flows. . See also LEMARIE-RIEUSSET P.G. 2002. Conversely if regularity holds, then in any interval from the smoothness in a compact connected set, the vorticity is supremum bounded. The above theorems in the book MAJDA A.J-BERTOZZI A. L. 2002 guarantee that the above conditions extent the local in time solution to global in time, that is to solutions (u, p, u_0, f, T) which is H^1 mild solution, **for any T** . Then applying the part (iv) of the PROPOSITION 4.1 above, we get that this solution is also smooth in the classical sense, for all $T > 0$, thus globally in time smooth.

PROPOSITION 4.7 (Condition for regularity)

The local solution $u(x,t)$, t in $[0, T^*)$ of the Euler or Navier-Stokes equations, with smooth compact connected support initial data, can be extended to $[0, T^*]$, where T^* is the maximal time that the local solution $u(x,t)$ is smooth in $[0, T^*)$, if and only if for the finite time interval $[0, T^*]$, there exist a bound $M > 0$, so that the **vorticity is bounded by M , supremum norm L^∞** in $[0, T^*]$:

$$\|\omega(x,t)\|_{L^\infty} \leq M \text{ for all } t \text{ in } [0, T^*) \quad (\text{eq. 18})$$

Then there is no maximal Blow-up time T^* , and the solution exists smooth in $[0, +\infty)$

Remark 4.6 Obviously if $\|\omega(x,t)\|_{L^\infty} \leq M$, then also the integral exists and is bounded: $\int_0^{T^*} \|\omega(x,t)\|_{L^\infty} dt \leq M_1$ and the previous proposition 4.6 applies. Conversely if regularity holds, then in any interval from smoothness in a compact connected set, the vorticity is supremum bounded.

1) PROPOSITION 4.8 (Condition for regularity)

The local solution $u(x,t)$, t in $[0,T^*)$ of the Euler or Navier-Stokes equations, with smooth compact connected support initial data, can be extended to $[0,T^*]$, where T^* is the maximal time that the local solution $u(x,t)$ is smooth in $[0,T^*)$, if and only if for the finite time interval $[0,T^*]$, there exist a bound $M>0$, so that the space accelerations are bounded by M , in the supremum norm L^∞ in $[0,T^*]$:

$$\|\nabla u(x,t)\|_{L^\infty} \leq M \text{ for all } t \text{ in } [0,T^*) \quad (\text{eq. 19})$$

Then there is no maximal Blow-up time T^* , and the solution exists smooth in $[0,+\infty)$

Remark 4.7 Direct from the inequality (eq.14) and the application of the proposition 4.5. Conversely if regularity holds, then in any finite time interval from smoothness, the accelerations are supremum bounded.

PROPOSITION 4.9 (FEFFERMAN C. L. 2006. Condition for regularity)

The local solution $u(x,t)$, t in $[0,T^*)$ of the Navier-Stokes equations with non-zero viscosity, and with smooth compact connected support initial data, can be extended to $[0,T^*]$, where T^* is the maximal time that the local solution $u(x,t)$ is smooth in $[0,T^*)$, if and only if

the velocities $\|u(x,t)\|$ do not get unbounded as $t \rightarrow T^*$.

Then there is no maximal Blow-up time T^* , and the solution exists smooth in $[0,+\infty)$.

Remark 4.8. This is mentioned in the Official formulation of the 4th Clay Millennium problem FEFFERMAN C. L. 2006 pp.2, line 1 from below: quote “...For the Navier-Stokes equations ($\nu>0$) , if there is a solution with a finite blowup time T , then the velocities $u_i(x,t)$, $1 \leq i \leq 3$ become unbounded near the blowup time.” The converse-negation of this is that if the velocities remain bounded near the T^* , then there is no Blowup at T^* and the solution is regular or global in time smooth. Conversely of course, if regularity holds, then in any finite time interval, because of the smoothness, the velocities, in a compact set are supremum bounded.

I did not find a dedicated such theorem in the books or papers that I studied, but since prof. C.L Fefferman , who wrote the official formulation of the 4th Clay Millennium problem, was careful to specify that is in the case of non-zero viscosity $\nu>0$, and not of the Euler equations as the other conditions, I assume that he is aware of a proof of it.

PROPOSITION 4.10. (Necessary condition for regularity)

Let us assume that the local solution $u(x,t)$, t in $[0,T^*)$ of the Navier-Stokes equations with non-zero viscosity, and with smooth compact connected support

initial data, can be extended to $[0, T^*]$, where T^* is the maximal time that the local solution $u(x, t)$ is smooth in $[0, T^*)$, in other words that are regular, then the trajectories-paths length $l(a, t)$ does not get unbounded as

$t \rightarrow T^*$.

Proof: Let us assume that the solutions is regular. Then also for all finite time intervals $[0, T]$, the velocities and the accelerations are bounded in the L_∞ , supremum norm, and this holds along all trajectory-paths too. Then also the length of the trajectories, as they are given by the formula

$$l(a_0, T) = \int_0^T \|u(x(a_0, t))\| dt \quad (\text{eq. 20})$$

are also bounded and finite (see e.g. APOSTOL T. 1974, theorem 6.6 p128 and theorem 6.17 p 135). Thus, if at a trajectory the lengths become unbounded as t goes to T^* , then there is a blow-up. QED.

PROPOSITION 4.11.(Physical meaningful inviscid solutions of the Euler equations are regular)

Let us consider the local solution $u(x, t)$, t in $[0, T^)$ of the Euler equations with zero viscosity, and with smooth compact connected support initial data. If we consider, because of zero-viscosity at every space point and at every time, as physical meaningful solutions those that also the time and space points energy dissipation density, due to viscosity, is zero or $\|\nabla u(x, t)\|_{L_\infty}^2 = 0$, then, they can be extended smooth to all times $[0, +\infty)$, in other words they are regular.*

Proof: Direct from the PROPOSITION 4.8. QED.

Remark 4.9.

Similar results about the local smooth solutions, hold also for the non-homogeneous case with external forcing which is nevertheless space-time smooth of bounded accumulation in finite time intervals. Thus an alternative formulation to see that the velocities and their gradient, or in other words up to their 1st derivatives and the external forcing also up to the 1st derivatives, control the global in time existence is the next proposition. See TAO. T. 2013 Corollary 5.8

PROPOSITION 4.12 (Maximum Cauchy development)

Let (u_0, f, T) be H^1 data. Then at least one of the following two statements hold:

1) There exists a mild H^1 solution (u, p, u_0, f, T) in $[0, T]$, with the given data.

2) There exists a blowup time $0 < T^* < T$ and an incomplete mild H^1 solution

(u, p, u_0, f, T^*) up to time T^* in $[0, T^*)$, defined as complete on every $[0, t]$, $t < T^*$ which blows up in the enstrophy H^1 norm in the sense that

$$\lim_{t \rightarrow T^*, t < T^*} \|u(x, t)\|_{H_x^1(\mathbb{R}^3)} = +\infty$$

Remark 4.10 The term “almost smooth” is defined in TAO, T. 2013, before Conjecture 1.13. The only thing that almost smooth solutions lack when compared to smooth solutions is a limited amount of time differentiability at the starting time $t = 0$;

The term *normalized pressure*, refers to the symmetry of the Euler and Navier-Stokes equations to substitute the pressure, with another that differs at a constant in space but variable in time measurable function. In particular normalized pressure is one that satisfies the (eq. 7) except for a measurable at a constant in space but variable in time measurable function. It is proved in TAO, T. 2013, at Lemma 4.1, that the pressure is normalizable (exists a normalized pressure) in almost smooth finite energy solutions, for almost all times. The viscosity coefficient in these theorems of the above paper by TAO has been normalized to $\nu=1$.

5. Conservation of the particles as a local structure of fluids in the context of continuous fluid mechanics. Proof of the regularity for fluids with conservation of particles as a local structure, and the hypotheses of the official formulation of the 4th Clay millennium problem, for the Euler and Navier-Stokes equations.

Remark 5.1 (Physical interpretation of the definition 5.1) The smoothness of the particle-trajectory mapping (or displacement transformation of the points), the smoothness of the velocity field and vorticity field, is a condition that involves statements in the orders of micro scales of the fluid, larger, equal and also by far smaller than the size of material molecules, atoms and particles, from which it consists. This is something that we tend to forget in continuous mechanics, because continuous mechanics was formulated before the discovery of the existence of material atoms. On the other-hand it is traditional to involve the atoms and particles of the fluid, mainly in mathematical models of statistical mechanics. Nevertheless, we may formulate properties of material fluids in the context of continuous fluid mechanics, that reflect approximately properties and behavior in the flow of the material atoms. This is in particular the DEFINITION 5.1. **For every atom or**

material particle of a material fluid, we may assume around it a ball of fixed radius, called particle range depending on the size of the atom or particle, that covers the particle and a little bit of the electromagnetic, gravitational or quantum vacuum field around it, that their velocities and space-time accelerations are affected by the motion of the molecule or particle. E.g. for the case water, we are speaking here for molecules of H_2O , that are estimated to have a diameter of 2.75 angstroms or $2r = 2.75 \cdot 10^{-10}$ meters, we may define as water molecule **particle range** the balls $B(r_0)$ of radius $r_0 = 3 \cdot 10^{-10}$ meters around the water molecule. As the fluid flows, especially in our case here of incompressible fluids, the shape and size of the molecules do not change much, neither there are significant differences of the velocities and space-time accelerations of parts of the molecule. Bounds δ_u, δ_ω of such differences remain constant as the fluid flows. **We may call this effect as the principle of conservation of particles as a local structure.** This principle must be posed in equal setting as the energy conservation and incompressibility together with the Navier-Stokes or Euler equations. Of course, if the fluid is say of solar plasma matter, such a description would not apply. Nevertheless, then incompressibility is hardly a property of it. But if we are talking about incompressible fluids that the molecule is conserved as well as the atoms and do not change atomic number (as e.g. in fusion or fission) then this principle is physically valid. The principle of conservation of particles as a local structure, blocks the self-similarity effects of concentrating the energy and turbulence in very small areas and creating thus a Blow-up. It is the missing invariant in the discussion of many researchers about supercritical, critical and subcritical invariants in scale transformations of the solutions.

The next DEFINITION 5.1 formulates precisely mathematically this principle for the case of incompressible fluids.

DEFINITION 5.1. (Conservation of particles as local structure in a fluid)

Let a smooth solution of the Euler or Navier-Stokes equations for incompressible fluids, that exists in the time interval $[0, T)$. We may assume initial data on all of R^3 or only on a connected compact support V_0 . For simplicity let us concentrate only on the latter simpler case. Let us denote by F the displacement transformation of the flow. Let us also denote by g the partial derivatives of 1st order in space and time, that is $|\partial_x^a \partial_t^b u(x)|$, $|\alpha| = 1$, $|b| \leq 1$, and call then space-time accelerations. We say that there is conservation of the particles in the interval $[0, T)$ in a derivatives homogenous setting, as a local structure of the solution if and only if:

There is a small radius r , and small constants $\delta_x, \delta_u, \delta_\omega, > 0$ so that for all t in $[0, T)$ there is a finite cover C_t (in the case of initial data on R^3 it is infinite cover, but

finite on any compact subset) of V_t , from balls $B(r)$ of radius r , called **ranges of the particles**, such that:

4) For an x_1 and x_2 in a ball $B(r)$ of V_s , s in $[0, T)$, $\|F(x_1) - F(x_2)\| \leq r + \delta_x$ for all $t \geq s$ in $[0, T)$.

5) For an x_1 and x_2 in a ball $B(r)$ of V_s , s in $[0, T)$, $\|u(F(x_1)) - u(F(x_2))\| \leq \delta_u$ for all $t \geq s$ in $[0, T)$.

6) For an x_1 and x_2 in a ball $B(r)$ of V_s , s in $[0, T)$, $\|g(F(x_1)) - g(F(x_2))\| \leq \delta_\omega$ for all $t \geq s$ in $[0, T)$.

If we state the same conditions 1) 2) 3) for all times t in $[0, +\infty)$, then we say that we have the **strong version** of the conservation of particles as local structure.

PROPOSITION 5.1 (Velocities on trajectories in finite time intervals with finite total variation, and bounded in the supremum norm uniformly in time.)

Let $u_t : V(t) \rightarrow \mathbb{R}^3$ be smooth local in time in $[0, T^*)$, velocity fields solutions of the Navier-Stokes or Euler equations, with compact connected support $V(0)$ initial data, finite initial energy $E(0)$ and conservation of particles in $[0, T^*)$ as a local structure. The $[0, T^*)$ is the maximal interval that the solutions are smooth. Then for t in $[0, T^*)$ and x in $V(t)$, the velocities are uniformly in time bounded in the supremum norm by a bound M independent of time t .

$$\|u(x, t)\|_{L_\infty} = \sup_{x \in V(t)} \|u(x, t)\| \leq M \text{ for all } t \text{ in } [0, T^*).$$

Therefore, the velocities on the trajectory paths, in finite time intervals are of bounded variation and the trajectories in finite time interval, have finite length.

1st Proof (Only for the Navier-Stokes Equations): Let us assume, that the velocities are unbounded in the supremum norm, as t converges to T^* . Then there is a sequence of times t_n with t_n converging to time T^* , and sequence of corresponding points $x_n(t_n)$, for which the norms of the velocities $\|u(x_n(t_n), t_n)\|$ converge to infinite.

$$\lim_{n \rightarrow +\infty} \|u(x_n(t_n), t_n)\| = +\infty. \quad (\text{eq.21})$$

From the hypothesis of the conservation of particles as a local structure of the smooth solution in $[0, T^*)$, for every t_n There is a finite cover C_{t_n} of particle ranges, of V_{t_n} so that $x_n(t_n)$ belongs to one such ball or particle-range $B_n(r)$ and for any other point $y(t_n)$ of $B_n(r)$, it holds that $\|u(x_n(t_n), t_n) - u(y(t_n), t_n)\| \leq \delta_u$. Therefore

$$\|u(x_n(t_n), t_n)\| - \delta_u \leq \|u(y(t_n), t_n)\| \leq \|u(x_n(t_n), t_n)\| + \delta_u \quad (\text{eq.22})$$

for all times t_n in $[0, T^*)$.

By integrating spatially on the ball $B_n(r)$, and taking the limit as $n \rightarrow +\infty$ we deduce that

$$\lim_{n \rightarrow +\infty} \int_{B_n} \|u\| dx = +\infty$$

But this also means as we realize easily, that also

$$\lim_{n \rightarrow +\infty} \int_{B_n} \|u\|^2 dx = +\infty \quad (\text{eq. 23})$$

Which nevertheless means that the total kinetic energy of this small, but finite and of constant radius, ball, converges to infinite, as t_n converges to T^* . This is impossible by the finiteness of the initial energy, and the conservation of energy. Therefore the velocities are bounded uniformly, in the supremum norm, in the time interval $[0, T^*)$.

Therefore the velocities on the trajectory paths, are also bounded in the supremum norm, uniformly in the time interval $[0, T^*)$. But this means by PROPOSITION 4.9 that the local smooth solution is regular, and globally in time smooth, which from PROPOSITION 4.8 means that the Jacobian of the 1st order derivatives of the velocities are also bounded in the supremum norm uniformly in time bounded in $[0, T^*)$. Which in its turn gives that the velocities are of bounded variation on the trajectory paths (see e.g. APOSTOL T. 1974, theorem 6.6 p128 and theorem 6.17 p 135) and that the trajectories in have also finite length in $[0, T^*)$, because the trajectory length is given by the formula $l(a_0, T) = \int_0^T \|u(x(a_0, t))\| dt$.

QED.

2nd Proof (Both for the Euler and Navier-Stokes equations): Instead of utilizing the condition 2) of the definition 5.1, we may utilize the condition 3). And we start assuming that the Jacobian of the velocities is unbounded in the supremum norm (instead of the velocities), as time goes to the Blow-up time T^* . Similarly we conclude that the energy dissipation density at a time on balls that are particle-ranges goes to infinite, giving the same for the total accumulative in time energy dissipation (see (eq. 11), which again is impossible from the finiteness of the initial energy and energy conservation. Then by PROPOSITION 4.8 we conclude that the solution is regular, and thus also that the velocities are bounded in the supremum norm, in all finite time intervals. Again, we deduce in the same way, that the total variation of the velocities is finite in finite time intervals and so are the lengths of the trajectories too.

QED.

PROPOSITION 5.2 (Global regularity as in the 4th Clay Millennium problem).

Let the Navier-Stokes or Euler equations with smooth compact connected initial data, finite initial energy and conservation of particles as local structure. Then the unique local in time solutions are also regular (are smooth globally in time).

Proof: We apply the PROPSOITION 5.1 above and the necessary and sufficient condition for regularity in PROPSOITION 4.9 (which is only for the Navier-Stokes equations). Furthermore, we apply the part of the 2d proof of the PROPOSITION 5.1, that concludes regularity from PROPSOITION 4.8 which holds for both the Euler and Navier-Stokes equations.

QED.

6. Bounds of measures of the turbulence from length of the trajectory paths, and the total variation of the velocities, space acceleration and vorticity. The concept of homogeneous smoothness.

Remark 6.1 In the next we define a **measure of the turbulence** of the trajectories, of the velocities, of space-time accelerations and of the vorticity, through the **total variation** of the component functions in finite time intervals. This is in the context of deterministic fluid dynamics and not stochastic fluid dynamics. We remark that in the case of a blowup the measures of turbulence below will become infinite.

DEFINITION 6.1 (The variation measure of turbulence)

Let smooth local in time in $[0, T]$ solutions of the Euler or Navier-Stokes equations. The total length $L(P)$ of a trajectory path P , in the time interval $[0, T]$ is defined as **the variation measure of turbulence of the displacements** on the trajectory P , in $[0, T]$. The total variation $TV(\|u\|)$ of the norm of the velocity $\|u\|$ on the trajectory P in $[0, T]$ is defined as **the variation measure of turbulence of the velocity** on the trajectory P in $[0, T]$. The total variation $TV(g)$ of the space-accelerations g (as in DEFINITION 5.1) on the trajectory P in $[0, T]$ is defined as **the variation measure of turbulence of the space-time accelerations** on the trajectory P in $[0, T]$. The total variation $TV(\|\omega\|)$ of the norm of the vorticity $\|\omega\|$ on the trajectory P in $[0, T]$ is defined as **the variation measure of turbulence of the vorticity** on the trajectory P in $[0, T]$.

PROPOSITION 6.1 Conservation in time of the boundedness of the maximum turbulence, that depend only on the initial data and time lapsed.

Let the Euler or Navier-Stokes equations with smooth compact connected initial data finite initial energy and conservation of the particles as a local structure. Then for all times t , there are bounds $M_1(t)$, $M_2(t)$, $M_3(t)$, so that the maximum turbulence of the trajectory paths, of the velocities and of the space accelerations are bounded respectively by the above universal bounds, that depend only on the initial data and the time lapsed.

Proof: From the PROPOSITIONS 5.1, 5.2 we deduce that the local in time smooth solutions are smooth for all times as they are regular. Then in any time interval $[0, T]$, the solutions are smooth, and thus from the PROPOSITION 4.8, the space acceleration g , are bounded in $[0, T]$, thus also as smooth functions their total variation $TV(g)$ is finite, and bounded. (see e.g. APOSTOL T. 1974, theorem 6.6 p128 and theorem 6.17 p 135). From the PROPOSITION 4.7, the vorticity is smooth and bounded in $[0, T]$, thus also as smooth bounded functions its total variation $TV(\|\omega\|)$ is finite, and bounded on the trajectories. From the PROPOSITION 4.9, the velocity is smooth and bounded in $[0, T]$, thus also as smooth bounded functions its total variation $TV(\|u\|)$ is finite, and bounded on the trajectories. From the PROPOSITION 4.10, the motion on trajectories is smooth and bounded in $[0, T]$, thus also as smooth bounded functions its total variation which is the length of the trajectory path $L(P)$ is finite, and bounded in $[0, T]$. In the previous theorems the bounds that we may denote them here by $M_1(t)$, $M_2(t)$, $M_3(t)$, respectively as in the statement of the current theorem, depend on the initial data, and the time interval $[0, T]$. QED.

Remark 6.2. (Homogeneity of smoothness relative to a property P.) There are many researchers that they consider that the local smooth solutions of the Euler or Navier-Stokes equations with smooth Schwartz initial data and finite initial energy, (even without the hypothesis of conservation of particles as a local structure) are general smooth functions. But it is not so! They are special smooth functions with the remarkable property that there are some critical properties P_i that if such a property holds in the time interval $[0, T)$ for the coordinate partial space-derivatives of 0, 1, or 2 order, then this property holds also for the other two orders of derivatives. In other words if it holds for the 2 order then it holds for the orders 0, 1 in $[0, T)$. If it holds for the order 1, then it holds for the orders 0, 2 in $[0, T]$. If it holds for the order 0, then it holds also for the orders 1, 2 in $[0, T]$. This pattern e.g. can be observed for the property P_1 of uniform boundedness in the supremum norm, in the interval $[0, T^*)$ in the PROPOSITIONS 4.5-4.10. But one might to try to prove it also for a second property P_2 which is the **finiteness of the total variation** of the coordinates of the

partial derivatives, or even other properties P_3 like **local in time Lipchitz conditions**. This creates a strong bond or coherence among the derivatives and might be called homogeneous smoothness relative to a property P . We may also notice that the formulation of the conservation of particles as local structure is in such a way, that as a property, it shows the same pattern of homogeneity of smoothness relative to the property of uniform in time bounds P_4 , 1), 2), 3) in the DEFINITION 5.1. It seem to me though that even this strong type of smoothness is not enough to derive the regularity, unless the homogeneity of smoothness is relative to the property P_4 , in other words the conservation of particles as a local structure.

7. Epilogue

I believe that the main reasons of the failure so far in proving of the 3D global regularity of incompressible flows, with reasonably smooth initial conditions like smooth Schwartz initial data, and finite initial energy, is hidden in the difference of the physical reality ontology that is closer to the ontology of statistical mechanics models and the ontology of the mathematical models of continuous fluid dynamics.

Although energy and momentum conservation and finiteness of the initial energy are easy to formulate in both types of models, the conservation of particles as type and size is traditionally formulated only in the context of statistical mechanics. By succeeding in formulating approximately in the context of the ontology of continuous fluid mechanics the conservation of particles during the flow, as local structure, we result in being able to prove the regularity in the case of 3 dimensions which is what most mathematicians were hoping that it holds.

References

1. APOSTOL T. 1974 "Mathematical Analysis" 2nd edition Addison Wesley Publishing Company. 1974
2. BEALE, J. T., KATO, T., AND MAJDA, A 1984., "Remarks on the breakdown of smooth solutions for the 3-D Euler equations," *Commun. Math. Phys.* **94**, 61–66, 1984.
3. BOGOVSKII M. E. 1980, Solution of some vector analysis problems connected with operators div and grad, *Trudy Sem. S. L. Sobolev* 80 (1) (1980) 5-40.

4. CANNON, J.R., KNIGHTLY, G.H. 1970 : A note on the Cauchy problem for the Navier-Stokes equations. *SIAM J. Appl. Math.* 18, 641-644(1970)
5. CHORIN A. J. 1968 Numerical solutions of the Navier-Stokes Equations *Mathematics of Computation* oct. 1968, vol 22, no 104 pp 745-762
6. CONSTANTIN P. 1995, A few results and open problems regarding incompressible fluids, *Notices of the AMS*, 42 (1995), 658-663.
7. CONSTANTIN P. 2001 , Some open problems and research directions in the mathematical study of fluid dynamics, in *Mathematics Unlimited–2001 and Beyond*, Springer Verlag, Berlin, 2001, 353–360.
8. CONSTANTIN P. 2007 On the Euler equations of incompressible fluids. *Bulletin of the AMS* Vol 44, no 4, October 2007, pp 603-621.
9. GATSKI T. B. GROSCH C. E. ROSE M. E. 1989 The Numerical Solution of the Navier-Stokes equations for 3-Dimensional, Unsteady, Incompressible Flows by Compact Support Schemes *Journal of computation Physics* 82 , 298-329 (1989) .
10. GERMAIN P. 2008 Strong solutions and weak-strong uniqueness for the
11. nonhomogeneous Navier-Stokes system *Journal d'Analyse Mathématique* January 2008, Volume 105, Issue 1, pp 169-196.
12. GIGA Y. 1983 Weak and Strong Solutions of the Navier-Stokes Initial Value Problem *Publ RIMS, Kyoto Univ.* 19 (1983), 887-910
13. GIGA, Y., INUI, K., MATSUI, S. 1999 : On the Cauchy problem for the Navier-Stokes equations with nondecaying initial data. *Advances in fluid dynamics* 4, Quad. Mat., Dept. Math., Seconda Univ. Napoli, Caserta, 27-68(1999).
14. GIGA, Y., MAHALOV, A., NICOLAENKO, B. 2004 : The Cauchy problem for the Navier-Stokes equations with spatially almost periodic initial data. *Hokkaido University preprint series in mathematics* 683, (2004)
15. GIGA, Y., MATSUI, S., SAWADA, O. 2001 : Global existence of two-dimensional Navier-Stokes flow with nondecaying initial velocity. *J. Math. Fluid Mech.* 3, no.3, 302- 315(2001)
16. GIGA, Y., MIYAKAWA, T. 1985 : Solutions in L^r of the Navier-Stokes initial value problem. *Arch. Rational Mech. Anal.* 89, no.3, 267-281(1985)

17. GIGA Y. SAWADA, O. 2003 On regularizing-decay rate estimates for solutions to the Navier-Stokes initial value problem, *Nonlinear Analysis and Applications, To V. Lakshmikantham on his 80th Birthday vol. I (2003)*, 549-562.
18. FEFFERMAN C. L. 2006 Existence and smoothness of the Navier-Stokes equation, *Millennium Prize Problems, Clay Math. Inst., Cambridge, MA, 2006*, 57-67.
19. FRIEDLANDER S., DENIS SERRE 2002 *Handbook of Mathematical Fluid Dynamics Vol 1,2, Gulf Professional Publishing, 2002*.
20. KATO, J. 2003 : The uniqueness of nondecaying solutions for the Navier-Stokes equations. *Arch. Ration. Mech. Anal.* 169, no.2, 159-175(2003)
21. KATO, T. 1983 : Strong L^p -solutions of the Navier-Stokes equation in $\mathbb{R}^m$, with applications to weak solutions. *Math. Z.* 187, no.4, 471-480(1984)
22. KATO, T., FUJITA, H. 1962 : On the nonstationary Navier-Stokes system. *Rend. Sem. Mat. Univ. Padova* 32, 243-260(1962).
23. KATO, T. PONCE, G. 1988 Commutator estimates and the Euler and Navier-Stokes equations, *CPAM* 41 (1988), 891-907.
24. KOCH, H., TATARU, D. 2001 Well-posedness for the Navier-Stokes equations, *Adv. Math.* 157 (2001), no. 1, 22-35.
25. KYRITSIS, K. 2017 "On the solution of the 4th Clay Millennium problem about the Navier-Stokes equations. Equivalence of smooth compact support and smooth Schwartz initial conditions". *Proceedings of the 1st international conference on quantitative, social, biomedical and economic issues ICQSBEI 2017 pp 147-172*
26. KYRITSIS K. E. 2017b An Axiomatic System for a Physical or Digital but Continuous 3-Dimensional Euclidean Geometry, Without Infinite Many Points. *World Journal of Research and Review (WJRR) ISSN:2455-3956, Volume-5, Issue-4, October 2017 Pages 31-43*
27. KYRITSIS, K. E. 2019b. Outline of the introduction to the natural differential and integral calculus. *Journal of Regional & Socio-Economic* . Vol. 9, Issue 3, p.p.70-89.
28. LADYZHENSKAYA, O. A. 1967 Uniqueness and smoothness of generalized solutions of Navier-Stokes equations, *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* 5 (1967), 169-185; *English transl., Sem. Math. Steklov Math. Inst., Leningrad* 5 (1969), 60-67.
29. LADYZHENSKAYA, O. 1969, The Mathematical Theory of Viscous Incompressible Flows (2nd edition), *Gordon and Breach, New York, 1969*.

30. LEMARIE-RIEUSSET, P.G. 2002 Recent developments in the Navier-Stokes problem. *Chapman & All/CRC* 2002
31. MAEKAWA, Y. TERASAWA, Y. 2006 : The Navier-Stokes equations with initial data in uniformly local L_p spaces. *Differential and integral equations: an international journal for theory & applications (Athens, Ohio : Khayyam Publ. Co.)*, Vol. 19, No. 4 (2006), p. 369-400
32. MAJDA, A.J-BERTOZZI, A. L. 2002 Vorticity and incompressible flow *Cambridge University texts in applied mathematics* 2002
33. MURIEL, A 2000 An Exact Solution of the 3-D Navier-Stokes Equation Department of Electrical Engineering Columbia University And Department of Philosophy Harvard University. Also A. Muriel and M. Dresden, *Physica D* 101, 299, 1997.
34. POLYANIN, A. D. ZAITSEV, V.F., MOUSSIAUX 2002. *Handbook of First Order Partial Differential Equations*, London : Taylor&Francis , ISBN0-415-27267-x
35. POLYANIN, A. D 2002. *Handbook of Linear Partial Differential Equations for Engineers and Scientist* , Boca Raton : Chapman &Hall /CRC press, ISBN 1-58488-299-9
36. PRODI, G 1959 Un teorema di unicit_a per le equazioni di Navier-Stokes, *Ann. Mat. Pura Appl.* 48 (1959), 173-182.
37. RUDIN, W. 1976 “Principles of Mathematical Analysis ” 3rd edition McGraw-Hill Inc. 1976
38. SARRA, SCOTT 2003 “The method of characteristics with applications to Conservation Laws”, *Journal of Online mathematics and its Applications*.
39. SERRIN, J 1963 The initial value problem for the Navier-Stokes equations, *In Nonlinear Problems (Proc. Sympos., Madison, Wis., pages 69{98. Univ. of Wisconsin Press, Madison, Wis., 1963.*
40. SAWADA, O. 2003 : On time-local solvability of the Navier-Stokes equations in Besov spaces. *Adv. Differential Equations* 8 no.4, 385-412(2003)
41. SPIVAK, M. 1965 “Calculus on Manifolds” *The Benjamin/Cumming publishing company* 1965.
42. TAO, T. 2013 Localisation and compactness properties of the Navier-Stokes global regularity problem. *Analysis & PDE* 6 (2013), 25-107

43.TAO, T. 2014 Finite time blowup for an averaged three-dimensional Navier-Stokes equation. *Cornell University Library arxiv.org*

CHAPTER 6

A SHORT AND SIMPLE SOLUTION OF THE MILLENNIUM PROBLEM ABOUT THE NAVIER-STOKES EQUATIONS AND SIMILARLY FOR THE EULER EQUATIONS.

Konstantinos E. Kyritsis*

* Associate Prof. of University of Ioannina. Dept. Accounting-Finance Psathaki
Preveza 48100

*Corresponding Author: ckiritsi@uoi.gr, C_kyrisis@yahoo.com

Abstract

In this chapter 6 the millennium problem is solved. without any additional hypotheses. This solution has also been published at least here

Kyritsis, K. (2022) A Short and Simple Solution of the Millennium Problem about the Navier-Stokes Equations and Similarly for the Euler Equations. *Journal of Applied Mathematics and Physics*, **10**, 2538-2560. doi: 10.4236/jamp.2022.108172. <https://www.scirp.org/journal/paperinformation.aspx?paperid=119569>

It is presented a very short solution to the 4th Millennium problem about the Navier-Stokes equations. The solution proves that there cannot be a blow up in finite or infinite time, and the local in time smooth solutions can be extended for all times, thus regularity. This happily is proved not only for the Navier-Stokes equations but also for the inviscid case of the Euler equations both for the periodic or non-periodic formulation and without external forcing (homogeneous case). The proof is based on an appropriate modified extension in the viscous case of the well-known Helmholtz-Kelvin-Stokes theorem of invariance of the circulation of velocity in the Euler inviscid flows. This is essentially a 1D line density of (rotatory) momentum conservation. We discover a similar 2D surface density of (rotatory) momentum conservation. These conservations are indispensable, besides to the ordinary momentum conservation, to prove that there cannot be a blow-up in finite time, of the point vorticities, thus regularity.

Keywords: *Incompressible flows, regularity, blow-up, Navier-Stokes equations, Clay millennium problem*

Mathematical Subject Classification: 76A02

The current chapter 6 and the solution it present have been already published as article in a Journal

Kyritsis, K. (2022) A Short and Simple Solution of the Millennium Problem about the Navier-Stokes Equations and Similarly for the Euler Equations. *Journal of Applied Mathematics and Physics*, **10**, 2538-2560. doi: 10.4236/jamp.2022.108172. <https://www.scirp.org/journal/paperinformation.aspx?paperid=119569>

1. Introduction

The Clay millennium problem about the Navier-Stokes equations is one of the 7 famous problem of mathematics that the Clay Mathematical Institute has set a high monetary award for its solution. It is considered a difficult problem as it has resisted solving it for almost a whole century. The Navier-Stokes equations are the equations that are considered to govern the flow of fluids, and had been formulated long ago in mathematical physics before it was known that matter consists from atoms. So actually, they formulate the old infinite divisible material fluids. Although it is known that under its assumptions of the millennium problem the Navier-Stokes equations have a unique smooth and local in time solution, it was not known if this solution can be extended smoothly and globally for all times, which would be called the **regularity of the Navier-Stokes equations in 3 dimensions**. The corresponding case of regularity in 2 dimensions has long ago been proved to hold but the 3-dimensions had resisted proving it. Of course, the natural outcome would be that regularity holds also in 3-Dimensions. Many people felt that this difficulty hides our lack of understanding of the laws of 3-dimensional flow of the incompressible fluids.

In this paper is presented a very short solution to the Clay Millennium problem about the Navier-Stokes equations. The solution proves that there cannot be a blow up in finite or infinite time, and the local in time smooth solutions can be extended for all times, thus regularity. This happily is proved not only for the Navier-Stokes equations but also for the inviscid case of the Euler equations both for the periodic or non-periodic formulation and without external forcing (homogeneous case). But it is also proved that once the hypotheses of external forcing of the millennium problem allow for the existence of a unique smooth solution local in time, then the same result

of regularity (no blow up) holds also for this inhomogeneous case. I try to keep the length of this paper as short as possible so as to encourage reading it, and make the solution as easy to be understood as much as possible.

My first attempt to solve the millennium problem about the regularity of the Navier-Stokes equations problem was during the spring 2013 (uploaded at that time see [4] Kyritsis K. October 2013). Later attempts and solutions were published between 2017 and 2022 (see references [7], [8], [9], [11], [10], [17]). All of them in the same direction of regularity and no Blow-up. But some of the proofs contained errors, that in the current paper have been eliminated and the solution shortened. In the current paper we prove also something more compared to my previous publications that the regularity holds also for the Euler inviscid equations, with the same hypotheses of the millennium problem putting $\nu=0$, for the viscosity coefficient.

The author has also solved the 3rd Millennium problem P vs NP in computational complexity with 3 different successive solutions each one simpler than the previous.

(see references [8], [10], [12], [13], [18])

This millennium problem seems by the title of the articles as if solved by other authors like [2] Durmagambetov Asset et al 2015 also [20] Moschandreou. T. 2021, and [23] Ramm G. A. 2021.

Nevertheless, in my assessment they do not really solve it but eventually prove something else. In [9] Durmagambetov Asset et al 2015, the authors do not utilize the strict hypotheses of the formulation of the millennium problem, and the existence in general of blows-ups that they prove is a rather known fact. In [22] Ramm G. A. the strict hypotheses of the formulation of the millennium problem are indeed utilized but the solution essentially gives the existence of a smooth solution locally in time. Because the local in time $[0, t_1]$ smooth solution that he produces does depend on the initial data, we cannot repeat it in $[t_1, t_2]$, $[t_2, t_3]$ till infinite with certainty because we cannot claim that $t_1 = t_2 - t_1 = t_2 - t_3$ etc. Thus, there is no really a proof for no blow up and regularity. On the other hand, in [23] Ramm G. A. 2021 he proves that any solution of the Navier-Stokes equations, with the hypotheses of the millennium problem it will blow-up in finite time. There is obviously the counter example of potential (irrotational) flows that it is known that they do not blow up, and plenty many other specific counter examples in various publications of various authors, that do not blow up. Thus, his solution cannot be correct (although I could not find the error in his arguments). And finally, in [20] Moschandreou T. the solution as he writes in the conclusions is regular but he leaves open that fact that for a set of measure zero of the 3-space there might be a blow-up in finite time. Thus, it does not really prove either regularity or the existence with certainty of a blow up.

2. The formulation of the millennium problem and the 4 sub-problems (A) , (B), (C), (D)

In this paragraph we highlight the basic parts of the standard formulation of the 4th Clay millennium problem.

The **Navier-Stokes** equations are given by (by $\mathbf{R}$ we denote the field of the real numbers, $\nu > 0$ is the density normalized viscosity coefficient)

$$\frac{\partial}{\partial t} u_i + \sum_{j=1}^n u_j \frac{\partial u_i}{\partial x_j} = -\frac{\partial p}{\partial x_i} + \nu \Delta u_i \quad (\mathbf{x} \in \mathbf{R}^3, t \geq 0, n=3) \quad (\text{eq.2.1})$$

$$\text{div} u = \sum_{i=1}^n \frac{\partial u_i}{\partial x_i} = 0 \quad (\mathbf{x} \in \mathbf{R}^3, t \geq 0, n=3) \quad (\text{eq.2.2})$$

with initial conditions $u(\mathbf{x}, 0) = u^0(\mathbf{x})$ $\mathbf{x} \in \mathbf{R}^3$ and $u^0(\mathbf{x})$ C^∞ divergence-free vector field on $\mathbf{R}^3$ (eq.2.3)

If $\nu = 0$ then we are taking about the **Euler** equations and **inviscid** case.

$\Delta = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2}$ is the Laplacian operator.

The Euler equations are (eq2.1), (eq2.2), (eq2.3) when $\nu = 0$.

It is reminded to the reader, that in the equations of Navier-Stokes, as in (eq. 2.1) the density ρ , is constant, it is custom to normalized to 1 and omit it.

For physically meaningful solutions we want to make sure that $u^0(\mathbf{x})$ does not grow large as $|\mathbf{x}| \rightarrow \infty$. This is set by defining $u^0(\mathbf{x})$, and $f(\mathbf{x}, t)$ and called in this paper **Schwartz initial conditions**, in other words

$$|\partial_x^\alpha u^0(\mathbf{x})| \leq C_{\alpha, K} (1 + |\mathbf{x}|)^{-K} \text{ on } \mathbf{R}^3 \text{ for any } \alpha \text{ and } K \quad (\text{eq.2.4})$$

(Schwartz used such functions to define the space of Schwartz distributions)

Remark 2.1. It is important to realize that smooth Schwartz initial velocities after (eq 2.4) will give that the initial vorticity $\omega_0 = \text{curl}(u^0)$, in its supremum norm, is bounded over all 3-space.

$$|\partial_x^\alpha \partial_t^m f(\mathbf{x}, t)| \leq C_{\alpha, m, K} (1 + |\mathbf{x}| + t)^{-K} \text{ on } \mathbf{R}^3 \times [0, +\infty)$$

for any α, m, K (eq.2.5)

We accept as physical meaningful solutions only if it satisfies

$$p, u \in C^\infty(\mathbb{R}^3 \times [0, \infty)) \quad (\text{eq.2.6})$$

and

$$\int_{\mathbb{R}^3} |u(x, t)|^2 dx < C \text{ for all } t \geq 0$$

$$(\text{Bounded or finite energy}) \quad (\text{eq.2.7})$$

Remark 2.2 It is important to realize that smooth external force (densities) with the Schwartz property as in (eq.2.5), have not only a rule for upper bounded spatial partial derivatives but also the same rule for time upper bounded partial derivatives.

Remark 2.3 We must stress here that imposing smoothness of the coordinate functions of velocities and external forces of the initial $t=0$ data and later time t data in the Cartesian coordinates plus and Schwartz condition as in (eq 2.5) is not equivalent with imposing similar such smoothness of the coordinate functions and conditions in the cylindrical or spherical coordinates. We will give in the paragraph 4, remark 4.5 an example of a strange blowup, where at any time $t>0$ the coordinates of the velocities are smooth and bounded in all space as functions in the polar coordinates and still the vorticity has infinite singularity at zero.

Alternatively, to rule out problems at infinity, we may look for spatially periodic solutions of (2.1), (2.2), (2.3). Thus we assume that $u^0(x)$, and $f(x, t)$ satisfy

$$u^0(x+e_j) = u^0(x), f(x+e_j, t) = f(x, t), p(x+e_j, 0) = p(x, 0), \text{ for } 1 \leq j \leq 3 \quad (\text{eq.2.8})$$

(e_j is the j th unit vector in $\mathbb{R}^3$)

In place of (2.4) and (2.5), we assume that $u^0(x)$, is smooth and that

$$|\partial_x^\alpha \partial_t^m f(x, t)| \leq C_{\alpha, m, K} (1+t)^{-K} \text{ on } \mathbb{R}^3 \times [0, +\infty) \text{ for any } \alpha, m, K \quad (\text{eq.2.9})$$

We then accept a solution of (2.1), (2.2), (2.3) as physically reasonable if it satisfies

$$u(x+e_j, t) = u(x, t), p(x+e_j, t) = p(x, t), \text{ on } \mathbb{R}^3 \times [0, +\infty) \text{ for } 1 \leq j \leq 3 \quad (\text{eq.2.10})$$

$$\text{and } p, u \in C^\infty(\mathbb{R}^3 \times [0, \infty)) \quad (\text{eq.2.11})$$

In the next paragraphs we may also write u_0 instead of u^0 for the initial data velocity.

We denote Euclidean balls by $B(a, r) := \{x \in \mathbb{R}^3 : \|x - a\| \leq r\}$, where $\|x\|$ is the Euclidean norm.

The 4 sub-problems or conjectures of the millennium problem are the next:

(Conjecture A) Existence and smoothness of Navier-Stokes solution on $\mathbb{R}^3$.

Take $\nu > 0$ and $n=3$. Let $u_0(x)$ be any smooth, divergent-free vector field satisfying (4). Take $f(x,t)$ to be identically zero. Then there exist smooth functions $p(x,t)$, $u(x,t)$ on $\mathbb{R}^3 \times [0, +\infty)$ that satisfy (2.1), (2.2), (2.3) , (2.6) , (2.7).

(Conjecture B) Existence and smoothness of Navier-Stokes solution on $\mathbb{R}^3/\mathbb{Z}^3$.

Take $\nu > 0$ and $n=3$. Let $u_0(x)$ be any smooth, divergent-free vector field satisfying (8); we take $f(x,t)$ to be identically zero. Then there exist smooth functions $p(x,t)$, $u(x,t)$ on $\mathbb{R}^3 \times [0, +\infty)$ that satisfy (2.1), (2.2), (2.3) , (2.10) , (2.11).

(Conjecture C) Breakdown of Navier-Stokes solution on $\mathbb{R}^3$

Take $\nu > 0$ and $n=3$. Then there exist a smooth, divergent-free vector field $u_0(x)$ on $\mathbb{R}^3$ and a smooth $f(x,t)$ on $\mathbb{R}^3 \times [0, +\infty)$ satisfying (4), (5) for which there exist no smooth solution $(p(x,t), u(x,t))$ of (2.1), (2.2), (2.3) , (2.6) , (2.7) on $\mathbb{R}^3 \times [0, +\infty)$.

(Conjecture D) Breakdown of Navier-Stokes solution on $\mathbb{R}^3/\mathbb{Z}^3$

Take $\nu > 0$ and $n=3$. Then there exist a smooth, divergent-free vector field $u_0(x)$ on $\mathbb{R}^3$ and a smooth $f(x,t)$ on $\mathbb{R}^3 \times [0, +\infty)$ satisfying (2.8), (2.9) for which there exist no smooth solution $(p(x,t), u(x,t))$ of (2.1), (2.2), (2.3) , (2.10) , (2.11) on $\mathbb{R}^3 \times [0, +\infty)$.

In the next the $\|\cdot\|_m$ is the corresponding Sobolev spaces norm and. We denote by $V^m = \{u \text{ in } H^m(\mathbb{R}^n) \text{ and } \operatorname{div} u = 0\}$ where $H^m(\mathbb{R}^n)$ are the Sobolev spaces with the L^2 norm.

Remark 2.4. It is stated in the same formal formulation of the Clay millennium problem by C. L. Fefferman see [3] Fefferman C.L. 2006 (see page 2nd line 5 from below) that the conjecture (A) has been proved to holds locally. “..if the time interval $[0, \infty)$, is replaced by a small time interval $[0, T)$, with T depending on the initial data....”. In other words there is $\infty > T > 0$, such that there exists a unique and smooth solution $u(x,t) \in C^\infty(\mathbb{R}^3 \times [0, T))$. See also [19] A.J. Majda-A.L. Bertozzi, Theorem 3.4 pp 104. In this paper, as it is standard almost everywhere, the term smooth refers to the space C^∞

In these next the $\|\cdot\|_m$ is the corresponding Sobolev spaces norm and. We denote by $V^m = \{u \text{ in } H^m(\mathbb{R}^n) \text{ and } \operatorname{div} u = 0\}$ where $H^m(\mathbb{R}^n)$ are the Sobolev spaces with the L^2 norm.

We must mention that in A.J. Majda-A.L. Bertozzi [19], Theorem 3.4 pp 104, Local in Time existence of Solutions to the Euler and Navier-Stokes equations it is proved that indeed if the initial velocities belong to V^m $m \geq [3/2]+2$ there exist unique smooth solutions locally in time $[0, t]$. Here, in the formulation of the millennium problem the hypotheses of smooth with Schwartz condition initial velocities satisfies this condition therefore we have the existence and uniqueness of smooth solution locally in time, both in the non-periodic and the periodic setting without external forcing (homogeneous case).

The existence and uniqueness of a smooth solutions locally in time is stated in the formulation by C.L. Fefferman [3] for the homogeneous cases and conjectures (A), (B). When a smooth Schwartz condition external force is added (inhomogeneous case), it is natural to expect that also there should exist a local in time unique smooth solution. But this I did not find to be stated in the A.J. Majda-A.L. Bertozzi [19], **so I will avoid assuming it.**

We state here also two, very well-known criteria of no blow-up and regularity.

In this theorem the $\|\cdot\|_m$ is the corresponding Sobolev spaces norm and. We denote by $V^m = \{u \in H^m(\mathbb{R}^n) \text{ and } \operatorname{div} u = 0\}$ where $H^m(\mathbb{R}^n)$ are the Sobolev spaces with the L^2 norm.

Theorem 2.1 Velocities Sobolev norm sufficient condition of regularity. *Given an initial condition $u_0 \in V^m$ $m \geq [3/2]+2=3.5$ e.g. $m=4$, then for any viscosity $\nu \geq 0$. there exists a maximal time T^* (possibly infinite) of existence of a unique smooth solution $u \in C([0, T^*]; V^m) \cap C^1([0, T^*]; V^{m-2})$ to the Euler or Navier-Stokes equation. Moreover, if $T^* < +\infty$ then necessarily $\lim_{t \rightarrow T^*} \|u(\cdot, t)\|_m = +\infty$.*

Proof: See A.J. Majda-A.L. Bertozzi [19], Corollary 3.2 pp 112).

QED

Remark 2.5 Obviously this proposition covers the periodic case too.

Theorem 2.2 Supremum of vorticity sufficient condition of regularity

Let the initial velocity $u_0 \in V^m$ $m \geq [3/2]+2$, e.g. $m=4$, so that there exists a classical solution $u \in C^1([0, T]; C^2 \cap V^m)$ to the 3D Euler or Navier-Stokes equations. Then :

(i) *If for any $T > 0$ there is $M_1 > 0$ such that the vorticity $\omega = \operatorname{curl}(u)$ satisfies*

$$\int_0^T |\omega(., \tau)|_{L^\infty} d\tau \leq M_1$$

Then the solution u exists globally in time, $u \in C^1([0, +\infty] ; C^2 \cap V^m)$

(ii) If the maximal time T^* of the existence of the solution $u \in C^1([0, T] ; C^2 \cap V^m)$ is finite,

then necessarily the vorticity accumulates so rapidly that

$$\lim_{t \rightarrow T^*} \int_0^T |\omega(., \tau)|_{L^\infty} d\tau = +\infty \quad (\text{eq. 2.12})$$

Proof: See A.J. Majda-A.L. Bertozzi [19] ,Theorem 3.6 pp 115, L^∞ vorticity control of regularity.

QED.

Remark 2.6 Obviously this proposition covers the periodic case too.

3. What is that we do not understand with the Navier-Stokes equations? The need for more consciousness for interpretations. Why we chose the geometric calculus approach for the solution?

It has been written in the initial formulation of the problem, that our difficulty of solving this millennium problem shows that there are several things that we do not understand very well in the Navier-Stokes equations. In this paragraph we will investigate this issue. We will explain also why the rather elementary geometric calculus approach is better so as to solve the millennium problem, compared to more advanced functional analysis.

1) One primary point, known but often forgotten is the next. The Euler and the Navier-Stokes equations are the equations that are considered to govern the flow of fluids, and had been formulated long ago in mathematical physics before it was known that matter consists from atoms. So actually, they formulated the old **infinite divisible material fluids**. After L. Boltzmann and the discovery of material atoms, the truer model is that of statistical mechanics. We may consider that the two different types of matter, a) infinite divisible b) made from finite atoms, behave the same as far as flows in fluid dynamics, and certainly there are many common properties but ultimately are mathematically and logically different. One example of the difference is that in the atomic structured material fluid model, the angular velocity of the spin e.g. of electrons, protons, neutrons which is about 1 terahertz (infrared range) can vary increase or decrease, independently from the vorticity,

which only the part of the angular velocity which is “geared to the environmental” rotation of the fluid. In the classical Weierstrass calculus of infinite divisible material fluids (Euler and Navier-Stokes equations) this distinction does not exist and all the angular velocity of a point is due to the vorticity. In [21] Muriel, A 2000 a corresponding to the millennium problem in statistical mechanics has been solved in the direction of regularity. Similarly, in [6] Kyritsis, K. November 2017 a solution of the current millennium problem has been proved in the direction of regularity, but only if adding an additional hypothesis to the initial formulation, that of existence of finite atomic particles that are conserved during the flow. Strictly speaking a mathematical model of the material fluids and their flow which will have a high degree of exactness should take in to account that matter consists of atoms, (the electron range of magnitudes is of the order 10^{-15} meters) and this it should avoid utilizing concepts of continuity and smoothness that use $\varepsilon > 0$ $\delta > 0$ in their definition smaller than 10^{-15} meters. To address this difficulty of our current (Weierstrass) calculus the author developed the Democritus digital and finite decimal differential calculus (see [16] Kyritsis K. 2019b , [15] Kyritsis K. 2017 B , [14] Kyritsis K. 2022) In this finite calculus, we define concepts, of seemingly infinitesimal numbers (they are finite), seemingly infinite numbers (they are finite) and feasible finite numbers, so as to develop a differential and integral calculus up to decimal numbers with only a fixed finite number decimal (decimal density of level of precision). Different levels of precision give different definitions of continuity and smoothness. These multi-precision levels Democritus calculi is what an applied mathematician is doing when applying the Newton-Leibniz and Weierstrass calculus with the infinite (and infinitesimals). The Democritus calculus strictly speaking is not logically equivalent to the Newton-Leibniz calculus or to the Weierstrass calculus. E.g. classical Weierstrass calculus continuity corresponds in the Democritus calculus of being continuous not only to a single precision level but to all possible precision levels. Because in the Democritus calculus continuity and smoothness is only up to a precision level, the turbulence can be defined in a way that in Weierstrass calculus cannot be defined. In a turbulent flow, the flow in the Democritus calculus may be smooth relative to a precision level but non-smooth relative to a coarser precision level (or the opposite) in the Weierstrass calculus this is impossible. Furthermore, now when a computer scientist is experimenting with computers to discover if in a flow there will be a blow up or not in finite time, within the Democritus calculus and its Navier-Stokes equations he will have an absolute proof and criterion. If the vorticity will become seemingly infinite (still finite) in a feasible finite time interval there is a blow up. If it becomes only feasible finite in any feasible finite time interval, there is no blow up. Of course blow-up in the Democritus calculus is not equivalent with a blow up in the Weierstrass calculus. Finally, with the Democritus

calculus the applied mathematician acquires the subjective quality of congruence. In other words, what he thinks, says and writes is what he acts and applies. With the infinite in the ontology of calculus this is not possible and it is unavoidable the incongruence, because infinite cannot be acted in the applications in a material reality where all are finite.

2) It is known that when the calculus (which is used in modeling the fluids) was discovered by Newton and Leibniz, the original mathematical ontology was utilizing infinitesimals, smaller than any positive real numbers but not zero. Then later with Weierstrass calculus this ontology was abandoned, we restricted ourselves to the real numbers only, and we utilized limits and convergence. So when we take the law of force (momentum conservation) of Newton $F=m*\gamma$ on a solid finite particle and then take the limit by shrinking it to a point to derive the Euler and Navier-Stokes equations, we must not forget, that originally the limit was not to a point but to an infinitesimal solid body particle. **This is not the same!** In [30x3] Kyritsis K. 2022, I have restored with strict mathematics the original ontology of infinitesimals of Newton-Leibniz , utilizing algebra of intervals (or inverses of ordinal numbers as J. H Conway has also done with the surreal numbers see [1] J H. Conway and [5] K Kyritsis ordinal real numbers 1,2,3). Then we have a two-density calculus with two different linearly ordered fields, a) the real numbers b) a larger such field of Newton-Leibniz fluxions, with infinitesimal, finite and infinite numbers. The topologies of convergence of a solid finite particle by shrinking it to a point or to an infinitesimal solid particle are different! And this affects the issue of vorticity and angular velocity of infinitesimal particle. When I was a University student, and I was learning about the equations of Navier-Stokes, I was satisfied to see that the simple law of force (momentum conservation) of Newton $F=m*\gamma$ was converted to the Navier-Stokes equations, but I was shocked to realize, that the rest of the independent information about the motion of the solid finite particle, namely its rotational momentum, was not shanked to an angular velocity ω of the infinitesimal solid particle. So, we see now that this is not reasonable in the Weisstrass calculus, which shrinks to a point, while it is possible in the older Newton-Leibniz calculus which shrinks to an infinitesimal solid body, and would lead to a different model of flows of fluids, with independent initial data of angular velocities, besides linear velocities and besides the derived from them vorticity.

3) In the current solution of the millennium problem, we may observe a 20%-80% Pareto rule. In other words, more than 80% of the equations utilized as governing equations of the flow, are those derived from fundamental theorem of the calculus, (in the form of Stokes theorem, divergence theorem, green theorem, Helmholtz-kelvin theorem, fundamental theorem of calculus etc.) and less that 30% the PDE of

the Navier-Stokes equations. So I might say that the main equations governing the phenomenon of flow is the machinery of exterior differential algebra (wedge product) differentiation (differential forms) etc. rather than simply PDE equations. For reasons of simplicity and because we are restricted here to only 3 spatial dimensions, we do not utilize the symbolism of the wedge products and differential forms, but only the Stokes theorem, divergence theorem etc.

4) These versions of the fundamental theorem of the calculus (Stokes theorem etc) lead to an extension of the law of momentum conservation of 3D fluid parts to a law of 1D line density (rotatory) momentum conservation (Theorem 4.1) and law of 2D surface density (rotatory) momentum conservation (Theorem 4.2). These laws are very valuable for infinite divisible fluids so valuable as the existence of finite atoms in the atomics structured fluids. Without these extra laws of momentum density conservation, we would have a hope to solve the millennium problem. As T. Tao had remarked, only an integral of 3D energy conservation and an integral of 3D momentum conservation is not adequate to derive that momentum point densities $\rho \cdot u$, or energy point densities $(1/2)\rho \cdot u^2$ will not blow up.

5) Besides the forgotten conservation law of finite particles, which unfortunately we cannot utilize in the case of infinite divisible fluids to solve the millennium problem, there are **two more forgotten laws of conservation or invariants**. The first of them is the obvious that during the flow, the physical measuring units' dimensions (dimensional analysis) of the involved physical quantities (mass density, velocity, vorticity, momentum, energy, force point density, pressure, etc.) are conserved. It is not very wise to eliminate the physical magnitudes interpretation and their dimensional analysis when trying to solve the millennium problem, because the dimensional analysis is a very simple and powerful interlink of the involved quantities and leads with the physical interpretation, to a transcendental shortcut to symbolic calculations. **By eliminating the dimensional analysis we lose part of the map to reach our goal.**

6) The 2nd forgotten conservation law or invariant, is related to the viscosity (friction). Because we do know that at each point (pointwise), the viscosity is only subtracting kinetic energy, with an irreversible way, and converting it to thermal energy, (negative energy point density), and this is preserved in the flow, (it can never convert thermal energy to macroscopic kinetic energy), we know that its sign does not change too it is a flow invariant, so the integrated 1D or 2D work density is always of the same sign (negative) and as sign an invariant of the flow. **The conservation or invariance of the sign of work density by the viscosity (friction)** is summarized in the lemma 3.1 below.

7) Finally we must not understate the elementary fact that the force densities F_p due to the pressures p , $F_p = -\nabla p$ are conservative, irrotational vector field, and they do not contribute to the increase or decrease of the rotational momentum and vorticity of the fluid during the flow. Because of this we get that the conserved 1D and 2D densities of momentum in Theorems 4.1 and 4.2 are only of the rotatory type.

8) Anyone who has spent time to try to prove existence of Blow up or regularity in the various physical quantities of the fluid like velocity, vorticity, acceleration, force density, momentum, angular momentum, energy etc. he will observe that in the arguments the regularity and uniform in time boundedness propagates easily from derivatives to lower order of differentiation, while the blowup propagates easily from the magnitudes to their derivatives. The converses are hard in proving. This is due to the usual properties of the calculus derivatives and integrals. The hard part of the proofs, must utilize forms of the fundamental theorem of the calculus like Stokes theorem, divergence theorem etc.

9) Based on the above 8 remarks about what is not very well understood with Navier-Stokes equations I decided that **elementary geometric calculus should be the appropriate to solve the millennium problem**, and this I did indeed.

Lemma 3.1 The viscosity sign forgotten invariant.

If we integrate the force density of the viscosity, over a line (1D work density) or surface (2D work density) or a volume (work) its sign will remain the same during the flow.

Proof: Because we do know that pointwise, the viscosity is only subtracting kinetic energy, with an irreversible way, and converting it to thermal energy, (negative energy point density), and this is preserved in the flow, (it can never convert thermal energy to macroscopic kinetic energy), we deduce that its sign does not change too it is a flow invariant, so the integrated 1D or 2D work density is always of the same sign (negative) and as sign an invariant of the flow.

QED.

4. The Helmholtz-Kelvin-Stokes theorem in the case of viscous flows. New monotone semi-invariants of viscous flows with the interpretation of average rotational momentum axial 1-D line densities.

Here we apply the idea that the most valuable equations that govern the flow of the fluid are not literally the Navier-Stokes equations but the invariants or semi-invariant properties of the flow, derived from the abstract multi-dimensional fundamental

theorems of calculus, in the forms of divergence theorems, Stokes theorems, Greens theorems etc. Actually, this is the mechanism of wedge-products and abstract algebra of differential forms which is beyond classical partial differential equations. We do not utilize though definitions and symbolism of wedge-products and differential forms in his paper so as to keep it elementary and easy to read. The main discovery of this paragraph is the **Helmholtz-Kelvin-Stokes theorems 4.3 in the case of viscous flows and the resulting general no-blow-up theorem 4.4 for the viscous flows without external forcing**. A blow-up when it occurs, it will occur at least as blow-up of the vorticity, or of $\rho \bullet \omega$. If we discover average value invariants of the flow with physical units dimensions $\rho \bullet \omega$, that in the limit can give also the point value of the $\rho \bullet \omega$, **and that are invariants independent from the size of averaging**, it is reasonable that we can deduce conclusions, if the point densities can blow-up or not.

Theorem 4.1 The Helmholtz-Kelvin-Stokes theorem in the case of inviscid Euler equations flows without external force or homogeneous case. (A 1D line density of rotatory momentum, conservation law).

Let initial data in R^3 so that they guarantee the existence of a unique smooth solution to the Euler equation in a local time interval $[0,T]$. Then at any time $t \in [0,T]$ the circulation $\Gamma(c)$ of the velocities on a closed smooth loop is equal to the flux of the vorticity on smooth surface S with boundary the loop c , and is constant and preserved as both loop and surface flow with the fluid. In symbols ($\rho=1$ is the density of the incompressible fluid)

$$\Gamma_{c(t)} = \rho \oint_{c=\partial S} u dl = \rho \iint_S \omega \bullet ds \quad (\text{eq. 4.1})$$

Proof:

See [19] Majda, A.J-Bertozzi, A. L. 2002, Proposition 1.11 and Corollary 1.3, in page 23. The proof is carried actually by integrating the Euler equations on a loop c and utilizing that the integral of the pressure forces (densities) defined as $-\nabla p$ are zero as it is a conservative (irrotational) field of force (densities). Then by applying also the Stokes theorem that makes the circulation of the velocity on a loop equal to the flux of the vorticity on a smooth surface with boundary the loop (see e.g., Wikipedia Stokes theorem https://en.wikipedia.org/wiki/Stokes%27_theorem) the claim is obtained. QED.

We may notice that this circulation and surface vorticity flux has physical measuring units $[\rho][\omega][s]^2 = [m][s]^{-3}[t]^{-1}[s]^2 = [m][s]^{-1}[t]^{-1} = [\text{moment_of_inertia}][\omega][s]^{-3}$ thus angular momentum point density. While the $\rho\omega$ has physical measuring units dimensions $[\rho][\omega] = [m][s]^{-3}[t]^{-1} =$

$[\text{moment_of_inertia}][\omega][s]^{-2}$ thus 2nd spatial derivative of rotational momentum of point density .

A blow-up when it occurs, it will occur at least as blow-up of the vorticity, or of $\rho\omega$. If we discover bounded average value invariants of the flow with physical units dimensions $\rho\omega$, that in the limit can give also the point value of the $\rho\omega$, and that are invariants and bounded independent from the size of averaging, it is reasonable that we can deduce conclusions, if the point densities can blow-up or not.

Here we convert the surface vorticity flux invariant of Helmholtz-Kelvin-Stokes to one with 3D integration which will be more convenient in the arguments as the volumes are preserved by incompressible flows and most important, the integration is 3-dimensional which can be utilized to define average values of the vorticity (flux) on 3D finite particles.

We will prove at first a lemma about the 3D volume integral of Theorem 4.2, and convergence of average values of vorticity, based on this 3D integral, to point values to vorticity.

We define an average value for the volume 3D integral of vorticity flux.

Definition 4.1

We define as average value on ball in of the vorticity ω , denoted by $\bar{\omega}_B$,the unique constant value of the vorticity on the interior of the ball that would give the same 3D flux of vorticity on the ball, $\rho \int_0^\pi \iint_S \bar{\omega} \cdot ds d\theta = \rho \int_0^\pi \iint_S \omega \cdot ds d\theta$. The integration on the surfaces S for the flux of the vorticity is on parallel circular discs in the ball. This average value $\bar{\omega}$ of the vorticity is of course the

$$||\bar{\omega}_B|| = \left| \frac{\rho \int_0^\pi \iint_S \omega \cdot ds d\theta}{|B|} \right| \quad (\text{eq.4.2})$$

and its direction is that of the vertical axis of the ball B

Where $|B|=(4/3)*\pi*r^2$ is the volume of the ball B , of radius r , and $||\bar{\omega}_B||$ is the Euclidean norm of the vector. A more detailed symbolism of the average vorticity is the $\bar{\omega}(x_t, t)_{B(r,t)}$

The numerator of this average value of vorticity has also the interpretation of **rotational momentum average axial density** on the ball B and relative to the axis a . A reason for this is that the physical dimensions of measuring units of this magnitude is that of rotational momentum line density. This is because the rotational momentum point density has physical dimensions $[moment_of_inertia]*[\omega]*[s]^(-3)=[m][s]^(-1)[t]^(-1)$, where $[m]$ for mass, $[s]$ for distance, $[t]$ for time, and this magnitude has physical units dimensions, $([\rho][\omega][s]^3)=[m][s]^(-1)[t]^(-1)[s]^3$, thus rotational momentum point density integrated on 1-d line axial density. And the full quotient therefore has physical units dimensions $[m][s]^(-3)[t]^(-1)=[\rho][\omega]$.

A blow-up when it occurs, it will occur at least as blow-up of the vorticity, or of $\rho \bullet \omega$. If we discover average value invariants of the flow with physical units dimensions $\rho \bullet \omega$, that in the limit can give also the point value of the $\rho \bullet \omega$, and that are invariants and bounded independent from the size of averaging, it is reasonable that we can deduce conclusions, if the point densities can blow-up or not.

Lemma 4.1 Let a ball B of radius r and center x , and the average vorticity $\bar{\omega}_B$ in it as in the Definition 4.1 so that its axis a that defines the average vorticity is also the axis of the point vorticity ω_x at the center x of the ball. By taking the limit of shrinking the ball to its center x , ($r > 0$), the average vorticity $\bar{\omega}_B$ converges to the point vorticity ω_x . In symbols $\lim_{r \rightarrow 0} \bar{\omega}_B = \omega_x$. If the axis a of the ball to estimate the average vorticity is not the axis of the point vorticity, then the limit of the average vorticity will be equal to the projection component $\omega_a(x,t)$ of the point vorticity $\omega(x,t)$ on the axis a .

Proof: We simply apply an appropriate 3-dimensional version, with iterated integrals of the 1-dimensional fundamental theorem of the calculus. QED.

Remark 4.1. Such a limit of 3D body to a point is the same as the limit that from the Newton equation of force $F=m\gamma$, We deduce the Navier-Stokes equations.

Since the flow of a fluid under the Euler or Navier-Stokes equations, with or without smooth Schwartz external force is a smooth and continuous mapping F , then such a limit will be conserved to still be a valid limit during the flow. In other words

$F(\lim_{B \rightarrow 0} \bar{\omega}_B) = \lim_{F(B) \rightarrow 0} F(\bar{\omega}_B)$ and $B \rightarrow 0$, implies $F_t(B) \rightarrow 0$. We define of course in

an obvious appropriate way the average vorticity $F_t(\bar{\omega}_B)$ as in definition 4.1, for the flow-image of a ball B after time t . Simply the disc surfaces will no longer be flat, and the loop no longer perfect circle. But the integrals in the definition will be the same. Constancy of the average vorticity on such surfaces will only be, up to its Euclidean norm and vertical angle to the surface. We must notice though that although a relation $F(\lim_{B \rightarrow 0} \bar{\omega}_B) = \lim_{B \rightarrow 0} F(\bar{\omega}_B)$ would hold, the value of this limit **will not be** the vorticity $\omega_{F(x)}$ at the flowed point! Unfortunately, the Lemma 4.2 holds not on arbitrary 3D shapes and arbitrary integration parametrization on it, but only when we start with standard 3D shapes like a sphere, a cylinder a cube etc. and the normal parametrization on them. The reason is that we need to take in to account in a normal way the average vorticity around a point in an unbiased way, that an arbitrary shape will not give.

Another important conservation point is that the relation of the vorticity ω_x being tangent to an axis a (or general curve) is conserved during inviscid Euler flows. It is the conservation of vorticity lines (See [19] Majda, A. J. –Bertozzi, A. L. 2002, Proposition 1.9 in page 21). Therefore for inviscid (and incompressible) flows the axis of the initial point vorticity $\omega(0)$, which is also the axis to estimate the average vorticity on the ball B , will still be after the flow and at time t , tangent to the point vorticity $\omega(t)$. But for general viscous flows this will not be so. Notice that such limits of average values would not work for the circulation of the velocity on a loop, as in the application of the iterated 1-dimensional fundamental theorem of the calculus would require boundaries of the integration.

Lemma 4.2 *Let the Euler or Navier-Stokes equations of incompressible fluids in the non-periodic or periodic setting, with smooth initial data and we assume that the initial data in the periodic or non-periodic case, are so that the supremum of the vorticity is finite denoted by F_ω on all 3-space at time $t=0$. Let the average vorticity, or average rotational momentum density, defined as in Definition 4.1 but with integration parametrization on any smooth 3D shape B of any size, that of course both as a diffeomorphic image of a spherical ball with its spherical coordinates integration parametrization. Then the average vorticity or average rotational momentum density is also upper bounded by the F_ω . In symbols*

$$\|\bar{\omega}_B\| = \left| \frac{\int_0^\pi \iint_S \omega \cdot ds d\theta}{|B|} \right| \leq F_\omega \quad (\text{eq. 4.3})$$

Proof: Since $\|\omega\| \leq F_\omega = \|(\omega/\|\omega\|)\|F_\omega$ in the flux-integration we have for the inner product of ω and the unit area vector n , $(\omega, n) \leq ((\omega/\|\omega\|)F_\omega, n) \leq F_\omega$. Thus in the integration we may factor out the F_ω

$$\left| \frac{\int_0^\pi \iint_S \omega \cdot ds d\theta}{|B|} \right| \leq \left| \frac{\int_0^\pi \iint_S F_\omega ds d\theta}{|B|} \right| = F_\omega \left| \frac{\int_0^\pi \iint_S ds d\theta}{|B|} \right| = F_\omega \left| \frac{|B|}{|B|} \right| = F_\omega. \quad \text{QED.}$$

Theorem 4.2 A 3-dimensionl integral version of the Helmholtz-Kelvin-Stokes theorem. (A 2D surface density of rotatory momentum, conservation law).

*Let initial data in R^3 so that they guarantee the existence of a unique smooth solution to the Euler equation in a local time interval $[0, T]$. Then at any time $t \in [0, T]$ let a sphere B of radius r (as in figure 4.) **considered as a finite particle**, then the azimuthal θ -angle, θ -integral on a meridian in spherical coordinates of the circulations $\Gamma(c)$ of the velocities on all closed longitude smooth loops parallel to the equatorial loop is equal to the same θ -integral of the surface flux of the vorticity on smooth flat disc surfaces S with boundary the loops c (as in figures 4.2) , and both integrals are constant and preserved as both surface and volume integrals during the flow with the fluid. In symbols ($\rho=1$ is the density of the incompressible fluid)*

$$\rho \int_0^\pi \oint_{c=\partial S} u dl = \rho \int_0^\pi \iint_S \omega \cdot ds \quad (\text{eq. 4.4})$$

After (eq. 4.2) $\|\bar{\omega}_B\| = \left| \frac{\int_0^\pi \iint_S \omega \cdot ds d\theta}{|B|} \right|$ it holds also

$$\text{for } t \in [0, T] \|\bar{\omega}_{B(0)}\| = \|\bar{\omega}_{B(t)}\| \quad (\text{eq. 4.5})$$




Figure 4.1

Proof: We simply take the θ -azimuthal angle θ -integral of both sides of the equation 4.1 in the theorem 4.1. Both sides are preserved during the flow and so is their θ -integrals too. We notice that the measuring physical units dimensions of the conserved quantity $\rho \int_0^\pi \oint_{c=\partial S} u dl d\theta$ is $[\text{mass}] * [\text{length}]^{(-3)} * [\text{velocity}] * [\text{length}]^{(2)} = [\text{mass}] * [\text{length}]^{(-2)} * [\text{velocity}]$ thus integration in 2-dimension surface of momentum 3D-point-density, or equivalently momentum 1D density
QED

Theorem 4.3. The Helmholtz-Kelvin-Stokes theorem in the case of viscous Navier-Stokes equations flows without external force (homogeneous case) .

Let initial data in R^3 so that they guarantee the existence of a unique smooth solution to the Navier-Stokes equation with viscosity coefficient $\nu > 0$, in a local time interval $[0, T]$. Then at any time $t \in [0, T]$ the circulation $\Gamma(c)$ of the velocities on a closed smooth loop is equal to the flux of the vorticity on smooth surface S with boundary the loop c , and is decreasing as both loop and surface flow with the fluid. In symbols ($\rho=1$ is the density of the incompressible fluid)

$$\rho \oint_{c=\partial S} u dl = \rho \iint_S \omega \cdot ds \quad (\text{eq. 4.1})$$

$$\text{and for } t \in [0, T] \quad \oint_{c=\partial S} u(0) dl > \oint_{c=\partial S} u(t) dl \quad (\text{eq. 4.6})$$

and similarly for the 3D volume integration as in Theorem 4.2

$$\text{for } t \in [0, T] \quad \rho \int_0^\pi \iint_S \omega(0) \cdot ds d\theta > \rho \int_0^\pi \iint_S \omega(t) \cdot ds d\theta \quad (\text{eq. 4.7})$$

$$\text{After (eq. 4.2) } \|\bar{\omega}_B\| = \left| \frac{\int_0^\pi \iint_S \omega \cdot ds d\theta}{|B|} \right| \quad \text{it holds also for initial finite spherical particles for } t \in [0, T] \quad \|\bar{\omega}_{B(0)}\| > \|\bar{\omega}_{B(t)}\| \quad (\text{eq. 4.8})$$

Proof: Again The (eq. 4.1) is nothing else of course but the Stokes theorem as in (eq 4.1)

We shall utilize here the next equation (See [19] Majda, A.J-Bertozzi, A. L. 2002, (eq 1.61) , in page 23.) in the case of viscous incompressible flows under the Navier-Stokes equations

$$\frac{d}{dt} \Gamma_{c(t)} = \frac{d}{dt} \oint_{c(t)} u dl = \nu \oint_{c(t)} \Delta u dl = - \nu \oint_{c(t)} \text{curl } \omega dl \quad (\text{eq. 4.9})$$

This equation is derived after applying as in Theorem 4.1 the loop integral of the circulation at the Navier-Stokes equations instead at the Euler equations taking the material-flow derivative outside the integral, and eliminating the conservative, irrotational part of the pressure forces as gradient of the pressure. Here the viscosity is not zero thus the left hand of the equations is not zero as in the case of Euler equations, where it is conserved. **The right-hand side is nothing else than the loop work density of the point density of the force of viscosity at any time t. And as the viscosity always subtracts energy, this right-hand side work density is always negative during the flow.** We notice after the **Lemma 3.1** that the viscosity force point density keeps constant sign on the trajectory path as orbital component during the flow and relative to the velocity on the trajectory. It is always as orbital component opposite to the motion and represents the always irreversible energy absorption and linear momentum and angular momentum decrease. Similarly, for any rotation of the fluid e.g. with axis the trajectory path. The viscosity force point density as component on the loop is always opposite to the rotation, it never converts thermal energy to add to linear or angular momentum. This opposite to motion monotonicity of the viscosity force density applies to the Navier-Stokes equations but also as opposite to rotation monotonicity in the vorticity equation $\frac{D\omega}{Dt} = \omega * \nabla u + \nu \Delta \omega$ (see [19] Majda, A.J-Bertozzi, A. L. 2002, (eq 1.33) and (eq 1.50) in pages 13 and 20) . So if we choose a direction of the loop so that the circulation integral on the right hand side is positive then this will have the same sign during the

flow (although different absolute value), and will make the left hand side of the (eq. 4.9) always negative during the flow. But this means from the left-hand side of the equation that the circulation of the velocity on the loop is always decreasing during the flow.

$$\frac{d}{dt} \oint_{c(t)} u dl < 0 \quad \text{for any } t \text{ in } [0, T] \quad (\text{eq. 4.10})$$

Thus (eq. 4.6) is proved, and (eq. 4.7) is direct consequence.

To prove the equation 4.8 we notice that due to incompressibility, the flow is volume preserving, thus $|B(x(t))| = |B(x(0))|$, and by dividing both sides of the equation 4.7, and after the definition

$$\|\bar{\omega}_B\| = \left| \frac{\int_0^\pi \iint_S \omega \cdot ds d\theta}{|B|} \right| \quad \text{it holds also}$$

$$\text{for } t \in [0, T] \quad \|\bar{\omega}_{B(0)}\| > \|\bar{\omega}_{B(t)}\| \quad (\text{eq. 4.8})$$

QED.

Remark 4.2. We can extend the results of the theorems 4.1, 4.3 with Euler or Navier-Stokes equations to similar ones in the inhomogeneous case with external forces F_{ext} , provided of course we have the existence and uniqueness of a smooth solution local in time. We would start from an equation

$$\frac{d}{dt} \Gamma_{c(t)} = \rho \frac{d}{dt} \oint_{c(t)} u dl = -\nu \rho \oint_{c(t)} \text{curl } \omega dl + \rho \oint_{c(t)} F_{ext} dl \leq \rho \oint_{c(t)} F_{ext} dl$$

Similarly

$$\frac{d}{dt} \rho \iint_S \omega \cdot ds = -\nu \rho \oint_{c(t)} \text{curl } \omega dl + \rho \iint_S \text{curl } F_{ext} \cdot ds$$

$$\leq \rho \iint_S \text{curl } F_{ext} \cdot ds \leq |S| M_0$$

since as in the proof of Theorem 4.3 the friction circulation term is always negative and due to the Schwartz conditions on the external force in space and time the constant M_0 is independent from space and time and the size of the surface of the loop in the integration. $|S|$ is the area of the flux integration.

Then from smoothness and elementary 1-dimensional calculus we would get an inequality like

$$\rho \left| \iint_S \omega(0) \cdot ds - \iint_S \omega(t) \cdot ds \right| \leq |S| M_1(F_{\text{ext}}, t)$$

where again due to the Schwartz conditions on the external force in space and time the constant M_1 is independent from space and the trajectories paths and depends only on the time and the external force.

Similarly by dividing the first equation by $|B|$ which does not change by time and integrating for 3D ball, we can result similarly to an inequality like

$$\left| \frac{\int_0^t \iint_S \omega(s) \cdot ds d\theta}{|B(t)|} - \frac{\int_0^t \iint_S \omega(0) \cdot ds d\theta}{|B(0)|} \right| \leq \frac{|B|}{|B|} M_1(F_{\text{ext}}, t) = M_1(F_{\text{ext}}, t) \quad \text{where again}$$

the constant M_2 is independent from space and the size of the ball and depends only on the time t .

Theorem 4.4 The no blow-up theorem in finite or infinite time in the Euler, Navier-Stokes, periodic or non-periodic and homogeneous cases.

Let the Euler or Navier-Stokes equations of incompressible fluids in the non-periodic or periodic setting (homogeneous case with no external forces), with

*a) smooth initial data and whatever else hypothesis is necessary so as, also to guarantee the **existence and uniqueness of smooth solutions** to the equations locally in time $[0, T)$.*

b) Furthermore we assume that the initial data in the periodic or non-periodic case, are such that the supremum of the vorticity, denoted by F_ω , is finite at $t=0$. (In the periodic case, smoothness of the initial velocities is adequate to derive it, while in the non-periodic setting smooth Schwartz initial velocities is adequate to derive it)

Then it holds that there cannot exist any finite or infinite time blow-up at the point vorticities during the flow.

Proof: The proof will be by contradiction. The main idea of the proof is to utilize that in the case of a blow-up the vorticity will converge to infinite, so it will become larger than an arbitrary lower bound $M + F_\omega$, $M > 0$, $F_\omega > 0$ and by approximating it with average flux vorticity of a 3D spherical particle, and tracing it back at the initial

conditions where all is bounded by F_ω , utilizing the semi-invariance of the average vorticity that we have proved, we will get that $F_\omega > M + F_\omega$.

So let us assume that there is a blow up, in a finite time or infinite time T^* , with the hypotheses of the theorem 4.x. Then from the Theorem 2.2 and (eq. 2.12) which is the well-known result of the control of regularity or blow up by the vorticity we get that ,

$$\lim_{t \rightarrow T^*} \int_0^T |\omega(\cdot, \tau)|_{L^\infty} d\tau = +\infty \quad (\text{eq. 2.12})$$

We conclude that there will exist an infinite sequence of points $\{x_{t_n}, n \text{ natural number}, 0 < t_n < T^*, \lim_{n \rightarrow \infty} t_n = T^*\}$ so that the point vorticity $\omega(x_{t_n})$ blows-up, or equivalently $\lim_{n \rightarrow \infty} \omega(x_{t_n}) = +\infty$. We do not need to assume them on the same trajectory. Therefore, for every positive arbitrary large real number M_0 , there is a n_0 such that for all natural numbers $n > n_0$, it holds that $\omega(x_{t_n}) > M_0$. We choose $M_0 = M_{00} + F_\omega$, for an arbitrary large positive number M_{00} . So

$$\omega(x_{t_n}) > M_{00} + F_\omega \quad (\text{eq. 4.11})$$

Now we approximate this point vorticity with an average flux vorticity on a 3D particle after Definition 4.1, theorem 4.2 and Lemma 4.1.

Let a spherical ball particle $B(r, x_{t_n})$ as in theorem 4.2. with center x_{t_n} and radius $r > 0$. After Definition 4.1, theorem 4.2 and Lemma 4.1. we have that

$$\lim_{r \rightarrow 0} \bar{\omega}_B = \omega_{x(t_n)}, \text{ with } \|\bar{\omega}_B\| = \left| \frac{\int_0^{2\pi} \iint_S \omega \cdot ds d\theta}{|B(r, x(t_n))|} \right| \quad (\text{eq. 4.2})$$

Therefore for arbitrary small positive number $\varepsilon > 0$, there is radius R , with

$$\bar{\omega}_{B(R)} > \omega_{x(t_n)} - \varepsilon \quad \text{or} \quad \left| \frac{\int_0^{2\pi} \iint_S \omega \cdot ds d\theta}{|B(r, x(t_n))|} \right| > \omega_{x(t_n)} - \varepsilon \quad (\text{eq. 4.12})$$

$$\text{Thus after (eq. 4.11)} \quad \left| \frac{\int_0^{2\pi} \iint_S \omega \cdot ds d\theta}{|B(R, x(t_n))|} \right| > M_{00} + F_\omega - \varepsilon \quad (\text{eq. 4.13})$$

Now we trace back on the trajectory of the x_{t_n} the parts of the (eq. 4.13)

At initial time $t=0$. We use the advantage that as the incompressible flow is volume preserving, the $|B(R, x_0)| = |B(R, x(t_n))|$. We also utilize theorems 4.2, 4.3, and (eq. 4.5), (eq. 4.8) , which prove that at the initial conditions $t=0$, this average vorticity is the same or higher than that at t_n .

$$\left| \frac{\int_0^{2\pi} \iint_S \omega \cdot ds d\theta}{|B(R, x(0))|} \right| \geq \left| \frac{\int_0^{2\pi} \iint_S \omega \cdot ds d\theta}{|B(R, x(t_n))|} \right|$$

We conclude that

$$\left| \frac{\int_0^{2\pi} \iint_S \omega \cdot ds d\theta}{|B(R, x(0))|} \right| > M_{00} + F_\omega - \varepsilon \quad (\text{eq. 4.14})$$

From the (eq. 4.14) and (eq. 4.3) of Lemma 4.2 we conclude that

$$F_\omega > M_{00} + F_\omega - \varepsilon \quad (\text{eq. 4.15})$$

But M_{00} was chosen in an independent way from $\varepsilon > 0$ to be arbitrary large, while $\varepsilon > 0$ can be chosen to be arbitrary small. Therefore, a contradiction. Thus there cannot be any blow-up either in finite or infinite time T^* . QED.

Remark 4.3. Infinite initial energy. We must remark that we did not utilize anywhere that the initial energy was finite, only that the vorticity initially has finite supremum. Thus this result of no-blow-up can be with infinite initial energy too. But when applying it to the millennium problem we do have there also that the initial energy is finite.

Remark 4.4. Inhomogeneous case. It is interesting to try to extend this result of no blowup, for the inhomogeneous case too of the Euler and Navier-Stokes equations and investigate where it would fail, if at all, provided of course we have the existence and uniqueness of a smooth solution local in time and also that the vorticity accumulation criterion of regularity and no-blow-up still holds with external forcing as in the formulation of the millennium problem. We would utilize the last inequality of remark 4.2

$\left| \frac{\int_0^\pi \iint_S \omega(t) \cdot ds d\theta}{|B(t)|} - \frac{\int_0^\pi \iint_S \omega(0) \cdot ds d\theta}{|B(0)|} \right| \leq M_1(F_{ext}, t)$ and we would anticipate for the choice of the constant M_0 in (eq 4.11) , $M_0 = M_{00} + M_1 + F_\omega$. We would reason similarly as in the proof of the Theorem 4.4 and we end to a same contradiction

$$F_\omega > M_{00} + F_\omega - \varepsilon$$

But since at least in the book [19] Majda, A.J-Bertozzi, A. L. 2002, that I took as reference on the subject, it does not claim the existence and uniqueness of a smooth solution locally in time, in the case of external forces, as we wrote in remark 2.4, I will avoid using it, and I remain only in the homogeneous case. Therefore, for the moment I will not spend space in this paper on the inhomogeneous case.

Remark 4.5. A strange blow up for any time $t > 0$ of initially smooth data. We might be curious to ask the question if it is possible, starting with zero initial velocities and pressures, to create an artificial blow-up only with external forcing. A good candidate is the perfect circular vortex, where all the trajectory paths are perfect circles, which is known that it is an instance of the solution of the Euler and Navier-Stokes equations. We can formulate the circular vortex in 3D with cylindrical or spherical coordinates. But for simplicity we will formulate it in 2 dimensions, in spite the fact that we do know that in 2D dimensions there cannot be a blow up under the hypotheses of the millennium problem. So with an external forcing also as perfect circular vortex that in polar coordinates are as follows

$$F_{r=0} , F_\theta = 2\rho/(1+\exp(r)) \quad (\text{eq 4.16})$$

we raise the absolute initial rest within finite time t the flow to a circular vortex which has velocities in polar coordinates

$$u_{r=0} , u_\theta = 2t/(1+\exp(r)) \quad (\text{eq 4.17})$$

Now it is elementary to show that

- 1) this flow follows the Euler and Navier-Stokes equations
- 2) Because $\text{curl} \omega = 0$, the viscosity has no effect it is as if an inviscid flow.

Where ω is the vorticity which is calculated in polar coordinates at the vertical z -axis by the formula

$$\omega_z = \frac{u_\theta}{r} + \frac{\partial u_\theta}{\partial r} - \frac{\partial u_r}{r \partial r} \quad (\text{eq. 4.18})$$

4) Although the velocity has smooth polar coordinates, the vorticity is in steady blow-up (singularity) at $r=0$ for any $t>0$. That is although at $t=0$ the initial data are smooth, for any $t>0$, there is a blow-up.

5) The 4) is so because the external forcing although it has smooth polar coordinates, in the Cartesian coordinates, it has $\text{curl}(F)=+\infty$, at $r=0$, thus it does not satisfy the smooth Schwartz condition external forcing of the millennium problem.

5. The solution of the Millennium Problem for the Navier-stokes Equations but Also for the Euler Equations.

We are now in a position to prove the Conjectures (A) and (B) , non-periodic and periodic setting , homogeneous case of the Millennium problem.

(Millennium Homogeneous Case A) Existence and smoothness of Navier-Stokes solution on $\mathbb{R}^3$.

Take $\nu>0$ and $n=3$. Let $u_0(x)$ be any smooth, divergent-free vector field satisfying (2.4). Take $f(x,t)$ to be identically zero. Then there exist smooth functions $p(x,t)$, $u(x,t)$ on $\mathbb{R}^3 \times [0, +\infty)$ that satisfy (2.1), (2.2), (2.3) , (2.6) , (2.7).

Proof: All the hypotheses of the no-blow-up theorem 4.4 are satisfied. After remark 2.4, with the current case of the millennium problem there exist indeed a unique smooth solution locally in time $[0,t]$ (after A.J. Majda-A.L. Bertozzi [19] ,Theorem 3.4 pp 104, Local in Time existence of Solutions to the Euler and Navier-Stokes equations) . And also, the Schwartz condition of the initial data, guarantees that the supremum of the vorticity, is finite at $t=0$. Therefore, we conclude by Theorem 4.4 that there cannot be any finite or infinite time blow-up. Thus from **Theorem 2.2 Supremum of vorticity sufficient condition of regularity** we conclude that this local in time $[0,t]$ solution , can be extended in $[0,+\infty)$.

QED

(Millennium Homogeneous Case B) Existence and smoothness of Navier-Stokes solution on $\mathbb{R}^3/\mathbb{Z}^3$.

Take $\nu>0$ and $n=3$. Let $u_0(x)$ be any smooth, divergent-free vector field satisfying (8); we take $f(x,t)$ to be identically zero. Then there exist smooth functions $p(x,t)$, $u(x,t)$ on $\mathbb{R}^3 \times [0, +\infty)$ that satisfy (2.1), (2.2), (2.3) , (2.10) , (2.11).

Proof: All the hypotheses of the no-blow-up theorem 4.4 are satisfied. After remark 2.4, with the current case of the millennium problem there exist indeed a unique

smooth solution locally in time $[0,t]$ (after A.J. Majda-A.L. Bertozzi [19] ,Theorem 3.4 pp 104, Local in Time existence of Solutions to the Euler and Navier-Stokes equations) . And also, the compactness of the 3D torus of the initial data, guarantees that the supremum of the vorticity, is finite at $t=0$. Therefore, we conclude by Theorem 4.4 that there cannot be any finite or infinite time blow-up. Thus from **Theorem 2.2 Supremum of vorticity sufficient condition of regularity** and **remark 2.6 (that the previous theorem covers the periodic setting too)** we conclude that this local in time $[0,t]$ solution , can be extended in $[0,+\infty)$.

QED

Remark 5.1. Now in the previous two Millennium cases we could as well take $\nu=0$, and we would have the same proofs and conclusions because the Theorem 4.4 of the no-blow-up covers too the **case of inviscid Euler equations flows**.

6. Epilogue

In this paper I solved the millennium problem about the Navier-Stokes equations in the homogeneous case without external forcing, and proved that there cannot be a blowup in finite or infinite time (regularity) both in the periodic and non-periodic setting without external forcing (homogeneous case). But it is also proved that 1) once the hypotheses of external forcing of the millennium problem allow for the existence of a unique smooth solution local in time, and also 2) the vorticity accumulation criterion for a regularity and no blow up also holds with external forcing as in the formulation of the millennium problem, then the same result of regularity (no blow up) holds also for this inhomogeneous case with external forcing. Furthermore, I proved also the by far more difficult same result for the Euler inviscid flows. I did so by utilizing (e.g. in in the inviscid case) that not only the momentum is conserved but also rotatory versions of the momentum 1D line and 2D surface densities are conserved. Then I extended the conservation in the case of viscous Navier-Stokes flows to monotone semi invariants, in other words that these densities are monotonously decreasing due to friction. This allowed me to prove with elementary geometric calculus that there cannot be any blow up (regularity). The solution of this millennium problem gave the opportunity to discover 2 new monotone semi invariants (1D and 2D densities of (rotatory type) momentum) for the viscous Navier-Stokes equations.

References

1. Conway J. H. On numbers and games (Surreal numbers) Academic press 1976.
2. Durmagambetov Asset Et Al 2015 “Navier Stokes equations Millennium prize problems” Natural science, 2015, 7, 88-89 <https://www.scirp.org/journal/ns/>
3. Fefferman C. L. 2006 Existence and smoothness of the Navier-Stokes equation, *Millennium Prize Problems, Clay Math. Inst., Cambridge, MA, 2006, 57-67.*
4. Kyritsis K. 2013. Proof of the existence of global in time smooth solutions of 3-dimensional incompressible inviscid smooth flow-solutions of the Euler equations and viscous smooth flow-solutions of the Navier-Stokes equations. (The solution of the 4th Clay millennium problem). An outline of this paper was presented in the 2nd Applied Innovation conference of TEI of Epirus in Arta 16-17 October 2013. (<http://innovation-conference-epirus.blogspot.gr/2015/01/2nd-international-conference-on-applied.html>) <http://libsearch.teiep.gr/Record/cris-123456789-1324>
5. Kyritsis, K. April 2017 Ordinal Real numbers 1,2,3 Conference: 1st international conference on quantitative, social, biomedical and economic issues 2017 - ICQSBEI 2017 ,Athens, Greece Volume: PROCEEDINGS pp 233-292 <https://books.google.gr/books?id=BSUsDwAAQBAJ&pg>, see also https://www.researchgate.net/publication/355155715_THE_CONTINUUM_OF_THE_SURREAL_NUMBERS_REVISITED_THE_SURREAL_NUMBERS_DEFINED_THROUGH_TRANSFINITE_FUNDAMENTAL_CAUCHY_SEQUENCES
6. Kyritsis, K. November 2017 “On the 4th Clay Millennium problem: Proof of the regularity of the solutions of the Euler and Navier-Stokes equations, based on the conservation of particles” *Journal of Scientific Research and Studies Vol 4 (11) , pp304-317, November 2017.*
7. Kyritsis. K. Feb 2018 A solution of the 4th Clay Millennium problem about the Navier-Stokes equations. Preprint 1) <http://cris.teiep.gr/jspui/handle/123456789/1628?locale=en> 2) https://www.researchgate.net/publication/323389082_A_SOLUTION_OF_THE_4_TH_CLAY_MILLENNIUM_PROBLEM_ABOUT_THE_NAVIER-STOKES_EQUATIONS
8. Kyritsis, K. February 2019 “The solutions of the 3rd and 4th Clay Millennium problems” <https://arxiv.org/abs/1902.07265>
9. Kyritsis, K. August 2021. A solution of the 4th Clay Millennium problem about the Navier-Stokes equations. *World Journal of Research and Review WJRR.org ISSN 2455-3956 Volume 13 Issue 20 August 21 pp 25-40*

10. Kyritsis, K. 2021A The Solutions of the 3rd and 4th Millennium Mathematical Problems The solutions of the Millennium Problem P vs NP in computational complexity and the Millennium problem in fluid dynamics LAP Lambert Academic Publishing (2021-11-25) ISBN-13 978-620-4-72562-8
11. Kyritsis, K. 2021B Solution of the Clay Millennium Problem about the Navier-Stokes Equations: A Recent Approach Chapter 4 in the book *New Visions in Science and Technology Vol. 6*, 2 October 2021 , Page 35-59
12. Kyritsis K. 2021 “Review of the solutions of the Clay millennium problem about $P \neq NP = \text{exptime}$ ”. September 2021 World Journal of Research and Review 13(3):21-26 <https://www.wjrr.org/vol-13issue-3>
13. Kyritsis K. 2021 Study on the Solution of the Clay Millennium Problem about the P vs NP: A Short Proof that $P \neq NP = \text{EXPTIME}$ in the Context of Deterministic Turing Machines
14. Author of chapter 5 in the book “New Visions in Science and Technology Vol.6, October 2, 2021 , Page 60-69
<https://doi.org/10.9734/bpi/nvst/v6/5176F> Published: 2021-10-02 Chapter in a book <https://stm.bookpi.org/NVST-V6/article/view/4135>
15. Kyritsis K. 2022 . The Newton-Leibniz number-fields of fluxions and infinitesimals and the new applied Democritus digital calculus of the 21st century A partial similarity of theory and practice captured as instances of the axiomatic system of abstract calculi over multi-density quantities. To appear as book in Lambert academic publications. See also
https://www.researchgate.net/publication/358425683_THE_NEWTON-LEIBNIZ_FIELDS_OF_FLUXIONS_AND_INFINITESIMALS_AND_THE_NEW_APPLIED_DEMOCRITUS_DIGITAL_CALCULUS_OF_THE_21ST_CENTURY_A_partial_similarity_of_theory_and_practice
16. Kyritsis K. 2017 B_An Axiomatic System for a Physical or Digital but Continuous 3-Dimensional Euclidean Geometry, Without Infinite Many Points. *World Journal of Research and Review (WJRR) ISSN:2455-3956, Volume-5, Issue-4, October 2017 Pages 31-43*
17. Kyritsis K. 2019B “An introduction to the natural differential and integral calculus, without the infinite” Journal of Regional & Socio-Economic Issues . Oct 2019, Vol. 9 Issue 3, p70-89.
18. Kyritsis K. 2022 A shorter solution to the Clay millennium problem about regularity of the Navier-Stokes equations February 2022 Journal of Scientific and Industrial Research Vol 9 Feb 2022(1):8-16

19. Kyritsis K. 2022b “A 3rd Shorter Solution to the 3rd Millennium problem P vs NP=Exptime” paper presented in the 6th International conference on quantitative, social, biomedical and economic issues, ICQSBE 2022 1st July 2022, <https://icqsbei2022.blogspot.com/2022/06/blog-post.html> See also https://www.researchgate.net/publication/358226092_A_3_RD_SHORTER_SOLUTION_OF_THE_CLAY_MILLENNIUM_PROBLEM_ABOUT_P_NP_EXPTIME
20. Majda, A.J-Bertozzi, A. L. 2002 Vorticity and incompressible flow *Cambridge University texts in applied mathematics* 2002.
21. Moschandreou. T. 2021 “No finite time Blow-up for 3D incompressible Navier-Stokes equations, via scaling invariance” *Mathematica and Statistics*, 9(3): 386-393. <https://doi.org/10.13189/ms.2021.090321>
22. Muriel, A 2000 An Exact Solution of the 3-D Navier-Stokes Equation Department of Electrical Engineering Columbia University And Department of Philosophy Harvard University. Also A. Muriel and M. Dresden, *Physica D* 101, 299, 1997.
23. Ramm G. A. 2019 Solution of the Navier-Stokes problem, *Applied mathematics Letters* 87, 2019, pp 160-164/
24. Ramm G. A. 2021, Navier-Stokes equations paradox , *Reports on Math. Phys.* (ROMP), 88 N1 (2021) pp 41-45.
25. Tao, T. 2013 Localisation and compactness properties of the Navier-Stokes global regularity problem. *Analysis & PDE* 6 (2013), 25-107
26. Tao, T. 2014 Finite time blowup for an averaged three-dimensional Navier-Stokes equation. *Cornell University Library arxiv.org*