

# POINTWISE MULTIPLIERS BETWEEN WEIGHTED COPSON AND CESÀRO FUNCTION SPACES

A.GOGATISHVILI, R.CH. MUSTAFAYEV, T. ÜNVER

**ABSTRACT.** In this paper the solution of the pointwise multiplier problem between weighted Copson function spaces  $\text{Cop}_{p_1, q_1}(u_1, v_1)$  and weighted Cesàro function spaces  $\text{Ces}_{p_2, q_2}(u_2, v_2)$  is presented, where  $p_1, p_2, q_1, q_2 \in (0, \infty)$ ,  $p_2 \leq q_2$  and  $u_1, u_2, v_1, v_2$  are weights on  $(0, \infty)$ .

## 1. INTRODUCTION

Various properties of the different spaces defined by the Cesàro operator have been studied extensively in the literature. The Cesàro sequence spaces  $\text{ces}_p$  and the Cesàro function spaces  $\text{Ces}_p$  have been introduced by Shiue in 1970 in [17] and [18], respectively. In 1971 Leibowitz proved that  $\text{ces}_1 = \{0\}$  and for  $1 < q < p \leq \infty$ ,  $\ell_p$  and  $\text{ces}_q$  sequence spaces are proper subspaces of  $\text{ces}_p$  [16]. Jagers has obtained the associate space for  $\text{ces}_p$ , ( $1 < p < \infty$ ) [14]. In [19], Sy, Zhang and Lee gave a description of dual spaces of  $\text{Ces}(p)$  spaces based on Jagers' result. In 1996 different, isomorphic description due to Bennett appeared in [4]. In [4, Theorem 21.1] Bennett observes that the classical Cesàro function space and the classical Copson function space coincide for  $p > 1$ . He also derives estimates for the norms of the corresponding inclusion operators. The same result, with different estimates, is due to Boas [6], who in fact obtained the integral analogue of the Askey-Boas Theorem [5, Lemma 6.18] and [1]. These results generalized in [13] using the blocking technique. In [2] they investigated dual spaces for  $\text{Ces}(p)$  for  $1 < p < \infty$ . Their description can be viewed as being analogous to one given for sequence spaces in [4] (For more detailed information about history of classical Cesàro spaces see recent survey paper [3]).

In this paper, we will describe the pointwise multiplier spaces of weighted Copson and Cesàro function spaces. The weighted Cesàro and Copson function spaces are defined in [8] as follows:

---

2010 *Mathematics Subject Classification.* Primary 26D10; Secondary 26D15.

*Key words and phrases.* Cesàro and Copson function spaces, embeddings, iterated Hardy inequalities, weights.

The research of A. Gogatishvili was partly supported by the grants P201-13-14743S of the Grant Agency of the Czech Republic and RVO: 67985840, by Shota Rustaveli National Science Foundation grants no. 31/48 (Operators in some function spaces and their applications in Fourier Analysis) and no. DI/9/5-100/13 (Function spaces, weighted inequalities for integral operators and problems of summability of Fourier series). The research of all authors was partly supported by the joint project between Academy of Sciences of Czech Republic and The Scientific and Technological Research Council of Turkey.

**Definition 1.1.** Let  $0 < p, q \leq \infty$ ,  $u \in \mathfrak{M}^+(0, \infty)$ ,  $v \in \mathcal{W}(0, \infty)$ . The weighted Cesàro and Copson spaces are defined by

$$\text{Ces}_{p,q}(u, v) := \left\{ f \in \mathfrak{M}^+(0, \infty) : \|f\|_{\text{Ces}_{p,q}(u, v)} := \left\| \|f\|_{p,v,(0,\cdot)} \right\|_{q,u,(0,\infty)} < \infty \right\},$$

and

$$\text{Cop}_{p,q}(u, v) := \left\{ f \in \mathfrak{M}^+(0, \infty) : \|f\|_{\text{Cop}_{p,q}(u, v)} := \left\| \|f\|_{p,v,(\cdot,\infty)} \right\|_{q,u,(0,\infty)} < \infty \right\},$$

respectively.

**Definition 1.2.** Let  $0 < q \leq \infty$ . We denote by  $\Omega_q$  the set of all functions  $u \in \mathfrak{M}^+(0, \infty)$  such that

$$0 < \|u\|_{q,(t,\infty)} < \infty, \quad t > 0,$$

and by  ${}^c\Omega_q$  the set of all functions  $u \in \mathfrak{M}^+(0, \infty)$  such that

$$0 < \|u\|_{q,(0,t)} < \infty, \quad t > 0.$$

Let  $v \in \mathcal{W}(0, \infty)$ . It is easy to see that  $\text{Ces}_{p,q}(u, v)$  and  $\text{Cop}_{p,q}(u, v)$  are quasi-normed vector spaces when  $u \in \Omega_q$  and  $u \in {}^c\Omega_q$ , respectively.

Note that the function spaces  $C$  and  $D$  defined by Grosse-Erdmann in [13] are related with this definition in the following way:

$$\text{Ces}_{p,q}(u, v) = C(p, q, u)_v \quad \text{and} \quad \text{Cop}_{p,q}(u, v) = D(p, q, u)_v.$$

We use the notations  $\text{Ces}_p(u) := \text{Ces}_{1,p}(u, \mathbf{1})$  and  $\text{Cop}_p(u) := \text{Cop}_{1,p}(u, \mathbf{1})$ . Obviously,  $\text{Ces}(p) = \text{Ces}_p(x^{-1})$  and  $\text{Cop}(p) = \text{Cop}_p(x^{-1})$ . In [15], Kamińska and Kubiak computed the dual norm of the Cesàro function space  $\text{Ces}_p(u)$ , generated by  $1 < p < \infty$  and an arbitrary positive weight  $u$ . A description presented in [15] resembles the approach of Jagers [14] for sequence spaces.

Let  $X$  and  $Y$  be a (quasi)-Banach spaces of Borel measurable functions on  $(0, \infty)$ . A *multiplier* from a space  $X$  into a space  $Y$  is a function  $f$  such that  $f \cdot g \in Y$  for all  $g \in X$ . In particular, the Köthe dual  $X'$  of  $X$  is defined as the space  $\mathcal{M}(X, L_1)$  of multipliers into  $L_1$ . The (linear) space of all such multipliers is denoted by  $\mathcal{M}(X, Y)$ , that is,

$$\mathcal{M}(X, Y) := \{f : f \cdot g \in Y \text{ for all } g \in X\}.$$

The  $\mathcal{M}(X, Y)$  becomes a quasi-normed space with the quantity

$$\|f\|_{\mathcal{M}(X, Y)} := \sup_{g \neq 0} \frac{\|fg\|_Y}{\|g\|_X}.$$

Given  $f \in \mathcal{W}(0, \infty)$ , if we define a weighted space by  $Y_f = \{g : f \cdot g \in Y\}$ , then

$$\|f\|_{\mathcal{M}(X, Y)} = \sup_{g \neq 0} \frac{\|g\|_{Y_f}}{\|g\|_X} = \|I\|_{X \rightarrow Y_f},$$

and we see that  $\|f\|_{\mathcal{M}(X, Y)} < \infty$  if and only if  $\|I\|_{X \rightarrow Y_f} < \infty$ .

The multiplier problem between  $\ell_s(w)$  and one of the  $\text{ces}_{p,q}(a, b)$  and  $\text{cop}_{p,q}(a, b)$  spaces was considered by Grosse-Erdmann [13]. It is mentioned in [13, p. 30] that multipliers between two spaces of type  $\text{ces}_{p,q}(a, b)$  and  $\text{cop}_{p,q}(a, b)$  are more difficult to treat.

In [8] the embeddings between the Copson and Cesàro function spaces and vice versa, that is, the embeddings

$$(1.1) \quad \text{Cop}_{p_1, q_1}(u_1, v_1) \hookrightarrow \text{Ces}_{p_2, q_2}(u_2, v_2),$$

$$(1.2) \quad \text{Ces}_{p_2, q_2}(u_2, v_2) \hookrightarrow \text{Cop}_{p_1, q_1}(u_1, v_1).$$

were characterized. Using these characterizations we give the solution to the multiplier problem between weighted Cesàro and Copson function spaces.

It is easy to see that the problem of characterization of the space of multipliers between  $\text{Cop}_{p_1, q_1}(u_1, v_1)$  and  $\text{Ces}_{p_2, q_2}(u_2, v_2)$  can be reduced to the characterization of the multiplier problem with three parameters and three weight functions (see, Proposition 3.5). So we will concentrate our attention on calculation of  $M(\text{Cop}_r(u), \text{Ces}_{p, q}(w, v))$ .

In order to describe the pointwise multipliers, we need to introduce the following weighted Cesàro-type spaces with three parameters:

**Definition 1.3.** Let  $0 < p, q, r \leq \infty$ ,  $u \in \mathfrak{M}^+(0, \infty)$ ,  $v, w \in \mathcal{W}(0, \infty)$ . The weighted Cesàro and Copson type spaces are defined by

$$\text{Ces}_{p, q, r}(u, v, w) := \left\{ f \in \mathfrak{M}^+(0, \infty) : \|f\|_{\text{Ces}_{p, q, r}(u, v, w)} := \left\| \left\| \|f\|_{p, w, (0, \cdot)} \right\|_{q, v, (0, \cdot)} \right\|_{r, u, (0, \infty)} < \infty \right\},$$

and

$$\text{Cop}_{p, q, r}(u, v, w) := \left\{ f \in \mathfrak{M}^+(0, \infty) : \|f\|_{\text{Cop}_{p, q, r}(u, v, w)} := \left\| \left\| \|f\|_{p, w, (\cdot, \infty)} \right\|_{q, v, (\cdot, \infty)} \right\|_{r, u, (0, \infty)} < \infty \right\},$$

respectively.

Throughout the paper we assume that  $I := (a, b) \subseteq (0, \infty)$ . By  $\mathfrak{M}(I)$  we denote the set of all measurable functions on  $I$ . The symbol  $\mathfrak{M}^+(I)$  stands for the collection of all  $f \in \mathfrak{M}(I)$  which are non-negative on  $I$ , while  $\mathfrak{M}^+(I; \downarrow)$  and  $\mathfrak{M}^+(I; \uparrow)$  are used to denote the subset of those functions which are non-increasing and non-decreasing on  $I$ , respectively. When  $I = (0, \infty)$ , we write simply  $\mathfrak{M}^+$ ,  $\mathfrak{M}^\downarrow$  and  $\mathfrak{M}^\uparrow$  instead of  $\mathfrak{M}^+(I)$ ,  $\mathfrak{M}^+(I; \downarrow)$  and  $\mathfrak{M}^+(I; \uparrow)$ , accordingly. The family of all weight functions (also called just weights) on  $I$ , that is, measurable, positive and finite a.e. on  $I$ , is given by  $\mathcal{W}(I)$ .

For  $p \in (0, \infty]$ , we define the functional  $\|\cdot\|_{p, I}$  on  $\mathfrak{M}(I)$  by

$$\|f\|_{p, I} := \begin{cases} \left( \int_I |f(x)|^p dx \right)^{\frac{1}{p}} & \text{if } p < \infty, \\ \text{ess sup}_I |f(x)| & \text{if } p = \infty. \end{cases}$$

If  $w \in \mathcal{W}(I)$ , then the weighted Lebesgue space  $L_p(w, I)$  is given by

$$L_p(w, I) \equiv L_{p, w}(I) := \{f \in \mathfrak{M}(I) : \|f\|_{p, w, I} := \|fw\|_{p, I} < \infty\},$$

and it is equipped with the quasi-norm  $\|\cdot\|_{p, w, I}$ . When  $I = (0, \infty)$ , we often write simply  $L_{p, w}$  and  $L_p(w)$  instead of  $L_{p, w}(I)$  and  $L_p(w, I)$ , respectively.

We adopt the following usual conventions.

**Convention 1.4.** (i) Throughout the paper we put  $0/0 = 0$ ,  $0 \cdot (\pm\infty) = 0$  and  $1/(\pm\infty) = 0$ .

(ii) We put

$$p' := \begin{cases} \frac{p}{1-p} & \text{if } 0 < p < 1, \\ \infty & \text{if } p = 1, \\ \frac{p}{p-1} & \text{if } 1 < p < \infty, \\ 1 & \text{if } p = \infty. \end{cases}$$

(iii) If  $I = (a, b) \subseteq \mathbb{R}$  and  $g$  is a monotone function on  $I$ , then by  $g(a)$  and  $g(b)$  we mean the limits  $\lim_{x \rightarrow a^+} g(x)$  and  $\lim_{x \rightarrow b^-} g(x)$ , respectively.

To state our results we use the notation  $p \rightarrow q$  for  $0 < p, q \leq \infty$  defined by

$$\frac{1}{p \rightarrow q} = \frac{1}{q} - \frac{1}{p} \quad \text{if } q < p,$$

and  $p \rightarrow q = \infty$  if  $q \geq p$  (see, for instance, [13, p.30]).

Throughout the paper, we always denote by  $c$  and  $C$  a positive constant, which is independent of main parameters but it may vary from line to line. However a constant with subscript or superscript such as  $c_1$  does not change in different occurrences. By  $a \lesssim b$ ,  $(b \gtrsim a)$  we mean that  $a \leq \lambda b$ , where  $\lambda > 0$  depends on inessential parameters. If  $a \lesssim b$  and  $b \lesssim a$ , we write  $a \approx b$  and say that  $a$  and  $b$  are equivalent. We will denote by  $\mathbf{1}$  the function  $\mathbf{1}(x) = 1$ ,  $x \in \mathbb{R}$ . Given two quasi-normed vector spaces  $X$  and  $Y$ , we write  $X = Y$  if  $X$  and  $Y$  are equal in the algebraic and the topological sense (their quasi-norms are equivalent). The symbol  $X \hookrightarrow Y$  ( $Y \hookleftarrow X$ ) means that  $X \subset Y$  and the natural embedding  $I$  of  $X$  in  $Y$  is continuous, that is, there exist a constant  $c > 0$  such that  $\|z\|_Y \leq c\|z\|_X$  for all  $z \in X$ . The best constant of the embedding  $X \hookrightarrow Y$  is  $\|I\|_{X \rightarrow Y}$ .

The paper is organized as follows. Some new "gluing" lemmas are presented in Section 2. The characterization of spaces of multipliers between weighted Cesàro and Copson function spaces are given in Section 3.

## 2. SOME NEW "GLUING" LEMMAS

In this section, we present some generalizations of "gluing" lemma, which is very useful and has independent interest (cf. [7, Lemma 2.2] and [9, Theorem 3.1]).

Recall that, if  $F \in \mathfrak{M}^+((0, \infty); \downarrow)$ , then

$$(2.1) \quad \underset{t \in (0, \infty)}{\text{ess sup}} F(t)G(t) = \underset{t \in (0, \infty)}{\text{ess sup}} F(t) \underset{\tau \in (0, t)}{\text{ess sup}} G(\tau);$$

likewise, when  $F \in \mathfrak{M}^+((0, \infty); \uparrow)$ , then

$$(2.2) \quad \underset{t \in (0, \infty)}{\text{ess sup}} F(t)G(t) = \underset{t \in (0, \infty)}{\text{ess sup}} F(t) \underset{\tau \in (t, \infty)}{\text{ess sup}} G(\tau)$$

(see, for instance, [11, p. 85]).

Given  $a \in \mathfrak{M}^+((0, \infty); \uparrow)$  denote by

$$\mathcal{A}(x, t) := \frac{a(x)}{a(x) + a(t)} \quad (x > 0, t > 0).$$

Observe that

$$\mathcal{A}(x, t) \approx \min \left\{ 1, \frac{a(x)}{a(t)} \right\}.$$

Moreover,

$$(2.3) \quad a(t)\mathcal{A}(x, t) = a(x)\mathcal{A}(t, x).$$

We say that a function  $f$  is  $a$ -quasiconcave if  $f$  is equivalent to an increasing function on  $(0, \infty)$  and  $f/a$  is equivalent to a decreasing function on  $(0, \infty)$ . It is easy to see that  $\mathcal{A}(x, t)$  is  $a$ -quasiconcave function of  $x$  for any fixed  $t > 0$ .

It have been shown in [11, p. 85] that the relations

$$(2.4) \quad \begin{aligned} \text{ess sup}_{t \in (0, \infty)} \mathcal{A}(x, t)g(t) &\approx \text{ess sup}_{t \in (0, \infty)} g(t) \min \left\{ 1, \frac{a(x)}{a(t)} \right\} \\ &= \text{ess sup}_{t \in (0, x)} a(t) \text{ess sup}_{\tau \in (t, \infty)} \frac{g(\tau)}{a(\tau)} \\ &= a(x) \text{ess sup}_{t \in (x, \infty)} \frac{1}{a(t)} \text{ess sup}_{\tau \in (0, t)} g(\tau) \end{aligned}$$

holds for any  $g \in \mathfrak{M}^+(0, \infty)$  and  $a \in \mathfrak{M}^+((0, \infty); \uparrow)$ . Consequently,  $\text{ess sup}_{t \in (0, \infty)} \mathcal{A}(x, t)g(t)$  is  $a$ -quasiconcave function.

**Lemma 2.1.** *Let  $a \in \mathcal{W}(0, \infty)$  be non-decreasing and  $g, h \in \mathfrak{M}^+(0, \infty)$ . Then*

$$\begin{aligned} \text{ess sup}_{x \in (0, \infty)} \left( \text{ess sup}_{t \in (0, \infty)} \mathcal{A}(x, t)g(t) \right) \left( \text{ess sup}_{t \in (0, \infty)} \mathcal{A}(t, x)h(t) \right) \\ \approx \text{ess sup}_{x \in (0, \infty)} g(x) \left( \text{ess sup}_{t \in (x, \infty)} h(t) \right) \\ + \text{ess sup}_{x \in (0, \infty)} a(x)^{-1} g(x) \left( \text{ess sup}_{t \in (0, x)} a(t)h(t) \right). \end{aligned}$$

*Proof.* Using the second relation in (2.4), by (2.1), (2.3), and (2.2), we obtain that

$$\begin{aligned} \text{ess sup}_{x \in (0, \infty)} \left( \text{ess sup}_{t \in (0, \infty)} \mathcal{A}(x, t)g(t) \right) \left( \text{ess sup}_{t \in (0, \infty)} \mathcal{A}(t, x)h(t) \right) \\ \approx \text{ess sup}_{x \in (0, \infty)} g(x) \left( \text{ess sup}_{t \in (0, \infty)} \mathcal{A}(t, x)h(t) \right). \end{aligned}$$

It remains to observe that

$$\text{ess sup}_{t \in (0, \infty)} \mathcal{A}(t, x)h(t) \approx a(x)^{-1} \text{ess sup}_{t \in (0, x)} a(t)h(t) + \text{ess sup}_{t \in (x, \infty)} h(t).$$

□

Assume that  $g \in \mathfrak{M}^+(0, \infty)$ ,  $a \in \mathfrak{M}^+((0, \infty); \uparrow)$  and  $0 < \beta < \infty$ . Recall that the function

$$\left( \int_0^\infty \mathcal{A}(x, t)^\beta g(t) dt \right)^{\frac{1}{\beta}}$$

is  $a$ -quasiconcave function, as well (see, for instance, [10, p. 318]).

**Lemma 2.2.** *Let  $\beta$  be a positive number. Suppose that  $g, h \in \mathfrak{M}^+(0, \infty)$  and  $a \in \mathcal{W}(0, \infty)$  is non-decreasing. Then*

$$\begin{aligned} & \text{ess sup}_{x \in (0, \infty)} \left( \text{ess sup}_{t \in (0, \infty)} \mathcal{A}(x, t) g(t) \right) \left( \int_0^\infty \mathcal{A}(t, x)^\beta h(t) dt \right)^{\frac{1}{\beta}} \\ & \approx \text{ess sup}_{x \in (0, \infty)} g(x) \left( \int_x^\infty h(t) dt \right)^{\frac{1}{\beta}} + \text{ess sup}_{x \in (0, \infty)} a(x)^{-1} g(x) \left( \int_0^x a(t)^\beta h(t) dt \right)^{\frac{1}{\beta}}. \end{aligned}$$

*Proof.* Using the second relation in (2.4), by (2.1), (2.3), and (2.2), we obtain that

$$\begin{aligned} & \text{ess sup}_{x \in (0, \infty)} \left( \text{ess sup}_{t \in (0, \infty)} \mathcal{A}(x, t) g(t) \right) \left( \int_0^\infty \mathcal{A}(t, x)^\beta h(t) dt \right)^{\frac{1}{\beta}} \\ & \approx \text{ess sup}_{x \in (0, \infty)} g(x) \left( \int_0^\infty \mathcal{A}(t, x)^\beta h(t) dt \right)^{\frac{1}{\beta}}. \end{aligned}$$

To complete the proof it remains to observe that

$$\left( \int_0^\infty \mathcal{A}(t, x)^\beta h(t) dt \right)^{\frac{1}{\beta}} \approx a(x)^{-1} \left( \int_0^x a(t)^\beta h(t) dt \right)^{\frac{1}{\beta}} + \left( \int_x^\infty h(t) dt \right)^{\frac{1}{\beta}}.$$

□

**Lemma 2.3.** *Let  $\beta$  be a positive number. Suppose that  $g, h \in \mathfrak{M}^+(0, \infty)$  and  $a \in \mathcal{W}(0, \infty)$  is non-decreasing. Then*

$$\begin{aligned} & \text{ess sup}_{x \in (0, \infty)} \left( \int_0^\infty \mathcal{A}(x, t)^\beta g(t) dt \right)^{\frac{1}{\beta}} \left( \text{ess sup}_{t \in (0, \infty)} \mathcal{A}(t, x) h(t) \right) \\ & \approx \text{ess sup}_{x \in (0, \infty)} h(x) \left( \int_0^x g(t) dt \right)^{\frac{1}{\beta}} + \text{ess sup}_{x \in (0, \infty)} a(x) h(x) \left( \int_x^\infty a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}}. \end{aligned}$$

*Proof.* Using identity (2.3), the statement easily follows from Lemma 2.2. □

We quote some known results. Proofs can be found in [12, Proposition 2.1].

**Definition 2.4.** Let  $N, M \in \overline{\mathbb{Z}}$ ,  $N < M$ . A positive almost non-increasing sequence  $\{\tau_k\}_{k=N}^M$  (that is, there exists  $K \geq 1$  such that  $\tau_{n+1} \leq K\tau_n$ ) is called *almost geometrically decreasing* if there are  $\alpha \in (1, \infty)$  and  $L \in \mathbb{N}$  such that

$$\alpha\tau_k \leq \tau_{k-L} \quad \text{for all } k \in \{N+L, \dots, M\}.$$

A positive almost non-decreasing sequence  $\{\sigma_k\}_{k=N}^M$  (that is, there exists  $K \geq 1$  such that  $\sigma_n \leq K\sigma_{n+1}$ ) is called *almost geometrically increasing* if there are  $\alpha \in (1, +\infty)$  and  $L \in \mathbb{N}$  such that

$$\sigma_k \geq \alpha\sigma_{k-L} \quad \text{for all } k \in \{N+L, \dots, M\}.$$

**Lemma 2.5.** Let  $q \in (0, \infty]$ ,  $N, M \in \overline{\mathbb{Z}}$ ,  $N \leq M$ ,  $\mathcal{Z} = \{N, N+1, \dots, M-1, M\}$  and let  $\{\tau_k\}_{k=N}^M$  be an almost geometrically decreasing sequence. Then

$$(2.5) \quad \left\| \left\{ \tau_k \sum_{m=N}^k a_m \right\} \right\|_{\ell^q(\mathcal{Z})} \approx \| \{\tau_k a_k\} \|_{\ell^q(\mathcal{Z})}$$

and

$$(2.6) \quad \left\| \left\{ \tau_k \sup_{N \leq m \leq k} a_m \right\} \right\|_{\ell^q(\mathcal{Z})} \approx \| \{\tau_k a_k\} \|_{\ell^q(\mathcal{Z})}$$

for all non-negative sequences  $\{a_k\}_{k=N}^M$ .

**Lemma 2.6.** Let  $q \in (0, +\infty]$ ,  $N, M \in \overline{\mathbb{Z}}$ ,  $N \leq M$  and let  $\{\sigma_k\}_{k=N}^M$  be an almost geometrically increasing sequence. Then

$$(2.7) \quad \left\| \left\{ \sigma_k \sum_{m=k}^M a_m \right\} \right\|_{\ell^q(\mathcal{Z})} \approx \| \{\sigma_k a_k\} \|_{\ell^q(\mathcal{Z})}$$

and

$$(2.8) \quad \left\| \left\{ \sigma_k \sup_{k \leq m \leq M} a_m \right\} \right\|_{\ell^q(\mathcal{Z})} \approx \| \{\sigma_k a_k\} \|_{\ell^q(\mathcal{Z})}$$

for all non-negative sequences  $\{a_k\}_{k=N}^M$ .

**Lemma 2.7.** Let  $\alpha$  and  $\beta$  be positive numbers. Suppose that  $g, h \in \mathfrak{M}^+(0, \infty)$  and  $a \in \mathcal{W}(0, \infty)$  is non-decreasing. Then

$$\begin{aligned} & \text{ess sup}_{x \in (0, \infty)} \left( \int_0^\infty \mathcal{A}(x, t)^\beta g(t) dt \right)^{\frac{1}{\beta}} \left( \int_0^\infty \mathcal{A}(t, x)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \\ & \approx \text{ess sup}_{x \in (0, \infty)} \left( \int_0^x g(t) dt \right)^{\frac{1}{\beta}} \left( \int_x^\infty h(t) dt \right)^{\frac{1}{\alpha}} \\ & \quad + \text{ess sup}_{x \in (0, \infty)} \left( \int_x^\infty a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} \left( \int_0^x a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}}. \end{aligned}$$

*Proof.* Denote by

$$\begin{aligned} A_5 &:= \text{ess sup}_{x \in (0, \infty)} \left( \int_0^x g(t) dt \right)^{\frac{1}{\beta}} \left( \int_x^\infty h(t) dt \right)^{\frac{1}{\alpha}}, \\ A_6 &:= \text{ess sup}_{x \in (0, \infty)} \left( \int_x^\infty a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} \left( \int_0^x a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}}. \end{aligned}$$

Obviously,

$$\begin{aligned} & \text{ess sup}_{x \in (0, \infty)} \left( \int_0^\infty \mathcal{A}(x, t)^\beta g(t) dt \right)^{\frac{1}{\beta}} \left( \int_0^\infty \mathcal{A}(t, x)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \\ & \approx \text{ess sup}_{x \in (0, \infty)} \left( \int_0^x g(t) dt + a(x)^\beta \int_x^\infty a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} \left( a(x)^{-\alpha} \int_0^x a(t)^\alpha h(t) dt + \int_x^\infty h(t) dt \right)^{\frac{1}{\alpha}} \\ & \approx A_5 + A_6 + B_5 + B_6, \end{aligned}$$

where

$$\begin{aligned} B_5 &:= \text{ess sup}_{x \in (0, \infty)} \left( \int_0^x g(t) dt \right)^{\frac{1}{\beta}} a(x)^{-1} \left( \int_0^x a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}}, \\ B_6 &:= \text{ess sup}_{x \in (0, \infty)} a(x) \left( \int_x^\infty a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} \left( \int_x^\infty h(t) dt \right)^{\frac{1}{\alpha}}. \end{aligned}$$

It is enough to show that  $B_i \lesssim A_5 + A_6$ ,  $i = 5, 6$ .

First, let us show that  $B_5 \lesssim A_5 + A_6$ . We will consider the case when  $\int_0^\infty g(t) dt < \infty$  (The case when  $\int_0^\infty g(t) dt = \infty$  is much simpler to treat). Define a sequence  $\{x_m\}_{m=-\infty}^M$  such that  $\int_0^{x_m} g(t) dt = 2^m$  if  $-\infty < m \leq M$  and  $2^M \leq \int_0^\infty g(t) dt < 2^{M+1}$ . Denote by  $x_{M+1} := \infty$ . Then, by (2.2) and Lemma 2.6, we have that

$$\begin{aligned} B_5 &= \text{ess sup}_{x \in (0, \infty)} \left( \int_0^x g(t) dt \right)^{\frac{1}{\beta}} \text{ess sup}_{y \in (x, \infty)} a(y)^{-1} \left( \int_0^y a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \\ &\approx \sup_{-\infty < m \leq M} 2^{\frac{m}{\beta}} \text{ess sup}_{y \in (x_m, \infty)} a(y)^{-1} \left( \int_0^y a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \\ &\approx \sup_{-\infty < m \leq M} 2^{\frac{m}{\beta}} \text{ess sup}_{y \in (x_m, x_{m+1})} a(y)^{-1} \left( \int_0^y a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}}. \end{aligned}$$

For every  $-\infty < m \leq M$ , there exists  $y_m \in (x_m, x_{m+1})$  such that

$$\text{ess sup}_{y \in (x_m, x_{m+1})} a(y)^{-1} \left( \int_0^y a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \leq 2a(y_m)^{-1} \left( \int_0^{y_m} a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}}.$$

Therefore,

$$\begin{aligned} B_5 &\lesssim \sup_{-\infty < m \leq M} 2^{\frac{m}{\beta}} a(y_m)^{-1} \left( \int_0^{y_m} a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \\ &\approx \sup_{-\infty < m \leq M} 2^{\frac{m}{\beta}} a(y_m)^{-1} \left( \int_0^{y_{m-2}} a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} + \sup_{-\infty < m \leq M} 2^{\frac{m}{\beta}} a(y_m)^{-1} \left( \int_{y_{m-2}}^{y_m} a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} =: I + II. \end{aligned}$$

Note that  $\int_{y_{m-2}}^{y_m} g(t) dt \approx 2^m$ ,  $-\infty < m \leq M$ . It yields that

$$\begin{aligned} I &\approx \sup_{-\infty < m \leq M} \left( \int_{y_{m-2}}^{y_m} g(t) dt \right)^{\frac{1}{\beta}} a(y_m)^{-1} \left( \int_0^{y_{m-2}} a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \\ &= \sup_{-\infty < m \leq M} \text{ess sup}_{y_{m-2} < x < y_m} \left( \int_x^{y_m} g(t) dt \right)^{\frac{1}{\beta}} a(y_m)^{-1} \left( \int_0^{y_{m-2}} a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \\ &\leq \sup_{-\infty < m \leq M} \text{ess sup}_{y_{m-2} < x < y_m} \left( \int_x^{y_m} a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} \left( \int_0^{y_{m-2}} a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \\ &\leq \sup_{-\infty < m \leq M} \text{ess sup}_{y_{m-2} < x < y_m} \left( \int_x^\infty a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} \left( \int_0^x a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \\ &\lesssim \text{ess sup}_{x \in (0, \infty)} \left( \int_x^\infty a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} \left( \int_0^x a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} = A_6. \end{aligned}$$

For  $II$  we have that

$$\begin{aligned} II &\approx \sup_{-\infty < m \leq M} \left( \int_{y_{m-4}}^{y_{m-2}} g(t) dt \right)^{\frac{1}{\beta}} a(y_m)^{-1} \left( \int_{y_{m-2}}^{y_m} a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \\ &= \sup_{-\infty < m \leq M} \text{ess sup}_{y_{m-4} < x < y_{m-2}} \left( \int_{y_{m-4}}^x g(t) dt \right)^{\frac{1}{\beta}} a(y_m)^{-1} \left( \int_{y_{m-2}}^{y_m} a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \\ &\leq \sup_{-\infty < m \leq M} \text{ess sup}_{y_{m-4} < x < y_{m-2}} \left( \int_{y_{m-4}}^x g(t) dt \right)^{\frac{1}{\beta}} \left( \int_{y_{m-2}}^{y_m} h(t) dt \right)^{\frac{1}{\alpha}} \\ &\leq \sup_{-\infty < m \leq M} \text{ess sup}_{y_{m-4} < x < y_{m-2}} \left( \int_{y_{m-4}}^x g(t) dt \right)^{\frac{1}{\beta}} \left( \int_x^{y_m} h(t) dt \right)^{\frac{1}{\alpha}} \\ &\leq \sup_{-\infty < m \leq M} \text{ess sup}_{y_{m-4} < x < y_{m-2}} \left( \int_0^x g(t) dt \right)^{\frac{1}{\beta}} \left( \int_x^\infty h(t) dt \right)^{\frac{1}{\alpha}} \\ &\leq \text{ess sup}_{x \in (0, \infty)} \left( \int_0^x g(t) dt \right)^{\frac{1}{\beta}} \left( \int_x^\infty h(t) dt \right)^{\frac{1}{\alpha}} = A_5. \end{aligned}$$

Combining, we get that  $B_5 \lesssim A_5 + A_6$ .

Now we show that  $B_6 \lesssim A_5 + A_6$ . Let  $\int_0^\infty a(t)^{-\beta} g(t) dt < \infty$  (It is much simpler to deal with the case when  $\int_0^\infty a(t)^{-\beta} g(t) dt = \infty$ ). Define a sequence  $\{x_m\}_{m=N}^\infty$  such that  $\int_{x_m}^\infty a(t)^{-\beta} g(t) dt = 2^{-m}$  if  $N \leq m < \infty$  and  $2^{-N} < \int_0^\infty a(t)^{-\beta} g(t) dt \leq 2^{-N+1}$ . Denote by  $x_{N-1} := 0$ .

By using Lemma 2.5, we find that

$$B_6 = \text{ess sup}_{x \in (0, \infty)} \left( \int_x^\infty a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} \text{ess sup}_{y \in (0, x)} a(y) \left( \int_y^\infty h(t) dt \right)^{\frac{1}{\alpha}}$$

$$\begin{aligned} &\approx \sup_{N \leq m < \infty} 2^{-\frac{m}{\beta}} \operatorname{ess\,sup}_{y \in (0, x_m)} a(y) \left( \int_y^\infty h(t) dt \right)^{\frac{1}{\alpha}} \\ &\approx \sup_{N \leq m < \infty} 2^{-\frac{m}{\beta}} \operatorname{ess\,sup}_{y \in (x_{m-1}, x_m)} a(y) \left( \int_y^\infty h(t) dt \right)^{\frac{1}{\alpha}}. \end{aligned}$$

For every  $N \leq m < \infty$ , there exists  $y_m \in (x_{m-1}, x_m)$  such that

$$\operatorname{ess\,sup}_{y \in (x_{m-1}, x_m)} a(y) \left( \int_y^\infty h(t) dt \right)^{\frac{1}{\alpha}} \leq 2a(y_m) \left( \int_{y_m}^\infty h(t) dt \right)^{\frac{1}{\alpha}}.$$

Hence

$$\begin{aligned} B_6 &\lesssim \sup_{N \leq m < \infty} 2^{-\frac{m}{\beta}} a(y_m) \left( \int_{y_m}^\infty h(t) dt \right)^{\frac{1}{\alpha}} \\ &\approx \sup_{N \leq m < \infty} 2^{-\frac{m}{\beta}} a(y_m) \left( \int_{y_m}^{y_{m+2}} h(t) dt \right)^{\frac{1}{\alpha}} + \sup_{N \leq m < \infty} 2^{-\frac{m}{\beta}} a(y_m) \left( \int_{y_{m+2}}^\infty h(t) dt \right)^{\frac{1}{\alpha}} =: III + IV. \end{aligned}$$

Since  $\int_{y_m}^{y_{m+2}} a(t)^{-\beta} g(t) dt \approx 2^{-m}$ , we have that

$$\begin{aligned} III &\approx \sup_{N \leq m < \infty} \left( \int_{y_{m+2}}^{y_{m+4}} a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} a(y_m) \left( \int_{y_m}^{y_{m+2}} h(t) dt \right)^{\frac{1}{\alpha}} \\ &= \sup_{N \leq m < \infty} \operatorname{ess\,sup}_{y_{m+2} < x < y_{m+4}} \left( \int_x^{y_{m+4}} a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} a(y_m) \left( \int_{y_m}^{y_{m+2}} h(t) dt \right)^{\frac{1}{\alpha}} \\ &\leq \sup_{N \leq m < \infty} \operatorname{ess\,sup}_{y_{m+2} < x < y_{m+4}} \left( \int_x^{y_{m+4}} a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} \left( \int_{y_m}^{y_{m+2}} a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \\ &\leq \sup_{N \leq m < \infty} \operatorname{ess\,sup}_{y_{m+2} < x < y_{m+4}} \left( \int_x^{y_{m+4}} a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} \left( \int_{y_m}^x a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \\ &\leq \sup_{N \leq m < \infty} \operatorname{ess\,sup}_{y_{m+2} < x < y_{m+4}} \left( \int_x^\infty a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} \left( \int_0^x a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} \\ &\approx \operatorname{ess\,sup}_{x \in (0, \infty)} \left( \int_x^\infty a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} \left( \int_0^x a(t)^\alpha h(t) dt \right)^{\frac{1}{\alpha}} = A_6. \end{aligned}$$

Moreover,

$$\begin{aligned} IV &\approx \sup_{N \leq m < \infty} \left( \int_{y_m}^{y_{m+2}} a(t)^{-\beta} g(t) dt \right)^{\frac{1}{\beta}} a(y_m) \left( \int_{y_{m+2}}^\infty h(t) dt \right)^{\frac{1}{\alpha}} \\ &\leq \sup_{N \leq m < \infty} \left( \int_{y_m}^{y_{m+2}} g(t) dt \right)^{\frac{1}{\beta}} \left( \int_{y_{m+2}}^\infty h(t) dt \right)^{\frac{1}{\alpha}} \\ &\leq \sup_{N \leq m < \infty} \left( \int_0^{y_{m+2}} g(t) dt \right)^{\frac{1}{\beta}} \left( \int_{y_{m+2}}^\infty h(t) dt \right)^{\frac{1}{\alpha}} \end{aligned}$$

$$\lesssim \operatorname{ess\,sup}_{x \in (0, \infty)} \left( \int_0^x g(t) dt \right)^{\frac{1}{\beta}} \left( \int_x^\infty h(t) dt \right)^{\frac{1}{\alpha}} = A_5.$$

Therefore, we obtain  $B_6 \lesssim A_5 + A_6$ . The proof is complete.  $\square$

**Lemma 2.8.** *Let  $\alpha, \beta, \gamma$  be positive numbers. Suppose that  $g, h \in \mathfrak{M}^+(0, \infty)$  and  $a \in \mathcal{W}(0, \infty)$  is non-decreasing. Then*

$$\begin{aligned} & \int_0^\infty \left( \int_0^\infty \mathcal{A}(x, t)^\alpha g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \int_0^\infty \mathcal{A}(t, x)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} g(x) dx \\ & \approx \int_0^\infty \left( \int_0^x g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \int_x^\infty h(t) dt \right)^{\frac{\gamma}{\beta}} g(x) dx \\ & \quad + \int_0^\infty \left( \int_x^\infty a(t)^{-\alpha} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \int_0^x a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} a(x)^{-\alpha} g(x) dx. \end{aligned}$$

*Proof.* Denote by

$$\begin{aligned} A_7 &:= \int_0^\infty \left( \int_0^x g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \int_x^\infty h(t) dt \right)^{\frac{\gamma}{\beta}} g(x) dx, \\ A_8 &:= \int_0^\infty \left( \int_x^\infty a(t)^{-\alpha} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \int_0^x a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} a(x)^{-\alpha} g(x) dx. \end{aligned}$$

Obviously,

$$\begin{aligned} & \int_0^\infty \left( \int_0^\infty \mathcal{A}(x, t)^\alpha g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \int_0^\infty \mathcal{A}(t, x)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} g(x) dx \\ & \approx \int_0^\infty \left( \int_0^x g(t) dt + a(x)^\alpha \int_x^\infty a(t)^{-\alpha} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( a(x)^{-\beta} \int_0^x a(t)^\beta h(t) dt + \int_x^\infty h(t) dt \right)^{\frac{\gamma}{\beta}} g(x) dx \\ & \approx A_7 + A_8 + B_7 + B_8, \end{aligned}$$

where

$$\begin{aligned} B_7 &:= \int_0^\infty \left( \int_0^x g(t) dt \right)^{\frac{\gamma}{\alpha}-1} a(x)^{-\gamma} \left( \int_0^x a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} g(x) dx, \\ B_8 &:= \int_0^\infty \left( \int_x^\infty a(t)^{-\alpha} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \int_x^\infty h(t) dt \right)^{\frac{\gamma}{\beta}} a(x)^{\gamma-\alpha} g(x) dx. \end{aligned}$$

It is enough to show that  $B_i \lesssim A_7 + A_8$ ,  $i = 7, 8$ .

First, let us show that  $B_7 \lesssim A_7 + A_8$ . Let  $\int_0^\infty g(t) dt < \infty$  (The case when  $\int_0^\infty g(t) dt = \infty$  is much simpler to treat). Define a sequence  $\{x_m\}_{m=-\infty}^M$  such that  $\int_0^{x_m} g(t) dt = 2^m$  if  $-\infty < m \leq M$  and  $2^M \leq \int_0^\infty g(t) dt < 2^{M+1}$ . Denote by  $x_{M+1} := \infty$ . Then we have by Lemma 2.6 that

$$B_7 \leq \int_0^\infty \left( \int_0^x g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \operatorname{ess\,sup}_{y \in (x, \infty)} a(y)^{-\gamma} \left( \int_0^y a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} \right) g(x) dx$$

$$\begin{aligned} &\approx \sum_{m=-\infty}^M 2^{m\frac{\gamma}{\alpha}} \operatorname{ess\,sup}_{y \in (x_m, \infty)} a(y)^{-\gamma} \left( \int_0^y a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} \\ &\approx \sum_{m=-\infty}^M 2^{m\frac{\gamma}{\alpha}} \operatorname{ess\,sup}_{y \in (x_m, x_{m+1})} a(y)^{-\gamma} \left( \int_0^y a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}}. \end{aligned}$$

For every  $-\infty < m \leq M$  there exists  $y_m \in (x_m, x_{m+1})$  such that

$$\operatorname{ess\,sup}_{y \in (x_m, x_{m+1})} a(y)^{-\gamma} \left( \int_0^y a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} \leq 2a(y_m)^{-\gamma} \left( \int_0^{y_m} a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}}.$$

Therefore,

$$\begin{aligned} B_7 &\lesssim \sum_{m=-\infty}^M 2^{m\frac{\gamma}{\alpha}} a(y_m)^{-\gamma} \left( \int_0^{y_m} a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} \\ &\approx \sum_{m=-\infty}^M 2^{m\frac{\gamma}{\alpha}} a(y_m)^{-\gamma} \left( \int_0^{y_{m-2}} a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} + \sum_{m=-\infty}^M 2^{m\frac{\gamma}{\alpha}} a(y_m)^{-\gamma} \left( \int_{y_{m-2}}^{y_m} a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} =: V + VI. \end{aligned}$$

Since  $\int_{y_{m-2}}^{y_m} \left( \int_x^{y_m} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} g(x) dx \approx 2^{m\frac{\gamma}{\alpha}}$  for  $-\infty < m \leq M$ , we have that

$$\begin{aligned} V &\approx \sum_{m=-\infty}^M \left( \int_{y_{m-2}}^{y_m} \left( \int_x^{y_m} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} g(x) dx \right) a(y_m)^{-\gamma} \left( \int_0^{y_{m-2}} a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} \\ &\leq \sum_{m=-\infty}^M \left( \int_{y_{m-2}}^{y_m} \left( \int_x^{y_m} a(t)^{-\alpha} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} a(x)^{-\alpha} g(x) dx \right) \left( \int_0^{y_{m-2}} a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} \\ &\leq \sum_{m=-\infty}^M \int_{y_{m-2}}^{y_m} \left( \int_x^\infty a(t)^{-\alpha} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \int_0^x a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} a(x)^{-\alpha} g(x) dx \\ &\lesssim \int_0^\infty \left( \int_x^\infty a(\tau)^{-\alpha} g(\tau) d\tau \right)^{\frac{\gamma}{\alpha}-1} \left( \int_0^x a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} a(x)^{-\alpha} g(x) dx = A_8. \end{aligned}$$

In view of  $\int_{y_{m-4}}^{y_{m-2}} \left( \int_{y_{m-4}}^x g(t) dt \right)^{\frac{\gamma}{\alpha}-1} g(x) dx \approx 2^{m\frac{\gamma}{\alpha}}$ ,  $-\infty < m \leq M$ , we get that

$$\begin{aligned} VI &\approx \sum_{m=-\infty}^M \left( \int_{y_{m-4}}^{y_{m-2}} \left( \int_{y_{m-4}}^x g(t) dt \right)^{\frac{\gamma}{\alpha}-1} g(x) dx \right) a(y_m)^{-\gamma} \left( \int_{y_{m-2}}^{y_m} a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} \\ &\leq \sum_{m=-\infty}^M \left( \int_{y_{m-4}}^{y_{m-2}} \left( \int_{y_{m-4}}^x g(t) dt \right)^{\frac{\gamma}{\alpha}-1} g(x) dx \right) \left( \int_{y_{m-2}}^{y_m} h(t) dt \right)^{\frac{\gamma}{\beta}} \\ &\leq \sum_{m=-\infty}^M \int_{y_{m-4}}^{y_{m-2}} \left( \int_0^x g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \int_x^\infty h(t) dt \right)^{\frac{\gamma}{\beta}} g(x) dx \end{aligned}$$

$$\lesssim \int_0^\infty \left( \int_0^x g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \int_x^\infty h(t) dt \right)^{\frac{\gamma}{\beta}} g(x) dx = A_7.$$

Combining, we get that  $B_7 \lesssim A_7 + A_8$ .

Now we show that  $B_8 \lesssim A_7 + A_8$ . Let  $\int_0^\infty a(t)^{-\alpha} g(t) dt < \infty$  (It is much simpler to deal with the case when  $\int_0^\infty a(t)^{-\alpha} g(t) dt = \infty$ ). Define a sequence  $\{x_m\}_{m=N}^\infty$  such that  $\int_{x_m}^\infty a(t)^{-\alpha} g(t) dt = 2^{-m}$  if  $N \leq m < \infty$  and  $2^{-N} < \int_0^\infty a(t)^{-\alpha} g(t) dt \leq 2^{-N+1}$ . Denote by  $x_{N-1} := 0$ . By using elementary calculations, in view of Lemma 2.5, we find that

$$\begin{aligned} B_8 &\leq \int_0^\infty \left( \int_x^\infty a(t)^{-\alpha} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \operatorname{ess\,sup}_{y \in (0,x)} a(y)^\gamma \left( \int_y^\infty h(t) dt \right)^{\frac{\gamma}{\beta}} \right) a(x)^{-\alpha} g(x) dx \\ &\approx \sum_{m=N}^\infty 2^{-m\frac{\gamma}{\alpha}} \operatorname{ess\,sup}_{y \in (0,x_m)} a(y)^\gamma \left( \int_y^\infty h(t) dt \right)^{\frac{\gamma}{\beta}} \\ &\approx \sum_{m=N}^\infty 2^{-m\frac{\gamma}{\alpha}} \operatorname{ess\,sup}_{y \in (x_{m-1},x_m)} a(y)^\gamma \left( \int_y^\infty h(t) dt \right)^{\frac{\gamma}{\beta}}. \end{aligned}$$

For every  $m = N, N+1, \dots$  there exists  $y_m \in (x_{m-1}, x_m)$  such that

$$\operatorname{ess\,sup}_{y \in (x_{m-1},x_m)} a(y)^\gamma \left( \int_y^\infty h(t) dt \right)^{\frac{\gamma}{\beta}} \leq 2a(y_m)^\gamma \left( \int_{y_m}^\infty h(t) dt \right)^{\frac{\gamma}{\beta}}.$$

Hence

$$\begin{aligned} B_8 &\lesssim \sum_{m=N}^\infty 2^{-m\frac{\gamma}{\alpha}} a(y_m)^\gamma \left( \int_{y_m}^\infty h(t) dt \right)^{\frac{\gamma}{\beta}} \\ &\approx \sum_{m=N}^\infty 2^{-m\frac{\gamma}{\alpha}} a(y_m)^\gamma \left( \int_{y_m}^{y_{m+2}} h(t) dt \right)^{\frac{\gamma}{\beta}} + \sum_{m=N}^\infty 2^{-m\frac{\gamma}{\alpha}} a(y_m)^\gamma \left( \int_{y_{m+2}}^\infty h(t) dt \right)^{\frac{\gamma}{\beta}} =: VII + VIII. \end{aligned}$$

Since  $\int_{y_{m+2}}^{y_{m+4}} \left( \int_x^{y_{m+4}} a(t)^{-\alpha} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} a(x)^{-\alpha} g(x) dx \approx 2^{-m\frac{\gamma}{\alpha}}$ ,  $N \leq m < \infty$ , we have that

$$\begin{aligned} VII &\approx \sum_{m=N}^\infty \left( \int_{y_{m+2}}^{y_{m+4}} \left( \int_x^{y_{m+4}} a(t)^{-\alpha} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} a(x)^{-\alpha} g(x) dx \right) a(y_m)^\gamma \left( \int_{y_m}^{y_{m+2}} h(t) dt \right)^{\frac{\gamma}{\beta}} \\ &\leq \sum_{m=N}^\infty \left( \int_{y_{m+2}}^{y_{m+4}} \left( \int_x^{y_{m+4}} a(t)^{-\alpha} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} a(x)^{-\alpha} g(x) dx \right) \left( \int_{y_m}^{y_{m+2}} a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} \\ &\leq \sum_{m=N}^\infty \int_{y_{m+2}}^{y_{m+4}} \left( \int_x^\infty a(t)^{-\alpha} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \int_0^x a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} a(x)^{-\alpha} g(x) dx \\ &\lesssim \int_0^\infty \left( \int_x^\infty a(\tau)^{-\alpha} g(\tau) d\tau \right)^{\frac{\gamma}{\alpha}-1} \left( \int_0^x a(t)^\beta h(t) dt \right)^{\frac{\gamma}{\beta}} a(x)^{-\alpha} g(x) dx = A_8. \end{aligned}$$

In view of  $\int_{y_m}^{y_{m+2}} \left( \int_{y_m}^x a(t)^{-\alpha} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} a(x)^{-\alpha} g(x) dx \approx 2^{-m\frac{\gamma}{\alpha}}$ ,  $N \leq m < \infty$ , we get that

$$\begin{aligned} VIII &\approx \sum_{m=N}^{\infty} \left( \int_{y_m}^{y_{m+2}} \left( \int_{y_m}^x a(t)^{-\alpha} g(t) dt \right)^{\frac{\gamma}{\alpha}-1} a(x)^{-\alpha} g(x) dx \right) a(y_m)^\gamma \left( \int_{y_{m+2}}^{\infty} h(t) dt \right)^{\frac{\gamma}{\beta}} \\ &\leq \sum_{m=N}^{\infty} \left( \int_{y_m}^{y_{m+2}} \left( \int_{y_m}^x g(t) dt \right)^{\frac{\gamma}{\alpha}-1} g(x) dx \right) \left( \int_{y_{m+2}}^{\infty} h(t) dt \right)^{\frac{\gamma}{\beta}} \\ &\leq \sum_{m=N}^{\infty} \int_{y_m}^{y_{m+2}} \left( \int_0^x g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \int_x^{\infty} h(t) dt \right)^{\frac{\gamma}{\beta}} g(x) dx \\ &\lesssim \int_0^{\infty} \left( \int_0^x g(t) dt \right)^{\frac{\gamma}{\alpha}-1} \left( \int_x^{\infty} h(t) dt \right)^{\frac{\gamma}{\beta}} g(x) dx = A_7. \end{aligned}$$

Therefore, we obtain  $B_8 \lesssim A_7 + A_8$ . The proof is complete.  $\square$

Finally, we recall the following statement from [8].

**Lemma 2.9.** [8, Lemma 3.13] *Let  $\beta$  be a positive number. Suppose that  $g, h \in \mathfrak{M}^+(0, \infty)$  and  $a \in \mathcal{W}(0, \infty)$  is non-decreasing. Then*

$$\begin{aligned} &\int_0^{\infty} \left( \int_0^{\infty} \mathcal{A}(x, t) g(t) dt \right)^{\beta-1} \left( \text{ess sup}_{t \in (0, \infty)} \mathcal{A}(t, x) h(t) \right)^\beta g(x) dx \\ &\quad \approx \int_0^{\infty} \left( \int_0^x g(t) dt \right)^{\beta-1} \left( \text{ess sup}_{t \in (x, \infty)} h(t) \right)^\beta g(x) dx \\ &\quad + \int_0^{\infty} \left( \int_x^{\infty} a(t)^{-1} g(t) dt \right)^{\beta-1} \left( \text{ess sup}_{t \in (0, x)} a(t) h(t) \right)^\beta a(x)^{-1} g(x) dx. \end{aligned}$$

### 3. MULTIPLIERS

To state further results we need the following definitions.

**Definition 3.1.** Let  $U$  be a continuous, strictly increasing function on  $[0, \infty)$  such that  $U(0) = 0$  and  $\lim_{t \rightarrow \infty} U(t) = \infty$ . Then we say that  $U$  is admissible.

Let  $U$  be an admissible function. We say that a  $U$ -quasiconcave function  $\varphi$  is non-degenerate if

$$\lim_{t \rightarrow 0+} \varphi(t) = \lim_{t \rightarrow \infty} \frac{1}{\varphi(t)} = \lim_{t \rightarrow \infty} \frac{\varphi(t)}{U(t)} = \lim_{t \rightarrow 0+} \frac{U(t)}{\varphi(t)} = 0.$$

The family of non-degenerate  $U$ -quasiconcave functions is denoted by  $\mathcal{Q}_U$ .

**Definition 3.2.** Let  $U$  be an admissible function, and let  $w$  be a non-negative measurable function on  $(0, \infty)$ . We say that the function  $\varphi$ , defined by

$$\varphi(t) = U(t) \int_0^{\infty} \frac{w(\tau) d\tau}{U(\tau) + U(t)}, \quad t \in (0, \infty),$$

is a fundamental function of  $w$  with respect to  $U$ . One will also say that  $w(\tau)d\tau$  is a representation measure of  $\varphi$  with respect to  $U$ .

**Remark 3.3.** Let  $\varphi$  be the fundamental function of  $w$  with respect to  $U$ . Assume that

$$\int_0^\infty \frac{w(\tau)d\tau}{U(\tau) + U(t)} < \infty, \quad t > 0, \quad \int_0^1 \frac{w(\tau)d\tau}{U(\tau)} = \int_1^\infty w(\tau)d\tau = \infty.$$

Then  $\varphi \in Q_U$ .

**Remark 3.4.** Suppose that  $\varphi(x) < \infty$  for all  $x \in (0, \infty)$ , where  $\varphi$  is defined by

$$\varphi(x) = \text{ess sup}_{t \in (0, x)} U(t) \text{ess sup}_{\tau \in (t, \infty)} \frac{w(\tau)}{U(\tau)}, \quad t \in (0, \infty).$$

If

$$\limsup_{t \rightarrow 0^+} w(t) = \limsup_{t \rightarrow +\infty} \frac{1}{w(t)} = \limsup_{t \rightarrow 0^+} \frac{U(t)}{w(t)} = \limsup_{t \rightarrow +\infty} \frac{w(t)}{U(t)} = 0,$$

then  $\varphi \in Q_U$ .

We are going to use the following operators

$$A_{q,p} : \mathfrak{M}^+(0, \infty) \rightarrow \mathfrak{M}^+(0, \infty) \quad \text{and} \quad A_{q,p}^* : \mathfrak{M}^+(0, \infty) \rightarrow \mathfrak{M}^+(0, \infty)$$

defined by

$$A_{q,p}(u)(x) := \left( \int_0^x u^q \right)^{-\frac{1}{p}} u^{\frac{q-p}{p}}(x) \quad \text{and} \quad A_{q,p}^*(u)(x) := \left( \int_x^\infty u^q \right)^{\frac{1}{p}} u^{\frac{p-q}{p}}(x)$$

for all  $x > 0$ .

In order to state our results we will use the following notations:

$$V(x) := \|v\|_{p', 0, x} \quad \text{and} \quad \mathcal{V}(x, t) := \frac{V(x)}{V(x) + V(t)} \quad (t > 0, x > 0).$$

The following simple statement allows us to reduce the characterization of the multiplier problem between weighted Copson function spaces  $\text{Cop}_{p_1, q_1}(u_1, v_1)$  and weighted Cesàro function spaces  $\text{Ces}_{p_2, q_2}(u_2, v_2)$  to the problem with three parameters and three weight functions.

**Proposition 3.5.** Let  $0 < p_1, p_2, q_1, q_2 < \infty$  and  $v_1, v_2 \in \mathcal{W}(0, \infty)$ ,  $u_1 \in {}^c\Omega_{q_1}$  and  $u_2 \in \Omega_{q_2}$ . Then

$$\|f\|_{M(\text{Cop}_{p_1, q_1}(u_1, v_1), \text{Ces}_{p_2, q_2}(u_2, v_2))} = \left\| f^{p_1} \right\|_{M\left(\text{Cop}_{\frac{q_1}{p_1}}(u_1^{p_1}), \text{Ces}_{\frac{p_2}{p_1}, \frac{q_2}{p_1}}(u_2^{p_1}, v_1^{-p_1} v_2^{p_1})\right)}^{\frac{1}{p_1}}$$

*Proof.* Indeed:

$$\|f\|_{M(\text{Cop}_{p_1, q_1}(u_1, v_1), \text{Ces}_{p_2, q_2}(u_2, v_2))} = \sup_{g \neq 0} \frac{\|fg\|_{\text{Ces}_{p_2, q_2}(u_2, v_2)}}{\|g\|_{\text{Cop}_{p_1, q_1}(u_1, v_1)}}$$

$$\begin{aligned}
&= \sup_{g \succ 0} \frac{\left( \int_0^\infty \left( \int_0^t f^{p_2} g^{p_2} v_2^{p_2} \right)^{\frac{q_2}{p_2}} u_2^{q_2}(t) dt \right)^{\frac{1}{q_2}}}{\left( \int_0^\infty \left( \int_t^\infty g^{p_1} v_1^{p_1} \right)^{\frac{q_1}{p_1}} u_1^{q_1}(t) dt \right)^{\frac{1}{q_1}}} \\
&= \sup_{h \succ 0} \frac{\left( \int_0^\infty \left( \int_0^t f^{p_2} h^{\frac{p_2}{p_1}} v_1^{-p_2} v_2^{p_2} \right)^{\frac{q_2}{p_2}} u_2^{q_2}(t) dt \right)^{\frac{1}{q_2}}}{\left( \int_0^\infty \left( \int_t^\infty h^{\frac{q_1}{p_1}} u_1^{q_1}(t) dt \right)^{\frac{1}{q_1}} \right)} \\
&= \sup_{h \succ 0} \frac{\left( \int_0^\infty \left( \int_0^t [f^{p_1} h]^{\frac{p_2}{p_1}} [v_1^{-p_1} v_2^{p_1}]^{\frac{p_2}{p_1}} \right)^{\frac{p_1 q_2}{p_2 p_1}} [u_2(t)^{p_1}]^{\frac{q_2}{p_1}} dt \right)^{\frac{p_1}{q_2 p_1}}}{\left( \int_0^\infty \left( \int_t^\infty h^{\frac{q_1}{p_1}} [u_1(t)^{p_1}]^{\frac{q_1}{p_1}} dt \right)^{\frac{p_1}{q_1 p_1}} \right)} \\
&= \|f^{p_1}\|_{M(\text{Cop}_{\frac{q_1}{p_1}}(u_1^{p_1}), \text{Ces}_{\frac{p_2}{p_1}, \frac{q_2}{p_1}}(u_2^{p_1}, v_1^{-p_1} v_2^{p_1}))}^{\frac{1}{p_1}}.
\end{aligned}$$

□

Now we present and prove our main results. We start with simple cases.

**Theorem 3.6.** *Let  $0 < q = p \leq r = 1$ ,  $v \in \mathcal{W}(0, \infty)$ ,  $u \in {}^c\Omega_1$  and  $w \in \Omega_q$ . Then*

$$M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v)) = L_{p'}(\omega),$$

where

$$\omega := A_{1,1}(u) \cdot A_{p,p}^*(w) \cdot v.$$

Moreover

$$\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} \approx \|f\|_{p', \omega, (0, \infty)}.$$

*Proof.* By [8, Theorem 2.1], we have that

$$\begin{aligned}
\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} &= \|I\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w, fv)} \\
&\approx \left\| \|u\|_{1, (0, \cdot)}^{-1} \|w\|_{p, (\cdot, \infty)} \right\|_{p', fv, (0, \infty)} \\
&= \|f\|_{p', \omega, (0, \infty)}.
\end{aligned}$$

□

**Theorem 3.7.** *Let  $0 < p, q, r < \infty$ ,  $q \neq p$  and  $p \leq r = 1$ . Let  $v \in \mathcal{W}(0, \infty)$ ,  $u \in {}^c\Omega_r$  and  $w \in \Omega_q$ .*

(i) *If  $1 \leq q$ , then*

$$M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v)) = \text{Ces}_{p', \infty}(\omega_1, \omega_2),$$

where

$$\omega_1 := A_{p,p}^*(w), \quad \omega_2 := A_{1,1}(u) \cdot v.$$

Moreover

$$\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} \approx \|f\|_{\text{Ces}_{p', \infty}(\omega_1, \omega_2)}.$$

(ii) If  $q < 1$ , then

$$M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v)) = \text{Ces}_{p',q'}(\omega_1, \omega_2),$$

where

$$\omega_1 := A_{q,1}^*(w), \quad \omega_2 := A_{1,1}(u) \cdot v.$$

Moreover

$$\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} \approx \|f\|_{\text{Ces}_{p',q'}(\omega_1, \omega_2)}.$$

*Proof.* (i) Let  $1 \leq q$ . By [8, Theorem 2.2, (i)], we get that

$$\begin{aligned} \|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} &= \|\mathbf{I}\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w, fv)} \\ &\approx \sup_{t \in (0, \infty)} \left\| \|u\|_{1,(0,\cdot)}^{-1} \right\|_{p', fv, (0,t)} \|w\|_{q,(t,\infty)} \\ &\approx \sup_{t \in (0, \infty)} \|f\|_{p', \|u\|_{1,(0,\cdot)}^{-1} v, (0,t)} \|w\|_{q,(t,\infty)} \\ &= \left\| \|f\|_{p', \omega_2, (0,\cdot)} \right\|_{\infty, \omega_1, (0,\infty)} \\ &= \|f\|_{\text{Ces}_{p',\infty}(\omega_1, \omega_2)}. \end{aligned}$$

(ii) Let  $q < 1$ . By [8, Theorem 2.2, (ii)], we obtain that

$$\begin{aligned} \|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} &= \|\mathbf{I}\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w, fv)} \\ &\approx \left( \int_{(0, \infty)} \left\| \|u\|_{1,(0,\cdot)}^{-1} \right\|_{p', fv, (0,t)}^{q'} d\left(-\|w\|_{q,(t,\infty)}^{q'}\right) \right)^{\frac{1}{q'}} \\ &= \left( \int_{(0, \infty)} \|f\|_{p', \|u\|_{1,(0,\cdot)}^{-1} v, (0,t)}^{q'} d\left(-\|w\|_{q,(t,\infty)}^{q'}\right) \right)^{\frac{1}{q'}} \\ &= \left\| \|f\|_{p', \omega_2, (0,\cdot)} \right\|_{q', \omega_1, (0,\infty)} \\ &= \|f\|_{\text{Ces}_{p',q'}(\omega_1, \omega_2)}. \end{aligned}$$

□

**Theorem 3.8.** Let  $0 < p, q, r < \infty$ ,  $r \neq 1$  and  $p = q \leq 1$ . Let  $v \in \mathcal{W}(0, \infty)$ ,  $u \in {}^c\Omega_r$  and  $w \in \Omega_q$ .

(i) If  $r \leq p$ , then

$$M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v)) = \text{Ces}_{p',\infty}(\omega_1, \omega_2),$$

where

$$\omega_1 := A_{r,r}(u), \quad \omega_2 := A_{p,p}^*(w) \cdot v.$$

Moreover,

$$\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} \approx \|f\|_{\text{Ces}_{p',\infty}(\omega_1, \omega_2)}.$$

(ii) If  $p < r$ , then

$$\begin{aligned} M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v)) \\ = \begin{cases} \text{Ces}_{p',r \rightarrow p}(\omega_1, \omega_2) \cap L_{p'}(\omega_2) & \text{if } \|u\|_{r,(0,\infty)} < \infty, \\ \text{Ces}_{p',r \rightarrow p}(\omega_1, \omega_2) & \text{if } \|u\|_{r,(0,\infty)} = \infty, \end{cases} \end{aligned}$$

where

$$\omega_1 := A_{r,p}(u), \quad \omega_2 := A_{p,p}^*(w)v.$$

Moreover,

$$\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} \approx \|f\|_{\text{Ces}_{p',r \rightarrow p}(\omega_1, \omega_2)} + \|u\|_{r,(0,\infty)} \|f\|_{p',\omega_2,(0,\infty)}.$$

*Proof.* (i) Let  $r \leq p$ . By [8, Theorem 2.3, (i)], we get that

$$\begin{aligned} \|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} &\approx \sup_{t \in (0, \infty)} \|u\|_{r,(0,t)}^{-1} \left\| \|w\|_{p,(\cdot,\infty)} \right\|_{1 \rightarrow p, f_v, (0,t)} \\ &= \sup_{t \in (0, \infty)} \|u\|_{r,(0,t)}^{-1} \|f\|_{p', \|w\|_{p,(\cdot,\infty)} v, (0,t)} \\ &= \left\| \|f\|_{p', \omega_2, (0,\cdot)} \right\|_{\infty, \omega_1, (0,\infty)} \\ &= \|f\|_{\text{Ces}_{p',\infty}(\omega_1, \omega_2)}. \end{aligned}$$

(ii) Let  $p < r$ . By [8, Theorem 2.3, (ii)], we get that

$$\begin{aligned} \|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} &\approx \left( \int_{(0,\infty)} \left\| \|w\|_{p,(\cdot,\infty)} \right\|_{1 \rightarrow p, f_v, (0,t)}^{r \rightarrow p} d(-\|u\|_{r,(0,t)}^{-r \rightarrow p}) \right)^{\frac{1}{r \rightarrow p}} \\ &\quad + \|u\|_{r,(0,\infty)}^{-1} \left\| \|w\|_{p,(\cdot,\infty)} \right\|_{1 \rightarrow p, f_v, (0,\infty)} \\ &= \left( \int_{(0,\infty)} \|f\|_{p', \|w\|_{p,(\cdot,\infty)} v, (0,t)}^{r \rightarrow p} d(-\|u\|_{r,(0,t)}^{-r \rightarrow p}) \right)^{\frac{1}{r \rightarrow p}} \\ &\quad + \|u\|_{r,(0,\infty)}^{-1} \|f\|_{p', \|w\|_{p,(\cdot,\infty)} v, (0,\infty)} \\ &= \|f\|_{\text{Ces}_{p',r \rightarrow p_2}(\omega_1, \omega_2)} + \|u\|_{r,(0,\infty)}^{-1} \|f\|_{p', \omega_2, (0,\infty)}. \end{aligned}$$

□

**Theorem 3.9.** Let  $0 < p, q, r < \infty$ ,  $p < 1$  and  $r \leq p < q$ . Let  $v \in \mathcal{W}(0, \infty)$ ,  $u \in {}^\circ\Omega_r$  and  $w \in \Omega_q$ . Assume that  $V$  is admissible and  $\varphi_1 \in Q_{V^{1/p'}}$ , where

$$\varphi_1(x) := \text{ess sup}_{t \in (0, \infty)} \mathcal{V}(x, t)V(t)\|u\|_{r,(0,t)}^{-1}.$$

(i) If  $1 \leq q$ , then

$$M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v)) = \text{Ces}_{p',\infty}(\omega, v),$$

where

$$\omega := A_{r,r}(u) \cdot A_{q,q}^*(w).$$

Moreover,

$$\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} \approx \|f\|_{\text{Ces}_{p',\infty}(\omega, v)}.$$

(ii) If  $q < 1$ , then

$$M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v)) = \text{Ces}_{p',\infty}(\omega_1, v) \cap \text{Ces}_{p',q',\infty}(\omega_2, \omega_3, v),$$

where

$$\omega_1 := A_{r,r}(u) \cdot A_{q,q}^*(w), \quad \omega_2 := A_{r,r}(u), \quad \omega_3 := A_{q,1}^*(w).$$

Moreover

$$\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w,v))} \approx \|f\|_{\text{Ces}_{p',\infty}(\omega_1, v)} + \|f\|_{\text{Ces}_{p',q',\infty}(\omega_2, \omega_3, v)}.$$

*Proof.* (i) By [8, Theorem 2.8, (i)], we have that

$$\|I\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w,v)} \approx \sup_{x \in (0,\infty)} \left( \text{ess sup}_{t \in (0,\infty)} \mathcal{V}(x,t) V(t) \|u\|_{r,(0,t)}^{-1} \right) \left( \sup_{t \in (0,\infty)} \mathcal{V}(t,x) \|w\|_{q,(t,\infty)} \right)$$

Using Lemma 2.1 with

$$a(t) = V(t), \quad g(t) = V(t) \|u\|_{r,(0,t)}^{-1}, \quad \text{and} \quad h(t) = \|w\|_{q,(t,\infty)} \quad (t > 0),$$

in view of (2.1), we obtain that

$$\|I\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w,v)} \approx \text{ess sup}_{x \in (0,\infty)} V(x) \|u\|_{r,(0,x)}^{-1} \|w\|_{q,(x,\infty)}.$$

Therefore,

$$\begin{aligned} \|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w,v))} &\approx \text{ess sup}_{x \in (0,\infty)} \|v f\|_{p',(0,x)} \|u\|_{r,(0,x)}^{-1} \|w\|_{q,(x,\infty)} \\ &= \text{ess sup}_{x \in (0,\infty)} \|f\|_{p',v,(0,x)} \|u\|_{r,(0,x)}^{-1} \|w\|_{q,(x,\infty)} \\ &= \|f\|_{\text{Ces}_{p',\infty}(\omega, v)}. \end{aligned}$$

(ii) By [8, Theorem 2.8, (ii)], we have that

$$\begin{aligned} &\|I\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w,v)} \\ &\approx \text{ess sup}_{x \in (0,\infty)} \left( \text{ess sup}_{t \in (0,\infty)} \mathcal{V}(x,t) V(t) \|u\|_{r,(0,t)}^{-1} \right) \left( \int_0^\infty \mathcal{V}(t,x)^{q'} \left( \int_t^\infty w^q dt \right)^{\frac{1}{q'}} \right). \end{aligned}$$

Using Lemma 2.2 with

$$\beta = q', \quad a(t) = V(t), \quad g(t) = V(t) \|u\|_{r,(0,t)}^{-1}, \quad \text{and} \quad h(t) = \left( \int_t^\infty w^q dt \right)^{\frac{1}{q'}} w^q(t) dt \quad (t > 0),$$

in view of (2.1), we arrive at

$$\begin{aligned} &\|I\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w,v)} \\ &\approx \text{ess sup}_{x \in (0,\infty)} V(x) \|u\|_{r,(0,x)}^{-1} \left( \int_x^\infty \left( \int_t^\infty w^q dt \right)^{\frac{1}{q'}} w^q(t) dt \right)^{\frac{1}{q'}} \\ &\quad + \text{ess sup}_{x \in (0,\infty)} \|u\|_{r,(0,x)}^{-1} \left( \int_0^x V(t)^{q'} \left( \int_t^\infty w^q dt \right)^{\frac{1}{q'}} w^q(t) dt \right)^{\frac{1}{q'}} \\ &\approx \text{ess sup}_{x \in (0,\infty)} V(x) \|u\|_{r,(0,x)}^{-1} \|w\|_{q,(x,\infty)} \\ &\quad + \text{ess sup}_{x \in (0,\infty)} \|u\|_{r,(0,x)}^{-1} \left( \int_0^x V(t)^{q'} \left( \int_t^\infty w^q dt \right)^{\frac{1}{q'}} w^q(t) dt \right)^{\frac{1}{q'}}. \end{aligned}$$

Hence

$$\begin{aligned}
& \|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w,v))} \\
& \approx \text{ess sup}_{x \in (0,\infty)} \|vf\|_{p',(0,x)} \|u\|_{r,(0,x)}^{-1} \|w\|_{q,(x,\infty)} \\
& \quad + \text{ess sup}_{x \in (0,\infty)} \|u\|_{r,(0,x)}^{-1} \left( \int_0^x \|vf\|_{p',(0,t)}^{q'} \left( \int_t^\infty w^q(t) dt \right)^{\frac{1}{q'}} \right) \\
& = \text{ess sup}_{x \in (0,\infty)} \|f\|_{p',v,(0,x)} \|u\|_{r,(0,x)}^{-1} \|w\|_{q,(x,\infty)} \\
& \quad + \text{ess sup}_{x \in (0,\infty)} \|u\|_{r,(0,x)}^{-1} \left( \int_0^x \|f\|_{p',v,(0,t)}^{q'} \left( \int_t^\infty w^q(t) dt \right)^{\frac{1}{q'}} \right) \\
& = \left\| \|f\|_{p',v,(0,x)} \right\|_{\infty, \omega_1, (0,\infty)} + \left\| \left\| \|f\|_{p',v,(0,t)} \right\|_{q', \omega_3, (0,x)} \right\|_{\infty, \omega_2, (0,\infty)} \\
& = \|f\|_{\text{Ces}_{p',\infty}(\omega_1, v)} + \|f\|_{\text{Ces}_{p',q',\infty}(\omega_2, \omega_3, v)}.
\end{aligned}$$

□

**Theorem 3.10.** Let  $0 < p, q, r < \infty$ ,  $p < 1$  and  $p < \min\{r, q\}$ . Let  $v \in \mathcal{W}(0, \infty)$ ,  $u \in {}^c\Omega_r$  and  $w \in \Omega_q$ . Assume that  $V$  is admissible and  $\varphi_2 \in Q_{V^{1/p'}}$ , where

$$\varphi_2(x) := \left( \int_{(0,\infty)} [\mathcal{V}(x,t)V(t)]^{r \rightarrow p} d(-\|u\|_{r,(0,t)}^{-r \rightarrow p}) \right)^{\frac{1}{r \rightarrow p}}.$$

(i) If  $\max\{1, r\} \leq q$ , then

$$\begin{aligned}
& M(\text{Cop}_r(u), \text{Ces}_{p,q}(w,v)) \\
& = \begin{cases} \text{Ces}_{p',r \rightarrow p,\infty}(\omega_1, \omega_2, v) \cap \text{Ces}_{p',\infty}(\omega_3, v) \cap \text{Ces}_{p',\infty}(\omega_1, v) & \text{if } \|u\|_{r,(0,\infty)} < \infty, \\ \text{Ces}_{p',r \rightarrow p,\infty}(\omega_1, \omega_2, v) \cap \text{Ces}_{p',\infty}(\omega_3, v) & \text{if } \|u\|_{r,(0,\infty)} = \infty. \end{cases}
\end{aligned}$$

where

$$\omega_1 := A_{q,q}^*(w), \quad \omega_2 := A_{r,p}(u), \quad \omega_3 := A_{r \rightarrow p, r \rightarrow p}^*(A_{r,p}(u)) \cdot A_{q,q}^*(w).$$

Moreover,

$$\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w,v))} \approx \|f\|_{\text{Ces}_{p',r \rightarrow p,\infty}(\omega_1, \omega_2, v)} + \|f\|_{\text{Ces}_{p',\infty}(\omega_3, v)} + \|u\|_{r,(0,\infty)}^{-1} \|f\|_{\text{Ces}_{p',\infty}(\omega_1, v)}.$$

(ii) If  $1 \leq q < r$ , then

$$\begin{aligned}
& M(\text{Cop}_r(u), \text{Ces}_{p,q}(w,v)) \\
& = \begin{cases} \text{Ces}_{p',r \rightarrow p,r \rightarrow q}(\omega_1, \omega_2, v) \cap \text{Ces}_{p',\infty,r \rightarrow q}(\omega_3, \omega_4, v) \cap \text{Ces}_{p',\infty}(\omega_4, v) & \text{if } \|u\|_{r,(0,\infty)} < \infty, \\ \text{Ces}_{p',r \rightarrow p,r \rightarrow q}(\omega_1, \omega_2, v) \cap \text{Ces}_{p',\infty,r \rightarrow q}(\omega_3, \omega_4, v) & \text{if } \|u\|_{r,(0,\infty)} = \infty, \end{cases}
\end{aligned}$$

where

$$\omega_1 := A_{q,r}^*(w), \quad \omega_2 := A_{r,p}(u), \quad \omega_3 := A_{q,q}^*(w), \quad \omega_4 := \left( A_{r \rightarrow p, r \rightarrow p}^*(A_{r,p}(u)) \right)^{\frac{r \rightarrow p}{q \rightarrow p}} \cdot A_{r,p}(u)^{\frac{r \rightarrow p}{r \rightarrow q}}.$$

Moreover

$$\begin{aligned} \|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} &\approx \|f\|_{\text{Ces}_{p', r \rightarrow p, r \rightarrow q}(\omega_1, \omega_2, v)} + \|f\|_{\text{Ces}_{p', \infty, r \rightarrow q}(\omega_3, \omega_4, v)} + \|u\|_{r, (0, \infty)}^{-1} \|f\|_{\text{Ces}_{p', \infty}(\omega_4, v)}. \end{aligned}$$

(iii) If  $r \leq q < 1$ , then

$$M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v)) = \begin{cases} \text{Ces}_{p', r \rightarrow p, \infty}(\omega_1, \omega_2, v) \cap \text{Ces}_{p', q', \infty}(\omega_3, \omega_4, v) \cap \text{Ces}_{p', q'}(\omega_4, v) & \text{if } \|u\|_{r, (0, \infty)} < \infty, \\ \text{Ces}_{p', r \rightarrow p, \infty}(\omega_1, \omega_2, v) \cap \text{Ces}_{p', q', \infty}(\omega_3, \omega_4, v) & \text{if } \|u\|_{q_1, (0, \infty)} = \infty, \end{cases}$$

where

$$\omega_1 := A_{q,q}^*(w), \quad \omega_2 := A_{r,p}(u), \quad \omega_3 := A_{r \rightarrow p, r \rightarrow p}^*(A_{r,p}(u)), \quad \omega_4 := A_{q,1}^*(w).$$

Moreover

$$\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} \approx \|f\|_{\text{Ces}_{p', r \rightarrow p, \infty}(\omega_1, \omega_2, v)} + \|f\|_{\text{Ces}_{p', q', \infty}(\omega_3, \omega_4, v)} + \|u\|_{r, (0, \infty)}^{-1} \|f\|_{\text{Ces}_{p', q'}(\omega_4, v)}.$$

(iv) If  $q < \min\{1, r\}$ , then

$$\begin{aligned} M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v)) &= \begin{cases} \text{Ces}_{p', r \rightarrow p, r \rightarrow q}(\omega_1, \omega_2, v) \cap \text{Ces}_{p', q', r \rightarrow q}(\omega_3, \omega_4, v) \cap \text{Ces}_{p', q'}(\omega_4, v) & \text{if } \|u\|_{r, (0, \infty)} < \infty, \\ \text{Ces}_{p', r \rightarrow p, r \rightarrow q}(\omega_1, \omega_2, v) \cap \text{Ces}_{p', q', r \rightarrow q}(\omega_3, \omega_4, v) & \text{if } \|u\|_{r, (0, \infty)} = \infty, \end{cases} \end{aligned}$$

where

$$\omega_1 := A_{q,r}^*(w), \quad \omega_2 := A_{r,p}(u), \quad \omega_3 := \left( A_{r \rightarrow p, r \rightarrow p}^*(A_{r,p}(u)) \right)^{\frac{r \rightarrow p}{q \rightarrow p}} \cdot A_{r,p}(u)^{\frac{r \rightarrow p}{r \rightarrow q}}, \quad \omega_4 := A_{q,1}^*(w).$$

Moreover

$$\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} \approx \|f\|_{\text{Ces}_{p', r \rightarrow p, r \rightarrow q}(\omega_1, \omega_2, v)} + \|f\|_{\text{Ces}_{p', q', r \rightarrow q}(\omega_3, \omega_4, v)} + \|u\|_{r, (0, \infty)}^{-1} \|f\|_{\text{Ces}_{p', q'}(\omega_4, v)}.$$

*Proof.* (i) Let  $\max\{1, r\} \leq q$ . Applying Lemma 2.3 with

$$\beta = r \rightarrow p, \quad a(t) = V(t), \quad g(t) = V(t)^{r \rightarrow p} \left( \int_0^t u^r \right)^{\frac{r}{r-p}} u^r(t) \text{ and } h(t) = \|w\|_{q, (t, \infty)} \quad (t > 0),$$

by [8, Theorem 2.9, (i)], we obtain that

$$\begin{aligned} \|I\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w, v)} &\approx \sup_{x \in (0, \infty)} \left( \int_{(0, x)} V(t)^{r \rightarrow p} d(-\|u\|_{r, (0, t)}^{-r \rightarrow p}) \right)^{\frac{1}{r \rightarrow p}} \|w\|_{q, (x, \infty)} \\ &\quad + \sup_{x \in (0, \infty)} \left( \int_{(x, \infty)} d(-\|u\|_{r, (0, t)}^{-r \rightarrow p}) \right)^{\frac{1}{r \rightarrow p}} V(x) \|w\|_{q, (x, \infty)} \\ &\quad + \|u\|_{r, (0, \infty)}^{-1} \sup_{t \in (0, \infty)} V(t) \|w\|_{q, (t, \infty)} \end{aligned}$$

Consequently,

$$\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} \approx \left\| \|f\|_{p', v, (0, \cdot)} \right\|_{r \rightarrow p, \omega_2, (0, \cdot)} + \left\| \|f\|_{p', v, (0, \cdot)} \right\|_{\infty, \omega_1, (0, \infty)}$$

$$\begin{aligned}
& + \|u\|_{r,(0,\infty)}^{-1} \left\| \|f\|_{p',v,(0,\cdot)} \right\|_{\infty,\omega_1,(0,\infty)} \\
& = \|f\|_{\text{Ces}_{p',r \rightarrow p,\infty}(\omega_1,\omega_2,v)} + \|f\|_{\text{Ces}_{p',\infty}(\omega_3,v)} \\
& \quad + \|u\|_{r,(0,\infty)}^{-1} \|f\|_{\text{Ces}_{p',\infty}(\omega_1,v)}.
\end{aligned}$$

(ii) Let  $1 \leq q < r$ . Applying Lemma 2.9 with

$$\beta = \frac{r \rightarrow q}{r \rightarrow p}, \quad a(x) = V(x)^{r \rightarrow p}, \quad g(t) = V(t)^{r \rightarrow p} \left( \int_0^t u^r \right)^{\frac{-r}{r-p}} u^r(t), \quad \text{and } h(t) = \|w\|_{q,(t,\infty)}^{r \rightarrow p},$$

noting that  $\beta - 1 = \frac{r \rightarrow q}{q \rightarrow p}$ , by [8, Theorem 2.9, (ii)], we obtain that

$$\begin{aligned}
& \|I\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w,v)} \\
& \approx \left( \int_{(0,\infty)} \left( \int_{(0,x)} V(t)^{r \rightarrow p} d\left(-\|u\|_{r,(0,t)}^{-r \rightarrow p}\right) \right)^{\frac{r \rightarrow q}{q \rightarrow p}} \|w\|_{q,(x,\infty)}^{r \rightarrow q} V(x)^{r \rightarrow p} d\left(-\|u\|_{r,(0,x)}^{-r \rightarrow p}\right) \right)^{\frac{1}{r \rightarrow q}} \\
& \quad + \left( \int_{(0,\infty)} \left( \sup_{t \in (0,x)} V(t)^{r \rightarrow p} \|w\|_{q,(t,\infty)}^{r \rightarrow p} \right)^{\frac{r \rightarrow q}{q \rightarrow p}} \left( \int_{(x,\infty)} d\left(-\|u\|_{r,(0,t)}^{-r \rightarrow p}\right) \right)^{\frac{r \rightarrow q}{q \rightarrow p}} d\left(-\|u\|_{r,(0,x)}^{-r \rightarrow p}\right) \right)^{\frac{1}{r \rightarrow q}} \\
& \quad + \|u\|_{r,(0,\infty)}^{-1} \sup_{t \in (0,\infty)} V(t) \|w\|_{q,(t,\infty)}.
\end{aligned}$$

Integrating by parts in the first integral on the right hand side, we arrive at

$$\begin{aligned}
& \|I\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w,v)} \\
& \approx \left( \int_{(0,\infty)} \left( \int_{(0,x)} V(t)^{r \rightarrow p} d\left(-\|u\|_{r,(0,t)}^{-r \rightarrow p}\right) \right)^{\frac{r \rightarrow q}{q \rightarrow p}} d\left(-\|w\|_{q,(x,\infty)}^{r \rightarrow q}\right) \right)^{\frac{1}{r \rightarrow q}} \\
& \quad + \left( \int_{(0,\infty)} \left( \sup_{t \in (0,x)} V(t)^{r \rightarrow p} \|w\|_{q,(t,\infty)}^{r \rightarrow p} \right)^{\frac{r \rightarrow q}{q \rightarrow p}} \left( \int_{(x,\infty)} d\left(-\|u\|_{r,(0,t)}^{-r \rightarrow p}\right) \right)^{\frac{r \rightarrow q}{q \rightarrow p}} d\left(-\|u\|_{r,(0,x)}^{-r \rightarrow p}\right) \right)^{\frac{1}{r \rightarrow q}} \\
& \quad + \|u\|_{r,(0,\infty)}^{-1} \sup_{t \in (0,\infty)} V(t) \|w\|_{q,(t,\infty)}.
\end{aligned}$$

Thus

$$\begin{aligned}
\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w,v))} & \approx \left\| \left\| \|f\|_{p',v,(0,\cdot)} \right\|_{r \rightarrow p,\omega_2,(0,\cdot)} \right\|_{r \rightarrow q,\omega_1,(0,\infty)} + \left\| \left\| \|f\|_{p',v,(0,\cdot)} \right\|_{\infty,\omega_4,(0,\cdot)} \right\|_{r \rightarrow q,\omega_3,(0,\infty)} \\
& \quad + \|u\|_{r,(0,\infty)}^{-1} \left\| \|f\|_{p',v,(0,\cdot)} \right\|_{\infty,\omega_4,(0,\infty)} \\
& = \|f\|_{\text{Ces}_{p',r \rightarrow p,r \rightarrow q}(\omega_1,\omega_2,v)} + \|f\|_{\text{Ces}_{p',\infty,r \rightarrow q}(\omega_3,\omega_4,v)} + \|u\|_{r,(0,\infty)}^{-1} \|f\|_{\text{Ces}_{p',\infty}(\omega_4,v)}.
\end{aligned}$$

(iii) Let  $r \leq q < 1$ . Applying Lemma 2.7 with

$$\alpha = q', \quad \beta = r \rightarrow p, \quad a(x) = V(x), \quad g(t) = V(t)^{r \rightarrow p} \left( \int_0^t u^r \right)^{\frac{-r}{r-p}} u^r(t), \quad \text{and } h(t) = \left( \int_t^\infty w^q \right)^{q'} w^q(t),$$

by [8, Theorem 2.9, (iii)], we obtain that

$$\begin{aligned} \|I\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w,v)} &\approx \sup_{x \in (0,\infty)} \left( \int_{(0,x)} V(t)^{r \rightarrow p} d(-\|u\|_{r,(0,t)}^{-r \rightarrow p}) \right)^{\frac{1}{r \rightarrow p}} \left( \int_{(x,\infty)} d(-\|w\|_{q,(t,\infty)}^{r \rightarrow q}) \right)^{\frac{1}{q'}} \\ &+ \sup_{x \in (0,\infty)} \left( \int_{(x,\infty)} d(-\|u\|_{r,(0,t)}^{-r \rightarrow p}) \right)^{\frac{1}{r \rightarrow p}} \left( \int_{(0,x)} V(t)^{q'} d(-\|w\|_{q,(t,\infty)}^{q'}) \right)^{\frac{1}{q'}} \\ &+ \|u\|_{r,(0,\infty)}^{-1} \left( \int_{(0,\infty)} V(t)^{q'} d(-\|w\|_{q,(t,\infty)}^{q'}) \right)^{\frac{1}{q'}}. \end{aligned}$$

Consequently

$$\begin{aligned} \|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w,v))} &\approx \left\| \left\| \|f\|_{p',v,(0,\cdot)} \right\|_{r \rightarrow p, \omega_2, (0,\cdot)} \right\|_{\infty, \omega_1, (0,\infty)} + \left\| \left\| \|f\|_{p',v,(0,\cdot)} \right\|_{q', \omega_4, (0,\cdot)} \right\|_{\infty, \omega_3, (0,\infty)} \\ &+ \|u\|_{r,(0,\infty)}^{-1} \left\| \|f\|_{p',v,(0,\cdot)} \right\|_{q', \omega_4, (0,\infty)} \\ &= \|f\|_{\text{Ces}_{p',r \rightarrow p,\infty}(\omega_1, \omega_2, v)} + \|f\|_{\text{Ces}_{p',q',\infty}(\omega_3, \omega_4, v)} + \|u\|_{r,(0,\infty)}^{-1} \|f\|_{\text{Ces}_{p',q'}(\omega_4, v)}. \end{aligned}$$

(iv) Let  $q < \min\{1, r\}$ . Applying Lemma 2.8 with

$$\alpha = r \rightarrow p, \beta = q', \gamma = r \rightarrow q,$$

and

$$a(x) = V(x), g(t) = V(t)^{r \rightarrow p} \left( \int_0^t u^r \right)^{\frac{-r}{r-p}} u^r(t), h(t) = \left( \int_t^\infty w^q \right)^{q'} w^q(t)$$

by [8, Theorem 2.9, (iv)], we obtain that

$$\begin{aligned} \|I\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w,v)} &\approx \left( \int_{(0,\infty)} \left( \int_{(0,x)} V(t)^{r \rightarrow p} d(-\|u\|_{r,(0,t)}^{-r \rightarrow p}) \right)^{\frac{r \rightarrow q}{r \rightarrow p}-1} \|w\|_{q,(x,\infty)}^{r \rightarrow q} V(x)^{r \rightarrow p} d(-\|u\|_{r,(0,x)}^{-r \rightarrow p}) \right)^{\frac{1}{r \rightarrow q}} \\ &+ \left( \int_{(0,\infty)} \left( \int_{(x,\infty)} d(-\|u\|_{r,(0,t)}^{-r \rightarrow p}) \right)^{\frac{r \rightarrow q}{r \rightarrow p}-1} \left( \int_{(0,x)} V(t)^{q'} d(-\|w\|_{q,(t,\infty)}^{q'}) \right)^{\frac{r \rightarrow q}{q'}} d(-\|u\|_{r,(0,x)}^{-r \rightarrow p}) \right)^{\frac{1}{r \rightarrow q}} \\ &+ \|u\|_{r,(0,\infty)}^{-1} \left( \int_{(0,\infty)} V(t)^{q'} d(-\|w\|_{q,(t,\infty)}^{q'}) \right)^{\frac{1}{q'}}. \end{aligned}$$

Integrating by parts at the first integral on the right hand side, we get that

$$\begin{aligned} \|I\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w,v)} &\approx \left( \int_{(0,\infty)} \left( \int_{(0,x)} V(t)^{r \rightarrow p} d(-\|u\|_{r,(0,t)}^{-r \rightarrow p}) \right)^{\frac{r \rightarrow q}{r \rightarrow p}} d(-\|w\|_{q,(x,\infty)}^{r \rightarrow q}) \right)^{\frac{1}{r \rightarrow q}} \\ &+ \left( \int_{(0,\infty)} \left( \int_{(x,\infty)} d(-\|u\|_{r,(0,t)}^{-r \rightarrow p}) \right)^{\frac{r \rightarrow q}{q \rightarrow p}} \left( \int_{(0,x)} V(t)^{q'} d(-\|w\|_{q,(t,\infty)}^{q'}) \right)^{\frac{r \rightarrow q}{q'}} d(-\|u\|_{r,(0,x)}^{-r \rightarrow p}) \right)^{\frac{1}{r \rightarrow q}} \\ &+ \|u\|_{r,(0,\infty)}^{-1} \left( \int_{(0,\infty)} V(t)^{q'} d(-\|w\|_{q,(t,\infty)}^{q'}) \right)^{\frac{1}{q'}}. \end{aligned}$$

Consequently

$$\begin{aligned} \|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w,v))} &\approx \left\| \left\| \|f\|_{p',v,(0,\cdot)} \right\|_{r \rightarrow p, \omega_2, (0,\cdot)} \right\|_{r \rightarrow q, \omega_1, (0,\infty)} + \left\| \left\| \|f\|_{p',v,(0,\cdot)} \right\|_{q',\omega_4, (0,\cdot)} \right\|_{\infty, \omega_3, (0,\infty)} \\ &\quad + \|u\|_{r,(0,\infty)}^{-1} \left\| \|f\|_{p',v,(0,\cdot)} \right\|_{q',\omega_4, (0,\infty)} \\ &= \|f\|_{\text{Ces}_{p',r \rightarrow p,r \rightarrow q}(\omega_1, \omega_2, v)} + \|f\|_{\text{Ces}_{p',q',r \rightarrow q}(\omega_3, \omega_4, v)} + \|u\|_{r,(0,\infty)}^{-1} \|f\|_{\text{Ces}_{p',q'}(\omega_4, v)}. \end{aligned}$$

□

**Theorem 3.11.** Let  $0 < r < p = 1 < q < \infty$ . Assume that  $v \in \mathcal{W}(0, \infty)$ ,  $u \in {}^c\Omega_r$  and  $w \in \Omega_q$ . Then

$$M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v)) = L_\infty(\omega),$$

where

$$\omega := A_{r,r}(u) \cdot A_{q,q}^*(w) \cdot v.$$

Moreover

$$\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} \approx \|f\|_{\infty, \omega, (0, \infty)}.$$

*Proof.* By [8, Theorem 2.10], on using (2.1), we obtain that

$$\begin{aligned} \|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} &\approx \sup_{t \in (0, \infty)} \left\| \|u\|_{r,(0,\cdot)}^{-1} \right\|_{\infty, f_v, (0,t)} \|w\|_{q,(t,\infty)} \\ &= \sup_{t \in (0, \infty)} f(t)v(t) \|u\|_{r,(0,t)}^{-1} \|w\|_{q,(t,\infty)} \\ &= \|f\|_{\infty, \omega, (0, \infty)}. \end{aligned}$$

□

**Theorem 3.12.** Let  $0 < r, q < \infty$ , and  $1 = p < \min\{r, q\}$ . Let  $u \in {}^c\Omega_r$  and  $w \in \Omega_q$ . Assume that  $v \in \mathcal{W}(0, \infty) \cap C(0, \infty)$  and  $0 < \|w^{-1}\|_{q',(x,\infty)} < \infty$ ,  $x > 0$ .

(i) If  $r \leq q$ , then

$$\begin{aligned} M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v)) \\ = \begin{cases} \text{Ces}_{\infty, r', \infty}(\omega_1, \omega_2, v) \cap L_\infty(\omega_3) \cap L_\infty(\omega_4) & \text{if } \|u\|_{r,(0,\infty)} < \infty, \\ \text{Ces}_{\infty, r', \infty}(\omega_1, \omega_2, v) \cap L_\infty(\omega_3) & \text{if } \|u\|_{r,(0,\infty)} = \infty, \end{cases} \end{aligned}$$

where

$$\omega_1 := A_{q,q}^*(w), \quad \omega_2 := A_{r,1}(u), \quad \omega_3(x) := v(x) \left\| A_{r',r'}^*(A_{r,1}(u))(t) \cdot \omega_1(t) \right\|_{\infty, (x,\infty)}, \quad \omega_4 := A_{q,q}^*(w) \cdot v.$$

Moreover

$$\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} \approx \|f\|_{\text{Ces}_{\infty, r', \infty}(\omega_1, \omega_2, v)} + \|f\|_{\infty, \omega_3, (0, \infty)} + \|u\|_{r,(0,\infty)}^{-1} \|f\|_{\infty, \omega_4, (0, \infty)}.$$

(ii) If  $q < r$ , then

$$\begin{aligned} M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v)) \\ = \begin{cases} \text{Ces}_{\infty, r', r \rightarrow q}(\omega_1, \omega_2, v) \cap \text{Ces}_{\infty, \infty, r \rightarrow q}(\omega_3, \omega_4, v) \cap L_\infty(\omega_5) & \text{if } \|u\|_{r,(0,\infty)} < \infty, \\ \text{Ces}_{\infty, r', r \rightarrow q}(\omega_1, \omega_2, v) \cap \text{Ces}_{\infty, \infty, r \rightarrow q}(\omega_3, \omega_4, v) & \text{if } \|u\|_{r,(0,\infty)} = \infty, \end{cases} \end{aligned}$$

where

$$\begin{aligned}\omega_1 &:= A_{q,r}^*(w), \quad \omega_2 := A_{r,1}(u), \quad \omega_3 := [A_{r',r'}^*(A_{r,1}(u))]^{\frac{r'}{q'}} \cdot [A_{r,1}(u)]^{\frac{r'}{r \rightarrow q}}, \\ \omega_4 &:= A_{q,q}^*(w), \quad \omega_5 := A_{q,q}^*(w) \cdot v.\end{aligned}$$

Moreover

$$\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} \approx \|f\|_{\text{Ces}_{\infty, r', r \rightarrow q}(\omega_1, \omega_2, v)} + \|f\|_{\text{Ces}_{\infty, \infty, r \rightarrow q}(\omega_3, \omega_4, v)} + \|u\|_{r, (0, \infty)}^{-1} \|f\|_{\infty, \omega_5, (0, \infty)}.$$

*Proof.* (i) Let  $r \leq q$ . Applying Lemma 2.3 with

$$\beta = r', \quad a(t) = V(t), \quad g(t) = V(t)^{r'} d\left(-\|u\|_{r, (0, t)}^{-r'}\right), \quad \text{and } h(t) = \|w\|_{q, (t, \infty)} \quad (t > 0),$$

by [8, Theorem 2.11, (i)], we obtain that

$$\begin{aligned}\|\mathbf{I}\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w, v)} &\approx \sup_{x \in (0, \infty)} \left( \int_{(0, x)} V(t)^{r'} d\left(-\|u\|_{r, (0, t)}^{-r'}\right) \right)^{\frac{1}{r'}} \|w\|_{q, (x, \infty)} \\ &\quad + \sup_{x \in (0, \infty)} V(x) \left( \int_{(x, \infty)} d\left(-\|u\|_{r, (0, t)}^{-r'}\right) \right)^{\frac{1}{r'}} \|w\|_{q, (x, \infty)} \\ &\quad + \|u\|_{r, (0, \infty)}^{-1} \sup_{x \in (0, \infty)} V(x) \|w\|_{q, (x, \infty)}.\end{aligned}$$

Consequently, in view of (2.1) and (2.2), we arrive at

$$\begin{aligned}\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w, v))} &\approx \left\| \left\| f \right\|_{\infty, v, (0, \cdot)} \right\|_{r', \omega_2, (0, \cdot)} + \|f\|_{\infty, \omega_3, (0, \infty)} + \|u\|_{r, (0, \infty)}^{-1} \|f\|_{\infty, \omega_4, (0, \infty)} \\ &= \|f\|_{\text{Ces}_{\infty, r', \infty}(\omega_1, \omega_2, v)} + \|f\|_{\infty, \omega_3, (0, \infty)} + \|u\|_{r, (0, \infty)}^{-1} \|f\|_{\infty, \omega_4, (0, \infty)}.\end{aligned}$$

(ii) Let  $q < r$ . Applying Lemma 2.9 with

$$\beta = \frac{r \rightarrow q}{r'}, \quad a(t) = V(t)^{r'}, \quad g(t) = V(t)^{r'} d\left(-\|u\|_{r, (0, t)}^{-r'}\right), \quad \text{and } h(t) = \|w\|_{q, (t, \infty)}^{r'} \quad (t > 0),$$

by [8, Theorem 2.11, (ii)], we obtain that

$$\begin{aligned}\|\mathbf{I}\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w, v)} &\approx \left( \int_{(0, \infty)} \left( \int_{(0, x)} V(t)^{r'} d\left(-\|u\|_{r, (0, t)}^{-r'}\right) \right)^{\frac{r \rightarrow q}{q'}} \|w\|_{q, (x, \infty)}^{r \rightarrow q} V(x)^{r'} d\left(-\|u\|_{r, (0, x)}^{-r'}\right) \right)^{\frac{1}{r \rightarrow q}} \\ &\quad + \left( \int_{(0, \infty)} \left( \sup_{t \in (0, x)} V(t)^{r'} \|w\|_{q, (t, \infty)}^{r'} \right)^{\frac{r \rightarrow q}{r'}} \left( \int_{(x, \infty)} d\left(-\|u\|_{r, (0, t)}^{-r'}\right) \right)^{\frac{r \rightarrow q}{q'}} d\left(-\|u\|_{r, (0, x)}^{-r'}\right) \right)^{\frac{1}{r \rightarrow q}} \\ &\quad + \|u\|_{r, (0, \infty)}^{-1} \sup_{t \in (0, \infty)} V(t) \|w\|_{q, (t, \infty)}.\end{aligned}$$

Integrating by parts at the first integral on the right hand side, we get that

$$\begin{aligned}\|\mathbf{I}\|_{\text{Cop}_r(u) \rightarrow \text{Ces}_{p,q}(w, v)} &\approx \left( \int_{(0, \infty)} \left( \int_{(0, x)} V(t)^{r'} d\left(-\|u\|_{r, (0, t)}^{-r'}\right) \right)^{\frac{r \rightarrow q}{r'}} d\left(-\|w\|_{q, (x, \infty)}^{r \rightarrow q}\right) \right)^{\frac{1}{r \rightarrow q}}$$

$$\begin{aligned}
& + \left( \int_{(0,\infty)} \left( \sup_{t \in (0,x)} V(t)^{r'} \|w\|_{q,(t,\infty)}^{r'} \right)^{\frac{r \rightarrow q}{r'}} \left( \int_{(x,\infty)} d(-\|u\|_{r,(0,t)}^{-r'})^{\frac{r \rightarrow q}{q'}} d(-\|u\|_{r,(0,x)}^{-r'})^{\frac{1}{r \rightarrow q}} \right. \right. \\
& \left. \left. + \|u\|_{r,(0,\infty)}^{-1} \sup_{t \in (0,\infty)} V(t) \|w\|_{q,(t,\infty)} \right) \right.
\end{aligned}$$

Therefore, in view of (2.1), we arrive at

$$\begin{aligned}
\|f\|_{M(\text{Cop}_r(u), \text{Ces}_{p,q}(w,v))} & \approx \left\| \left\| \|f\|_{\infty, v, (0,\cdot)} \right\|_{r', \omega_2, (0,\cdot)} \right\|_{r \rightarrow q, \omega_1, (0,\infty)} + \left\| \left\| \|f\|_{\infty, v, (0,\cdot)} \right\|_{\infty, \omega_4, (0,\cdot)} \right\|_{r \rightarrow q, \omega_3, (0,\infty)} \\
& \quad + \|u\|_{r,(0,\infty)}^{-1} \|f\|_{\infty, \omega_5} \\
& = \|f\|_{\text{Ces}_{\infty, r', r \rightarrow q}(\omega_1, \omega_2, v)} + \|f\|_{\text{Ces}_{\infty, \infty, r \rightarrow q}(\omega_3, \omega_4, v)} + \|u\|_{r,(0,\infty)}^{-1} \|f\|_{\infty, \omega_5, (0,\infty)}.
\end{aligned}$$

□

## REFERENCES

- [1] R. Askey and R.P. Boas Jr., *Some integrability theorems for power series with positive coefficients*, Mathematical Essays Dedicated to A. J. Macintyre, Ohio Univ. Press, Athens, Ohio, 1970, pp. 23–32. MR0277956 (43 #3689)
- [2] S.V. Astashkin and L. Maligranda, *Structure of Cesàro function spaces*, Indag. Math. (N.S.) **20** (2009), no. 3, 329–379, DOI 10.1016/S0019-3577(10)00002-9. MR2639977 (2011c:46056)
- [3] ———, *Structure of Cesàro function spaces: a survey*, Banach Center Publ. **102** (2014), 13–40.
- [4] G. Bennett, *Factorizing the classical inequalities*, Mem. Amer. Math. Soc. **120** (1996), no. 576, viii+130, DOI 10.1090/memo/0576. MR1317938 (96h:26020)
- [5] R.P. Boas Jr., *Integrability theorems for trigonometric transforms*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 38, Springer-Verlag New York Inc., New York, 1967. MR0219973 (36 #3043)
- [6] ———, *Some integral inequalities related to Hardy's inequality*, J. Analyse Math. **23** (1970), 53–63. MR0274685 (43 #447)
- [7] A. Gogatishvili, A. Kufner, and L.-E. Persson, *Some new scales of weight characterizations of the class  $B_p$* , Acta Math. Hungar. **123** (2009), no. 4, 365–377. MR2506756
- [8] A. Gogatishvili, R.Ch. Mustafayev, and T. Ünver, *Embeddings between weighted Copson and Cesàro function spaces*, Czechoslovak Math. J. **67(142)** (2017), no. 4, 1105–1132. MR3736022
- [9] A. Gogatishvili, L.-E. Persson, V. D. Stepanov, and P. Wall, *Some scales of equivalent conditions to characterize the Stieltjes inequality: the case  $q < p$* , Math. Nachr. **287** (2014), no. 2-3, 242–253, DOI 10.1002/mana.201200118. MR3163577
- [10] A. Gogatishvili and L. Pick, *Discretization and anti-discretization of rearrangement-invariant norms*, Publ. Mat. **47** (2003), no. 2, 311–358. MR2006487
- [11] ———, *Embeddings and duality theorems for weak classical Lorentz spaces*, Canad. Math. Bull. **49** (2006), no. 1, 82–95. MR2198721
- [12] M.L. Goldman, H.P. Heinig, and V.D. Stepanov, *On the principle of duality in Lorentz spaces*, Canad. J. Math. **48** (1996), no. 5, 959–979. MR1414066
- [13] K.-G. Grosse-Erdmann, *The blocking technique, weighted mean operators and Hardy's inequality*, Lecture Notes in Mathematics, vol. 1679, Springer-Verlag, Berlin, 1998. MR1611898 (99d:26024)
- [14] A.A. Jagers, *A note on Cesàro sequence spaces*, Nieuw Arch. Wisk. (3) **22** (1974), 113–124. MR0348444 (50 #942)
- [15] A. Kamińska and D. Kubiak, *On the dual of Cesàro function space*, Nonlinear Anal. **75** (2012), no. 5, 2760–2773, DOI 10.1016/j.na.2011.11.019. MR2878472 (2012m:46034)
- [16] G.M. Leibowitz, *A note on the Cesàro sequence spaces*, Tamkang J. Math. **2** (1971), 151–157.
- [17] J.-S. Shiue, *On the Cesàro sequence spaces*, Tamkang J. Math. **1** (1970), no. 1, 19–25.

- [18] ———, *A note on Cesàro function space*, Tamkang J. Math. **1** (1970), no. 2, 91–95. MR0276751 (43 #2491)
- [19] P.W. Sy, W.Y. Zhang, and P.Y. Lee, *The dual of Cesàro function spaces*, Glas. Mat. Ser. III **22(42)** (1987), no. 1, 103–112 (English, with Serbo-Croatian summary). MR940098 (89g:46059)

Amiran Gogatishvili

Institute of Mathematics, Academy of Sciences of the Czech Republic, Žitná 25, 115 67 Praha 1,  
Czech Republic

E-mail: gogatish@math.cas.cz

Rza Mustafayev

Department of Mathematics, Faculty of Science, Karamanoglu Mehmetbey University, Karaman, 70100, Turkey

E-mail: rzamustafayev@gmail.com

Tugce Ünver

Department of Mathematics, Faculty of Science and Arts, Kirikkale University, 71450 Yahsihan,  
Kirikkale, Turkey

E-mail: tugceunver@gmail.com